Dosage compensation and meiotic sex chromosome inactivation are maintained in the absence of selection

Darren J. Parker[1,2,*], Zoé Dumas[2], Rocío Gómez Lencero[3], Jean-Marc Aury[4], Marjorie Labedan[2], Patrick Tran Van[2], Benjamin Istace[4], Corinne Cruaud[5], Karine Labadie[5], Benjamin Noel[4], Susana Freitas[2], Jelisaveta Djordjevic[2], and Tanja Schwander[2,*]

1. School of Natural and Environmental Sciences, Bangor University, Bangor, UK.
2. Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland.
3. Department of Biology, Universidad Autónoma de Madrid, 28049 Madrid, Spain
4. Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Université Evry, Université Paris-Saclay, 91057 Evry, France
5. Genoscope, Institut François Jacob, CEA, CNRS, Université Evry, Université Paris-Saclay, 91057 Evry, France

* co-corresponding authors: Darren J. Parker (d.parker@bangor.ac.uk) and Tanja Schwander (tanja.schwander@unil.ch)

Keywords: parthenogenesis, Dosage compensation, meiotic sex chromosome inactivation

Classification: Biological Sciences / Evolution

Abstract

Dosage compensation and meiotic sex chromosome inactivation (MSCI) are key mechanisms regulating gene expression from the X chromosome in male-heterogametic species. While the convergent evolution of these mechanisms is well-documented, their evolutionary fate under relaxed selection remains poorly understood. Here, we test whether dosage compensation and MSCI persist following three independent transitions to parthenogenesis in stick insects, where selection on male phenotypes is relaxed. Using rare males occasionally produced by parthenogenetic females, chromosome-level genome assemblies, RNA-seq from multiple tissues, and immunocytochemistry, we find that dosage compensation is fully conserved across all nine studied somatic tissues. This is even the case in the oldest, approximately 1.5 million years old all-female lineage, and for tissue-specific genes for which dosage variation is not expected to be very deleterious. Meiotic X inactivation in the germline is also conserved. Surprisingly however, expression data and cytological markers indicate that MSCI signatures are even stronger in parthenogenetic males, a pattern likely driven by prolonged autosomal transcription during meiosis. These results indicate that X-targeting dosage compensation and MSCI are highly stable over evolutionary time and may be maintained in all-female lineages by a combination of evolutionary constraint, pleiotropy, or very weak selection, whereas autosomal expression during meiosis shifts rapidly under relaxed selection.

Significance Statement:

Sex chromosomes are regulated by specialized mechanisms that balance gene expression between males and females in somatic tissues (dosage compensation) and silence the X chromosome during male meiosis (meiotic sex chromosome inactivation, MSCI). These processes are thought to decay when they become redundant, yet suitable systems to test this prediction are scarce. We studied rare males from parthenogenetic stick insect lineages that have been evolving without sex for up to 1.5 million years. Surprisingly, both dosage compensation and MSCI remain fully functional, despite relaxed selection. Instead, we find altered autosomal expression during meiosis. Our results reveal that sex chromosome regulation is evolutionarily stable and constrained, persisting even after the loss of its purpose.

Introduction

Sexual reproduction is key to the diversity of life and the main driver of species' lifecycles. In eukaryotic species with distinct male and female sexes, sex determination is frequently governed by sex chromosomes such as X and Y chromosomes in male heterogametic species (1). Because males carry only a single X chromosome while females carry two, X chromosome expression is regulated by highly specialized mechanisms: dosage compensation (in somatic cells) and meiotic sex chromosome inactivation (MSCI, in germ cells). Dosage compensation is a mechanism that balances the expression of X-linked and autosomal genes independently of the number of X chromosome copies. It has evolved independently across a wide range of taxa (2–4) including mammals (5), reptiles (6), fish (7, 8), nematodes (9), and insects (10–16). However, the X is typically not dosage compensated in gonads, where expression from the sex chromosome(s) is further often suppressed via MSCI (2, 12, 13, 17, 18). Specifically, MSCI involves the inactivation of heteromorphic sex chromosomes during meiosis through the addition of repressive chromatin marks, likely due to the inability of the X and the degraded (or completely lacking) Y to synapse at the onset of meiosis (19).

While far from ubiquitous (4, 20), dosage compensation and MSCI are typically maintained across clades for millions of years. Furthermore, a breadth of studies on both mechanisms indicate that they can evolve convergently as new sex chromosomes arise, involving similar or different mechanisms. This is revealed for example by research in dipterans, where X chromosomes have evolved independently from ancestral autosomes, with different dosage compensation mechanisms emerging in parallel (21–28). Similarly, the independent evolution of MSCI mechanisms in divergent taxa involves both shared and lineage-specific components (e.g. (29–31)).

In contrast to the evidence for the convergent *de novo* evolution of dosage compensation and MSCI, it remains unknown how ancestral dosage compensation and MSCI systems evolve once they become redundant, which would be the case upon the evolution of a novel sex chromosome system. This is in part because suitable model systems with sex chromosome turnovers and accompanying evolution of X-targeting regulatory mechanisms remain understudied. Furthermore, sex chromosome turnover events in species with established dosage compensation and MSCI systems are expected to be rare because the maintenance of active systems could cause extensive misregulation of gene expression on the newly-formed sex chromosomes (32–34). A unique opportunity to overcome such constraints is to study the fate and potential decay of dosage compensation and MSCI in all-female parthenogenetic species. Following a transition to parthenogenesis, the X chromosome effectively adopts an autosomal inheritance pattern as it no longer varies in copy numbers since all individuals are female and carry two X copies. As a result, if dosage compensation or MSCI is costly (because of the cost of the machinery or due to maladaptive expression in females) we would expect selection to reduce a species' ability to perform these mechanisms following a transition to parthenogenesis. In the absence of such costs in females, dosage compensation and MSCI would be expected to decay and vestigialize as a consequence of drift. A reduction of the two X regulatory mechanisms is difficult to observe in parthenogenetic species as we need to examine expression in both sexes. This barrier can be overcome in some systems if rare males are produced, a phenomenon frequently observed in parthenogenetic arthropods (35). The production of rare males would be impossible for

species where the perturbation of dosage compensation is lethal in males, as is the case for certain mutations in the *Drosophila* fly dosage compensation machinery (e.g., (36)). The suppression of dosage compensation does however not affect male viability in *Anopheles* mosquitoes, and instead causes a developmental delay (25). This variation between taxa illustrates that inference regarding the functioning and maintenance of dosage compensation cannot be based on the observation of male phenotypes in parthenogenetic species. With respect to X regulation in the germline, X-autosome translocations negatively affect fertility in *Drosophila* flies (37, 38), where X regulation happens via a mechanism that differs from classical MSCI as described in mammals and other insect taxa (17, 39). For other insects, fertility consequences of MSCI perturbation are unknown. As a consequence, fertility or sperm viability phenotypes in rare males do not provide information about the functioning of MSCI.

Here we take advantage of such rare males to examine dosage compensation and MSCI in three species of parthenogenetic *Timema* stick insects, using a combination of new chromosome-level genome assemblies, RNA sequencing of eight different tissues, and cell biology. *Timema* have had multiple independent transitions to parthenogenetic reproduction over the last 0.5-1.5 million years (40–43). Previously we have found that sexual *Timema* species show complete dosage compensation in somatic tissues throughout development, and MSCI in the male germline (11, 17, 44). This targeted regulation of the X chromosome likely involves general molecular mechanisms shared across insects (17) given that the stick insect X chromosome likely evolved prior to the diversification of insects over 450 mya (33). We can examine changes in dosage compensation and MSCI in parthenogenetic *Timema* as they occasionally produce rare males via two different processes. Some males arise from aneuploidy of the X chromosome (45), which since *Timema* have an XX/X0 sex-determination system, results in the production of a male. Others develop as a consequence of very rare instances of sex within parthenogenetic populations (46). Field and experimental evidence indicates that males produced by parthenogenetic *Timema* are exceedingly rare and exhibit reduced reproductive performance (45). During extensive field collections conducted over multiple years (2007–2012), only 11 males were recovered across several independently derived asexual lineages, despite the simultaneous collection of very large numbers of females. Functional assays of sperm performance further demonstrate that while rare males are capable of fertilization when mated as the sole partner of virgin females from closely related sexual species, reproductive success is low and they fail entirely under sperm competition (45). Thus, male production in parthenogenetic *Timema* is both extremely infrequent and associated with reduced reproductive performance. These features indicate relaxed selection on male-specific traits, while still allowing the occasional availability of males for mechanistic analyses of dosage compensation and meiotic sex chromosome inactivation. Contrary to our expectations we find that the targeted regulation of the X chromosome underlying dosage compensation and MSCI has been fully conserved in each of the three parthenogenetic lineages, with relaxed selection affecting autosomal rather than X-linked expression in male meiotic cells. These results indicate that selection against X-regulatory mechanisms is weak or that changes to dosage compensation and MSCI mechanisms are highly constrained.

Results and Discussion

Genome assemblies reveal a highly conserved X chromosome contrasted by autosomal fissions and fusions

We generated chromosome-level genome assemblies for two of the three sexual species of *Timema* used in this study, *T. cristinae* and *T. podura*, to complement the available assembly of *T. poppense* (Table S1). In both new assemblies, the number of large scaffolds assembled (hereafter referred to as chromosomes) matched previous karyotype information (40) and coverage of male- vs female-derived short reads were used to identify the X chromosome (Supplemental Info). Annotation using a combination of gene predictions, protein, and RNAseq evidence from each species and its parthenogenetic sister species and inference of 8118 one-to-one orthologs (see Methods) revealed that the X chromosome content has not changed in any of the sexual or asexual species, as expected given the long-term conservation of X chromosome content in stick insects (11, 47). Using comparisons with a stick insect (*Bacillus rossius*) from the Euphasmatodea clade (which split over 120 million years ago from *Timema* (48), we further find that even X chromosome synteny is highly conserved, while many rearrangements are apparent on the autosomes (Fig. 1). Specifically, the change in karyotype from 14 to 12 chromosomes during *Timema* evolution (40) involved at least two fissions and three fusions involving *T. podura* chromosomes 2, 3, 5 and 10 (Fig. 1, S1).

Fig. 1 | Autosomal genomic instability contrasts with X chromosome conservation. A. Simplified phylogeny of the used species, highlighting the independently evolved parthenogenetic *Timema* species and their closest sexual relatives. *Timema* are the sister clade to all other stick insects (including B. *rossius*), and branches reflecting this split have been shortened for display. B. Chord diagram showing synteny between the assemblies of sexual species, based on 4241 one-to-one orthologs across the focal *Timema* genomes and a distantly related stick insect (*Bacillus rossius*). See Fig. S1 for synteny based on 8118 one-to-one orthologs across *Timema* only.

Dosage compensation in somatic tissues is not reduced in parthenogenetic species

Dosage compensation between the sexes in parthenogenetic species was very similar to that observed in sexual species. In all seven non-reproductive tissues (antennae, brain, defence glands, gut, fat body, femur, and tarsi) we observe almost complete dosage compensation in all species (Fig. 2). As expected, we observe some variation in the expression of X-linked genes between the sexes for specific tissues and species (resulting in small but significant differences in expression of autosomal versus X-linked genes). Such variation is typical in studies of dosage compensation (e.g., (13, 49, 50) and can be interpreted as sex-specific regulation of individual genes rather than a consequence of variable or incomplete dosage compensation. If dosage compensation was reduced in parthenogenetic species, we would expect to see a larger deviation from complete dosage compensation in parthenogenetic species and an accompanying lower male-to-female expression ratio for the X in parthenogenetic species when compared to the sexual sister species for tissues with almost complete dosage compensation. We do not observe this, instead finding only small variations in X-linked expression with no consistent pattern (Fig. 2, S2-S11, Table S2), showing that dosage compensation is maintained across tissues even under relaxed selection for up to 1.5 million years. This conclusion extends to more in depth analyses: the amount of dosage compensation is consistent between the sexual and parthenogenetic species on a gene-by-gene basis (Fig. S2-S4), and we also find no evidence for decreased efficiency of dosage compensation for genes with tissue-specific expression (Fig. S5-S8), which are expected to be relatively less constrained than housekeeping genes (51).

Fig. 2 | Male to female expression ratios for the X and autosomes in non-reproductive tissues reveal the maintenance of dosage compensation under relaxed selection. Dashed lines represent a two- fold reduction in expression in males (as expected if there was no dosage compensation). A. Sexual species = *T. cristinae*, parthenogenetic species = *T. monikense*. B. Sexual species = *T. podura*, parthenogenetic species = *T. genevievae*. C. Sexual species = *T. poppense*, parthenogenetic species = *T. douglasi.* Asterisks indicate the significance (FDR) of Wilcoxon tests (***<0.001, **<0.01, *<0.05, NS > 0.05).

208
Parthenogenetic species no longer require sex-specific regulation of X chromosome copy
number across individuals. Under such conditions, the strength of selection acting on dosage
compensation mechanisms is expected to be substantially reduced. Whether this relaxation
should lead to detectable evolutionary decay depends on several unknown parameters,
including the number of loci involved, mutational target size, and the degree of pleiotropy. If
the machinery is costly or maladaptive in females, selection could favor its reduction.
Alternatively, if costs are negligible, its evolution would be governed largely by drift. The
absence of detectable change over up to 1.5 million years therefore indicates either unusually
slow neutral decay, functional constraint, or the action of weak purifying selection, but does
not allow us to distinguish conclusively among these alternatives. Support for functional
constraints because of pleiotropic effects in females stems from evidence for the maintenance
of dosage compensation-like mechanisms targeting the ancestral X in *Drosophila*
melanogaster fruit flies (52) even tens of myr after a sex chromosome turnover, which reverted
the ancestral X (now referred to as the dot) back to an autosome (15). Similar to our present
study in parthenogenetic species, such reversals should result in relaxed selection for dosage
compensation on the ancestral X (dot) chromosome. Yet, the dot is still regulated in a
chromosome-specific manner, distinct from the other autosomes and the novel X
chromosome. Thus, the dot is regulated via the protein POF (52), which might reflect vestiges
of the former dosage compensation machinery when the dot was still the X chromosome.
228
A second reason to expect changes in dosage compensation could be if dosage compensation
mechanisms result in maladaptive gene expression in females, i.e., partially unresolved sexual
antagonism (53). This factor may not have a strong influence in *Timema* as unresolved
antagonisms are expected to be transient (54) and the stick insect X chromosome is very old
and highly conserved (Fig. 1) and shares content with the X chromosome in other insect orders
that diverged over 450 million years ago (33). Any maladaptive gene expression in females
as a result of sexual conflict over dosage compensation would therefore be expected to be
very small.
237
While our results show that dosage compensation is maintained in parthenogenetic species
because selection in females against dosage compensation components is weak or because
changes to the underlying mechanisms are constrained, we cannot formally exclude that
perhaps not enough time has passed since the evolution of parthenogenesis to observe any
changes to dosage compensation. Particularly under a fully neutral-drift scenario, timescales
to observe decay could be longer for complex regulatory systems such as DC and MSCI (55,
56). Similarly, under a selective scenario (where there is selection against male-essential
components of dosage compensation in females) a lack of time may be a plausible explanation
in the young lineage *T. douglasi* and for the lack of apparent changes in dosage compensation
in whole-body samples of males of parthenogenetic aphid lineages (57), however, it is very
unlikely for the oldest species used here (*T. genevievae*) with an age estimate of over 1.5
million years (41, 42, 55).
250

A stronger signature of MSCI in parthenogenetic than sexual species driven by autosomal mis-expression during meiosis

We have shown previously that the X chromosome is consistently inactivated throughout male
meiosis in sexual *Timema* species (17, 44). Accordingly, we observe very low expression of
X-linked genes in the testes, much lower than the expectation for a lack of dosage
compensation (Fig. 3). We do however still observe some expression of the X in males, as
expected given the presence of non-meiotic cells in testes. Surprisingly, we see a consistently
lower expression of X-linked genes in males from parthenogenetic than sexual species (Fig.
3, S12). This could indicate variable cellular composition of testes such as a higher proportion
of meiotic cells in parthenogenetic than sexual males, or, more likely, that autosomal
expression is more persistent across meiosis in parthenogenetic as compared to sexual
males. In sexual *Timema* males, there are two bursts of autosomal transcription during meiosis
(44): a first at the end of prophase I (specifically, during pachytene and diplotene stages), and
a second during meiosis II at the spermatid stage.
266
To detect potential changes to these temporal patterns of autosomal expression in
parthenogenetic males, we performed immunolabeling of testis tissue of *T. monikense* and *T.*
douglasi using RNA polymerase II, a conserved transcriptional marker. This labeling revealed
that autosomal expression during prophase I is longer in parthenogenetic than sexual males.
In the former, the RNA polymerase II signal already appears at the onset of prophase I (i.e.,
during the zygotene stage; Fig. 4, S13) and is maintained until diplotene (Fig. 4, S13), while
in the latter it only appears at pachytene (44). Since males of parthenogenetic species are not
exposed to selection, this extension of the autosomal transcription burst during prophase I is
most likely a consequence of drift, i.e., it reflects autosomal mis-expression during meiosis in
parthenogenetic males.
277

Fig. 3 | A stronger signature of MSCI in parthenogenetic than sexual species. Gene
expression in reproductive tissues: Log$_2$ of male to female expression ratio for the X and
autosomes in testes and female gonads. Species codes: Tce = *T. cristinae*, Tms = *T.*
monikense, Tpa = *T. podura,* Tge = *T. genevievae,* Tps = *T. poppense,* Tdi = *T. douglasi.*
Asterisks indicate the significance (FDR) of Wilcoxon tests (***<0.001, **<0.01, *<0.05, NS >
0.05). Dashed lines represent a two- fold reduction in expression in males (as expected if there
was no dosage compensation).

Fig. 4 | Misregulation of autosomal gene expression in meiotic cells of rare males produced by parthenogenetic females. RNApol-II distribution in meiocytes of sexual (*T. californicum*) and parthenogenetically produced (*T. monikensis*) males. Projections of stack images acquired across squashed meiocytes at the stages indicated, stained with DAPI (blue) and double immunolabeled for SMC3 (green) and RNApol-II (magenta). RNApol-II labelling is already present at leptotene and early zygotene in parthenogenetic *T. monikensis* males, while it only appears during late zygotene in sexual *T. californicum* males, as shown previously (Dumas et al 2025). Note that the used immunolabeling allows for inference of qualitative (presence-absence) rather than quantitative (intensity of signals) differences (Fig. S13). Scale bar: 10 µm. Corresponding image analyses for males of *T. douglasi* are available in Fig. S13).

Conclusion

Our results demonstrate that both dosage compensation and meiotic sex chromosome inactivation (MSCI) are robustly maintained in parthenogenetic stick insects, despite the long-term absence of selective pressures typically associated with male phenotypes. This maintenance suggests that the molecular mechanisms underlying dosage compensation and MSCI are either functionally constrained (e.g. due to pleiotropic effects in females), maintained by weak purifying selection (for example through sufficiently frequent reproduction of rare males), or decay too slowly as a consequence of drift for phenotypic effects to be observed in the stick insects. Nevertheless, the enhanced signature of MSCI in parthenogenetic males appears to be driven by extended autosomal transcription during meiosis, revealing an unexpected consequence of relaxed selection on gene expression. Together, these findings highlight the stability and evolutionary inertia of complex regulatory processes, even in the face of dramatic shifts in reproductive mode.

Methods

Reference genome assemblies

Reference genome assemblies were produced for *T. cristinae* (Bioproject PRJNA1268975/ Accession JBOIUC000000000) and *T. podura* (Bioproject PRJNA1268975/ Accession JBOIUD000000000) from wild collected females (GPS coordinates 34°30'53.2"N, 119°46'45.3"W and 33°47'47.0"N, 116°46'34.6"W). Contigs for the *T. cristinae* assembly were based on PacBio sequencing (43x) and on a combination of Nanopore and Illumina sequencing for *T. podura* (sequenced at 85x respectively 62x coverage). Scaffolding was done using a Hi-C library (sequenced at approximately 69 x and 53 x coverage for *T. cristinae* and *T. podura* respectively) to produce the final reference assembly.

DNA extraction and library preparation

To extract high molecular weight (HMW) DNA, we flash-froze a single female (without gut) in liquid nitrogen and ground it using a Cryomill (Retsch). We then extracted HMW DNA using a G/20 Genomic Tips kit (Qiagen) following the manufacturer's protocols and checked DNA integrity on a pulse field agarose gel.

For *T. cristinae*, a PacBio library was prepared using the Template Prep Kit 1.0 following manufacturer's instructions and sequenced on a single cell of the PacBio SMRT™ system. For *T. podura*, four ONT libraries were prepared following Oxford Nanopore instructions. A first library was prepared using the SQK-LSK108 ligation sequencing kit and was loaded on a MinION R9.4 Flow Cells, and then three libraries were prepared using the SQK-LSK109 ligation sequencing kit and were loaded on three PromethION R9.4.1 Flow Cells. A PCR free Illumina library (based on the same HMW extraction) was prepared using the Kapa Hyper Prep Kit (Roche, Basel, Switzerland), following manufacturer instructions. The library was quantified by qPCR using the KAPA Library Quantification Kit for Illumina Libraries (Roche), and the library profile was assessed using a High Sensitivity DNA kit on an Agilent Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA). The library was then sequenced on an Illumina

343 HiSeq 4000 instrument (Illumina, San Diego, CA, USA), using 150 base-length read chemistry
344 in a paired-end mode.
345
346 A Hi-C library was prepared for each species using the Proximo Hi-C Kit. We generated
347 ground tissue for cross-linking using Cryomill (Retsch) and following manufacturer
348 instructions, using a different female than the one used for Nanopore and HiSeq sequencing,
349 from the same natural population. Library construction and sequencing (250 million read pairs)
350 was outsourced to Phase Genomics (Seattle).
351
352
353 **Assembly pipelines**
354
We used different strategies for genome assembly for the two species. For *T. cristinae* raw
PacBio reads were assembled into contigs using Canu (58) with the parameter
correctedErrorRate=0.105. For *T. podura* raw Oxford Nanopore reads were filtered using
Filtlong v0.2.0 (https://github.com/rrwick/Filtlong) with the parameters --min_length 1000 --
keep_percent 90 --target_bases 69050000000. The filtered Nanopore reads were then
assembled into contigs using Flye v2.8.1 (59) with --genome-size 1.381g. All Nanopore reads
were mapped against the contigs using minimap2 v2.19 (60) with the parameters -c -x map-
ont and a first step of polishing was performed using Racon v1.4.3 (61). Three additional
rounds of polishing were conducted: Illumina short reads were aligned to the contigs using
BWA mem v0.7.17 (62) and polishing was performed using Pilon v1.23 (63).
365
Both assemblies were then decontaminated using BlobTools v1.0 (64) under the taxrule
"bestsumorder". Hit files were generated after a blastn v2.10.1+ against the NBCI nt database,
searching for hits with an e-value below 1e-25 (parameters: -max_target_seqs 10 -max_hsps
1 -evalue 1e-25). Contigs without hits to metazoans were removed from the assemblies.
Haplotypic duplications were then filtered out as follows. Filtered reads were then mapped
against the decontaminated genomes using minimap2 and haplotigs were detected with Purge
Haplotigs v1.1.1 (65) using the parameters -l 3 -m 27 -h 190 -j 101 --editor-coarse-resolution
250000 for *T. cristinae* and -l 3 -m 45 -h 195 -j 101 --editor-coarse-resolution 50000 for *T.*
podura.
375
Hi-C reads from each species were then mapped against their respective haploid genome
using Juicer v1.6 (66) with the restriction site Sau3AI. Finally, chromosome-level scaffolding
was performed using 3D-DNA v180922 (67) with species-specific parameters, as
recommended by the authors. The resulting Hi-C contact matrices were visualized with
Juicebox, and minor polishing was conducted in accordance with these recommendations.
The completeness of both genome assemblies was assessed with BUSCO v5.1.2 (68) against
the *insecta_odb10* dataset using the --long and --augustus parameters.

Annotation
384
We annotated the *T. cristinae* and *T. podura* genomes in a similar way as we previously
annotated the *T. poppense* genome (17), using a combination of *ab initio* gene prediction,
protein homology, and RNA-seq using the Braker2 pipeline (v. 2.1.6) (69). Firstly, each
genome assembly was soft-masked using RepeatModeler (v. 2.0.2, options: -LTRStruct, -

engine ncbi) (70) and RepeatMasker (v. 4.1.2, options: -engine ncbi, -xsmall) (71). For protein
evidence, we used the arthropod protein sequences from OrthoDB (v.10.1) (72) and the
predicted protein sequences for *Timema* from our previous genome assemblies. For RNAseq
evidence, we used the RNAseq libraries generated here for *T. cristinae* and *T. monikense*
along with publicly available data (163 libraries generated here, 43 public (Bioproject
accessions: PRJNA380865, PRJNA679785, PRJNA1295360, PRJNA504764 (42, 43, 73))) to
annotate the *T. cristinae* genome and for *T. podura* and *T. genevievae* (119 libraries generated
here, 29 public (Bioproject accessions: PRJNA380865, PRJNA679785, PRJNA1295360 (42,
43))) to annotate the *T. podura* genome. This is a total of 354 RNAseq libraries (330 paired-
end, 24 single-end) covering over 19 different life stages, tissue, and sex combinations for
each species (see Table S3). Reads were quality trimmed with Trimmomatic (v. 0.39, options:
ILLUMINACLIP:3:25:6 LEADING:9 TRAILING:9 SLIDINGWINDOW:4:15 MINLEN:80) (74)
before mapping to the genome assembly with STAR (v. 2.7.8a, options: --twopassMode Basic)
(75). Braker2 was run using protein evidence and RNAseq separately with the gene predictors
Augustus (v. 3.4.0) (76) and Genemark (v. 4.72) (77). Following the RNAseq run, UTR
predictions were added to the RNAseq gene predictions using GUSHR (v. 1.0) in Braker2 (--
addUTR=on). The separate gene predictions were then merged using TSEBRA (v1.0.3) using
the pref_braker1.cfg configuration file, which weights RNAseq evidence more strongly than
the default option. We then ran BUSCO (v. 5.3.2, insecta_odb10) on the gene regions
annotated by Braker2 and on the whole genome assembly. Any genes found by BUSCO but
missed by Braker were then added to the annotation (69 genes for *T. cristinae* and 19 genes
for *T. podura*). ncRNA genes were predicted using Infernal (v. 1.1.2, minimum e-value 1e-10)
(78). GO-terms for protein-coding gene predictions were obtained using blastP within
OmicsBox (v3.1.2, default parameters) to blast the nr *Drosophila melanogaster* database
(taxonomy filter: 7227). 1-to-1 orthologs between were obtained for protein-coding genes
using OrthoFinder (version 2.5.5) (79) with default options.

Identification of the X chromosome

We used a coverage approach to identify X-linked scaffolds in our new genome assemblies.
To do this we mapped publicly available Illumina genome reads from males and females (4
males and 5 females for *T. cristinae* and 4 males and 2 females for *T. podura* (PRJNA670663,
PRJNA725673 (11, 43)) to the respective genome assemblies and their publicly available
mtDNA genomes (Accessions) using BWA-MEM (v. 0.7.15). Multi- mapping and poor quality
alignments were filtered (removing reads with XA:Z or SA:Z tags or a mapq <30). PCR
duplicates were removed with Picard (v. 2.9.0) (http://broadinstitute.github.io/picard/).
Coverage was then estimated for 100000 bp sliding windows using BEDTools (v. 2.26.0) (80).
To compare coverage between males and females, coverage was first summed for all male
and all female libraries per window. Male and female coverage was then normalized by modal
coverage to adjust for differences in overall coverage. X- linked windows were then identified
using the $\log_2$ ratio of male to female coverage. Autosomal windows should have equal
coverage in males and females (log2 ratio of male to female coverage ≈ 0) and X-linked
windows should have half the coverage in males as in females (log2 ratio of male to female
coverage ≈ -1). This approach identified one large scaffold in each assembly as the X
chromosome (Fig. S14-16).

We then repeated these analyses using genome reads from each of the parthenogenetic
species. Female reads were publicly available (5 for *T. monikense*, 6 for *T. genevievae*, and
5 for *T. douglasi*, PRJNA670663 (43)) as were the two male samples for *T. douglasi*
(PRJNA808673, (81)). For *T. monikense* and *T. genevievae* we also generated new Illumina
libraries for males (2 for *T. genevievae* and 4 for *T. monikense*) and an additional female
library for *T. genevievae* (Accession number: PRJNA1293928) as described previously (43)
using the carcasses after tissue dissection described below. Reads were then mapped to the
genome of each sexual sister species as described above (the sexual species genome for *T.*
douglasi is our previously published genome *T. poppense,* Accession number:
PRJNA1126215 (17)). Each of the newly generated samples were checked for normal
coverage distributions before calculating coverage ratios.
445

RNA extraction and sequencing

447
We generated a total of 282 RNA-seq libraries for eight female tissues (antennae, brain,
defence glands, fat body, femur, tarsi, and gonads) and nine male tissues (the same as the
female plus the male accessory glands) for *T. cristinae*, *T. podura*, *T. monikense*, and *T.*
genevievae following our previously developed methodology (17) (see Table S3 for details).
Individuals were collected as nymphs in the field and reared to adulthood in the laboratory.
Prior to dissection all individuals were isolated in Petri dishes and fed on an artificial diet (a
mix of flour, sugar, and water) for two days, allowing their gut contents to be cleared of plant
material. Insects were then anesthetized with CO_2, dissected, and tissues individually flash-
frozen in liquid nitrogen before being stored at -80°C.
457
TRIzol solution (1 mL) and a small amount of ceramic beads (Sigmund Lindner) were added
to each tissue. Samples were homogenized using a tissue homogenizer (Precellys Evolution;
Bertin Technologies). Chloroform (200 µL) was added to each sample and samples were then
vortexed for 15 seconds and centrifuged for 25 minutes at 12,000 rpm at 4°C. The upper
phase containing the RNA was then transferred to a new 1.5 mL tube with the addition of
isopropanol (650 µL). The samples were vortexed and placed at -20°C overnight. Samples
were then centrifuged for 30 minutes at 12,000 rpm at 4°C. The liquid supernatants were
removed, and the RNA pellet underwent two washes using 80% and 70% ethanol. Each wash
was followed by a 5-minute centrifugation step at 12,000 rpm. Finally, the RNA pellet was
resuspended in nuclease-free water and quantified using a fluorescent RNA-binding dye
(QuantiFluor RNA System) and nanodrop (DS-11 FX).
469
470 Library preparation using NEBNext (New England BioLabs) and sequencing on an Illumina
471 NovaSeq 6000 platform with 100 bp paired-end sequencing (36.2 million read pairs per
472 sample on average) was outsourced to a sequencing facility (Fasteris, Geneva).
473

Gene expression analyses

475
We mapped RNAseq reads from *T. cristinae*, *T. podura*, *T. monikense*, and *T. genevievae*
generated here and reads for *T. poppense* and *T. douglasi* we generated previously (17) to
their respective sexual species genomes as described above. This gave 4 replicates for each

of the 16 sex and tissue combinations for each of the six species, except for the male samples
of parthenogenetic species (*T. monikense, T. genevievae,* and *T. douglasi*) where we had
fewer replicates due to the difficulty of obtaining these specimens (n = 2-4 see Table S4 for
details). Read counts were obtained using HTSeq v.0.9.1 (82), with the following options
(htseq-count --order = name --type = exon --idattr = gene_id --stranded = reverse). Expression
analyses were performed using the Bioconductor package EdgeR (v 3.42.4) (83), and done
separately for each species and tissue. Normalization factors for each library were computed
using the TMM method and were used to calculate normalized expression levels (FPKM,
Fragments Per Kilobase of transcript per Million mapped reads). mtDNA, rRNA, tRNA and
genes with low expression (less than 2 FPKM in 2 or more libraries per sex in either the sexual
or parthenogenetic sister species) were excluded. This filtering was done to ensure that genes
used for gene expression comparisons between sexual and parthenogenetic sister species
and between males and females were the same. To examine dosage compensation, we used
the $\log_2$ ratio of mean male expression level to female expression level and used Wilcoxon
tests (adjusted for multiple testing using Benjamini and Hochberg's (84) algorithm) to
determine if differences in expression between the autosomes and the X chromosome were
significant.
496
497   ***Immunolabeling of parthenogenetic male meiotic cells***
498
In order to determine the dynamics of autosomal gene expression during meiosis in
parthenogenetic males, we followed the same protocol previously used to characterize
autosomal gene expression during meiosis in sexual males (44). In brief, we performed double
immunolabeling of SMC3_488 (#AB201542, AbCam) as a marker of cohesin axes (this allows
for the assessment of synapsis progression, and hence, cell cycle stage) and RNA polymerase
II phosphorylated at serine 2 (p-RNApol-II (ab193468, AbCam); an indicator of transcription).
Gonads from adult *T. monikense* and *T. douglasi* males were dissected in 1X PBS, then fixed
in paraformaldehyde (2%) and Triton X-100 (0.1%) solution for 15 minutes. Gonads were then
squashed on slides coated with poly-L-lysine, and flash frozen in liquid nitrogen. Slides were
incubated in PBS 1x for 20 minutes at room temperature, then blocked with a BSA 3% solution
(prepared by dissolving 3g of BSA in 100mL of PBS 1X) for 20 minutes. Slides were then
incubated overnight at 4°C in a humid chamber with the primary (p-RNA-Pol II), diluted in BSA
3% (1:100), and washed 3x for 5 minutes in PBS 1X. The secondary antibody, anti-rabbit
Alexa 594 (711-585-152, Jackson), diluted in BSA 3% (1:100), was applied, and samples were
incubated for 45 minutes at room temperature. Further washes were conducted as described
above, followed by an extended 10-minute wash in PBS 1X, and by blocking using a 5%
Normal Rabbit Serum (NRS) solution (X090210-8, Agilent Technology) for 30 minutes at room
temperature. Another 5-minute wash in PBS 1x was performed to remove excess blocking
solution. Samples were then incubated with the labelled antibody SMC3_488 (#AB201542,
AbCam) at a 1:100 dilution for 1 hour at room temperature, followed by a final series of washes
and staining with DAPI (D9542, Sigma-Aldrich) for 3 minutes at room temperature. Finally,
after a 5-minute wash in 1x PBS, two drops of Vectashield were added to the slide, which was
then covered with a coverslip, and sealed with nail varnish for imaging. Image acquisition was
performed at the Cellular Imaging Facility (CIF, University of Lausanne) using a Zeiss LSM
880 Airyscan.

Data

Genome assemblies have been deposited at DDBJ/ENA/GenBank under the accessions JBOIUC000000000 (*T. cristinae*) and JBOIUD000000000 (*T. podura*). The Illumina and PromethION sequencing data for *T. podura* are available in the European Nucleotide Archive under the following project PRJEB94285. Male genomic reads have been deposited at SRA under the following project PRJNA1293928. RNAseq reads have been deposited at SRA under the following projects: PRJNA1320679, PRJNA1327219. Code and annotations are available here: https://github.com/DarrenJParker/Timema_DC_MSCI_code and will be archived upon acceptance.

Acknowledgments

This work was supported by the European Research Council Consolidator Grant 864672 ("No Sex No Conflict", T.S.), Swiss FNS grant 31003A_182495 (T.S.), the Genoscope, the Commissariat à l'Énergie Atomique et aux Énergies Alternatives (CEA) and France Génomique (ANR-10-INBS-09–08). We thank the Lausanne Genomic Technologies Facility for PacBio library preparation and sequencing.

References

1. D. Bachtrog, *et al.*, Sex determination: why so many ways of doing it? *PLoS Biol.* **12**,
e1001899 (2014).

2. C. M. Disteche, Dosage compensation of the sex chromosomes. *Annu. Rev. Genet.* **46**,
537–560 (2012).

3. L. Gu, J. R. Walters, Evolution of sex chromosome dosage compensation in animals: a
beautiful theory, undermined by facts and bedeviled by details. *Genome Biol. Evol.* **9**,
2461–2476 (2017).

4. J. E. Mank, Sex chromosome dosage compensation: definitely not for everyone. *Trends*
Genet. **29**, 677–683 (2013).

5. N. Brockdorff, B. M. Turner, Dosage compensation in mammals. *Cold Spring Harb.*
Perspect. Biol. **7**, a019406 (2015).

6. R. Marin, *et al.*, Convergent origination of a Drosophila-like dosage compensation
mechanism in a reptile lineage. *Genome Res.* **27**, 1974–1987 (2017).

7. D. C. H. Metzger, B. A. Sandkam, I. Darolti, J. E. Mank, Rapid Evolution of Complete
Dosage Compensation in Poecilia. *Genome Biol. Evol.* **13** (2021).

8. I. Darolti, *et al.*, Extreme heterogeneity in sex chromosome differentiation and dosage
compensation in livebearers. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 19031–19036 (2019).

9. B. J. Meyer, Sex in the worm: counting and compensating X-chromosome dose. *Trends*
Genet. **16**, 247–253 (2000).

10. P. Chauhan, *et al.*, Genome assembly, sex-biased gene expression and dosage
compensation in the damselfly *Ischnura elegans*. *Genomics* **113**, 1828–1837 (2021).

11. D. J. Parker, K. S. Jaron, Z. Dumas, M. Robinson-Rechavi, T. Schwander, X
chromosomes show relaxed selection and complete somatic dosage compensation
across Timema stick insect species. *J. Evol. Biol.* (2022).
https://doi.org/10.1111/jeb.14075.

12. A. Pal, B. Vicoso, The X Chromosome of Hemipteran insects: conservation, dosage
compensation and sex-biased expression. *Genome Biol. Evol.* **7**, 3259–3268 (2015).

13. J. G. Rayner, T. J. Hitchcock, N. W. Bailey, Variable dosage compensation is
associated with female consequences of an X-linked, male-beneficial mutation. *Proc.*
Biol. Sci. **288**, 20210355 (2021).

14. E. G. Prince, D. Kirkland, J. P. Demuth, Hyperexpression of the X chromosome in both
sexes results in extensive female bias of X-linked genes in the flour beetle. *Genome*
Biol. Evol. **2**, 336–346 (2010).

15. B. Vicoso, D. Bachtrog, Numerous transitions of sex chromosomes in Diptera. *PLoS*
Biol. **13**, e1002078 (2015).

16. T. Conrad, A. Akhtar, Dosage compensation in *Drosophila melanogaster*: epigenetic
fine-tuning of chromosome-wide transcription. *Nat. Rev. Genet.* **13**, 123–134 (2012).

17. J. Djordjevic, *et al.*, Dynamics of X chromosome hyper-expression and inactivation in
male tissues during stick insect development. *PLoS Genet.* **21**, e1011615 (2025).

18. J. E. Mank, H. Ellegren, All dosage compensation is local: gene-by-gene regulation of
sex-biased expression on the chicken Z chromosome. *Heredity* **102**, 312–320 (2009).

19. P. S. Burgoyne, S. K. Mahadevaiah, J. M. A. Turner, The consequences of asynapsis
for mammalian meiosis. *Nat. Rev. Genet.* **10**, 207–216 (2009).

20. B. D. McKee, M. A. Handel, Sex chromosomes, recombination, and chromatin
conformation. *Chromosoma* **102**, 71–80 (1993).

21. A. I. Kalita, C. I. Keller Valsecchi, Dosage compensation in non-model insects -
progress and perspectives. *Trends Genet.* (2024).
https://doi.org/10.1016/j.tig.2024.08.010.

22. J. B. Pease, M. W. Hahn, Sex chromosomes evolved from independent ancestral
linkage groups in winged insects. *Mol. Biol. Evol.* **29**, 1645–1653 (2012).

23. F. N. Hamada, P. J. Park, P. R. Gordadze, M. I. Kuroda, Global regulation of X
chromosomal genes by the MSL complex in Drosophila melanogaster. *Genes Dev.* **19**,
2289–2294 (2005).

24. T. Straub, G. D. Gilfillan, V. K. Maier, P. B. Becker, The Drosophila MSL complex
activates the transcription of target genes. *Genes Dev.* **19**, 2284–2288 (2005).

25. A. I. Kalita, *et al.*, The sex-specific factor SOA controls dosage compensation in
Anopheles mosquitoes. *Nature* **623**, 175–182 (2023).

26. E. Krzywinska, P. Ribeca, L. Ferretti, A. Hammond, J. Krzywinski, A novel factor
modulating X chromosome dosage compensation in Anopheles. *Curr. Biol.* **33**, 4697–
4703.e4 (2023).

27. R. J. Davis, E. J. Belikoff, E. H. Scholl, F. Li, M. J. Scott, No blokes is essential for male
viability and X chromosome gene expression in the Australian sheep blowfly. *Curr. Biol.*
28, 1987–1992.e3 (2018).

28. M. A. Toups, B. Vicoso, Insect sex chromosome evolution: conservation, turnover, and
mechanisms of dosage compensation. *Curr. Opin. Insect Sci.* 101411 (2025).

29. N. Page, *et al.*, Single-cell profiling of Anopheles gambiae spermatogenesis defines the
onset of meiotic silencing and premeiotic overexpression of the X chromosome.
Commun. Biol. **6**, 850 (2023).

30. S. Schoenmakers, *et al.*, Female meiotic sex chromosome inactivation in chicken. *PLoS*
Genet. **5**, e1000466 (2009).

31. S. H. Namekawa, J. T. Lee, XY and ZW: is meiotic sex chromosome inactivation the
rule in evolution? *PLoS Genet.* **5**, e1000493 (2009).

32. R. L. Kelley, *et al.*, Expression of msl-2 causes assembly of dosage compensation
regulators on the X chromosomes and female lethality in Drosophila. *Cell* **81**, 867–877
(1995).

33. M. A. Toups, B. Vicoso, The X chromosome of insects likely predates the origin of class
Insecta. *Evolution* **77**, 2504–2511 (2023).

34. J. T. Lee, Disruption of imprinted X inactivation by parent-of-origin effects at Tsix. *Cell*
103, 17–27 (2000).

35. C. J. van der Kooi, T. Schwander, On the fate of sexual traits under asexuality. *Biol.*
Rev. Camb. Philos. Soc. **89**, 805–819 (2014).

36. J. M. Belote, J. C. Lucchesi, Control of X chromosome transcription by the maleless
gene in Drosophila. *Nature* **285**, 573–575 (1980).

37. E. Lifschytz, D. L. Lindsley, The role of X-chromosome inactivation during
spermatogenesis (Drosophila-allocycly-chromosome evolution-male sterility-dosage
compensation). *Proc. Natl. Acad. Sci. U. S. A.* **69**, 182–186 (1972).

38. Z. Bao, A. Belyi, E. Argyridou, J. Parsch, Disruption of meiotic sex chromosome
inactivation by X-autosome translocations in Drosophila melanogaster. *Genetics* **232**,
iyag011 (2026).

39. C. C. Avelino, *et al.*, Meiotic Sex Chromosome Inactivation: Conservation across the
Drosophila genus. *PLoS Genet.* **21**, e1011511 (2025).

40. T. Schwander, B. J. Crespi, Multiple direct transitions from sexual reproduction to
apomictic parthenogenesis in *Timema* stick insects. *Evolution* **63**, 84–103 (2009).

41. T. Schwander, L. Henry, B. J. Crespi, Molecular evidence for ancient asexuality in
Timema stick insects. *Curr. Biol.* **21**, 1129–1134 (2011).

42. J. Bast, *et al.*, Consequences of asexuality in natural populations: insights from stick
insects. *Mol. Biol. Evol.* **35**, 1668–1677 (2018).

43. K. S. Jaron, *et al.*, Convergent consequences of parthenogenesis on stick insect
genomes. *Science Advances* **8**, eabg3842 (2022).

44. Z. Dumas, W. Toubiana, M. Delattre, T. Schwander, Dynamics of recombination, X
inactivation and centromere proteins during stick insect spermatogenesis. *bioRxiv*
2024.09.05.611401 (2024).

45. T. Schwander, B. J. Crespi, R. Gries, G. Gries, Neutral and selection-driven decay of
sexual traits in asexual stick insects. *Proc. Biol. Sci.* **280**, 20130823 (2013).

46. S. Freitas, D. J. Parker, M. Labédan, Z. Dumas, T. Schwander, Evidence for cryptic sex
in parthenogenetic stick insects of the genus Timema. *Proceedings of the Royal Society*
B: Biological Sciences **290**, 20230404 (2023).

47. S. Simon, *et al.*, Old World and New World Phasmatodea: phylogenomics resolve the
evolutionary history of stick and leaf Insects. *Frontiers in Ecology and Evolution* **7**, 345
(2019).

48. S. Bank, S. Bradler, A second view on the evolution of flight in stick and leaf insects
(Phasmatodea). *BMC Ecol. Evol.* **22**, 62 (2022).

49. X. Li, J. E. Mank, L. Ban, The grasshopper genome reveals long-term gene content
conservation of the X Chromosome and temporal variation in X Chromosome evolution.
Genome Res. **34**, 997–1007 (2024).

50. Q.-L. Hu, *et al.*, Chromosome-level assembly, dosage compensation and sex-biased
gene expression in the small brown planthopper, Laodelphax striatellus. *Genome Biol.*
Evol. **14**, evac160 (2022).

51. A. M. Rice, Y. Li, P. Donnelly, A. McLysaght, Evolution of dosage-sensitive genes by
tissue-restricted expression changes. *Genome Biol. Evol.* **17**, evaf132 (2025).

52. J. Larsson, M. J. Svensson, P. Stenberg, M. Mäkitalo, Painting of fourth in genus
Drosophila suggests autosome-specific gene regulation. *Proc. Natl. Acad. Sci. U. S. A.*
101, 9728–9733 (2004).

53. J. E. Mank, D. J. Hosken, N. Wedell, Some inconvenient truths about sex chromosome
dosage compensation and the potential role of sexual conflict. *Evolution* **65**, 2133–2144
(2011).

54. E. Flintham, V. Savolainen, S. P. Otto, M. Reuter, C. Mullon, The maintenance of
genetic polymorphism underlying sexually antagonistic traits. *Evol. Lett.* **9**, 259–272
(2025).

55. D. C. Lahti, *et al.*, Relaxed selection in the wild. *Trends Ecol. Evol.* **24**, 487–496 (2009).

56. D. J. Futuyma, Evolutionary constraint and ecological consequences. *Evolution* **64**,
1865–1884 (2010).

57. H. Defendini, *et al.*, The release of sexual conflict after sex loss is associated with
evolutionary changes in gene expression. *Proc. Biol. Sci.* **292**, 20242631 (2025).

58. S. Koren, *et al.*, Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. *Genome Res.* **27**, 722–736 (2017).

59. M. Kolmogorov, J. Yuan, Y. Lin, P. A. Pevzner, Assembly of long, error-prone reads
using repeat graphs. *Nat. Biotechnol.* **37**, 540–546 (2019).

60. H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

61. R. Vaser, I. Sović, N. Nagarajan, M. Šikić, Fast and accurate de novo genome assembly
from long uncorrected reads. *Genome Res.* **27**, 737–746 (2017).

62. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. *arXiv* 1303.3997 (2013).

63. B. J. Walker, *et al.*, Pilon: an integrated tool for comprehensive microbial variant
detection and genome assembly improvement. *PLoS One* **9**, e112963 (2014).

64. D. R. Laetsch, M. L. Blaxter, BlobTools: Interrogation of genome assemblies.
F1000Res. **6**, 1287 (2017).

65. M. J. Roach, S. A. Schmidt, A. R. Borneman, Purge Haplotigs: allelic contig
reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 460
(2018).

66. N. C. Durand, *et al.*, Juicer provides a one-click system for analyzing loop-resolution Hi-
C experiments. *Cell Syst.* **3**, 95–98 (2016).

67. O. Dudchenko, *et al.*, De novo assembly of the Aedes aegypti genome using Hi-C yields
chromosome-length scaffolds. *Science* **356**, 92–95 (2017).

68. M. Seppey, M. Manni, E. M. Zdobnov, BUSCO: Assessing genome assembly and
annotation completeness. *Methods Mol. Biol.* **1962**, 227–245 (2019).

69. T. Brůna, K. J. Hoff, A. Lomsadze, M. Stanke, M. Borodovsky, BRAKER2: automatic
eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a
protein database. *NAR Genom Bioinform* **3**, lqaa108 (2021).

70. J. M. Flynn, *et al.*, RepeatModeler2 for automated genomic discovery of transposable
element families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457 (2020).

71. A. Smith, R. Hubley, P. Green, RepeatMasker Open-4.0. *RepeatMasker Open-4.0*
(2013).

72. E. V. Kriventseva, *et al.*, OrthoDB v10: sampling the diversity of animal, plant, fungal,
protist, bacterial and viral genomes for evolutionary and functional annotations of
orthologs. *Nucleic Acids Res.* **47**, D807–D811 (2019).

73. D. J. Parker, J. Djordjevic, T. Schwander, Olfactory proteins in *Timema* stick insects.
Frontiers in Ecology and Evolution **7**, 101 (2019).

74. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

75. A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

76. M. Stanke, *et al.*, AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic
Acids Res.* **34**, W435–9 (2006).

77. T. Brůna, A. Lomsadze, M. Borodovsky, GeneMark-EP+: eukaryotic gene prediction
with self-training in the space of genes and proteins. *NAR Genom Bioinform* **2**, lqaa026
(2020).

78. E. P. Nawrocki, S. R. Eddy, Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics **29**, 2933–2935 (2013).

79. D. M. Emms, S. Kelly, OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).

80. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**, 841–842 (2010).

81. C. Larose, G. Lavanchy, S. Freitas, D. J. Parker, T. Schwander, Facultative
parthenogenesis in an asexual stick insect: re-evolution of sex or vestigial sexual
capacity? *bioRxiv* 2022.03.25.485836 (2022).

82. S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).

83. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–
140 (2010).

84. Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: A practical and powerful
approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289–300
(1995).