Aging-enhanced accumulation of fibroblasts excludes oligodendrocytes in demyelinated lesions

Brian M. Lozinski[1,], Charlotte D'Mello[1], Yifei Dong[1,2], George S. Melchor[3], Samira Ghorbani[1,4],

Cenxiao Li[1], Marlene Morch[1], Dorsa Moezzi[1], Rajiv Jain[1], Parisa Etemadi[1], Chang-Chun Ling[5],

Jeffrey K. Huang[3], Carlos Camara-Lemarroy[1], V. Wee Yong[1]*

[1]Department of Clinical Neurosciences and Hotchkiss Brain Institute, University of Calgary, Canada

[2]Department of Biochemistry, Microbiology and Immunology, University of Saskatchewan, Canada

[3]Department of Biology, Georgetown University, Washington DC, USA

[4]Department of Biology, University of Toronto Mississauga, Canada

[5]Department of Chemistry, University of Calgary, Canada

*Correspondence:

V. Wee Yong, PhD

Professor

University of Calgary

Calgary, Alberta T2N 4N1, Canada

vyong@ucalgary.ca

403-220-3544

<u>Abstract</u>

Fibroblast dysregulation contributes to aberrant repair and pathological fibrosis. Emerging

evidence suggest that fibroblasts accumulate in lesions following central nervous system injury,

but whether and how they influence oligodendrocytes and myeloid cell responses, especially in

aging, is uncertain. Here we report that fibroblasts infiltrate the parenchyma of spinal cord white

matter (SCWM) lesions after lysolecithin-induced demyelination. Transcriptomic analysis

reveals diverse activation states of fibroblasts in lesions and communication networks between

fibroblasts, microglia/macrophages and oligodendrocyte precursor cells (OPCs). The invasion of

fibroblasts is facilitated by microglia/macrophages, and areas of fibroblast accumulation are

devoid of OPCs. Fibroblast density is exacerbated with aging leading to greater fibrosis in

lesions. Finally, we found fibroblasts in multiple sclerosis lesions with predicted communication

networks between microglia/macrophages and OPCs. These results demonstrate a role of

fibroblasts in demyelination-associated neuropathology, which is exacerbated by aging, and

highlight the importance of regulating fibroblasts to promote effective CNS repair.

<u>Introduction</u>

Fibroblasts are a heterogeneous population of cells with important roles in extracellular matrix

(ECM) synthesis, maintenance of cellular niches, tissue regeneration and wound healing[1]. To

carry out their roles in homeostasis and injury fibroblasts form numerous interactions with other

cell types including immune cells such as macrophages and T cells. Fibroblasts associated with

the central nervous system (CNS) primarily reside in barriers including the meninges,

perivascular space, and choroid plexus where they contribute to tissue homeostasis[2] and

immune regulation[3,4]. Recent studies suggest that in response to injury, such as spinal cord

trauma and experimental autoimmune encephalomyelitis (EAE), fibroblasts accumulate in the

CNS parenchyma and contribute to fibrotic scaring[5–9]. This results from the ability of fibroblasts

to respond to inflammatory and injury mediated signals promoting their activation. Activated

fibroblasts differentiate into an ECM-producing smooth muscle actin (SMA)-expressing

myofibroblast phenotype which can contribute either to tissue regeneration or fibrosis[10]. Thus,

while fibroblast activation is necessary to maintain tissue integrity, it may also impair

regenerative processes[5,11,12].

Remyelination is the most common and robust form of CNS regeneration[13].

Remyelination occurs when demyelination, such as that seen in multiple sclerosis (MS),

stimulates the migration, proliferation, and differentiation of oligodendrocyte progenitor cells

(OPCs) into mature myelinating oligodendrocytes[14]. Restoring myelin to denuded axons

reconstitutes saltatory conduction, axonal trophic support, and produces a mechanical barrier to

injury leading to axonal preservation and functional improvement[13,15,16]. However, remyelination

often fails due to extrinsic variables in the environment, such as inhibitory ECM, and intrinsic

factors including age-related cellular dysfunctions[16,17].

Age is a major contributor to disease worsening across a wide range of conditions,

including MS and fibrotic disorders[18–20]. For example, fibroblasts in the skin and skeletal

muscles exhibit age-related dysregulation, characterized by altered lineage markers, increased

expression of inflammation-associated genes, and elevated ECM production[21–23]. In MS, OPCs

also show signs of age-related impairment, including upregulation of senescence markers,

epigenetic dysfunction, reduced responsiveness to growth factors, and impaired differentiation

and remyelination capacity[24–26]. While age-related remyelination decline is well recognized, the

impact of aging on fibroblast responses to CNS injury remains poorly understood.

In this study, we characterized the response of fibroblasts in lesions following focal

lysolecithin (LPC)-mediated demyelination of the spinal cord white matter (SCWM), and further

evaluated the influence of aging on the fibrotic response. With spatial transcriptomics of LPC

tissue and cells in culture, we interrogated interactions between fibroblasts,

microglia/macrophages and OPCs. Finally, we utilized a publicly available transcriptomic

dataset to show that fibroblasts are present in MS lesions, and describe communication patterns

between fibroblasts, microglia/macrophages and OPCs in MS. Our findings highlight the

necessity of considering invasion of fibroblasts into the CNS parenchyma in lesion evolution and

recovery.

Results

Fibroblasts are localized within demyelinated lesions

Although fibroblasts are restricted to CNS barriers in homeostasis, they are reported to be

elevated in the parenchyma in models of traumatic spinal cord injury, stroke and autoimmune

neuroinflammation[5,12,27]. While there is a growing appreciation for the functional connectivity of

fibroblasts with other CNS resident cells, their roles in the regenerative processes of the CNS,

primarily remyelination, remain poorly understood. The LPC model of demyelination provides an

optimal system for studying neuroinflammation and remyelination due to the precise spatial

location and temporal evolution of lesions[28,29]. LPC was stereotaxically injected into the ventral

spinal cord white matter (SCWM) of mice resulting in a focal lesion characterized by damaged

myelin (disrupted immunoreactivity for myelin basic protein, MBP), swollen axons as determined

by neurofilament heavy chain staining (NFH), and GFAP-positive reactive astrocytes (Fig 1a). In

PDGFRβTdTomato transgenic mice commonly used to study fibroblast responses in peripheral

organs[30–32],we found TdTomato expression amongst the disrupted MBP-positive myelin debris

and in close proximity to Iba1-positive microglia/macrophages (Fig 1b). As well, TdTomato and

PDGFRβ immunofluorescence overlapped in LPC lesions giving confidence to the specificity of

expression (Fig 1c). Further, we found that PDGFRβ was correspondent with a number of other

markers of fibroblasts including collagen type 1 alpha 1 (Col1a1) and SMA for activated

fibroblasts (Fig 1d).

We examined the fibroblast response at 7 and 21 days post injury (dpi) as periods when

tissue regenerative responses begin to be robust and when they are concluding,

respectively[17,28] (Fig 1e-j). While the total lesion volume was reduced from day 7 to 21 (Fig 1e,

h), the total TdTomato-positive volume was comparable between 7 to 21 dpi (Fig 1f, i).

Consequently, the proportion of the lesion occupied by fibroblasts was relatively greater at 21

compared to 7 dpi (Fig 1g, j). We then asked if fibroblasts differentiate into contractile ECM

producing myofibroblasts using markers of ECM expression (fibronectin and periostin) and cell

contractility (SMA) (Fig 1k). Approximately 40% of TdTomato-positive fibroblasts overlapped

with these markers at 7 dpi (Fig 1l-n). In comparison, at day 21 TdTomato-labelled fibroblasts

showed increased co-expression of SMA and periostin but decreased co-expression of

fibronectin (Fig 1l-n).

We validated the fibroblast response in LPC lesions using single-nuclear RNA

sequencing (snRNAseq) dataset at 5, 10, and 20 dpi (Fig 2a-c). Vascular and mesenchymal

cells in LPC lesions were clustered and annotated into five groups: fibroblasts (*Col3a1, Col1a2,*

Col1a1, Bnc2, Dcn, and Foxp2), endothelial cells (*Flt1, Pecam1, Ly6c1, Adgrl4),* pericytes

(*Abcc9, Cspg4, Lin7a),* smooth muscle cell *(SMC) (Acta2, Myh11, Pdlim3, Lmod1),* and

ependymal cells (*Adamts20, Cfap54, Dnah12, Agbl4, Cfap299)* (Fig 2a, b). *Pdgfrb* was

expressed in fibroblasts, pericytes, and SMCs at variable levels (Fig 2a). We found that

fibroblasts were significantly expanded in LPC lesions at all timepoints compared to naïve and

non-lesioned samples (Fig 2b, c). Fibroblasts constituted ~75% of the vascular and

mesenchymal cells 5 and 10 dpi, and approximately 60% at 20 dpi (Fig 2d). Further evaluation

of the vascular and mesenchymal cell populations can be found elsewhere[33].

Collectively, our analysis of the fibroblast response to LPC injury using

immunohistochemistry of the PDGFRβ-TdTomato reporter mice and snRNAseq of wildtype

demyelinated mice confirms that activated fibroblasts expand rapidly and persist into late stages

of LPC injury.

Spatial transcriptomics of fibroblasts in LPC lesions

To further investigate fibroblasts in the LPC lesion environment we used the 1000-plex Mouse

Neuroscience Panel with the in-situ single cell CosMx spatial platform from Nanostring on three

sham and two LPC (14 dpi) lesioned spinal cords with 1 field of view (FOV) per section that

demarcates the lesion area (Fig 3a, b). Day 14 LPC lesions were used as this is an intermediate

period with ongoing remyelination[28,34]. We identified a total of 2954 segmented cells with (1828

cells from LPC and 1126 cells from sham controls). Unsupervised clustering of all cells revealed

12 clusters which were then mapped to the Azimuth mouse motor cortex reference map (Fig.

3c, d). This delineated the following cell types: astrocytes (*Gja1, Slc4a4, Slc6a1*),

oligodendrocytes (*Mag, Mog, Myrf*), OPCs (*Pdgfra, Cspg5, Vcan*), microglia/macrophages

(*Csf1r, Hexb, Csf3r*), vascular leptomeningeal cells (VLMCs) (*Dcn, Vtn, Igfbp7*), endothelial cells

(*Cldn5, Pecam1, Flt1*) and pericytes (*Rgs5, Vtn, Myl9*) (Fig 3d, e). A separate identity was

created for fibroblasts as cells which expressed *Col1a2* and *Pdgfrb* (Fig 3d, e). The UMAP in

Fig. 3c shows an overlay of the clusters as coloured by cells from Sham and LPC mice. We

identified a total of 84 *Col1a2* and *Pdgfrb* double-positive fibroblasts, 5 in sham controls and 79

in LPC lesions (Fig 3f, g). The top genes for fibroblasts aside from *Pdgfrb* and *Col1a2* were

Col3a1, *Col1a1, Vtn* and *Dcn* (Fig 3e). The spatial analysis of LPC lesions affirmed that

fibroblasts were 16 times more abundant in the parenchyma of LPC lesions at 14 dpi compared

to sham (Fig 3g), and that fibroblasts were in close proximity to macrophages and OPCs (Fig

3b). Collectively, the spatial transcriptomic dataset shows an increased presence of fibroblasts

in the injured CNS parenchyma during ongoing tissue regeneration and remyelination.

Fibroblast interactions in LPC lesions

We analyzed the spatial dataset for ligand-receptor (L-R) interactions and thus potential

functions of fibroblasts within LPC lesions. Using the Cellchat package[35] we examined L-R

interactions between fibroblasts, microglia/macrophages, and OPCs (Supplementary Fig 1a-d).

We chose to investigate fibroblast interactions with OPCs as the latter are critical for

remyelination, and with microglia/macrophages as they are the dominant immune cells in LPC

lesions. Each interaction is assigned a probability value followed by a permutation test which

then allows us to make inferences about biologically significant cell-cell communication. Circle

plots and heatmaps delineated the aggregated cell-cell communication network between

fibroblasts, OPCs and macrophages in terms of number of interactions and total interaction

strength (Supplementary Fig 1a-d). Fibroblasts interacted most with OPCs, followed by

fibroblasts with themselves (Supplementary Fig 1a, c). In terms of total strength of interactions

this was determined to be highest between fibroblasts and microglia/macrophages

(Supplementary Fig 1b, d). Fibroblast derived ligands affected receptors involved in signaling

crucial for phagocytosis (*Pros1-Axl, Pros1-Mer*tk), macrophage polarization (*Spp1-Cd44, Apoe-*

Trem2/Tyrobp), cell adhesion (*Col1a1-Itgav/Itgb8*) and OPC survival (*Ptn-Ptprz1, Bdnf-Ntrk2*)

(Supplementary Fig 1f; Supplementary Fig 2a). Conversely, fibroblasts were predicted to

respond to L-R interactions that affect their survival, proliferation, differentiation, and migration

such as *Pdgfd-Pdgfrb*, *Fgf2-Fgfr2* and *Ptprs-Ntrk3* (Supplementary Fig 1g; Supplementary Fig

2b).

Predicted L-R interactions were also categorized into functionally related signaling

pathways. Fibroblasts were highlighted as the major driver of outgoing signaling in LPC lesions

among these three cell groups (Fig 4a; Supplementary Fig 1e). The signaling pathways that

contributed the most to outgoing fibroblast signaling included apolipoprotein E (APOE),

COLLAGEN, amyloid precursor protein (APP), prosaposin (PSAP) and FN1 (Fig 4a;

Supplementary Fig 3a). The outgoing microglia/macrophage signals were mainly contributed by

APOE, osteopontin (SPP1), colony stimulating factor (CSF) and GALANIN pathways. The

outgoing signaling from OPCs were mainly from pleiotrophin (PTN), Glutamate, neural cell

adhesion molecule (NCAM), and PDGF pathways (Fig 4a; Supplementary Fig 3a).

Microglia/macrophages were the primary recipient of L-R signaling through pathways such as

APOE, SPP1, and protein S (PROS) (Fig 4b; Supplementary Fig 3b). As well, OPCs responded

to COLLAGEN, PTN, and FN1 (Fig 4b; Supplementary Fig 3b). Although fibroblasts were not

identified as a dominant receiver of signaling through the cell types assessed they were inferred

to receive signaling predominantly from COLLAGEN, GTPase-activating protein (GAP) and

PDGF signaling pathways (Fig 4b; Supplementary Fig 1e).

To further contextualize the communication between fibroblasts, microglia/macrophages

and OPCs, we utilized Cellchat's pattern recognition feature. This allows for the identification of

global communication patterns to explore how the different cell groups and signaling pathways

coordinate to function. This analysis uncovered three outgoing and three incoming signaling

patterns that were dominated by individual cell groups (Fig 4c-f). The outgoing signaling from

fibroblasts was characterized by pattern 1 which represented multiple pathways including

COLLAGEN, vitronectin (VTN), and SPP1, while pattern 2 (which included signaling pathways

such as endothelin (EDN), PTN, and VEGF) was driven by OPCs, and pattern 3 (which included

signaling pathways such as leptin (LEP), GALININ, and CSF) by microglia/macrophages (Fig

4c, d). On the other hand, the communication pattern of target cells showed that incoming

fibroblast signaling was dominated by pattern 3 which includes PDGF, nerve growth factor

(NGF), and LEP pathways, while pattern 1 drove signaling (which included signaling pathways

such as PTN, FN1, and EDN) in OPCs, and pattern 2 (which included signaling pathways such

as APOE, APP, and CSF) drove signaling in microglia/macrophages (Fig 4e, f). Altogether this

highlights the involvement of fibroblasts in the LPC environment as a major source of input for

microglia/macrophages and OPCs.

Fibroblasts respond to microglia/macrophages

Since the spatial data analyses suggest that fibroblasts engaged in reciprocal communication

with microglia/macrophages in the 14 dpi LPC lesion, we hypothesized that one cell type might

help with the recruitment of the other into the area of injury. Indeed, Iba1+

microglia/macrophages and PDGFRβ+ fibroblasts were closely associated in day 7 LPC lesions

(Fig 5). We thus examined 3 dpi and found that microglia/macrophages began accumulating in

the LPC lesion at day 3 before peaking at 7 days (Fig 5a-d). Similarly, the fibroblast response

peaked at day 7 (Fig 5b, d). However, there was no observable PDGFRβ immunoreactivity at

day 3 indicating that fibroblasts do not become elevated until after microglia/macrophage

accumulation had begun (Fig 5a, d).

To corroborate the possibility that microglia/macrophages attract fibroblasts into LPC

lesions, meningeal fibroblasts were added to the upper compartment of a Boyden chamber with

bone marrow derived macrophages (BMDMs) or medium alone in the lower chamber (Fig 5e).

After 24 hours there was a 3-fold increase in the number of fibroblasts that migrated across the

membrane when cultured with BMDMs compared to the basal rate of migration (Fig 5f, g).

These results suggest a role for macrophages in the promotion of fibroblast elevation in CNS

lesions.

Fibroblasts spatially exclude OPCs

The CosMx Cellchat analyses suggest the interaction of fibroblasts with OPCs which we

examined further. We found that at 14 dpi Olig2+ oligodendrocytes and Olig2+PDGFRα+ OPCs

seemed to be restricted from regions of the LPC lesion that were occupied by PDGFRβ+

fibroblasts (Fig 6a). To determine whether this was previously surviving cells or newly

responding OPCs, we utilized NG2CreER:MAPTmGFP mice that label newly differentiated

oligodendrocytes with a membrane bound green fluorescent protein (GFP)[15,16]. Using this

method, we found that areas within 14 dpi LPC lesions that were occupied by fibroblasts, based

on PDGFRβ immunoreactivity, contained little GFP (Fig 6b, c). The difference in GFP was not

due to axonal preservation as there was no difference in the density of NFH-positive axons in

the fibroblast occupied region compared to the rest of the LPC lesion (Supplementary Fig. 4a,

b). These results suggest that fibroblasts have the potential to regionally restrict OPC functions

in LPC lesions.

To test this further we established cultures of meningeal fibroblasts in 96-well plates,

upon which OPCs were then seeded (Fig 6d). No difference in total O4+ OPCs was observed

when comparing OPC alone with those in co-cultured wells (Fig 6e). However, there was a

significant decrease in the number of O4+MBP+ mature oligodendrocytes in co-cultured wells

(Fig 6f). As well, oligodendrocytes in co-cultured wells displayed less complex morphological

phenotypes than control cells grown without fibroblasts (Fig 6g). In controls, mature

oligodendrocytes were able to spread in approximately even distribution around the cell soma

(Fig 6d). In comparison, in co-culture, mature oligodendrocytes occupied only areas without

fibroblasts and had more irregular orientations (Fig 6d). The restriction of mean outgrowth of

oligodendrocytes by fibroblasts occurred regardless of the density of fibroblasts (Supplementary

Fig 5a, b). As well, the inhibitory effect of fibroblasts was consistent regardless of the order that

cells were seeded and was not recapitulated by culturing OPCs with fibroblast conditioned

media (Supplementary Fig 5c, d). Collectively, this data highlights the inhibitory effect of

fibroblasts on OPC differentiation.

Aging enhances fibroblast properties in LPC lesions

Age is a critical factor for many biological processes including tissue regeneration[36]. Aging is

known to affect OPC and microglia/macrophage dynamics and is a critical factor in

remyelination potential[17,18,37,38]. It is not known how the fibroblast response to CNS injury is

affected by age and whether this impacts downstream processes. To test how age affects

fibroblasts in LPC lesions, young (2-3 month) and middle-aged (12 month) mice were injected

with LPC in the ventral SCWM. Tissue was collected 7, 14 and 21 dpi and immunostained for

MBP and PDGFRβ (Fig 7a). Age did not affect lesion volume at 7 dpi but LPC lesions in middle-

aged mice were significantly larger at 14 and 21 dpi (Fig 7b, d). Age also did not affect the

PDGFRβ+ volume at 7 and 14 dpi, but did cause a significant increase at 21dpi in middle-aged

lesions (Fig 7c, e). When expressed in proportion to the lesion volume, the middle-aged LPC

lesion has more fibroblasts than young LPC lesions at 7 and 21 dpi (Fig 7f).

We investigated the effect of age on the expression of phenotypic markers in fibroblasts

in LPC lesions. Fibroblasts in day 7 LPC lesions in middle-aged demyelinated mice displayed

elevated levels of activated myofibroblast marker SMA than those in their young counterparts

(Fig 8a). While we did not find quantitative differences in Col1a1 or laminin at 7 dpi, these were

elevated at 21 dpi (Fig 8b, c). Finally, we found significantly fewer Olig2+ oligodendrocyte

lineage cells within the fibroblast-occupied regions of middle-aged compared to young mice (Fig

8d). Overall, we found elevated ECM levels in middle-aged mice that corresponded with a

reduction in oligodendrocyte lineage cells in lysolecithin injury.

Fibroblasts are present in MS lesions

We extended from murine data into the human demyelinating condition, MS. Frozen brain

sections from autopsied MS cases were stained for PDGFRβ as well as CD45 to identify

leukocytes, and DAPI for cell nuclei (Fig 9a). We found that PDGFRβ+ cells were found in

perivascular areas in close proximity to CD45+ leukocytes (Fig 9a).

Next, we reanalyzed previously published snRNAseq datasets from human brain tissues

of MS lesion rim, core and periplaque; and white matter tissue from neurologically healthy

controls[39]. Unsupervised clustering identified 25 clusters from the 66,432 total cells including

leukocytes, Bergmann glia, dendritic cells, vascular endothelial cells, OPCs, oligodendrocytes,

astrocytes, ependymal cells, microglia/macrophages, and neurons (Fig 9b). We highlighted a

cluster of 174 cells that expressed fibroblast related genes[40] such as *PDGFRB*, *COL1A2*,

COL15A1, *CEMIP*, *DCN*, and *FBLN1* (Fig 9c). From this cluster we focused on *PDGFRB*+ cells

as our population of interest that were positive for the markers *PDGFRA*, *COL1A2*, *TGFBR3*,

FBLN1, *DCN*, *COL15A1*, and *LAMC3* (Fig 9d; Supplementary Fig 6). Importantly, genes

associated with pericytes (*RGS5*), and smooth muscle cells (*TAGLN*) were not appreciably

expressed in our population of interest (Supplementary Fig 7a)[41]. The *PDGFRB*+ fibroblasts

were separated from the other cells in the dataset and re-clustered. This delineated 2

subpopulations (Fig 9e). Subcluster 1 expressed genes associated with stress response

(*HSPB1*, *HMGB1*) while subcluster 0 expressed genes associated with ECM, cell proliferation,

and migration (*LAMA2*, *NTRK3*) (Fig 9e; Supplementary Fig 7b, c). Assessment of these

clusters via region showed that subcluster 0 was present in chronic active and chronic inactive

edges and also the lesion core, but only minimally observed in control white matter or

periplaque area; subcluster 1 was elevated only in the lesion core (Fig 9f).

Fibroblast interactions in MS lesions

We interrogated the interactions of fibroblasts with other cell types in MS lesions using CellChat

L-R interaction analysis. Similar to LPC lesions, fibroblasts, microglia/macrophages and OPCs

all communicated with each other (Supplementary Fig 8a-d). Fibroblast outgoing signals

primarily involved ECM-receptor interactions while most incoming signals were via secreted

signaling (Supplementary Fig 8e; Supplementary Fig 9 a, b). This is seen in the L-R pairings

engaged by fibroblast derived ligands such as *COL1A2-ITGAV/ITGB9* and *LAMA2-CD44*

(Supplementary Fig 8f; Supplementary Fig 9 a, b). As well, incoming secreted L-R interactions

that fibroblasts were predicted to respond to included *FGF1-FGFR4* and *PDGFB-PDGFRA*

(Supplementary Fig 8g; Supplementary Fig 9 a, b).

Fibroblast L-R pairs were further categorized into 76 signaling pathways such as

transforming growth factor beta (TGFB), SPP1, PDGF, CSF, FN1, and COLLAGEN

(Supplementary Fig 10a). Fibroblasts contributed to pathways that drove microglia/macrophage

signaling including ephrin receptor A (EPHA), GAP, COLLAGEN, LAMININ, CSF, and CXC

motif chemokine ligand (CXCL) (Figure 10a; Supplementary Fig 9c; Supplementary Fig 10a).

Additionally, fibroblasts participated in signaling pathways that signaled at OPCs including bone

morphogenic protein (BMP), WNT, FGF, COLLAGEN, FN1, and LAMININ (Figure 10a;

Supplementary Fig 9c; Supplementary Fig 10a).

Communication patterns between cell populations highlighted three outgoing and three

incoming patterns (Fig 10a, b). Outgoing pattern one was primarily driven by OPCs. This pattern

was represented by pathways such as EPHA, VCAM, and NOTCH (Fig 10a). Pattern two was

driven mainly by microglia/macrophages defined by signaling pathways such as SPP1, PDGF,

and ANNEXIN (Fig 10a). The third outgoing pattern was defined primarily by fibroblast signaling.

This pattern was driven by pathways such as GAP, KIT, and plasminogen activator urokinase

(PLAU) pathways (Fig 10a). In the reverse, fibroblasts responded to incoming pattern three

represented by VCAM, neurotrophin (NT) and adhesion G protein-coupled receptor A2

(ADGRA) pathways (Fig 10b). OPCs responded to incoming pattern one that was driven by

signaling pathways including LAMININ, EDN, and WNT (Fig 10b). Microglia/macrophages

responded to pattern two that was defined by pathways including CSF, PLAU and TGFB (Fig

10b).

Comparison of the signaling pathways in LPC and MS lesions revealed 16.3% overlap of

outgoing fibroblast pathways and 15.3% of shared incoming pathways (Fig 10c, d). Shared

outgoing pathways included PDGF, PTN and COL1A1, and shared incoming pathways included

FGF, COL1A1 and VEGF (Fig 10c, d). Some unique fibroblast signaling pathways in MS

included outgoing PLAU and TGFB signaling and incoming WNT and TGFB signaling (Fig 10c,

d). Within these pathways we identified that there were 16 conserved outgoing fibroblast L-R

interactions and 10 conserved incoming fibroblast L-R interactions; these include fibroblast

derived FGF1 and PTN signaling to FGFR2 and PTPRZ1 respectively and, fibroblast CD44 and

NTRK3 responding to COL1A1 and PTPRS respectively (Supplementary Fig 10c, d). There

were similarities in the L-R interactions of fibroblasts in MS and LPC lesions such as outgoing L-

R interactions (*PROS1-MERTK*, *PTN-PTPRZ1*, *FGF1-FGFR2*, and *COL1A1-CD44*), and

incoming interactions (*FGF2-FGFR2*, *PTPRS-NTRK3*, *COL1A1-ITGAV/ITGB8*, and *COL4A5-*

CD44) (Fig 10c, d; Supplementary Fig 2; Supplementary Fig 3; Supplementary Fig 9a, b;

Supplementary Fig 10c, d).

Overall, it appears that microglia/macrophages and OPCs employ multiple signaling

pathways compared to fibroblasts. Furthermore, cross-referencing the outgoing and incoming

patterns highlighted that some pathways such as VCAM are shared by all three cell groups

assessed with microglia/macrophages and OPCs both contributing to outgoing VCAM signaling

and fibroblasts as sole responder (Supplementary Fig 10a, b). Taken together these data

highlight not only the presence of fibroblasts in MS lesions but also their prominent interactions

with microglia/macrophages and OPCs in their environment.

<u>Discussion</u>

The accumulation of fibroblasts in the parenchyma of CNS lesions has received scant attention

due to the lack of a prominent marker that clearly distinguishes this cell population, and because

antigens that label them (e.g. PDGFRβ, NG2 or GLAST) are also present on other cell types

such as pericytes and smooth muscle cells[4]. Nonetheless, combining markers such as PDGFRβ

and Col1a1 has elevated confidence in detecting fibroblasts and an increasing literature now

implicate them in models of traumatic CNS injury and MS[5,7–9,12]. Previously, PDGFRβ positive

cells were described in the core of the scar resulting from traumatic spinal cord injury[12,42]

although their cellular identity (e.g. fibroblasts or pericytes) could not be definitively discerned.

Additional recent reports describe perivascular fibroblasts with angiogenic and fibrogenic

functions[43]. In the current study, the presence of fibroblasts using a PDGFRβ-TdTomato

reporter[30,31,44] and additional staining markers (Col1a1, SMA, fibronectin, periostin) affirm the

presence of fibroblasts in the parenchyma of CNS lesions of LPC injury and MS. Highly

multiplexed RNA in situ hybridization (RNA-ISH, CosMx), which revealed additional fibroblast

transcriptome (*Col1a2, Col3a1, Vtn, Dcn, Cfh* and *Fbn1*), showed that there was a 16-fold

increase in fibroblasts in the LPC lesion compared to sham controls. Moreover, the absence of

Rgs5, a commonly used pericyte marker, lend further support to the identification of fibroblasts.

In the LPC model, fibroblasts are present in day 7 lesions, a period when the repopulation of

OPCs is at its peak and remyelination is beginning[28,34]. The period between day 7 and day 21 is

when OPCs differentiate and remyelination proceeds progressively albeit incomplete[17,34]. We

find that fibroblasts persist in the LPC lesion from 7 to 21 days with no resolution.

The accumulation of fibroblasts in LPC injury appears to be facilitated by

microglia/macrophages as suggested by the temporal relationship where fibroblasts do not

become elevated until after onset of neuroinflammation, and the in vitro results that show

macrophages to be chemotactic for fibroblasts. Transcriptomic data with Cellchat analyses

suggest microglia/macrophages and fibroblasts communicate through numerous signaling

pathways/molecules including PDGF which is a known chemotactic signal for fibroblasts[45].

Indeed, macrophage-derived PDGF has been proposed as one half of a reciprocal cell-circuit

with fibroblast derived M-CSF that is critical for regulating tissue fibroblast responses[46,47].

Further studies will be required to determine the predominant microglia/macrophage-derived

signals that promote accumulation of fibroblasts in LPC lesions.

Fibroblasts have been reported to be inhibitors of axon regrowth[11]. However, little is

known about the potential effect of fibroblast accumulation on the most common form of CNS

regeneration, remyelination. Analysis of L-R interactions predicted from our CosMx dataset

suggests that fibroblasts are a potent driver of signaling in LPC demyelination. Importantly for

remyelination, we found that the many of the fibroblast-derived ligands that can affect OPCs are

ECM genes including *Fn1* and *Col1a1*[43]. Accumulation of inhibitory ECM components impair

OPC function and reduce remyelination[16,48]. Aside from its role as a structural protein,

fibronectin can form aggregates that impair OPC maturation[49]. The effect of collagens on OPC

functions are less understood, although impaired migration may occur[5]. ECM components can

also impede OPC function through indirect effects including by stimulating

microglia/macrophages and lymphocytes that kill OPCs[49]. Furthermore, how potential fibroblast-

led signaling interactions continue to influence remyelination requires further investigation, as

fibroblasts are transcriptionally dynamic throughout remyelination[33].

It is also possible that fibroblasts provide a physical obstacle to oligodendrocyte

processes that are contacting axons for remyelination. This is supported by our in vivo results

that regions of an LPC lesion occupied by fibroblasts have few OPCs or newly formed myelin

compared to the unoccupied areas of LPC lesions. Furthermore, in vitro co-culture of fibroblasts

with OPCs show a density dependent curve of fibroblast inhibition. Additional experiments

focusing on the depletion or reduction of fibroblasts within LPC lesions will provide clarity to the

mechanisms regulating the inhibition of oligodendrocyte lineage cells by fibroblasts.

We did attempt to deplete fibroblasts using PDGFRβCre:Rosa26iDTR mice but these

cells remained persistent in lesions (data not shown). In a report in EAE, reducing fibroblast

proliferation by fibroblast-specific expression of the herpes simplex virus thymidine kinase gene

produced a modest improvement in clinical disability and an elevation of oligodendrocyte

lineage cells[5].

Age is a major contributor to overall morbidity and the reduced efficiency of many

biological processes[36] including the formation of oligodendrocytes[50] and remyelination[17,51]. We

found a greater number of fibroblasts in LPC lesions of middle-aged compared to young mice.

While there is a reduced rate of accumulation of microglia/macrophages within lesions of aging

compared to young mice, these cells in aging animals in 'inflammaging' are more capable of

producing inflammatory cytokines, chemokines and other molecules[52,53]. Indeed, a greater

number of mononuclear phagocytes in aging CNS lesions are positive for *Spp1* which encodes

for the glycoprotein osteopontin[18]. *Spp1* positive macrophages have been shown to drive the

activation of myofibroblasts leading to increased tissue fibrosis[54]. We found in both our CosMx

dataset and in the reanalysis of MS snRNAseq data that *Spp1*-positive microglia/macrophages

are present in both LPC and MS lesions. Thus, the increased presence of fibroblasts in middle-

aged LPC lesions may contribute to worse remyelination outcomes noted with aging as we find

that there are fewer oligodendrocyte lineage cells in fibroblast occupied regions. As well,

fibroblasts in lesions of middle-aged relative to young mice express greater amounts of ECM

components which may contribute to already existing OPC dysregulation.

To extend from animal models to MS, we stained and identified PDGFRβ positive cells in

close proximity to CD45 positive leukocytes. We next analysed a single-nucleus RNA

sequencing dataset of MS[39] and found a small population of presumptive fibroblasts; it is

possible that the cell isolation process was not optimized for fibroblasts and thus did not

adequately capture the entirety of the number of these cells in lesions. Nonetheless, we

identified a population of *PDGFRβ* positive cells that expressed genes associated with

fibroblasts, and similar to those found in LPC lesions, such as *COL1A2, COL15A1, DCN and*

FBLN1. It is noteworthy that these cells are found in lesions but minimally in the periplaque area

or neurologically normal white matter, indicating their association with pathology in MS. As with

LPC injury, the Cellchat analyses of MS brain cells indicate the immense potential of fibroblasts

to interact with microglia/macrophages and oligodendrocyte lineage cells.

Fibroblasts have been identified in a number of neurological disorders such as

Alzheimer's Disease and stroke[4,40]. By using multiple markers simultaneously, it is likely that

fibroblasts will be detected in the parenchyma of the CNS of most neurological conditions. While

their numbers in the CNS parenchyma do not appear high (an estimated 5-10% on LPC lesions

by spatial transcriptomics), their potent capacity to exclude oligodendrocytes from lesions and

their many potential interactions with microglia/macrophages and CNS cells position them to be

important regulators of pathophysiology and recovery.

In conclusion, we provide evidence that fibroblasts accumulate in the parenchyma of

demyelinated lesions including in MS. We show that fibroblast-occupied areas lack new

myelinating cells, and that fibroblasts inhibit OPC differentiation in vitro. Importantly, this

process is exacerbated by age leading to greater fibrosis of the lesion. These results highlight

the necessity of targeting fibroblasts to mediate regenerative processes and improve outcomes

in neurological conditions.

Acknowledgments

We acknowledge operating grant support from MS Canada, USA Department of Defense

Multiple Sclerosis Research Program, and the Canadian Institutes of Health Research (CIHR).

BML and DM received PhD studentship awards, and RJ a postdoctoral fellowship, from MS

Canada. YD and SG were recipients of postdoctoral awards from CIHR. MM received a

postdoctoral fellowship from Lundbeck Foundation, Denmark.

Contributions of authors

BML conducted the majority of the experiments and acquired and analysed the results; he wrote

the initial drafts of the manuscript. CDM contributed bioinformatics and transcriptomic expertise,

and led the acquisition of home grown or public domain transcriptomic results. YD, GSM, SG,

CL, MM, DM, and RJ provided tissue samples for the manuscript; GSM also provided single-

nucleus RNA sequencing results and figure panels. PE contributed to datasets of the aging

experiments. CCL (Ling) contributed expertise in chemistry of the extracellular matrix. JKH

contributed datasets of single-nucleus RNA sequencing in lysolecithin-demyelinated mice. CCL

(Camara-Lemarroy) contributed datasets of spatial transcriptomics of lysolecithin lesions. All

authors edited and approved the manuscript. VWY oversaw the entire project including the

acquisition of funds and finalizing the manuscript.

Competing interests

The authors declare no conflict of interest.

<u>Materials and Methods</u>

MS specimens

Postmortem frozen brain tissues from people with MS were obtained from The Multiple

Sclerosis and Parkinson's Tissue Bank situated at Imperial College, London

(https://www.imperial.ac.uk/ medicine/multiple-sclerosis-and-parkinsons-tissue-bank). All

samples at their local sites were collected with full informed consent for autopsy. Human tissue

use in Calgary for research was approved by the Conjoint Health Research Ethics Board at the

University of Calgary (Ethics ID REB15-0444). MS sections were characterized using luxol fast

blue (LFB) and hematoxylin & eosin (H&E) to identify lesions as previously described[55].

Mice

All experiments were conducted with ethics approval from the Animal Care Committee at the

University of Calgary under regulations of the Canadian Council of Animal Care. Female

C57BL/6J mice were from Jackson laboratories. CX3CR1-CreER (JAX 021160), iDTR

[Rosa26^{iDTR} (JAX 007900)], PDGFRβ-P2A-CreERT2 (JAX 030201), Ai9 [Rosa26-TdTomato

(JAX 007905)], NG2-CreER (JAX 008538), and Tau-mGFP (JAX 021162) mice were from The

Jackson Laboratory. PDGFRβ-P2A-CreERT2, Ai9 [Rosa26-TdTomato] mice were bred in the

single barrier mouse unit at the University of Calgary to produce female PDGFRβ^{TdTom} mice for

fate mapping experiments. NG2-CreER and Tau(Mapt)-mGFP mice were bred in the single

barrier mouse unit at the University of Calgary to produce female NG2-Mapt[mGFP] mice for OPC

maturation experiments. All young female mice were 6-10 weeks of age, and all middle-aged

female mice were 48-52 weeks of age. Male and female CD1 pups P0-2 were used for OPC

and meningeal fibroblast cultures. C57BL6 female mice 6-10 weeks of age were used for bone

marrow-derived macrophages (BMDM) cultures. All mice were maintained on a 12-h light/dark

cycle with food (Pico-Vac Mouse Diet 20) and water given ad libitum.

Spinal cord surgery

Lysolecithin/lysophosphatidylcholine (LPC) demyelination was accomplished as previously

described[56]. Mice were anaesthetized with intraperitoneal injections of ketamine (100 mgkg-1)

and xylazine (10 mgkg-1). Buprenorphine (0.05 mgkg-1) was given as an analgesic. The

surgery site was shaved and disinfected with 70% ethanol and betadine. Mice were positioned

on a stereotaxic frame and a 2-3 cm incision was made between the shoulder blades. The

intervertebral space between T3 and T4 was identified, blunt dissected apart, and the exposed

dura was removed using a 30-gauge metal needle. Using a 32-gauge needle attached to a 10

µL Hamilton syringe, 0.5 µL of 1% w/v LPC (Sigma-Aldrich, L1381) was injected into the ventral

column of the spinal cord at a rate of 0.25 µLmin-1 for 2 min. The needle was left for 2 min

following the injection to avoid back flow. Muscle and skin were then sutured, and mice placed

in a thermally controlled environment for recovery.

Spinal cord tissue isolation

LPC mice were anaesthetized as described above. Mice were then transcardially perfused with

15 mL of PBS. Spinal cords were removed drop fixed in 4% paraformaldehyde (PFA) at 4°C

overnight. Tissue was cryoprotected in 30% w/v sucrose solution for 72-hours then frozen in

FSC 22 frozen section media (Leica). Coronal sections were generated with a cryostat

(ThermoFisher Scientific) onto Superfrost Plus microscope slides (VWR) and stored at -20°C

until analysis.

Immunofluorescence staining

Slides were thawed for 10 minutes, rehydrated for 10 minutes with PBS, and permeabilized with

0.2% TritonX-100 in PBS for 10 minutes. Tissue was blocked for 1 hour at RT with horse serum

blocking solution (0.01 M PBS, 10% horse serum, 1% bovine serum albumin (BSA), 0.1% cold

fish skin gelatin, 0.1% Triton-X100, and 0.05% Tween-20). Primary staining was done overnight

at RT in buffer (PBS, 1% BSA, 0.1% cold fish stain gelation, 0.1% Triton X-100). Slides were

washed 2 x 5 minutes with PBS, and stained with TrueBlack Lipofuscin Autofluorescence

Quencher (Biotium) for 2 min at room temperature. Slides were washed 3 x 5 minutes, then

incubated with secondary antibodies and 1 µgmL-1 of DAPI resuspended in buffer for 1 hour at

RT. Slides were washed 3 x 5 minutes, and mounted with Fluoromount-G solution

(SouthernBiotech). Antibodies used: anti-mouse myelin basic protein (MBP, BioLegend, PA1-

10008), anti-mouse POSTN (R&D Systems, AF2955), anti-mouse αSMA-Cy3 (Millipore,

C6198), anti-mouse FN1 (Millipore, AB2033), anti-CD45 (Thermofisher, MA5-17687), rabbit

anti-mouse pan-laminin antibody (a gift from L. Sorokin, University of Münster, Münster,

Germany; 1:1000), anti-mouse/human PDGFRβ (R&D Systems, AF1042), anti-mouse Iba1

(Wako, 019-19741), anti-mouse CD45-AF488 (Biolegend, 103122), Col1a1 (Invitrogen, PA1-

26204), anti-mouse GFAP (Biolegend, PCK-591P), anti-mouse NFH (RPCA-NF-H), anti-mouse

PDGFRα (R&D Systems, AF1062), anti-mouse Olig2 (Millipore, ab9610), GFP (Aveslab, GFP-

1020), anti-mouse sulfatide O4 (R&D Systems, MAB1326), anti-mouse Arg1 (BioLegend,

678802), and anti-mouse MHCII (Thermofisher, 13-5321-82). Secondary antibodies from

Jackson ImmunoResearch were used at 1:400 dilution: AF488 donkey anti-mouse IgM, AF488

donkey anti-rabbit IgG, AF488 donkey anti-rat IgG, AF488 donkey anti-goat IgG, cyanine Cy3

donkey anti-goat IgG, cyanine Cy3 donkey anti-chicken IgY, AF647 donkey anti-rat IgG, and

AF647 donkey anti-rabbit IgG.

Confocal immunofluorescence microscopy

Images were acquired on a Leica TCS Sp8 laser confocal microscope or Leica Sp8 Falcon.

Leica TCS images were taken using a ×25/0.5 NA water objective. The 405-nm, 488-nm, 552-

nm and 640-nm lasers were used and detected by two low-dark-current Hamamatsu PMT

detectors and two high-sensitivity hybrid detectors. Leica SP8 Falcon images were taken using

a x20/0.8 NA air objective with 405nm and white light laser (470-670nm), and detected by one

low-dark-current Hamamatsu PMT detector, one cooled high sensitivity, single molecule

detection hybrid detector, and two high sensitivity hybrid detectors. All images were acquired

using the following parameters: in 8 bits, in a z-stack using unidirectional scanning, 1 times

frame averaging, 1 airy unit pinhole, ×1 zoom, 0.57 µm or 1 µm per optical section, and 2,048 x

2,048 or 1024 x 1024 pixels x–y resolution. Equal laser, gain and offset settings to maximize

contrast and minimize saturation were consistently used for all samples within each set of

experiments. All lesions were imaged for volume measurements and sections at the lesion

epicenter were used for all other analysis.

Cell culture

Mouse meningeal fibroblasts

Meninges from P0-2 CD1 pups were collected in RPMI and digested for 18 minutes at 37°C in 1

µgmL-1 DNase I (Sigma) and 3.7 µgmL-1 collagenase D (Sigma) triturating every 6 minutes.

Meninges were centrifuged at 300xg for 3 minutes and cells were resuspended in growth

medium (DMEM (Gibco) supplemented with 10% FBS, 1% non-essential amino-acids (NEAA),

1% GlutaMAX, 1% sodium pyruvate, and 1% penicillin-streptomycin (all from Gibco). Cells were

incubated in T-75 flasks at 37°C with 5% CO2. Medium replaced every 3 days and cells were

passaged when 90% confluent. Cells were used after at least one passage.

Mouse oligodendrocyte progenitor cells (OPCs)

Mouse OPCs were cultured as previously described[16]. Briefly, cortices from P0 CD1 pups were

removed and dissociated with papain (Worthington), DNase (Sigma), and L-cysteine (Sigma) for

30 minutes at 37°C. Cells were cultured in mixed glial culture growth medium in poly-L-lysine

coated T-75 culture flasks. Medium changes were performed 3–4 h, and days 3 and 6.

Supplemented medium was added from day 6. On day 9, flasks were shaken for 45 minutes

and medium replaced with 10-12mL of insulin supplemented mixed glial culture medium. Flasks

were shaken at overnight at 37°C, 5% CO2 and the supernatant was transferred to a 100 mm

tissue culture dish for incubation at 37°C and 5% CO2 for 30 min with gentle swirling by hand

after 15 minutes. OPC enriched medium was collected, centrifuged at 300xg for 10 min, and

seeded at 3,200 cells/cm^2 density in oligodendrocyte differentiating medium. Plates were left at

37°C and 8.5% CO2 for 24–72 h. 96-well flat bottom black/clear plates pre-coated with poly-L-

lysine was used.

Mouse bone marrow-derived macrophages (BMDMs)

Cells were collected by flushing femurs with DMEM supplemented with 10% FBS, 1% penicillin-

streptomycin and 1% glutamax (all Gibco). Cells were centrifuged at 300xg x 10 minutes,

supernatant was discarded, and cells resuspended in BMDM culture medium (10% supernatant

from L929 cell line enriched in macrophage-colony stimulating factor, 10% FBS, 1% penicillin-

streptomycin, 2% minimum essential medium MEM non-essential amino acid solution (all

except L929 from Gibco). Cells were cultured in 100 mm culture plates at 37°C with 8.5% CO2

for 9 days. Medium was half changed day 5 and full changed day 9. Day 9, BMDMs were

collected using cold PBS and scraping cells. Cells were collected into 50 mL tubes, spun at

300xg for 10 minutes, and resuspended in 1 mL of BMDM culture medium for counting.

Cocultures

OPCs were added at a density of 31,250 cells/cm^2 to poly-L-lysine coated 96-well plates.

Meningeal fibroblasts were added at densities of 31.25, 312.5, 3125, 31250, 312500, and 10000

cells/cm^2 as listed. OPCs and meningeal fibroblasts were added together, or one added 2 hours

before the other as defined in the experiment, in 100 µL of OPC differentiating medium.

Cocultures were incubated at 37°C with 8.5% CO_2 for 72 hours. Cells were fixed with 4% PFA

for 10 minutes then stored in PBS until staining. Tissue was blocked with Licor Blocking Buffer

for 1 hour at RT. Anti-O4 primary antibody was added in Licor antibody dilution buffer (1:250)

overnight at 4°C. Cells were washed 3 x 5 minutes with PBS then donkey anti-mouse IgM

AF488 secondary antibodies were added (1:200) for 1 hour at RT. Cells were washed 3 x 5

minutes each in PBS and permeabilized for 5 minutes with 0.02% Triton-X100 in PBS. Cells

were washed once with Licor blocking buffer, then incubated overnight at 4°C with anti-

PDGFRβ (1:50; R&D systems;AF1042) and anti-MBP (1:100; Abcam; ab7349). Cells were

washed in PBS-Tween 0.1% and stained with secondary antibodies (1:200) for 1 hour at RT on

a shaker. Cells were washed 3 x 5 minutes each and suspended in PBS with 1 µg/mL DAPI.

Plates were kept covered at 4°C until imaged.

Transwell migration assay

BMDMs were plated in 24 well plates at a density of 260,000 cells/cm2 overnight at 37°C with

8.5% CO2. Medium with 1% FBS was added for 24 hours prior to experiments. Meningeal

fibroblasts (150,000 cells/cm2) were added to transwell inserts (Corning, 29442-120) for 16-18

hours, then fixed and stained. The upper chamber was cleaned using a cotton swab, inserts

were fixed with acid alcohol (1% HCl in 95% ethanol) for 15 minutes then washed in PBS for 5

minutes. Inserts were transferred to hematoxylin for 15 minutes, and washed in PBS for 1

minutes. Inserts were mounted and imaged on an Olympus VS110 slide scanner on a 10x/0.4

NA objective and VS-ASW-S5 (V2.9) with Batch Converter software.

Fluorescent image analysis

Leica Application Suite X was used for image acquisition, ImageJ for image threshold and

particle analysis, and QuPath 0.3.1 for transwell and slide scanner image analysis. Maximum-

intensity projections were created for each channel/marker. All images were blinded prior to

analysis. ROI for lesions were drawn using DAPI or MBP. Fibroblast regions were defined using

PDGFRβ. Threshold intensity was determined using at least 3 images. The Analyze Particles

function was then used to create a mask and to quantify the positive signals in each ROI using

size exclusion of 2-infinity. Threshold intensity, size exclusion, and circularity settings for particle

analysis were kept constant across all samples for each experimental set. Oligodendrocytes

were counted manually using Olig2 and DAPI as a guide. When analyzing lesion spread

samples were excluded if more than 20% of sections were unable to be analyzed due to tearing,

folding, or other loss.

Quantitative fluorescence microscopy analysis

Plates were imaged using a ×10/0.5 NA air objective on an ImageXpress Micro XLS High-

Content Analysis system (Molecular Devices). The following filters were used for detection:

DAPI, excitation of 387/11 and emission of 447/60; FITC, excitation of 482/35 and emission of

536/40; Cy3, excitation of 531/40 and emission of 593/40; and Cy5, excitation of 628/40 and

emission of 692/40. For quantification, 9–12 FOVs were imaged per well. Multiwavelength cell

scoring analysis in the MetaXpress High-Content Image Acquisition and Analysis software

(Molecular Devices) was used to quantify cell survival from the fluorescence microscopy images

gathered. For fold change values, all numbers were divided by the mean of the control samples.

The fold-change values from all samples and experiments were then plotted for statistical

analysis. For the representative images shown, each channel/marker for the sample was

merged and displayed using pseudocolours.

CosMx spatial molecular imaging

Slides with FFPE sections as prepared above were sent to NanoString for analysis using the

CosMx platform. Sample processing, staining, imaging, and cell segmentation were performed

as previously described[57]. The 1000-plex CosMx Mouse Neuro probe panel was used along

with markers for morphology and cell segmentation: GFAP, Histone_H3, 18S- rRNA and DAPI.

A 0.51mm FOV was centered on the lesion area. The CosMx optical system has an

epifluorescent configuration based on a customized water objective (13×, NA 0.82), and uses

widefield illumination, with a mix of lasers and light-emitting diodes (385 nm, 488 nm, 530 nm,

590 nm, 647 nm) that allow imaging of DAPI, Alexa Fluor-488, Atto-532, Dyomics Dy-605 and

Alexa Fluor-647, as well as removal of photocleavable dye components. Nine Z-stack images

(0.8 um step size) of each FOV were acquired and then fluorophores on the reporter probes

were UV cleaved and washed off with Strip Wash buffer. This procedure was repeated for the

remaining 15 reporter pools, and the 16 cycles of reporter hybridization-imaging was repeated 8

times to increase RNA detection sensitivity. Raw image processing and feature extraction were

performed using an in-house SMI data processing pipeline which includes registration, feature

detection, and localization[57]. A cell segmentation pipeline using machine learning algorithm was

used to accurately assign transcripts to cell locations and subcellular compartments as

previously described[58].

CosMx data processing

Data from the CosMx platform was formulated into an object for Seurat v5 package [59] in R

programming environment. Data from 2 LPC and 3 Sham brain sections with FOVs from the

right hemisphere were used. Data was normalized using SCTransform v2. An unsupervised

analysis and generation of a final UMAP with 20 PCs and resolution = 0.7 was done. The

package Harmony [60] was used to integrate 'Group' data (group.by.vars = group (LPS or Sham).

Robust Cell Type Decomposition (RCTD) was used to map the cluster profiles to the Azimuth

mouse motor cortex reference map (azimuth.hubmapconsortium.org). The following cell types

were identified: astrocytes, endothelial cells, neurons, macrophages, oligodendrocytes,

oligodendrocyte precursor cells (OPCs), vascular leptomeningeal cells (VLMCs) and pericytes.

A separate identity was created for fibroblasts as cells which expressed *Col1a2* and *Pdgfrb*, and

OPCs as being *Pdgfra* and *Sox6* positive. The FindAllMarkers function with default Wilcoxon

rank sum test was used to identify distinguishing markers for each cluster. The DoHeatmap

function with downsampling to 1000 was used to generate a heatmap with top 5 genes for each

cluster. Data from macrophages, fibroblasts and OPCs from LPC sections were subset from the

main dataset and used with the package CellChat v2 to assess ligand-receptor interactions[61].

To assess significant ligand-receptor interactions the computeCommunProb command was

used with the following parameters (type="truncatedMean", distance.use=TRUE,

contact.dependent=TRUE, contact.range=50).

Analysis of mouse single nuclei dataset from LPC lesions

LPC-lesioned tissue dataset from Melchor et al. (accession number GSE278643) was used to

validate fibroblast population increases in LPC. Briefly, processed data were directly

downloaded, and all quality control, dimensionality reduction, and clustering were performed

using Seurat (v5.1.0) in R (v4.4.1) as per original publication. All samples (naïve spinal cord

tissue (n=2), LPC and sham 5dpl (n=3, n=1), 10dpl (n=2, n=1), and 20dpl (n=3, n=1) were

utilized. To cluster the data, we used the SCTransform v2 function[62,63]. The count matrix was log

normalized using the NormalizeData function to identify cluster gene markers. All cell

populations were annotated based off of well-known and calculated marker genes through the

FindAllMarkers function. Vascular and mesenchymal cells (*Pdgfrb*, *Cfap43*, *Pecam1*, *Cspg4*,

Col1a2) were subset based off marker genes. The subset vascular/mesenchymal dataset was

reclustered (65 PCs and resolution = 2.2) and filtered following the original publication, count

matrices were renormalized using the NormalizeData function, and clusters were manually

annotated based on the expression of cell type-specific genes.

Analysis of publicly available human single nuclei dataset from MS patients

Fastq files from the human MS single nuclei transcriptomic study by Absinta et al. (accession

number GSE180759) were downloaded and demultiplexed using 10x Genomics Cellranger

version 5.0 pipeline. The final expression matrix with gene counts from n=3 controls and n=5

patients was then analysed using the bioinformatics package Seurat v5[59]. Metadata defining

tissue area was included in the object. Before running QC metrics the expression matrix

comprised 34,131 features and 134,932 cells. Data was filtered for parameters: gene present in

> 3 cells and cells with < 150,000 nCount_RNA and percent of mitochondrial genes < 10%. Post

filtering, the expression matrix contained 33,454 features and 128,343 cells. Data was

normalized and scaled using the SCTransform v2 function with percent.mt being regressed out.

A PCA-reduction was performed and 30 significant dimensions were considered to generate a

UMAP with all cells in the dataset. Clusters were determined using the FindClusters function

which implements the Louvain algorithm for modularity optimization and with resolution of 0.3.

Cluster annotation was done manually based on the expression of lineage specific hallmark

genes. Differentially expressed genes for one cluster (versus all cells in other clusters) was

determined by the default Wilcoxon rank sum test. PDGFRB+ cells from the initial fibroblast

cluster were then subset and re-clustered. Data from microglia/macrophages, OPCs and

PDGFRB+ fibroblasts were subset from the main data and used with the package CellChat2 v2

to assess ligand-receptor interactions[61]. To assess significant ligand-receptor interactions the

computeCommunProb command was used with the following parameters

(type="truncatedMean", trim = 0.05, population.size=TRUE).

Statistical analysis

Data was collated using Microsoft Excel. Graphs were generated using GraphPad Prism 8. Data

shown are the individual data points, whereby each point on a bar graph represents a biological

(for in vivo experiments) or technical replicate (for in vitro experiments), and the mean. No

statistical methods were used to predetermine sample sizes, but sample sizes were similar to

those previously reported[28,55,64]. Experimental groups were randomly assigned. Data collection

and analysis were not performed blind to the conditions of the experiment (unless otherwise

stated), as all image and data analyses were completed with the same acquisition conditions

and analysis thresholds. Statistical tests are listed in figure legends.

Legend to Figures

Fig 1. Fibroblasts localize in LPC demyelinated SCWM. a-d. Immunofluorescence images of

day 7 LPC lesions labeled with: **(a)** DAPI, MBP, NFH and GFAP in wildtype mice; **(b)** DAPI,

Iba1 and TdTomato (TdTom) in PDGFRβ-TdTom mice; **(c)** DAPI, TdTom and PDGFRβ in

PDGFRβ-TdTom mice; **(d)** DAPI, PDGFRβ, COL1A1 and SMA in wildtype mice. Scale bar =

100 μm, inset = 25 μm. **e-j.** Graphs comparing day 7 and day 21 **(e, h)** lesion volume, **(f, i)**

TdTom positive volume, **(g, j)** proportion of LPC lesion that is TdTom positive. **k.** Representative

images of day 7 and 21 LPC lesions labeling for PDGFRβ-TdTom and **(left)** periostin (POSTN),

(middle) fibronectin (FN1), and **(right)** SMA. Scale bar = 100 μm. **l-n.** Graphs comparing the

proportion of day 7 and 21 lesions positive for **(l)** POSTN, **(m)** FN1, **(n)** SMA. Data was acquired

from two experiments; n=8 per experimental group. Significance is indicated as *P<0.05,

**P<0.01, ***P<0.001, ****P<0.0001; unpaired *t*-test, comparing day 7 and day 21 LPC lesions.

Data presented as the mean.

Fig 2. Single-nucleus RNA sequencing confirms the prominence of fibroblasts in LPC

lesions. a. Dotplot of broadly annotated vascular and mesenchymal cell populations

highlighting marker genes used for annotation. **B.** UMAP plot of broadly annotated vascular and

mesenchymal cell populations (*left*), and at the different timepoints investigated post-

demyelination (**c**). **d.** Bar plot showing the relative frequencies of broadly annotated

vascular/mesenchymal subpopulations in each experimental timepoint, highlighting the

prominence of fibroblasts in lesioned samples.

Fig 3. Spatial transcriptomics of fibroblasts in demyelinated lesions. **a, b.** Representative

images of annotated cells in **(a)** sham and **(b)** LPC demyelinated spinal cords at 14 dpi. Lesion

edge indicated by white dashed line. **c, d.** UMAP visualization of 2954 cells from LPC and sham

colored by **(c)** experimental group and **(d)** mapped cell clusters (20 principal components,

resolution=0.7). **e.** Heatmap of top 5 differentially expressed genes across annotated cell

groups as determined using FindAllMarkers (default Wilcoxon Rank Sum test, min. logfc

threshold = 0.25). **f, g.** Bar graph showing **(f)** the total number of cells, and **(g)** the fold change

in cell groups between LPC and sham groups. Abbreviations: M/M – microglia/macrophages;

OL – oligodendrocytes; OPCs – oligodendrocyte precursor cells; EC – endothelial cells; VLMC –

vascular leptomeningeal cells.

Fig 4. Inferred Incoming and outgoing signaling patterns between fibroblasts, OPCs and

macrophages in day 14 LPC lesions. Heatmap of **(a)** inferred outgoing and **(b)** incoming

signaling pathways as assessed for the 3 cell groups. Color bar in the heatmap represents the

relative signaling strength of a pathway across all cell groups (values are row-scaled). Top

colored bar plot shows the total signaling strength of a cell group by summarizing all pathways

displayed in the heatmap. The right grey bar plot shows the total signaling strength of a pathway

by summarizing all cell groups displayed in the heatmap. Heatmap of **(c)** outgoing and **(e)**

incoming signaling pathways that contribute to each communication pattern. River plots showing

the cell source for **(d)** outgoing and **(f)** incoming communication patterns and the pathways

associated with them. The thickness of the flow indicates contribution of cell group or signaling

pathway to each pattern.

Fig 5. Post inflammation fibroblast response stimulated by microglia/macrophages. **a, b.**

Representative image of **(a)** day 3 and **(b)** day 7 LPC lesion labelling for DAPI, PDGFRβ, and

Iba1. Scale bar = 100 μm. **b.** Representative images of LPC lesioned SCWM day 3, 7, 14, and

21 day post injury labelling for DAPI, PDGFRβ, and Iba1. Scale bar = 100 μm (insert = 25 μm).

c, d. Graphs indicating day 3, 7, 14, and 21 day **(c)** Iba1 positive % area of lesion and **(d)**

PDGFRβ positive % area of lesion. **e.** Schematic of experimental design. Meningeal fibroblasts

were added to upper compartment and BMDMs or medium alone added to the lower

compartment. **f.** Representative images of hematoxylin stained transwell inserts from BMDM

and medium control groups. **g.** Graph comparing the amount of hematoxylin stained meningeal

fibroblasts migrated across the transwell membrane in BMDM and medium alone control. **(c, d)**

n=5 (D3), 10 (D7), 7 (D14), 5 (D21); **(g)** n = 4 for each experimental group; Significance

indicated as *P<0.05, **P<0.01, one-way ANOVA comparing time points (D3, 7, 14, or 21) **(c,**

d); or two-tailed unpaired *t*-test **(g).** All data presented as the mean.

Fig 6. Fibroblasts spatially exclude OPCs in LPC lesions and in vitro. **a.** Representative

images of day 14 LPC lesion stained for PDGFRβ, Olig2, and PDGFRα. **b.** Representative

images of day 14 LPC lesion in NG2CreER:MAPTmGFP mice labelling for DAPI, PDGFRβ,

NG2-GFP. **c.** Graph comparing GFP positive area in day 14 LPC lesions. PDGFRβ positive

area of LPC lesion (red) and PDGFRβ negative area of LPC lesion (white). n = 9 , Significance

indicated ***P<0.001, two-tailed unpaired *t*-test. Data presented as mean. **d.** Representative

image of OPC alone (left) and OPC-fibroblast co-cultures (right) labelled for nuclei (blue), O4

(green), MBP (red), and PDGFRβ (white). **e-g.** Graphs indicating fold change in **(e)** O4 positive

OPCs, **(f)** O4 positive MBP positive mature oligodendrocytes, and **(g)** mean processes per cell

comparing co-cultures with OPC alone controls. n = 2 independent experiments with 4

replicates for e-g. Significance indicated as *P<0.05, **P<0.01, two-tailed, unpaired *t*-test, Data

presented as mean.

Fig 7. Age exacerbates fibroblast response to demyelination. **a.** Representative

immunofluorescence images of day 7 LPC lesions in young and middle-aged mice labeled with

DAPI and MBP (red) or PDGFRβ (white). Lesion is indicated with white dashed line. **b-f.** Graphs

comparing spinal cord spread of **(b, d)** total lesion volume, **(c, e)** PDGFRβ positive

immunofluorescence, and **(f)** proportion of lesion volume that is PDGFRβ positive at days 7, 14,

and 21 after LPC injury. For young group, n = 4 (D7), 4 (D14), 5 (D21; for middle-aged group n

= 5 for each time point. Significance is indicated as *P<0.05, **P<0.01; unpaired, *t*-test,

comparing young and middle-aged groups. Data is presented as the mean.

Fig 8. Age enhances fibroblast properties in demyelinated lesions. **a.** Representative

immunofluorescence images of day 7 LPC lesions in young and middle-aged mice labeled for

DAPI, PDGFRβ and SMA; and quantification of the proportion of PDGFRβ overlapping with

SMA at 7 and 21 dpi in young and middle-aged mice. **b.** Representative immunofluorescence

images of day 7 LPC lesion of young mice labeled for DAPI, PDGFRβ, and Col1a1; and

quantification of Col1a1 positive area at 7 and 21 dpi in young and middle-aged mice. **c.**

Representative immunofluorescence images of day 7 LPC lesion in young mice labeled for

DAPI and LAMA (laminin); and quantification of LAMA positive area at 7 and 21 dpi in young

and middle-aged mice. **d.** Representative immunofluorescence images of day 7 LPC lesion in

young mice labeled for DAPI, Olig2, and PDGFRβ; graphs comparing the number of Olig2

positive cells within PDGFRβ positive regions of the LPC lesion at 7 and 21 days post LPC

injury. In the graphs, each dot is an individual mouse. Significance is indicated as *P<0.05,

**P<0.01, ***P<0.001; unpaired, *t*-test, comparing young and middle-aged groups. Data

presented as mean.

Fig 9. Fibroblasts in MS lesions. **a.** Representative immunofluorescence images of MS

lesions labeled for DAPI, CD45, and PDGFRβ. This is observed in 5 different MS cases (not

shown). **b.** UMAP visualization of 66,432 cells from MS lesions and their cell annotation[39]. **c.**

Dot plot indicating markers used to identify fibroblast population. Size of the dot indicates the

percentage of cells while the color encodes the average expression levels across all cells. **d.**

Violin plots depicting log normalized gene expression of fibroblast markers comparing PDGFRβ

positive fibroblast population to other cell groups. **e, f.** UMAP visualization of **(e)** subclusters of

PDGFRβ positive fibroblast population with top 10 DEGs, and **(f)** distribution of PDGFRβ

positive fibroblast subclusters across MS lesion types and frontal white matter from

neurologically normal controls.

Fig 10. Fibroblast interactions in MS lesions. **a, b.** Heatmap and river plots indicating cell

and signaling pathways associated with outgoing and incoming communication patterns in MS

lesions. The thickness of the flow indicates contribution of cell group or signaling pathway to

each pattern. **c, d.** Venn diagrams of (l) outgoing and (m) incoming signaling pathways

comparing MS and LPC lesions.

Legend to Supplementary figures

Supp Fig 1. Ligand – receptor (L-R) interactions between fibroblasts, OPCs, and

microglia/macrophages in day 14 LPC lesions (related to main Figure 4). a, c. Number of

significant ligand-receptor between any pair of two cell populations shown as **(a)** circle plot and

(c) heatmap. **b, d.** Interaction strength between ligand-receptor of any pair of two cell

populations shown as **(b)** circle plot and **(d)** heatmap. In the circle plots, the circle sizes are

proportional to the number of cells in each group; edge weights are proportional to the

interaction strength. **e. Scatter plot** of incoming and outgoing interaction strength showing

fibroblasts as major outgoing signaling source. Dot size is proportional to the number of inferred

links (outgoing and incoming) associated with each cell group. **f, g**. Chord plot showing L-R

interactions with fibroblasts as **(f)** source of signal.

Supp Fig 2. Incoming and outgoing fibroblast L-R interactions in day 14 LPC lesions

(related to main Figure 4). a, b. Bubble plot delineating L-R interactions from one cell group

(i.e. sender) to another (i.e. target) as indicated in the plot.

Supp Fig 3. Cell contributions to communication networks (related to main Figure 4). a, b.

Dot plots indicating cell type contribution to **(a)** outgoing and **(b)** incoming communication

patterns.

Supp Fig 4. Axon densities are unchanged in fibroblast occupied regions. a.

Representative image of a day 21 LPC lesion stained for DAPI, NFH, and PDGFRβ. **b.**

Quantification comparing NFH positive axon density in fibroblast occupied areas with the rest of

the LPC lesion. Sample sizes n = 7 across two experiments. Two-tailed, unpaired *t*-test

comparing groups. Data are the mean ± s.d.

Supp Fig 5. Fibroblast density is associated with reduced OPC maturation (related to

main Figure 5). a. Quantification of total DAPI+ cells with 10, 100, 1k, 10k, 100k fibroblasts

added. **b.** Mean outgrowth per cell with 10, 100, 1k, 10k, 100k fibroblasts added. **c.** Graphs

comparing OPCs cultured alone, with fibroblast conditioned medium (CM), or with fibroblasts

showing mean outgrowth per cell. **d.** Graphs comparing OPCs cultured alone, added before

fibroblasts, and added after fibroblasts showing mean outgrowth per cell. Significance is

indicated as *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. Sample sizes n = as shown

number of technical replicates. Kruskal-Wallis followed by a Dunn's multiple comparisons. Mean

shown as horizontal line.

Supp Fig 6. Main cell types as assessed in MS lesions (related to main Figure 9). a.

Heatmap of top 5 differentially expressed genes across annotated cell groups.

Supp Fig 7. Non-fibroblast related marker expression across cell clusters assessed in MS

lesions (related to main Figure 9). a. Violin plots of *TAGLN* and *RGS5* normalised gene

expression in snRNAseq dataset. **b, c.** String-db plot of DEG between fibroblast **(b)** subcluster

0, and **(c)** subclusters 1.

Supp Fig 8. L-R interactions between fibroblasts, M/M and OPcs in MS lesions (related to

main Figure 10). a, c. Number of significant ligand-receptor between any pair of two cell

populations shown as **(a)** circle plot and **(c)** heatmap. **b, d.** Interaction strength between ligand-

receptor any pair of two cell populations shown as **(b)** circle plot and **(d)** heatmap. In the circle

plots, the circle sizes are proportional to the number of cells in each group; edge weights are

proportional to the interaction strength. **e.** Schematic indicating inferred outgoing and incoming

fibroblast signals and their distribution across signaling databases. **f, g.** Chord plot showing L-R

interactions with fibroblasts as **(f)** source of signals, and **(g)** receiver of signals.

Supp Fig 9. Inferred Fibroblast communication patterns in MS (related to main Figure 10).

a, b. Bubble plots delineating L-R interactions from one cell group (i.e. sender) to another (i.e.

target) as assessed with different signaling databases indicated in the plot **(a)** ECM-receptor

and **(b)** secreted signaling databases. **c, d.** Dot plots of **(c)** outgoing and **(d)** incoming

communication patterns.

Supp Fig 10. Inferred Communication pattern contributions in MS lesions (related to main

Figure 10) a, b. Heatmap of **(a)** inferred outgoing and **(b)** incoming signaling pathways as

assessed for the 3 cell groups. Colour bar in the heatmap represents the relative signaling

strength of a pathway across all cell groups (values are row-scaled). Top colored bar plot shows

the total signaling strength of a cell group by summarizing all pathways displayed in the

heatmap. The right grey bar plot shows the total signaling strength of a pathway by summarizing

all cell groups displayed in the heatmap. **c, d.** Venn-diagrams comparing LPC (CosMx) and MS

(snRNAseq) **(c)** outgoing and **(d)** incoming fibroblast L-R interactions.

References

1. Plikus, M. V. *et al.* Fibroblasts: Origins, definitions, and functions in health and disease. *Cell* **184**, 3852–3872 (2021).

2. Zhou, X. *et al.* Microenvironmental sensing by fibroblasts controls macrophage population size. *Proc Natl Acad Sci U S A* **119**, e2205360119 (2022).

3. Plikus, M. V. *et al.* Regeneration of fat cells from myofibroblasts during wound healing. *Science* **355**, 748–752 (2017).

4. Dorrier, C. E., Jones, H. E., Pintarić, L., Siegenthaler, J. A. & Daneman, R. Emerging roles for CNS fibroblasts in health, injury and disease. *Nature Reviews Neuroscience* **23**, 23–34 (2022).

5. Dorrier, C. E. *et al.* CNS fibroblasts form a fibrotic scar in response to immune cell infiltration. *Nature Neuroscience* **24**, 234–244 (2021).

6. Bolte, A. C. *et al.* The meningeal transcriptional response to traumatic brain injury and aging. *eLife* **12**, 1–38 (2023).

7. Yahn, S. L. *et al.* Neurobiology of Disease Fibrotic scar after experimental autoimmune encephalomyelitis inhibits oligodendrocyte di ff erentiation. *Neurobiology of Disease* **134**, 104674 (2020).

8. Liu, X. *et al.* Reactive Fibroblasts in Response to Optic Nerve Crush Injury. *Mol Neurobiol* **58**, 1392–1403 (2021).

9. Şekerdağ-Kılıç, E. *et al.* Perivascular PDGFRB+ cells accompany lesion formation and clinical evolution differentially in two different EAE models. *Mult Scler Relat Disord* **69**, 104428 (2023).

10. Goritz, C. *et al.* A Pericyte Origin of Spinal Cord Scar Tissue. *Science* **333**, 238–243 (2011).

11. Dias, D. O. *et al.* Reducing Pericyte-Derived Scarring Promotes Recovery after Spinal Cord Injury. *Cell* **173**, 153–165 (2018).

12. Dias, D. O. *et al.* Pericyte-derived fibrotic scarring is conserved across diverse central

nervous system lesions. *Nat Commun* **12**, 5501 (2021).

13. Franklin, R. J. M. & Simons, M. CNS remyelination and inflammation: From basic

mechanisms to therapeutic opportunities. *Neuron* **110**, 3549–3565 (2022).

14. Lubetzki, C., Zalc, B., Williams, A., Stadelmann, C. & Stankoff, B. Remyelination in multiple

sclerosis : from basic science to clinical translation. *The Lancet Neurology* **19**, 678–688

(2020).

15. Mei, F. *et al.* Accelerated remyelination during inflammatory demyelination prevents axonal

loss and improves functional recovery. *eLife* **5**, 1–21 (2016).

16. Ghorbani, S. *et al.* Versican promotes T helper 17 cytotoxic inflammation and impedes

oligodendrocyte precursor cell remyelination. *Nature Communications* **13**, 1–18 (2022).

17. Rawji, K. S. *et al.* Niacin-mediated rejuvenation of macrophage/microglia enhances

remyelination of the aging central nervous system. *Acta Neuropathologica* **139**, 893–909

(2020).

18. Dong, Y. *et al.* Single-cell and spatial RNA sequencing identify perturbators of microglial

functions with aging. *Nature Aging* (2022) doi:10.1038/s43587-022-00205-z.

19. Koch, M., Mostert, J., Heersema, D. & De Keyser, J. Progression in multiple sclerosis:

Further evidence of an age dependent process. *Journal of the Neurological Sciences* **255**,

35–41 (2007).

20. Brack, A. S. *et al.* Increased Wnt signaling during aging alters muscle stem cell fate and

increases fibrosis. *Science* **317**, 807–810 (2007).

21. Chambers, E. S. *et al.* Recruitment of inflammatory monocytes by senescent fibroblasts

inhibits antigen-specific tissue immunity during human aging. *Nature Aging* **1**, (2021).

22. Salzer, M. C. *et al.* Identity Noise and Adipogenic Traits Characterize Dermal Fibroblast

Aging. *Cell* **175**, 1575-1590.e22 (2018).

23. Mahmoudi, S. *et al. Heterogeneity in Old Fibroblasts Is Linked to Variability in*

Reprogramming and Wound Healing. Nature vol. 574 (Springer US, 2019).

24. Shen, S. *et al.* Age-dependent epigenetic control of differentiation inhibitors is critical for

remyelination efficiency. *Nature Neuroscience* **11**, 1024–1034 (2008).

25. Nicaise, A. M. *et al.* Cellular senescence in progenitor cells contributes to diminished

remyelination potential in progressive multiple sclerosis. *Proceedings of the National*

Academy of Sciences of the United States of America **116**, 9030–9039 (2019).

26. Neumann, B. *et al.* Metformin Restores CNS Remyelination Capacity by Rejuvenating Aged

Stem Cells. *Cell Stem Cell* **25**, 473-485.e8 (2019).

27. Anderson, M. A. *et al.* Astrocyte scar formation aids central nervous system axon

regeneration. *Nature* **532**, 195–200 (2016).

28. Plemel, J. R. *et al.* Microglia response following acute demyelination is heterogeneous and

limits infiltrating macrophage dispersion. *Science Advances* **6**, eaay6324 (2020).

29. Lozinski, B. M. *et al.* Exercise rapidly alters proteomes in mice following spinal cord

demyelination. *Scientific Reports* **11**, (2021).

30. Henderson, N. C. *et al.* Targeting of αv integrin identifies a core molecular pathway that

regulates fibrosis in several organs. *Nature Medicine* **19**, 1617–1624 (2013).

31. Buhl, E. M. *et al.* Dysregulated mesenchymal PDGFR-β drives kidney fibrosis. *EMBO*

Molecular Medicine **12**, (2020).

32. Holl, D. *et al.* Distinct origin and region-dependent contribution of stromal fibroblasts to

fibrosis following traumatic injury in mice. *Nat Neurosci* (2024) doi:10.1038/s41593-024-

01678-4.

33. Melchor, G. S., Baydyuk, M., Manavi, Z., Hu, J. & Huang, J. K. Dissecting the evolving

cellular landscape of a remyelinating microenvironment. 2024.12.25.630253 Preprint at

https://doi.org/10.1101/2024.12.25.630253 (2024).

34. Jensen, S. K. *et al.* Multimodal Enhancement of Remyelination by Exercise with a Pivotal

Role for Oligodendroglial PGC1α. *Cell Reports* **24**, 3167–3179 (2018).

35. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat Commun*

12, 1088 (2021).

36. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. Hallmarks of aging:

An expanding universe. *Cell* **186**, 243–278 (2023).

37. Segel, M. *et al.* Niche stiffness underlies the ageing of central nervous system progenitor

cells. *Nature* **573**, 130–134 (2019).

38. Goldschmidt, T., Antel, J., Konig, F. B., Bruck, W. & Kuhlmann, T. Remyelination capacity of

the MS brain decreases with disease chronicity. *Neurology* **72**, 1914–1921 (2009).

39. Absinta, M. *et al.* A lymphocyte – microglia – astrocyte axis in chronic active multiple

sclerosis. *Nature* (2021) doi:10.1038/s41586-021-03892-7.

40. Yang, A. C. *et al.* A human brain vascular atlas reveals diverse mediators of Alzheimer's

risk. *Nature* **603**, 885–892 (2022).

41. Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain

vasculature. *Nature* **554**, 475–480 (2018).

42. Cregg, J. M. *et al.* Functional Regeneration Beyond the Glial Scar. *Exp Neurol* **0**, 197–207

(2014).

43. Bernier, L.-P. *et al.* Brain pericytes and perivascular fibroblasts are stromal progenitors with

dual functions in cerebrovascular regeneration after stroke. *Nat Neurosci* **28**, 517–535

(2025).

44. Rustenhoven, J. *et al.* Functional characterization of the dural sinuses as a neuroimmune

interface. *Cell* **184**, 1000-1016.e27 (2021).

45. Donovan, J., Shiwen, X., Norman, J. & Abraham, D. Platelet-derived growth factor alpha

and beta receptors have overlapping functional activities towards fibroblasts. *Fibrogenesis*

Tissue Repair **6**, 10 (2013).

46. Zhou, X. *et al.* Circuit Design Features of a Stable Two-Cell System. *Cell* **172**, 744-757.e17

(2018).

47. Adler, M. *et al.* Principles of Cell Circuits for Tissue Repair and Fibrosis. *iScience* **23**,

100841 (2020).

48. Stephenson, E. L. *et al.* Chondroitin sulfate proteoglycans as novel drivers of leucocyte

infiltration in multiple sclerosis. *Brain* 1094–1110 (2018) doi:10.1093/brain/awy033.

49. Ghorbani, S. & Yong, V. W. The extracellular matrix as modifier of neuroinflammation and

remyelination in multiple sclerosis. *Brain* **144**, 1958–1973 (2021).

50. Wang, F. *et al.* Myelin degeneration and diminished myelin renewal contribute to age-

related deficits in memory. *Nature Neuroscience* **23**, 481–486 (2020).

51. Shields, S. A., Gilson, J. M., Blakemore, W. F. & Franklin, R. J. M. Remyelination occurs as

extensively but more slowly in old rats compared to young rats following gliotoxin-induced

CNS demyelination. *Glia* **28**, 77–83 (1999).

52. Rawji, K. S. *et al.* Immunosenescence of microglia and macrophages: Impact on the ageing

central nervous system. *Brain* **139**, 653–661 (2016).

53. Müller, L. & Di Benedetto, S. Aging brain: exploring the interplay between bone marrow

aging, immunosenescence, and neuroinflammation. *Front Immunol* **15**, 1393324 (2024).

54. Hoeft, K. *et al.* Platelet-instructed SPP1+ macrophages drive myofibroblast activation in

fibrosis in a CXCL4-dependent manner. *Cell Reports* **42**, 112131 (2023).

55. Dong, Y. *et al.* Oxidized phosphatidylcholines found in multiple sclerosis lesions mediate

neurodegeneration and are neutralized by microglia. *Nature Neuroscience* **24**, 489–503

(2021).

56. Keough, M. B., Jensen, S. K. & Wee Yong, V. Experimental demyelination and

remyelination of murine spinal cord by focal injection of lysolecithin. *Journal of Visualized*

Experiments **2015**, 1–8 (2015).

57. He, S. *et al.* High-plex imaging of RNA and proteins at subcellular resolution in fixed tissue
by spatial molecular imaging. *Nat Biotechnol* **40**, 1794–1806 (2022).

58. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nat Methods* **18**, 100–106 (2021).

59. Hao, Y. *et al.* Dictionary learning for integrative, multimodal and scalable single-cell
analysis. *Nat Biotechnol* **42**, 293–304 (2024).

60. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with Harmony.
Nat Methods **16**, 1289–1296 (2019).

61. Jin, S., Plikus, M. V. & Nie, Q. CellChat for systematic analysis of cell–cell communication
from single-cell transcriptomics. *Nat Protoc* **20**, 180–219 (2025).

62. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. *Genome Biol* **20**, 296 (2019).

63. Choudhary, S. & Satija, R. Comparison and evaluation of statistical error models for scRNA-
seq. *Genome Biol* **23**, 27 (2022).

64. Jain, R. W., Elliott, D. A. & Yong, V. W. Single Cell Analysis of High-Parameter Histology
Images Using Histoflow Cytometry. *The Journal of Immunology* **210**, 2038–2049 (2023).

1009

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

Fig 9

Fig 10