Concurrent category-selective neural activity across the ventral occipito-temporal cortex supports a non-hierarchical view of human visual recognition

Corentin Jacques[1], Jacques Jonas[1,2] , Sophie Colnat-Coulbois[3], Bruno Rossion*[1,2]

1   Université de Lorraine, CNRS, IMoPA, F-54000 Nancy, France
2   Université de Lorraine, CHRU-Nancy, Service de Neurologie, F-54000 Nancy, France
3   Université de Lorraine, CHRU-Nancy, Service de Neurochirurgie, F-54000 Nancy, France

*Corresponding Author:
Bruno Rossion
@mail: bruno.rossion@univ-lorraine.fr

IMoPA, UMR 7365, CNRS - Université de Lorraine
Pavillon Krug (1er étage - entrée CC-1), Hôpital Central
CHRU Nancy - University Hospital of Nancy
29 Avenue du Maréchal de Lattre de Tassigny
54000 NANCY FRANCE

Authors contribution: Corentin Jacques : Conceptualization, Methodology, Software, Formal analysis, Visualization, Writing - Original Draft, Writing - Review & Editing ; **Jacques Jonas** : Conceptualization, Methodology, Investigation, Resources, Writing - Original Draft, Writing - Review & Editing; **Sophie Colnat-Coulbois** : Resources, Investigation; **Bruno Rossion** : Conceptualization, Methodology, Project administration, Supervision, Funding acquisition, Writing - Original Draft, Writing - Review & Editing .

Competing Interest Statement : The authors declare no competing interests.

Keywords: human face recognition; cortical hierarchy, SEEG; ventral occipito-temporal cortex; high-frequency broadband gamma.

ABSTRACT

Visual recognition is a fundamental human brain function, supported by a network of regions in the ventral occipito-temporal cortex (VOTC). This network is thought to be organized hierarchically, with definite processing stages increasing in invariance and time-course from posterior to anterior cortical regions. Here we provide a stringent test of this view by measuring category-selective neural activity to natural images of faces across the VOTC with electrophysiological intracerebral recordings in a large human sample (N=140; >11000 recording sites). Face-selective high frequency broadband (30-160 Hz) neural activity is distributed across the VOTC, with right-hemispheric dominance and regional peaks of activity. Crucially, while a progressive increase in degree of category-selectivity is found along the postero-anterior axis, neural activity occurs largely concurrently (~100ms onset – ~450ms offset) across all VOTC regions. These observations challenge the standard hierarchical view of neural organization of visual object recognition in the human association cortex, calling for alternative models of this key brain function.

INTRODUCTION

Recognition, i.e., reproducible discrimination, of signals from the sensory environment is a fundamental function of all central nervous systems (Edelman, 2004). Vision is considered the dominant sensory modality for most primates, who have high visual acuity and excellent binocular vision for recognizing their environment (Jacobs, 1999). In the primate order, visual recognition is supported by a wide bilateral network of brain regions located in the occipital and temporal cortices (Conway, 2018; DiCarlo et al., 2012; Ungerleider and Bell, 2011). Based essentially on non-human primates research, this cortical network is generally conceived as being organized hierarchically, with definite processing stages increasing progressively in complexity of representation from posterior to anterior cortical regions, leading ultimately to rich invariant visual representations readily available for memory associations (Conway, 2018; DiCarlo et al., 2012; Freiwald, 2020; Grill-Spector et al., 2017; Hubel and Wiesel, 1968; Issa et al., 2018; Marr, 1982; Riesenhuber and Poggio, 1999; Tsao, 2014; Van Essen et al., 1992; Zeki and Shipp, 1988). However, beyond primary sensory brain structures (i.e., retina, lateral geniculate nucleus, primary visual cortex), direct evidence for this hierarchical organization of visual recognition in the primate brain, in particular the human association cortex (Krienen and Buckner, 2020), is lacking, and the hierarchical view remains subject to debate (Bullier, 2001; Mumford, 1992; Rossion, 2022). Here we provide important information to evaluate the hierarchical view of visual recognition through an extensive characterization of the time-course of category-selective brain activity in the human association cortex.

To do so, we take advantage of the high spatial and temporal resolution provided by intracerebral electrophysiological (iEEG) recordings in a particularly large sample of individual human brains (N=140) implanted from posterior to anterior regions of their ventral occipito-temporal cortex (VOTC; >11000 intracerebral recording sites). With iEEG frequency-tagging, we isolate category-selective high frequency ('high frequency broadband', 30-160 Hz) neural activity to natural images of faces – arguably the most familiar and ecologically valid stimulus in the human environment-, to characterize its time-course across the whole VOTC. This allows testing two essential features of a hierarchical organization (Bullier and Nowak, 1995; Schmolesky et al., 1998): (1) an increase in representation complexity or abstraction along the hierarchy and (2) a progressive increase in response timing at each level of the hierarchy, its inputs being driven by the output from the previous level.

Relative to natural images of various object shapes, we find consistent face-selective neural activity throughout the VOTC, with a progressive increase in face-selectivity, or abstraction, along the postero-anterior axis, reaching a high proportion of exclusive response to faces in the most anterior regions of the ventral temporal lobe. While these findings apparently support a hierarchical organization of visual recognition, face-selective neural activity occurs largely concurrently (~100ms onset – ~450ms offset latency) across the VOTC, challenging a hierarchical organization of visual recognition in the human association cortex.

RESULTS

Neural signal was measured in 140 human participants implanted with intracerebral electrodes (**Figure 1A**) while they viewed sequences of variable natural images of objects presented at a rapid periodic rate (6 Hz, or 167ms SOA, one fixation/image). Highly variable natural images of faces appeared every 5 stimuli (i.e., 6/5 = 1.2 Hz; **Figure 1B-C**). This stimulation mode has been validated in tens of studies with multiple recording techniques (e.g., EEG (Rossion et al., 2015) ; MEG (Hauk et al., 2021); intracerebral recordings (Jacques et al., 2022; Jonas et al., 2016a); fMRI (Gao et al., 2018)), providing high signal-to-noise ratio (SNR) face-selective activity. In this stimulation mode, the periodic modulation of the high frequency broadband (HFB, 30-160 Hz, **Figure 1D**) signal allows to objectively (i.e., at pre-determined frequencies (Norcia et al., 2015; Regan, 1989)) identify and quantify both the neural activity common to faces and nonface objects at 6 Hz and harmonics, as well as the face-selective (i.e., differential) activity identified at the frequency of face stimulation (1.2 Hz) and specific harmonics (e.g., 2.4 Hz, 3.6 Hz,…; **Figure 1E**) in the frequency-domain (Jacques et al., 2022; Retter and Rossion, 2016) (z > 3.1; p < 0.001). The use of HFB signal, considered as a local signal (relative to higher SNR phase-locked low frequency signals, (Jacques et al., 2022)) correlated with local population spiking activity (Manning et al., 2009) as well as BOLD signal (Jacques et al., 2016b), was intended to ensure maximal signal independence between VOTC regions investigated. Potential dependencies due to electrical field propagation were further limited by the use of local bipolar referencing.

Face-selective activity can also be characterized in the time-domain, which allows visualizing periodic activity common to faces and nonface objects (every 0.167 s) and any significant deviation from this activity (i.e., face-selective activity; every 0.833 s; **Figure 1F**). In the time-domain, the face-selective signal can be isolated by filtering-out the 6 Hz signal

common to face and non-face objects (i.e., notch filter at 6 Hz and harmonics) (**Figure 1G**) (e.g., (Jacques et al., 2016a; Quek and Rossion, 2017; Retter and Rossion, 2016; Rossion et al., 2015)).

Figure 1. Recording and quantifying time-domain face-selective activity in the VOTC. **A.** SEEG (depth) electrode arrays (white circles) shown on the reconstructed white matter surface of one of the participants (ventral view of the left hemisphere). Electrodes penetrate both gyral and sulcal cortical tissues. **B.** The frequency-tagging paradigm to quantify face-selective neural activity: images of nonface objects appear at a rate of six stimuli per second (6 Hz) with variable face images presented every five stimuli (i.e., every 0.835s). Each stimulation sequence lasts for 65s (2s showed here). **C.** Representative examples of natural face images used in the study (actual images not shown for copyright reasons). **D**. Top: example raw intracranial EEG signal measured at the bipolar recording contacts shown in panel A (red). The signal is shown from -1.5 to 20 s relative to the onset of a stimulation sequence. The time-series displayed is an average of 2 sequences. Above the time-series, red vertical ticks indicate the appearance of face image every 0.835s and small black vertical ticks indicate the appearance of non-face objects every 0.167s. Middle: time by frequency representation of SEEG data in the HFB range (30-160 Hz). The plot shows the percent signal change at each frequency relative to a pre-stimulus baseline period (-1.6s to -0.3s). Periodic burst of HFB activity at the frequency of face stimulation (i.e., 1.2 Hz) are visible. Bottom: modulation of HFB amplitude over time obtained by averaging time-frequency signals across 30-160 Hz. **E.** HFB signal is transformed in the frequency domain to quantify face-selective amplitude as the sum of 12 face-selective frequency harmonics. **F.** Time-domain averaged HFB response to face images shows both the periodic response to non-face objects at 6 HZ (cycle duration = 0.167 s) and the larger face-selective response starting ~0.1 s after face onset. Shaded area around the

curve is standard error of the mean across face trials. **G.** Mean face responses from two separate example recording contacts (in PTL and ATL) in which the 6 Hz response to non-face objects has (black traces) or has not (blue traces) been filtered-out.

Visualization of time-courses in contacts with significant face-selective activity reveals that face-selectivity is manifested either as a periodically larger response to faces compared to nonface objects (**Figure 1F,G, Figure 2**) or more rarely as a periodically smaller response to faces (**Figure S4**). Since category-selectivity is usually defined as a larger response for a given category compared to control categories in electrophysiological and neuroimaging studies, in the current study we focus on those larger activity for faces. Nevertheless, we provide illustrations and timing analyses for contacts showing response decrease in supplemental material (**Figures S5 and S7**). We identified 444 VOTC recording contacts in the gray matter and medial temporal lobe that showed a significant HFB face-selective response increase in 101 participants (among 8278 bipolar contacts located in gray matter or medial temporal lobe in the VOTC of 140 participants; 4680 in the left hemisphere and 3598 in the right hemisphere, **Table S1**). The proportion of face-selective contacts was significantly larger in the right (6.5%, 234/3598) than in the left hemisphere (4.5%, 210/4680, $p < 0.001$, 2-tailed permutation test).

Face-selective contacts displayed in the Talairach space for group visualization and analyses (**Figure 3A**), indicate that HFB face-selective activity was distributed across the VOTC, from the occipital lobe (OCC) to the anterior temporal lobe (ATL, 95% of contacts located in [-92 to -5] TAL y coordinate, **Figure 3A**), although not reaching the temporal pole (TP). Face-selective activity was also measured in subcortical structures of the medial temporal lobe (MTL, amygdala- AMG). The bulk of the activity stretched from the IOG, through the FG and adjacent sulci (OTS and COS), up to the antFG+ (antFG and surrounding sulci: antOTS and antCOS). The highest number and proportion of face-selective contacts (**Figure 3B,C,** contacts labeled according to each participants individual anatomy as in (Jacques et al., 2022; Jonas et al., 2016a), **Figure S1**) were observed in bilateral latFG and antOTS as well as right IOG.

Figure 2: **Time-domain face-selective periodic increases.** Left column: Time by frequency response (percent signal change, psc; see scale value at the top of each plot and color scale at the bottom) in the HFB range (30-160 Hz) averaged over face-selective contacts in each main VOTC region (OCC: occipital, bottom; PTL: Posterior temporal lobe, middle; ATL: anterior temporal lobe, top). For each recording contact, the time-frequency data was segmented in epochs of about 3 face cycles (i.e. 3 x 0.833 s), averaged by contacts and then averaged over the three groups of contacts. Frequency axis is on the left. Green traces are the HFB amplitude envelope obtained by averaging over the 30-160 Hz range. Amplitude (psc) axis is on the right, show in green. Right column: Frequency spectra averaged over the corresponding groups of recording contacts and showing the face-selective response (red circles) at multiples of 1.2 Hz (i.e., face stimulation frequency) and visual response at 6 Hz (black square, other harmonics not shown). This highlights the sharp decrease in general visual response (i.e., 6Hz) relative to face-selective response from posterior to anterior VOTC.

Figure 3: Spatial organization and increase in abstraction of face-selective HFB activity in VOTC. **A**. Map of all VOTC recording contacts across the 140 individual brains displayed in the Talairach space in a transparent reconstructed cortical surface of the Colin27 brain (ventral view). Each circle represents a single recording contact. Face-selective contacts are color-coded according to their anatomical location in the original individual anatomy. White-filled circles correspond to contacts without significant face-selective activity. Values along the y-axis of the Talairach coordinate system (postero-anterior) are shown near the interhemispheric fissure. **B**. VOTC maps of the local proportion of face-selective contacts relative to the number of recorded contacts. Black contour lines delineate local proportions significantly above zero (*p* < 0.01, percentile bootstrap). **C**. The number of face-selective contact is show for each anatomical region (region defined in each individual participant) and hemisphere. **D**. Face-selectivity index (FSI) along the postero-anterior axis collapsed along the X dimension (medio-lateral). The shaded area shows the 99% confidence interval computed using a percentile bootstrap. **E**. Map of the proportion of face-exclusive (i.e., face-selective without significant response to nonface-objects at 6Hz and harmonics) relative to face-selective contacts across VOTC.

Increase in selectivity ('abstraction') from posterior to anterior VOTC

Quantifying separately the amplitude of the face-selective response (i.e., 1.2Hz and harmonics, excluding harmonics of 6 Hz) and the amplitude the general visual response (i.e., manifested for all presented images, at 6Hz and harmonics) for each contact allows computing an index of face-selectivity (face-selectivity index – FSI) that reflects the magnitude of the face-

selective response relative to the overall visual responsiveness of the cortex around the recording contact. The FSI varies from 0 (no face-selective response) to 1 (only face-selective response, no general visual response) and can also be thought of as a proxy for the level of abstraction exhibited by the local neural population (i.e., the degree to which the neural population represents faces independently of generic visual input). FSI increased from posterior VOTC to anterior VOTC: OCC = 0.72, 99% confidence interval: [0.65 0.79]; PTL = 0.88, [0.86 0.89]; ATL = 0.88, [0.86 0.9]; OCC vs. PTL: p < 0.0002, 2-tailed bootstrap test; PTL vs. ATL: p = 0.45; OCC vs. ATL: p < 0.0002. Along the same lines, computing the FSI as a function of the position along postero-anterior axis (Y Talairach dimension, collapsing contacts over both hemispheres, **Figure 3D**) confirms a strong increase in the FSI from posterior to anterior VOTC (from <0.4 to >0.9), most pronounced over the posterior half of the VOTC. Additionally, we also quantified the proportion of face-selective contacts that are 'face-exclusive' (i.e. face-selective contacts that do not display any significant response to non-face objects measured at 6 Hz and harmonics with z < 1.64; p > 0.05). Face-exclusive contacts composed 38% of all face-selective contacts (169/444) with a marked increase in proportion from OCC (18%), to PTL (26%) and ATL (51%), reaching ~80% in the more anterior parts of ATL (**Figure 3E**).

Concurrent face-selective activity across VOTC

To achieve the most comprehensive and robust possible overview of the timing of face-selective activity in VOTC, we started by examining the 3 main VOTC regions (i.e., OCC, PTL, ATL, excluding TP and MTL for which there were only few significant contacts), including all significant face-selective contacts showing a response increase (**Figure 3A**). Timing was characterized by 4 parameters: (1) latency of response onset and (2) offset, as well as (3) response overlap and (4) response correlation across VOTC regions (**Figure S2**). Analyses focus on comparing these parameters (within hemispheres) to specifically test the hierarchical processing hypothesis.

We first considered onset latencies of face-selective signals (**Figure 4**, see **Figure S3A** for non-normalized time-series, **Figure S4** and supplementary results for contacts exhibiting face-selective response decrease). These were overall similar across VOTC main regions, ranging from 81 ms (95% confidence interval: [65-97] ms) in the right PTL region to 112 ms ([100-122] ms) in the left PTL (**Figure 4**), with no clear pattern of inter-hemispheric or postero-anterior differences. Statistically comparing regions within hemispheres revealed no significant differences of onset latency (all p's > 0.27, two-tailed permutation test, fdr-corrected, **Figure**

4B). Second, the latency at which neural activity returned to baseline level (i.e., offset latency) tended to slightly increase from OCC to ATL, ranging from 354 ms ([290-415] ms) in left OCC to 532 ms ([454-620] ms) in right ATL (**Figure 4B**). However, due to a large variability across contacts within each region, these differences in offset latency were not significant (p's > 0.1, fdr-corrected), except when comparing OCC to ATL in the right hemisphere (393 ms, [349-425] ms vs. 532 ms ([454-620] ms for OCC and ATL respectively; p = 0.034, two-tailed permutation test). Collapsing contacts across hemisphere confirmed the significant increase in the offset latency from OCC (380 ms, [345-417] ms) to PTL (454 ms, [407-483] ms; OCC vs PTL, p = 0.02, two-tailed permutation test, fdr-corrected) and from PTL to ATL (522 ms, [497-618] ms; PTL vs. ATL., p = 0.0024). Together with the similar onset latencies across main VOTC regions, this indicates that overall, response duration increases as a function of postero-anterior VOTC location. Fourth, we examined the temporal overlap between two given regions by computing the percentage of the area under the response curve (AUC) of the overlap between region A and B (determined using onset and offset latencies) relative to the total AUC of region A (see methods, **Figure S2**).

Interestingly, despite the differences in offset latencies across VOTC, on average when comparing two regions, the AUC that falls within the temporal overlap between these regions represents 95% of the total AUC for each region, ranging from 87% (percentage of the right ATL AUC occupied by the AUC of its temporal overlap with the right OCC, see **Figure 4B**) to 100% (e.g., percentage of the left OCC AUC occupied by the AUC of its temporal overlap with the left ATL, **Figure 4B**). This indicates that, even in case of relatively large differences in offset latency (e.g., 354ms for left OCC and 495ms for left ATL), the portion of the left ATL AUC during non-overlapping window (e.g., between 354ms to 495ms for left OCC vs. ATL) is negligible (12%) compared to the AUC observed during the temporal overlap (91 ms to 354 ms, 88% of the left ATL AUC occupied by the AUC of its temporal overlap with the left OCC). Moreover, these observations indicate that the vast majority of the response in a region located anteriorly (e.g., ATL) to another region (e.g., PTL) takes place during the time-window of the posterior region.

Figure 4: **Concurrent face-selective activity across VOTC. A.** Mean time-domain face-selective HFB activity in each VOTC region (OCC, PTL, ATL) and hemisphere. HFB time-series were filtered to remove the general visual response at 6 Hz and harmonics so only face-selective signal remains. The maximum amplitude of each averaged waveform was normalized to 1 for visualization purposes only (see Figure S4 for non-normalized waveforms). Shaded area represents the standard error of the mean between contacts. Colored vertical lines indicate onset and offset latencies for each VOTC region with shaded horizontal bar around each line representing a 95% confidence interval (percentile bootstrap). **B**. Timing parameters split by VOTC region and hemisphere: onset (top left) and offset (top right) latency of face-selective response, correlation of time-series between regions (bottom left) and between-region area under the curve (AUC) overlap (bottom right). For the AUC overlap, the arrow shows the directionality of the computed overlap. For instance, the arrow from ATL to OCC indicates that we are describing the percentage of the total AUC of ATL (measured between onset and offset latencies) occupied by the AUC of its temporal overlap (also determined using onset and offset) with OCC. Acronyms: LH: left hemisphere, RH: right hemisphere.

Fourth, as an additional measure of temporal overlap we considered the correlations between the mean time-series recorded in each region, which were highly correlated (**Figure 4B**), ranging from Pearson's r=0.94 when comparing right OCC to right ATL, to r=0.99 when comparing right OCC to right PTL or left PTL to left ATL. This means that the time-series across

regions share from 88% to 98% of their variance. Importantly, statistically comparing the correlations computed between-regions to correlations computed *within* each region revealed no significant difference (there were only trends for differences when comparing OCC to ATL and PTL to ATL in the right hemisphere (within OCC and within ATL: r=0.99 and 0.98 respectively, between OCC and ATL: 0.94, *p* = 0.06, one-tailed percentile boostrap test, fdr-corrected; within PTL and within ATL: r=0.99 and 0.98 respectively, between PTL and ATL: 0.94, *p* = 0.06). This indicates that the time-domain activity measured in signal+ contacts across different VOTC regions are not significantly different from the correlations expected within signals from a given region itself.

In a separate analysis (**Figure S5**), we also fully replicate the above observations but focusing on timing properties in 3 main face-selective regions in VOTC: IOG, latFG and antFG+ (i.e., antFG, antOTS, antCOS).

In addition to the group analyses described above, we performed a within-participants comparison of onset latencies in participants that had face-selective contacts in at least 2 different main VOTC regions (e.g., OCC and PTL). While within-participants statistics allow disregarding between-participants variability, it was done here at the expense of sample size since only a small subset of participants have face-selective contacts located in multiple face-selective regions. Onset latencies were computed for individual recording contacts and averaged by region within participants. These comparison did not reveal significant differences in onset latencies between main VOTC regions collapsed across hemispheres (OCC minus PTL: N=13, mean difference = -8 ms, 95% confidence interval: [-14 14]ms, *p*=0.27, 2-tailed permutation test; PTL minus ATL, N=19, mean difference = +14 ms, 95% CI: [-20 20]ms, *p*=0.18; OCC minus ATL, N=9, mean difference = +17 ms, 95% CI: [-27 27]ms, *p*=0.24) or between VOTC main face-selective regions (IOG minus latFG: N=9, mean difference = 0 ms, 95% CI: [-12 12]ms, *p*=0.88, 2-tailed permutation test; latFG minus antFG+, N=19, mean difference = +15 ms, 95% CI: [-20 20]ms, *p*=0.14; OCC minus ATL, N=7, mean difference = +8 ms, 95% CI: [-24 24]ms, *p*=0.52).

Concurrent functional connectivity between face-selective regions

Last, we estimated the functional connectivity between pairs of main face-selective regions (IOG, latFG, antFG+) by correlating single-trial face-selective amplitude measured in different VOTC regions (IOG, latFG, antFG+), across time. Our use of local bipolar referencing of electrophysiological signals ensures that responses in each region are maximally local and that

any significant correlation between regions cannot be attributed to the reference itself. Directionality in connectivity (e.g. connectivity from IOG to latFG or the reverse) was estimated by correlating face-selective amplitude measured at different time-lags (-150 to 150ms) between the compared regions (Kadipasaoglu et al., 2017). The properties of the functional connectivity patterns were largely similar across pairs or regions (**Figure 5**), although correlations were weaker between IOG and antFG (**Figure 5C, F**; especially in the left hemisphere where only 1 participant was included). Overall, correlations were present before response onset, manifested by weaker scattered yet largely symmetrical small correlation clusters. Stronger, more consistent and clustered between-region functional connectivity appeared at response onset, i.e. around 80-100ms, and was maximal at response peak between 150 and 200 ms (see Figure 4A), remaining significant for the remaining of the response. Most importantly, correlations measured after response onset were consistently centered on the diagonal (0 ms lag), further supporting concurrent activity of the VOTC regions and suggesting that these regions receive common correlated concurrent inputs possibly from lower-level visual cortex. Despite correlations being centered on the diagonal, correlation coefficients symmetrically spread around the diagonal, resulting either from reciprocal connections or from short (<30-40ms, likely due to temporal smoothing caused by the time-frequency analyses) or longer (up to 150ms) duration of temporal auto-correlation of the electrophysiological signal. These broader/longer correlations around the diagonal, which were most pronounced between left IOG-latFG and left/right latFG-antFG (Figure 5B, D-E), appeared after 200 ms and were likely related to sustained and stable face-selective neural responses at these latencies which are correlated in amplitude across regions.

Figure 5: Concurrent functional connectivity between face-selective regions. (A) Group-averaged Pearson's correlations between single-trials face-selective amplitude measured at left IOG and left latFG computed across time, representing functional connectivity between these 2 regions. Black contour lines indicate significant positive correlations ($p < 0.01$, fdr-corrected). Correlations were computed across -150 to 150 ms lags between regions to infer direction of connectivity. The black dashed diagonal line represents a 0 ms time-lag between regions. Correlations centered above the diagonal would indicate that face-selective activity in latFG correlates with but precedes activity in IOG, suggesting information flow from latFG to IOG, and the reverse for correlations centered below the diagonal. B-F: functional connectivity between pairs of face-selective regions. Same conventions as in panel A.

Mapping face-selective onset latency in Talairach space

Next, we aimed to explore the onset latencies of face-selectivity HFB signal at a more local scale, taking advantage of the dense sampling of the VOTC to compute maps in the Talairach space. Due to the lower sample size in any given local measurement compared to when using larger anatomically-defined regions, and to limit noise in the latency estimates, we did not take into account contacts with low split-half correlation (SHC: $r < 0.4$; removing contacts with SHC < 0.4 had no effect when comparing timing properties across main VOTC regions). Moreover, we only included local VOTC volumes (i.e., 'voxels') with more than 5 contacts. For contacts showing face-selective response increase (contacts with response decrease reported in **Figure S6**), face-selective response latency was lowest in regions exhibiting

the most consistent face-selectivity: right IOG (right IOG: mean = 100 ms, range = [83 - 118ms], **Figure 6A**) and laFG (mean = 95 ms, range = [87 - 110ms]), as well as left latFG (mean = 103 ms, range: [99 - 112ms]), and the anterior portion of the right antFG+ area (mean = 113 ms, range: [95 - 132ms]). Between and around those regions, latency was slightly higher (e.g.: posterior FG: mean = 123 ms, range = [102 - 137ms]), especially in the antMTG region (mean = 148 ms, range = [132 - 172ms]). Examining the postero-anterior profile of response latency (**Figure 6B**) reveals relatively constant response latencies (90-110ms) in the OCC (Talairach Y coordinates: - 90 to -65 mm]) and PTL regions (Talairach Y coordinates: -65 to -40 mm) up to the ATL region (Y > -40 mm) where the response latency increased, especially in the right hemisphere (i.e., 20-40 ms increase). Face-selective response onset latency in the right ATL   (Y Talairach coordinates - 30.5 to -27.5 and -18.5 to -15.5) was significantly above what would be expected by chance ($p <$ 0.05, 2-tailed, fdr-corrected) if the antero-posterior position had no influence on face-selective response onset latency   (**Figure 6B**, randomization test where the pairing between each recording contact location and its time-domain response was shuffled). Since the signal-to-noise ratio (SNR) and associated split-half reliability tends to be lower in the ATL (Jacques et al., 2022) and the estimated onset latency tends to increase at lower SNR, we computed the same analyses keeping only contacts with a high split-half correlation (SHC) above r = 0.7 (**Figure 6B, Figure S7**). As expected, due to the exclusion of lower SNR contacts, the estimated mean latency in the [-27.5 to -12.5 mm] Y coordinate interval was reduced by about 10-20 ms (compared to when using a threshold of SHC > 0.4) and was no longer significantly above chance. Additionally, given the relatively large distance between the ATL and posterior occipital regions (>50 mm), the slight increase in response onset latency from posterior to anterior VOTC can largely be accounted for by the predicted increase in response latency due to simple conduction delay (likely between 1.5 and 5 m/s (van Blooijs et al., 2023), **Figure 6B**), even with direct cortico-cortical connections from earlier visual cortical areas.

Figure 6: **Mapping concurrent face-selective response onset latency in VOTC. A**. Face-selective onset latency map across VOTC. **B.** Variation of face-selective response latency along the postero-anterior axis. Each data point represents the onset latency measured from the time-series averaged over contacts collapsed across the medio-lateral X dimension in each hemisphere within 20 mm segments (in the Y dimension). Thick lines are estimated onset latencies and shaded areas show the 99% confidence intervals expected under the null hypothesis that the postero-anterior location has no influence on the onset latency. Dark blue (full lines) is used for latencies computed when excluding contacts with SHC < 0.4 (as in panel A) and gray (dashed lines) are for latencies computed when excluding contacts with SHC < 0.7. Green line and shaded area show the expected increase in response onset latency based on simple conduction delays (van Blooijs et al., 2023) due to increasing distance from the occipital region, with reference to the latency averaged over the 2 most posterior face-selective response measured here. The dotted line is the mean expected conduction velocity for direct cortico-cortical connections (~3.2 m/s) and the shaded area represents the minimum (1.7 m/s) and maximum (5 m/s) expected conduction velocity (i.e., based on measurements from other brain regions (van Blooijs et al., 2023)).

Discussion

Here we provide a large-scale mapping of the temporal properties of neural activity supporting visual recognition in the human VOTC with electrophysiological intracerebral recording. Faces are used as the key target category for visual recognition not only because of their highly significant social value for humans, but also due to their wide distribution of representation across the (bilateral) VOTC, allowing to compare the time-course of category-selectivity across multiple regions. In addition, the cortical network of face-selective regions is the most extensively studied within the occipital and temporal lobes and is thought to reflect the manifestation of a canonical set of operations that reveal general principles of how primate visual recognition works (Conway, 2018; DiCarlo et al., 2012).

We report two major findings. First, a progressive increase in face-selectivity along the postero-anterior axis. Second, a concurrent face-selective neural activity, with similar onset latencies (80-100ms) and largely overlapping and correlated time courses across the whole VOTC, spanning around 90 mm of cortical territory along the postero-anterior axis. Our

observations suggest that while the human VOTC may apparently be functionally organized in a hierarchy of regions with increasingly more abstract representations, these regions of the association cortex do not appear to be activated sequentially (or even in cascade) but rather concurrently.

1. Time course of VOTC face-selective activity and speed of face categorization

The onset of category-selectivity for faces identified here in humans at around 80-100 ms at the group level is early, supporting an initial fast feedforward sweep of activity for rapid face categorization (Cauchoix et al., 2014; Crouzet and Thorpe, 2010; Retter et al., 2020; Serre et al., 2007). This onset latency, with a peak of activity reached at about 200 ms, is in line with multiple sources of evidence (speed of gaze to lateralized natural images of faces (Crouzet and Thorpe, 2010); scalp EEG studies (Retter and Rossion, 2016; Rossion and Caharel, 2011; Rousselet et al., 2008) ; intracranial EEG studies (Allison et al., 1999; Jacques et al., 2016b; Kadipasaoglu et al., 2017; Liu et al., 2009); single neuron activity in human face-selective regions (Laurent et al., 2025; Quian Quiroga et al., 2023)). Here we extend these findings by providing comprehensive measurements of response onset latencies across the whole VOTC. Importantly, we measure the onset of face-*selectivity*, not the absolute response onset latency to face images that could be driven by unspecific lower-level features (Cao et al., 2025; Schrouff et al., 2020). Investigating category-selective activity is critical because it may be temporally dissociated from general visual responses in the same brain areas (Jiang et al., 2011) and is directly related to visual recognition function (Jonas and Rossion, 2021; Rangarajan et al., 2014; Volfart et al., 2022).

Beyond onset latencies, we also report response durations that are in line with values obtained from the scalp (EEG) with the same paradigm (about 420ms (Retter and Rossion, 2016; Rossion et al., 2015)). Despite a slight increase in duration of the sustained low amplitude part of the time-series along postero-anterior VOTC axis, the overwhelming majority of signal amplitude (87 to 100%) and variability (88 to 98%) occurs concurrently, i.e., it temporally overlaps across brain regions. Given the timing similarity between the present intracerebral data and scalp EEG data obtained in neurotypical individuals with the same approach (Retter and Rossion, 2016; Rossion et al., 2015), there appear to be no delay potentially attributable to the specific population tested here, i.e., patients with intractable refractory epilepsy (Allison et al., 1999). While we cannot exclude overall lower response amplitudes of face-selective activity

in this specific population relative to normal individuals and/or potential shifts of hemispheric lateralization at the individual level depending on hemispheric seizure localization, this would not affect our conclusions. Finally, whereas face-selectivity is measured here during temporally fixed face stimulation (every 833 ms), observers perform an orthogonal task, are unaware of the periodicity of face stimulation and unable to determine it due to the rapid base stimulation rate. Most importantly, both the amplitude and time course of face-selective activity are unaffected by the temporal distance between face stimuli/ratio of faces *vs.* objects (Retter & Rossion, 2016), or the temporal periodicity or predictability of face presentation (Quek and Rossion, 2017).

2. A progressive posterior to anterior gradient of category-selective abstraction in the VOTC

Our findings indicate a progressive increase in category-selectivity from posterior to anterior VOTC. More specifically, while category-selective populations of neurons in posterior VOTC regions tend to respond to all visual stimuli, as evidenced by high 6 Hz amplitude, this common visual response gradually decreases along the postero-anterior axis (**Figures 2, 3D**), leading to a high proportion of exclusive response to faces in the most anterior regions of the ventral temporal lobe (**Figure 3E**). The postero-anterior increase in face-selectivity is only partially in line with fMRI observations that have reported both larger face-selectivity in the fusiform gyrus ('Fusiform Face Area', FFA) than in the posteriorly located inferior occipital gyrus (IOG; 'Occipital Face Area', OFA) (Tsao et al., 2008; Weiner and Grill-Spector, 2010) , as well as the opposite finding (Rossion et al., 2012) (see also recent evidence (Chen et al., 2023) of larger face-selectivity in the posterior than middle fusiform gyrus), potentially depending on the general size/amplitude of activity in a given ROI and how selectivity indexes are computed. Most importantly, fMRI studies are limited to comprehensively explore the VOTC due to the large magnetic susceptibility artifacts (low SNR) in the ventral ATL (Ojemann et al., 1997; Rossion et al., 2024). To the best of our knowledge, previous intracranial studies have not described such a comprehensive pattern of variation in face-selectivity across the whole VOTC, due to limited sampling (spatially or limited to the gyral surface), lack of mapping and/or face-selective response quantification, or focus on medio-lateral variations of selectivity (Allison et al., 1999; Engell and McCarthy, 2014; Jacques et al., 2016b; Jonas et al., 2016a; Kadipasaoglu et al., 2017, 2016; Rangarajan et al., 2014; Schrouff et al., 2020; Vidal et al., 2010).

This increase in selectivity for faces, as well as the increase in the proportion of face-exclusive activity in the postero-anterior VOTC axis, cannot be attributed to a mere reduced ability of anterior regions to generate responses at a fast presentation rate (6 Hz) (Jonas et al., 2016a). Rather, these observations suggest an increase in 'abstraction' of visual neural representations, with a large proportion of neuronal populations in anterior VOTC regions exhibiting similar activity to different faces images independently of the context in which they appear, while being much less responsive to other complex visual stimulation, or even responding only to faces. These findings are compatible with standard hierarchical models of visual processing, such as proposed originally by Hubel and Wiesel (1962) (Hubel and Wiesel, 1962) and extended to a hierarchy of stages for building increasingly complex invariant visual object or face representations (Conway, 2018; DiCarlo et al., 2012; Duchaine and Yovel, 2015; Fairhall and Ishai, 2007; Freiwald, 2020; Grill-Spector et al., 2017; Issa et al., 2018; Marr, 1982; Riesenhuber and Poggio, 1999; Tsao, 2014; Van Essen et al., 1992), as well as with the view that this hierarchy is implemented anatomically along the postero-anterior axis of the VOTC.

3. Concurrent face-selectivity across VOTC: evidence for non-hierarchical organization of visual association cortex

While the increase in category-selectivity from posterior to anterior VOTC as described above is compatible with a hierarchical model of visual recognition, the remarkably similar time-course of face-selective visual neural activity across VOTC regions is difficult to reconcile with the temporal properties implied by such model. In the classical hierarchical view of visual processing (Conway, 2018; DiCarlo et al., 2012; Duchaine and Yovel, 2015; Fairhall and Ishai, 2007; Freiwald, 2020; Grill-Spector et al., 2017; Issa et al., 2018; Marr, 1982; Riesenhuber and Poggio, 1999; Tsao, 2014; Van Essen et al., 1992), representations at a given level of the hierarchy are built from combination of inputs received from a lower level, implying serial ordering and thus a time difference between successive levels (Bullier, 2001; Bullier and Nowak, 1995; DiCarlo et al., 2012; Schmolesky et al., 1998; Thorpe and Fabre-Thorpe, 2001).

Hierarchical models do not imply that a process should be completed at a given stage before the next one is initiated. Yet, even in a hierarchical cascade of cortical processes and representations (DiCarlo et al., 2012), and regardless of acknowledged feedback/reentrant loops (Edelman and Gally, 2013; Issa et al., 2018; Kar et al., 2019; Lamme and Roelfsema, 2000) there should be delays in onset times, and only partial temporal overlap between increasing

processing level of the hierarchy (Bullier and Nowak, 1995; Issa et al., 2018; Schmolesky et al., 1998). Based on findings from electrophysiological as well as cortico-cortical evoked potential (CCEP) measurements, the delay between each level should be at least 10-30ms depending on (multi)synaptic transmission delays and conduction delays in short- or long-range connections (Conner et al., 2011; Nowak and Bullier, 1997; Schmolesky et al., 1998; van Blooijs et al., 2023). Importantly, this standard view is also how human, and non-human primate, cortical recognition of faces is generally conceived (Conway, 2018; Duchaine and Yovel, 2015; Fairhall and Ishai, 2007; Grill-Spector et al., 2017; Issa et al., 2018; Schweinberger and Neumann, 2016; Tsao, 2014), with 50-100 ms time difference postulated between stages (Sadeh et al., 2010; Schweinberger and Neumann, 2016).

In contrast to this hierarchical view, the present findings of concurrent face-selective activity across the human VOTC support a non-hierarchical organization of this recognition process. More specifically, our observations suggest that, rather than being successively activated along a posterior-axis, multiple face-selective populations of neurons spread across the whole VOTC receive direct and parallel sensory inputs from low-level visual areas (e.g., V1), through so-called bypass pathways. Such pathways have long been described in anatomical studies of the visual system of macaque monkeys (Conway, 2018; Van Essen and Maunsell, 1983; Zeki and Shipp, 1988). In humans, specifically for face recognition, both lesion studies and time-resolved fMRI investigations have provided indirect evidence for bypass cortical pathways, reporting face-selective activity in the midFusiform gyrus (i.e. 'Fusiform face Area', FFA) in the absence of (lesion (Rossion et al., 2003; Steeves et al., 2006; Weiner et al., 2016)), or before (Jiang et al., 2011), any category-selective activity in the posteriorly located IOG. Moreover, while an intracranial recording study with a limited sample focusing largely on occipital and posterior temporal face-selective regions has provided mixed outcomes regarding the onset of selective response to faces (delayed onset in the left latFG relative to IOG in n=4; no delay in the right hemisphere n=3), CCEP in the same study (Kadipasaoglu et al., 2017) also suggested independent and parallel signal propagations between early visual areas and both face-selective regions (i.e. latFG/FFA and IOG). Critically, this study did not report face-selective onset latencies in the ATL, where ~200 face-selective contacts were measured in the current study. Further evidence from Diffusion Tensor Imaging (DTI), suggest independent direct connections from early visual cortex to face-selective regions in the IOG and post/mid-fusiform gyrus (Bryant et al., 2019; Finzi et al., 2021; Kim et al., 2006; Wang et al., 2020; Weiner et al., 2016).

More generally, human V1 even appears to show (hominoid-specific) connectivity with lateral, inferior, and anterior temporal regions, beyond the retinotopically organized cortical areas (Bryant et al., 2019).

While these observations provide an anatomical basis for concurrent onset times in face-selective regions of the human VOTC, the present electrophysiological data, collected from a particularly large sample and with an objective face-selective measure, go beyond by showing for the first time that (1) even the most anterior temporal population of neurons (ATL) respond selectively to faces as early as posterior VOTC regions and (2) that all of these regions' time-courses overlap in time remarkably. The current findings also support direct connections between early visual cortex and face-selective regions around the anterior fusiform gyrus, anterior OTS and anterior COS (antFG+) in the ATL (likely via the same fiber bundles as for posterior face-selective regions: the inferior longitudinal fasciculus and inferior fronto-occipital fasciculus) where DTI measurements are limited by magnetic susceptibility artifacts. Importantly, while we report a slight overall increase in response onset latency in the ATL compared to more posterior regions, this increase can be accounted for by the predicted conduction delay given the large distance of the ATL from early visual areas in the posterior occipital cortex and the likely reliance on long range white matter fibers (~1.5 to 5 m/s or mm/ms (van Blooijs et al., 2023), **Figure 6B**). This further supports the view that there is no need to postulate mandatory intermediate hierarchical stage/relays between posterior and anterior regions. A number of factors can affect transmission speed, as determined using CCEP in humans, among which the number of fibers connecting different regions (Conner et al., 2011) and which is likely to vary between face-selective regions (Finzi et al., 2021; Wang et al., 2020). Together with concurrent onset latencies, the large overlap in time-courses suggests parallel category-selective processing, likely to be initiated independently by sensory inputs from low-level visual cortex, as well as ongoing reentrant interactions (i.e., dynamic recursive/recurrent exchange of signals (Edelman and Gally, 2013; Kar et al., 2019; Lamme and Roelfsema, 2000)) between these regions for ~300ms (Issa et al., 2018; Kadipasaoglu et al., 2017). Moreover, our finding of differences across face-selective regions in the duration of the low amplitude sustained part of the response may reflect different patterns of connectivity between these regions and other (sub)cortical regions to which they are connected to. In particular, the longest duration in the ATL may potentially be linked to the specific connectivity of this region

with medial temporal lobe and other structures involved in semantic or episodic memory formation or retrieval (Persichetti et al., 2021).

4. Implications for models of visual recognition

How would sensory inputs be recognized as faces in such a non-hierarchical human system? In a nutshell, through synaptic sensory inputs originating from low-level visual cortex (e.g., V1) successfully triggering (post-synaptic) activity of sufficiently large face-selective populations of neurons in parallel throughout the VOTC. Despite temporal overlap, and concurrent process, one should acknowledge that there are genuine differences in degree of selectivity, as disclosed here, as well as receptive fields sizes, foveal bias and ipsilateral sensitivity across face-selective regions of the human VOTC (Finzi et al., 2021; Kay et al., 2015). These functional differences between regions provide a substantial advantage for visual recognition since it increases the probability of variable incoming sensory inputs from early visual cortices to trigger face-selective populations of neurons distributed along the VOTC. Importantly, in such a non-hierarchical system, there is no requirement for normalization processes across definite successive stages to build 'invariant' representations from inputs varying in size, orientation, lighting, etc. Instead, through hebbian learning mechanisms, views that are more commonly experienced (e.g., full-front faces at conversational distances) would be represented by larger populations of neurons than rarely seen views, providing a faster accumulation of category-selective neural activity (i.e., evidence accumulation) in this network and a more efficient recognition function for sensory inputs corresponding to these views (Perrett et al., 1998).

To be clear, such a non-hierarchical organization of the human ventral cortical face network corresponds fully to a feedforward/bottom-up view of visual recognition (at least for recognition of relatively clear views of faces) enriched by concurrent reentrant exchanges of signals within VOTC regions for a few hundreds of milliseconds. That is, putative 'top-down' signals from population of neurons in parietal or prefrontal cortices (Bar, 2003; Kar and DiCarlo, 2021) are not necessary for fast automatic recognition of faces. Yet, this simple view rests on a conceptual reinterpretation of functional organization of the human VOTC: rather than constituting a series of processing stages with increasing levels of complexity to derive visual representations (i.e., build the most veridical images of the world), face-selective populations of neurons are conceptualized as memory representations ('cortical memories' (Fuster, 1995))

distributed throughout the VOTC. That is, these populations of neurons, constrained by cytoarchitecture-specific white matter connections from birth (Kubota et al., 2025), have learned (from early developmental stages (Kosakowski et al., 2022)) through Hebbian mechanisms to discharge selectively to faces, with temporal synchrony strengthening their connections. Recognition, which 'simply' involves successful – (often) concurrent - matching to these cortical memories in the human association cortex is achieved in a bottom-up fashion, with the 'up' component being critical in recognizing sometimes ambiguous or degraded sensory inputs as faces (Cavanagh, 1991).

Summary and conclusions

In summary, our large-scale human intracerebral investigation of category-selective activity in human VOTC reveals that while there is an increase in selectivity/abstraction in face-selective activity from posterior to anterior VOTC, in line with a hierarchical view of visual recognition, response timing indicate that category-selective neural processes occur largely concurrently across the whole VOTC. These findings suggest that visual recognition is supported by category-selective neural populations spread across VOTC receiving direct concurrent inputs from early visual cortex, without the need to rely on intermediate and successive hierarchical relays.

MATERIALS AND METHODS

Participants

The study included 140 participants (71 females, mean age: 33.0±9.2 years; 123 right-handed, 3 ambidextrous) undergoing clinical intracerebral evaluation with depth electrodes (stereotactic electroencephalography or SEEG**, Figure 1A**) for refractory partial epilepsy. Participants were included in the study if they had at least one intracerebral electrode implanted in the ventral occipito-temporal cortex. All participants gave written consent to participate to the study, which was approved by a national human investigation committee certified by the French Ministry of Health (Institutional Review Board: IORG0009855)

Fast periodic visual stimulation paradigm

A well validated fast periodic visual stimulation (FPVS) paradigm with natural images was used to elicit face-selective neural activity in iEEG with high signal to-noise ratio (SNR) (see

(Rossion et al., 2015) for the original description of the paradigm in EEG; see (Jonas et al., 2016a; Rossion et al., 2018) for its validity in iEEG).

Stimuli. Two hundred grayscale natural images of various non-face objects (from 14 non-face categories: cats, dogs, horses, birds, flowers, fruits, vegetables, houseplants, phones, chairs, cameras, dishes, guitars, lamps) and 50 grayscale natural images of faces were used (Jacques et al., 2022; Jonas et al., 2016a; Rossion et al., 2015). Each image contained an unsegmented object or face near the center, these stimuli differing in terms of size, viewpoint, lighting conditions and background. Images were equalized for mean pixel luminance and contrast, but low-level visual cues associated with the faces and visual objects remained highly variable, naturally eliminating the systematic contribution of low-level visual cues to the recorded face-selective neural activity (Gao et al., 2018; Rossion et al., 2015).

Procedure. Participants viewed continuous sequences of natural images of objects presented at a fast rate of 6 Hz (i.e., stimulus onset asynchrony of 167ms) through sinusoidal contrast modulation. This relatively fast rate allows only one fixation per stimulus and is largely sufficient to elicit maximal face-selective activity (Retter and Rossion, 2016). Images of faces appear periodically as every 5th stimulus, so that neural activity that is common to faces and nonface stimuli is manifested at 6 Hz and harmonics, while differential (i.e., selective) reliable activity to faces are expressed at 1.2 Hz (i.e., 6 Hz/5) (see **Figure 1B,C**). All images were randomly selected from their respective categories (face and nonface), with the constrain that no image could be immediately repeated. A stimulation sequence lasted 70 s: 66 s of stimulation at full-contrast flanked by 2 s of fade-in and fade-out, where contrast gradually increased or decreased, respectively. During a sequence, participants were instructed to fixate a small black cross which was presented continuously at the center of the stimuli and to detect brief (500 ms) color-changes (black to red) of this fixation-cross. Among the 140 participants, participants viewed either 2 sequences (69 participants), 3 sequences (6 participants), 4 sequences (55 participants), 5 sequences (1 participant), 6 sequences (3 participants), 8 or more sequences (6 participants). No participant had seizures in the 2 hours preceding the recordings.

Intracerebral electrode implantation and SEEG recording

Intracerebral electrodes (Dixi Medical, Besançon, France) were stereotactically implanted within the participants' brains for clinical purposes, i.e., to delineate their seizure onset zones (Talairach and Bancaud, 1973) and to functionally map the surrounding cortex for

potential epilepsy surgery. Each 0.8 mm diameter intracerebral electrode contains 5-15 independent recording contacts of 2 mm in length separated by 1.5 mm from edge to edge (**Figure 1A**). A total of 997 electrode arrays were implanted in the VOTC of the 140 participants. These electrodes contained 11121 individual recording contacts in the VOTC (i.e., in the gray/white matter or medial temporal lobe-MTL; 6295 and 4826 contacts in the left and right hemisphere respectively). Intracerebral EEG was sampled at either 500 or 512 Hz and referenced to either a midline prefrontal scalp electrode or an intracerebral contact in the white matter. SEEG signal was re-referenced offline to bipolar reference to limit dependencies between neighboring contacts (Hagen et al., 2025). Specifically, the signal at a given recording contact was computed as the signal measured at that contact (i.e., with the recording reference) minus the signal at the directly adjacent contact located more medially on the same SEEG electrode array. Since SEEG field potentials are computed using pairs of adjacent contacts, each electrode array contains 1 contact less than in the original recording. All subsequent analyses were performed on bipolar-referenced signal in the set of bipolar contacts as described just above.

Contact localization in the individual anatomy. The position of each contact relative to brain anatomy was determined in each participant's own brain by coregistration of the post-operative CT-scan with a T1-weighted MRI of the patient's head. Anatomical labels of bipolar contacts were determined using the anatomical location of the 'active' contact. In cases where the active contact was in the white matter and the 'reference' contact was in the gray matter, the active contact was labeled according to the anatomical location of the reference contact. Bipolar contacts in which both the active and reference contacts were in the white matter were excluded from analyses. To accurately assign an anatomical label to each contact, we used the same topographic parcellation of the VOTC as in (Jacques et al., 2022; Jonas et al., 2016b) (**Table S1, Figure S1**).

SEEG signal processing and analyses

High frequency broadband (HFB) preprocessing. Segments of iEEG corresponding to FPVS stimulation sequences were extracted (74-second segments, -2s to +72s, **Figure 1D**, top) and notched-filtered to remove 50Hz line noise and 2 harmonics (100 and 150Hz). Variation in signal amplitude as a function of time and frequency was estimated by a Morlet wavelet transform applied on each SEEG 74-second segment from frequencies of 30 to 160 Hz, in 2 Hz

increments (**Figure 1D**, middle). The number of cycles (i.e., central frequency) of the wavelet was adapted as a function of frequency from 2 cycles at the lowest frequency to 9 cycles at the highest frequency. The temporal smoothing resulting from the wavelet transform was minimal: wavelet of 20 ms of full width at half maximum (FWHM) across the frequency range (i.e. median of FWHM computed at each frequency bin), ensuring that timing information is accurate up to 10 ms which corresponds to half of the FWHM. The wavelet transform was computed on each time-sample and the resulting amplitude envelope was downsampled by a factor of 6 (i.e., to a 85.3 Hz sampling rate) to save disk space and computation time. However, for timing analyses on face-selective contacts, the original recording sampling rate (500 or 512 Hz) was used to preserve temporal resolution. Resulting time-by-frequency SEEG segment were normalized to obtain, for each frequency bin, the percentage of signal change (PSC) generated by the stimulus onset relative to the mean amplitude in a pre-stimulus time-window (-1600 ms to -300 ms relative to the onset of the stimulation sequence). The PSC was then averaged across frequencies (between 30 Hz and 160 Hz) to obtain 74-seconds segments of time-varying HFB amplitude envelope (**Figure 1D**, bottom).

HFB frequency-domain processing. For each recording contact, 74s segments (i.e., 2-8 per participant) of HFB PSC envelope were averaged in the time-domain and cropped to contain an integer number of 1.2 Hz cycles beginning 2 seconds after the onset of the FPVS sequence (right at the end of the fade-in period) until approximately 68 seconds, before stimulus fade-out (79 face cycles ≈ 65.8 s). An FFT was then applied to the resulting cropped HFB envelope to obtain the amplitude spectrum in the frequency domain (**Figure 1E**). The frequency-tagging approach used here allows identifying and separating two types of neural activity: (1) a general visual response occurring at the base stimulation frequency (6 Hz) and its harmonics, as well as (2) a face-selective activity at 1.2 Hz and its harmonics (Jacques et al., 2022; Jonas et al., 2016a; Rossion et al., 2018). Face-selective activity significantly above noise level at the face stimulation frequency (1.2 Hz) and its harmonics (2.4, 3.6 Hz, etc.) were determined as follows (Hagen et al., 2025; Jacques et al., 2022): (1) the FFT spectrum was cut into 4 segments centered at the face frequency and harmonics, from the 1st until the 4th (1.2 Hz until 4.8 Hz), and surrounded by 25 neighboring bins on each side; (2) the amplitude values in these 4 segments of FFT spectra were summed; (3) the summed FFT spectrum was transformed into a Z-score. Z-scores were computed as the difference between the amplitude at the face frequency bin and the mean amplitude of surrounding bins divided by the standard deviation of

amplitudes in the surrounding bins. A contact was considered as showing a face-selective response in HFB if the Z-score at the frequency bin of face stimulation exceeded 3.1 (i.e., *p* < 0.001 one-tailed: signal>noise).

HFB time-domain preprocessing. Recording contacts with a significant face-selective response (based on frequency-domain analyses) were further processed in the time-domain. Starting from the 74-seconds HFB amplitude segments (see 'HFB preprocessing'), time-series were processed in the following way: (1) an FFT notch filter (filter width = 0.07Hz) was applied to remove the general visual response at 6Hz and 3 additional harmonics (i.e. 6, 12, 18, 24 Hz); (2) time-series were segmented in 1.17 s epochs centered on the onset of each face (i.e. [-2 to + 5] 6Hz-cycles relative to face onset) in the FPVS sequences; (3) resulting epochs were averaged (see **Figure 1F** for an example averaged time-course without the notch filtering of the general visual response). Unless noted otherwise, averaged time-series per contact were baseline-corrected by subtracting the mean amplitude in a [-0.166 to 0 s] time-window relative to face onset. Considering time-series for each contact revealed that face-selective activity manifest either as a periodically *larger* response to faces compared to non-face objects, or as a periodically *smaller* response to faces (**Figure S4**). Response increase/decrease was defined as a function of whether the amplitude in the 0.15s to 0.35s time-window was respectively larger or smaller than the amplitude in the time-window just preceding the onset of the face (1 cycle at 6Hz, i.e. [-0.167s to 0s]). During the FPVS sequences, images are presented using a sinusoidal modulation of contrast, rather than an abrupt onset (**Figure 1A**). Hence, to account for the delay with which the images become visible/perceivable in this setting, the face onset time was shifted forward by 33 ms, as established from a direct comparison of sine wave to square wave/abrupt stimulus presentation in a full EEG study (Retter and Rossion, 2016) and a subset of participants of the present study. This corresponds to 4 frames at 120Hz screen refresh rate and to a face contrast of about 35% of the maximal (Retter and Rossion, 2016).

Group visualization and analyses in Talairach space. For group mapping and visualization, anatomical MRIs were spatially normalized to determine the Talairach (TAL) coordinates of VOTC intracerebral contacts. The cortical surface used to display group contact locations and maps was obtained from segmenting the Collin27 brain from AFNI (Cox, 1996), which is aligned to the TAL space. We used TAL transformed coordinates to compute maps of the local proportion of face-selective intracerebral contacts across the VOTC. Local proportion of contacts was computed in volumes (i.e., 'voxels') of size 15 x 15 x 100 mm (respectively for the X: left –

right, Y: posterior – anterior, and Z: inferior – superior dimensions) by steps of 3 x 3 x 100 mm over the whole VOTC. A large voxel size in the Z dimension enabled collapsing across contacts along the inferior-superior dimension. For each voxel, we extracted the following information across all participants in our sample: (1) number of recorded contacts located within the voxel across all participants; (2) number of significant face-selective contacts. From these values, for each voxel we computed the proportion of face-selective contacts as the number of significant contacts within the voxel divided by the total number of recorded contacts in that voxel. Then, for each voxel we determined whether the proportion of significant contacts was significantly above zero using a percentile bootstrap procedure, as follows: (1) within each voxel, sample as many contacts as the number of recorded contacts, with replacement; (2) for this bootstrap sample, determine the proportion of significant contacts and store this value; (3) repeat steps (1) and (2) 2,000 times to generate a distribution of bootstrap proportions; and (4) estimate the *p*-value as the fraction of bootstrap proportions equal to zero.

Quantification of response amplitudes. Amplitude quantification was performed on face-selective contacts. We first computed baseline-subtracted amplitudes in the frequency domain as the difference between the amplitude at each frequency bin and the average of 48 corresponding surrounding bins (up to 25 bins on each side, i.e., 50 bins, excluding the 2 bins directly adjacent to the bin of interest, i.e., 48 bins). Then, for each contact, face-selective amplitude was quantified as the sum of the baseline-subtracted amplitudes at the face frequency from the 1st until the 14th harmonic (1.2 Hz until 16.8 Hz), excluding the 5th and 10th harmonics (6 Hz and 12 Hz) that coincided with the base frequency (Jonas et al., 2016a). General visual response amplitude was quantified separately as the sum of the baseline-subtracted amplitudes at the base frequency from the 1st until the 3rd harmonic (6 Hz until 18 Hz).

Face-selectivity index. We computed an index of face-selectivity for face-selective contacts by taking the ratio of the face-selective amplitude (i.e., at 1.2Hz and harmonics, see above) to the sum of the face-selective and general visual amplitude (i.e., 6Hz and harmonics). This index provides an additional quantification of face-selectivity by taking into account the magnitude of response to non-face stimuli. It varies from 0 (no face-selective response) to 1 (face-selective response only, no general visual response). Face-selectivity indices were either computed at the scale of main VOTC regions or computed in Talairach space as a function of the posterior-anterior axis (Y TAL dimension). In the latter case, face-selectivity indices were computed along

the Y dimension in segments of 15 mm and by steps of 3 mm, collapsing contacts across the X (lateral-medial, collapsed across both hemispheres) and Z (inferior-superior) dimensions. Face-selective and general visual activity were first averaged across contacts (i.e., within regions or voxels) before computing the index. This avoided obtaining indices outside the [0 - 1] range if, for instance, the denominator is smaller than 1.

HFB response timing parameters. In a first group timing analysis, the averaged HFB time-domain activity per recording contact (see '*HFB time-domain preprocessing*', **Figure 1G**) were grouped by main VOTC region (OCC, PTL, and ATL) and hemisphere. The timing of face-selective activity was characterized using 4 parameters: (1) latency of response onset and (2) offset, as well as (3) response overlap and (4) response correlation across VOTC regions (**Figure S2**). The first two parameters, onset and offset latencies of face-selective response, were computed per group of contacts using a bootstrapping approach in the following way (**Figure S2A**): (1) randomly sample contacts with replacement; (2) averaging HFB time-series from sampled contacts; (3) converting this averaged HFB time-series to Z-score values by subtracting the mean amplitude in the baseline window of the time-series (i.e. before face onset: [-0.166 to 0s]) and dividing by the standard deviation of the amplitude in the same baseline window; (4) converting the z-score values to *p*-values, apply an FDR correction and compute onset latency as the first time-point at which the $p < 0.05$ (two-tailed), for at least 30 ms. Offset latency was the first time bin (after the onset) at which the response was no longer significantly different from baseline; (5) store onset and offset latencies for the current bootstrap sample; (6) repeat steps (1) to (5) 4000 times; (7) use the obtained bootstrap distributions of onsets/offsets to compute the mean and 95% confidence interval for these 2 parameters. Onset and offset latencies were compared across regions within hemispheres and signal type using a permutation test, randomly shuffling (5000 times) the assignment of contacts in the two regions compared. Third, we estimated the overlap between time-series for pairs of regions (e.g., region A and B) within hemispheres (**Figure S2B**). The overlap is asymmetrical and is calculated separately for region A and region B as the ratio between the area under the curve (AUC) of the overlap between regions A and region B (i.e., summing the amplitude between the maximum of onset A and B and the minimum of offset A and B) and the total AUC for region A (for overlap of region A to B) or for region B (for overlap of region B to A). Fourth, for each signal type we computed Pearson correlations between the time-series obtained in the three main VOTC regions, within each hemisphere (**Figure S2C**). For each pair of regions, we determined whether the between-region

correlation coefficient was significantly lower than within-regions correlations using a bootstrapping approach: (1) compute the *between-region correlation* by repeatedly (5000 times) randomly sampling half of the available time-series from each of the two regions, averaging the time-series for each region, calculate the Pearson correlation between these averaged time-series and take the mean over the obtained 5000 correlation coefficient; (2) compute the within-region correlation for each region in a similar manner by repeatedly (5000 times) randomly splitting the available time-series in two groups, average the time-series for each group, correlate time-series across the two groups and take the mean over the obtained 5000 correlation coefficient.

Functional connectivity between VOTC regions. Connectivity between regions was computed using participants that had face-selective contact (s) in at least 2 of the dominant face-selective regions in the same hemisphere: right/left IOG, latFG, antFG+. We used largely the same approach as in (Kadipasaoglu et al., 2017) , except for the statistics. Starting from the single-trial HFB time-domain activity per recording contact, we performed the following steps: (1) dowsampling of the time-series by a factor of 2 to save computation time; (2) baseline-correction of each trial by subtracting the mean amplitude in the -166 to 0 ms; (3) subtract from each single-trial, the mean across trials to obtain trial-by-trial variability around the mean. The single-trial amplitudes were correlated between pairs of electrodes at each time point from -100 to 550 ms to compute the instantaneous functional connectivity between regions. To explore the directionality in the connectivity, we also computed the across-trial correlations after introducing variable temporal lags between regions (from -150 to 150 ms). For each pair of contacts, we lagged the time-series at one of the contact before computing the correlation. This allows determining directionality between regions if the amplitude of face-selective activity at a given time point in one region correlates with amplitude in the other region at an earlier or later time-point. Correlations were summarized in lagged-correlation matrices (**Figure 5**). In each participant and each pair of regions, correlations were computed across all possible pairs of face-selective contacts, averaged across contacts and then averaged across participants to obtain group estimates. Statistics were performed using a randomization test to determine the null distribution, by running 1000 times the same analyses steps after having randomized the order of trials between pairs of electrodes being correlated. Group-level correlations were tested against the null distributions at p < 0.01 fdr-corrected.

HFB onset latency map. In addition to computing timing parameters across main VOTC regions, we computed a group map of local face-selective response onset latency across VOTC in Talairach space. We averaged HFB time-series from contiguous face-selective contacts located within voxels of 20 x 20 x 100 mm (swept across VOTC in steps of 3 x 3 x 100 mm; see above *'Group visualization and analyses in Talairach space')* and computed onset latency of face-selective response for the averaged time-series in the same way as for the analyses in main VOTC regions (see *'HFB response timing parameters',* **Figure S2**). Due to the lower sample size in any given local measurement (i.e., voxel) compared to when using larger anatomically-defined regions and to limit noise in the latency estimates, we rejected recording contacts with split-half correlation below r = 0.4. Split-half correlation was computed for each face-selective contact by repeatedly (5000 times) randomly splitting the available single face trials (i.e., short time-series evoked by individual face presentations) in 2 groups, averaging the time-series for each group, calculate the Pearson correlation between these averaged time-series and take the mean over the obtained 5000 correlation coefficients. In additions, we only considered voxels with more than 5 contacts. We also examined the poster-anterior variation of face-selective response latency by averaging HFB time-series from contacts located in voxels of 20 mm (along the Y dimension) collapsing contacts across the X (lateral-medial, separately for each hemisphere) and Z (inferior-superior) dimensions. Voxels were swept across the postero-anterior axis in steps of 3 mm. We used a randomization procedure to determine whether the measured postero-anterior profile of face-selective response onset latency deviate from what would be expected under the (null) hypothesis that the postero-anterior VOTC location has no influence on the onset latency. Specifically, we built a random null distribution by repeatedly (5000 times) computed the postero-anterior profile of onset latency after having randomly shuffled the association between contacts (and associated averaged time-series) and their TAL coordinates. P-values were determined for each voxel as the fraction of the null distribution that was higher/lower than the actual measured onset latencies and were corrected for multiple comparisons in each hemisphere using FDR (Benjamini and Hochberg, 1995).

Acknowledgments/Funding

This research was supported by ERC AdG HUMANFACE 101055175

References

Allison, T., Puce, A., Spencer, D.D., McCarthy, G., Belger, A., 1999. Electrophysiological studies of human face perception. I: Potential generated in occiptotemporal cortex by face and non-face stimuli. Cerebral Cortex 9, 415–430.

Bar, M., 2003. A cortical mechanism for triggering top-down facilitation in visual object recognition. J Cogn Neurosci 15, 600–609. https://doi.org/10.1162/089892903321662976

Benjamini, Y., Hochberg, Y., 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B (Methodological) 57, 289–300.

Bryant, K.L., Glasser, M.F., Li, L., Bae, J.J., Jacquez, N.J., Preuss, T.M., 2019. Organization of extrastriate and temporal cortex in chimpanzees compared to humans and macaques. Cortex 118, 223–243. https://doi.org/10.1016/j.cortex.2019.02.010

Bullier, J., 2001. Integrated model of visual processing. Brain Res Brain Res Rev 36, 96–107.

Bullier, J., Nowak, L.G., 1995. Parallel Versus Serial Processing - New Vistas on the Distributed Organization of the Visual-System. Curr Opin Neurobiol 5, 497–503.

Cao, R., Zhang, J., Zheng, J., Wang, Y., Brunner, P., Willie, J.T., Wang, S., 2025. A neural computational framework for face processing in the human temporal lobe. Current Biology. https://doi.org/10.1016/j.cub.2025.02.063

Cauchoix, M., Barragan-Jason, G., Serre, T., Barbeau, E.J., 2014. The neural dynamics of face detection in the wild revealed by MVPA. The Journal of Neuroscience 34, 846–854. https://doi.org/10.1523/JNEUROSCI.3030-13.2014

Cavanagh, P., 1991. What's up in top-down processing, in: Gorea, A. (Ed.), Representations of Vision: Trends and Tacit Assumptions in Vision Research. Cambridge University Press.

Chen, X., Liu, X., Parker, B.J., Zhen, Z., Weiner, K.S., 2023. Functionally and structurally distinct fusiform face area (s) in over 1000 participants. Neuroimage 265, 119765. https://doi.org/10.1016/j.neuroimage.2022.119765

Conner, C.R., Ellmore, T.M., DiSano, M.A., Pieters, T.A., Potter, A.W., Tandon, N., 2011. Anatomic and electro-physiologic connectivity of the language system: A combined DTI-CCEP study. Comput Biol Med 41, 1100–1109. https://doi.org/10.1016/j.compbiomed.2011.07.008

Conway, B.R., 2018. The organization and operation of inferior temporal cortex. Annu Rev Vis Sci. https://doi.org/10.1146/annurev-vision-091517-034202

Cox, R.W., 1996. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. Comput Biomed Res 29, 162–173. https://doi.org/10.1006/cbmr.1996.0014

Crouzet, S.M., Thorpe, S.J., 2010. Fast saccades toward faces: Face detection in just 100 ms. J Vis 10, (4) 16, 1-17. https://doi.org/10.1167/10.4.16.

DiCarlo, J.J., Zoccolan, D., Rust, N.C., 2012. How does the brain solve visual object recognition? Neuron 73, 415–434. https://doi.org/10.1016/j.neuron.2012.01.010

Duchaine, B., Yovel, G., 2015. A Revised Neural Framework for Face Processing. Annu Rev Vis Sci 1, 393–416. https://doi.org/10.1146/annurev-vision-082114-035518

Edelman, G.M., 2004. Biochemistry and the Sciences of Recognition. Journal of Biological Chemistry. https://doi.org/10.1074/jbc.X400001200

Edelman, G.M., Gally, J.A., 2013. Reentry: A key mechanism for integration of brain function. Front Integr Neurosci. https://doi.org/10.3389/fnint.2013.00063

Engell, A.D., McCarthy, G., 2014. Face, eye, and body selective responses in fusiform gyrus and adjacent cortex: an intracranial EEG study. Front Hum Neurosci 8, 642. https://doi.org/10.3389/fnhum.2014.00642

Fairhall, S.L., Ishai, A., 2007. Effective connectivity within the distributed cortical network for face perception. Cerebral Cortex 17, 2400–2406. https://doi.org/10.1093/cercor/bhl148

Finzi, D., Gomez, J., Nordt, M., Rezai, A.A., Poltoratski, S., Grill-Spector, K., 2021. Differential spatial computations in ventral and lateral face-selective regions are scaffolded by structural connections. Nat Commun 12, 2278. https://doi.org/10.1038/s41467-021-22524-2

Freiwald, W.A., 2020. The neural mechanisms of face processing: cells, areas, networks, and models. Curr Opin Neurobiol 60, 184–191. https://doi.org/10.1016/j.conb.2019.12.007

Fuster, J., 1995. Memory in the Cerebral Cortex: An Empirical Approach to Neural Networks in the Human and Nonhuman Primates., MIT Press. MIT Press, Cambridge, MA.

Gao, X., Gentile, F., Rossion, B., 2018. Fast periodic stimulation (FPS): a highly effective approach in fMRI brain mapping. Brain Struct Funct 223, 2433–2454. https://doi.org/10.1007/s00429-018-1630-4

Grill-Spector, K., Weiner, K.S., Kay, K.N., Gomez, J., 2017. The Functional Neuroanatomy of Human Face Perception. Annu Rev Vis Sci. https://doi.org/10.1146/annurev-vision-102016-061214

Hagen, S., Jacques, C., Ranta, R., Koessler, L., Maillard, L., Colnat-Coulbois, S., Rossion, B., Jonas, J., 2025. Intracranial EEG referencing for large-scale category-selective mapping in the human ventral occipito-temporal cortex. Imaging Neuroscience 3. https://doi.org/10.1162/imag_a_00479

Hauk, O., Rice, G.E., Volfart, A., Magnabosco, F., Ralph, M.A.L., Rossion, B., 2021. Face-selective responses in combined EEG/MEG recordings with fast periodic visual stimulation (FPVS). Neuroimage 242, 118460. https://doi.org/10.1016/j.neuroimage.2021.118460

Hubel, D.H., Wiesel, T.N., 1968. Receptive fields and functional architecture of monkey striate cortex. Journal of Physiology 195, 215–243.

Hubel, D.H., Wiesel, T.N., 1962. Receptive Fields, Binocular Interaction and Functional Architecture in the Cat's Visual Cortex. J. Physiol. 160, 106–154.

Issa, E.B., Cadieu, C.F., DiCarlo, J.J., 2018. Neural dynamics at successive stages of the ventral visual stream are consistent with hierarchical error signals. Elife 7, e42870. https://doi.org/10.7554/eLife.42870

Jacobs, G.H., 1999. Vision and Behavior in Primates, in: Archer, S.N., Djamgoz, M.B.A., Loew, E.R., Partridge, J.C., Vallerga, S. (Eds.), Adaptive Mechanisms in the Ecology of Vision. Springer Netherlands, Dordrecht, pp. 629–650. https://doi.org/10.1007/978-94-017-0619-3_21

Jacques, C., Jonas, J., Colnat-Coulbois, S., Maillard, L., Rossion, B., 2022. Low and high frequency intracranial neural signals match in the human associative cortex. Elife 11, e76544. https://doi.org/10.7554/eLife.76544

Jacques, C., Retter, T.L., Rossion, B., 2016a. A single glance at natural face images generate larger and qualitatively different category-selective spatio-temporal signatures than other ecologically-relevant categories in the human brain. Neuroimage 137, 21–33. https://doi.org/10.1016/j.neuroimage.2016.04.045

Jacques, C., Witthoft, N., Weiner, K.S., Foster, B.L., Rangarajan, V., Hermes, D., Miller, K.J., Parvizi, J., Grill-spector, K., 2016b. Corresponding ECoG and fMRI category-selective signals in human ventral temporal cortex. Neuropsychologia 83, 14–28. https://doi.org/10.1016/j.neuropsychologia.2015.07.024

Jiang, F., Dricot, L., Weber, J., Righi, G., Tarr, M.J., Goebel, R., Rossion, B., 2011. Face categorization in visual scenes may start in a higher order area of the right fusiform gyrus:

Evidence from dynamic visual stimulation in neuroimaging. J Neurophysiol 106, 2720–2736. https://doi.org/10.1152/jn.00672.2010

Jonas, J., Jacques, C., Liu-shuang, J., Brissart, H., Colnat-coulbois, S., Maillard, L., Rossion, B., 2016a. A face-selective ventral occipito-temporal map of the human brain with intracerebral potentials. Proc Natl Acad Sci U S A 113, E4088--E4097. https://doi.org/10.1073/pnas.1522033113

Jonas, J., Jacques, C., Liu-shuang, J., Brissart, H., Colnat-coulbois, S., Maillard, L., Rossion, B., 2016b. A face-selective ventral occipito-temporal map of the human brain with intracerebral potentials. Proc Natl Acad Sci U S A 113, E4088--E4097. https://doi.org/10.1073/pnas.1522033113

Jonas, J., Rossion, B., 2021. Intracerebral electrical stimulation to understand the neural basis of human face identity recognition. Eur J Neurosci 54, 4197–4211. https://doi.org/10.1111/ejn.15235

Kadipasaoglu, C.M., Conner, C.R., Baboyan, V.G., Rollo, M., Pieters, T.A., Tandon, N., 2017. Network dynamics of human face perception. PLoS One 12, e0188834. https://doi.org/10.1371/journal.pone.0188834

Kadipasaoglu, C.M., Conner, C.R., Whaley, M.L., Baboyan, V.G., Tandon, N., 2016. Category-Selectivity in Human Visual Cortex Follows Cortical Topology: A Grouped icEEG Study. PLoS One 11, e0157109. https://doi.org/10.1371/journal.pone.0157109

Kar, K., DiCarlo, J.J., 2021. Fast Recurrent Processing via Ventrolateral Prefrontal Cortex Is Needed by the Primate Ventral Stream for Robust Core Visual Object Recognition. Neuron 109, 164-176.e5. https://doi.org/10.1016/j.neuron.2020.09.035

Kar, K., Kubilius, J., Schmidt, K., Issa, E.B., DiCarlo, J.J., 2019. Evidence that recurrent circuits are critical to the ventral stream's execution of core object recognition behavior. Nat Neurosci 22, 974–983. https://doi.org/10.1038/s41593-019-0392-5

Kay, K.N., Weiner, K.S., Grill-Spector, K., 2015. Attention reduces spatial uncertainty in human ventral temporal cortex. Current Biology 25, 595–600. https://doi.org/10.1016/j.cub.2014.12.050

Kim, M., Ducros, M., Carlson, T., Ronen, I., He, S., Ugurbil, K., Kim, D.S., 2006. Anatomical correlates of the functional organization in the human occipitotemporal cortex. Magn Reson Imaging 24, 583–590. https://doi.org/10.1016/j.mri.2005.12.005

Kosakowski, H.L., Cohen, M.A., Takahashi, A., Keil, B., Kanwisher, N., Saxe, R., 2022. Selective responses to faces, scenes, and bodies in the ventral visual pathway of infants. Current Biology 32, 265-274.e5. https://doi.org/10.1016/j.cub.2021.10.064

Krienen, F.M., Buckner, R.L., 2020. Chapter 35 - Human Association Cortex: Expanded, Untethered, Neotenous, and Plastic, in: Kaas, J.H. (Ed.), Evolutionary Neuroscience (Second Edition). Academic Press, London, pp. 845–860. https://doi.org/https://doi.org/10.1016/B978-0-12-820584-6.00035-0

Kubota, E., Yan, X., Tung, S., Fascendini, B., Tyagi, C., Duhameau, S., Ortiz, D., Grotheer, M., Natu, V.S., Keil, B., Grill-Spector, K., 2025. White matter connections of human ventral temporal cortex are organized by cytoarchitecture, eccentricity and category-selectivity from birth. Nat Hum Behav. https://doi.org/10.1038/s41562-025-02116-6

Lamme, V.A.F., Roelfsema, P.R., 2000. The distinct modes of vision offered by feedforward and recurrent processing. Trends Neurosci 23, 571–579.

Laurent, M.-A., Jacques, C., Yan, X., Jurczynski, P., Colnat-Coulbois, S., Maillard, L., Cam, S. Le, Ranta, R., Cottereau, B.R., Koessler, L., Jonas, J., Rossion, B., 2025. A tight relationship between BOLD fMRI activation/deactivation and increase/decrease in single neuron responses in human association cortex. https://doi.org/10.7554/elife.104779.1

Liu, H., Agam, Y., Madsen, J.R., Kreiman, G., 2009. Timing, timing, timing: fast decoding of object information from intracranial field potentials in human visual cortex. Neuron 62, 281–90. https://doi.org/10.1016/j.neuron.2009.02.025

Manning, J.R., Jacobs, J., Fried, I., Kahana, M.J., 2009. Broadband shifts in local field potential power spectra are correlated with single-neuron spiking in humans. Journal of Neuroscience 29, 13613–13620. https://doi.org/10.1523/JNEUROSCI.2041-09.2009

Marr, D., 1982. Vision: A Computational Investigation into the Human Representation and Processing of Visual Information, W. H. Free. ed. W. H. Freeman and Co, San Francisco, CA, USA.

Mumford, D., 1992. On the computational architecture of the neocortex II The role of cortico-cortical loops, Biol. Cybern.

Norcia, A.M., Appelbaum, L.G., Ales, J.M., Cottereau, B.R., Rossion, B., 2015. The steady-state visual evoked potential in vision research: A review. J Vis 15, (6) 4, 1--46. https://doi.org/10.1167/15.6.4

Nowak, L.G., Bullier, J., 1997. The Timing of Information Transfer in the Visual System, in: Rockland, K.S., Kaas, J.H., Peters, A. (Eds.), Extrastriate Cortex in Primates. Springer US, Boston, MA, pp. 205–241. https://doi.org/10.1007/978-1-4757-9625-4_5

Ojemann, J.G., Akbudak, E., Snyder, A.Z., McKinstry, R.C., Raichle, M.E., Conturo, T.E., 1997. Anatomic localization and quantitative analysis of gradient refocused echo-planar fMRI susceptibility artifacts. Neuroimage 6, 156–167. https://doi.org/10.1006/nimg.1997.0289

Perrett, D.I., Oram, M.W., Ashbridge, E., 1998. Evidence accumulation in cell populations responsive to faces: an account of generalisation of recognition without mental transformations. Cognition 67, 111–145.

Persichetti, A.S., Denning, J.M., Gotts, S.J., Martin, A., 2021. A data-driven functional mapping of the anterior temporal lobes. Journal of Neuroscience 41, 6038–6049. https://doi.org/10.1523/JNEUROSCI.0456-21.2021

Quek, G.L., Rossion, B., 2017. Category-selective human brain processes elicited in fast periodic visual stimulation streams are immune to temporal predictability. Neuropsychologia 104, 182–200. https://doi.org/10.1016/j.neuropsychologia.2017.08.010

Quian Quiroga, R., Boscaglia, M., Jonas, J., Rey, H.G., Yan, X., Maillard, L., Colnat-Coulbois, S., Koessler, L., Rossion, B., 2023. Single neuron responses underlying face recognition in the human midfusiform face-selective cortex. Nat Commun 14, 5661. https://doi.org/10.1038/s41467-023-41323-5

Rangarajan, V., Hermes, D., Foster, B.L., Weiner, K.S., Jacques, C., Grill-Spector, K., Parvizi, J., 2014. Electrical Stimulation of the Left and Right Human Fusiform Gyrus Causes Different Effects in Conscious Face Perception. The Journal of Neuroscience 34, 12828–12836. https://doi.org/10.1523/JNEUROSCI.0527-14.2014

Regan, D., 1989. Human Brain Electrophysiology: Evoked Potentials and Evoked Magnetic Fields in Science and Medicine, Elsevier. ed. New York, NY, USA.

Retter, T.L., Jiang, F., Webster, M.A., Rossion, B., 2020. All-or-none face categorization in the human brain. Neuroimage 213, 116685. https://doi.org/10.1016/j.neuroimage.2020.116685

Retter, T.L., Rossion, B., 2016. Uncovering the neural magnitude and spatio-temporal dynamics of natural image categorization in a fast visual stream. Neuropsychologia 91, 9–28. https://doi.org/10.1016/j.neuropsychologia.2016.07.028

Riesenhuber, M., Poggio, T., 1999. Hierarchical models of object recognition in cortex. Nat Neurosci 2, 1019–1025.

Rossion, B., 2022. Twenty years of investigation with the case of prosopagnosia PS to understand human face identity recognition.Part II: Neural basis. Neuropsychologia 173, 108279. https://doi.org/10.1016/j.neuropsychologia.2022.108279

Rossion, B., Caharel, S., 2011. ERP evidence for the speed of face categorization in the human brain: Disentangling the contribution of low-level visual cues from face perception. Vision Res 51, 1297–1311. https://doi.org/10.1016/j.visres.2011.04.003

Rossion, B., Caldara, R., Seghier, M., Schuller, A.M., Lazeyras, F., Mayer, E., 2003. A network of occipito-temporal face-sensitive areas besides the right middle fusiform gyrus is necessary for normal face processing. Brain 126, 2381–2395. https://doi.org/10.1093/brain/awg241

Rossion, B., Hanseeuw, B., Dricot, L., 2012. Defining face perception areas in the human brain: a large-scale factorial fMRI face localizer analysis. Brain Cogn 79, 138–157. https://doi.org/10.1016/j.bandc.2012.01.001

Rossion, B., Jacques, C., Jonas, J., 2024. The anterior fusiform gyrus: The ghost in the cortical face machine. Neurosci Biobehav Rev. https://doi.org/https://doi.org/10.1016/j.neubiorev.2024.105535

Rossion, B., Jacques, C., Jonas, J., 2018. Mapping face categorization in the human ventral occipito-temporal cortex with direct neural intracranial recordings. Ann N Y Acad Sci 1426, 5–24. https://doi.org/10.1111/nyas.13596

Rossion, B., Jacques, C., Liu-shuang, J., 2015. Fast periodic presentation of natural images reveals a robust face-selective electrophysiological response in the human brain. J Vis 15, (1) 18, 1--18. https://doi.org/10/1167.15.1.18.

Rousselet, G., Husk, J.S., Bennett, P.J., Sekuler, A.B., 2008. Time course and robustness of ERP object and face differences. J Vis 8, (12) 3, 1-18. https://doi.org/10.1167/8.12.3.

Sadeh, B., Podlipsky, I., Zhdanov, A., Yovel, G., 2010. Event-related potential and functional MRI measures of face-selectivity are highly correlated: a simultaneous ERP-fMRI investigation. Hum Brain Mapp 31, 1490–1501. https://doi.org/10.1002/hbm.20952

Schmolesky, M.T., Wang, Y.C., Hanes, D.P., Thompson, K.G., Leutgeb, S., Schall, J.D., Leventhal, A.G., 1998. Signal timing across the macaque visual system. J Neurophysiol 79, 3272–3278.

Schrouff, J., Raccah, O., Baek, S., Rangarajan, V., Salehi, S., Mourão-Miranda, J., Helili, Z., Daitch, A.L., Parvizi, J., 2020. Fast temporal dynamics and causal relevance of face processing in the

human temporal cortex. Nat Commun 11, 656. https://doi.org/10.1038/s41467-020-14432-8

Schweinberger, S.R., Neumann, M.F., 2016. Repetition effects in human ERPs to faces. Cortex 80, 141–153. https://doi.org/10.1016/j.cortex.2015.11.001

Serre, T., Oliva, A., Poggio, T., 2007. A feedforward architecture accounts for rapid categorization. Proceedings of the National Academy of Sciences 104, 6424–6429. https://doi.org/10.1073/pnas.0700622104

Steeves, J.K., Culham, J.C., Duchaine, B.C., Pratesi, C.C., Valyear, K.F., Schindler, I., Humphrey, G.K., Milner, A.D., Goodale, M.A., 2006. The fusiform face area is not sufficient for face recognition: evidence from a patient with dense prosopagnosia and no occipital face area. Neuropsychologia 44, 594–609.

Talairach, J., Bancaud, J., 1973. Stereotaxic approach to epilepsy. Methodology of anatomo-functional stereotaxic investigations. Prog Neurol Surg 5, 297–354. https://doi.org/https://doi.org/10.1159/000394343

Thorpe, S.J., Fabre-Thorpe, M., 2001. Seeking Categories in the Brain. Science (1979) 291, 260–263. https://doi.org/10.1126/science.1058249

Tsao, D., 2014. The macaque face patch system: A window into object representation. Cold Spring Harb Symp Quant Biol 79, 109–114. https://doi.org/10.1101/sqb.2014.79.024950

Tsao, D.Y., Moeller, S., Freiwald, W. a, 2008. Comparing face patch systems in macaques and humans. Proc Natl Acad Sci U S A 105, 19514–19519. https://doi.org/10.1073/pnas.0809662105

Ungerleider, L.G., Bell, A.H., 2011. Uncovering the visual " alphabet": Advances in our understanding of object perception. Vision Res. https://doi.org/10.1016/j.visres.2010.10.002

van Blooijs, D., van den Boom, M.A., van der Aar, J.F., Huiskamp, G.M., Castegnaro, G., Demuru, M., Zweiphenning, W.J.E.M., van Eijsden, P., Miller, K.J., Leijten, F.S.S., Hermes, D., 2023. Developmental trajectory of transmission speed in the human brain. Nat Neurosci 26, 537–541. https://doi.org/10.1038/s41593-023-01272-0

Van Essen, D.C., Anderson, C.H., Felleman, D.J., 1992. Information processing in the primate visual system: an integrated systems perspective. Science (1979) 255, 419–23.

Van Essen, D.C., Maunsell, J.H.R., 1983. Hierarchical organization and functional streams in the visual cortex. Trends Neurosci 6, 370–375. https://doi.org/https://doi.org/10.1016/0166-2236 (83)90167-4

Vidal, J.R., Ossandón, T., Jerbi, K., Dalal, S.S., Minotti, L., Ryvlin, P., Kahane, P., Lachaux, J.-P., 2010. Category-Specific Visual Responses: An Intracranial Study Comparing Gamma, Beta, Alpha, and ERP Response Selectivity. Front Hum Neurosci 4, 195. https://doi.org/10.3389/fnhum.2010.00195

Volfart, A., Yan, X., Maillard, L., Colnat-Coulbois, S., Hossu, G., Rossion, B., Jonas, J., 2022. Intracerebral electrical stimulation of the right anterior fusiform gyrus impairs human face identity recognition. Neuroimage 250, 118932. https://doi.org/10.1016/j.neuroimage.2022.118932

Wang, Y., Metoki, A., Smith, D. V., Medaglia, J.D., Zang, Y., Benear, S., Popal, H., Lin, Y., Olson, I.R., 2020. Multimodal mapping of the face connectome. Nat Hum Behav 4, 397–411. https://doi.org/10.1038/s41562-019-0811-3

Weiner, K.S., Grill-Spector, K., 2010. Sparsely-distributed organization of face and limb activations in human ventral temporal cortex. Neuroimage 52, 1559–1573. https://doi.org/10.1016/j.neuroimage.2010.04.262

Weiner, K.S., Jonas, J., Gomez, J., Maillard, L., Brissart, H., Hossu, G., Jacques, C., Loftus, D., Colnat-Coulbois, S., Stigliani, A., Barnett, M.A., Grill-Spector, K., Rossion, B., 2016. The Face-Processing Network Is Resilient to Focal Resection of Human Visual Cortex. The Journal of neuroscience 36, 8425–8440. https://doi.org/10.1523/JNEUROSCI.4509-15.2016

Zeki, S., Shipp, S., 1988. The functional logic of cortical connections. Nature 335, 311–317. https://doi.org/10.1038/335311a0