The PPARβ/delta-induced mesenchymal stromal cell secretome has cytoprotective effects via ANGPTL4 in a pre-clinical model of acute lung inflammation

Courteney Tunstead[1], Molly Dunlop[1], Sinéad Ryan[1], Evelina Volkova[1], Evangeline Johnston[1], Sabrina Batah[2], Claudia C. Dos Santos[2], Bairbre McNicholas [3,4], Claire Masterson[3,4], John G. Laffey[3,4] & Karen English[1,5,*]

[1]*Cellular Immunology Lab, Department of Biology, Maynooth University, Maynooth, Co. Kildare, Ireland.*

[2] *Keenan Research Centre for Biomedical Research, St. Michael's Hospital, Toronto, Canada.*

[3]*Anesthesia and Intensive Care Medicine, School of Medicine, College of Medicine Nursing and Health Sciences, University of Galway, Galway, Ireland.*

[4]*Anesthesia and Intensive Care Medicine, Galway University Hospitals, Saolta University Hospitals Groups, Galway, Ireland.*

[5]*Kathleen Lonsdale Institute for Human Health Research, Maynooth University, Maynooth, Co. Kildare, Ireland.*

[*]*Corresponding author*

Corresponding author information

Prof. Karen English

karen.english@mu.ie

Keywords: MSC, ARDS, Lung reapir, ANGPTL4, PPARβ/δ

ABSTRACT

Rationale: Human bone marrow-derived mesenchymal stromal cells (hBM-MSCs) are known

to exert immunomodulatory and pro-reparative effects *in vivo*. This makes hBM-MSCs an

enticing therapeutic candidate for inflammatory diseases, such as acute respiratory distress

syndrome (ARDS). The ARDS microenvironment is complex and contains an abundance of

free fatty acids (FFAs); which are known to differentially impact MSC functionality. PPARβ/δ

is a ubiquitously expressed nuclear receptor that is activated in response to FFA-binding.

PPARβ/δ has been shown to impact the therapeutic efficacy of mouse MSCs. **Objective:** This

study sought to investigate the impact of PPARβ/δ-modulation on human MSC functionality

in vitro and *in vivo.* **Methods:** hBM-MSCs were exposed to a synthetic PPARβ/δ

agonist/antagonist in the presence or absence of ARDS patient serum and the

immunomodulatory and pro-reparative capacity of the MSC secretome was investigated using

in vitro assays and a pre-clinical model of LPS-induced acute lung inflammation (ALI).

Results: Our results highlighted enhanced pro-reparative capacity of PPARβ/δ-agonised hBM-

MSCs secretome in CALU-3 lung epithelial cells, mediated by MSC derived angiopoietin-like

4 (ANGPTL4). PPARβ/δ-induced ANGPTL4-high MSC secretome facilitated enhanced

endothelial barrier integrity in the lungs of ALI mice. Therapeutic effects of PPARβ/δ-

agonised hBM-MSCs secretome were further enhanced by licensing MSCs with human ARDS

patient serum. ARDS-licensed PPARβ/δ-induced ANGPTL4-high MSC secretome had

reduced clinical score and weight loss. The role ANGPL4 in these protective effects was

confirmed using an anti-ANGPTL4 antibody. **Conclusion:** These findings conclude that the

MSC secretome therapeutic effects can be enhanced both *in vitro* and *in vivo* through licensing

strategies that upregulate the angiogenic factor ANGPTL4.

INTRODUCTION

Cell therapy, particularly mesenchymal stromal cell (MSC) therapy, has long been thought of as a therapeutic candidate in the treatment of many inflammatory diseases. This is largely due to their immunomodulatory and cytoprotective capacity [1]. One of the conditions that MSCs have been considered for is acute respiratory distress syndrome (ARDS); an inflammatory, degenerative condition of the lung [2, 3]. MSCs have been trialled and tested for ARDS over the last 20 years, but clinical trials have shown poor outcomes; with ~60% non-responders [4-6]. The environment into which MSCs are administered can alter their functional capacity [7-9]. There is limited understanding of how the ARDS microenvironment may impact MSCs. The complex ARDS microenvironment contains inflammatory-mediators, microorganisms, and free fatty acids (FFAs). These components have all been shown to impact MSC therapeutic capacity in various ways [10, 11]. Our previous work highlighted the differential effect of the hypo-inflammatory vs hyper-inflammatory ARDS microenvironment on MSC therapeutic efficacy with a clear correlation between high levels of inflammation, and enhanced MSC-driven repair [12]. We, and others, have also demonstrated that FFAs can influence MSC-induced immunosuppressive effects [13, 14]. In this study, we wanted to investigate the influence of FFAs present in the ARDS patient microenvironment, and the impact they have on hBM-MSCs in a pre-clinical model of acute lung inflammation (ALI).

The peroxisome proliferator-activated receptor-beta/delta (PPARβ/δ) is a ubiquitously-expressed nuclear receptor, present on all nucleated cells including hBM-MSCs, that is activated in response to FFA exposure [15, 16]. In recent years, PPARβ/δ has been shown to impact MSC therapeutic efficacy [17-27]. Interestingly, there are differential findings with some studies showing that deficiency of PPARβ/δ enhances MSC immunosuppressive effects

[21, 22], while other studies demonstrate similar immunosuppressive effects mediated by both

naive and PPARβ/δ-deficient mouse MSCs [24]. Studies have shown enhanced

immunomodulatory capacity of T-cells by PPARβ/δ-inhibited mouse MSCs [21, 22].

Differentially, PPARβ/δ antagonism of MSCs impaired MSC therapeutic effects while priming

of mouse MSCs with a PPARβ/δ agonist enhanced MSC cardioprotective effects *in vivo* [25].

Moreover, cardioprotective effects of MSCs were shown to be PPARβ/δ-dependent as the

therapeutic effect was lost when PPARβ/δ-deficient mouse MSCs were administered [24].

These findings suggest that natural ligands of PPARβ/δ may have positive or detrimental

effects on MSC efficacy *in vivo*. Due to the abundance of FFAs in the ARDS

microenvironment, we sought to investigate the impact of PPARβ/δ (via synthetic agonism or

antagonism) on the human BM-MSC secretome; and report the impact this had on

immunomodulation and cytoprotection, in a model of ALI.

MATERIALS AND METHODS

hBM-MSC cell culture

Human bone marrow-derived MSCs (hBM-MSCs; RoosterBio™) were cultured in Dulbecco's

Modified Eagle Medium (DMEM) containing 10% foetal bovine serum (FBS) and 1%

Penicillin streptomycin (cDMEM). The cells were cultured for 5 days at 37°C + 5% CO_2. All

experiments are a representation of three independent MSC donors.

Generation and concentration of MSC-CM

hBM-MSCs (passage 2-5) from three independent donors were seeded at a density of $1x10^5$

per well in a 6-well plate in 1ml of cDMEM and allowed to adhere. Once adhered, the cells

were stimulated with 1µM of PPARβ/**δ** agonist (GW0742, Tocris™) or antagonist (GSK3787,

Tocris™), as previously described [12]. 24hrs-post stimulation, the cells were washed with

PBS, and the media was replaced with serum-free DMEM. This was left for a further 24hrs to

allow for the generation of MSC-CM. When including ARDS serum-exposure, the

agonist/antagonist were washed from the cells, and 20% ARDS serum was added for 24hrs.

This was then washed away, and serum-free DMEM was added for a further 24hrs to allow for

the generation of MSC-CM. All MSC-CM was harvested and stored at -20°C for future

experiments. MSC-CM was then concentrated, as required, as previously described [12].

Gene & Protein expression

For gene expression, RNA was isolated and normalised to 100 ng/µl. cDNA was generated

using the QuantBio™ cDNA Synthesis kit (as per manufacturer's instructions). Real-time

reverse transcription polymerase chain reaction (RT-PCR) was run using PerfeCta SYBR

Green FastMix (QuantBio™) and pre-made Sigma™ primers (supplementary table 1). Relative

expression was quantified in relation to *hprt* by calculating the 2-ΔΔCT values. For protein

expression, 96-well half-area ELISA plates (COSTAR™) were coated as per manufacturers

guidelines (human ANGPTL4 and VEGF (R&D™)) and run as previously described [12].

Sequencing & metabolomic analysis

hBM-MSCs were seeded at a density of $1x10^5$ in a 6-well plate and allowed to adhere. Once

attached, the cells were exposed to 1μM of PPARβ/**δ** agonist (GW0742, Tocris™) or

antagonist (GSK3787, Tocris™) and incubated for 6hrs. This dose was chosen based on

previous research from our group [19]. The cells were harvested in TRIzol for RNA isolation.

RNA was isolated through a series of chloroform and isopropanol extractions. RNA

concentration and purity was assessed using Nanodrop 2000 (Thermo Fisher Scientific) and

samples were sent to Novogene™ for sequencing (GEO accession no.: GSE281162). Processed

RNA sequencing datasets, provided by Novogene™, were then used to identify differentially

expressed genes (DEGs) of interest. Volcano plots highlighting the most significantly enriched

genes were generated using Flourish™ software. Data was also collected from publicly

available RNA sequencing (GEO accession no.: GSE185263) and metabolomic datasets (Study

ID: ST000042; doi: 10.21228/M8SG64). Normalised values were taken from both studies and

data was plotted using GraphPad Prism™ (version 10) software.

Scratch Assay

CALU-3 lung epithelial cells (cultured as previously described) [12] were seeded at a density

of $5x10^5$ in a 6-well plate. Once 80% confluent, a single vertical scratch was made down the

centre of the well using a p200 tip. The cells were then washed with PBS to remove debris, and

MSC-CM was added for a period of 6 - 48hrs. The cells were then fixed with 10% formalin,

before allowing to dry. Once dry, the cells were stained with crystal violet, allowed to dry, and

imaged. To avoid bias, a marker was used to denote an area in the upper, central and lower

regions of each scratch and images were taken from these points (supplementary figure 1).

Mitomycin C (Sigma™) was added to assess the impact of proliferation at a concentration of

5 µg/ml for 2hrs prior to the scratch assay.

Scratch Assay Analysis

Three images, taken at each of the three marked points, were taken per well; for each of the

three independent MSC donors (i.e. 3 images per well x 3 MSC donors = 9 images per group).

Fiji/ImageJ™ software was then used to divide each picture into four distinct quadrants in an

unbiased manner (Analyse -> Tools -> Grid -> set area per point to 44444 inches squared).

This grid includes the addition of 4 horizontal lines imputed on the image. Using the straight-

line feature in Fiji/ImageJ™ software, four measurements were then taken per picture at each

of the horizontal lines. This provided 4 measurements per picture, at unbiased points, to get a

clear depiction of each percentage wound closure. The average wound closure per donor was

calculated, and divided by the baseline scratch value, to get the final percentage wound closure

(supplementary figure 1).

Study cohorts and ethical approval for patient samples

Ethical approval was obtained from Galway University Medical Ethics Committee and

Maynooth University Ethics Committee. All participants provided full consent.

ARDS serum extraction

Serum samples from both male and female patients, aged 40-80 years old, were taken from patients with SARS-CoV-2-induced ARDS, from the patient pools we previously characterised and described [12].

ANGPTL4 Neutralisation

MSC-CM was exposed to 5 µg/ml of ANGPTL4 neutralising antibody (Cell Sciences™), or 5 µg/ml IgG2aκ isotype control (Rockland Immunochemicals™). This was incubated at room temperature for 1hr prior to use. This concentration was utilised as per manufacturer guidelines for our study.

ALI mouse model

Male and female C57BL6/J mice (Charles River), aged 8+ weeks were exposed to 2mg/kg of endotoxin (Sigma, 0001:B4 Serotype), or PBS control administered intratracheally (IT) with a total of 42 animals with sample size determined based on similar studies in the published literature. For IT administration, mice were placed in an isoflurane chamber with a steady flow of both isoflurane and oxygen for 30-45 seconds, depending on the weight of the mouse. Mice were then placed on the IT stand, and a nose cone of isoflurane was used in increments throughout the administration to ensure sufficient anaesthesia. Mice were randomized to groups according to cage mate sex and ID sequence. A single animal represents the experimental unit; and within each cage, there were mice receiving different treatments to account for potential confounders with numbers per group outlined in each figure legend. There was no specific inclusion or exclusion criteria used for this study; any animals not accounted for across all groups were due to required euthanasia prior to experimental end-point and limitation on the amount of ARDS patient serum available. Animals were identified using a numerical ID for the purpose of this study. Although information regarding groupings and

treatments was available to the lead researchers, the numerical IDs were used for data harvest

and data analysis to prevent bias in result interpretation. MSC-CM was prepared as described

in the methods section. 500µl of PPARβ/δ-modulated MSC-CM, i.e. the equivalent of what

would be produced from 5×10^4 (low dose) of hBM-MSCs, was then concentrated to a volume

of 30µl and administered intranasally (IN) 4hrs post-endotoxin challenge under light isoflurane

anaesthesia for 15-30 seconds. An Evan's Blue permeability assay was carried out by

administration of an intravenous (IV) tail vein injection of sterile-filtered Evan's Blue Dye

(10%) 1hr prior to harvesting. The mice were sacrificed using an intraperitoneal (IP) injection

of pentobarbital. Absorbance readings were taken to assess the total Evan's Blue (620nm and

740nm) in bronchoalveolar lavage fluid (BALF) blood and whole lungs. Measurements taken

from the mice included (a) pro-inflammatory cytokines screening of the BALF, (b) Evan's

Blue permeability absorbances/images, and (c) clinical scoring (based on HPRA-approved

scoring methodology) and weight loss. Statistical analysis was performed using GraphPad

Prism 10 software and statistical tests used have been highlighted in the figure legends of each

figure. The work has been reported in line with the ARRIVE guidelines 2.0.

Statistical Analysis

Power calculations were performed to guide sample size, and data were analyzed using

GraphPad Prism 10 software. One-way ANOVA, or where appropriate a Two-way ANOVA

followed by the post hoc Tukey's multiple comparison test, was used to assess significance.

For figure 2C, a t-test was carried out to identify significance in DEGs. All data are presented

as mean ± SEM and the n numbers are outlined in each figure legend.

RESULTS

The MSC$_{PPARβ/δ(+)}$ secretome enhances pro-reparative capacity in CALU-3 lung epithelial cells via migration and proliferation *in vitro*

A scratch assay in CALU-3 cells was used to investigate the impact of PPARβ/δ agonism (+) or antagonism (-) on MSC conditioned medium (MSC-CM) capacity to promote wound healing *in vitro* (figure 1A). Our data highlighted a significant increase in wound closure in both MSC-CM$_{Naive}$ and MSC-CM$_{PPARβ/δ(+)}$ groups (figure 1B, C). Interestingly, MSC-CM$_{PPARβ/δ(-)}$ promoted wound healing when compared to control medium but did not significantly enhance CALU-3 wound closure. While there was no statistical difference between MSC-CM$_{Naive}$ and MSC-CM$_{PPARβ/δ(-)}$ (figure 1B), the percentage wound closure in the MSC-CM$_{PPARβ/δ(-)}$ group was less than in the MSC-CM$_{Naive}$ group. There is a possibility that PPARβ/δ antagonism could lead to the production of inhibitory factors in the conditioned medium, however, our RNA-Seq data set did not identify any known inhibitory factors (see figure 2, table 2). It is known that wound closure can involve proliferation and or migration [28, 29], thus this experiment was repeated using Mitomycin C; a known proliferation inhibitor [30]. While the addition of Mitomycin C significantly reduced wound closure in the MSC-CM$_{PPARβ/δ(+)}$ group at the 48hr time point, the percentage wound healing in the presence of Mitomycin C remained higher than the medium control group suggesting that both proliferation and migration may have a combination effect (figure 1D). Interestingly, significantly more wound repair was observed in the MSC-CM$_{PPARβ/δ(+)}$ group compared to the MSC-CM$_{Naive}$ group at 48hrs, while the MSC-CM$_{PPARβ/δ(-)}$ group did not inhibit or enhance would healing (figure 1D). *TOP2A is* a gene largely indicative of cell proliferation [31] and has previously been used as an indicator for cell proliferation in wound healing. Gene expression analysis of *TOP2A* in CALU-3 cells highlighted a significant difference between the MSC-CM$_{PPARβ/δ(+)}$ and MSC-CM$_{PPARβ/δ(-)}$

groups, with the MSC-CM$_{PPARβ/δ(+)}$ group having significantly more proliferative capacity than

the MSC-CM$_{PPARβ/δ(-)}$ group (figure 1E). Importantly, Mitoymcin C significantly reduced

TOP2A expression in the MSC-CM$_{Naive}$ and MSC-CM$_{PPARβ/δ(-)}$ groups (figure 1E).

Analysis of MSC$_{PPARβ/δ(+)}$ RNA sequencing highlighted upregulation of ANGPTL4

To try to identify the potential mechanisms associated with enhanced wound healing capacity

in MSC$_{PPARβ/δ(+)}$, RNA sequencing (Novogene™; GEO accession no.: GSE281162) was

performed on MSC$_{PPARβ/δ(+)}$, MSC$_{PPARβ/δ(-)}$ and MSC$_{Naive}$ (figure 2A). The data highlighted a

variety of differentially expressed genes (DEGs) between the three groups. The DEGs numbers

were visualised using a Venn diagram (figure 2B), showing 170 DEGs specific to the MSC$_{Naive}$

group, 128 specific to the MSC$_{PPARβ/δ(+)}$ group, and 189 specific to the MSC$_{PPARβ/δ(-)}$ group.

Volcano plot visualisation highlighted several of the most significantly altered DEGs in each

group (figure 2C). The MSC$_{PPARβ/δ(+)}$ group had a significant enrichment of genes such as

PLIN2, ANGPTL4, FAM156A, CPT1A, CAT, NADK2, TMED6, and *LDHAL6B.* This group

also showed significant downregulation in genes such as *MAGED4* and *FAM157B* (table 1).

The MSC$_{PPARβ/δ(-)}$ group had a significant enrichment of genes such as *IKBKGP1, SNRPGP2,*

LINC01159, MIR3648-1, RARRES1, MALAT1 and *RNF217-AS1.* This group also showed

significant downregulation in genes such as *RGPD5, PFN1P3, MIR3936HG, U2AF1* and

TSNAXIP1 (table 2). Fragments per kilobase of exon per million mapped fragments (FPKMs)

of two known PPARβ/**δ** target genes (angiopoietin like protein 4 (ANGPTL4) and pyruvate

dehydrogenase kinase 4 (PDK4)) were plotted to confirm the data was as expected (figure 2D,

E), and this was confirmed by qRT-PCR (figure 2F, G).

MSC-associated genes/proteins are unaffected by PPARβ/δ agonism or antagonism;

except ANGPTL4

Given the functional enhancement of MSC-CM$_{PPAR\beta/\delta(+)}$ promotion of wound healing observed

in figure 1, the production of known cytoprotective factors released by MSCs were assessed.

While MSC$_{Naive}$ produced ANGPTL4, VEGF and MIF, MSC$_{PPAR\beta/\delta(+)}$ only produced

significantly higher levels of ANGPTL4 (figure 3A-C). This aligns with the RNA-sequencing

data as *ANGPTL4* was the only cytoprotective factor differentially expressed in MSC$_{PPAR\beta/\delta(+)}$.

Along with this, PPARβ/**δ** agonism/antagonism did not affect expression of typical MSC-

associated immunomodulatory factors such as *PTGS2, IL-6, IDO, PTGES* or *TGFβ (figure 3D-*

H).

The secretome from MSC-CM$_{PPAR\beta/\delta(+)}$ enhances wound healing in CALU-3 lung

epithelial cells in an ANGPTL4-dependent manner

Other studies have highlighted the role of a PPARβ/**δ** agonist [18], and ANGPTL4 in

promoting repair in other tissues [29, 32]. It has also been shown that ANGPTL4 can play a

significant role in proliferation and migration [28]. To determine if the increased levels of

ANGPTL4 present in MSC-CM$_{PPAR\beta/\delta(+)}$ played a role in the enhanced pro-reparative capacity,

we used an anti-ANGPTL4 neutralising-antibody to eliminate the functional capacity of

ANGPTL4 in the MSC-CM (figure 4A). Importantly, ANGPTL4-neutralisation but not the

isotype control antibody, abrogated the enhanced wound closure observed in the MSC-

CM$_{PPAR\beta/\delta(+)}$ group (figure 4B-C). Addition of the ANGPTL4 neutralising antibody interfered

with ANGPTL4 protein levels as quantified by ELISA (figure 4D).

Ligands and co-activators of PPARβ/δ are increased, while co-repressors of PPARβ/δ are

decreased, in ARDS patients

Thus far, our data has shown that PPARβ/**δ** agonism leads to enhanced cytoprotective effects

of MSC-CM. Previously, it was deduced that MSC-CM derived from ARDS-serum licensed

MSCs has superior therapeutic effects in a pre-clinical model of ALI [12]. To ensure translational relevance of our findings from this study in human conditions such as ARDS, a publicly available metabolomics dataset (Study ID: ST000042; doi: 10.21228/M8SG64) was mined. From this metabolomics dataset we identified that PPARβ/δ ligands are more prevalent in ARDS. Higher levels of linoleate, palmitoleic acid and arachidonate were present in the bronchoalveolar lavage fluid (BALF) from ARDS patients, in comparison to healthy controls (figure 5A). Using a publicly available RNA sequencing dataset (GEO accession no.: GSE185263), it was shown that several PPARβ/δ-associated co-activators that were increased in the blood of ARDS patients when compared to Non-ARDS controls (figure 5B). This included a significant increase in *NCOA1* and *NCOA3,* a trend toward increased *PPARGC1A* and *NCOA2,* and no difference in *CREBBP* (figure 5C-G). Along with this, several PPARβ/δ-associated co-repressors were decreased in the blood of the same ARDS patients (figure 5B). This included a significant reduction in *SIRT1, NCOR1* and *NCOR2* (figure 5H-J). Given that *ANGPTL4* was significantly upregulated in response to PPARβ/δ agonism in our own sequencing dataset (figure 2), levels of ANGPTL4 were further assessed upon MSC-exposure to ARDS patient serum. In comparison to the PPARβ/δ agonist, exposure of MSCs to ARDS patient serum significantly increased MSC secretion of ANGPTL4. Addition of both the PPARβ/δ agonist and ARDS patient serum further enhanced MSC production of ANGPTL4 (figure 5K). Interestingly, addition of the PPARβ/δ antagonist could not block the ARDS serum induction of ANGPTL4 secretion by MSCs. This is largely down to PPARβ/δ having multiple binding-sites [33], whereby both agonists and antagonists can bind simultaneously; with agonism out-competing antagonism (supplementary figure 2).

MSC-CM~PPARβ/δ(+)~ provides enhanced lung endothelial barrier function in ALI

Next, we investigated the protective effects of MSC-CM$_{PPAR\beta/\delta(+)}$ in a preclinical model of LPS

induced-ALI. First we set out to determine the number of MSCs that could provide a

therapeutic effect in a mouse model of ALI, comparing two different doses of MSCs.

Administration of high dose MSCs ($1x10^5$) led to significant decrease in TNFα, IL-1β and IL-

6 in the BALF of ALI mice, while low dose MSCs (5×10^4) could only decrease TNFα and

led to increased IL-6 in the BALF (supplementary figure 3). Following this, MSC-CM that was

generated from the equivalent of the low dose was used; with the idea that using a low dose

would allow us to detect any enhanced therapeutic efficacy following PPARβ/**δ** agonism or

antagonism. This showed that MSC-CM$_{PPAR\beta/\delta(+)}$ but not MSC-CM$_{PPAR\beta/\delta(-)}$ significantly

improved the therapeutic efficacy of the non-efficacious dose of MSC-CM. Lungs of mice that

received MSC-CM$_{PPAR\beta/\delta(+)}$ exhibited enhanced vascular and epithelial barrier function

measured using Evans Blue dye administration (figure 6A, B). Although modest, this result

was statistically powered. This effect was somewhat impacted by pre-exposing the MSC-CM

to anti-ANGPTL4 neutralising antibody, but this was not significantly different from the MSC-

CM$_{PPAR\beta/\delta(+)}$ group (figure 6A, B). Indeed MSC-CM contains other growth factors and soluble

mediators that may also play a role in promoting lung repair and we cannot rule out the

possibility that other factors may contribute the protective effects observed. MSC-CM$_{PPAR\beta/\delta(+)}$

did not significantly reduce the level of TNFα or IL-6 in the BALF of LPS-exposed mice

(figure 6C-D). Administration of MSC-CM$_{PPAR\beta/\delta(+)}$ did not decrease weight loss or clinical

score in LPS-exposed mice (figure 6E-F).

ARDS-licensed MSC-CM$_{PPAR\beta/\delta(+)}$ provide protection in ALI in an ANGPTL4 dependent

manner

Given our findings that MSC-CM$_{PPAR\beta/\delta(+)}$ could enhance vascular and epithelial barrier

function but could not decrease BALF pro-inflammatory cytokines or reduce weight loss and

clinical score, we sought to investigate if we could increase the efficacy of MSC-CM.

Interestingly exposure to ARDS patient serum, along with PPARβ/δ agonism, allowed for a

10-fold increased induction of ANGPTL4 in MSC-CM compared to PPARβ/δ agonist alone

(figure 5K). Other proteins examined including IL-6, IL-8, VEGF, IL-1β and MIF

(supplementary figure 4) were not increased in MSCs that had been exposed to ARDS patient

serum, along with PPARβ/δ agonism. We hypothesised that ARDS-licensed MSC-

CM$_{PPAR\beta/\delta(+)}$ would provide greater protection in ALI. In this case, the combination of ARDS

serum and PPARβ/δ agonism led to enhanced therapeutic effects with reduction in TNFα and

IL-6 in the BALF (figure 7A-B). Importantly, anti-ANGPTL4 reduced the anti-inflammatory

effects mediated by the ARDS-licensed MSC-CM$_{PPAR\beta/\delta(+)}$ (figure 7A-B).

In line with this, there was a significant reduction in percentage weight-loss in mice exposed

to ARDS-licensed MSC-CM$_{PPAR\beta/\delta(+)}$ (figure 7C), and again, this effect was abrogated in the

presence of the anti-ANGPTL4-neutralising antibody (figure 7C). Interestingly, we saw a non-

significant trend toward a reduced clinical score in mice exposed to ARDS-licensed MSC-

CM$_{PPAR\beta/\delta(+)}$ (figure 7D). ANGPTL4-neutralisation appeared to abrogate this effect (figure 7D).

DISCUSSION

There are two primary functions of MSCs in the context of cell therapy: immunomodulation

and promotion of repair [1]. It is known that the environment can largely dictate MSC response

[7, 11, 34]. In the context of ARDS, the complex patient microenvironment; containing

inflammatory-agents, microorganisms and FFAs, can notably impact MSC functionality [10, 35]. We previously highlighted that higher levels of inflammation in the ARDS patient microenvironment can promote VEGF-driven repair [12]. Abreu *et al.* (2020) investigated the detrimental impact of fungal toxins present in the lung on MSC viability and showed a drastic increase in MSC cell-death in response to this microenvironmental element [7]. However, there has been little focus on how the FFAs in the ARDS patient microenvironment may impact MSCs [36]. FFAs are known natural ligands of PPARβ/δ [37-39]. Importantly, studies have already shown that modulation of this receptor can alter MSC functionality [17, 20, 21, 23, 25]. Given this, we wanted to investigate the functional impact of PPARβ/δ-modulation on both primary functions of MSCs; immunomodulation and/or pro-reparative capacity.

In the case of pro-reparative capacity, PPARβ/δ-activation enhanced MSC-CM mediated wound repair in CALU-3 lung epithelial cells (figure 1B, C). A time-lapse experiment, using a mitomycin C proliferation inhibitor, highlighted that this was likely a combination effect of both migration and proliferation (figure 1D, E). In the context of promoting tissue repair, MSC secretion of VEGF [40-43], and HGF [44, 45] among other factors have been shown to play key roles. However, PPARβ/δ-activation did not lead to differential expression of VEGF or HGF. Interestingly, ANGPTL4 was significantly increased in PPARβ/δ agonised MSCs. It has already been shown in several studies that ANGPTL4 can promote both migration and proliferation in endothelial cells [28, 29]. Ziveri *et al.* (2024) recently published a comprehensive study comparing the role of ANGPTL4 cytoprotection in skin and brain endothelial cells. ANGPTL4 in the brain endothelium protects against bacterial sepsis, and addition of ANGPTL4 to the skin endothelium, which naturally has lower levels of ANGPTL4, leads to protection against bacterial sepsis [27]. ANGPTL4 has also been shown to improve cardiac function in a mouse model of myocardial infarction [46]. Moreover, human MSCs produced increased levels of ANGPTL4 following co-culture with THP-1 macrophages

in vitro [46]. Based on the identification of increased expression of ANGPTL4 in PPARβ/δ

agonised MSCs in our RNA-seq data set, we used an anti-ANGPTL4 antibody to neutralise the

ANGPTL4 in the MSC secretome, before addition to the CALU-3 lung epithelial cells (figure

4B-D). This abrogated the enhanced wound closure we had originally observed in our MSC-

$CM_{PPARβ/δ(+)}$ group.

We identified increased levels of PPARβ/δ ligands and co-activators in ARDS patient

samples using publicly available datasets. Exposure to ARDS patient serum significantly

increased ANGPTL4 levels produced by MSCs. A combination of MSC exposure to ARDS

patient serum and a PPARβ/δ agonist further increased ANGPTL4 levels. This proves that

MSCs are receptive to FFAs/other PPARβ/δ ligands presents in the ARDS microenvironment.

This aligns to our previous study where we identified that human BM-MSCs produce

significantly increased levels of ANGPTL4 in response to the FFA palmitate [14].

To enhance the clinical relevance of our findings, we showed that $MSC-CM_{PPARβ/δ(+)}$

mediated protective effects in ALI significantly enhancing vascular and epithelial barrier

integrity *in vivo*. Indeed, other studies have demonstrated that MSC-CM can mediate

therapeutic effects *in vivo* in ALI models [47] including our own study [12]. Our previous study

showed that MSC-CM generated from the equivalent of $5x10^4$ MSCs was ineffective in the

treatment of ALI; but licensing $5x10^4$ MSCs with ARDS patient serum enhanced the

therapeutic efficacy [12]. Using this approach of MSC-CM generated from the non-efficacious

dose of $5x10^4$ MSCs, we investigated the potential for enhanced efficacy of MSC-CM

generated from $5x10^4$ MSCs that had been exposed to a PPARβ/δ agonist or antagonist. MSC-

$CM_{PPARβ/δ(+)}$ provided enhanced efficacy over and above $MSC-CM_{PPARβ/δ(-)}$ in a pre-clinical

model of ALI. While there was a significant difference in the capacity for $MSC-CM_{PPARβ/δ(+)}$

in enhancing barrier function, addition of the ANGPTL4-neutralising antibody failed to

significantly impair this effect. This may be associated with growth factors or mediators other

than ANGPTL4 present in the MSC-CM$_{PPAR\beta/\delta(+)}$ that may be contributing to the protective

effects observed. Our study demonstrated that ARDS patient serum induced significantly

higher levels of ANGPTL4 in MSC-CM and that combination with a PPARβ/**δ** agonist further

increased it. While MSC-CM$_{PPAR\beta/\delta(+)}$ did not significantly reduce proinflammatory cytokines

in the BALF, ARDS licensed MSC-CM$_{PPAR\beta/\delta(+)}$ reduced TNFα and IL-6 in the BALF of ALI

mice. By combining the PPARβ/**δ** agonist and licensing with ARDS patient serum, we created

MSC-CM with greater immunosuppressive capacity when compared to MSC-CM$_{PPAR\beta/\delta(+)}$

alone. This is likely associated with the ~10-fold increase in ANGPTL4 levels when using the

combined licensing approach as addition of the ANGPTL4 neutralising antibody reversed the

trend. The limitation to these findings is the small numbers of mice (n=3/group) included in

the study for the ARDS licensed MSC-CM$_{PPAR\beta/\delta(+)}$ treatment group. This was associated with

a limited volume of ARDS patient serum available for the study.

Here we provide novel findings to demonstrate that human BM-MSCs are responsive

to FFAs (ligands that bind PPARβ/**δ**) present in the ARDS patient environment, leading to

significantly increased levels of ANGPTL4 production. For the first time, we show that MSC-

derived ANGPTL4 plays an important role in promoting epithelial repair and enhancing

vascular barrier integrity. As most published studies have focused on mouse MSCs, we have

also addressed the gap in the knowledge of how PPARβ/**δ** influences human BM-MSC

cytoprotective and immunosuppressive functions. In this study, focusing on ALI, we did not

detect any significant positive or negative effects associated with PPARβ/**δ** antagonism in

hBM-MSCs. However, PPARβ/**δ** agonism enhanced MSC-CM promotion of wound healing

in vitro and vascular and barrier integrity *in vivo*. Furthermore, the combined approach of

PPARβ/**δ** agonism and licensing with ARDS patient serum produced an MSC-CM that could

suppress pro-inflammatory cytokines in the BALF of ALI mice.

Declarations

This manuscript has been uploaded to Bioxriv as a pre-print.

Author Contributions

C.T. performed research, data analysis, and study design and wrote the manuscript. M.D., S.R., E.V., E.J., S.B. and C.M. provided expertise and or performed research and data analysis. B.McN and J.G.L. provided patient samples and data for the study. C.C.D.S and J.G.L. contributed to study design. K.E. designed and supervised the study and wrote the manuscript. All authors approved the final manuscript.

Funding

This research was supported by a Science Foundation Ireland Award to K.E. and J.G.L (20/FFP-A/8948). This publication has emanated from research supported in part by the National Irish Covid Biobank, through which our patient samples were collected and stored.

Competing interests

The authors declare no competing interests.

Ethical Approval and HPRA Compliance

Ethical approval was granted by the Maynooth University research ethics committee (BRESC-2022-2453953) granted 17th February 2022 under the title: Cell based therapeutics for acute respiratory distress syndrome for the animal studies and by Galway University Hospitals Research Ethics Committee and by the Maynooth University research ethics committee (BSRESC-2022-2482563) granted 14th October 2022 under the title: Human serum

Sepsis/ARDS MSC. HPRA approval was granted under the project authorisation(s)

AE19124/P031/P037.

RoosterBio (company where the human BM-MSCs were purchased) has confirmed that there

was initial ethical approval for collection of human cells, and that the donors had signed

informed consent. RoosterBio state "RoosterBio sources commercially available in vitro

research only human bone marrow aspirate from qualified donors. All human bone marrow

aspirate collections are from healthy adult consented donors. Collection protocols and the

donor-informed consent document are approved by an Institutional Review Board (IRB)"

Use of AI

The authors declare that they have not used AI-generated work in this manuscript.

Data availability statement

The RNA-sequencing dataset generated in our lab is available through GEO accession no.:

GSE281162. The other publicly available data sets are available online as outlined in the

Methods section. The remaining data supporting the findings of this study are available upon

request from the corresponding author.

Acknowledgements

The authors are grateful to the ARDS patients who donated the samples used in this study. We

would like to thank Rebecca Kerrigan, Deirdre Daly and Gillian O'Meara for the outstanding

care of the animals used in this study. Schematic figures contained in this manuscript have been

created using biorender.com.

Table 1: DEG functions in MSC_{PPARβ/δ(+)} group.

Gene name	Up/Down	Involved in
plin2	Up	Lipid metabolism
angptl4	Up	Lipid metabolism and angiogenesis
fam156A	Up	Enables protein binding
cpt1A	Up	Lipid metabolism
cat	Up	Protection against oxidative stress
nadk2	Up	Protection against oxidative stress
tmed6	Up	Trafficking and transport
ldhal6B	Up	Lactate metabolism
maged4	Down	Cell cycle regulation
fam157B	Down	Non-coding

502 **Table 2: DEG functions in MSC_{PPARβ/δ(-)} group.**

Gene name	Up/Down	Involved in
ikbkgp1	Up	NFκB signalling
snrpgp2	Up	RNA processing
linc01159	Up	Cell differentiation
mir3648-1	Up	Cell differentiation
rarres1	Up	Cell differentiation & retinoid signalling
malat1	Up	Cell differentiation
rnf217-as1	Up	Cell differentiation

rgpd5	Down	GTPase regulation
pfn1p3	Down	Cell motility/migration
mir3936hg	Down	Proliferation and apoptosis
u2af1	Down	ROS production/oxidative stress
tsnaxip1	Down	RNA processing and transport

REFERENCES

1.   Faircloth, T.U., et al., *Vascular endothelial growth factor secretion and immunosuppression are distinct potency mechanisms of human bone marrow mesenchymal stromal cells.* Stem Cells, 2024. **42**(8): p. 736-751.

2.   Matthay, M.A. and G.A. Zimmerman, *Acute lung injury and the acute respiratory distress syndrome: four decades of inquiry into pathogenesis and rational management.* Am J Respir Cell Mol Biol, 2005. **33**(4): p. 319-27.

3.   Yang, X., et al., *Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study.* Lancet Respir Med, 2020. **8**(5): p. 475-481.

4.   Kaffash Farkhad, N., et al., *Mesenchymal stromal cell therapy for COVID-19-induced ARDS patients: a successful phase 1, control-placebo group, clinical trial.* Stem Cell Res Ther, 2022. **13**(1): p. 283.

5.   Bowdish, M.E., et al., *A Randomized Trial of Mesenchymal Stromal Cells for Moderate to Severe Acute Respiratory Distress Syndrome from COVID-19.* Am J Respir Crit Care Med, 2023. **207**(3): p. 261-270.

6.   Gorman, E., et al., *Repair of acute respiratory distress syndrome by stromal cell*
administration (REALIST) trial: A phase 1 trial. EClinicalMedicine, 2021. **41**: p.
101167.

7.   Abreu, S.C., et al., *Differential effects of the cystic fibrosis lung inflammatory*
environment on mesenchymal stromal cells. Am J Physiol Lung Cell Mol Physiol,
2020. **319**(6): p. L908-L925.

8.   Hawthorne, I.J., et al., *Human macrophage migration inhibitory factor potentiates*
mesenchymal stromal cell efficacy in a clinically relevant model of allergic asthma.
Mol Ther, 2023. **31**(11): p. 3243-3258.

9.   Corbett, J.M., et al., *Cyclosporine A and IFNγ licencing enhances human*
mesenchymal stromal cell potency in a humanised mouse model of acute graft versus
host disease. Stem Cell Res Ther, 2021. **12**(1): p. 238.

10.   Dunbar, H., et al., *The Inflammatory Lung Microenvironment; a Key Mediator in*
MSC Licensing. Cells, 2021. **10**(11).

11.   Rolandsson Enes, S., et al., *Healthy versus inflamed lung environments differentially*
affect mesenchymal stromal cells, Eur Respir J, 2021. **58**(4).

12.   Tunstead, C., et al., *The ARDS microenvironment enhances MSC-induced repair via*
VEGF in experimental acute lung inflammation. Mol Ther, 2024. **32**(10): p. 3422-
3432.

13.   Boland, L., et al., *IFN-γ and TNF-α Pre-licensing Protects Mesenchymal Stromal*
Cells from the Pro-inflammatory Effects of Palmitate. Mol Ther, 2018. **26**(3): p. 860-
873.

14.   Tunstead, C., Bitterlich, L.B.*, Ankrum, J.A., Hogand, A.E. and English, K.,
Palmitate enhances MSC immunomodulation of human macrophages via the
ceramide/CCL2 axis in vitro. Stem Cell Research and Therapy, 2025. 16(435).

15.   Liu, Y., et al., *The Role of PPAR-δ in Metabolism, Inflammation, and Cancer: Many*

548   *Characters of a Critical Transcription Factor.* Int J Mol Sci, 2018. **19**(11).

549   16.   Kota, B.P., T.H. Huang, and B.D. Roufogalis, *An overview on biological mechanisms*

550   *of PPARs.* Pharmacol Res, 2005. **51**(2): p. 85-94.

551   17.   Lim, J. and E.K. Park, *Effect of fibroblast growth factor-2 and retinoic acid on*

552   *lineage commitment of bone marrow mesenchymal stem cells.* Tissue Eng Regen

553   Med, 2016. **13**(1): p. 47-56.

554   18.   Park, J.R., et al., *Effects of Peroxisome Proliferator-Activated Receptor-δ Agonist on*

555   *Cardiac Healing after Myocardial Infarction.* PLoS One, 2016. **11**(2): p. e0148510.

556   19.   Carty, F., et al., *IFN-γ and PPARδ influence the efficacy and retention of multipotent*

557   *adult progenitor cells in graft vs host disease.* Stem Cells Transl Med, 2021. **10**(11):

558   p. 1561-1574.

559   20.   Heck, B.E., et al., *PPAR-δ Agonist With Mesenchymal Stem Cells Induces Type II*

560   *Collagen-Producing Chondrocytes in Human Arthritic Synovial Fluid.* Cell

561   Transplant, 2017. **26**(8): p. 1405-1417.

562   21.   Luz-Crawford, P., et al., *PPARβ/δ directs the therapeutic potential of mesenchymal*

563   *stem cells in arthritis.* Ann Rheum Dis, 2016. **75**(12): p. 2166-2174.

564   22.   Contreras-Lopez, R.A., Elizondo-Vega, R., Torres, M.J. *et al*, *PPARβ/δ-dependent*

565   *MSC metabolism determines their immunoregulatory properties.* 2020: p. 11423

566   23.   Kim, D.H., et al., *PPAR-δ agonist affects adipo-chondrogenic differentiation of*

567   *human mesenchymal stem cells through the expression of PPAR-γ.* Regen Ther, 2020.

568   **15**: p. 103-111.

569   24.   Nernpermpisooth, N., et al., *PPARβ/δ Is Required for Mesenchymal Stem Cell*

570   *Cardioprotective Effects Independently of Their Anti-inflammatory Properties in*

571   *Myocardial Ischemia-Reperfusion Injury.* Front Cardiovasc Med, 2021. **8**: p. 681002.

25.   Sarre, C., et al., *PPARβ/δ priming enhances the anti-apoptotic and therapeutic*

properties of mesenchymal stromal cells in myocardial ischemia-reperfusion injury.

Stem Cell Res Ther, 2022. **13**(1): p. 167.

26.   Djouad, F., et al., *PPARβ/δ: A master regulator of mesenchymal stem cell functions.*

Biochimie, 2017. **136**: p. 55-58.

27.   Ziveri, J., et al., *Angiopoietin-like 4 protects against endothelial dysfunction during*

bacterial sepsis. Nat Microbiol, 2024. **9**(9): p. 2434-2447.

28.   Goh, Y.Y., et al., *Angiopoietin-like 4 interacts with integrins beta1 and beta5 to*

modulate keratinocyte migration. Am J Pathol, 2010. **177**(6): p. 2791-803.

29.   Zhan, W., et al., *ANGPTL4 attenuates palmitic acid-induced endothelial cell injury by*

increasing autophagy. Cell Signal, 2022. **98**: p. 110410.

30.   An, Q., et al., *In vitro effects of mitomycin C on the proliferation of the non-small-cell*

lung cancer line A549. Int J Clin Exp Med, 2015. **8**(11): p. 20516-23.

31.   Neubauer, E., et al., *Comparative evaluation of three proliferation markers, Ki-67,*

TOP2A, and RacGAP1, in bronchopulmonary neuroendocrine neoplasms: Issues and

prospects. Oncotarget, 2016. **7**(27): p. 41959-41973.

32.   Huang, W., et al., *ANGPTL4 induces Kupffer cell M2 polarization to mitigate acute*

rejection in liver transplantation. Sci Rep, 2025. **15**(1): p. 986.

33.   Batista, F.A., et al., *Structural insights into human peroxisome proliferator activated*

receptor delta (PPAR-delta) selective ligand binding. PLoS One, 2012. **7**(5): p.

e33643.

34.   Kornicka, K., J. Houston, and K. Marycz, *Dysfunction of Mesenchymal Stem Cells*

Isolated from Metabolic Syndrome and Type 2 Diabetic Patients as Result of

Oxidative Stress and Autophagy may Limit Their Potential Therapeutic Use. Stem

Cell Rev Rep, 2018. **14**(3): p. 337-345.

35.   Lu, P., et al., *Serum Free Fatty Acid Concentration Predicts ARDS after Off-Pump*

598   *CABG: A Prospective Observational Study.* Lung, 2024. **202**(5): p. 523-532.

599   36.   Boland, L., et al., *Translating MSC Therapy in the Age of Obesity.* Front Immunol,

600   2022. **13**: p. 943333.

601   37.   Grygiel-Górniak, B., *Peroxisome proliferator-activated receptors and their ligands:*

602   *nutritional and clinical implications--a review.* Nutr J, 2014. **13**: p. 17.

603   38.   Mirza, A.Z., I.I. Althagafi, and H. Shamshad, *Role of PPAR receptor in different*

604   *diseases and their ligands: Physiological importance and clinical implications.* Eur J

605   Med Chem, 2019. **166**: p. 502-513.

606   39.   Capelli, D., et al., *Structural basis for PPAR partial or full activation revealed by a*

607   *novel ligand binding mode.* Sci Rep, 2016. **6**: p. 34792.

608   40.   Yang, Y., et al., *Synergism of MSC-secreted HGF and VEGF in stabilising*

609   *endothelial barrier function upon lipopolysaccharide stimulation via the Rac1*

610   *pathway.* Stem Cell Res Ther, 2015. **6**: p. 250.

611   41.   Zisa, D., et al., *Vascular endothelial growth factor (VEGF) as a key therapeutic*

612   *trophic factor in bone marrow mesenchymal stem cell-mediated cardiac repair.*

613   Biochem Biophys Res Commun, 2009. **390**(3): p. 834-8.

614   42.   Yang, Y., et al., *The Vascular Endothelial Growth Factors-Expressing Character of*

615   *Mesenchymal Stem Cells Plays a Positive Role in Treatment of Acute Lung Injury In*

616   *Vivo.* Mediators Inflamm, 2016. **2016**: p. 2347938.

617   43.   Dunbar, H., et al., *The VEGF-Mediated Cytoprotective Ability of MIF-Licensed*

618   *Mesenchymal Stromal Cells in House Dust Mite-Induced Epithelial Damage.* Eur J

619   Immunol, 2025. **55**(1): p. e202451205.

620   44.   Cahill, E.F., et al., *Hepatocyte Growth Factor Is Required for Mesenchymal Stromal*

621   *Cell Protection Against Bleomycin-Induced Pulmonary Fibrosis*. Stem Cells Transl

622   Med, 2016. **5**(10): p. 1307-1318.

623   45.   Kennelly, H., B.P. Mahon, and K. English, *Human mesenchymal stromal cells exert*

624   *HGF dependent cytoprotective effects in a human relevant pre-clinical model of*

625   *COPD*. Sci Rep, 2016. **6**: p. 38207.

626   46.   Cho, D.I., et al., *Antiinflammatory activity of ANGPTL4 facilitates macrophage*

627   *polarization to induce cardiac repair*. JCI Insight, 2019. **4**(16).

628   47.   Curley, G.F., et al., *Effects of intratracheal mesenchymal stromal cell therapy during*

629   *recovery and resolution after ventilator-induced lung injury*. Anesthesiology, 2013.

630   **118**(4): p. 924-32.

Figure Legends

Figure 1: Secretome from PPARβ/δ agonised hBM-MSCs enhanced wound healing in

CALU-3 lung epithelial cells. (A) Experimental Design. A CALU-3 scratch assay was run, in

combination with MSC-CM from naive, PPARβ/δ agonized (+) or antagonized (-) hBM-

MSCs. (B) Images of the wound healing assay at baseline or following 48hrs fixed and stained

with crystal violet dye and (C) quantified using ImageJ software (one-way ANOVA followed

by Tukey's post-hoc test, n=3). White bar is CALU-3 cells in media alone. (D) Percentage (%)

wound healing in scratched CALU-3 cells was recorded at different time points (6, 24 and

48hrs). Mitomycin C (MMC, denoted as circular data points) was added to CALU-3 cells 2hr

prior to the scratch assay. Conditioned medium from naive, PPARβ/δ agonized (+) or

antagonized (-) hBM-MSCs was added immediately after the scratch. Wound closure (%) was

quantified using imageJ software (two-way ANOVA followed by Tukey's post-hoc test, n=3).

(E) Expression of the cell proliferation gene *TOP2A* was analysed by RT-PCR at a 48hr time-

point (one-way ANOVA followed by Tukey's post-hoc test, n=3). Replicates are a

representation of 3 individual hBM-MSC donors. Data is presented as mean ± SEM; *$p<0.05$,

**$p<0.01$, ***$p<0.001$, ****$p<0.0001$.

Figure 2: RNA-Seq identified enrichment of ANGPTL4 in PPARβ/δ agonised hBM-

MSCs. (A) Experimental design: hBM-MSCs exposed to 1uM of PPARβ/δ agonist or

antagonist were sent to Novogene™ for RNA sequencing. (B) Venn diagrams showing the

number of differentially expressed genes (DEGs) in naive, PPARβ/δ agonized (+) or

antagonized (-) hBM-MSCs. (C) Volcano plots generated using Flourish™ software highlight

the most significant DEGs Log10(p-value). Fragments per kilobase of transcript per million

mapped reads (FPKM) was plotted for two known PPARβ/δ target genes in the sequencing

dataset: (D) *ANGPTL4* and (E) *PDK4*. Expression of *ANGPTL4* (F) and *PDK4* (G) in hBM-

MSCs exposed to 1uM of PPARβ/δ agonist or antagonist for 6hrs was confirmed by real-time

RT-PCR (one-way ANOVA followed by Tukey's post-hoc test, n=3). Replicates are a

representation of 3 individual hBM-MSC donors. Data is presented as mean ± SEM; **$p<0.01$,

***$p<0.001$, ****$p<0.0001$.

Figure 3: PPARβ/δ(+) hBM-MSCs show enhanced expression of ANGPTL4, but not

other traditional MSC-associated genes or proteins. Supernatant from hBM-MSCs exposed

to 1uM of PPARβ/δ agonist or antagonist were harvested before cells were collected in TRIzol

for gene expression studies. Protein expression of (A) ANGPTL4, (B) VEGF and (C) MIF

were assessed by ELISA. Gene expression of (D) *PTGS2,* (E) *IL-6,* (F) *IDO,* (G) *PTGES* and

(H) *TGFβ* were assessed by real-time RT-PCR (one-way ANOVA followed by Tukey's post-

hoc test, n=3). Replicates are a representation of 3 individual hBM-MSC donors. Data is

presented as mean ± SEM; *$p<0.05$.

Figure 4: The secretome from PPARβ/δ agonised hBM-MSCs mediates enhanced wound repair via ANGPTL4. (A) Experimental design. Using an ANGPTL4 neutralisation approach, we assessed wound closure of CALU-3 lung epithelial cells. (B) Images taken at baseline or 48hr post scratch. Scratched CALU-3 cells were incubated with isotype control or anti-ANGPTL4 antibody before addition of conditioned medium. Medium only served as the control. (C) % wound closure quantified using ImageJ. (D) Levels of ANGPTL4 produced by naive hBM-MSCs or exposed to 1uM of PPARβ/δ agonist or antagonist in the presence of the isotype control of anti-ANGPTL4 antibody were measured by ELISA (one-way ANOVA followed by Tukey's post-hoc test, n=3). Replicates are a representation of 3 individual hBM-MSC donors. ND denotes not detected. Data is presented as mean ± SEM; **p<0.01, ***p<0.001, ****p<0.0001.

Figure 5: ARDS patients have higher levels of PPARβ/δ ligands and co-activators. (A) A publicly available dataset (Study ID: ST000042; doi: 10.21228/M8SG64) was mined for PPARβ/δ ligands present in the ARDS lung. Levels of Arachidonate, Palmitoleic acid and Linoleate were elevated in ARDS bronchoalveolar lavage fluid compared to healthy controls. (B) An additional RNA-Seq dataset (GEO accession no.: GSE185263), based on ARDS serum, highlighted an increase in PPARβ/δ-associated co-activators in both SARS-CoV-2 ARDS and Classical ARDS patient serum compared to Non-ARDS controls. RNA-Seq data counts for PPARβ/δ-associated co-activator genes (C) *PPARGC1A*, (D) *NCOA1*, (E) *NCOA2*, (F) *NCOA3* and (G) *CREBBP;* and co-repressor genes (H) *SIRT1*, (I) *NCOR1* and (J) *NCOR2* were plotted comparing expression between SARS-CoV-2 ARDS, Classical ARDS and Non-ARDS controls (one-way ANOVA followed by Tukey's post-hoc test, n=6-58). Each data point (A-J) is from an individual ARDS patient, or healthy control. (K) ANGPTL4 production from 3

individual hBM-MSC donors exposed to PPARβ/**δ** agonist (+) or antagonist (-) in the presence

or absence of 20% ARDS patient serum was measured by ELISA. Data is presented as mean

± SEM; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 6: MSC-CM$_{PPARβ/δ(+)}$ supports enhanced endothelial barrier function in ALI. Mice

were exposed to 2mg/kg LPS IT, or PBS control. Some groups received MSC-CM$_{PPARβ/δ(-)}$ or

MSC-CM$_{PPARβ/δ(+)}$ + isotype control or MSC-CM$_{PPARβ/δ(+)}$ + anti-ANGPTL4. (A) Lungs were

harvested and imaged after Evan's Blue Dye administration highlighting increased vascular

permeability (visible blue dye). (B) Quantification of the amount of Evan's Blue dye leakage

from the lungs of mice exposed to LPS +/- treatments of interest (n=5/group, or n=3 for PBS

control group). Levels of pro-inflammatory cytokines in the bronchoalveolar lavage fluid (C)

TNFα and (D) IL-6 were assessed by ELISA (one-way ANOVA followed by Tukey's post-

hoc test, n=4-6/group). (E) % weigh loss and (F) Clinical scores (n=4-6/group) were recorded

over the course of the model. Data points are a representation of individual mice. Data is

presented as mean ± SEM; *p<0.05, **p<0.01.

Figure 7: ARDS-licensed MSC-CM$_{PPARβ/δ(+)}$ provide protection in ALI via ANGPTL4.

Mice were exposed to 2mg/kg LPS IT, or PBS control. Some groups received ARDS licensed

MSC-CM$_{PPARβ/δ(+)}$ + isotype control or ARDS-licensed MSC-CM$_{PPARβ/δ(+)}$ + anti-ANGPTL4.

Levels of bronchoalveolar lavage fluid pro-inflammatory cytokines (A) TNFα and (B) IL-6

were assessed by ELISA (one-way ANOVA followed by Tukey's post-hoc test, n=3-6/group).

(C) % weigh loss and (D) Clinical scores (n=3-6/group) were recorded over the course of the

model. Data points represent individual mice. Data is presented as mean ± SEM; *p<0.05,

***p<0.001.

Graphical Abstract

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7