Parametric models for predicting nonstationary spike-spike correlations with local field potentials

Zeinab Tajik-Mansouri[1,4*], James P. Dion Jr.[2], Monty A. Escabí[1-4], and Ian H. Stevenson[1,2,4]

[1] *Department of Biomedical Engineering, University of Connecticut*
[2] *Department of Psychological Sciences, University of Connecticut*
[3] *Department of Electrical and Computer Engineering, University of Connecticut*
[4] *Connecticut Institute for Brain and Cognitive Science, University of Connecticut*

* Corresponding Author: zeinab.tajik@uconn.edu

Abstract

Correlations between the spiking of pairs of neurons are often used to study the brain's representation of sensory or motor variables and neural circuit function and dysfunction. Previous statistical techniques have shown how time-averaged spike-spike correlations can be predicted by the time-averaged relationships between the individual neurons and the local field potential (LFP). However, spiking and LFP are both nonstationary, and spike-spike correlations have nonstationary structure that cannot be accounted for by time-averaged approaches. Here we develop parametric models that predict spike-spike correlations using a small number of LFP-based predictors, and we then apply these models to the problem of tracking changes in spike-spike correlations over time. Parametric models allow for flexibility in the choice of which LFP recording channels and frequency bands to use for prediction, and coefficients directly indicate which LFP features drive correlated spiking. Here we demonstrate our methods in simulation and test the models on experimental data from large-scale multi-electrode recordings in the mouse hippocampus and visual cortex. In single time windows, we find that our parametric models can be as accurate as previous nonparametric approaches, while also being flexible and interpretable. We then demonstrate how parametric models can be applied to describe nonstationary spike-spike correlations measured in sequential time windows. We find that although the patterns of both cortical and hippocampal spike-spike correlations vary over time, these changes are, at least partially, predicted by models that assume a fixed spike-field relationship. This approach may thus help to better understand how the dynamics of spike-spike correlations are related to functional brain states. Since spike-spike correlations are increasingly used as features for decoding external variables from neural activity, these models may also have the potential to improve the accuracy of adaptive decoders and brain machine interfaces.

Introduction

Correlations between the spiking activities of pairs of neurons (spike-spike correlations) are key indicators of neural processing (Eggermont, 1990; deCharms and Merzenich, 1996; Salinas and Sejnowski, 2001; Cohen and Kohn, 2011; Panzeri et al., 2022) and dysfunction (Uhlhaas and Singer, 2006). Correlations can indicate similar stimulus-response properties (signal correlations), but can also occur when neurons have shared relationships to intrinsic variables such as local field potentials (LFP). Distinguishing between signal correlations and other shared membrane fluctuations (noise correlations) allows researchers to better predict neural responses and better decode external variables from neural responses (Stevenson et al., 2012). However, one challenge in interpreting neural correlations is that they are typically not fixed but change over time due to changes in the firing rates of the individual neurons and broader changes in stimuli, task, or attentional state (Murthy and Fetz, 1996; Hatsopoulos et al., 1998; Baker et al., 2001; Cohen and Maunsell, 2009; Canolty et al., 2012; Mizuseki and Buzsaki, 2014). Here we aim to develop

46 statistical models to better predict and track time-varying spike-spike correlations by accounting for the
47 relationships between spiking and simultaneously-recorded local field potentials (LFP).
48
Measuring spike-spike correlations and interpreting their causes can be statistically challenging
(Stevenson et al., 2008; Grün, 2009; Gerkin et al., 2013; Harrison et al., 2013). Spike-spike correlations
may be due to synaptic connections between the observed neurons and other direct physical links (Moore
et al., 1970; Fetz et al., 1991; Swadlow and Gusev, 2001), as well as, indirect effects such as shared
synaptic input from other neurons (Swadlow et al., 1998; Trepka et al., 2022) or otherwise correlated input
currents (Haider et al., 2016). When measuring correlations in short time windows or when the firing rates
of the neurons are low, the number of coincident spikes is small and estimates of correlation have high
uncertainty. Additionally, correlations between neurons vary substantially from one pair of neurons to the
next – in both the strength of the correlations and their timescales. Here we propose one strategy for
modeling these noisy, nonstationary correlations based on parametric models that use features of the LFP
as predictors. The relationship between spikes and LFP is complex (Buzsáki et al., 2012; Pesaran et al.,
2018; Watson et al., 2018), but broadly speaking, spikes are often coupled to LFPs and occur at specific
phases of LFP oscillations. At the same time, spike-spike correlations often have strong oscillatory pattern,
and spike-spike correlations appear to be related to the spike-field relationships of the individual neurons
(Goldberg et al., 2004; Denker et al., 2011). As LFP spectra change due to brain state, stimuli, and
behavior, we expect some of these changes to translate into predictable changes in spike-spike correlations
over time.
66
Many previous studies have used the LFP to predict the activity of single neurons (Harris et al., 2003;
Canolty et al., 2010; Kelly et al., 2010), and these studies often find that observed spike-spike correlations
are consistent with the model predictions (Zhou et al., 2015; Cui et al., 2016). By modeling the observed
correlations directly, previous studies have also developed extensive methods for estimating high-
dimensional correlations (Yatsenko et al., 2015; Kass et al., 2023; Nejatbakhsh et al., 2023), separating
stimulus and noise correlations (Gawne and Richmond, 1993; Josić et al., 2009; Sadeghi et al., 2019;
Keeley et al., 2020), testing for the presence of fast correlations (Amarasingham et al., 2012; Spivak et
al., 2022) and putative synapses (Barthó et al., 2004; Kobayashi et al., 2019; Ren et al., 2020), or detecting
changes in correlations over time (Lepage et al., 2013). Here we focus on direct, model-based descriptions
of the spike-spike correlations that account for the contribution of time-varying LFP. Data analytic tools
that can account for the detailed contribution of LFP to spike-spike correlations may be useful in allowing
neuroscientists to better understand how and why these correlations change over time and to identify the
specific contributions of different anatomical sources and frequency bands. Since LFP-driven correlations
are often present alongside stimulus-driven correlations or fast synaptic effects, these data analysis tools
may also have the potential to lead to improvements in decoding and more accurate detection of synaptic
connections.
83
Our aim in this work is to predict cross-correlations between spiking neurons based on their individual
spike-LFP coupling. Here we first introduce a generalized bilinear modeling framework that can take
covariates derived from LFP and predict the observed correlations. We use simulations to illustrate the
basic framework and then apply our methods to experimental data from pairs of neurons recorded in mouse
hippocampus and cortex. Since the LFP covariates can take many forms and can be based on either single
or multiple recording sites, we compare models with different types of covariates and with different
optimization strategies. We also compare our parametric models with previously-developed non-
parametric methods that predict time-averaged spike-spike correlations based on the spike-triggered

average LFP (STA). After applying our methods to time-averaged data, we extend our parametric model framework to the case where both correlations and covariates are calculated over multiple short-time windows. We find that, due to time-varying LFP statistics, the predictions of our models change dynamically over time and partially account for the observed nonstationarity in spike-spike correlations.

Methods

Predicting cross-correlograms with LFP features

Given the binned spiking of the two neurons n_1 and n_2 over time, the time-averaged cross-correlogram is given by

$$c(\tau) = \sum_t n_1(t)n_2(t + \tau)$$

Our goal here is to develop predictions $\hat{c}(\tau)$ based on observations of the local field potential y. In previous work, Halliday et al. (1995) developed an STA Method for constructing predictions based on the concept of the partial correlation or partial spectra. With this approach, we first estimate the spike-triggered average (STA) local field potential for each neuron

$$sta(\tau) = \frac{1}{N}\sum_t n(t)y(t + \tau)$$

where N is number of spikes. The method of partial spectra then constructs the prediction in the frequency domain by taking the cross-spectrum of the STAs for each neuron and whitening it by the spectrum of the LFP Y.

$$\hat{c}(\tau) = 1 + F^{-1}\left(\frac{STA_1(f)STA_2^*(f)}{Y^2(f)}\right)$$

Here STA_1 and STA_2 denote the Fourier transforms of the STAs for the two neurons, and Y denotes the Fourier transform of the LFP computed over a common set of frequencies f. F^{-1} denotes the inverse Fourier transform and $\cdot^*$ denotes complex conjugation. Although this model often fits well in practice (Goldberg et al., 2004; Middleton et al., 2012), predictions can sometimes go wrong, especially when the firing rates of the neurons are low or when the LFP power is low at certain frequencies.

Here we consider alternative approaches that take two general forms: an indirect model-based approach and a direct model-based approach. The indirect approach follows the STA Method and fits the correlogram by first describing the rates of the individual neurons λ_1 and λ_2 based on their relationship with the LFP. Then, in a second stage, we use the single-neuron predictions to construct a prediction of the correlation

$$\hat{c}(\tau) = \sum_t \lambda_1(t)\lambda_2(t + \tau)$$

133   This indirect strategy has been used by several previous studies to understand to what extent rate models
134   of individual neurons and populations of neurons can explain the observed correlations (Cui et al., 2016).
135   In the special case where the rate models are linear, the full prediction can be described in closed form.
136   Suppose we represent the LFP using a set of covariates X we have $\lambda_1(t) = x^T(t)\beta_1$ and $\lambda_2(t) = x^T(t)\beta_2$,
137   with coefficients given by

139
$$\beta = (X^TX)^{-1}X^Tn$$

141   and the correlogram prediction given by

143
$$\hat{c}_{linear}(\tau) = \beta_1^T C(\tau)\beta_2$$

where C is a cross-covariance matrix $\sum_t x^T(t)x(t+\tau)$. The LFP covariates X can take different forms
(see below) and can include, for example, band-pass filtered LFP or analytic signals at one or more
recording sites.

In practice, the linear rate estimates λ are poor descriptions of nonlinear spiking activity, so some
additional modeling is required to make $\hat{c}$ a better match of the observed correlation. In the results here
(see below), we consider the specific case where we start with $\lambda_i(t) = x^T(t)\beta_i$, and the correlogram
prediction is then modified by an additional output nonlinearity. We use Poisson regression,

154
$$f_{indirect}(\tau) = \exp(\gamma_0 + \gamma_1 \hat{c}_{linear}(\tau))$$

and assume that the number of coincident counts observed in the correlogram is distributed following
$c(\tau) \sim Poisson(f(\tau))$. Our indirect model-based approach thus works by first fitting linear models to the
rates with coefficients β, then, in a second step, fitting an output nonlinearity with coefficients γ to better
match the correlogram. As with the STA method, this approach is based on first describing the
relationships between each of the neurons and the LFP. The βs in our model represent a type of whitened
spike-triggered average of the predictors X. However, with this approach we use the time-domain rather
than frequency-domain. Additionally, we can compare the impact of different choices for X, and, if
needed, use tools from regression and model selection to prevent overfitting.

We also consider a direct approach for cases where we may not need or want descriptions of the individual
neurons or their rates over time. In this case, we fit a generalized bilinear model (GBLM)

168
$$f_{direct}(\tau) = \exp(\alpha_1^T C(\tau)\alpha_2)$$

170   where rather than predicting spike-LFP coupling for each neuron (with β_1 and β_2), we aim to reconstruct
171   the spike-spike correlation directly by using coefficients $\alpha = \{\alpha_1, \alpha_2\}$. Here we optimize by coordinate
172   ascent – iteratively fitting α_1 with fixed α_2 then fitting α_2 with fixed α_1 – on the log likelihood, assuming
173   correlogram observations are Poisson distributed with mean f_{direct}.

175   To evaluate the accuracy of model fits we use the pseudo-R^2

177
$$R^2 = \frac{D_0 - D}{D_0}$$

179 Where $D. = 2(\mathcal{L}_s - \mathcal{L}.)$ denotes the model deviance for the null (D_0) and fitted (D) models with log-
180 likelihood $\mathcal{L} \propto \sum_\tau (c(\tau) \log f(\tau) - f(\tau))$. For the null model, $f(\tau)$ is a constant $\overline{c(\tau)}$, and $\mathcal{L}_s$ denotes the
181 log-likelihood of the saturated model with $f(\tau) = c(\tau)$.

LFP Representations

185 One potential advantage of both the indirect and direct modeling approaches described here is that we can
186 choose an appropriate, compact representation of the LFP to use for our predictor X. Previous work has
187 found that neurons often couple to specific LFP frequency bands and phase-lock to different phases within
188 each band. This observation motivates a generic representation of the form

$$x_i(t) = y(t) * F_i(t)$$

where F_i denotes a bandpass filter applied to the LFP $y(t)$, and we choose a set of filters $i \in \{1 \cdots p\}$ to
account for the sensitivity of the two neurons whose spike-spike correlation we aim to model. For
modeling single-neuron responses, previous studies have used these types of representations to explain
sensitivity to broadband LFP (Truccolo et al., 2005), beta-band rhythms in cortex (Kelly et al., 2010),
theta rhythms in hippocampus (Harris et al., 2003), and sensitivity across bands and phases(Cui et al.,
2016). Here we consider two models: 1) a compact model based on the Generalized Phase (Davis et al.,
2020), and 2) a complex Morlet filterbank model that can capture more-detailed frequency-specific
effects.

Briefly, the Generalized Phase (GP) approach (Davis et al., 2020) constructs a broadband analytic signal
with a smoothly-varying phase. Recent work (Davis et al., 2022) has shown that the broadband phase is
often a better predictor of spiking than the phase of traditional narrowband signals (e.g. theta or beta).
Here we consider signals in the band (5-40 Hz, 4th-order Butterworth), and the representation takes the
form

$$\boldsymbol{x}(t) = [1 \; Re\big(gp(t)\big) \; Im\big(gp(t)\big)]$$

where $gp(t)$ denotes a complex-valued analytic signal (Davis et al., 2022).

The complex Morelet filterbank has also been previously used for modeling spike-LFP coupling (Cui et
al., 2016). Here we use 7 complex-valued Morelet wavelet filters with log-spaced center frequencies
varying from 1 to 64 Hz (approximately equal quality factor) plus a real-valued DC. After including an
intercept term, this gives a representation $\boldsymbol{X}$ with 16 predictors.

Note that we can also construct predictors that include features from multiple LFP channels, e.g.

$$x_{ij}(t) = y_j(t) * F_i(t)$$

220 where x_{ij} denotes the output of the ith filter for the jth LFP channel. Here we consider GP and filterbank
221 models that use either one LFP channel (the closest channel to neuron 1) or models with two LFP channels
222 (the closest channel to neuron 1 and the closest channel to neuron 2).

Time-varying prediction

To characterize nonstationary changes in the spike-spike correlation, we measure the correlogram in short-term moving windows. For a time window $w = (t_{start}, t_{end})$, we have observations $c_w(\tau) = \sum_{t \in w} n_1(t)n_2(t + \tau)$, and stacking the results from each window generates a data matrix for the overall recording.

Here we are particularly interested in whether modeling LFP allows for dynamic description of the spike-spike correlation. In addition to the global models described above, we thus also consider local observations and models based on calculating and predicting the spike-spike correlation within a moving window. For a time window $w = (t_{start}, t_{end})$, we have observations $c_w(\tau)$ and predictions $\hat{c}_w(\tau)$ or $f_w(\tau)$ based on the local LFP cross-covariance $\boldsymbol{C}_w(\tau) = \sum_{t \in w} \boldsymbol{x}^T(t)\boldsymbol{x}(t + \tau)$. Even when the parameters $\{\boldsymbol{\beta}, \boldsymbol{\gamma}\}$ or $\boldsymbol{\alpha}$ are fixed, the predictions of both the indirect and direct models described above will change with the spectrum of the LFP.

In general, we find that such stationary models do not accurately track the time-varying correlograms due to the time-varying rates of the two neurons. Rate changes lead to shifts in the baseline of the correlogram that a stationary model cannot track. To better match the data, we consider two methods for tracking these changes. In a first approach, we maintain fixed parameters, but modify predictions with an offset of the form

$$f_{w,indirect}(\tau) = \exp(\gamma_0 + \gamma_1 \hat{c}_w(\tau) + \log(N_{w1}N_{w2} + 1))$$

where N_{wi} denotes the total number of observed spikes in the window w, $N_{wi} = \sum_{t \in w} n_i(t)$. When $\boldsymbol{\gamma} = 0$ this offset model gives an additively-smoothed description of the expected baseline correlation that varies with w but is flat over τ. For general γ, the offset acts as a gain on $f(\tau)$ that varies from window to window based on the observed rates of the two neurons.

In a second approach, we consider a fully-adaptive version of the indirect model where the parameters $\boldsymbol{\gamma}$ vary along with the moving window. This approach mirrors previous work on Poisson adaptive filtering and smoothing for point-processes (Eden et al., 2004). Briefly, we consider

$$f_{w,adaptive}(\tau) = \exp(\gamma_{w0} + \gamma_{w1} \hat{c}_w(\tau))$$

where $\boldsymbol{\gamma}$ is updated at each window according to the observed local correlations. Here we track the posterior mean $\boldsymbol{\gamma}_w = [\gamma_{w0}, \gamma_{w1}]$ and covariance $\boldsymbol{\Sigma}_w$ of the parameters using a forward-filtering approach under the assumption that $\boldsymbol{\gamma}$ evolves according to $\boldsymbol{\gamma}_{w+1} \sim N(\boldsymbol{\gamma}_w, \boldsymbol{\sigma}^T I)$. This Poisson adaptive filter has forward updates

$$\boldsymbol{\Sigma}_w^{-1} = (\boldsymbol{\Sigma}_{w-1}^{-1} + \boldsymbol{\sigma}^T I) + \sum_\tau \boldsymbol{z}_w^T(\tau)f_{w-1}(\tau)\boldsymbol{z}_w(\tau)$$

$$\boldsymbol{\gamma}_w = \boldsymbol{\gamma}_{w-1} + \boldsymbol{\Sigma}_w \sum_\tau \boldsymbol{z}_w^T(\tau)(c_w(\tau) - f_{w-1}(\tau))$$

267 where $\mathbf{z}_w = [1 \; \hat{c}_w]$. In this setting, the only additional parameter needed to model the dynamic correlation
268 is the process noise $\boldsymbol{\sigma}$, which describes how variable the baseline γ_{w0} and gain γ_{w1} are expected to be
269 from window to window.

Quantifying stability of cross-correlations over time
272 To measure the extent of nonstationarity of spike-spike correlations between neuron pairs, we use a low-
273 rank Poisson decomposition and row/column-wise approximations of $c_w(\tau)$. Namely, we fit models of
274 the form

$$c_w \sim Poisson(\exp(\boldsymbol{AB}))$$

where measured $\boldsymbol{c}_w$ is the $[W \times M]$ matrix of count observations in W non-overlapping windows at M
delays, with entries ij corresponding to $c_{w_i}(\tau_j)$. The observations are reconstructed based on the
combination of a $[W \times k]$ matrix $\boldsymbol{A}$ and a $[k \times M]$ matrix $\boldsymbol{B}$. Here we use $k \leq 2$ for the low-rank models,
and find approximate maximum likelihood estimates of $\boldsymbol{A}$ and $\boldsymbol{B}$ by coordinate ascent (i.e. alternately
optimizing $\boldsymbol{A}$ with fixed $\boldsymbol{B}$, and optimizing $\boldsymbol{B}$ with fixed $\boldsymbol{A}$). We also consider the simplified row-
approximation $\exp(a1^T)$ where $a_i = \log\left(\overline{c_{w=w_i}(\tau)}\right)$ and column-approximation $\exp(1b^T)$ where $b_j =$
$\log\left(\overline{c_w(\tau = \tau_j)}\right)$. In this case, the predictions are based on the row- and column-wise averages. We then
evaluate goodness-of-fit for all models using the pseudo-R^2 as above.

Neural Data

To evaluate and test our models, we use multi-region recordings from the Allen Institute for Brain
Science's Visual Coding Neuropixels dataset (https://portal.brain-map.org/explore/circuits). Detailed
procedures for surgery, stimulus presentation, and preprocessing are documented in Siegle et al. (2021).
Briefly, recordings were obtained from head-fixed, freely-running mice exposed to a variety of visual
stimuli, including Gabor patches, full-field drifting gratings, moving dots, and naturalistic images and
videos. Our analysis focuses on single-unit data sorted with Kilosort 2, along with simultaneously-
recorded local field potentials obtained from the same probe and region as the units. LFP data was low-
pass filtered and sampled at 1250 Hz. We restrict our analysis to units with SNR>2 and with more than
1000 spikes. Additionally, we only evaluate recordings with >2hrs of simultaneous spikes and LFP, with
electrodes in either the hippocampus or visual cortex, and with $\geq$10 units per probe in the specified area.

Altogether we analyze data from 53 unique recordings, with 44 probes in hippocampus and 51 probes in
cortex meeting our criteria. On average, there were 25 units that met criteria per probe in hippocampus
and 54 units per probe in cortex across the recordings with 15,000 possible pairs of simultaneously-
recorded hippocampal neurons and 86,000 possible pairs of cortex neurons in total. For our population
analyses below, we randomly-sampled 25 pairs from each probe with sufficiently-high correlation strength
index (CSI) compared to the jittered data, giving n=1100 hippocampal pairs and n =1400 cortical pairs.

Code for the analysis and simulations described here is available at
https://github.com/nztmansouri/nonstationary-correlogram.

Results

Here we consider models of spike-spike coupling based on simultaneous observations of the local field
potential. Intuitively, if the spikes from neuron 1 and 2 have common drive from the LFP, we expect the

correlation between neurons 1 and 2 to be partially determined by their separate spike-field relationships.
To illustrate how spike-spike correlations can arise from separate spike-field relationships, we first
introduce a basic simulation where two Poisson neurons are linearly-coupled to a common local field.
Neuron 1 tends to spike at the troughs of the LFP while neuron 2 tends to spike at the peaks (Fig. 1, top).
The spike-field relationships are summarized here by the spike-triggered average (STA), and the spike-
triggered average is a cross-correlation between the spikes and LFP (De Boer and Kuyper, 1968). At the
same time, the two neurons' separate spike-field relationships give rise to a spike-spike relationship, where
the cross-correlation between the two neurons is reduced at delays/intervals near 0, and increased at delays
near one-half of the LFP period (Fig 1, middle).
321
Some analyses have framed the interpretation of spike-spike correlations as one where the observed spike-
spike correlation should be distinguished from the partial correlation between the two neurons (Rosenberg
et al., 1989, 1998; Eichler et al., 2003). Using the partial correlation rather than the direct pairwise
correlation can generate estimates of the excess correlation between the neurons after other variables have
been taken into account. However, the partial correlation approach also allows the observed spike-spike
correlation to be predicted. When predicting spike-spike cross-correlations at multiple delays given a
single LFP, the partial correlation approach works by using the spike-triggered-average LFP (STA) for
each of the two neurons. The predicted correlation is then based on the normalized product of the STAs
in the frequency domain (see Methods). This STA method (Halliday et al., 1995) can often accurately
describe the spike-spike correlations observed *in vivo* (Goldberg et al., 2004; Middleton et al., 2012). In
Fig 1, the STA method accurately predicts the spike-spike correlation of the simulated neurons.
333
One potential challenge in predicting spike-spike correlations, in general, is that the LFP may not be
stationary over the course of the recording. For a stationary LFP, the STAs and effect of the LFP on the
spike-spike correlation will be constant over time as long as the spike-field relationships are fixed.
However, for a nonstationary LFP, these quantities can both change with the LFP. Even if the spike-field
relationships are fixed, a changing LFP can result in both changing STAs and changing correlations
between neurons. In our simulation, we illustrate two potential patterns of changes where the LFP is either
modulated in frequency (Fig. 1A) or amplitude (Fig. 1B). In this case, the STA method accurately predicts
spike-spike correlations in short windows, but only when the STA is calculated within the same windows.
Additionally, although the STA for the time-averaged data (Fig. 1, bottom) can predict the observed time-
average spike-spike correlogram, the time-averaged STA does not match the STA for short-windows. The
STA method provides a nonparametric approach where data directly defines the predictions. Here we
consider alternative parametric approaches for predicting spike-spike correlations, and demonstrate how
these approaches might be useful for predicting time-varying spike-spike correlations.
347

356
To illustrate both the parametric and nonparametric approaches for predicting spike-spike correlations,
we consider one example pair of neurons recorded from the hippocampus of an awake mouse along with
simultaneously-observed LFP (Fig 2). Like the simulation, these real neurons are coupled to the LFP and
tend to spike at the peaks and troughs of the LFP signal, respectively (Fig 2A). As with many hippocampal
recordings, the LFP signal here has substantial power in the theta band (~8 Hz), and this oscillation is
reflected in the spike-spike correlation. As in the simulation above, the STA method predicts the spike-
spike correlation by first measuring the STAs of the two neurons, then normalizing the product of the
STAs by the power spectrum of the LFP in the frequency domain (Fig 2B). Similar to previous studies
(Goldberg et al., 2004), the STA method accurately predicts the correlation between this particular pair of
neurons (Fig 2C).
367
As an alternative to the nonparametric STA method, we introduce a parametric approach where an explicit
representation of the LFP allows the prediction problem to be framed using a fixed set of predictors

weighted by coefficients. Briefly, this approach works by modeling the spike-spike correlations with a
bilinear model – where coefficients for both neurons weight the cross-correlation of LFP predictors,
followed by a nonlinearity that matches the observed cross-correlation (see Methods). To illustrate this
parametric approach, we consider two representations for the LFP here: (1) a generalized phase
representation (Fig 2D-F) and (2) a wavelet filterbank representation (Fig 2G-I). The generalized phase
representation extracts two predictors from the LFP, which correspond to the real and imaginary parts of
the broadband analytic signal. This model only has two coefficients for each neuron (Figure 2E), but
combining these coefficients with the cross-correlation of the predictors can yield predictions of the spike-
spike correlation nearly as accurate as those of the nonparametric STA method. Similarly, we can consider
separating the LFP into narrower frequency bands (Fig 2G). Here we use a wavelet filterbank model with
seven narrow-band, complex Morlet wavelet filters (center frequencies starting from 1 Hz and increasing
to 64 Hz, log-spaced octave wide filters) to generate predictors. This model has 15 fixed parameters for
each neuron (Fig 2H), and predicts the spike-spike correlation accurately as well (Fig 2I). The differences
between the coefficients of the two neurons reflect the difference in the neurons' spike-field relationships,
including their amplitude, phase, and sensitivity to specific frequency bands.
385
There are several approaches to optimize the coefficients of the parametric models. Here we consider an
indirect method and a direct method (see Methods). In the indirect method, the per-neuron coefficients
and the nonlinearity are optimized sequentially, with the per-neuron coefficients fit based on linear
prediction of the spikes of each neuron. In the direct method, the coefficients of a generalized bilinear
model are optimized directly to predict the observed cross-correlation. Results for the indirect method are
shown in Fig 2. In both approaches, the key feature of the parametric models is that they explicitly
distinguish between LFP-based predictors and parameters that describe the spike-field or correlation-field
relationships. This distinction allows us to directly extend the models to include multiple LFP recording
channels, and to make predictions of spike-spike correlations for arbitrary LFP without needing to track
or update the STA based on newly-observed spikes.

*Figure 2. Methods and models for predicting spike-spike correlations from LFP. **A)** An example segment (2 s) of an LFP signal recorded in mouse hippocampus with two simultaneously-recorded neurons coupled to the LFP peaks and troughs. Nonparametric methods for predicting spike-spike correlations are based on the data directly **(B-C)**, while here we consider parametric, bilinear models with an explicit representation of predictors **(D-I)**. **B)** A nonparametric STA method. STAs (left) for neuron 1 (blue) and 2 (orange) are normalized by the LFP power spectrum (right) to generate the predicted spike-spike correlation, with the observed (gray) and predicted (red) correlations shown in (C). **D)** A generalized phase representation where the bandpass-filtered LFP is used to calculate a complex analytic signal (real part shown in black and imaginary part shown in gray) with smoothly-varying phase (depicted on a color scale below). **E)** The predictions of the parametric model (shown in **F**) are based on the cross-correlation of the predictors (left) weighted by coefficients for each neuron (right). **G)** A wavelet filterbank representation splits the LFP into narrow bands (here the real part is shown), with y-axis indicating the central frequencies of each complex filter. **H)** This representation generates predictors (left) again weighted by coefficients (right) to generate predictions of the spike-spike correlation (**I**).*

Although both the nonparametric STA method and the parametric models accurately fit this example pair of neurons, the models do not always explain the variability in the data for different pairs of neurons. One challenge in measuring goodness-of-fit for these models is that different pairs of neurons have correlations of different strengths (Fig 3A-B), and for weak correlations the variance of the cross-correlation at different intervals is mostly driven by noise/chance. To quantify the correlation strength, we use a correlation strength index (CSI) based on the ratio of the variance and the mean of the spike-spike cross-correlations over different delays. Since independent Poisson neurons will have cross-correlations with a variance equal to the mean, a CSI of 1 represents a case where we would not expect the model to predict any meaningful structure in the cross-correlation. Here we measure the CSI of many simultaneously-recorded pairs of neurons in mouse hippocampus (n=1100 pairs) and cortex (n=1400 pairs) from the Allen Institute's Visual Coding Neuropixels dataset. We find a wide range of CSIs with strong, highly-structured cross-correlations (Fig 3A-B, top) as well as weaker cross-correlations with no clear variation across different delays (Fig 3A-B, bottom). The distribution of CSIs is highly-skewed with a median of 2.90 for

hippocampus and 2.65 for cortex (Fig 3C-D). For comparison, we also evaluate the CSI for a null model of each pair of neurons where we jitter the neural spike times (normally-distributed jitter with 1s s.d.). Since neuron pairs with low CSIs (similar to the CSI for the jittered data) do not have substantial structure for the LFP-based models to capture, we focus our analysis on the pairs with CSIs higher than the decision boundary between the jittered and observed CSI distributions. We fit our models to a random sample of 25 pairs from each recording probe with n=1100 pairs from hippocampus and n=1400 pairs from cortex. As expected, we find that the goodness-of-fit (pseudo-R^2) is correlated with the CSI (Fig 3C-D).

Figure 3. *Example cross-correlograms for neuron pairs from the hippocampus **(A)** and cortex **(B)**. Observed correlations (gray) are shown normalized by the correlations of a jittered null model (blue). The prediction of the indirect model with generalized phase representation of the LFP is shown in red. Histograms of observed correlation strength index (CSI) for all correlogram samples are shown in gray alongside jittered correlograms (blue) for neuron pairs in the hippocampus **(C)** and cortex **(D)**. Scatter plots show the model pseudo-R^2 with respect to CSI for a sample of neuron pairs with CSIs above that of the jittered data.*

Using this sample of high-CSI pairs, we compare the performance of the nonparametric STA method to that of several parametric models: 1) a generalized phase (GP) model based on the LFP channel closest to one neuron in the pair, 2) a GP model based on two LFP channels – the closest to neuron 1 and the closest to neuron 2, 3) a wavelet filterbank model of one LFP channel, and 4) a GP model based on one LFP channel fit by direct optimization of the bilinear coefficients. These comparisons vary in terms of the

representation of LFP predictors (GP vs filterbank), the number of LFP channels used, and the method of optimizing the coefficients (indirect vs direct). They also differ in the number of parameters (the 1-channel GP indirect model has 8 parameters, the 2-channel GP indirect model has 12, the 1-channel filterbank model has 32, and the 1-channel GP direct model has 6).

Fitting each of these models to the high-CSI pairs from hippocampus and cortex (Fig 4A and B), we find that the models are all correlated in their goodness-of-fits (pseudo-R^2), with the results depending on the CSI of the data. However, there are some differences in performance across models. For hippocampus, the 1-channel GP direct model (pseudo-R^2=0.022 [0.008, 0.069], median and inter-quartile range), 1-channel filterbank model (pseudo-R^2=0.015 [0.003, 0.068]), and STA method (pseudo-R^2=0.014 [0.003, 0.070]) all perform well, while the 1-channel (pseudo-R^2=0.008 [0.002, 0.040]) and 2-channel (pseudo-R^2=0.010 [0.002, 0.048]) GP indirect models perform slightly worse (Fig 4A). For cortex, the 1-channel filterbank model (pseudo-R^2=0.026 [0.005, 0.085]), and STA method (pseudo-R^2=0.027 [0.006, 0.088]) perform best, while the 1-channel (pseudo-R^2=0.004 [0.001, 0.012]) and 2-channel (pseudo-R^2=0.003 [0.001, 0.012]) GP indirect and GP direct (pseudo-R^2=0.011 [0.005, 0.026]) models perform slightly worse (Fig 4B). The absolute R^2 values may be somewhat misleading since the goodness-of-fit is highly dependent on the CSI (Fig 4, insets). The models, overall, can accurately capture structure in the cross-correlations when there is structure present (e.g. high-CSI examples in Fig 3).

Figure 4. *Goodness-of-fit for five models of spike-spike correlations based on LFP: STA method, 1-channel generalized phase model with indirect optimization of the coefficients, 1-channel filterbank model with indirect optimization of the coefficients, 2-channel generalized phase model with indirect optimization of the coefficients, and 1-channel generalized phase model with direct optimization of the coefficients. Scatter plots comparing pseudo-R^2 for different pairs of models for **A)** hippocampus and **B)** cortex. Insets show dependence of pseudo-R^2 on data CSI. Boxplots compare the model pseudo-R^2 measured across different CSI ranges. High, Med, and Low CSI denote the first, second, and third CSI terciles, respectively. Boxplot horizontal black lines denote the median pseudo-R^2, and the box ranges indicate the interquartile ranges for goodness-of-fits.*

As in the simulation, a major consideration when predicting spike-spike correlations from LFP is that the LFP is not stationary. LFP power spectra change over time, and we expect these changes to be reflected in changing spike-spike correlations. To determine the nonstationarity of spike-spike correlations, we first

measure correlograms for the high-CSI neuron pairs in short time windows of 60s (Fig 5A-B). Most pairs
show dynamic patterns where the correlation increases and decreases over time, and the correlations at
different intervals/delays are not fixed over the course of the recording. Here we are measuring the
"unnormalized" correlogram based on spike counts and without adjusting for changes in firing rates over
time. When the firing rates of the two neurons vary over the course of the recording, the cross-correlation
naturally changes in magnitude. Such changes are not strictly due to changes in LFP, since they can occur
even if the neurons are statistically independent (e.g. two independent inhomogeneous Poisson processes).
484
To test whether the observed changes in the correlograms over the course of the recording go beyond
simple rate changes, we here fit four Poisson matrix decompositions (see Methods). Specifically, we fit:
1) a row-decomposition that assumes an interval-dependent correlation that is fixed over time, 2) a
column-decomposition that assumes a time-dependent correlation that is constant for all intervals, 3) a
rank-1 decomposition that assumes the matrix can be decomposed as a product of one pattern over time
and one over interval, and 4) a rank-2 decomposition. We measure pseudo-R^2 for each of these matrix
decompositions as a function of the window size (Fig 5C-D). We find that the row decomposition does
not accurately track the unnormalized correlograms, while the other three decompositions do well across
different window sizes. Most of the variability in these matrices is explained by a column effect where
rate fluctuations drive changes in the baseline cross-correlation over the course of the recording. However,
a rank-1 decomposition does slightly better by assuming one pattern of correlation across different
intervals, and a second pattern that acts as a gain for different time windows. These decompositions are
consistent with the idea that there are fixed patterns of correlations that change due to the changing firing
rates of the two neurons. However, we find that a rank-2 decomposition does lead to small improvements
in the reconstruction relative to the rank-1 decomposition, suggesting that there may be some additional
nonstationarity beyond what results from changing firing rates.
501
The goodness-of-fit increases with the window size for all decompositions (Fig 5C). For short time
windows (< ~10s) the pseudo-R^2 for all reconstructions is relatively low, likely due to the fact that the CSI
is also lower at this resolution. The improvement provided by the rank-2 decomposition compared to the
rank-1 decomposition gives an approximate lower-bound for how much the correlation pattern changes
over the course of the recording. For comparison, we also evaluate the improvement from using the rank-
2 vs rank-1 decomposition for jittered data (Gaussian noise used to jitter spike times with 1s s.d.). The
improvement from using the rank-2 vs rank-1 decomposition for the observed data is higher than the
improvement with the jittered data for all window sizes. For 60-s windows (Fig 5C-D, inset), the rank-2
decomposition shows an improvement over the rank-1 decomposition of 1.4% (median) and 1.3% for
cortex and hippocampus, respectively, and the improvement is higher than that of the jittered data (0.6%
and 0.5% for cortex and hippocampus). As with the case of predicting stationary spike-spike correlations,
CSI is also an important consideration when evaluating nonstationary correlations. This analysis suggests
that in some cases there is nonstationarity in spike-spike correlations, and thus our goal here is to test
whether these changes can be predicted by the LFP.

Figure 5. Measuring nonstationarity in spike-spike correlations with matrix decomposition. To illustrate how correlograms change over the course of the recording, we calculate correlations in short time windows. Here, heatmaps indicate correlations at different delays (y-axis) calculated in 60-s time windows over the course of the ~3hr recording (x-axis). Example neuron pairs with time-varying correlograms are shown for the hippocampus (**A**, left) and cortex (**B**, left), along with a low-rank (rank=1) matrix decomposition (A, right for hippocampus, and B, right for cortex). For hippocampus (**C**, left) and cortex (**D**, left), pseudo-R^2 is shown for the row (blue), column (red), rank-1 (yellow), and rank-2 (green) decompositions as a function of window size. Improvement of the decomposition when using rank-2 compared to rank-1 with both the jittered (green) and the observed correlations (gray). The red arrow indicates a window size of ~60s, which corresponds to the inset histogram.

To illustrate how changes in the LFP spectrum over time might lead to changes in the spike-spike correlations, here we consider one example pair of neurons from the hippocampus (Fig 6). As the theta-band power and peak frequency vary over the course of the recording (Fig 6A), the spike-spike cross-correlation for this pair of neurons tends to vary as well (Fig 6A, middle). We fit our indirect generalized phase model using this LFP channel and find that when we apply the predictions in short 60-s windows (see Methods, Fig 6A, bottom), the model accurately tracks changes in the correlogram, which correspond to changes in both the LFP's peak frequency (Fig 6B) and amplitude (Fig 6C).

Figure 6. *Example and model fits illustrating how nonstationary spike-spike correlations are related to changes in the LFP spectra, with LFP and neuron pair from the hippocampus.* ***A***. *Top, spectrogram of the LFP signal. Middle, observed correlograms calculated in 60s short-time windows (50-s overlap). Bottom, time-varying correlogram predictions from the indirect generalized phased model.* ***B***. *Top, the LFP's PSD for two time points (1=blue and 2=orange) indicating that the dominant frequency of time 1 is higher than that of time 2. In the bottom left and right, the observed correlograms (gray) and corresponding model fits in blue (bottom left) and orange (bottom right) also illustrate the changes in frequency (i.e. by counting the number of peaks if we consider the fluctuations as pseudo-sinusoid).* ***C***. *Top, the LFP's PSD for two different time points, in which time 3 (green) has a higher amplitude than time 4 (purple). In the bottom left, the observed correlogram (gray) has higher amplitude (model fit in green) than the observed correlogram (gray) in the bottom right (model fit in purple). Both fits use the generalized phase model.*

As with modeling the stationary correlograms, there are multiple possible approaches when describing the time-varying correlograms that differ in predictors (i.e. generalized phase vs filterbank), optimization strategies (i.e. indirect vs direct) and number of parameters. One major challenge, however, is in how to account for the time-varying rates of the two neurons and the column-wise structure of the matrices described above. Here, when fitting the time-varying indirect model, we assume that there is a known

549 count offset for each window (see Methods). This approach may be appropriate for offline modeling, but
550 a different strategy may be needed when correlations are predicted in real-time. To illustrate one possible
551 approach, we consider an adaptive filtering model to track nonstationary spike-spike correlations from
552 streaming data.
553
Adaptive filtering has been previously applied to the estimation of changing neural tuning properties
(Eden et al., 2004) and synaptic connections (Stevenson and Kording, 2011) from spike data. Here we
consider the adaptive filtering approach for LFP-driven spike-spike correlations. Briefly, we use the
predictors from the indirect model and add an output nonlinearity with time-varying parameters for the
baseline correlation and gain (see Methods). We maintain a distribution over parameters at each time
window, and update the mean and covariance as new observations come in. To validate this approach,
here we use experimentally-observed hippocampal LFP to simulate two neurons with a spike-spike
correlogram with known time-varying gain (Fig 7). The adaptive filtering model appropriately
accumulates uncertainty even when spike observations are missing, and makes predictions based on the
LFP. When there are true changes in the baseline or gain, the adaptive filter can track these changes, and
distinguish these effects from changes in the LFP itself. Unlike the indirect model described above, the
adaptive filter does not require a separate offset when accounting for changes in the column-wise structure.
Instead, the column-wise structure is captured by a baseline that is updated based on new observations as
they occur.

568
569
570 ***Figure 7.*** *Adaptive filtering model for time-varying correlogram data. Here we use the observed LFP to simulate time-varying spike-spike*
571 *correlations with known changes in the parameters. We simulate Poisson spiking with a fixed baseline and variable gain (in green in top two*
572 *rows, respectively) as well as some missing data (time points 8-17 min). The adaptive filter tracks the observed spiking (middle row) over*
573 *time and updates parameter estimates based on the observations (top two rows, gray bands). The adaptive filter can fill-in missing data (red*
574 *arrows at time points 8 and 17 min.), and accurately tracks imposed changes on the gain (step-wise changes indicated by arrows at time*

points between 20 and 40 min). The correlations estimated by the adaptive filter (row above bottom) follow the true correlations (bottom

row) when the ground truth is known. Note that some changes in the data are driven by changes in the LFP (e.g. yellow arrow) rather than

changes in the adaptive parameters.

Figure 8. *Goodness-of-fit for LFP-based models of time-varying spike-spike correlations. For hippocampus (left) and cortex (right), we fit*

cross-correlations measured in 60-s windows. We compare a stationary model that assumes a fixed spike-spike correlation over the recording

(generalized phase, indirect optimization) to a nonstationary model that includes the local variations in LFP (generalized phase, indirect

optimization), as well as an adaptive filtering model that tracks time-varying baseline and gain. Boxes denote inter-quartile ranges (i.e.

middle 50% of observed model R^2 values) and whiskers denote extrema.

To evaluate the goodness-of-fit for these nonstationary LFP-driven models of spike-spike correlations, we
again calculate pseudo-R^2 after fitting the high-CSI pairs from hippocampus and cortex. We compare the
nonstationary, indirect generalized phase model and the adaptive filtering model against a stationary
model where the same LFP-based prediction (indirect GP) is applied to all time windows. Generally, we
find that the nonstationary models are substantially more accurate than the stationary model, and the
indirect approach with column-wise rate offsets performs slightly better than an adaptive filtering
approach with running parameter estimates. Using 60-s windows and hippocampal neuron pairs, we
obtained median pseudo-R^2 for the stationary (0.001), nonstationary indirect (0.45), and adaptive filtering
models (0.35). For cortical neuron pairs, we also observed median pseudo-R^2 for the stationary (0.0001),
nonstationary indirect (0.41), and adaptive filtering models (0.31). Thus for both hippocampal and cortical
neural pairs, the LFP can provide an accurate prediction of time-varying cross-correlations.

596
597
Discussion

Here we developed parametric models to predict nonstationary spike-spike correlations based on
simultaneously-observed LFP. Using experimental data from mouse hippocampus and cortex, we showed
how our models, with relatively few parameters, often perform as well as a previous nonparametric
approach based on the spike-triggered average LFP (STA). Parametric models may also prove to be more
interpretable due to the fact that they can specifically describe the contributions from distinct LFP channels
and bands. We then applied these models to the problem of predicting time-varying spike-spike
correlations. We found that most of the time-varying structure is due to time-varying firing rates of the
neurons, and most of the structure in cross-correlogram matrices is accounted for by a rank-1
decomposition. However, there is additional variability over time that is not well-explained by time-
varying firing rates when combined with the average correlogram, and this structure can be partially-
predicted by parametric models based on the LFP.

Spike-spike correlations are major tools for characterizing neural processing, and many studies have
documented the dependence of correlations on stimuli (Kohn and Smith, 2005), attention (Cohen and
Maunsell, 2009), brain state (Ecker et al., 2014), distance between neurons , anatomical location (Senzai
et al., 2019), and neuromodulation (Minces et al., 2017), among many other factors. At the same time,
correlations are often used to assess potential synaptic interaction between neurons (Moore et al., 1970;
Fetz et al., 1991) or functional connectivity more broadly (Gerstein and Perkel, 1969; Stevenson et al.,
2008; Magrans de Abril et al., 2018). Some of these relationships, such as the effect of brain state, may
be directly attributable to the LFP, while others, such as synaptic effects, are likely distinct from LFP-
driven correlations. In previous work on detecting and modeling synaptic effects, researchers have
explicitly attempted to distinguish between slow, time-varying baseline changes and fast synaptic effects
(Ren et al., 2020; Wei and Stevenson, 2021). However, modeling the contribution of the LFP to spike-
spike correlations, and how this contribution changes over time, may allow for more precise identification
of both LFP-driven effects and the contributions of other factors.

There are several limitations to the results presented here. One consideration is the presence of spike
bleed-through, since low-pass filtering of the wideband extracellular signal does not completely remove
the effects of spikes on the LFP (David et al., 2010; Ray, 2015; Banaie Boroujeni et al., 2020). Here we
have not attempted to correct for these effects, thus spike bleed-through may affect the predicted
correlations for delay intervals close to 0. However, since the parametric methods are based on even-
narrower filtering of the LFP, it is unlikely to play a major role in the results here, especially when $\tau \gg 0$.
We have also focused exclusively on pairwise, single-trial correlations and time-domain descriptions of
the spike-LFP relationship. There may be some benefits to extending our models to the population level
(Yatsenko et al., 2015), using multiple trials to distinguish between spike count correlations and firing rate
correlations (Vinci et al., 2016), or considering nonstationarity in the frequency domain (Zhan et al.,
2006).

There are many existing statistical tools for detecting the presence of and changes in correlations (Grün,
2009; Harrison et al., 2013; Lepage et al., 2013; Zhou et al., 2015). When repeated trials are available,
one common approach is to use shuffled correlations (Perkel et al., 1967; Eggermont, 1990; Chen et al.,
2012), and with single trials the jittered correlogram can provide a null model. Here we have focused
primarily on modeling single-trial data and changes over time with LFP-based predictors. Our parametric
approach is, most directly, an alternative to nonparametric methods of linking the LFP to spike-spike
correlations (Eggermont and Smith, 1995; Halliday et al., 1995; Goldberg et al., 2004; Middleton et al.,
2012). However, the parametric models developed here could also be used for hypothesis testing using

model comparison. When evaluating nonstationarity, it may be useful to more precisely distinguish
between situations where spike-spike correlations change due to: 1) common dependence on a changing
LFP, 2) changing spike-field coupling, and 3) changes unexplained by the LFP, such as plasticity in a
monosynaptic connection between the neurons.
649
Finally, one important application of the parametric models here may be in improving neural decoding.
Modeling correlations improves decoding (Stevenson et al., 2012), and the presence of correlations may
have a beneficial effect on readout by downstream neurons (Valente et al., 2021). Observed correlations
can be also be directly used by decoders to predict stimuli (Sadeghi et al., 2019; Zhai et al., 2020), brain
state (Benisty et al., 2023), and behavior (Maynard et al., 1999) more accurately than when using firing
rates alone. At the same time, LFP has been used as a signal for decoding both by itself and in addition to
spike signals (i.e. joint decoding). These studies have found that joint decoding generally leads to
improvements, but the individual contributions from LFP vs. from spikes are partially redundant, just as
the contributions from each of multiple neurons can be partially redundant. By modeling the link between
LFP and observed spikes, as well as the link between LFP and spike-spike correlations, the tools developed
here may allow for easier disentangling of these contributions.
661

Acknowledgements

This material is based upon work supported by the National Science Foundation under Grant 1651396. Thanks to the Allen Institute for Brain Science for sharing their datasets and for supporting open science.

References

Amarasingham A, Harrison MT, Hatsopoulos NG, Geman S (2012) Conditional modeling and the jitter method of spike resampling. Journal of Neurophysiology 107:517–531.

Baker SN, Spinks R, Jackson A, Lemon RN (2001) Synchronization in Monkey Motor Cortex During a Precision Grip Task. I. Task-Dependent Modulation in Single-Unit Synchrony. Journal of Neurophysiology 85:869–885.

Banaie Boroujeni K, Tiesinga P, Womelsdorf T (2020) Adaptive spike-artifact removal from local field potentials uncovers prominent beta and gamma band neuronal synchronization. Journal of Neuroscience Methods 330:108485.

Barthó P, Hirase H, Monconduit L, Zugaro M, Harris KD, Buzsáki G (2004) Characterization of neocortical principal cells and interneurons by network interactions and extracellular features. Journal of neurophysiology 92:600–608.

Benisty H, Barson D, Moberly AH, Lohani S, Tang L, Coifman RR, Crair MC, Mishne G, Cardin JA, Higley MJ (2023) Rapid fluctuations in functional connectivity of cortical networks encode spontaneous behavior. Nat Neurosci:1–11.

Buzsáki G, Anastassiou CA, Koch C (2012) The origin of extracellular fields and currents — EEG, ECoG, LFP and spikes. Nature Reviews Neuroscience 13:407–420.

Canolty RT, Ganguly K, Carmena JM (2012) Task-Dependent Changes in Cross-Level Coupling between Single Neurons and Oscillatory Activity in Multiscale Networks. PLOS Computational Biology 8:e1002809.

Canolty RT, Ganguly K, Kennerley SW, Cadieu CF, Koepsell K, Wallis JD, Carmena JM (2010) Oscillatory phase coupling coordinates anatomically dispersed functional cell assemblies. Proceedings of the National Academy of Sciences 107:17356.

Chen C, Read HL, Escabí MA (2012) Precise Feature Based Time Scales and Frequency Decorrelation Lead to a Sparse Auditory Code. J Neurosci 32:8454–8468.

Cohen MR, Kohn A (2011) Measuring and interpreting neuronal correlations. Nature neuroscience 14:811–819.

Cohen MR, Maunsell JHR (2009) Attention improves performance primarily by reducing interneuronal correlations. Nature neuroscience 12:1594–1600.

Cui Y, Liu LD, McFarland JM, Pack CC, Butts DA (2016) Inferring Cortical Variability from Local Field Potentials. J Neurosci 36:4121–4135.

David SV, Malaval N, Shamma SA (2010) Decoupling Action Potential Bias from Cortical Local Field
Potentials. Computational Intelligence and Neuroscience 2010:e393019.

Davis ZW, Muller L, Martinez-Trujillo J, Sejnowski T, Reynolds JH (2020) Spontaneous travelling
cortical waves gate perception in behaving primates. Nature 587:432–436.

Davis ZW, Muller L, Reynolds J (2022) Spontaneous Spiking Is Governed By Broadband Fluctuations. J
Neurosci Available at: https://www.jneurosci.org/content/early/2022/05/20/JNEUROSCI.1899-
21.2022 [Accessed June 2, 2022].

De Boer E, Kuyper P (1968) Triggered Correlation. IEEE Transactions on Biomedical Engineering BME-
15:169–179.

deCharms RC, Merzenich MM (1996) Primary cortical representation of sounds by the coordination of
action-potential timing. Nature 381:610–613.

Denker M, Roux S, Lindén H, Diesmann M, Riehle A, Grün S (2011) The Local Field Potential Reflects
Surplus Spike Synchrony. Cerebral Cortex 21:2681–2695.

Ecker AS, Berens P, Cotton RJ, Subramaniyan M, Denfield GH, Cadwell CR, Smirnakis SM, Bethge M,
Tolias AS (2014) State Dependence of Noise Correlations in Macaque Primary Visual Cortex.
Neuron 82:235–248.

Eden UT, Frank LM, Barbieri R, Solo V, Brown EN (2004) Dynamic analysis of neural encoding by point
process adaptive filtering. Neural computation 16:971–998.

Eggermont JJ (1990) The Correlative Brain. In: The Correlative Brain: Theory and Experiment in Neural
Interaction (Eggermont JJ, ed), pp 267–281. Berlin, Heidelberg: Springer. Available at:
https://doi.org/10.1007/978-3-642-51033-5_15 [Accessed July 25, 2025].

Eggermont JJ, Smith GM (1995) Synchrony between single-unit activity and local field potentials in
relation to periodicity coding in primary auditory cortex. Journal of Neurophysiology 73:227–245.

Eichler M, Dahlhaus R, Sandkühler J (2003) Partial correlation analysis for the identification of synaptic
connections. Biol Cybern 89:289–302.

Fetz E, Toyama K, Smith W (1991) Synaptic interactions between cortical neurons. In: Normal and
Altered States of Function. Cerebral cortex (Peters A, Jones EG, eds), pp 1–47. Boston, MA:
Springer.

Gerkin RC, Tripathy SJ, Urban NN (2013) Origins of correlated spiking in the mammalian olfactory bulb.
Proceedings of the National Academy of Sciences 110:17083–17088.

Gerstein GL, Perkel DH (1969) Simultaneously Recorded Trains of Action Potentials: Analysis and
Functional Interpretation. Science 164:828–830.

Goldberg JA, Rokni U, Boraud T, Vaadia E, Bergman H (2004) Spike Synchronization in the Cortex-Basal Ganglia Networks of Parkinsonian Primates Reflects Global Dynamics of the Local Field Potentials. J Neurosci 24:6003–6010.

Grün S (2009) Data-Driven Significance Estimation for Precise Spike Correlation. Journal of Neurophysiology 101:1126–1140.

Haider B, Schulz DPA, Häusser M, Carandini M (2016) Millisecond Coupling of Local Field Potentials to Synaptic Currents in the Awake Visual Cortex. Neuron 90:35–42.

Halliday DM, Rosenberg JR, Amjad AM, Breeze P, Conway BA, Farmer SF (1995) A framework for the analysis of mixed time series/point process data—Theory and application to the study of physiological tremor, single motor unit discharges and electromyograms. Progress in Biophysics and Molecular Biology 64:237–278.

Harris KD, Csicsvari J, Hirase H, Dragoi G, Buzsáki G (2003) Organization of cell assemblies in the hippocampus. Nature 424:552–556.

Harrison MT, Amarasingham A, Kass RE (2013) Statistical Identication of Synchronous Spiking. In: Spike Timing (DiLorenzo PM, Victor JD, eds). CRC Press.

Hatsopoulos NG, Ojakangas CL, Paninski L, Donoghue JP (1998) Information about movement direction obtained from synchronous activity of motor cortical neurons. Proceedings of the National Academy of Sciences 95:15706–15711.

Kass RE, Bong H, Olarinre M, Xin Q, Urban KN (2023) Identification of interacting neural populations: methods and statistical considerations. Journal of Neurophysiology 130:475–496.

Kelly RC, Smith MA, Kass RE, Lee TS (2010) Local field potentials indicate network state and account for neuronal response variability. Journal of Computational Neuroscience 29:567–579.

Kobayashi R, Kurita S, Kurth A, Kitano K, Mizuseki K, Diesmann M, Richmond BJ, Shinomoto S (2019) Reconstructing neuronal circuitry from parallel spike trains. Nature Communications 10:1–13.

Kohn A, Smith MA (2005) Stimulus dependence of neuronal correlation in primary visual cortex of the macaque. The Journal of Neuroscience 25:3661–3673.

Lepage KQ, Gregoriou GG, Kramer MA, Aoi M, Gotts SJ, Eden UT, Desimone R (2013) A procedure for testing across-condition rhythmic spike-field association change. Journal of Neuroscience Methods 213:43–62.

Magrans de Abril I, Yoshimoto J, Doya K (2018) Connectivity inference from neural recording data: Challenges, mathematical bases and research directions. Neural Networks 102:120–137.

Maynard EM, Hatsopoulos NG, Ojakangas CL, Acuna BD, Sanes JN, Normann R a, Donoghue JP (1999) Neuronal interactions improve cortical population coding of movement direction. The Journal of neuroscience : the official journal of the Society for Neuroscience 19:8083–8093.

Middleton JW, Omar C, Doiron B, Simons DJ (2012) Neural Correlation Is Stimulus Modulated by Feedforward Inhibitory Circuitry. Journal of Neuroscience 32:506–518.

Minces V, Pinto L, Dan Y, Chiba AA (2017) Cholinergic shaping of neural correlations. Proceedings of the National Academy of Sciences 114:5725–5730.

Mizuseki K, Buzsaki G (2014) Theta oscillations decrease spike synchrony in the hippocampus and entorhinal cortex. Philosophical Transactions of the Royal Society B: Biological Sciences 369:20120530.

Moore GP, Segundo JP, Perkel DH, Levitan H (1970) Statistical Signs of Synaptic Interaction in Neurons. Biophysical Journal 10:876.

Murthy VN, Fetz EE (1996) Synchronization of neurons during local field potential oscillations in sensorimotor cortex of awake monkeys. Journal of Neurophysiology 76:3968–3982.

Nejatbakhsh A, Garon I, Williams A (2023) Estimating Noise Correlations Across Continuous Conditions With Wishart Processes. Advances in Neural Information Processing Systems 36:54032–54045.

Panzeri S, Moroni M, Safaai H, Harvey CD (2022) The structures and functions of correlations in neural population codes. Nat Rev Neurosci 23:551–567.

Perkel DH, Gerstein GL, Moore GP (1967) Neuronal Spike Trains and Stochastic Point Processes: II. Simultaneous Spike Trains. Biophysical Journal 7:419.

Pesaran B, Vinck M, Einevoll GT, Sirota A, Fries P, Siegel M, Truccolo W, Schroeder CE, Srinivasan R (2018) Investigating large-scale brain dynamics using field potential recordings: analysis and interpretation. Nat Neurosci 21:903–919.

Ray S (2015) Challenges in the quantification and interpretation of spike-LFP relationships. Current Opinion in Neurobiology 31:111–118.

Ren N, Ito S, Hafizi H, Beggs JM, Stevenson IH (2020) Model-based detection of putative synaptic connections from spike recordings with latency and type constraints. Journal of Neurophysiology 124:1588–1604.

Rosenberg JR, Amjad AM, Breeze P, Brillinger DR, Halliday DM (1989) The Fourier approach to the identification of functional coupling between neuronal spike trains. Prog Biophys Mol Biol 53:1–31.

Rosenberg JR, Halliday DM, Breeze P, Conway BA (1998) Identification of patterns of neuronal connectivity—partial spectra, partial coherence, and neuronal interactions. Journal of Neuroscience Methods 83:57–72.

Sadeghi M, Zhai X, Stevenson IH, Escabí MA (2019) A neural ensemble correlation code for sound category identification. PLoS Biology 17.

Salinas E, Sejnowski TJ (2001) Correlated neuronal activity and the flow of neural information. Nature Reviews Neuroscience 2:539–550.

Senzai Y, Fernandez-Ruiz A, Buzsáki G (2019) Layer-Specific Physiological Features and Interlaminar Interactions in the Primary Visual Cortex of the Mouse. Neuron 101:500-513.e5.

Siegle JH et al. (2021) Survey of spiking in the mouse visual system reveals functional hierarchy. Nature:1–7.

Smith MA, Kohn A (2008) Spatial and temporal scales of neuronal correlation in primary visual cortex. Journal of Neuroscience 28:12591–12603.

Spivak L, Levi A, Sloin HE, Someck S, Stark E (2022) Deconvolution improves the detection and quantification of spike transmission gain from spike trains. Commun Biol 5:1–17.

Stevenson IH, Kording KP (2011) Inferring spike-timing-dependent plasticity from spike train data. In: Advances in Neural Information Processing Systems (Shawe-Taylor J, Zemel RS, Bartlett P, Pereira FCN, Weinberger KQ, eds), pp 1–9.

Stevenson IH, London BM, Oby ER, Sachs NA, Reimer J, Englitz B, David SV, Shamma SA, Blanche TJ, Mizuseki K, Zandvakili A, Hatsopoulos NG, Miller LE, Kording KP (2012) Functional Connectivity and Tuning Curves in Populations of Simultaneously Recorded Neurons. PLoS Computational Biology 8:e1002775.

Stevenson IH, Rebesco JM, Miller LE, Körding KP (2008) Inferring functional connections between neurons. Current Opinion in Neurobiology 18:582–588.

Swadlow HA, Beloozerova IN, Sirota MG (1998) Sharp, local synchrony among putative feed-forward inhibitory interneurons of rabbit somatosensory cortex. Journal of Neurophysiology 79:567–582.

Swadlow HA, Gusev AG (2001) The impact of "bursting" thalamic impulses at a neocortical synapse. Nature neuroscience 4:402–408.

Trepka EB, Zhu S, Xia R, Chen X, Moore T (2022) Functional interactions among neurons within single columns of macaque V1 Vinck M, Behrens TE, Kremkow J, eds. eLife 11:e79322.

Truccolo W, Eden UT, Fellows MR, Donoghue JP, Brown EN, John P (2005) A Point Process Framework for Relating Neural Spiking Activity to Spiking History, Neural Ensemble, and Extrinsic Covariate Effects. Journal of Neurophysiology 93:1074–1089.

Uhlhaas PJ, Singer W (2006) Neural Synchrony in Brain Disorders: Relevance for Cognitive Dysfunctions and Pathophysiology. Neuron 52:155–168.

Valente M, Pica G, Bondanelli G, Moroni M, Runyan CA, Morcos AS, Harvey CD, Panzeri S (2021) Correlations enhance the behavioral readout of neural population activity in association cortex. Nat Neurosci 24:975–986.

Vinci G, Ventura V, Smith MA, Kass RE (2016) Separating Spike Count Correlation from Firing Rate
Correlation. Neural Computation 28:849–881.

Watson BO, Ding M, Buzsáki G (2018) Temporal coupling of field potentials and action potentials in the
neocortex. European Journal of Neuroscience 48:2482–2497.

Wei G, Stevenson IH (2021) Tracking Fast and Slow Changes in Synaptic Weights From Simultaneously
Observed Pre- and Postsynaptic Spiking. Neural computation 33:2682–2709.

Yatsenko D, Josić K, Ecker AS, Froudarakis E, Cotton RJ, Tolias AS (2015) Improved Estimation and
Interpretation of Correlations in Neural Circuits. PLOS Computational Biology 11:e1004083.

Zhai X, Khatami F, Sadeghi M, He F, Read HL, Stevenson IH, Escabí MA (2020) Distinct neural
ensemble response statistics are associated with recognition and discrimination of natural sound
textures. Proceedings of the National Academy of Sciences 117:31482–31493.

Zhan Y, Halliday D, Jiang P, Liu X, Feng J (2006) Detecting time-dependent coherence between non-
stationary electrophysiological signals—A combined statistical and time–frequency approach.
Journal of Neuroscience Methods 156:322–332.

Zhou P, Burton SD, Snyder AC, Smith MA, Urban NN, Kass RE (2015) Establishing a Statistical Link
between Network Oscillations and Neural Synchrony. PLOS Computational Biology
11:e1004549.

846