Density dependence promotes species coexistence and provides a unifying explanation for distinct productivity-diversity relationships

Liang Xu[1], Christopher A. Klausmeier[2], Emily Zakem[1]

[1] Department of Global Ecology, Carnegie Institution for Science, Stanford, CA, USA
[2] W. K. Kellogg Biological Station, Department of Plant Biology & Integrative Biology, Program in Ecology, Evolution & Behavior, Michigan State University, USA

Correspondence: Liang Xu (lxu@carnegiescience.edu) | Emily Zakem (ezakem@carnegiescience.edu)
Email addresses of other coauthors: klausme1@msu.edu

Running head: Density-dependent mortality promotes coexistence
Keywords: density-dependent mortality, coexistence, resource competition theory, resource consumer model, negative density dependence, productivity-diversity curve
The type of article: Letter
The number of words in the abstract: 149
The number of words in the main text without citing references in text: 4973
The number of words in Text box 1: 435
The number of words in Text box 2: 347
The number of references: 49
The number of figures: 4

Statement of authorship*:* LX and EZ conceived the study. CAK helped refine the idea. LX performed the modeling work, theoretical derivation, and simulations, and analyzed the simulated data. LX, CAK and EZ discussed biological interpretation. LX wrote the first draft of the manuscript, and all authors contributed substantially to revisions.

Data accessibility*:* All data used in this study is generated by simulation. The simulation model and code can be accessed at https://github.com/xl0418/DensityDependenceOnZNGI

Abstract:

Understanding diversity patterns in complex communities, such as microbial consortia, requires a mechanistic framework appropriate for many species. Negative density dependence is often utilized in complex ecosystem models, typically as a density-dependent mortality term for a population, but its full impact on community structure remains unclear. Here we use mechanistic population models of resource consumption to examine the effects of negative density dependence and develop a tractable framework for understanding diversity patterns in complex systems. To provide mechanistic grounding, we quantify how density-dependent mortality expands coexistence zones along resource gradients in simple communities using graphical analysis. We then derive an analytical, ecologically insightful formula predicting species abundances in subsets (guilds) of complex communities, in which many species share a resource or predator. Finally, we use the formula to explain how distinct relationships between productivity and diversity emerge from the resulting mechanistic framework, providing insights into previously unreconciled observed patterns.

1. Introduction

Since Hutchinson articulated the "paradox of the plankton" (Hutchinson 1961), numerous hypotheses have been proposed to explain the coexistence of many plankton species competing for a limited number of resources. Resource competition theory (RCT), a foundational framework in ecology, has deepened our understanding of plankton diversity (Tilman 1982; Litchman & Klausmeier 2001; Klausmeier *et al.* 2004; Litchman & Klausmeier 2008). However, RCT analysis is typically restricted to simplified food web structures (Holt 1977; Holt *et al.* 1994; Chase & Leibold 2003), where its graphical techniques are not easily extended to more than a few (< 3) resources and species. Microbial communities, which contain thousands of species (or species-like equivalents) consuming diverse substrates (Moran *et al.* 2022), therefore present a significant challenge to the application of this theoretical framework. To understand species coexistence and diversity patterns in such complex communities, we need a general, comprehensive, and mechanistic framework applicable to diverse systems with many resources and species.

Any such framework must consider the impact of negative density dependence, the suppression of a species' net growth rate as its density increases due to multiple potential biological processes, such as viral lysis (Winter *et al.* 2010; Thingstad 2022), host-specific parasitism (McPeek 1998), and soil pathogen interactions (Bagchi *et al.* 2014; Comita *et al.* 2014; LaManna *et al.* 2017a; Adler *et al.* 2018; Tuck *et al.* 2018). Though phenomenological, density-dependent mortality in mathematical descriptions of population growth captures processes that are too uncertain or complex to model explicitly, and so is often utilized in studies of complex ecosystems (Bagchi *et al.* 2014; Fricke & Wright 2017; Adler *et al.* 2018; Bendik

& Dries 2018; Xu *et al.* 2022). Additionally, density-dependent mortality is a well-established

mechanism for resolving the paradox of the plankton because it facilitates the persistence of

inferior competitors (Chesson 2000; Thingstad 2000).

Although density-dependent mortality is widely invoked to explain biodiversity patterns

(Holt & Lawton 1993; Winter *et al.* 2010; Bagchi *et al.* 2014; Comita *et al.* 2014; LaManna *et*

al. 2017b), theoretical studies, grounded in mechanistic resource competition models, remain

limited relative to empirical studies. Thus, a key question remains: how does density

dependence shape species coexistence and relative abundance patterns along resource gradients?

Although theory has shown that density dependence can stabilize coexistence and allow more

species to persist than there are limiting resources (Leibold 1998; Adler *et al.* 2007; McPeek

2012, 2019), most analyses rely on low-dimensional isocline approaches, leaving its role in

more diverse systems poorly understood.

A broader question is how biodiversity responds to productivity and how this relationship

varies. Different productivity–diversity relationships (PDRs) have been observed, with

monotonic increases and unimodal (hump-shaped) curves common (Mittelbach *et al.* 2001).

For example, Fraser et al. (2015), analyzing 30 grassland sites on six continents, found strong

support for a unimodal PDR, with richness peaking at intermediate productivity at regional and

global scales. However, Adler et al. (2011) sampled 48 herbaceous plant communities across

five continents and found no consistent relationship: while some individual sites showed linear

or hump-shaped trends, the overall pattern lacked coherence.

These observed PDRs are not easily explained by classic resource competition theory

alone, since the number of distinct resources sets a limit on the number of stably coexisting

species, and it is unlikely that resource diversity increases along productivity gradients at the

same pace as species diversity. Additional mechanisms are therefore required. Proposed

mechanistic explanations include spatial or temporal heterogeneity (e.g., due to dispersal or

time-varying resource supply) (Abrams 1988; Leibold 1996), sampling limitations (Pastor *et al.*

1996), and spatial scale dependency (Steiner & Leibold 2004), but any one of these mechanisms

cannot explain why multiple distinct patterns may arise. Density-dependent mortality may

constitute another potential mechanistic explanation, but this has not yet been rigorously

evaluated. Exploring this potential requires a theoretical framework that can mechanistically

link PDRs to trophic interactions and ecosystem structure that is applicable to a large number

of species. Ideally, such a framework would also explain how different PDRs may emerge from

a general description of ecosystem dynamics.

Here, we develop a unified framework that links density dependence to both species

coexistence and PDRs. We begin by extending classic resource competition theory to reveal

how density dependence modifies coexistence criteria, reshaping resource consumption vectors

and supply points. We use well-established, simple food web modules (Holt 1997) to provide

this mechanistic grounding, illustrating how density dependence enables more species to

coexist on limited resources (Section 2). Building on this foundation, we then derive general

formulas for species abundances within analytically tractable subsets of a complex community:

a bottom-up (one shared resource) or a top-down (one shared predator) guild (Section 3). These

formulas are applicable to systems with many species, such as diverse microbial consortia with

one dominant predator. Finally, we demonstrate how this framework, combined with trait trade-

offs, captures multiple PDRs in consistency with observations, which unifies coexistence

theory and PDRs within one mechanistic description (Section 4).

2. Graphical analysis of simple systems

We begin by exploring how density-dependent mortality affects species coexistence in

simple systems using graphical analysis, building on previous work (Leibold 1996; Chase &

Leibold 2003). Specifically, we quantify how density-dependent mortality expands the

coexistence domain in three well-known food web modules: (1) two species competing for two

substitutable resources, (2) three generalists competing for two resources, and (3) a diamond

food web with two prey species and one shared predator.

2.1 Two species and two resources

123   We first consider a food web of two species competing for two nutritionally substitutable

124   resources. Assuming linear functional responses, species i with density N_i grows at rate μ_{ij}

125   on resources R_j, supplied with concentration s_j and dilution rate a:

$$\frac{1}{N_1}\frac{dN_1}{dt} = \mu_{11}R_1 + \mu_{12}R_2 - m - m_q N_1 \qquad 1a$$

$$\frac{1}{N_2}\frac{dN_2}{dt} = \mu_{21}R_1 + \mu_{22}R_2 - m - m_q N_2 \qquad 1b$$

$$\frac{dR_1}{dt} = a(s_1 - R_1) - V_{11}N_1 R_1 - V_{21}N_2 R_1 \qquad 1c$$

$$\frac{dR_2}{dt} = a(s_2 - R_2) - V_{12}N_1 R_2 - V_{22}N_2 R_2. \qquad 1d$$

Here V_{ij} are consumption rates with yields $Y_{ij} = \mu_{ij}/V_{ij}$. Density-dependent mortality for

species i with density N_i is modeled as $m_{qi}N_i^{x-1}$, where $x > 1$. For simplicity, we use

quadratic mortality ($x = 2$) in the main text. We generally assume equal linear mortality rates

(m) and quadratic mortality constants (m_q). Species-specific cases can be rescaled to the

general framework, providing quantitative but not qualitative variability to results

(Supplementary Material).

Without density-dependent mortality ($m_q = 0$), coexistence requires each species to be

competitively superior on one limiting resource, represented by intersecting ZNGIs (zero net

growth isoclines) (Ryabov & Blasius 2011; McPeek 2014; Kleinhesselink & Adler 2015; Letten

et al. 2017) (Fig. 1b). Density-dependent mortality does not alter invasion criteria—resource

levels must still lie above a species' invasion ZNGI where $N_i = 0$—but shifts equilibrium

resource levels ($\widehat{N}_1 > 0$) to depend on species densities and thus on supply rates. Equilibrium

ZNGIs are:

$$\mu_{11}\widehat{R}_1 + \mu_{12}\widehat{R}_2 = m + m_q\widehat{N}_1 \qquad\qquad 2a$$

$$\mu_{21}\widehat{R}_1 + \mu_{22}\widehat{R}_2 = m + m_q\widehat{N}_2. \qquad\qquad 2b$$

where equilibrium conditions are denoted by $\widehat{\blacksquare}$'s. Unlike invasion ZNGIs, these equilibrium

ZNGIs with $\widehat{N}_1 > 0$ vary with biomass and elevate equilibrium resource levels (Fig. S2). As

supply increases, mortality rises with density, raising resources. Strong competitors benefit

from increased growth but also suffer higher mortality, while weaker ones, at lower densities,

experience less mortality. This asymmetry broadens the coexistence domain beyond what is

possible under pure resource competition (Leibold 1998; McPeek 2019).

We next quantitatively analyze the impact of density-dependent mortality on the boundaries

between competitive exclusion and coexistence regions using mutual invasibility analysis

(Chesson 2000) (Fig. 1c) (Supplementary Material). Density-dependent mortality significantly

widens coexistence ranges and produces nonlinear boundaries between competitive exclusion

and coexistence regions (Fig. 1c). The lower boundary of the coexistence zone curves

downward compared to the curve of $m_q = 0$. This indicates that species 2 can invade at a lower

supply rate of resource 2 when species 1 dominates, since species 1 cannot deplete resources to

its invasion ZNGI ($\mu_{11}R_1 + \mu_{12}R_2 = m$) but only to the higher equilibrium ZNGI $\mu_{11}\hat{R}_1 +$

$\mu_{12}\hat{R}_2 = m + m_q\hat{N}_1$. The same principle applies to the upper boundary of the coexistence zone

when species 1 can invade a community with the resident species 2.

In general, a larger m_q results in a larger coexistence zone, indicating that stronger density-

dependent mortality promotes species coexistence across an expanded range of resource supply

rates (Fig. 1c). Box 1 summarizes the four possible competitive outcomes.

Density-dependent mortality reduces the likelihood of priority effects

Notably, density-dependent mortality allows coexistence where two generalists would

otherwise exhibit priority effects (Chesson 2000; Fukami 2015; Grainger *et al.* 2019). In Fig.

1d-f, we selected consumption rates for the two species where each species consumes more of

its less preferred resource ($\frac{v_{12}}{v_{11}} > \frac{v_{22}}{v_{21}}$), normally yielding first-come-first-survive without density

dependent mortality. The striped area on the resource plane in Fig. 1e represents where priority

effects occur: the species that initially establishes itself in the community excludes the other.

Without density-dependent mortality, this region expands linearly with resource supply, with

no intersection of the boundaries (Fig. 1e). In contrast, with density-dependent mortality, the

invasion boundaries intersect at sufficiently high resource supply levels (Fig. 1f). Thus, low

supply maintains the priority effect, but high supply enables coexistence. This finding suggests

that both supply magnitude and ratio govern the ecological outcomes.

2.2 Three species and two resources

We next examine how density-dependent mortality can expand not only the domain of

coexistence but also increase the total number of coexisting species. We consider a food web

with three species and two substitutable resources, with two generalists (species 1 and 2) with

advantages on different resources, and one generalist (species 3) having intermediate

advantages on both resources. Without density-dependent mortality, the system exhibits two

distinct regions of pairwise coexistence. As in the above example, density-dependent mortality

expands the coexistence regions by bending the invasion boundaries (Fig. 1g-i). In this case,

the two coexistence zones may overlap at high resource supply rates, leading to coexistence of

all three species (Fig. 1i).

2.3 The diamond food web: Two species, one predator, and one resource

We finally demonstrate how density-dependent mortality promotes species coexistence in a

different classic system: the "diamond" food web with two species N_i competing for one

resource R and consumed by a common predator P (apparent competition (Holt 1977))as:

190
$$\frac{1}{P}\frac{dP}{dt} = f_1 N_1 + f_2 N_2 - m_p \qquad 3a$$

191
$$\frac{1}{N_1}\frac{dN_1}{dt} = \mu_1 R - m - g_1 P - m_q N_1 \qquad 3b$$

192
$$\frac{1}{N_2}\frac{dN_2}{dt} = \mu_2 R - m - g_2 P - m_q N_2 \qquad 3c$$

193
$$\frac{dR}{dt} = a(s - R) - V_1 N_1 R - V_2 N_2 R, \qquad 3d$$

where f_i is predator growth at predation rate g_i on species i, and m_p denotes the mortality

rate of the predator. Without density-dependent mortality, coexistence requires the two species'

ZNGIs to intersect on the predator-resource plane (Fig. 1k). The stability of this equilibrium

depends on the condition that the species with superior resource-competitive capabilities is

more poorly defended against the predator, while the worse resource competitor is better-

defended against the predator (Leibold 1996). Here, we assume that species 1 is the better

resource competitor and species 2 is better defended against the predator, i.e., $\mu_1 > \mu_2, g_1 >$

g_2.

With density-dependent mortality, the two critical resource supply concentrations are given

by:

204
$$s_{21} = \frac{(g_1 - g_2)m - m_q \dfrac{m_p g_2}{f_1}}{g_1 \mu_2 - g_2 \mu_1} \left(1 + \frac{V_1 m_p}{a f_1} \right),$$
     4

205   and

206
$$s_{12} = \frac{(g_1 - g_2)m + m_q \dfrac{m_p g_1}{f_2}}{g_1 \mu_2 - g_2 \mu_1} \left(1 + \frac{V_2 m_p}{a f_2} \right).$$
     5

where s_{ij} denotes the threshold of the supply rate for species i invading species j

(Supplementary Material). Increasing m_q increases the interval between s_{21} and s_{12} ,

promoting coexistence. We note that the invasion point of species 2 is still bounded by a

minimum supply, which is determined by the intercept of its ZNGI and the supply axis. Unlike

the previous cases (Fig. 1a-i), further increasing resource supply does not necessarily result in

coexistence, as it ultimately goes beyond the coexistence interval, favoring the superior

apparent competitor, species 2 (Fig. 1l). Thus, although density dependence continues to

promote species coexistence, its effect is weakened by apparent competition as resource supply

increases. The community composition then shifts towards species with higher predator

resistance.

Note that the case of no density-dependent mortality ($m_q = 0$) is a special case of these

general equations (Supplementary Material). The four possible competitive outcomes are

summarized in Box 1 (Fig. 1i-l).

3. Control formulas for complex systems

We have quantified how density-dependent effects promote species coexistence, allowing more species to coexist than the number of limiting factors in simple systems. However, graphical analysis becomes impractical for quantitatively extending this approach to communities with more than a few species. We need a more general quantitative approach to mechanistically understand how diversity changes with resource productivity across different food-web structures in more complex communities.

To address this, we consider analytically tractable subsets of communities with many coexisting species. Specifically, we consider two types of subsets, which we refer to as guilds: (1) a bottom-up guild, in which multiple species compete for a single shared resource, while their interactions with higher-level predators remain unspecified (Fig. 2a); and (2) a top-down guild, in which multiple species are preyed upon by a single shared predator, with unspecified details of their resource use (Fig. 2b). This approach aligns with a common and ecologically realistic strategy of analyzing food-web modules—simplified sub-networks where species interactions are governed by basic trophic relationships, such as resource consumption and predation (Holt 1997)—while still allowing considerable complexity in the rest of the food web. For example, tropical forests, which harbor high plant diversity competing for similar resources (Gentry 1988), and marine phytoplankton, which compete for a limited set of inorganic nutrients (Hutchinson 1961), both exemplify bottom-up guilds. Top-down control is evident in freshwater lakes, where invertebrate and vertebrate predators structure distinct prey assemblages (Werner & McPeek 1994). Similarly, microorganisms of comparable size often share predators due to size-based predation (Clauset & Erwin 2008; Ward *et al.* 2012; Follett *et*

al. 2022). Considering these guilds, each with a single shared limiting factor, permits analytical

treatment that would be intractable in fully specified complex networks.

We next develop analytical expressions to characterize coexistence patterns of these guilds,

which allows us to later explore mechanistic relationships between productivity and diversity.

We present a general control formula that results from our derivations of the two guild-specific

control formulas. This general form provides ecological insight into coexistence dynamics,

including the impact of density-dependent mortality (Section 3.1). We then present the specifics

of the bottom-up formula (Section 3.2), noting that the corresponding specifics of the top-down

formula follow a similar derivation and structure (Box 2). We use the bottom-up formula to link

coexistence and productivity-diversity patterns within one framework (Section 3.3).

3.1 The general control formula

We derive control formulas for each of the guilds, which results in one general control

formula for species abundances (see Box 2 for derivation overview and SM section 1). The

resulting general formula describes the steady-state density $\widehat{N}_i$ of species i as:

256
$$\widehat{N}_i = \overline{N} + \frac{(G_i - \overline{G})}{m_q}$$

$$= [\text{average equilibrium density}] + \frac{[\text{ growth advantage/disadvantage of species } i]}{[\text{strength of density-dependent mortality}]} \quad 6$$

The quantity $\overline{N}$ is the appropriately weighted average density, set by the predator or resource

(details below). Thus, $\overline{N}$ is set by the species of the guild that can coexist (i.e., have a positive

steady-state density) at equilibrium, which may be the complete set or a subset of all possible

species. G_i is the net growth rate of species $i,$ which we define as the net growth excluding

the density-dependent term (i.e., $G_i = \sum_h \mu_{ih} R_h - m - \sum_h g_{hi} P_h$). Finally, $\overline{G}$ is the weighted

average growth rate of sustained species in the community, with a specific form determined by

the guild type.

Equation 6 thus explains the equilibrium density of a species as determined by three

interdependent components. The first component, the average density of the community ($\overline{N}$), is

intrinsically linked to the shared limiting factor (resource or predator) defined by the guild type.

The second component is the discrepancy between the net growth rate (G_i) of the focal species

and the weighted community average. When the value of the focal species' G_i surpasses that

of the community average ($\bar{G}$), it represents a relative competitive advantage that elevates the

species' abundance above the average. The third component is the strength of density-

dependent mortality ($\frac{1}{m_q}$), which scales the impact of the second component. When density-

dependent mortality is negligible ($m_q = 0$), all coexisting species must have the same net

growth rate ($G_i - \bar{G} = 0$). In contrast, when density dependence is strong, species with net

growth rates lower than the community average may persist, increasing diversity of the guild

at equilibrium.

3.2 The specific control formula for a bottom-up guild

In a bottom-up guild, multiple consumer species compete for and are predominantly limited

by a single shared resource. Predation is unspecified and thus not the main constraint on total

consumer density (Fig.3a). The system is described as:

280
$$\frac{1}{N_i}\frac{dN_i}{dt} = \mu_i R - m - \sum_h g_{hi} P_h - m_q N_i \qquad\qquad 7a$$

281
$$\frac{dR}{dt} = a(s - R) - \sum_i V_i N_i R. \qquad\qquad 7b$$

282   with parameters as defined above (Eqn. 1). The specific form of the control formula (Eq. 6) for

283   the bottom-up guild is

$$\widehat{N}_i = \frac{a(s-\widehat{R})}{\sum_j^n V_j \widehat{R}} + \frac{1}{m_q}\left(G_i - \frac{\sum_j^n V_j G_j}{\sum_j^n V_j}\right) \qquad 8$$

where the average density is $\bar{N} = \frac{a(s-\widehat{R})}{\sum_j^n V_j \widehat{R}} = \frac{\sum_j^n V_j \widehat{N}_j}{\sum_j^n V_j}$, representing the resource consumed ($a(s -$

$\widehat{R}$)) divided by the total consumption rate of all sustained species ($\sum_j^n V_j \widehat{R}$). Though predation

is not specified, predation does indirectly impact the effective resource level $\widehat{R}$, because

predation reduces prey densities and thereby indirectly influences the equilibrium resource

concentration ($\widehat{R} = \frac{1}{\mu_i}(m + \sum_h g_{hi} \widehat{P}_h + m_q \widehat{N}_i)$). The average community growth rate is

weighted by each species' resource consumption rate $\bar{G} = \frac{\sum_j^n V_j G_j}{\sum_j^n V_j}$, where $G_j = \mu_j R - m -$

$\sum_h g_{hk} P_h$. Thus, Eqn. 8 reveals a trade-off between species density and diversity (which also

applies to the top-down guild). As the strength of density-dependent mortality (m_q) increases,

more species can coexist, but the average density per species declines, as made clear by the

simpler derivation noted in Box 2.

3.3 Linking coexistence and productivity-diversity relationships

The control formula, using the specifics for the bottom-up guild, links resource supply,

traits, and species abundances (and thus diversity) mechanistically, allowing prediction of

diversity patterns as a function of resource supply. For example, Eqn. 8 can be used to determine

the minimum resource supply rate required for a species to persist, as follows. First, we note

that the equilibrium resource concentration is associated with the resource supply rate s (Eqn.

7b and 15) as:

$$s = \widehat{R} + \frac{1}{am_q}\sum_j V_j G_j \widehat{R}. \qquad 9$$

303   given an existing set of species (1 to j). We next consider how species i may invade relative

304   to this rate. Imposing $\widehat{N}_i > 0$ in Eqn. 8, the threshold supply rate for species i corresponds to

305   the condition under which its equilibrium density becomes positive:

$$s_{min,i} = \hat{R} + \frac{1}{am_q} \sum_j V_j \left(G_j - G_i \right) \hat{R} \qquad 10$$

This expression defines the minimum supply rate necessary for species i to persist, given the

presence of other coexisting species. The summation term captures the cumulative effect of

fitness differences weighted by resource uptake rates between species i and other species that

coexist at lower supply rates. Note that $\hat{R}$ is the same in Eqn. 9 and 10 when species i is just

invading ($\hat{N}_i \approx 0$), and thus that $s - \hat{R}$ quantifies the amount of resource consumed by the

community. If $s > s_{min,i}$, the consumed resource exceeds the minimum requirement, allowing

species i to invade and coexist with the established community.

Furthermore, we note that $s = s_{min,i} - \frac{1}{am_q} \sum_j V_j\, G_i \hat{R}$. Thus, when G_i remains positive

across the supply gradient, the coexistence criterion is always satisfied. Conversely, if G_i becomes

negative, the criterion is violated. This yields a tractable condition for quantifying coexistence and

diversity for any specified set of species. Therefore, by examining how species' net growth rates

vary with resource supply, we can directly link resource-driven productivity to biodiversity patterns,

as we next demonstrate.

4. Mechanistic explanations for distinct productivity-diversity relationships

Our mechanistic framework links resource supply rate, which governs the productivity of

the ecosystem, to species richness and diversity. Here, we explore how this provides a

mechanistic basis for the two most commonly observed PDRs: increasing and unimodal. We

identify the factors that determine whether one or the other should emerge and use our results

to interpret observations in new ways.

4.1 Increasing productivity-diversity relationship

We continue to use the control formula, specific for a bottom-up guild, to show how diversity increases with productivity via the resource supply rate. Critically, this makes the assumption that predation has negligible impact on species coexistence patterns. This may, for example, describe phytoplankton communities competing for scarce nutrients with little grazing pressure in oligotrophic systems (Reynolds 1984).

We illustrate the ecological dynamics underlying the increasing PDR clearly by using a specific example of this guild structure with simple tradeoffs between species traits. In a purely bottom-up guild with negligible predation, the net growth rate is $G_i = \mu_i R - m$, which is a monotonically increasing function of the equilibrium resource concentration $\hat{R}$. With density-dependent mortality ($m_q > 0$), the equilibrium resource concentration increases with the supply rate (Supplementary Material, Fig.S1, S3):

$$\frac{d\hat{R}}{ds} > 0. \qquad\qquad 11$$

which in turn gives

$$\frac{dG_i}{ds} = \mu_i \frac{d\hat{R}}{ds} > 0 \qquad\qquad 12$$

Thus, G_i increases with the supply rate, and the coexistence condition $s > s_{min,i}$ always holds once species i has successfully invaded. This mechanism produces an increasing PDR (Fig. 3b).

When density-dependent mortality is negligible ($m_q = 0$), the denominator diverges in Eqn. 10, and any differences in the net growth rates G are effectively amplified, implying that no finite supply rate can support the persistence of an inferior competitor. From the perspective of resource competition, the species with the lowest R^* of the community can deplete the

resource to its R^*, which does not increase with the supply rate. Thus, it prevents the other

species from invading.

More generally, this framework indicates that an increasing PDR is expected whenever the

net growth rate G remains stable or increases with productivity for the majority of species.

This outcome is fundamentally shaped by density-dependent mortality, which acts as the key

regulatory mechanism enabling the pattern. Importantly, this outcome is not restricted to a

purely bottom-up guild without predators. Even when predation is present, increasing PDRs

can still emerge provided that apparent competition is weak, such as if predation is sufficiently

specialized to avoid driving species to extinction through shared predators (Holt 1977). Thus,

increasing PDRs arise not only in pure bottom-up systems but also in predator–prey systems

with specialized or low rates of predation (Fig. 4a, Supplementary Material section 3.5). We

next explore how apparent competition via predation qualitatively changes the shape of PDR.

4.2 Unimodal productivity-diversity relationship

We next illustrate how a qualitatively different, unimodal PDR emerges from the same

general description of ecosystem dynamics, but when the details regarding predation are

different. Specifically, we add a shared predator to the above bottom-up guild, resulting in the

diamond food web structure, the simple but ecologically relevant case where prey species share

a single resource and a common predator (explored using graphical analysis in Section 2.3)

(Levin 1970; Holt *et al.* 1994; Leibold 1996). Thus, in contrast to the bottom-up guild, apparent

competition from shared predation is significant. As in Section 2.3, we assume a trade-off

between intrinsic growth rate and vulnerability to predation, reflecting a range of strategies

from efficient resource exploiters (high μ, high g) to predator-resistant species (low μ, low

g).

In this system, the net growth rate $G_i = \mu_i R - m - g_i P$ is no longer monotonic with

resource supply rate. The derivative of G_i with respect to s takes the form

373
$$\frac{dG_i}{ds} = \left(\mu_i - g_i \frac{\sum_j f_j \mu_j}{\sum_j f_j g_j} \right) \cdot \frac{d\hat{R}}{ds} \qquad 13$$

indicating that G exhibits variable responses to changes in the supply rate s and species

diversity (Supplementary Material).

As above, increasing the supply rate increases the resource concentration, allowing more

species to invade. However, the term $\mu_i - g_i \dfrac{\sum_j f_j \mu_j}{\sum_j f_j g_j}$ decreases when more species that are

more resistant to the predator (with sequentially smaller μ and g than the i^{th} species)

invade (Supplementary Material). Consequently, as the resource supply rate continues to

increase, more predation-resistant species are established, reducing G_i. Eventually, G_i

becomes negative, leading to the exclusion of fast-growing but predator-vulnerable species, and

the coexistence criterion $s > s_{min,i}$ is no longer satisfied. This process unfolds sequentially

along the supply gradient.

Meanwhile, a shared predator can only persist once total prey biomass exceeds a threshold,

$\sum_i f_i N_i > m_p$ (Eqn. 16 in Box 2). Our framework therefore indicates that a sufficient resource

supply is needed to enable predator persistence, leading to apparent competition, which

weakens density dependence among prey. As a result, diversity peaks at intermediate

productivity, giving rise to a unimodal productivity–diversity relationship.

4.3 The emergence of multiple PDRs within one framework

Thus far, we have used the control formula to explain the emergence of both increasing

and unimodal PDRs. Density-dependent mortality is a key mechanism for both, but whether an

increasing or unimodal pattern emerges depends on the community structure, specifically the

strength of apparent competition from a common predator. A structural shift between the two

PDRs could thus result from ecological or evolutionary changes that lead to predator extinction,

colonization, or increasing predator specialization. This suggests a testable hypothesis: in

systems where predation pressure declines (e.g., through predator loss), diversity should shift

to increase monotonically with productivity. We are not aware of any experimental study that

directly tests this prediction, but this hypothesis could be tested with ecosystem manipulation

experiments where predators are added or removed systematically.

More generally, our framework indicates that the strength of apparent competition from

shared predation, not solely its presence or absence, has a strong control on the emergent PDR.

We next illustrate the transition from the increasing to the unimodal PDR as a function of this

control. We simulate ecosystems where species compete for a single limiting resource while

varying the number of predator species (Fig. 4a). We assume that each predator targets a distinct

group of prey (Fig. S5), with adjacent predators sometimes overlapping in prey species. When

the number of predators is small (<10 in our simulations), apparent competition from shared

predation is strong, and the unimodal productivity–diversity curve emerges (Fig.4a). As the

number of predators increases and their diets become more specialized, apparent competition

is reduced, and the curve shifts toward an increasing pattern, eventually converging to the

pattern seen in bottom-up guilds where predation is entirely negligible. Thus, species-specific

predation (when the number of predators reaches 100 in our simulations) also generates an

increasing PDR. Stronger density dependent mortality modulates the quantification of this

curve and generally promotes higher species diversity at lower levels of productivity (Fig. S6).

4.4 Insights into observed patterns

Finally, we relate our framework and analysis of PDRs to observations. We analyze two

plant community datasets that report contrasting PDRs (Fig. 4, (Adler *et al.* 2011; Fraser *et al.*

2015)). We pool all data together across sites, with total biomass used as a proxy for

productivity (Supplementary Material). We find that statistically, the data from Adler et al. 2011

supports a significantly increasing PDR (the dashed line in Fig. 4b), whereas the data from

Fraser et al. 2015 exhibits a unimodal (hump-shaped) relationship (the dashed line in Fig. 4c).

However, despite these differences, the patterns of the two datasets pooled across sites seem

visually indistinguishable, highlighting the difficulty of discerning the PDR based on empirical

data alone. Additionally, both original studies found that within-site regressions reveal a

diversity of patterns, which we also recovered, with increasing ("positive linear" in Fig. 4bc)

and unimodal ("concave-down") patterns as well as decreasing and concave-up patterns, and

that many sites indicated no significant relationship.

Our framework provides insight into these variable PDRs (Fig. 4). For one, our simulations

help to make it obvious that sampling only part of the productivity gradient can produce diverse

PDR patterns (Fig. 4a). For example, a decreasing relationship would be inferred if only the

"right-hand-side" component of the unimodal curve were observed. This sub-sampling could

explain much of the within-site PDR variability. Overall, our simulations suggest three general

patterns. First, unimodal PDRs are most likely to emerge when productivity spans a wide range,

because the persistence of a common predator requires a sufficient level of productivity.

Resolving the unimodal curve empirically thus requires sampling across a sufficiently broad

range, including sufficiently low, intermediate, and high levels of productivity. Second, increasing PDRs are more likely to be observed at low productivity levels in food webs, even with common predators (representing the "left-hand side" of a unimodal curve), as well as across a wide range in productivity in food webs with specialized predators or no predators. Third, declining PDRs tend to occur at relatively high productivity levels, where apparent competition is dominant throughout the entire range. In summary, we posit that density-dependent mortality may be the mechanism underlying many emergent PDRs, with their differences controlled by the strength of apparent competition from shared predation.

5. Outlook

We have presented a quantitative framework for understanding how density-dependent mortality shapes community structure and diversity. By extending resource competition theory, we quantified the specifics of how density dependence reshapes the boundary between competitive exclusion and coexistence, enlarging coexistence zones along resource gradients. Moving beyond graphical analyses of a few species, we derived a general control formula for two fundamental types of guilds within complex ecosystems: bottom-up guilds, where species share a resource, and top-down guilds, where species share a predator. This formula predicts species abundance distributions and attributes the abundance of any species to three mechanistic components: the average biomass set by the shared resource or predator, its deviation from the community-averaged net growth rate, and the strength of density-dependent mortality.

Our modular approach, examining tractable subsets (guilds) of complex ecosystems, also

456 has practical implications. Empirically, the net growth rate can be estimated as the coefficient

457 of the linear term in fits of species densities from time-series observations. For example (Eqn.

458 14), the net growth rate G_i corresponds to the linear term in the following formulation:

459
$$\frac{dN_i}{dt} = G_i N_i - m_q N_i^2.$$
14

These parameters can in principle be estimated from time-series data without requiring detailed

knowledge of specific trophic interactions. This allows our control formula to forecast the

population levels of coexisting species using empirical datasets.

We used our control formula to provide a unifying explanation for the emergence of

multiple diversity–productivity curves, depending on the food web details. We found that

density-dependent mortality is a mechanism that may underly PDRs, and we outlined the

dynamics by which this mechanism may result in the emergence of the two most commonly

observed forms – increasing and unimodal curves. This obviates the need to rely on

environmental heterogeneity and other explanations as mechanisms (though they may also

contribute). When predation is negligible or highly specialized, and thus apparent competition

is negligible, increasing resource supply allows more species to persist, resulting in a monotonic

increase in species richness with productivity. In contrast, when density is additionally

constrained by shared predation, generating apparent competition, diversity peaks at

intermediate productivity.

More generally, our results contribute to the necessary work of extending theoretical

analysis of ecosystem structure to complex ecosystems, such as microbial communities with

thousands of species. The control formula provides a tractable framework for predicting species

abundances based on traits and measurable quantities, rather than phenomenological interaction

coefficients used by the Lotka-Volterra equations. In this way, our mechanistic framework

paves the way towards predictive capability in complex ecosystems.

Acknowledgements: EJZ acknowledges support from NSF (Grant #2125142) and the Simons

Foundation Early Career Investigator in Aquatic Microbial Ecology and Evolution Award. LX

and EZ acknowledge the Carnegie Institution for Science for funding and support. CAK was

supported by National Science Foundation grant EF-2124800.

References

1.
Abrams, P.A. (1988). Resource Productivity-Consumer Species Diversity: Simple Models of
Competition in Spatially Heterogeneous Environments. *Ecology*, 69, 1418-1433.
2.
Adler, P.B., HilleRisLambers, J. & Levine, J.M. (2007). A niche for neutrality. *Ecology Letters*, 10, 95-
104.
3.
Adler, P.B., Seabloom, E.W., Borer, E.T., Hillebrand, H., Hautier, Y., Hector, A. *et al.* (2011).
Productivity Is a Poor Predictor of Plant Species Richness. *Science*, 333, 1750-1753.
4.
Adler, P.B., Smull, D., Beard, K.H., Choi, R.T., Furniss, T., Kulmatiski, A. *et al.* (2018). Competition and
coexistence in plant communities: intraspecific competition is stronger than interspecific
competition. *Ecol Lett*, 21, 1319-1329.
5.
Bagchi, R., Gallery, R.E., Gripenberg, S., Gurr, S.J., Narayan, L., Addis, C.E. *et al.* (2014). Pathogens
and insect herbivores drive rainforest plant diversity and composition. *Nature*, 506, 85-88.
6.
Bendik, N.F. & Dries, L.A. (2018). Density-dependent and density-independent drivers of
population change in Barton Springs salamanders. *Ecol Evol*, 8, 5912-5923.
7.
Chase, J.M. & Leibold, M.A. (2003). *Ecological Niches: Linking Classical and Contemporary*
Approaches. University of Chicago Press.
8.
Chesson, P. (2000). Mechanisms of Maintenance of Species Diversity. *Annual Review of Ecology*
and Systematics, 31, 343-366.
9.
Clauset, A. & Erwin, D.H. (2008). The evolution and distribution of species body size. *Science*, 321,

399-401.

10.

Comita, L.S., Queenborough, S.A., Murphy, S.J., Eck, J.L., Xu, K., Krishnadas, M. *et al.* (2014). Testing predictions of the Janzen-Connell hypothesis: a meta-analysis of experimental evidence for distance- and density-dependent seed and seedling survival. *J Ecol*, 102, 845-856.

11.

Follett, C.L., Dutkiewicz, S., Ribalet, F., Zakem, E., Caron, D., Armbrust, E.V. *et al.* (2022). Trophic interactions with heterotrophic bacteria limit the range of Prochlorococcus. *Proc Natl Acad Sci U S A*, 119.

12.

Fraser, L.H., Pither, J., Jentsch, A., Sternberg, M., Zobel, M., Askarizadeh, D. *et al.* (2015). Worldwide evidence of a unimodal relationship between productivity and plant species richness. *Science*, 349, 302-305.

13.

Fricke, E.C. & Wright, S.J. (2017). Measuring the demographic impact of conspecific negative density dependence. *Oecologia*, 184, 259-266.

14.

Fukami, T. (2015). Historical Contingency in Community Assembly: Integrating Niches, Species Pools, and Priority Effects. *Annual Review of Ecology, Evolution, and Systematics*, 46, 1-23.

15.

Gentry, A.H. (1988). Tree species richness of upper Amazonian forests. *Proceedings of the National Academy of Sciences*, 85, 156-159.

16.

Grainger, T.N., Letten, A.D., Gilbert, B. & Fukami, T. (2019). Applying modern coexistence theory to priority effects. *Proceedings of the National Academy of Sciences*, 116, 6205-6210.

17.

Holt, R.D. (1977). Predation, apparent competition, and the structure of prey communities. *Theoretical Population Biology*, 12, 197-229.

18.

Holt, R.D. (1997). Community Modules. In: *Multitrophic Interactions in Terrestrial Systems* (eds. Grange, AC & Brown, VK). Wiley, British Ecological Society, pp. 549-551.

19.

Holt, R.D., Grover, J. & Tilman, D. (1994). Simple Rules for Interspecific Dominance in Systems with Exploitative and Apparent Competition. *The American Naturalist*, 144, 741-771.

20.

Holt, R.D. & Lawton, J.H. (1993). Apparent Competition and Enemy-Free Space in Insect Host-Parasitoid Communities. *The American Naturalist*, 142, 623-645.

21.

Hutchinson, G.E. (1961). The Paradox of the Plankton. *The American Naturalist*, 95, 137-145.

22.

Klausmeier, C.A., Litchman, E. & Levin, S.A. (2004). Phytoplankton growth and stoichiometry under multiple nutrient limitation. *Limnology and Oceanography*, 49, 1463-1470.

23.

Kleinhesselink, A.R. & Adler, P.B. (2015). Indirect Effects of Environmental Change in Resource
Competition Models. *The American Naturalist*, 186, 766-776.
24.
LaManna, J.A., Belote, R.T., Burkle, L.A., Catano, C.P. & Myers, J.A. (2017a). Negative density
dependence mediates biodiversity-productivity relationships across scales. *Nat Ecol Evol*,
1, 1107-1115.
25.
LaManna, J.A., Mangan, S.A., Alonso, A., Bourg, N.A., Brockelman, W.Y., Bunyavejchewin, S. *et al.*
(2017b). Plant diversity increases with the strength of negative density dependence at the
global scale. *Science*, 356, 1389-1392.
26.
Leibold, M.A. (1996). A Graphical Model of Keystone Predators in Food Webs: Trophic Regulation
of Abundance, Incidence, and Diversity Patterns in Communities. *The American Naturalist*,
147, 784-812.
27.
Leibold, M.A. (1998). Similarity and local co-existence of species in regional biotas. *Evolutionary*
Ecology, 12, 95-110.
28.
Letten, A.D., Ke, P.-J. & Fukami, T. (2017). Linking modern coexistence theory and contemporary
niche theory. *Ecological Monographs*, 87, 161-177.
29.
Levin, S.A. (1970). Community Equilibria and Stability, and an Extension of the Competitive
Exclusion Principle. *The American Naturalist*, 104, 413-423.
30.
Litchman, E. & Klausmeier, C.A. (2001). Competition of Phytoplankton under Fluctuating Light. *The*
American Naturalist, 157, 170-187.
31.
Litchman, E. & Klausmeier, C.A. (2008). Trait-Based Community Ecology of Phytoplankton. *Annual*
Review of Ecology, Evolution, and Systematics, 39, 615-639.
32.
McPeek, M.A. (1998). The Consquences of Changing the Top Predator in a Food Web: A
Comparative Experimental Approach. *Ecological Monographs*, 68, 1-23.
33.
McPeek, M.A. (2012). Intraspecific density dependence and a guild of consumers coexisting on
one resource. *Ecology*, 93, 2728-2735.
34.
McPeek, M.A. (2014). Keystone and Intraguild Predation, Intraspecific Density Dependence, and a
Guild of Coexisting Consumers. *The American Naturalist*, 183, E1-E16.
35.
McPeek, M.A. (2019). Mechanisms influencing the coexistence of multiple consumers and multiple
resources: resource and apparent competition. *Ecological Monographs*, 89, e01328.
36.
Mittelbach, G.G., Steiner, C.F., Scheiner, S.M., Gross, K.L., Reynolds, H.L., Waide, R.B. *et al.* (2001).
WHAT IS THE OBSERVED RELATIONSHIP BETWEEN SPECIES RICHNESS AND

601   PRODUCTIVITY? *Ecology*, 82, 2381-2396.

602   37.

603   Moran, M.A., Kujawinski, E.B., Schroer, W.F., Amin, S.A., Bates, N.R., Bertrand, E.M. *et al.* (2022).

604   Microbial metabolites in the marine carbon cycle. *Nat Microbiol*, 7, 508-523.

605   38.

606   Pastor, J., Downing, A. & Erickson, H.E. (1996). Species-Area Curves and Diversity-Productivity

607   Relationships in Beaver Meadows of Voyageurs National Park, Minnesota, USA. *Oikos*, 77,

608   399-406.

609   39.

610   Reynolds, C.S. (1984). *The Ecology of Freshwater Phytoplankton*. Cambridge University Press.

611   40.

612   Ryabov, A.B. & Blasius, B. (2011). A graphical theory of competition on spatial resource gradients.

613   *Ecology Letters*, 14, 220-228.

614   41.

615   Steiner, C.F. & Leibold, M.A. (2004). CYCLIC ASSEMBLY TRAJECTORIES AND SCALE-DEPENDENT

616   PRODUCTIVITY–DIVERSITY RELATIONSHIPS. *Ecology*, 85, 107-113.

617   42.

618   Thingstad, T.F. (2000). Elements of a theory for the mechanisms controlling abundance, diversity,

619   and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnology and*

620   *Oceanography*, 45, 1320-1328.

621   43.

622   Thingstad, T.F. (2022). Competition–defense trade-offs in the microbial world. *Proceedings of the*

623   *National Academy of Sciences*, 119, e2213092119.

624   44.

625   Tilman, D. (1982). Resource competition and community structure. *Monogr Popul Biol*, 17, 1-296.

626   45.

627   Tuck, S.L., Porter, J., Rees, M. & Turnbull, L.A. (2018). Strong responses from weakly interacting

628   species. *Ecol Lett*, 21, 1845-1852.

629   46.

630   Ward, B.A., Dutkiewicz, S., Jahn, O. & Follows, M.J. (2012). A size-structured food-web model for

631   the global ocean. *Limnology and Oceanography*, 57, 1877-1891.

632   47.

633   Werner, E.E. & McPeek, M.A. (1994). Direct and Indirect Effects of Predators on Two Anuran Species

634   along an Environmental Gradient. *Ecology*, 75, 1368-1382.

635   48.

636   Winter, C., Bouvier, T., Weinbauer, M.G. & Thingstad, T.F. (2010). Trade-offs between competition

637   and defense specialists among unicellular planktonic organisms: the "killing the winner"

638   hypothesis revisited. *Microbiol Mol Biol Rev*, 74, 42-57.

639   49.

640   Xu, L., Clark, A.T., Rees, M. & Turnbull, L.A. (2022). Estimating competition in metacommunities:

641   accounting for biases caused by dispersal. *Methods in Ecology and Evolution*, 00.

Food web structure	Without density dependent mortality	With density dependent mortality

Figure 1. : Illustration of the range of resource supply rates for the species coexistence with and without the density-dependent effect. The impact vectors of species on resources are denoted by c_1, c_2, c_3. (a), (d), (g), (j). The food web structures. (b). The two generalists can coexist when $m_q = 0$; (d). Increasing m_q expands the range of resource supply rates for species coexistence. The model parameters are $\mu_{11} = 1.0, \mu_{21} = 0.2, \mu_{12} = 0.5, \mu_{22} = 0.5, V_{11} = 3.3, V_{21} = 0.67, V_{12} = 1.67, V_{22} = 1.67, m = 0.1, a = 1$; (e). The two generalists cannot coexist when $m_q = 0$ because their impact vectors indicate a priority effect; (f). The effect of density dependence facilitates coexistence at higher supply rates. For the supply rates vector falls in the striped zone, the priority effect emerges. The model parameters are $\mu_{11} = 1.0, \mu_{21} = 0.2, \mu_{12} = 0.5, \mu_{22} = 0.5, V_{11} = 1.1, V_{21} = 1.8, V_{12} = 2.2, V_{22} = 0.6, m = 0.1, a = 1$. (h). When $m_q = 0$, at most two species can coexist in the supply rate space. The outcome of species interaction depends on where the vector of resource productivities falls.; (i). When $m_q = 0.01$, the two pairwise coexistence zones overlap, indicating an area where three species can coexist. The model parameters are $a = 1, \mu_{11} = 0.27, \mu_{22} = 0.24, \mu_{13} = 0.15, \mu_{23} = 0.15, V_{11} = 0.9, V_{22} = 0.8, V_{12} = 0.5, V_{22} = 0.5, m = 0.1, a = 1$. (k-l). In the diamond food web, two species can coexist at a larger range of supply rates with increase of m_q; (k).$m_q = 0$; (l). $m_q = 0.01$; The boundaries are given by Eqns. 4 and 5. The model parameters are $\mu_1 = 2.8, \mu_2 = 1.2, V_1 = 4.2, V_2 = 1.8, g_1 = 8.7, g_2 = 1.725, f_1 = 5.8, f_2 = 1.15, m = 0.1, m_p = 0.1, a = 1$.

Figure 2: Two types of guilds illustrating species interactions within and between trophic levels. The negative density dependence is denoted by the red arrows. (a). The bottom-up guild assumes that consumer species are mainly limited by food availability of one common resource; (b) The top-down guild assumes that consumer species are restricted by the common predator. Predators are denoted by P; consumer species are denoted by N; resources are denoted by R.

Figure 3: Negative density dependence generates contrasting productivity–diversity relationships. (a) Simulation of a bottom-up guild with 100 species ranked by growth rates (color gradient). (b) In the presence of density-dependent mortality, species richness increases with resource supply. (c) Simulation of a diamond food web with 100 species, sharing a common predator and resource. Species are ranked by growth and predation rates. (d) Under density-dependent mortality, the resulting productivity–diversity relationship becomes unimodal. As in Section 2, species are ordered by decreasing intrinsic growth rates ($\mu_i > \mu_{i+1} > \cdots > \mu_k$) for simulations in (a) (b) and with additional condition $\frac{\mu_i}{g_i} < \cdots < \frac{\mu_k}{g_k}$ for simulations in (c) and (d).

Figure 4: Density-dependent mortality generates productivity–diversity relationships (PDRs). (a). Simulated PDRs in a food web with 100 prey species and varying number of predators (NOP=0,…, 100). Different PDR shapes emerge at different ranges of resource supply rates (indicated by colored lines). Solid green lines: increasing PDRs; solid yellow lines: unimodal PDRs; solid brown lines: decreasing PDRs; solid blue lines: concave up PDRs. (b); (c). Empirical PDRs compiled from Adler et al. 2011 and Fraser et al. 2015 (see Supplementary Material). Solid lines show within-site regressions; dashed lines indicate regressions pooled across sites. Inset panels show the distribution of observed PDR types: NS = non-significant, Linear+ = positive linear, Linear− = negative linear, Concave+ = anti-unimodal, and Concave− = unimodal relationships.

Text box 1:

In a food web with two species competing for two nutritionally substitutable resources, density-dependent mortality affects the four regions of the resource plane as follows (Fig. 1a-c):

- **Region of no species**: This zone is bounded by the invasion ZNGIs of the two species and two resource axes. In this zone, the supply of both resources is severely limited and cannot support the persistence of either species. This zone is not changed when incorporating density-dependent mortality because the minimum requirement of resources is not impacted: the invasion ZNGIs remain constant as Eqn. 1a-b do not depend on m_q when $N_i = 0$.

- **Regions of competitive exclusion**: In these regions, resource 1 (or 2) is sufficiently supplied, which allows species 1 (or 2) to maintain its presence. In contrast, the limited supply of resource 2 (or 1) hinders the survival of species 2 (or 1), resulting in its exclusion. The presence of density-dependent mortality narrows this zone and allows species 2 (or 1) to eventually persist with the increase of the supply concentration of resource 2 (or 1).

- **Region of coexistence**: This region is bounded by two boundaries characterized by adequate resource supply levels for both resources. Consequently, it creates an environment where both species can coexist. Density-dependent mortality significantly expands this zone (Fig. 1c).

In a food web with two species N_i competing for one resource R and consumed by a common predator P, density-dependent mortality affects the four regions of the resource plane as follows (Fig. 1j-l):

- **Region of no species**: When $s < R_1^* = \frac{m}{\mu_1}$, resource supply is insufficient to meet the minimum requirement for either species, even for the best resource exploiter. No species can persist in this range of resource supply rate. Incorporating density-dependent mortality does not change this threshold.

- **Region of species 1 winning**: When $R_1^* < s < s_{21}$, species 1 can persist, but not species 2. Specifically, in the range of $R_1^* < s < R_2^* = \frac{m}{\mu_2}$, species 2 cannot persist because the resource supply concentration is below its minimum requirement. When $R_2^* \leq s < s_{21}$, species 2 is excluded by competition with species 1. Density-dependent mortality decreases s_{21} (Eqn. 4), which allows species 2 to invade at a lower resource supply compared to the case of no density-dependent mortality.

- **Region of coexistence**: When $s_{21} < s < s_{12}$, both species can coexist. The effect of density-dependent mortality expands the coexistence region through increasing s_{12} and decreasing s_{21} with increasing m_q (Fig. 1k,l).

- **Region of species 2 winning**: In the last region where $s > s_{12}$, only species 2 can survive. Increasing resource supply raises the total density of consumer species, resulting in an increase in the density of the shared predator. This apparent competition leads to the exclusion of species 1.

Text box 2: The control formula derivation and detail

Derivation overview: There are two ways to derive the density of species at equilibrium. The most direct approach is to solve Eqn. 7 for equilibrium. Specifically, at equilibrium, the density of the consumer can be obtained by setting Eqn. 7a to 0:

$$N_i = \frac{1}{m_{qi}}\left(\mu_i(R_1, \cdots, R_l) - m_i - g_i(P_1, \cdots, P_h)\right) \quad 15$$

This equation shows that consumer density increases with gains from resources but decreases due to linear mortality and predation. Moreover, density is inversely proportional to the nonlinear mortality rate. While this formulation is broadly applicable in general food webs, it does not reveal how interactions among species within trophic levels contribute to changes in density.

To address this, we instead apply a contrast-based decomposition. In this approach, we derive formulas specifically for the bottom-up and top-down guilds (see Supplementary Material Section 1) and then present a general control expression as the shared functional form. The resulting equation differs from Eqn. 15—particularly in the weighting terms. The advantage of this derivation – our control formula – provides a richer ecological interpretation, as discussed in the main text and summarized in Eqn. 8 and 18.

The control formula for top-down trophic guilds: Both the bottom-up and top-down guild control formula have the same functional form, the general control formula. In the main text, we provide the specifics for the bottom-up guild, and here we provide the analogues specifics for the tow-down guild. In a top-down trophic guild, all species are preyed upon and largely controlled by a common predator, with unspecified resources that do not constrain total density of consumer species, given by (Fig. 2b):

$$\frac{1}{P}\frac{dP}{dt} = \sum_i f_i N_i - m_p \quad 16$$

$$\frac{1}{N_i}\frac{dN_i}{dt} = \sum_h \mu_{ih} R_h - g_i P - m - m_q N_i. \quad 17$$

In this case, the explicit form of the control formula Eq. 7 is given by

$$\widehat{N}_i = \frac{m_p}{\sum_j^n f_j} + \frac{1}{m_q}\left(G_i - \frac{\sum_j^n f_j G_j}{\sum_j^n f_j}\right) \quad 18$$

where the average biomass density reflects the control from the predator as $\overline{N} = \frac{m_p}{\sum_j^n f_j} = \frac{\sum_j^n f_j \widehat{N}_j}{\sum_j^n f_j}$, which is the subsistence concentration of the predator, or the mean density $\widehat{N}_j$ weighted by the predation rate f_j. The average community growth rate is weighted by the predator's assimilation (consumption) rate on each species f as $\overline{G} = \frac{\sum_j^n f_j G_j}{\sum_j^n f_j}$, where $G_i = \sum_h \mu_{ih} R_h - g_i P - m$. Note that we assume a static (non-oscillatory) equilibrium state, though oscillatory equilibria can be treated by averaging species densities over cycles.