Terrain ruggedness and human activities influence the distribution of Grauer's

gorillas (*Gorilla beringei graueri*) and eastern chimpanzees (*Pan troglodytes*

***schweinfurthii*) in the Tayna Nature Reserve, Democratic Republic of Congo**

Alice Michel[1], Jackson Kabuyaya Mbeke[2], Benezeth Kambale Visando[2,3], Sonya Kahlenberg[2], Katie

Fawcett[2], Damien Caillaud[1]

[1]Department of Anthropology, University of California, Davis, USA

[2]Gorilla Rehabilitation and Conservation Education Center (GRACE), Kasugho, North Kivu, DRC

[3]Tayna Nature Reserve, Kasugho, North Kivu, DRC

Corresponding author email address: aljmichel@ucdavis.edu

Corresponding author address:

Alice Michel

Department of Anthropology

1 Shields Ave

University of California, Davis,

Davis, CA 95616 USA

Keywords: Great apes, survey, transect, critically endangered, gorilla, chimpanzee, distribution

Abstract

The Albertine Rift Hotspot in east-central Africa is one of the most biodiverse regions in the world. It hosts a number of conservation-priority species, including Critically Endangered Grauer's gorillas and Endangered eastern chimpanzees. Yet, prolonged insecurity in the region has made wildlife monitoring and conservation challenging. This is particularly true of the northeastern limits of the range of Grauer's gorillas, where wide stretches of unprotected forest could harbor behaviorally and genetically unique peripheral populations of great apes and other species. Here, we developed a rapid wildlife survey method to map population distributions and monitor the impact of various human activities in the Tayna Nature Reserve in eastern Democratic Republic of Congo. Despite a lack of surveys since the early 2000s, we find that Grauer's gorillas and eastern chimpanzees, as well as a number of smaller-bodied species, have persisted in Tayna. Observations of gorilla and chimpanzee signs were highly heterogeneous. They were less dense near human settlements and artisanal mines and peaked in areas with rugged terrain. The association with rugged terrain may be a result of historical anthropogenic pressure, behavioral avoidance of human activities, and/or ecological preference. Maintaining connectivity between patches of suitable great ape habitat will be critical for long-term great ape conservation in the Albertine Rift region. Our findings also demonstrate the utility of priority species monitoring driven by local communities for conservation initiatives.

1. Introduction

Habitat loss, overhunting, and climate change are the predominant threats to terrestrial wildlife globally (Hogue and Breon, 2022). Yet wildlife populations vary in their response to human activities, depending on the type of activity, local context, and species. For example, whether hunting is sustainable depends on local ecological processes and prey behavioral responses (Ripple et al., 2016). In some cases, wildlife persists in or recolonizes areas with high hunting pressure (*e.g.*, African forest elephant, *Loxodontis cyclotis*, in a logging concession in Gabon; Van Vliet and Nasi, 2008; cougar, *Puma concolor,* in hunting areas in the USA; Robinson et al., 2008; African wild dog, *Lycaon pictus* in the buffer zone around Hwange National Park, Zimbabwe; van der Meer et al., 2014), while in other cases, wildlife avoids them (*e.g.,* African forest elephant in high human use areas within Dzanga-Sangha Protected Area Complex, Central African Republic; Blom et al., 2005; black bear, *Ursus americanus*, and elk, *Cervus canadensis*, in hunting areas in the USA; Stillfried et al., 2015; Spitz et al., 2019). Other activities that do not directly target wildlife may still cause avoidance, as for leopards (*Panthera pardus*) in pastoral lands in Kenya (Van Cleave et al., 2018) and chimpanzees (*Pan troglodytes*) in logging concessions in northern Congo (Morgan et al., 2018). Populations may even exhibit selective preference for human-dominated habitats: leopards in the Kalahari are more active near farms than protected areas (Van der Weyde et al., 2018) and chimpanzees raid crops in human-dominated landscapes (Hockings and McLennan, 2012; Krief et al., 2014). Finally, some species are ecological generalists (Ramiadantsoa et al., 2018) or specialists that thrive in secondary habitats created, for example, by selective logging, including wildlife targeted by hunting like western lowland gorillas (*Gorilla gorilla gorilla*; Morgan et al., 2018). Identifying how specific threats and habitat preferences interact at the population level is important to the design of conservation plans (Martin, 1998).

Tropical forests, the most biodiverse terrestrial biome, are facing escalating habitat degradation, forest fragmentation, and pressure from bushmeat hunting (Abernethy et al., 2013; Aleman et al., 2018; Taubert et al., 2018). The Albertine Rift in tropical east-central Africa includes 40% of the continent's mammalian taxa in less than one percent of its total land area (Mittermeier et al., 2004). Though 60% of

the region is under some degree of protection, it faces numerous conservation challenges, including armed

conflicts, deforestation for charcoal, agriculture, mining, and oil extraction, and climate change (Plumptre

et al., 2017). These processes interrelate, amplifying pressure on wildlife; for example, mining and

logging increase hunting pressure in remote forests to feed workers, and hunters use new access roads that

fragment forests to supply growing urban demand for bushmeat (Edwards et al., 2014; Laurance et al.,

2006; Spira et al., 2019). As a result, keystone vertebrate species in the Albertine Rift, including forest

elephants (*Loxodonta cyclotis*), forest buffalo (*Syncerus cafer nanus*), eastern chimpanzees (*Pan*

troglodytes schweinfurthii), and gorillas (*Gorilla beringei*), have experienced population declines and

local extirpations, with long-term consequences on forest ecosystem composition and structure

(Mittermeier et al., 2004; Poulsen et al., 2018). In particular, the critically endangered Grauer's gorilla

(*Gorilla beringei graueri*), endemic to eastern Democratic Republic of Congo (DRC), declined from

~17,000 individuals in the mid-1990s (Hall et al., 1998) to between 4,500 to 9,500 in 2019 (Plumptre et

al., 2021, 2016). In addition to overall loss of numbers, the population has become fragmented (Plumptre

et al., 2021, 2016). Sympatric endangered eastern chimpanzees have experienced a similar situation

(Plumptre et al., 2021, 2016).

However, only a fraction of the range of Grauer's gorillas has been surveyed in the past 20 years

(Plumptre et al., 2021, 2016). In particular, there is a lack of data from the northeastern part of their range

due to regional insecurity and logistical challenges (Hart and Hall, 1996; Plumptre et al., 2015). Recent

range-wide population assessments have relied on extrapolation of occupancy models across different

landscapes, with data centered on protected forests (Plumptre et al., 2021, 2016). This approach may miss

localized effects, especially in areas not well represented by the data in the model such as at the edge of a

species' range, where environmental, anthropogenic, ecological, and behavioral patterns may diverge

from the range center (Sexton et al., 2009).

Because great apes are rare, elusive, and inhabit dense forests, surveying them requires a proxy,

typically the nests that adult individuals construct in new locations most nights (Kühl et al., 2008). To

estimate absolute animal density, surveys commonly rely on "standing crop" nest counts along distance

transects (Kühl et al., 2008). Field teams walk along randomly placed transect lines and record the

perpendicular distance between each nest site and the transect line (Buckland et al., 2015). Given known

nest production and decay rates, nest density can be converted to individual animal density (Buckland et

al., 2015). However, nest decay rates vary significantly among habitats, seasons, height, and the plant

materials used for construction, rendering density estimates based on rates calculated at another site or in

a different habitat unreliable (Morgan et al., 2016; Romani et al., 2023; Tutin et al., 1995; Walsh and

White, 2005). Survey-specific nest decay rates can be obtained with additional effort by locating and

revisiting 50-150 fresh nests, ideally per species, habitat, and nest type stratum (Bessone et al., 2021;

Kouakou et al., 2009; Laing et al., 2003). Similarly, the marked nest count method circumvents the issue

of decay rates entirely, by re-walking transects and only considering fresh nests, but this not only requires

additional time and effort but the appearance along transects of at least 60 fresh nests over the survey

period (Kühl et al., 2008). Both of these options can be difficult, and even impossible, at low great ape

density. Because the production and decay rates of other signs such as footprints, feeding remains, and

dung are unknown, distance analyses are typically restricted to nest count data, leaving aside data on other

signs even though such data are often collected together with nest data (*e.g.*, Plumptre et al., 2021).

A related challenge of the transect method in tropical forests is efficiency. Walking along a

perfectly straight line and measuring precise perpendicular distances from that line is difficult and time

consuming when the vegetation is dense. Field teams can often only walk a maximum of 1-2 km/day

along line transects in central Africa (Caillaud, personal communication). When animal density is low,

the sampling effort required to reach the recommended minimum of 60-80 groups of nests can be

considerable (Kühl et al., 2008). Taking the example of Grauer's gorillas, the exceptional 689 km, multi-

year transect survey conducted by Plumptre et al (2021) only found 118 groups of nests. This means that

observing 60-80 groups of nests required 350-467 km of transects. To increase sample size, authors

sometimes analyze individual nests instead of groups (Carvalho et al., 2013; Haurez et al., 2014; Kouakou

et al., 2009; Stokes et al.; 2010; van der Hoek et al., 2024). However, since nests from a given nest group

are not statistically independent units, this approach underestimates the variance of nest density and the

width of confidence intervals (Buckland et al., 2001; Haurez et al., 2014; Kouakou et al., 2009; Morgan et

al., 2006; Stokes et al.; 2010). This leads to inflated type 1 error when animal densities are compared

across sites, species, or time periods (Buckland et al., 2001).

122   Here, we propose an efficient and rapid method to survey wildlife distribution patterns that

provides relative rather than absolute abundance estimates and is designed to be implemented by

members of local communities. The resulting animal distribution maps and models of ecological

covariates are not subject to errors induced by inaccurate nest decay rates, production rates, average group

size estimates, or low sample size of groups of nests. Because this method, termed the "fast-transect"

method, does not attempt to estimate absolute animal density, it can incorporate a variety of signs of

animal presence in addition to nests to map animal distribution patterns, like dungs, feeding remains,

tracks, and trails. Compared to transect analyses based on groups of nests, it uses more data, which

increases the power of statistical analyses (Buckland et al., 2001). Importantly, fast-transect data

collection does not require the same level of training as stranding crop nest counts, and can be performed

by members of communities with limited access to formal education.

133   The 899 km^2 Tayna Nature Reserve (TNR; **Fig. 1**) lies at the northeastern edge of the historical

range of Grauer's gorillas, at the border of the Lubero and Walikale territories in North Kivu, eastern

DRC. The only published wildlife survey of TNR gorillas covered the central and southern portion of the

reserve in 2003 (Mehlman, 2008). Two surveys were attempted since then, in 2006 and 2013, but neither

was completed due to security issues (Nixon, 2013). Nevertheless, Plumptre et al. (2015) suggested a

crash of the gorilla population in TNR by 89% between 2007 and 2013 based on nest encounter rates.

Additional surveys are urgently needed to determine if a viable population of Grauer's gorillas persists in

TNR. If so, it would be one of the furthest north and east populations of Grauer's gorillas left (aside from

the few individuals at Mount Tshiaberimu, Virunga National Park, DRC; Iyer et al., 2023), which would

warrant improved protection of the reserve itself, as well as the community conservation forests to the

west, the Usala Corridor, that extends towards Maiko National Park 78 km away.

Here, we investigate the spatial distribution, terrain covariates, and impacts of anthropogenic

disturbances on Grauer's gorillas, eastern chimpanzees, guenons (*Cercopithecus spp.*), duikers

(*Cephalophus spp.* and *Philantomba monticola*), and red river hogs (*Potamochoerus porcus*) in TNR. We

conducted a 400-km fast-transect survey covering the entire reserve and leveraged generalized additive

models to examine how wildlife distribution related to terrain and human activities. Data were collected

by five teams of four people recruited from local communities living in and around TNR. We

hypothesized that, if still present, Grauer's gorillas and eastern chimpanzees would occur at lower

densities near human settlements and forest activities such as artisanal mining and snare-trapping. Great

apes are hunted for bushmeat (Ziegler et al., 2016), and even small-scale mining increases hunting

pressure (Edwards et al., 2014; Spira et al., 2019). While it was unclear if large-scale trends applied

locally, previous landscape-scale surveys found negative relationships between great ape density and

distance to villages and mines (Plumptre et al., 2021, 2016). We expected the density of other wildlife

species to also be impacted by human activities, but we predicted that the more resilient, faster-

reproducing red river hogs and duikers would be less affected than slow-reproducing great apes and

guenons (Estrada et al., 2017). Finally, we hypothesized that habitat suitability would vary across the

heterogeneous terrain of TNR due to long-term human impacts and ecological gradients, such as shifts in

plant community composition with altitude (Imani et al., 2016). As such, we predicted that wildlife

distribution would relate to terrain, with different patterns for different species.

2. Methods

2.1. Study area

The Tayna Nature Reserve (TNR; 0° 20' 36" S, 28° 43' 0" E; **Fig. 1**) was created in 2006 at the border of the Lubero and Walikale territories, North Kivu, eastern Democratic Republic of Congo. It encompasses 899 km^2 of closed- and open-canopy primary forest. About 7% of its surface is agricultural land (Global Forest Watch, 2025). It lies in the transition zone between the Congo Basin and the Albertine Rift, generating a heterogeneous mix of high-altitude terrain up to 2,028 m above sea level (asl) stretching along the east of the reserve, with rugged hills and valleys near the center, and lower, flatter areas to the northwest dipping to 919 m asl (**Fig. 2A-B**). The forests of TNR extend continuously to the west to Maiko National Park. Vegetation is typical of low- to medium-altitude primary and secondary equatorial forest, though botanical surveys have not yet been undertaken. Rainfall averages 1,800-2,300 mm per year, with the rainiest season from October to November and a second rainy season from March to April (Congo Basin Forest Partnership, 2006). Temperature ranges from 13-34 ˚C, with a daytime average of 19 ˚C (Gorilla Rehabilitation and Conservation Education Center, unpublished data). The largest human settlements around TNR are to the northeast. Additional villages are found along the eastern and mid-western boundaries, and there are small dispersed settlements and encampments within the reserve (**Fig. 2C**). The survey was completed between September 12 and December 24, 2020.

2.2. Fast-Transect Survey

Compared to distance transects (Buckland et al., 2015), fast-transects do not require measuring perpendicular distances between transect line and signs. Instead, observers are positioned on the transect line and on either side of it, sweeping a ~100m strip of forest and searching for all signs of the species of interest. An important requirement for fast-transects is that the survey line be unbiased by habitat type (unlike reconnaissance surveys), so field teams follow a predefined compass bearing to the transect endpoint. The teams do not need to strictly adhere to this line, though, and can walk around localized

obstacles such as patches of dense vegetation or fallen trees, as long as these deviations (typically less

than 20m) are limited. This makes the data collection faster than traditional distance transects.

We designated 200 linear transects in a horizontally-staggered gridded pattern across the reserve

(**Fig. 1**). Each transect was 2 km long, oriented north to south. Transects were spaced 1.5 km apart east to

west and 750 m north to south. Each team comprised four members, all hired and trained locally: one

team leader, one trail opener who walked ahead and cut minimally through the dense vegetation, and two

observers on either side of the leader. The team leader carried a Garmin 64 handheld GPS, which

automatically recorded points every 50 m. Waypoints for the start and end of each transect and any

obstacles encountered were recorded in the GPS. The two observers "swept" about 50 m to the left and

right of the leader for signs, covering a ~100 m wide strip over each 2-km transect. As our goal was to

estimate relative sign abundances, we did not need these two observers to sweep exactly 50 m of forest.

Rather, they were trained to be consistent in the way they were walking and looking for signs. Observers

remained within sight of the team leader, who recorded GPS points at sign locations. All signs of animal

presence, including nest groups, footprints, feeding remains, dungs, burrows, vocalizations, and direct

observations, were recorded and attributed to a particular species. We followed the method used by

Plumptre et al. (2021) to assign great ape nest sites to gorillas or chimpanzees based on nest height, odor,

dung, hairs, and trails. In addition, we recorded signs of human activities, such as camps, paths, snares,

artisanal mines (operational and abandoned), logging, and crop fields. When signs of equivalent age, type,

and species occurred within 50 m, only the first sign was considered, to reduce statistical non-

independence issues caused by the clustering of signs produced by the same group of animals. Factors

like rain could alter the decay time and detectability of signs, potentially inflating the between-transect

variation in relative sign density. Our statistical approach explicitly modelled between-transect variation

to limit the impact these factors might have had on the precision of our relative density estimates (Wood,

2017).

All data were collected strictly non-invasively in accordance with Congolese laws and regulations

for wildlife research by local community members hired based on their tracking skills. Disturbance to

wildlife during field work was negligible. Team members received extensive training in the fast-transect

methodology at GRACE (Gorilla Rehabilitation and Conservation Education Centre), a research and

conservation center located 7 km east of TNR.

2.3. Animal population distribution models

2.3.1. Data preparation

To evaluate spatial heterogeneity in animal density in TNR, we modeled the probability of

occurrence of animal signs using presence and absence points along transects and absence points

generated randomly, as recommended by Barbet-Massin et al. (2012). To reflect the true sampling area ,

GPS track points were interpolated to a polyline and clipped to the start and end of each transect using

QGIS. We excluded detours around obstacles and GPS errors (deviations over 50 m). Transect polylines

were buffered by 50 m on each side, resulting in 100 m wide polygons covering the area "swept" by field

teams along each transect. We retained only observations located within these polygons for our models,

corresponding to a total survey area of 41.7 km^2. We created *pseudo*-absence points by randomly placing

10,000 points within the polygons, as recommended for analysis of species distribution using generalized

linear and additive models (Barbet-Massin et al., 2012). These *pseudo*-absence points were generated

using function *spsample* in the 'sp' R package (Bivand et al., 2013; Pebesma and Bivand, 2005),

corresponding to a density of 240 points per square kilometer of transect strip.

2.3.2. Modeling framework

We opted for generalized additive models (GAMs) to model our presence/absence data because

of their ability to incorporate nonlinear effects and two-dimensional (spatial) effects (Wood, 2017).

Models were fit in the R package 'mgcv' with a binomial distribution and logit link function. We used

restricted maximum-likelihood to find the optimum smoothing parameters (Wood, 2018). Models were fit

for each of the following animals: Grauer's gorillas, eastern chimpanzees, guenons, duikers, and red river

hogs. We modelled animal sign presence/absence in relation to four smoothed predictor variables:

altitude, ruggedness, proximity to human settlements, and mining density (**Fig. 2**). All smooth terms were

thin-plate regression splines with a basis dimension restricted to five to avoid overfitting. Because we had

a large number of absence points (n = 10,000), model estimated probabilities (p) were very low (less than

0.01). At low p and high n, the binomial distribution converges towards the Poisson distribution (Poisson,

1837). As a consequence, our model's probability estimates scaled perfectly with sign densities, and

therefore, were proportional to animal densities. To add interpretability, we converted model-predicted

probability of occurrence to relative density, measured in observed signs/km^2, by multiplying the model

prediction at each 300 x 300 m raster cell by the average sign density (total number of observed signs

over 41.7 km^2 of transect polygons), divided by the average predicted probability of sign occurrence

across TNR. Hereafter we report modelling results in relative density (predicted number of observed

signs/km^2).

2.3.3. Predictor Variables

To measure the altitude and terrain ruggedness in the region, we downloaded the Shuttle Radar

Topography Mission 1 Arc-Second Global 30 m resolution digital elevation model (DEM) for the tile at

1˚S, 28˚E (USGS, 2015). Altitude (meters asl) was taken at the coordinates of each data point and pseudo-

absence point. The terrain ruggedness index was computed as the mean of the absolute differences

between the altitude at each DEM raster cell, in meters, and its eight adjacent cells (Wilson et al., 2007),

averaged over a radius of one kilometer surrounding each data point. We chose one kilometer to reflect

the average daily distance travelled by Grauer's gorillas at low-altitude during the rainy season (van der

Hoek et al., 2021a).

To evaluate the impact of human settlements, we calculated a "proximity index" that considers

the distance to settlements as well as their relative size, since settlements in TNR ranged from camps to

small villages. To do this, we drew polygons around settlements, fields, and camps visible in satellite

imagery (Google Earth V7.3.3.7786, 2021) in TNR and the surrounding five km around it using QGIS

(v.3.20). We converted the multi-polygon vector layer to a five-meter resolution raster using the R

package 'fasterize' (R Core Team, 2023; Ross, 2020). Next, we converted the raster to a spatial points

object, resulting in a grid of points every five meters over the settlement areas, using R functions

rasterToPoints and *SpatialPoints* in the packages 'raster' package (Hijmans et al., 2013) and 'sp'

(Pebesma and Bivand, 2005), respectively. We then interpolated the density of settlement points across

TNR at the same resolution as the DEM (30 m) using kernel density estimation (KDE) with the R

function *kernelUD* from package 'adehabitatHR', setting the smoothing parameter, h, to 1000 m

(Calenge, 2006). Finally, we rescaled this settlement proximity function by dividing by its maximum

value so it ranged between zero (far from settlements) and one (proximate to large settlements). This

index captured the relative size of settlements as well as distance to them. Although the smoothing

parameter (h) had a strong effect on KDE (Chen, 2017), its exact value did not have a major importance

because the predictor variables in the GAM were allowed to have non-linear effects. Our only

requirement for the KDE was that it captured some spatial variation in settlement density and gave more

weight to larger settlements (Wood, 2017).

Along fast-transects, we observed signs of artisanal gold mining, consisting of one to a few small

holes dug in the forest with an adjacent rudimentary camp that were either active or abandoned. To

interpolate mining density across TNR, we fit a binomial GAM to the mining points and 10,000 pseudo-

absence points using bivariate thin-plate splines to model the isotropic effects of latitude and longitude.

This model predicted significant spatial variation in mining. The smoothed latitude-longitude predictor

variable significantly influenced the probability of mining presence ($p = 0.02$). We converted these values

to relative mining density by dividing them by their mean, and then multiplying resulting values by the

observed mining density in transect strips (0.55 observed mining signs/km^2) (**Fig. 2D**). We attempted the

same approach to model the density of snares found along transects. However, latitude and longitude did

not significantly influence snare density ($p = 0.8$). As a consequence, we could not include snare density

as a predictor variable in our animal distribution models (Bryn et al., 2021).

2.3.4. Testing and accounting for spatial autocorrelation

Spatial autocorrelation is important to consider in species distribution modelling (Gaspard et al.,

2019). If left unaccounted for, it violates model assumptions of nonindependence and can cause spurious

association with covariates (Dormann et al., 2007). Our data collection protocol only considered

successive observations over 50 m apart, which likely reduced the risk of spatial autocorrelation. To

determine whether this was sufficient, we evaluated spatial autocorrelation using semi-variograms of

GAM residuals for points within five km of one another, with the R package 'gstat' (Pebesma, 2004). We

did not detect significant spatial autocorrelation for any of the animal species considered (**Fig. S1**). Thus,

we did not incorporate spatial autocorrelation in GAMs.

2.3.5. Model diagnostics

We used two strategies to assess the fit of GAMs. First, we mapped predicted relative density for

each species using the *predict* function in the 'raster' package, taking as predictor variables a raster stack

of the terrain and anthropogenic variables in our model (Hijmans et al., 2013). We plotted observations of

animal signs on top of this map to visually inspect model fit. Second, we assessed model fit by binning

the predicted probabilities from the GAM into 25 equal-sized groups. Bins had equal counts to avoid

differences in variance, so their widths varied. To evaluate alignment between predictions and

observations, we calculated the observed presence rate for each bin (i.e., the proportion of points with

detected animal sign) and standard error, and plotted these against the mean predicted probability.

310
311 **Fig. 1.** Situated in the eastern DRC, Tayna Nature Reserve (yellow, lower inset) lies in an ecological transition zone
312 between the Albertine rift mountains to the east and the Congo Basin rainforest to the west. Black vertical lines
313 correspond to the 196 line transects included in this survey, extracted from GPS tracks that recorded GPS points
314 every 50 m. Grey lines correspond to paths walked to reach transects (grey lines). The background forest cover map
315 is a composite satellite image from Google Earth (Maxar Technologies, Landsat/Copernicus 2020).
316

317
318 **Fig. 2**. Maps of predictor variables for wildlife species distribution models in Tayna Nature Reserve. Altitude and
319 TRI were calculated from a 30 m resolution digital elevation model (USGS, 2015; see Methods). Calculation of the
320 Settlement Index and Mine density are described in the Methods section.

3. Results

3.1. Fast-transect survey

The five survey teams completed 196 of the 200 planned transects. Four transects were inaccessible. Teams covered one or two transects per day and seven to twelve transects per two-week patrol over 22 total patrols, equating to a survey effort of 162 work days, excluding travel and rest days (76 actual days because of contemporaneous deployment). The average pace of fast-transects was 2.4 km/day (SD = 0.08 km/day across teams).

We recorded 87 signs of Grauer's gorilla and 151 signs of eastern chimpanzees both on and off transects in TNR. Signs included nest sites, tracks, trails, feeding remains, dung, vocalizations, and visual (for chimpanzees) observations. Among these signs, 41 gorilla signs and 119 chimpanzee signs were found on fast-transects (**Table 1**). There were 0.99 gorilla signs observed per km^2 of transect strip and 2.87 chimpanzee signs/km^2, or 0.10 gorilla signs per km of fast-transect walked and 0.30 chimpanzee signs/km. In addition to great apes, we recorded signs of cercopithecine monkeys, duikers, at least two pangolin species (*Smutsia gigantea* and *Phataginus tricuspis* and/or *P. tetradactyla*), red-river hog, brush-tailed porcupine (*Atherurus africanus*), forest buffalo (*Syncerus caffer nanus*), aardvark (*Orycteropus afer*), and leopard (*Panthera pardus*) (**Table 1**). There were no signs of forest elephants (*Loxodonta cyclotis*).

During the survey in TNR, both on and off transects, we found 292 gorilla nests in 40 nest sites (including 93 nests in 16 sites along transects) and 280 chimpanzee nests in 69 nest sites (including 209 nests in 55 sites along transects; **Fig. S2**). Excluding 23 solitary nests (15 attributed to chimpanzees and 8 to gorillas), the average nest group size was 8.89 (SD = 5.72) for gorillas and 4.91 (SD = 2.88) for chimpanzees (**Fig. S3**).

343 **Table 1.** Species identified along transects in Tayna Nature Reserve (TNR). Animal signs were recorded on 196
344 transects that covered 41.7 km^2 of the 899 km^2 of TNR.

Wildlife species	Seen	Heard	Tracks, trails, dung, and food remains	Sleeping sites, *i.e.*, nest sites or burrows, etc
Gorilla beringei graueri	0	1	24	16
Pan troglodytes schweinfurthii	1	18	45	55
Cercopithecus spp.	65	108	28	-
Cephalophus spp. & Philantomba monticola	11	0	380	18
Potamochoerus porcus	0	0	223	5
Atherurus africanus	1	1	196	7
Phataginus spp. & Smutsia gigantea	0	0	12	34
Syncerus caffer nanus	0	0	19	-
Panthera pardus	0	0	5	-
Orycteropus afer	0	0	1	5

3.2. Population distribution models

3.2.1. Grauer's Gorillas

The relative density of gorilla signs observed in TNR was positively associated with terrain

ruggedness (GAM, edf = 2.88, χ^2 = 14.50, p = 0.003), but not related to altitude (edf = 1.00, χ^2 = 1.84, p

= 0.18). It was negatively associated with the anthropogenic factors settlement index (edf = 1.00, χ^2 =

9.98, p = 0.002) and mine density (edf = 1.00, χ^2 = 12.37, p < 0.001; **Table 2**). Model fitted values

aligned visually with off-transect observations of gorilla signs (**Fig. 3a**). There was a clear positive

relationship between observations and model predictions, indicating that the model was able to capture

population distribution (**Fig. S4a**).

The model showed that gorilla sign relative density was maximized at ruggedness values above

7.5, which was above the average ruggedness of TNR (7.04, SD = 1.71, range = 3.04 – 12.90, median =

6.60), and plateaued thereafter (**Fig. 4a**). The settlement index was negatively associated with gorilla sign

density, with gorilla sign density declining to below average at values just above the median settlement

index in TNR (0.006, median = 0.004; **Fig. 4a**). The effect of mine density was negative, and no signs of

gorillas were found in mining hotspots (**Figs. 2 and 3**).

3.2.2. Eastern chimpanzees

The density of chimpanzee signs was positively associated with terrain ruggedness (GAM, edf =

1.92, χ^2 = 45.33, p < 0.001) but not altitude (edf = 1.00, χ^2 = 2.89, p = 0.09), and negatively associated

with the settlement index (edf = 1.00, χ^2 = 4.12, p = 0.04) and mining density (edf = 1.88, χ^2 = 5.94, p =

0.05; **Table 2**; **Fig. 3b**). Chimpanzee sign density was predicted to peak at maximal ruggedness (TRI >

12; **Fig. 4b**). Compared to gorilla signs, chimpanzee signs were more uniformly distributed across TNR,

including in the far north where great ape signs were never attributed to gorillas (**Fig. 4a**). The fitted

GAM for observed chimpanzee signs predicted well the distribution of observations (**Fig. S4b**).

Table 2. Generalized additive models of great ape sign density in TNR

Gorilla beringei graueri				
Term	*Edf*	**Ref df**	χ^2	*p*
Thin-plate smoothed terms				
Altitude	1.000	1.001	1.838	0.175
Terrain ruggedness index	2.879	3.285	14.504	**0.003**
Settlement index	1.000	1.000	9.977	**0.002**
Mining density	1.000	1.000	12.373	**<0.001**
n = 10041 *Deviance explained = 19.6%*				
Pan troglodytes schweinfurthii				
Term	*Edf*	**Ref df**	χ^2	*p*
Thin-plate smoothed terms				
Altitude	1.000	1.001	2.886	0.089
Terrain ruggedness index	1.918	2.371	45.327	**<0.001**
Settlement index	1.000	1.000	4.124	**0.042**
Mining density	1.883	2.298	5.936	**0.047**
n = 10119 *Deviance explained = 8.63%*				

372
373 **Fig. 3.** Predicted relative densities of animal populations in observed signs/km^2 in TNR based on GAM probability
374 of occurrences (see Methods). The average density of signs of gorillas, chimpanzees, monkeys, duikers, and red-
375 river hogs on transects were 0.99, 2.87, 4.15, 8.52, and 4.46 per km^2, respectively. Animal sign observations on
376 transects are marked in black circles, which were the presence points in the model. Off-transect sign observations
377 are shown in blue.

Fig. 4. Partial effects of altitude, terrain ruggedness, settlement proximity, and mine density on predicted animal sign density in observed signs per km² (see Methods). The dashed line shows the mean sign density for each population. Gray lines show the upper and lower 95% confidence intervals around the main effect prediction (black line).

3.2.3. Monkeys

The survey recorded sightings, vocalizations, prints, and food remains from guenons. From direct

observations only, we detected at least five different species: red-tailed monkey (*Cercopithecus ascanius*),

blue monkey (*C. mitis*), owl-faced monkey (*C. hamlyni*), mountain monkey (*C. lhoesti*), and Dent's mona

monkey (*C. denti*). Considering signs of all monkey species combined, they were less spatially variable

than those of great apes (**Fig. 3c**) and were not statistically associated with variation in altitude (GAM *edf*

$= 1.88$, $\chi^2 = 1.66$, $p = 0.58$), ruggedness (*edf* $= 1.00$, $\chi^2 = 0.76$, $p = 0.38$), nor mine density (*edf* $= 1.00$, χ^2

$= 0.52$, $p = 0.47$). Settlement proximity had a significant negative effect on the relative density of monkey

signs observed (*edf* $= 1.003$, $\chi^2 = 4.77$, $p = 0.03$; **Fig. 4c**). As expected from the relatively even sign

distribution, the predictive capability of the GAM for monkey signs was low (**Fig. S4c**).

3.2.4. Duikers

Signs and sightings of commonly-called "red duikers" (duikers of similar size with similar

footprints and dung, such as black-fronted duiker, *Cephalophus nigrifrons*, Weyn's duiker, *C. weynsi*, bay

duiker, *C. dorsalis*, and white-bellied duiker, *C. leucogaster*), yellow-backed duiker (*Cephalophus*

silvicultor), and blue duiker (*Philantomba monticola*) were observed widely across TNR (**Fig. 3d**).

Duiker signs consisted of prints, trails, food remains, and dung. The significant factors influencing the

distribution of duiker species' signs combined were settlement index (GAM *edf* $= 1.77$, $\chi^2 = 9.27$, p =

0.01) and altitude (*edf* $= 1.00$, $\chi^2 = 4.28$, p = 0.04). More duiker signs were recorded at low values of

both. Relative density appeared to decline at high mining density, but this effect was not significant (*edf* =

1.56, $\chi^2 = 4.53$, p = 0.07; **Fig. 4d**). There was a clear positive relationship between observations and

model predictions, indicating that the model was able to capture population distribution (**Fig. S4d**).

3.2.5. Red-river hogs

Red-river hogs left signs including trails, prints, dung, and food remains on transects. They were

heard and seen once (three individuals) off-transect. The settlement index had a strong negative effect on

the distribution of observed red river hog signs (GAM $edf = 1.000$, $\chi^2 = 20.58$, $p < 0.001$; **Figs. 3e and**

4e). Terrain also mattered, with sign density peaking just below the average ruggedness ($edf = 2.64$, $\chi^2 =$

8.95, $p = 0.04$; **Fig. 4e**). There was a clear positive relationship between observations and model

predictions, indicating sufficient predictive power of the model (**Fig. S4e**).

4. Discussion

We used a novel field data collection method, termed "fast-transects", to survey Tayna Nature Reserve (TNR) in north Kivu, DRC, in 2020. Our survey identified populations of Critically Endangered Grauer's gorillas and Endangered eastern chimpanzees. Observed signs of gorillas and chimpanzees were positively associated with increasing terrain ruggedness, but not altitude, and negatively influenced by proximity to human settlements. Gorilla signs also showed a negative association with artisanal mining in TNR. Signs of duikers, monkeys, and red river hogs were distributed more homogeneously throughout the reserve than either of the great apes. Nevertheless, all species groups were negatively impacted by proximity to human settlements. The survey also detected signs from other wildlife species, including forest buffalo, leopard, pangolins, and aardvark.

4.1. Comparison with prior surveys

This is the first survey to cover the entirety of TNR. We found that the Grauer's gorilla population in TNR is restricted to the rugged mountains and valleys in the center and south of the reserve. Gorillas were present in this sector in the early 2000s, based on partially-completed surveys (Mehlman, 2008; Nixon, 2013; Plumptre et al., 2015). The gorilla population seems to have persisted since then, at least in the central and southern sector. This may indicate that poaching of gorillas has been minimal since the legal establishment of TNR in 2006, although firm conclusions cannot be drawn in the absence of data from the 2000s from the vast northern sector of the reserve (Mehlman, 2008; Nixon, 2013; Plumptre et al., 2015). Like gorilla signs, chimpanzee signs were predominantly found in the center and south of TNR, but, contrastingly, chimpanzee signs were also found in the northern sector. Neither the distribution nor abundance of chimpanzees was reported in previous surveys (Mehlman, 2008; Nixon, 2013), but the encounter rate with nests appeared to have declined by over 80% from 2006 to the uncompleted 2013 survey (Nixon, 2013; Plumptre et al., 2015). Other species had not been formally surveyed previously, though the 2013 survey team in the center of TNR reported that no monkeys were seen or even heard (Nixon, 2013).

4.2. Influence on terrain on species distribution

Gorilla and chimpanzee signs in TNR were positively associated with ruggedness, but not altitude. Rugged terrain may provide foraging opportunities or refuge against predators and competitors, or the preference may reflect inaccessibility to humans. This finding may be a general characteristic of Grauer's gorillas and eastern chimpanzee in this region, as Plumptre et al. (2021) found a positive association between terrain steepness, a variable that likely correlates with terrain ruggedness, and great ape abundance in Kahuzi-Biega National Park and the Oku Community Reserve. In the tropics, plant diversity tends to be maximized in mountainous ridges and valleys (Baldeck et al., 2013). Great apes have diverse diets, comprising terrestrial herbaceous vegetation, seasonal fruit, roots, and more (*e.g.,* Basabose, 2002; Michel et al., 2022; van der Hoek et al., 2021b), and they may select areas with diverse plant communities where fallback foods and preferred foods co-occur within short distances. An alternative possibility is that great apes were extirpated from flatter areas due to hunting, or that they avoid such areas because of current or past human activities.

We did not detect an effect of terrain ruggedness or altitude on guenon relative density. If the negative association between flat terrain and great ape signs reflects hunting pressure, guenons may rebound quicker due to their larger population sizes and faster reproductive rates (Ingram et al., 2021), or their presence in risky habitats may reflect territorial exclusion or source-sink dynamics, where "source" rugged habitats are already at ecological carrying capacity (Novaro et al., 2000). It is also possible that terrain variability does not influence food resources for Cercopithecine monkeys or their perception of safety as they are more arboreal than great apes (Clark and Kaplin, 2024; Tuyisingize et al., 2023). Importantly, in our analysis we combined monkey species because of identification uncertainty for indirect signs, potentially smoothing over interspecific variation (Chapman et al., 2004). Duiker and red river hog signs were more common in low-altitude and less rugged areas, respectively. Little is known of the ecological preferences of either animal group in the region, so more research would be needed to

formulate and test hypotheses explaining this pattern (but see Bobo et al., 2025; van der Hoek et al.,

2023).

4.3. Influence of human presence & activities on species distribution

Human settlements and signs of artisanal mining negatively affected the relative density of

Grauer's gorillas and, to a slightly lower extent, eastern chimpanzees. Negative associations between

great apes and both small-scale mining and villages have been reported in the Grauer's gorilla range at

large (Plumptre et al., 2021), as well as for closely related great apes in west central Africa (Morgan et al.,

2018; Ordaz-Németh et al., 2021; Strindberg et al., 2018). Notably, chimpanzees were found in the

northern sector of TNR in areas where gorillas were not found. This may reflect their flexible multi-male

fission-fusion social system and smaller group sizes than gorillas, which could render them more tolerant

to hunting pressure. Other aspects of eastern chimpanzee behavior demonstrate their adaptability. For

example, chimpanzees living adjacent to croplands have been observed raiding crop fields at night (Krief

et al., 2014).

Relative densities of monkeys, duikers, and red river hogs were also lower near human

settlements. Regionally, duikers are the most commonly consumed bushmeat, with red river hogs second

(Batumike et al., 2021; Mbete et al., 2011; van Vliet et al., 2015). Artisanal mining did not appear to have

a significant effect on these taxa. This could reflect lower hunting pressure near mines than near more

permanent settlements.

Local community culture, including preferences, dietary taboos, traditional forest products, and

hunting tactics, result in prioritized offtake of different wildlife species and could drive the observed

pattern (Malimbo et al., 2020). While cultural transitions are likely eroding this source of variation

(Chausson et al., 2019; Mbete et al., 2011), traditional hunting practices are still important in this region.

The long-term absence of great apes from a significant part of the TNR forest may have

consequences on ecosystem functions (Abernethy et al., 2013). For example, in Nigeria, primate-

dispersed tree seedlings are rarer in areas where hunting is not regulated, compared to protected areas

491 (Effiom et al., 2013). Whether the structure of the TNR forest is affected by the heterogeneous

492 distribution of apes requires further research.

494 *4.4. Methodological limitations*

495  Fast-transects provided an efficient, rapid means to map the distribution of wildlife activity at the

496 population level. When the density of the type of animal sign used for a distance analysis is low, making

497 derived population densities too imprecise, fast-transects provide a practical solution. Relative to distance

498 transects, fast-transects enable more data to be collected in less time, reducing financial and logistical

499 costs to conservation organizations and reducing the risk that insecurity or unforeseen circumstances

500 prevent survey completion. If we had used the traditional method of standing crop nest counts on distance

501 transects, we would have only been able to analyze data from 16 gorilla nest sites and 55 chimpanzee nest

502 sites, at best. As Kühl et al (2008) recommend a minimum of 60-80 nest sites to calculate great ape

503 density using this method, it is clear that relying on nests only would have prevented us from testing most

504 of our hypotheses regarding great ape distribution. Instead, the fast-transect method allowed us to analyze

505 a total of 160 great ape signs. Combined with powerful non-linear models, the larger dataset allowed us to

506 detect significant effects of terrain and human activity on great ape distribution.

507  The main limitation of the fast-transect method is its inability to estimate absolute animal

508 densities and population sizes (Buckland et al., 2015). However, even distance transects based on

509 standing crop nest counts often fail at providing reliable absolute estimates, due to their reliance on

510 published nest decay rates from other geographical areas. A variation of the distance transect method

511 requiring marking observed nests and revisiting transects regularly reduced this bias, but this requires a

512 substantial data collection effort often incompatible with large geographical areas and impossible when

513 nest density is too low (Kouakou et al., 2009; Spehar et al., 2010). Because we did not attempt to estimate

514 absolute densities, we could analyze all sign types. Since each sign has particular biases (*e.g.,* dung;

515 Kamgaing et al., 2018; and vocalizations; Croes et al., 2007), combining multiple sign types helps build a

more accurate model of the distribution of wildlife rather than a particular behavior (*e.g.*, nesting or

alarm-calling).

4.5 Conservation perspectives & conclusions

The persistence of gorillas and chimpanzees in TNR underscores the importance of community

land protection in the eastern Albertine Rift biodiversity hotspot. Our survey results provide a much-

needed baseline for future surveys in TNR. Repeated fast-transects, currently underway, will track

changes in animal relative density patterns, letting managers cater development, sensitization, and

education programs to specific communities and identified threats. TNR now has field teams collecting

data on the movement patterns and diet of gorilla groups in the central rugged region, where this survey

identified multiple groups (Mbeke, personal communication). These continuous activities offer long-term

employment to community members living in closest proximity to wildlife (Mbeke, personal

communication).

As a peripheral population, TNR gorillas are more exposed to the risks of habitat fragmentation

and genetic isolation (van der Valk et al., 2018). Maintaining connectivity between TNR and other

wildlife populations is critical, beginning with the forested landscape that stretches westward toward

Maiko National Park. The status of great apes and other wildlife populations within this remote area, the

Usala Corridor, remains uncertain. Building on this study, a fast-transect survey is currently underway in

this region. Usala features heterogeneous terrain, yet it has a different pattern of anthropogenic pressures

compared to TNR. A comparative study could help disentangle whether the association between great

apes and rugged terrain observed in TNR is due to ecological factors, human avoidance, or historical

extirpation.

Globally, forest fragmentation is increasing at an alarming rate, causing biodiversity loss and

disrupting ecological processes (Taubert et al., 2018). The present survey is only the first step towards the

long-term protection of TNR. Moving forward, the continuation of monitoring activities and the

development of research will maintain the engagement of local communities and support long-term

conservation outcomes.

Acknowledgments

We are grateful to the many field assistants and trackers hired from the local villages surrounding Tayna Nature Reserve, without whom this study would not have been possible, including the following field team leaders: Arnold Kakule Vahwere, Affable Kakule Pilipili, Faustin Kambale Kinaba, Celestin Kasereka Kunemutumba, Moise Kalingini, Faustin Muhindo Kibwana, and Evariste Muhindo Katsongo. We thank the GRACE DRC team for managing ongoing survey logistics, in particular Augustin Kambere Mbangi. We thank Finley for his support. Two anonymous reviewers provided valuable suggestions. This work was funded by the Disney Conservation Fund's Reverse the Decline Initiative, the Margot Marsh Biodiversity Foundation, and the University of California, Davis.

CRediT authorship contribution statement

Alice Michel: Data curation, Formal analysis, Investigation, Methodology, Software, Visualization, Writing – original draft, Writing – review & editing. **Jackson Kabuyaya Mbeke:** Conceptualization, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing – review & editing. **Benezeth Kambale Visando:** Data curation, Investigation, Project administration, Resources, Supervision, Writing – review & editing. **Sonya Kahlenberg:** Conceptualization, Funding acquisition, Methodology, Project administration, Supervision, Writing – review & editing. **Katie Fawcett:** Funding acquisition, Project administration, Supervision, Writing – review & editing. **Damien Caillaud:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Validation, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data Availability

All data and code are available in the A.P.E.S. database (Kühl et al., 2007).

Appendix A: Supplementary Materials

Separate Word document containing supplementary figures.

References

Abernethy, K.A., Coad, L., Taylor, G., Lee, M.E., Maisels, F., 2013. Extent and ecological consequences

of hunting in Central African rainforests in the twenty-first century. Philos. Trans. R. Soc. B Biol.

Sci. 368. https://doi.org/10.1098/rstb.2012.0303

Aleman, J.C., Jarzyna, M.A., Staver, A.C., 2018. Forest extent and deforestation in tropical Africa since

1900. Nat. Ecol. Evol. 2, 26–33. https://doi.org/10.1038/s41559-017-0406-1

Baldeck, C.A., Harms, K.E., Yavitt, J.B., John, R., Turner, B.L., Valencia, R., Navarrete, H., Davies, S.J.,

Chuyong, G.B., Kenfack, D., Thomas, D.W., Madawala, S., Gunatilleke, N., Gunatilleke, S.,

Bunyavejchewin, S., Kiratiprayoon, S., Yaacob, A., Supardi, M.N.N., Dalling, J.W., 2013. Soil

resources and topography shape local tree community structure in tropical forests. Proc. R. Soc. B

Biol. Sci. 280, 20122532. https://doi.org/10.1098/rspb.2012.2532

Barbet-Massin, M., Jiguet, F., Albert, C.H., Thuiller, W., 2012. Selecting pseudo-absences for species

distribution models: How, where and how many? Methods Ecol. Evol. 3, 327–338.

https://doi.org/10.1111/j.2041-210X.2011.00172.x

Basabose, A.K., 2002. Diet composition of chimpanzees inhabiting the montane forest of Kahuzi,

Democratic Republic of Congo. Am. J. Primatol. 58, 1–21. https://doi.org/10.1002/ajp.10049

Batumike, R., Imani, G., Urom, C., Cuni-Sanchez, A., 2021. Bushmeat hunting around Lomami National

Park, Democratic Republic of the Congo. Oryx 55, 421–431.

https://doi.org/10.1017/S0030605319001017

Bessone, M., Booto, L., Santos, A.R., Kühl, H.S., Fruth, B., 2021. No time to rest: How the effects of

climate change on nest decay threaten the conservation of apes in the wild. PLOS ONE 16,

e0252527. https://doi.org/10.1371/journal.pone.0252527

Bivand, R.S., Pebesma, Edzer J, Gómez-Rubio, V., Pebesma, E., 2013. Applied Spatial Data Analysis

with R, 2nd ed. Springer, NY.

Blom, A., van Zalinge, R., Heitkönig, I.M.A., Prins, H.H.T., 2005. Factors influencing the distribution of

large mammals within a protected central African forest. Oryx 39, 381–388.

https://doi.org/10.1017/S0030605305001080

Bobo, D.N.B., Bessone, M., Cleva, R., Fruth, B., 2025. Spatial Ecology and Home Range Drivers of the

Weyns's Duiker (*Cephalophus weynsi*) in Rainforest Habitat. Afr. J. Ecol. 63, e70041.

Bryn, A., Bekkby, T., Rinde, E., Gundersen, H., Halvorsen, R., 2021. Reliability in Distribution

Modeling—A Synthesis and Step-by-Step Guidelines for Improved Practice. Front. Ecol. Evol. 9.

https://doi.org/10.3389/fevo.2021.658713

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L., Thomas, L., 2001.

Introduction to distance sampling: estimating abundance of biological populations. Oxford

University Press.

Buckland, S.T., Rexstad, E.A., Marques, T.A., Oedekoven, C.S., 2015. Distance Sampling: Methods and

Applications. Springer, NY.

Calenge, C., 2006. The package "adehabitat" for the R software: a tool for the analysis of space and

habitat use by animals. Ecol. Model. 197, 516–519.

Carvalho, J.S., Marques, T.A., Vicente, L., 2013. Population Status of Pan troglodytes verus in Lagoas de

Cufada Natural Park, Guinea-Bissau. PLOS ONE 8, e71527.

https://doi.org/10.1371/journal.pone.0071527

Chapman, C.A., Chapman, L.J., Cords, M., Gathua, J.M., Gautier-Hion, A., Lambert, J.E., Rode, K.,

Tutin, C.E.G., White, L.J.T., 2004. Variation in the Diets of Cercopithecus Species: Differences

within Forests, among Forests, and across Species, in: Glenn, M.E., Cords, M. (Eds.), The

Guenons: Diversity and Adaptation in African Monkeys. Springer US, Boston, MA, pp. 325–350.

https://doi.org/10.1007/0-306-48417-X_22

Chausson, A.M., Rowcliffe, J.M., Escouflaire, L., Wieland, M., Wright, J.H., 2019. Understanding the

sociocultural drivers of urban bushmeat consumption for behavior change interventions in Pointe

Noire, Republic of Congo. Hum. Ecol. 47, 179–191. https://doi.org/10.1007/s10745-019-0061-z

Chen, Y.C., 2017. A tutorial on kernel density estimation and recent advances. Biostat. Epidemiol. 1,

161–187. https://doi.org/10.1080/24709360.2017.1396742

Clark, A., Kaplin, B.A., 2024. Preliminary study of Dent's monkey (*Cercopithecus denti*) living in a

forest fragment in Rwanda, highlighting dietary flexibility in guenons. Afr. J. Ecol. 62, e13228.

Congo Basin Forest Partnership, 2006. Maiko-Tayna-Kahuzi-Biega Landscape, The Forests of the Congo

Basin: State of the Forest 2006. Johannesburg, South Africa.

Croes, B.M., Laurance, W.F., Lahm, S.A., Tchignoumba, L., Alonso, A., Lee, M.E., Campbell, P., Buij,

R., 2007. The Influence of Hunting on Antipredator Behavior in Central African Monkeys and

Duikers. Biotropica 39, 257–263. https://doi.org/10.1111/j.1744-7429.2006.00247.x

Cuni-Sanchez, A., Imani, G., Bulonvu, F., Batumike, R., Baruka, G., Burgess, N.D., Klein, J.A.,

Marchant, R., 2019. Social perceptions of forest ecosystem services in the Democratic Republic

of Congo. Hum. Ecol. 47, 839–853. https://doi.org/10.1007/s10745-019-00115-6

Dormann, C.F., McPherson, J.M., Araújo, M.B., Bivand, R., Bolliger, J., Carl, G., G. Davies, R., Hirzel,

A., Jetz, W., Daniel Kissling, W., Kühn, I., Ohlemüller, R., R. Peres-Neto, P., Reineking, B.,

Schröder, B., M. Schurr, F., Wilson, R., 2007. Methods to account for spatial autocorrelation in

the analysis of species distributional data: A review. Ecography 30, 609–628.

https://doi.org/10.1111/j.2007.0906-7590.05171.x

Edwards, D.P., Sloan, S., Weng, L., Dirks, P., Sayer, J., Laurance, W.F., 2014. Mining and the African

environment. Conserv. Lett. 7, 302–311. https://doi.org/10.1111/conl.12076

Effiom, E.O., Nuñez-Iturri, G., Smith, H.G., Ottosson, U., Olsson, O., 2013. Bushmeat hunting changes

regeneration of African rainforests. Proc. R. Soc. B Biol. Sci. 280, 20130246.

https://doi.org/10.1098/rspb.2013.0246

Estrada, A., Garber, P.A., Rylands, A.B., Roos, C., Fernandez-Duque, E., Fiore, A.D., Anne-Isola

Nekaris, K., Nijman, V., Heymann, E.W., Lambert, J.E., Rovero, F., Barelli, C., Setchell, J.M.,

Gillespie, T.R., Mittermeier, R.A., Arregoitia, L.V., de Guinea, M., Gouveia, S., Dobrovolski, R.,

Shanee, S., Shanee, N., Boyle, S.A., Fuentes, A., MacKinnon, K.C., Amato, K.R., Meyer, A.L.S.,

Wich, S., Sussman, R.W., Pan, R., Kone, I., Li, B., 2017. Impending extinction crisis of the

world's primates: Why primates matter. Sci. Adv. 3. https://doi.org/10.1126/sciadv.1600946

Gaspard, G., Kim, D., Chun, Y., 2019. Residual spatial autocorrelation in macroecological and

biogeographical modeling: a review. J. Ecol. Environ. 43, 19. https://doi.org/10.1186/s41610-

019-0118-3

Global Forest Watch, 2025. https://gfw.global/4n8GhNa

Google Earth V7.3.3.7786, 2021. Tayna, Democratic Republic of Congo. 2/10/2020. 0˚21'53"S

28˚46'02"E, Eye alt 38.69 mi.

Hall, J.S., Saltonstall, K., Inogwabini, B.I., Omari, I., 1998. Distribution, abundance and conservation

status of Grauer's gorilla. Oryx 32, 122–130. https://doi.org/10.1046/j.1365-3008.1998.d01-22.x

Hart, J.A., Hall, J.S., 1996. Status of eastern Zaire's forest parks and reserves. Conserv. Biol. 10, 316–

327.

Haurez, B., Petre, C.-A., Vermeulen, C., Tagg, N., Doucet, J.-L., 2014. Western lowland gorilla density

and nesting behavior in a Gabonese forest logged for 25 years: implications for gorilla

conservation. Biodivers. Conserv. 23, 2669–2687. https://doi.org/10.1007/s10531-014-0743-1

Hijmans, R.J., van Etten, J., Mattiuzzi, M., Sumner, M., Greenberg, J.A., Lamigueiro, O.P., Bevan, A.,

Racine, E.B., Shortridge, A., 2013. Raster package in R. [Computer software].

Hockings, K.J., McLennan, M.R., 2012. From Forest to Farm: Systematic Review of Cultivar Feeding by

Chimpanzees – Management Implications for Wildlife in Anthropogenic Landscapes. PLOS ONE 7,

e33391. https://doi.org/10.1371/journal.pone.0033391

Hogue, A.S., Breon, K., 2022. The greatest threats to species. Conserv. Sci. Pract. 4, e12670.

https://doi.org/10.1111/csp2.12670

Imani, G., Zapfack, L., Kalume, J., Riera, B., Cirimwami, L., Boyemba, F., 2016. Woody vegetation

groups and diversity along the altitudinal gradient in mountain forest: case study of Kahuzi-Biega

National Park and its surroundings, RD Congo. J. Biodivers. Environ. Sci. 8, 134–150.

Ingram, D.J., Coad, L., Milner-Gulland, E.J., Parry, L., Wilkie, D., Bakarr, M.I., Benítez-López, A.,

Bennett, E.L., Bodmer, R., Cowlishaw, G., Bizri, H.R.E., Eves, H.E., Fa, J.E., Golden, C.D.,

Iponga, D.M., Minh, N.V., Morcatty, T.Q., Mwinyihali, R., Nasi, R., Nijman, V., Ntiamoa-Baidu,

Y., Pattiselanno, F., Peres, C.A., Rao, M., Robinson, J.G., Rowcliffe, J.M., Stafford, C., Supuma,

M., Tarla, F.N., van Vliet, N., Wieland, M., Abernethy, K., 2021. Wild Meat Is Still on the Menu:

Progress in Wild Meat Research, Policy, and Practice from 2002 to 2020. Annu. Rev. Environ.

Resour. 46, 221–254. https://doi.org/10.1146/annurev-environ-041020-063132

Iyer, N., Kahlenberg, S.M., Williamson, E.A., Kyungu, J.-C., Syaluha, E.K., Mbeke, J.K., de Merode, E.,

Caillaud, D., 2023. Viability analysis for population reinforcement of Grauer's gorillas at Mount

Tshiaberimu, Democratic Republic of the Congo. J. Wildl. Manag. 87, e22384.

https://doi.org/10.1002/jwmg.22384

Kamgaing, T.O.W., Bobo, K.S., Djekda, D., Azobou, K.B.V., Hamadjida, B.R., Balangounde, M.Y.,

Simo, K.J., Yasuoka, H., 2018. Population density estimates of forest duikers (*Philantomba*

monticola & *Cephalophus* spp.) differ greatly between survey methods. Afr. J. Ecol. 56, 908–916.

https://doi.org/10.1111/aje.12518

Kouakou, C.Y., Boesch, C., Kühl, H., 2009. Estimating chimpanzee population size with nest counts:

validating methods in Taï National Park. Am. J. Primatol. 71, 447–457.

https://doi.org/10.1002/ajp.20673

Krief, S., Cibot, M., Bortolamiol, S., Seguya, A., Krief, J.-M., Masi, S., 2014. Wild Chimpanzees on the

Edge: Nocturnal Activities in Croplands. PLOS ONE 9, e109925.

https://doi.org/10.1371/journal.pone.0109925

Kühl, H., Maisels, F., Ancrenaz, M., Williamson, E.A., 2008. Best Practice Guidelines for Surveys and

Monitoring of Great Ape Populations, Occasional Paper of the IUCN Species Survival

Commission No. 36.

Kühl, H., Williamson, L., Sanz, C.M., Morgan, D., Boesch, C., 2007. Launch of A.P.E.S. Database.

Gorilla J. 34, 20–21.

Laing, S.E., Buckland, S.T., Burn, R.W., Lambie, D., Amphlett, A., 2003. Dung and nest surveys:

Estimating decay rates. J. Appl. Ecol. 40, 1102–1111. https://doi.org/10.1111/j.1365-

2664.2003.00861.x

Laurance, W.F., Croes, B.M., Tchignoumba, L., Lahm, S.A., Alonso, A., Lee, M.E., Campbell, P.,

Ondzeano, C., 2006. Impacts of Roads and Hunting on Central African Rainforest Mammals.

Conserv. Biol. 20, 1251–1261. https://doi.org/10.1111/j.1523-1739.2006.00420.x

Malimbo, D.K., Nyumu, J.K., Vitekere, K., Mapoli, M., Visando, B., Mbumba, J., Tungaluna, G.C.,

Tarla, F., Steve, N., 2020. Local perception on the exploitation, the current state and taboos

related to pangolins (Pholidota, Mammalia) by the communities living in the Tayna Nature

Reserve and Its surroundings (RNT) North Kivu DRC. J. Geosci. Environ. Prot. 08, 18.

https://doi.org/10.4236/gep.2020.812002

Martin, K., 1998. The role of animal behavior studies in wildlife science and management. Wildl. Soc.

Bull. 1973-2006 26, 911–920.

Mbete, R.A., Banga-Mboko, H., Racey, P., Mfoukou-Ntsakala, A., Nganga, I., Vermeulen, C., Doucet,

J.L., Hornick, J.L., Leroy, P., 2011. Household bushmeat consumption in Brazzaville, the

Republic of the Congo. Trop. Conserv. Sci. 4, 187–202.

https://doi.org/10.1177/194008291100400207

Mehlman, P.T., 2008. Current status of wild gorilla populations and strategies for their conservation, in:

Conservation in the 21st Century: Gorillas as a Case Study. Springer, NY, pp. 3–54.

https://doi.org/10.1007/978-0-387-70721-1_1

Michel, A., Minocher, R., Niehoff, P.-P., Li, Y., Nota, K., Gadhvi, M.A., Su, J., Iyer, N., Porter, A.,

Ngobobo-As-Ibungu, U., Binyinyi, E., Nishuli Pekeyake, R., Parducci, L., Caillaud, D.,

Guschanski, K., 2022. Isolated Grauer's gorilla populations differ in diet and gut microbiome.

Mol. Ecol. https://doi.org/10.1111/mec.16663

Mittermeier, R., Gil, P., Hoffmann, M., Pilgrim, J., Brooks, T., Mittermeier, C., Da Fonseca, G., 2004.

Hotspots revisited: Earth's biologically richest and most threatened ecoregions. CEMEX, Mexico

City, Mexico.

Morgan, D., Mundry, R., Sanz, C., Ayina, C.E., Strindberg, S., Lonsdorf, E., Kühl, H.S., 2018. African

apes coexisting with logging: Comparing chimpanzee (*Pan troglodytes troglodytes*) and gorilla

(*Gorilla gorilla gorilla*) resource needs and responses to forestry activities. Biol. Conserv. 218,

277–286. https://doi.org/10.1016/j.biocon.2017.10.026

Morgan, D., Sanz, C., Onononga, J.R., Strindberg, S., 2006. Ape Abundance and Habitat Use in the

Goualougo Triangle, Republic of Congo. Int. J. Primatol. 27, 147–179.

https://doi.org/10.1007/s10764-005-9013-0

Morgan, D., Sanz, C., Onononga, J.R., Strindberg, S., 2016. Factors Influencing the Survival of

Sympatric Gorilla (Gorilla gorilla gorilla) and Chimpanzee (Pan troglodytes troglodytes) Nests.

Int. J. Primatol. 37, 718–737. https://doi.org/10.1007/s10764-016-9934-9

Nixon, S., 2013. A Preliminary Assessment of Great Apes in the Tayna Nature Reserve, Eastern

Democratic Republic of Congo. Fauna & Flora International.

Novaro, A.J., Redford, K.H., Bodmer, R.E., 2000. Effect of Hunting in Source-Sink Systems in the

Neotropics. Conserv. Biol. 14, 713–721. https://doi.org/10.1046/j.1523-1739.2000.98452.x

Ordaz-Németh, I., Sop, T., Amarasekaran, B., Bachmann, M., Boesch, C., Brncic, T., Caillaud, D.,

Campbell, G., Carvalho, J., Chancellor, R., Davenport, T.R.B., Dowd, D., Eno-Nku, M., Ganas-

Swaray, J., Granier, N., Greengrass, E., Heinicke, S., Herbinger, I., Inkamba-Nkulu, C., Iyenguet,

F., Junker, J., Bobo, K.S., Lushimba, A., Maisels, F., Malanda, G.A.F., McCarthy, M.S.,

Motsaba, P., Moustgaard, J., Murai, M., Ndokoue, B., Nixon, S., Nseme, R.A., Nzooh, Z., Pintea,

L., Plumptre, A.J., Roy, J., Rundus, A., Sanderson, J., Serckx, A., Strindberg, S., Tweh, C.,

Vanleeuwe, H., Vosper, A., Waltert, M., Williamson, E.A., Wilson, M., Mundry, R., Kühl, H.S.,

2021. Range-wide indicators of African great ape density distribution. Am. J. Primatol.

https://doi.org/10.1002/ajp.23338

Pebesma, E., Bivand, R.S., 2005. S classes and methods for spatial data: the sp package. [Computer

software].

Pebesma, E.J., 2004. Multivariable geostatistics in S: the gstat package. Comput. Geosci. 30, 683–691.

Plumptre, A.J., Ayebare, S., Segan, D., Watson, J., Kujirakwinja, D., 2017. Conservation Action Plan for

the Albertine Rift. [Unpublished Report for Wildlife Conservation Society and its Partners].

https://doi.org/10.13140/RG.2.2.15701.32485

Plumptre, A.J., Kirkby, A., Spira, C., Kivono, J., Mitamba, G., Ngoy, E., Nishuli, R., Strindberg, S.,

Maisels, F., Buckland, S., Ormsby, L., Kujirakwinja, D., 2021. Changes in Grauer's gorilla

(*Gorilla beringei graueri*) and other primate populations in the Kahuzi-Biega National Park and

Oku Community Reserve, the heart of Grauer's gorilla global range. Am. J. Primatol. 83.

https://doi.org/10.1002/ajp.23288

Plumptre, A.J., Nixon, S., Critchlow, R., Vieilledent, G., Nishuli, R., Kirkby, A., Williamson, E.A., Hall,

J.S., Kujirakwinja, D., 2015. Status of Grauer's Gorilla and Chimpanzees in Eastern Democratic

Republic of Congo: Historical and Current Distribution and Abundance. Wildlife Conservation

Society, Fauna & Flora International and Institut Congolais pour la Conservation de la Nature.

https://doi.org/10.19121/2015.report.dmx3094100000

Plumptre, A.J., Nixon, S., Kujirakwinja, D.K., Vieilledent, G., Critchlow, R., Williamson, E.A., Nishuli,

R., Kirkby, A.E., Hall, J.S., 2016. Catastrophic decline of world's largest primate: 80% loss of

Grauer's gorilla (*Gorilla beringei graueri*) population justifies critically endangered status. PLOS

ONE 11. https://doi.org/10.1371/journal.pone.0162697

Poisson, S.-D., 1837. Recherches sur la probabilité des jugements en matière criminelle et en matière

civile: précédées des règles générales du calcul des probabilités. Bachelier.

Poulsen, J.R., Rosin, C., Meier, A., Mills, E., Nuñez, C.L., Koerner, S.E., Blanchard, E., Callejas, J.,

Moore, S., Sowers, M., 2018. Ecological consequences of forest elephant declines for

Afrotropical forests. Conserv. Biol. 32, 559–567. https://doi.org/10.1111/cobi.13035

R Core Team, 2023. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria [Computer software]. https://www.R-project.org/

Ramiadantsoa, T., Hanski, I., Ovaskainen, O., 2018. Responses of generalist and specialist species to

fragmented landscapes. Theor. Pop. Bio. 124, 31–40. https://doi.org/10.1016/j.tpb.2018.08.001

Ripple, W.J., Abernethy, K., Betts, M.G., Chapron, G., Dirzo, R., Galetti, M., Levi, T., Lindsey, P.A.,

Macdonald, D.W., Machovina, B., Newsome, T.M., Peres, C.A., Wallach, A.D., Wolf, C., 2016.

Bushmeat hunting and extinction risk to the world's mammals. R. Soc. Open Sci. 3.

https://doi.org/10.1098/rsos.160498

Robinson, H.S., Wielgus, R.B., Cooley, H.S., Cooley, S.W., 2008. Sink populations in carnivore

management: cougar demography and immigration in a hunted population. Ecol. Appl. 18, 1028–

1037. https://doi.org/10.1890/07-0352.1

Romani, T., Mundry, R., Shaban, G.M., Konarzewski, M., Namaganda, M., Hobaiter, C., Gruber, T.,

Hicks, T.C., 2023. Decay rates of arboreal and terrestrial nests of Eastern chimpanzees (Pan

troglodytes schweinfurthii) in the Bugoma Central Forest Reserve, Uganda: Implications for

population size estimates. Am. J. Primatol. 85, e23536. https://doi.org/10.1002/ajp.23536

Ross, N., 2020. fasterize: Fast polygon to raster conversion. R Package Version 1. [Computer software].

Sexton, J.P., McIntyre, P.J., Angert, A.L., Rice, K.J., 2009. Evolution and Ecology of Species Range

Limits. Annu. Rev. Ecol. Evol. Syst. 40, 415–436.

https://doi.org/10.1146/annurev.ecolsys.110308.120317

Spehar, S.N., Mathewson, P.D., Wich, S.A., Marshall, A.J., Kühl, H., Meijaard, E., 2010. Estimating

Orangutan Densities Using the Standing Crop and Marked Nest Count Methods: Lessons Learned

for Conservation: Assessment of Orangutan Nest Count Methods. Biotropica 42, 748–757.

https://doi.org/10.1111/j.1744-7429.2010.00651.x

Spira, C., Kirkby, A., Kujirakwinja, D., Plumptre, A.J., 2019. The socio-economics of artisanal mining

and bushmeat hunting around protected areas: Kahuzi-Biega National Park and Itombwe Nature

Reserve, eastern Democratic Republic of Congo. Oryx 53, 136–144.

https://doi.org/10.1017/S003060531600171X

Spitz, D.B., Rowland, M.M., Clark, D.A., Wisdom, M.J., Smith, J.B., Brown, C.L., Levi, T., 2019.

Behavioral changes and nutritional consequences to elk (*Cervus canadensis*) avoiding perceived

risk from human hunters. Ecosphere 10, e02864. https://doi.org/10.1002/ecs2.2864

Stillfried, M., Belant, J.L., Svoboda, N.J., Beyer, D.E., Kramer-Schadt, S., 2015. When top predators

become prey: Black bears alter movement behaviour in response to hunting pressure. Behav.

Processes 120, 30–39. https://doi.org/10.1016/j.beproc.2015.08.003

Stokes, E.J., Strindberg, S., Bakabana, P.C., Elkan, P.W., Iyenguet, F.C., Madzoké, B., Malanda, G.A.F.,

Mowawa, B.S., Moukoumbou, C., Ouakabadio, F.K., Rainey, H.J., 2010. Monitoring great Ape

and elephant abundance at large spatial scales: Measuring effectiveness of a conservation

landscape. PLOS ONE 5. https://doi.org/10.1371/journal.pone.0010294

Strindberg, S., Maisels, F., Williamson, E.A., Blake, S., Stokes, E.J., Aba'a, R., Abitsi, G., Agbor, A.,

Ambahe, R.D., Bakabana, P.C., Bechem, M., Berlemont, A., De Semboli, B.B., Boundja, P.R.,

Bout, N., Breuer, T., Campbell, G., De Wachter, P., Akou, M.E., Mba, F.E., Feistner, A.T.C.,

Fosso, B., Fotso, R., Greer, D., Inkamba-Nkulu, C., Iyenguet, C.F., Jeffery, K.J., Kokangoye, M.,

Kühl, H.S., Latour, S., Madzoke, B., Makoumbou, C., Malanda, G.A.F., Malonga, R., Mbolo, V.,

Morgan, D.B., Motsaba, P., Moukala, G., Mowawa, B.S., Murai, M., Ndzai, C., Nishihara, T.,

Nzooh, Z., Pintea, L., Pokempner, A., Rainey, H.J., Rayden, T., Ruffler, H., Sanz, C.M., Todd,

A., Vanleeuwe, H., Vosper, A., Warren, Y., Wilkie, D.S., 2018. Guns, germs, and trees determine

density and distribution of gorillas and chimpanzees in Western Equatorial Africa. Sci. Adv. 4.

https://doi.org/10.1126/sciadv.aar2964

Taubert, F., Fischer, R., Groeneveld, J., Lehmann, S., Müller, M.S., Rödig, E., Wiegand, T., Huth, A.,

2018. Global patterns of tropical forest fragmentation. Nature 554, 519–522.

https://doi.org/10.1038/nature25508

Tutin, C.E.G., Parnell, R.J., White, L.J.T., Fernandez, M., 1995. Nest building by lowland gorillas in the

Lopé Reserve, Gabon: Environmental influences and implications for censusing. Int. J. Primatol.

16, 53–76. https://doi.org/10.1007/BF02700153

Tuyisingize, D., Kaplin, B.A., Eckardt, W., Musana, A., Caillaud, D., 2023. Distribution and conservation

status of the golden monkey *Cercopithecus mitis kandti* in Rwanda. Oryx 57, 98–106.

https://doi.org/10.1017/S0030605321001009

USGS, 2015. Shuttle radar topography mission (SRTM) 1 Arc-Second global. US Geol. Surv.

Van Cleave, E.K., Bidner, L.R., Ford, A.T., Caillaud, D., Wilmers, C.C., Isbell, L.A., 2018. Diel patterns

of movement activity and habitat use by leopards (*Panthera pardus pardus*) living in a human-

dominated landscape in central Kenya. Biol. Conserv. 226, 224–237.

https://doi.org/10.1016/j.biocon.2018.08.003

van der Hoek, Y., Binyinyi, E., Ngobobo, U., Stoinski, T.S., Caillaud, D., 2023. Diversity and diel

activity patterns of terrestrial mammals in the Nkuba Conservation Area, Democratic Republic of

the Congo. Oryx 57, 107–117. https://doi.org/10.1017/S003060532100106X

van der Hoek, Y., Binyinyi, E., Ngobobo, U., Stoinski, T.S., Caillaud, D., 2021a. Daily travel distances of

unhabituated Grauer's gorillas (*Gorilla beringei graueri*) in a low elevation forest. Folia Primatol.

92, 112–125. https://doi.org/10.1159/000514626

van der Hoek, Y., Pazo, W.D., Binyinyi, E., Ngobobo, U., Stoinski, T.S., Caillaud, D., 2021b. Diet of

Grauer's gorillas (*Gorilla beringei graueri*) in a low-elevation forest. Folia Primatol. 92, 126–

138. https://doi.org/10.1159/000515377

van der Hoek, Y., Van de Perre, F., Fastré, C., Kirkpatrick, L., Ngobobo, U., Tokunda, R., Binyinyi, E.,

Stoinski, T.S., 2024. Short-term trends in great ape density in a community-based conservation

area in the eastern Democratic Republic of the Congo. Ecol. Solut. Evid. 5, e12379.

https://doi.org/10.1002/2688-8319.12379

van der Meer, E., Fritz, H., Blinston, P., Rasmussen, G.S.A., 2014. Ecological trap in the buffer zone of a

protected area: effects of indirect anthropogenic mortality on the African wild dog Lycaon pictus.

Oryx 48, 285–293. https://doi.org/10.1017/S0030605312001366

van der Valk, T., Sandoval-Castellanos, E., Caillaud, D., Ngobobo, U., Binyinyi, E., Nishuli, R., Stoinski,

T., Gilissen, E., Sonet, G., Semal, P., Kalthoff, D.C., Dalén, L., Guschanski, K., 2018. Significant

loss of mitochondrial diversity within the last century due to extinction of peripheral populations

in eastern gorillas. Sci. Rep. 8, 1–10. https://doi.org/10.1038/s41598-018-24497-7

Van der Weyde, L.K., Mbisana, C., Klein, R., 2018. Multi-species occupancy modelling of a carnivore

guild in wildlife management areas in the Kalahari. Biol. Conserv. 220, 21–28.

https://doi.org/10.1016/j.biocon.2018.01.033

van Vliet, N., Nasi, R., 2008. Mammal distribution in a Central African logging concession area.

Biodivers. Conserv. 17, 1241–1249. https://doi.org/10.1007/s10531-007-9300-5

van Vliet, N., Nebesse, C., Nasi, R., 2015. Bushmeat consumption among rural and urban children from

Province Orientale, Democratic Republic of Congo. Oryx 49, 165–174.

https://doi.org/10.1017/S0030605313000549

Walsh, P.D., White, L.J.T., 2005. Evaluating the steady state assumption: Simulations of gorilla nest

decay. Ecol. Appl. 15, 1342–1350. https://doi.org/10.1890/03-5283

Wilson, M.F.J., O'Connell, B., Brown, C., Guinan, J.C., Grehan, A.J., 2007. Multiscale terrain analysis of

multibeam bathymetry data for habitat mapping on the continental slope. Mar. Geod. 30, 3–35.

Wood, S.N., 2018. Mixed GAM computation vehicle with GCV/AIC/REML smoothness estimation and

GAMMs by REML/PQL. R Package Version 1–8.

Wood, S.N., 2017. Generalized Additive Models: An Introduction with R, Second Edition. CRC Press.

Ziegler, S., Fa, J.E., Wohlfart, C., Streit, B., Jacob, S., Wegmann, M., 2016. Mapping bushmeat hunting

pressure in Central Africa. Biotropica 48, 405–412. https://doi.org/10.1111/btp.12286