Antiparallel Cell Circulation Emerging from Self-Aligned Tension Gradients

Ryunosuke Karimata[1], Hidenori Hashimura[2], Shuhei A. Horiguchi[3], Taihei Fujimori[2,4], Satoshi Sawai[2,5], Satoru Okuda[3, 6*]

[1] Division of Nano Life Science, Graduate School of Frontier Science Initiative, Kanazawa University, Kanazawa 920-1192, Japan

[2] Graduate School of Arts and Sciences, The University of Tokyo, Tokyo 153-8902, Japan

[3] Nano Life Science Institute, Kanazawa University, Kanazawa 920-1192, Japan

[4] Chan Zuckerberg Biohub, San Francisco, CA 94158, USA

[5] Research Center for Complex Systems Biology, Universal Biology Institute, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo 153-8902, Japan

[6] Sapiens Life Sciences, Evolution and Medicine Research Center, Kanazawa University, Kanazawa 920-8640, Japan

* Corresponding author: satokuda@staff.kanazawa-u.ac.jp

Abstract

Active tissues exhibit diverse collective dynamics, yet the cell–cell interactions that generate ordered microscopic flows remain poorly understood. Here, we show that antiparallel cell circulation can emerge from self-aligned, polarity-dependent tension gradients. Using a minimal vertex model of confluent tissues, we studied polar cells that align their polarity with their own velocity and impose polarity-dependent tension gradients along cell–cell contacts, without relying on substrate traction. This behavior can be generalized as a minimal interaction in which forces transmitted between cells act with opposite signs, reminiscent of action–reaction forces, organizing cells into stable interlocking antiparallel lanes. In mixtures of motile and nonmotile cells, this circulation drives phase separation, in which motile cells spontaneously form persistent domains. Accordingly, we identified similar antiparallel circulation patterns in two-dimensional aggregates of *Dictyostelium discoideum*, supporting the biological relevance of the mechanism. Together, these results demonstrate that self-aligned tension gradients provide a robust and underappreciated route to dynamic microscopic pattern formation in multicellular systems.

I. INTRODUCTION

Ordered patterns are a hallmark of active matter, appearing in diverse systems such as self-propelled colloids [1–3], active droplets [4,5], bacterial suspensions [6–9], and living cell assemblies [10–12]. Such systems can exhibit diverse spatiotemporal order [7,8,13,14], demonstrating how internally generated active forces can coordinate matter far from equilibrium. A natural mechanistic question is how these forces are exchanged and transmitted between constituents. This issue is especially relevant in densely packed biological assemblies, where the dominant mechanical interactions often arise not from interactions with the surrounding environment, but from force exchange and transmission via direct physical contacts between constituents. A simplified setting that highlights such contact-mediated force transmission is provided by motor-filament assays such as actin–myosin and microtubule–kinesin systems: cytoskeletal filaments linked by motor proteins slide past one another [15]. Similar contact-mediated force exchange also occurs at cell–cell junctions in confluent tissues. Yet how inter-constituent force exchange shapes spatiotemporal organization at the single-cell scale in confluent tissues remains poorly understood.

Epithelial tissues are a good example of cell-based active systems, exhibiting a variety of dynamic patterns, such as flocking [16–19] and vortex formation [19–23]. Often, cells migrate on frictional substrates or basement membranes, generating motion via multiple types of active driving forces. The emergence of patterns driven by self-propulsion at the cell–substrate interface has been widely investigated in both theoretical and experimental studies [22,24–28]. A distinct, less explored possibility arises from active interfacial tensions at cell–cell contacts, such as epithelial junctions [29–31]. Because interfacial tensions exert equal and opposite forces on the two neighboring cells, they obey action–reaction symmetry and provide a natural route to internally coordinated motion without requiring net external forces. Importantly, these tensions can become directionally biased by intracellular polarity, producing polarity-aligned interfacial tension gradients that drive directional cell migration and complex collective dynamics in confluent tissues [29–31].

Polarity regulation plays a central role in determining how mechanical interactions are translated into collective cell behavior. Two primary mechanisms have been considered: mutual alignment, in which neighboring cells directly coordinate their polarities [17,18,32,33], and self-alignment, where polarity aligns with each cell's own movement [24,34–37]. Mutual alignment can directly prescribe the local arrangement of cell polarities at the single-cell level, whereas in self-alignment, polarity is generated intrinsically and does not by itself specify how neighboring cells arrange or orient [35,37,38]. In systems dominated by self-alignment, cell-scale order is therefore likely to emerge from additional intercellular mechanical interactions. Interfacial tension gradients offer a natural mechanism for such coupling: by biasing active tensions along cell–cell contacts, they

generate directional yet force-balanced interactions that feed back onto cell rearrangements and motion. Whether these self-aligned tension gradients alone can generate robust, cell-level dynamic patterns remains an open question.

Here, we explore single-cell-scale dynamic patterns emerging from collective motion driven by polarized, force-balanced interactions. We focus on a minimal self-alignment framework, where the polarity of each cell aligns with its own movement, implemented within a two-dimensional (2D) vertex model of confluent tissues. In this setting, such force-balanced interactions arise from polarity-dependent interfacial tensions and simplified attractive–repulsive forces along cell–cell contacts, which drive collective motion through internal rearrangements. This minimal mechanism produces a unique, previously unreported antiparallel circulation pattern at the level of constituent cells. Guided by this prediction, we observe a similar circulation pattern in aggregating *Dictyostelium discoideum*, and a simple action–reaction argument explains how neighboring circulations adopt opposite orientations. Together, these results show that polarized force-balanced interactions provide a robust and underappreciated route for generating dynamic microscopic patterns in multicellular systems.

II. MODEL

We now consider a confluent tissue composed of N_c cells in a 2D periodic box: $0 < x < L_x$ and $0 < y < L_y$ [Fig. 1(a)]. The tissue consists of N_m motile-type cells, indexed by $\mathcal{M} \coloneqq \{1, 2, \cdots, N_m\}$ and $N_n \coloneqq N_c - N_m$, and nonmotile-type cells, indexed by $\mathcal{N} \coloneqq \{N_m + 1, \cdots, N_c\}$, with a mixing ratio $\phi \coloneqq N_m / N_c$. We describe multicellular dynamics using a 2D vertex model [39–42], specified by the vertex positions $\{r_i\}$, and the polarity orientations of motile cells $\{\theta_\alpha\}_{\alpha \in \mathcal{M}}$, where $q_\alpha = (\cos \theta_\alpha, \sin \theta_\alpha)$.

We first assume self-alignment, meaning that the polarity of each motile-type cell aligns with the direction of its own velocity, consistent with previous theoretical and experimental studies [24,26,34–37,43–46]. Additionally, accounting for rotational noise, the time evolution of the orientation of the polarity, denoted by $\{\theta_\alpha\}_{\alpha \in \mathcal{M}}$, is given by

$$\frac{d\theta_\alpha}{dt} = J \sin(\psi_\alpha - \theta_\alpha) + w_\alpha, \tag{1}$$

where J is the alignment rate and ψ_α is the orientation of the velocity of the α-th cell. Additionally, w_α represents white Gaussian noise, satisfying $\langle w_\alpha(t) \rangle = 0$ and $\langle w_\alpha(t) w_{\alpha'}(t') \rangle = 2D_r \delta_{\alpha\alpha'} \delta(t - t')$, where D_r is the noise intensity. From Eq. (1), the characteristic time scale on which polarity aligns with velocity is determined by J and D_r.

96   We next describe the multicellular dynamics driven by self-aligned tension gradients. A cell

97   monolayer is represented as a polygonal tiling, where each polygon corresponds to a cell, and

98   vertices and edges are shared among adjacent cells [Fig. 1(a)]. Cell rearrangements occur through T1

99   transitions when the length of an edge becomes smaller than the threshold length Δl [Fig. 1(b)].

100   The positions of the vertices $\{r_i\}$ evolve under overdamped dynamics,

$$\eta \frac{\mathrm{d}r_i}{\mathrm{d}t} = -\frac{\partial U}{\partial r_i}, \tag{2}$$

101   where η is the substrate friction coefficient. The left-hand side represents the frictional drag on

102   vertex i, and the right-hand side gives the mechanical force derived from the effective energy, U. U

103   is given by a standard vertex-model functional widely employed in previous studies [40,41,47]:

$$U = \sum_{\alpha=1}^{N_c} \left[\frac{K_A}{2}(A_\alpha - A_0)^2 + \frac{K_P}{2}(P_\alpha - P_0)^2 + \sum_{\varepsilon \in \mathcal{E}_\alpha} \gamma_{\alpha\varepsilon} l_\varepsilon \right]. \tag{3}$$

The first term represents the area elastic energy, where K_A is the area elasticity modulus, A_α is the

area of the α-th cell, and A_0 is the preferred cell area. The second term represents the perimeter

elastic energy, with K_P the perimeter-elastic modulus, P_α the perimeter of the α-th cell, and P_0

the preferred perimeter. The third term accounts for interfacial energy, where $\gamma_{\alpha\varepsilon}$ is the tension on

the ε-th edge of the α-th cell, l_ε is the length of the ε-th edge, and $\mathcal{E}_\alpha$ is the set of edges forming

the α-th cell.

Our model assumes that each motile cell generates a driving force by establishing a polarity-

aligned interfacial tension gradient along its edges, consistent with previous studies of polarity-

dependent interfacial tensions in epithelial systems [29–31,48,49]. The tension, $\gamma_{\alpha\varepsilon}$ for $\varepsilon \in \mathcal{E}_\alpha$, is

defined as

$$\gamma_{\alpha\varepsilon} = \begin{cases} -\gamma^p \cos \varphi_{\alpha\varepsilon} & (\alpha \in \mathcal{M}) \\ 0 & (\alpha \in \mathcal{N}) \end{cases}, \tag{4}$$

where γ^p represents the strength of polarized tension, and $\varphi_{\alpha\varepsilon}$ denotes the angle between the

polarity vector $q_\alpha = (\cos \theta_\alpha, \sin \theta_\alpha)$ and the vector from the center of the α-th cell to the midpoint

of the ε-th edge [Fig. 1(d)]. Through Eq. (4), each motile cell generates a gradient of interfacial

tension—higher at the trailing side ($\cos \varphi_{\alpha\varepsilon} < 0$) and lower at the leading side ($\cos \varphi_{\alpha\varepsilon} > 0$), thereby

producing a polarity-directed active force within the tissue. Although $\gamma_{\alpha\varepsilon}$ depends implicitly on $\{r_i\}$

and $\{\theta_\alpha\}$ through $\varphi_{\alpha\varepsilon}$, we assume that tensions are set autonomously by cellular activity rather than

arising from mechanical equilibration. Thus, when calculating forces from Eq. (2), we treat $\gamma_{\alpha\varepsilon}$ as

constant with respect to vertex positions (i.e., $\partial \gamma_{\alpha\varepsilon}/\partial r_i = 0$). This assumption ensures that the

tension gradient acts as an active force, a detailed-balance-breaking force while still satisfying force

balance.

All parameters are nondimensionalized by the unit length $\sqrt{A_0}$, unit time $\eta/(K_A A_0)$, and unit

energy $K_A A_0{}^2$. The parameters, $K_p = 1$, $\gamma^p = 0.20$, $N_c = 32 \times 32$, and $\phi = 0.50$, are fixed in all

simulations, unless otherwise specified, while J and D_r are varied. To ensure that varying the

values of J and D_r does not affect the tissue fluidity, the nondimensionalized preferred perimeter

$p_0 := P_0/\sqrt{A_0}$, known as the target shape index, is fixed at $p_0 = 3.90$, meaning that the system

remains in a fluid-like state [50] (see Appendix A). Equations (1) and (2) are solved numerically

using the Euler method from $t = 0$ to $t = 10^4$ with a time step of $\Delta t = 0.01$. The threshold

length of T1 transitions is set to $\Delta l = 0.05$.

III. RESULTS

A. Self-aligned tension gradients induce an antiparallel circulation pattern

By varying the alignment rate, J, and noise intensity, D_r, we first found that the cells underwent

phase separation when $J > D_r$: motile cells spontaneously cohered and formed domains surrounded

by nonmotile cells [Fig. 2(a) top row and Movie S1]. To quantify the phase separation, we introduced

the separation degree, μ, as an order parameter,

$$\mu := 1 - \frac{1}{\phi}\left(\frac{1}{N_n}\sum_{\alpha \in \mathcal{N}} \frac{n_\alpha^m}{n_\alpha^m + n_\alpha^n}\right), \tag{5}$$

where n_α^m and n_α^n are the numbers of motile-type and nonmotile-type cells adjoining the α-th cell,

respectively; $\mu \approx 1$ indicates phase separation, while $\mu \approx 0$ indicates a homogeneous system.

During phase separation, μ increases and eventually reaches a plateau value, whereas it

remains near zero when no separation occurs [Fig. 2(b)]. The plateau value approaches unity for

sufficiently large J and small D_r values [Fig. 2(c)].

When $J < D_r$, phase separation does not occur; even when a pre-separated structure is used as

the initial condition, the system reverts to a disordered configuration (see Appendix B). Separation

persists even at $D_r = 0$ as long as $J \neq 0$, whereas no separation is observed when $J = 0$ and $D_r \neq$

0. These results demonstrate that self-alignment is essential for phase separation [Fig. 2(c)].

Moreover, for small D_r, μ exhibits a steep transition with respect to J, revealing a critical point,

whereas the transition becomes more gradual as D_r increases [Fig. 2(d)]. These findings indicate

that phase separation arises from polarity alignment coupled to active forces and is therefore

qualitatively distinct from thermally driven phase separation.

We next examined the cell movement and polarity dynamics within the domains formed under
phase-separation conditions. We found that motile cells self-organize into a vortex-like pattern [Fig.
2(a) bottom row and Movie S2] (see Appendix C), characterized by concentric alignment of both
velocity and polarity. Cells sharing the same velocity and polarity direction assemble into chain-like
arrangements, which we refer to as circulations. These circulations are organized into concentric,
onion-like layers, with adjacent circulations aligned and moving in antiparallel directions [Fig. 2(e)].

To quantify the emergence of antiparallel circulations, we performed simulations starting from a
circular domain composed of motile cells with random initial polarity directions. Under a phase-
separation condition with $J = 5.0$ and $D_{\mathrm{r}} = 0.01$, we calculated the statistics of cell movement
directions and speeds [Fig. 2(f) and 2(g)]. The results showed that cells predominantly move either
clockwise or counterclockwise, exhibiting maximal velocity in the tangential directions,
demonstrating that cells migrate in a concentric antiparallel manner. We further evaluated the
persistence time of cell migration along the circulation, denoted τ_{α} (see Appendix C), for $D_{\mathrm{r}} =$
0.01 and $D_{\mathrm{r}} = 1.00$. This analysis revealed that cells migrate more persistently at lower noise [Fig.
2(h)], indicating that circulation persists for significant durations under low-noise conditions.

Furthermore, to quantify the emergence of antiparallel alignment between adjacent circulations,
we introduce a pairwise order parameter S, defined as

$$S := \frac{1}{N_{\mathrm{m}}} \sum_{\alpha \in \mathcal{M}} \left[\frac{1}{n_{\alpha}^{\mathrm{m}}} \sum_{\alpha' \in \mathcal{M}_{\alpha}} \chi_{\alpha\alpha'} \mathrm{sgn}(\boldsymbol{q}_{\alpha} \cdot \boldsymbol{q}_{\alpha'}) \right], \tag{6}$$

169 where, $\mathcal{M}_{\alpha}$ denotes the set of motile-type cells adjoining the α-th motile-type cell. The geometric
170 factor $\chi_{\alpha\alpha'}$ is given by

$$\chi_{\alpha\alpha'} := \begin{cases} -1 & -\frac{\sqrt{3}}{2} < \frac{(\boldsymbol{r}_{\alpha'} - \boldsymbol{r}_{\alpha}) \cdot \boldsymbol{q}_{\alpha}}{\|\boldsymbol{r}_{\alpha'} - \boldsymbol{r}_{\alpha}\|} < \frac{\sqrt{3}}{2}, \\ 1 & \text{otherwise} \end{cases} \tag{7}$$

where $\boldsymbol{r}_{\alpha}$ is the centroid position of the α-th motile-type cell. Thus, $\chi_{\alpha\alpha'} = 1$ when the
neighboring α'-th cell lines within a forward or backward $\pi/3$ cone relative to the polarity
direction, and $\chi_{\alpha\alpha'} = -1$ otherwise [Fig. 2(i)]. Values of $S \approx 1$ indicate predominantly
antiparallel alignment within each domain, whereas $S \approx 0$ corresponds to random polarity.

By averaging S over the time interval $t = 0.95T$ to $t = T$, we confirm that phase separation
is accompanied by the emergence of strong antiparallel alignment [Fig. 2(j)]. This alignment, and the
resulting structure, emerges and remains stable under low-noise conditions, whereas it appears only
transiently or disappears entirely when noise is higher. These results demonstrate that motile cells
within the separated domains robustly form continuous, concentric antiparallel circulation patterns.

B. Ordering leads to dynamic scaling with 1/3 power-law domain growth

To investigate the macroscopic behavior of the observed phase separation, we first calculate the mean-square displacement (MSD) of motile-type cells [Fig. 3(a)] (see Appendix D). At short timescales, cells exhibit ballistic motion, characterized by an MSD $\propto \tau^2$. At longer timescales, the MSD follows a subdiffusive power law, MSD $\propto \tau^\lambda$ with $\lambda < 1$, for large J values, indicating constrained motion within a phase-separated structure. In contrast, for small J, the MSD displays normal diffusion, MSD $\propto \tau^1$, reflecting noise-dominated behavior. These results show that motile-type cells exhibit subdiffusive dynamics characteristic of phase-separating systems.

We next calculated the typical domain length, $R(t)$, to characterize the growth dynamics of the phase-separated domains [Fig 3(b)] (see Appendix D). In the parameter range where phase separation occurs, $R(t)$ grows according to a $t^{1/3}$ power law once separation is fully established, that is, when the alignment rate is moderate ($J \lesssim 7$) and noise is low ($D_r \lesssim 0.5$). Outside this range, the growth becomes slower, following a $t^{1/4}$ power law [Fig. 3(c)]. In both the 1/3 and 1/4 regimes, spatial correlation functions at different times collapse onto a single curve when distances are rescaled by the instantaneous domain length $R(t)$ [Fig. 3(d)]. This collapse indicates that the system obeys dynamic scaling, evolving self-similarly in time, with coarsening governed by a single characteristic growing length scale [51].

Our model exhibits a growth exponent of $\nu = 1/3$ in the domain-growth law $R \propto t^\nu$ within the low-noise regime. A similar exponent arises in motility-induced phase separation (MIPS) driven by run-and-tumble motion [52,53]. In contrast, $\nu = 1/4$ is commonly observed in MIPS driven by active Brownian motion [1,53,54] and in the standard Vicsek model [55,56]. The difference in scaling exponents suggests distinct mechanisms of domain coarsening. The $\nu = 1/4$ exponent is typically associated with domain coalescence mediated by diffusion–coalescence processes [57–60], although the microscopic origin of the 1/4 law in the Vicsek model remains debated. By comparison, the $\nu = 1/3$ exponent is characteristic of diffusion-limited growth [61–63], indicating that in our system, domain coarsening is primarily controlled by diffusive transport along the interfaces of motile-cell domains.

Moreover, to characterize cell dynamics during phase separation, we quantified the average speed of motile cells. We found that their speed increases over time when phase separation occurs [Fig. 3(e), left], whereas it remains approximately constant in the absence of separation [Fig. 3(e), right]. This behavior indicates that the mechanism driving phase separation in our system differs fundamentally from MIPS, where aggregation arises because repeated collisions slow particles down and reduce their effective speed [53].

C. Pairwise translational cell motion leads to persistent circulation

To clarify the elementary processes of cell circulation, we examined pairwise cell movements. We simulated the behavior of a simplified system composed of two motile-type cells, A and B, surrounded by nonmotile-type cells and found that a persistent pairwise motion of the motile-type cells emerged [Fig. 4(a) and Movie S3]. To quantify the persistence of this motion, we calculated the autocorrelation function of the average velocity of cells A and B, $C_{\bar{v}}(\tau) = \langle \bar{v}(t) \cdot \bar{v}(t + \tau) \rangle / \langle \bar{v}(t) \cdot \bar{v}(t) \rangle$, under low-noise ($D_{\mathrm{r}} = 0.01$) and high-noise ($D_{\mathrm{r}} = 1.00$) conditions at $J = 5.0$. We found that the correlation time was markedly longer under low-noise conditions than under the high-noise conditions [Fig. 4(b)], consistent with the persistence-time analysis of circulation obtained earlier [Figs. 2(h) and 3(f)].

To understand how the paired configuration persists, we analyzed the frequency of polarity alignment between the two motile cells and their interfacial tension [Fig. 4(c)]. Alignment was assessed by the inner product of the polarity vectors of cells A and B: $\boldsymbol{q}_A \cdot \boldsymbol{q}_B$. Under high-noise conditions, the frequency of aligned states increases as $\boldsymbol{q}_A \cdot \boldsymbol{q}_B$ increases; however, even at strong alignment, the interfacial tension remains positive. In contrast, under low-noise conditions, the frequency of the aligned configuration is higher than that under high-noise conditions. Remarkably, as alignment approaches the fully aligned state ($\boldsymbol{q}_A \cdot \boldsymbol{q}_B \sim 1$), the interfacial tension approaches zero, indicating that positive and negative tensions exerted by the rear and front cells cancel each other at their interface. This cancellation mechanism is consistent with a previous report of chain-like collective migration driven by a tension gradient [31].

These findings indicate that once a head-to-tail configuration forms, the pair behaves as a mechanically self-stabilizing unit. The persistence of polarity in low-noise environments prevents the cancellation of tensions from being disrupted, allowing the pair to migrate coherently until perturbed by noise, collisions, or shape fluctuations.

D. Self-aligned tension gradient induces stable antiparallel motion of the circulation flows

To clarify how the combination of tension gradient and self-alignment gives rise to antiparallel circulation, we constructed a simple mathematical model. Specifically, to analyze the distinct contributions of tension-gradient-driven interactions from self-propelled forces, we introduced both terms into a minimal two-rail system and compared their effects.

We consider two parallel circulation rails composed of motile cells [Fig. 5(a)]. Motivated by the

translational migration of motile cells observed in Fig. 4(a), we assume that within each rail, all cells share a common velocity and polarity direction. We denote the velocity and polarity of the i-th rail ($i = 1,2$) along the x-th axis by v_i and q_i, respectively. The polarity q_i is treated as a scalar defined by $q_i = \cos\theta_i$, where θ_i is the angle between the polarity vector and the rail direction. Each rail experiences friction against the substrate with a coefficient, η, fixed to the unit for simplicity. Under this setup, the motion equation of the i-th rail is written as

$$0 = -v_i + q_i + \beta q_j, \tag{8}$$

where $i, j = 1,2$ and $j \neq i$. The first term represents the substrate friction force. The second term represents a polarity-dependent self-propelled force. The third term represents a reaction force from another rail, proportional to the polarity of the j-th rail and controlled by β.

To understand the basic nature of the two-rail model, we first consider the case where the velocity is aligned with polarity ($\text{sgn}(v_i) = \text{sgn}(q_i)$). In this case, Eq. (8) describes several situations that combine driving forces with polarity regulation. When $\beta = 1$, Eq. (8) immediately yields $v_1 = v_2$ (i.e., q_1 and q_2 are aligned in the same direction), corresponding to parallel motion. This situation violates force balance and corresponds to the combined case with mutual alignment and self-propelled driving forces, as defined in previous studies [17,32]. In contrast, when $\beta = -1$, Eq. (8) immediately yields $v_1 = -v_2$, corresponding to antiparallel motion. In this situation, the action–reaction forces generated by q_1 and q_2, respectively, are in force balance, and the system corresponds to the self-aligned tension-gradient case, defined in our vertex model. These results demonstrate that antiparallel circulation arises naturally from the force-balanced nature of tension gradients.

To further examine the effects of self-alignment on circulation patterns, we next incorporate polarity regulation into the simple model. The time evolution of θ_i is given by

$$\frac{d\theta_i}{dt} = -Jv_i \sin\theta_i, \tag{9}$$

where J denotes the alignment rate. Combining Eqs. (8) and (9), we obtain the time evolution of the polarity q_i:

$$\frac{dq_i}{dt} = J(q_i + \beta q_j)(1 - q_i^2). \tag{10}$$

Solving Eq. (10) yields four fixed points of the polarity pair $(\pm 1, \pm 1)$ and $(\pm 1, \mp 1)$, corresponding to the parallel and antiparallel configurations, respectively. The points that satisfy $q_i + \beta q_j = 0$ are also fixed points; however, they correspond to $v_i = 0$. Therefore, we exclude these cases from consideration.

Using Eq. (10), we determined which fixed point the dynamics converge to and evaluated the

steady-state value of $v_1 v_2$ in the (q_1, q_2) phase space [color-coded in Fig. 5(b)]. Moreover, to

investigate the linear stability of the fixed points, we analytically calculated the eigenvalues λ_1 and

λ_2 of the Jacobian matrix $\mathbf{J}$ of Eq. (10) evaluated at each fixed point. We found that λ_1 and λ_2

reflect the stabilities of the parallel and antiparallel configurations, respectively. The Jacobian matrix

is $\mathbf{J} = J(-2\beta - 2)\mathbf{I}$ at $(q_1, q_2) = (\pm 1, \pm 1)$ and $\mathbf{J} = J(2\beta - 2)\mathbf{I}$ at $(q_1, q_2) = (\pm 1, \mp 1)$, where $\mathbf{I}$

denotes the identity matrix. Thus, the eigenvalues depend on β as $\lambda_1 = J(-2\beta - 2)$ and $\lambda_2 =$

$J(2\beta - 2)$ [Fig. 5(c)].

When the driving force is purely tension-gradient-based ($\beta = -1$), the polarity dynamics

converge to the antiparallel configuration [Fig. 5(b), left]. In this case, the antiparallel configuration

is linearly stable ($\lambda_2 = -4J < 0$), whereas the parallel configuration is not ($\lambda_1 = 0$) [Fig. 5(c)]. This

behavior is consistent with our vertex model, in which force-balanced tension gradients combined

with self-alignment give rise to antiparallel circulation. In contrast, when motion is driven solely by

self-propulsion ($\beta = 0$), the dynamics can converge to either the parallel or antiparallel configuration

depending on initial conditions [Fig. 5(b), middle], and both are equally stable ($\lambda_1 = \lambda_2 = -2J < 0$)

[Fig. 5(c)]. Finally, when the interaction corresponds to mutual alignment ($\beta = 1$), the polarity

dynamics converge to the parallel configuration [Fig. 5(b), right]; the parallel configuration is

linearly stable ($\lambda_1 = -4J < 0$), whereas the antiparallel configuration is not ($\lambda_2 = 0$) [Fig. 5(c)].

Therefore, antiparallel patterns do not emerge through mutual alignment alone.

These results demonstrate that the antiparallel pattern is a characteristic feature emerging from

self-aligned, force-balanced driving forces, distinct from dynamics governed by self-propulsion or

mutual alignment. Importantly, this mechanism is not limited to the vertex-model framework or to a

specific tension-gradient-driven interaction; rather, it may apply more generally when constituents

form head-to-tail units and interact through locally force-balanced couplings under self-alignment.

E. *D. discoideum* demonstrates an antiparallel circulation pattern

To examine whether antiparallel circulation arises in biological systems, we analyzed the collective

cell movements of *D. discoideum*. These cells are monopolar and motile, and they exhibit vortex-like

motion [20] even in the absence of chemotactic signals [64]. Their polarity is established through

front-to-back cell–cell contacts [64], which is consistent with the polarity–tension coupling assumed

in our model. This suggested that an antiparallel circulating state might exist but remain overlooked

in three-dimensional environments, where cell trajectories are more complex. To clearly identify cell

movements, we confined differentiated cells to quasi-2D geometry by seeding cells in a custom-

made 2.5-μm-thick chamber placed between a glass coverslip and polydimethylsiloxane [Fig. 6(a)]

(see Appendix E) [65,66]. The cells formed a disk-like aggregate [Fig. 6(b)], within which we

observed a circulation pattern strikingly similar to that predicted by our simulations [Fig. 6(c) and

Movie S4].

To quantify this behavior, we measured the direction, speed, and persistence time of cell motion

using the same definitions applied in our simulations [compare Fig. 6(d)–(f) with Fig. 3(f)–(h)]. For

persistence time, we further evaluated the significance by comparing the observed data with

randomized trajectories generated by bootstrap resampling. These analyses show that the quantitative

features of cell motion in the aggregates are qualitatively consistent with those from our model. This

finding demonstrates that *D. discoideum* cells exhibit continuous, concentric antiparallel circulation

patterns similar to those predicted in our simulations. We note that the circulations were not always

single-cell-wide lanes but sometimes consisted of several cells, likely reflecting side-by-side

adhesion [67], an effect not included in our model, leading to broader circulation bands than those

observed in the vertex simulations.

Furthermore, when *D. discoideum* cells were seeded at low density, we frequently observed

persistent translational motion in which two cells aligned in a head-to-tail configuration and migrated

together as a pair [Fig. 6(g) and Movie S5], comparable to the configuration observed in our

simulations [Fig. 4(a)]. We quantified this translational alignment using the inner product of the

normalized velocity vectors of cells A and B, $\hat{\boldsymbol{v}}_A \cdot \hat{\boldsymbol{v}}_B$ [Fig. 6(h)]. Fully aligned states ($\hat{\boldsymbol{v}}_A \cdot \hat{\boldsymbol{v}}_B \sim 1$)

were frequent, consistent with the simulations [Fig. 4(c)]. This configuration persists until the pair

encounters another cluster—typically a large one, at which point the pair merges or reorganizes. This

behavior resembles the migration patterns of 2D cell clusters under an imposed gradient of the

chemoattractant, cAMP [64]. These results demonstrate that *D. discoideum* cells can induce a head-

to-tail configuration as a mechanically self-stabilizing unit, suggesting that a similar mechanism

predicted by the model could underlie the antiparallel circulations observed in *D. discoideum*.

IV. DISCUSSION

Our results show that antiparallel cell circulation can emerge from polarized, force-balanced

interactions, providing a distinct alternative to the widely studied picture of self-propelled cells

driven by substrate traction. In many active-matter models of multiple cells, motility is represented

as a polarity-dependent self-propulsion force that is balanced by friction at the cell–substrate

interface. In our framework, in contrast, the driving forces are polarity-dependent interfacial forces

(or, more generally, polarized cell–cell interactions) that act along contacts between cells. These

forces are locally force-balanced, where each interface exerts equal and opposite forces on its

neighboring cells—yet their polarization, together with polarity–velocity alignment, generates

sustained microscopic circulation. This mechanism shows that, when combined with self-alignment, alternative driving modes with different patterns of force transfer between neighboring cells can generate robust dynamic patterns at the cell level in multicellular systems.

This difference in how forces are applied is reflected in the emergent patterns. Existing self-alignment and mutual alignment, including Vicsek-type models, can generate vortices [68–70], unidirectional circulation [71,72], or flocking states [16–19], but the circulation within each structure is typically single-directional and often relies on center-of-mass propulsion or geometric confinement. In contrast, our model produces interdigitated antiparallel circulation, where neighboring lanes of cells move in opposite directions while the tissue remains confluent and mechanically compact. To the best of our knowledge, such constituent-level antiparallel circulation has not been reported in models based on MIPS, mutual alignment, or conventional self-alignment rules. This highlights that polarized force-balanced interactions, combined with self-alignment of polarity to velocity, define a qualitatively different route to organized flow patterns in cell collectives.

The same mechanism that produces antiparallel circulation also leads to a distinct form of phase separation. In MIPS, clusters form because particles repeatedly slow down upon encountering dense regions; reduced speed promotes accumulation. In our system, the opposite trend appears as the separation degree, μ, increases [Fig. 2(b)], the speed of motile cells also increases [Fig. 3(e)]. Phase separation is therefore circulation-driven, rather than slowdown-driven, establishing positive feedback between microscopic circulation and mesoscopic domain growth. Moreover, although mutual alignment often exhibits phase separation through large density fluctuations reminiscent of gas–liquid separation [73], our system maintains nearly constant density within compact domains, and domain growth is constrained by diffusion [Fig. 3(b) and 3(c)]. We refer to this distinct mechanism as circulation-induced phase separation (CirPS).

An important aspect of our results is that the mechanism is robust to how polarized force-balanced interactions are realized. We first introduced them through polarity-biased interfacial tensions but then showed that simplified force-balanced attractive–repulsive interactions between neighboring cell rails can generate qualitatively similar antiparallel circulations. This indicates that the phenomenon does not depend on the detailed form of interfacial tension; instead, the combination of (i) polarity bias and (ii) local force balance at cell–cell contacts is essential. We therefore expect that other realizations of polarized force-balanced interactions, for example, those arising from active stress patterns or directed shape changes, may likewise produce circulation-driven pattern formation.

The cellular behaviors underpinning CirPS—persistent flows and local rearrangements—are frequently observed in multicellular systems, suggesting that similar mechanisms may operate *in*

vivo. In our experiments, we observed antiparallel circulations in 2D aggregates of *D. discoideum* [Fig. 6], consistent with the model prediction. Cell reorientations are thought to be important for breaking up *Dictyostelium* aggregates and regulating aggregate size and sorting; our findings raise the possibility that antiparallel circulation and CirPS contribute to these processes. Similar circulation-based mechanisms could also influence the breakup or remodeling of tumor clusters and other dense cell assemblies. Overall, our work identifies polarized force-balanced interactions as a minimal and robust route to antiparallel cell circulation and CirPS, broadening the landscape of active-matter mechanisms that can underlie organization and remodeling in multicellular systems.

DATA AVAILABILITY

The data that support the findings of this article are openly available [citation number to be inserted here].

ACKNOWLEDGEMENTS

We thank all the members of the Okuda Laboratory for their discussions. This work was supported by the WISE Program for Nano-Precision Medicine, Science, and Technology of Kanazawa University, Ministry of Education, Culture, Sports, Science and Technology (MEXT); the Japan Science and Technology Agency (JST), CREST [Grant No. JPMJCR1921, JPMJCR24B2, JPMJCR1923]; the Japan Society for the Promotion of Science (JSPS), KAKENHI [Grant No. 21K15081, 22H05170, 23H00384, 23H04304, 24H01398, 24H01937, 25K01118, 25H01363, 25H01771, 25K22468]; the Japanese Agency for Medical Research and Development (AMED), a Program for Technological Innovation of Regenerative Medicine Grant [23bm0704065h0003], and the World Premier International Research Center Initiative, MEXT, Japan.

References

[1] G. S. Redner, M. F. Hagan, and A. Baskaran, Structure and dynamics of a phase-separating active colloidal fluid, Phys. Rev. Lett. **110**, 055701 (2013).

[2] I. Buttinoni, J. Bialké, F. Kümmel, H. Löwen, C. Bechinger, and T. Speck, Dynamical clustering and phase separation in suspensions of self-propelled colloidal particles, Phys. Rev. Lett. **110**, 238301 (2013).

[3] A. Zöttl and H. Stark, Emergent behavior in active colloids, Journal of Physics: Condensed Matter **28**, (2016).

[4] C. C. Maass, C. Krüger, S. Herminghaus, and C. Bahr, Swimming droplets, Annu. Rev. Condens. Matter Phys. **7**, 171 (2016).

[5] C. Jin, C. Krüger, and C. C. Maass, Chemotaxis and autochemotaxis of self-propelling droplet swimmers, Proc. Natl. Acad. Sci. U. S. A. **114**, 5089 (2017).

[6] C. Dombrowski, L. Cisneros, S. Chatkaew, R. E. Goldstein, and J. O. Kessler, Self-concentration and large-scale coherence in bacterial dynamics, Phys. Rev. Lett. **93**, 098103 (2004).

[7] H. H. Wensink, J. Dunkel, S. Heidenreich, K. Drescher, R. E. Goldstein, H. Löwen, and J. M. Yeomans, Meso-scale turbulence in living fluids, Proc. Natl. Acad. Sci. U. S. A. **109**, 14308 (2012).

[8] J. Dunkel, S. Heidenreich, K. Drescher, H. H. Wensink, M. Bär, and R. E. Goldstein, Fluid dynamics of bacterial turbulence, Phys. Rev. Lett. **110**, 228102 (2013).

[9] E. Lushi, H. Wioland, and R. E. Goldstein, Fluid flows created by swimming bacteria drive self-organization in confined suspensions, Proc. Natl. Acad. Sci. U. S. A. **111**, 9733 (2014).

[10] W. Xi, T. B. Saw, D. Delacour, C. T. Lim, and B. Ladoux, Material approaches to active tissue mechanics, Nat. Rev. Mater. **4**, 23 (2018).

[11] R. Alert and X. Trepat, Living cells on the move, Phys. Today **74**, 30 (2021).

[12] S. Shankar, A. Souslov, M. J. Bowick, M. C. Marchetti, and V. Vitelli, Topological active matter, Nat. Rev. Phys. **4**, 380 (2022).

[13] A. Doostmohammadi, J. Ignés-Mullol, J. M. Yeomans, and F. Sagués, Active nematics, Nat. Commun. **9**, 1 (2018).

[14] R. Alert, J. Casademunt, and J.-F. Joanny, Active turbulence, Annu. Rev. Condens. Matter Phys. **13**, 143 (2022).

[15] T. Sanchez, D. T. N. Chen, S. J. DeCamp, M. Heymann, and Z. Dogic, Spontaneous motion in hierarchically assembled active matter, Nature **491**, 431 (2012).

[16] C. W. Reynolds, Flocks, herds and schools: A distributed behavioral model, Comput. Graph. (ACM) **21**, 25 (1987).

[17] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen I., and O. Shochet, Novel type of phase transition in a system of self-driven particles, Phys. Rev. Lett. **75**, 1226 (1995).

[18] M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, M. Rao, and R. A. Simha, Hydrodynamics of soft active matter, Rev. Mod. Phys. **85**, 1143 (2013).

[19] Y. Shen, J. O'Byrne, A. Schoenit, A. Maitra, R.-M. Mège, R. Voituriez, and B. Ladoux, Flocking and giant fluctuations in epithelial active solids, Proc. Natl. Acad. Sci. U. S. A. **122**, e2421327122 (2025).

[20] W.-J. Rappel, A. Nicol, A. Sarkissian, H. Levine, and W. F. Loomis, Self-organized vortex state in two-DimensionalDictyosteliumDynamics, Phys. Rev. Lett. **83**, 1247 (1999).

[21] K. Doxzen, S. R. K. Vedula, M. C. Leong, H. Hirata, N. S. Gov, A. J. Kabla, B. Ladoux, and C. T. Lim, Guidance of collective cell migration by substrate geometry, Integr. Biol. (Camb.) **5**,

446   1026 (2013).

[22] C. Blanch-Mercader, V. Yashunsky, S. Garcia, G. Duclos, L. Giomi, and P. Silberzan, Turbulent dynamics of epithelial cell cultures, Phys. Rev. Lett. **120**, 208101 (2018).

[23] S.-Z. Lin, W.-Y. Zhang, D. Bi, B. Li, and X.-Q. Feng, Energetics of mesoscale cell turbulence in two-dimensional monolayers, Commun. Phys. **4**, (2021).

[24] B. Szabó, G. J. Szöllösi, B. Gönci, Z. Jurányi, D. Selmeczi, and T. Vicsek, Phase transition in the collective migration of tissue cells: experiment and model, Phys. Rev. E Stat. Nonlin. Soft Matter Phys. **74**, 061908 (2006).

[25] S.-Z. Lin, S. Ye, G.-K. Xu, B. Li, and X.-Q. Feng, Dynamic Migration Modes of Collective Cells, Biophys. J. **115**, 1826 (2018).

[26] D. Sarkar, G. Gompper, and J. Elgeti, A minimal model for structure, dynamics, and tension of monolayered cell colonies, Commun. Phys. **4**, (2021).

[27] T. H. Tan, A. Amiri, I. Seijo-Barandiaran, M. F. Staddon, A. Materne, S. Tomas, C. Duclut, M. Popović, A. Grapin-Botton, and F. Jülicher, Emergent chirality in active solid rotation of pancreas spheres, PRX Life **2**, (2024).

[28] W. Wang, R. A. Law, E. Perez Ipiña, K. Konstantopoulos, and B. A. Camley, Confinement, Jamming, and Adhesion in Cancer Cells Dissociating from a Collectively Invading Strand, PRX Life **3**, 013012 (2025).

[29] K. Sato, T. Hiraiwa, E. Maekawa, A. Isomura, T. Shibata, and E. Kuranaga, Left-right asymmetric cell intercalation drives directional collective cell movement in epithelial morphogenesis, Nat. Commun. **6**, 10074 (2015).

[30] S. Okuda, E. Kuranaga, and K. Sato, Apical Junctional Fluctuations Lead to Cell Flow while Maintaining Epithelial Integrity, Biophys. J. **116**, 1159 (2019).

[31] S. Okuda and K. Sato, Polarized interfacial tension induces collective migration of cells, as a cluster, in a 3D tissue, Biophys. J. **121**, 1856 (2022).

[32] J. Toner, Y. Tu, and S. Ramaswamy, Hydrodynamics and phases of flocks, Ann. Phys. (N. Y.) **318**, 170 (2005).

[33] S. Ramaswamy, The mechanics and statistics of active matter, Annu. Rev. Condens. Matter Phys. **1**, 323 (2010).

[34] M. Basan, J. Elgeti, E. Hannezo, W.-J. Rappel, and H. Levine, Alignment of cellular motility forces with tissue flow as a mechanism for efficient wound healing, Proc. Natl. Acad. Sci. U. S. A. **110**, 2452 (2013).

[35] B. A. Camley, Y. Zhang, Y. Zhao, B. Li, E. Ben-Jacob, H. Levine, and W.-J. Rappel, Polarity mechanisms such as contact inhibition of locomotion regulate persistent rotational motion of mammalian cells on micropatterns, Proc. Natl. Acad. Sci. U. S. A. **111**, 14770 (2014).

[36] C. Malinverno et al., Endocytic reawakening of motility in jammed epithelia, Nat. Mater. **16**, 587 (2017).

[37] P. Baconnier, O. Dauchot, V. Démery, G. Düring, S. Henkes, C. Huepe, and A. Shee, Self-

aligning polar active matter, Rev. Mod. Phys. **97**, (2025).

[38] V. Petrolli et al., Confinement-induced transition between wavelike collective cell migration
modes, Phys. Rev. Lett. **122**, 168101 (2019).

[39] H. Honda and G. Eguchi, How much does the cell boundary contract in a monolayered cell
sheet?, J. Theor. Biol. **84**, 575 (1980).

[40] R. Farhadifar, J.-C. Röper, B. Aigouy, S. Eaton, and F. Jülicher, The influence of cell mechanics,
cell-cell interactions, and proliferation on epithelial packing, Curr. Biol. **17**, 2095 (2007).

[41] A. G. Fletcher, M. Osterfield, R. E. Baker, and S. Y. Shvartsman, Vertex models of epithelial
morphogenesis, Biophys. J. **106**, 2291 (2014).

[42] S. Alt, P. Ganguly, and G. Salbreux, Vertex models: from cell mechanics to tissue
morphogenesis, Philos. Trans. R. Soc. Lond. B Biol. Sci. **372**, (2017).

[43] S. Henkes, Y. Fily, and M. C. Marchetti, Active jamming: self-propelled soft particles at high
density, Phys. Rev. E Stat. Nonlin. Soft Matter Phys. **84**, 040301 (2011).

[44] B. A. Camley and W.-J. Rappel, Velocity alignment leads to high persistence in confined cells,
Phys. Rev. E Stat. Nonlin. Soft Matter Phys. **89**, 062705 (2014).

[45] K.-D. N. T. Lam, M. Schindler, and O. Dauchot, Self-propelled hard disks: implicit alignment
and transition to collective motion, New J. Phys. **17**, 113056 (2015).

[46] F. Giavazzi Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e
Medicina Traslazionale, 20090 Segrate, Italy. roberto. cerbino@unimi. it, M. Paoluzzi, M.
Macchi, D. Bi, G. Scita, M. L. Manning, R. Cerbino, and M. C. Marchetti, Flocking transitions
in confluent tissues, Soft Matter **14**, 3471 (2018).

[47] G. Erdemci-Tandogan and M. L. Manning, Effect of cellular rearrangement time delays on the
rheology of vertex models for confluent tissues, PLoS Comput. Biol. **17**, e1009049 (2021).

[48] M. Bergert, A. Erzberger, R. A. Desai, I. M. Aspalter, A. C. Oates, G. Charras, G. Salbreux, and
E. K. Paluch, Force transmission during adhesion-independent migration, Nat. Cell Biol. **17**,
524 (2015).

[49] K. Sato, A cell membrane model that reproduces cortical flow-driven cell migration and
collective movement, Front. Cell Dev. Biol. **11**, 1126819 (2023).

[50] D. Bi, X. Yang, M. C. Marchetti, and M. L. Manning, Motility-driven glass and jamming
transitions in biological tissues, Phys. Rev. X. **6**, (2016).

[51] A. J. Bray, Theory of phase-ordering kinetics, Adv. Phys. **51**, 481 (2002).

[52] A. G. Thompson, J. Tailleur, M. E. Cates, and R. A. Blythe, Lattice models of nonequilibrium
bacterial dynamics, J. Stat. Mech. **2011**, P02029 (2011).

[53] M. E. Cates and J. Tailleur, Motility-Induced Phase Separation, Annual Review of Condensed
Matter Physics **6**, 219 (2015).

[54] J. Stenhammar, D. Marenduzzo, R. J. Allen, and M. E. Cates, Phase behaviour of active
Brownian particles: the role of dimensionality, Soft Matter **10**, 1489 (2014).

[55] S. Dey, D. Das, and R. Rajesh, Spatial structures and giant number fluctuations in models of

active matter, Phys. Rev. Lett. **108**, 238001 (2012).

[56] N. Katyal, S. Dey, D. Das, and S. Puri, Coarsening dynamics in the Vicsek model of active matter, Eur. Phys. J. E Soft Matter **43**, 10 (2020).

[57] H. Furukawa, Kinetics of droplet growth in systems with arbitrary degeneracy, Phys. Rev. A **29**, 2160 (1984).

[58] M. Kolb, Unified Description of Static and Dynamic Scaling for Kinetic Cluster Formation, Phys. Rev. Lett. **53**, 1653 (1984).

[59] P. Meakin, Diffusion-limited droplet coalescence, Physica A **165**, 1 (1990).

[60] C. P. Beatrici, R. M. C. de Almeida, and L. G. Brunnet, Mean-cluster approach indicates cell sorting time scales are determined by collective dynamics, Phys. Rev. E. **95**, 032402 (2017).

[61] I. M. Lifshitz and V. V. Slyozov, The kinetics of precipitation from supersaturated solid solutions, J. Phys. Chem. Solids **19**, 35 (1961).

[62] T. M. Rogers and R. C. Desai, Numerical study of late-stage coarsening for off-critical quenches in the Cahn-Hilliard equation of phase separation, Phys. Rev. B Condens. Matter **39**, 11956 (1989).

[63] A. Nakajima and S. Ishihara, Kinetics of the cellular Potts model revisited, New J. Phys. **13**, 033035 (2011).

[64] T. Fujimori, A. Nakajima, N. Shimada, and S. Sawai, Tissue self-organization based on collective cell migration by contact activation of locomotion and chemotaxis, Proc. Natl. Acad. Sci. U. S. A. **116**, 4291 (2019).

[65] T. Fujimori, H. Hashimura, and S. Sawai, Imaging-based analysis of cell-cell contact-dependent migration in Dictyostelium, Methods Mol. Biol. **2828**, 23 (2024).

[66] H. Hashimura, S. Kuwana, H. Nakagawa, K. Abe, T. Adachi, T. Sugita, S. Fujishiro, G. Honda, and S. Sawai, Multi-color fluorescence live-cell imaging in Dictyostelium discoideum, Cell Struct. Funct. **49**, 135 (2024).

[67] K. Müller and G. Gerisch, A specific glycoprotein as the target site of adhesion blocking Fab in aggregating Dictyostelium cells, Nature **274**, 445 (1978).

[68] A. Costanzo and C. K. Hemelrijk, Spontaneous emergence of milling (vortex state) in a Vicsek-like model, J. Phys. D Appl. Phys. **51**, 134004 (2018).

[69] X. Wang, P.-C. Chen, K. Kroy, V. Holubec, and F. Cichos, Spontaneous vortex formation by microswimmers with retarded attractions, Nat. Commun. **14**, 56 (2023).

[70] H. Xu and Y. Wu, Self-enhanced mobility enables vortex pattern formation in living matter, Nature **627**, 553 (2024).

[71] A. C. Quillen, J. P. Smucker, and A. Peshkov, Boids in a loop: Self-propelled particles within a flexible boundary, Phys. Rev. E. **101**, 052618 (2020).

[72] Y. Wu, J. Li, D. Chen, X. Li, and H.-T. Zhang, Pattern phase transitions in 3D minimal Vicsek model, EPL **134**, 50004 (2021).

[73] A. P. Solon, H. Chaté, and J. Tailleur, From phase to microphase separation in flocking models:

560   the essential role of nonequilibrium fluctuations, Phys. Rev. Lett. **114**, 068101 (2015).

561 [74] D. Bi, J. H. Lopez, J. M. Schwarz, and M. L. Manning, A density-independent rigidity transition

562   in biological tissues, Nat. Phys. **11**, 1074 (2015).

563 [75] B. Loewe, M. Chiang, D. Marenduzzo, and M. C. Marchetti, Solid-liquid transition of

564   deformable and overlapping active particles, Phys. Rev. Lett. **125**, 038003 (2020).

565 [76] M. Chiang, A. Hopkins, B. Loewe, D. Marenduzzo, and M. C. Marchetti, Multiphase field

566   model of cells on a substrate: From three dimensional to two dimensional, Phys. Rev. E. **110**,

567   044403 (2024).

568 [77] See Supplemental Material at [URL] for further details on the theoretical model and numerical

569   simulations.

570 [78] D. A. Huse, Corrections to late-stage behavior in spinodal decomposition: Lifshitz-Slyozov

571   scaling and Monte Carlo simulations, Phys. Rev. B Condens. Matter **34**, 7845 (1986).

572 [79] M. Durand, Large-scale simulations of biological cell sorting driven by differential adhesion

573   follow diffusion-limited domain coalescence regime, PLoS Comput. Biol. **17**, e1008576 (2021).

Appendices

APPENDIX A: EFFECTS OF TISSUE FLUIDITY ON PHASE SEPARATION

Varying cell properties, such as deformability, motility, and noise, can induce a jamming transition, during which tissue fluidity changes drastically [50,74–76]. To investigate the effects of tissue fluidity on phase separation, we first reproduced the jamming transition using our model. Here, we considered a tissue composed entirely of motile-type cells without self-alignment. In this model, the shape index p_0, the strength of the tension gradient γ^p, and the noise intensity D_r can be used to tune tissue fluidity, which corresponds to cell deformability, motility, and noise. For simplicity, the noise intensity was fixed at $D_r = 1$, and the mixing ratio and alignment rate were set to $\phi = 1$ and $J = 0$, respectively. To quantitatively assess tissue fluidity, we calculated the effective diffusion coefficient $D_{\text{eff}} := D_s/D_0$, where D_s is the self-diffusion coefficient defined as $D_s := \text{MSD}(\tau)/(4\tau)$ averaged for $\tau > 0.45T$, and D_0 is the free diffusion coefficient given by $D_0 := 2\pi^2(\gamma^p/\eta)^2/D_r$ [31,50] (see the Supplemental material [77] for the detailed calculation of D_0). Here, $\text{MSD}(\tau)$ denotes the mean-square displacement of motile-type cells at a time shift of τ (see Appendix C). We classified systems with $D_{\text{eff}} < 10^{-3}$ as being a solid-like state and those with $D_{\text{eff}} \geq 10^{-3}$ as fluid-like. As a result, we found that tissue fluidity transitioned between solid-like and fluid-like states depended on the values of p_0 and γ^p, indicating the occurrence of a jamming transition [Fig. 7(a)].

We next considered a mixture of motile-type cells with self-alignment and nonmotile-type cells. In this case, the mixing ratio and noise intensity were set to $\phi = 0.5$ and $D_r = 1$. By varying p_0 and J, we observed that phase separation occurred only when the tissue was in a fluid-like state [Fig. 7(b)]. Based on these findings, all numerical simulations presented in the main text were performed under fluid-like conditions, as phase separation does not occur in the solid-like state.

APPENDIX B: DEPENDENCE ON THE INITIAL CONDITIONS

To investigate the dependence of phase separation on the initial conditions, we conducted simulations using both homogeneous and pre-separated initial states. The separated state was prepared by placing a circular domain of motile-type cells at the center of the system. When $J > D_r$, phase separation occurred regardless of the initial condition, although the degree of separation, μ varied depending on the initial state [Fig. 8(a)]. In contrast, when $J < D_r$, the system eventually relaxed into a homogeneous state, irrespective of the initial condition, and μ converged to zero [Fig. 8(b)]. These results indicate that the occurrence of phase separation is not influenced by the initial condition. However, when phase separation does occur, the degree of separation is influenced by the

608 initial condition.

APPENDIX C: CALCULATION OF THE CIRCULATION VELOCITY AND THE PERSISTENT TIME OF THE CIRCULATION DIRECTION

612 To quantify the antiparallel circulation pattern, we calculated the speed and direction of motile-type

613 cell motion with respect to the center of circulation $\widetilde{\boldsymbol{R}}(t)$ for each domain, which was determined

614 from the velocity profile at each time t as follows. Using the velocity vector, $\boldsymbol{v}_\alpha$ and the position

615 vector, $\boldsymbol{r}_\alpha - \widetilde{\boldsymbol{R}}$, which connects the circulation center and the centroid of the α-th cell, $\widetilde{\boldsymbol{R}}(t)$, is

616 defined as the point $\boldsymbol{R}'$ that minimizes the following function,

$$\sum_{\alpha=1}^{N_c} [\boldsymbol{v}_\alpha \cdot (\boldsymbol{r}_\alpha - \boldsymbol{R}')]^2. \tag{11}$$

617 The optimal circulation center, $\widetilde{\boldsymbol{R}}$ can be written explicitly as

$$\widetilde{\boldsymbol{R}} = \left(\sum_{\alpha=1}^{N_c} \boldsymbol{v}_\alpha \boldsymbol{v}_\alpha^\top \right)^{-1} \left(\sum_{\alpha=1}^{N_c} \boldsymbol{v}_\alpha \boldsymbol{v}_\alpha^\top \boldsymbol{r}_\alpha \right). \tag{11}$$

618 Next, the circulation velocity v_α^{circ} of the α-th cell at each time point was calculated as the

619 projection of its velocity vector onto the tangential direction of the circular path. This is defined as

$$v_\alpha^{\text{circ}} := \frac{\boldsymbol{v}_\alpha \wedge (\boldsymbol{r}_\alpha - \widetilde{\boldsymbol{R}})}{\|\boldsymbol{r}_\alpha - \widetilde{\boldsymbol{R}}\|} = \frac{v_{\alpha x}(r_{\alpha y} - \tilde{R}_y) - v_{\alpha y}(r_{\alpha x} - \tilde{R}_x)}{\sqrt{(r_{\alpha x} - \tilde{R}_x)^2 + (r_{\alpha y} - \tilde{R}_y)^2}}. \tag{13}$$

620 Here, v_α^{circ} is positive when the cell is circulating in the clockwise direction. Figures 2(a) and 4(c)

621 show the spatial pattern of the circulation direction, as indicated by the sign of v_α^{circ} at a single time

622 point. These findings clearly exhibit an antiparallel circulation pattern.

623 To investigate the frequency of the reversal movement of the cells, we calculated the persistent

624 time of the circulation direction. Using the autocorrelation function

$$\rho_\alpha(\tau) = \langle \text{sgn}[v_\alpha^{\text{circ}}(t)] \text{sgn}[v_\alpha^{\text{circ}}(t+\tau)] \rangle_t, \tag{14}$$

625 the persistent time, τ_α is defined as the first zero-crossing point, i.e.,

$$\tau_\alpha = \min\{ \tau \mid \rho_\alpha(\tau) \le 0 \}. \tag{15}$$

APPENDIX D: MACROSCOPIC BEHAVIOR OF THE CELLS AND DOMAINS

628 We calculated the MSD of motile-type cells during the time interval, $T/2 \le t \le T$ as follows,

$$\text{MSD}(\tau) := \frac{1}{(T/2) - \tau} \int_{T/2}^{T-\tau} dt \left[\frac{1}{N_{\text{m}}} \sum_{\alpha \in \mathcal{M}} \|\boldsymbol{r}_\alpha(\tau + t) - \boldsymbol{r}_\alpha(t)\|^2 \right], \tag{16}$$

where τ is a time shift within the range, $\tau \in [0, T/2)$. For short timescales, the ballistic diffusion
regime ($\text{MSD} \propto \tau^2$) expanded as the system transitioned from a homogeneous to a phase separation
state by varying the alignment rate [Fig. 3(a)]. This expansion is attributed to the increased
persistence of motile-type cells when $J/D_{\text{r}} > 1$ [44].

For longer timescales, we observed subdiffusive behavior, characterized by $\text{MSD} \propto \tau^\lambda$ with
$\lambda < 1$, when J was sufficiently large, indicating restricted motion of motile-type cells under
complete phase separation. In contrast, when J was small, we found normal diffusion behavior,
characterized by $\text{MSD} \propto \tau^1$, where motion was primarily governed by noise.

Next, we examined the time evolution of the typical domain length. To simplify the calculation,
we first define the "existence probability" $f_\alpha(\boldsymbol{x}, t)$ of the α-th cell at coordinate $\boldsymbol{x}$ as

$$f_\alpha(\boldsymbol{x}, t) := \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{\|\boldsymbol{x} - \boldsymbol{r}_\alpha(t)\|^2}{2\sigma^2} \right], \tag{17}$$

639 where σ is the standard deviation of f_α, regarded as a spatial scale of a single cell. We set $\sigma =$
640 $1/2\sqrt{\pi}$ for all cells, which is equal to half of the radius of the circle with unit area. Using f_α, we
641 introduced the distribution function of motile-type cells, $f(\boldsymbol{x}, t)$, defined as

$$f(\boldsymbol{x}, t) := \frac{1}{N_{\text{c}}} \sum_{\alpha=1}^{N_{\text{c}}} \omega_\alpha f_\alpha(\boldsymbol{x}, t), \tag{18}$$

642 where ω_α is given by

$$\omega_\alpha = \begin{cases} 1 & (\alpha \in \mathcal{M}) \\ -1 & (\alpha \in \mathcal{N}) \end{cases}. \tag{19}$$

643 Note that f can be negative when nonmotile-type cells are near coordinate $\boldsymbol{x}$. Using f, we defined
644 the spatial correlation function of cell types $C(r, t)$ [63,78,79] as

$$C(r, t) := \langle s(\boldsymbol{x}_0, t) s(\boldsymbol{x}_0 + \boldsymbol{x}, t) \rangle - \langle s(\boldsymbol{x}_0, t) \rangle^2, \tag{20}$$

645 where $r = \|\boldsymbol{x}\|$ and $s(\boldsymbol{x}, t) = \text{sgn} \, f(\boldsymbol{x}, t)$. The average $\langle \cdot \rangle$ is taken over the coordinate $\boldsymbol{x}_0$ and
646 the displacement $\boldsymbol{x}$ with the given distance of $r = \|\boldsymbol{x}\|$. The domain length $R(t)$ is defined as the
647 first zero-crossing point of the correlation function, $C(r, t)$, i.e.,

$$R(t) := \min\{ r \mid C(r, t) \le 0 \}. \tag{21}$$

We first confirmed that $C(r, t)$ for different time points converged to the same curve as a
function of $r/R(t)$ [Fig. 3(d)]. Furthermore, domain length follows $R \propto t^\nu$ with growth exponent
ν, which indicates the mechanisms of domain growth. Under conditions in which the noise intensity

is very low ($D_r \lesssim 0.25$) and the alignment rate is moderate ($J \lesssim 7$), $\nu \approx 1/3$ [Figs. 3(b) and 3(c)],

suggesting that the domain growth is governed by diffusion [62]. As the noise intensity or the

alignment rate increases ($J \gtrsim 7$), the growth exponent rapidly shifts to $\nu \approx 1/4$ [Figs. 3(b) and

3(c)], suggesting that the domain growth is dominated by direct domain coalescence (diffusion-

coalescence) processes [57–60].

APPENDIX E: OBSERVATION OF *DICTYOSTELIUM DISCOIDEUM* AGGREGATION

The *D. discoideum* Ax4 cell line carrying the plasma membrane marker PKBR1(N150)-mScarlet-I-

2x at the *act5* locus [66] was used. The cell culture, development, microfabrication, and observation

were performed as described previously [64,66]. To image a monolayer cell population [Fig. 6(b)],

cells dissociated from the slugs were suspended at a density of 2.2×10^5 cells/mL in phosphate

buffer (PB: 12 mM KH_2PO_4, 8 mM Na_2HPO_4, pH 6.5). To observe the translational motion of two

cells [Fig. 6(g)], cells were suspended at a density of 5×10^4 cells/mL in PB. Then, 0.4 mL of cell

suspension was loaded into a microfluidic chamber with a height of 2.5 µm [Fig. 6(b)] or 4 [Fig.

6(g)]. Fluorescence images were acquired at 22°C using an inverted microscope (IX83,

Olympus/Evident) equipped with a multibeam confocal scanning unit (CSU-W1, Yokogawa) and

CMOS cameras (ORCA-FusionBT, Hamamatsu Photonics).

The membrane images of *D. discoideum* were collected at 5-second intervals to obtain $I = 361$

frames [Fig. 6(b)]. The raw images were binarized using Trainable Weka Segmentation in Fiji, and

cells were detected by connected component labeling. Hereafter, cells were represented by their

centroid positions. A pair of cells in consecutive frames was identified as the same cell if the

opponent had the smallest distance among other cells in the same frame. In this way, we obtained

$N_c = 507$ trajectories $\{r_\alpha(i)\}_{\alpha=1,\cdots,N_c, i=1,\cdots,I}$, which included more than 100 frames. Note that some

cells were occasionally not detected so that a true trajectory with full length could be divided into

multiple trajectories with shorter lengths. Then, cells were assigned with velocity vectors, $\{v_\alpha(i)\}$

at each time point. We applied a mean filter with a length of 10 frames before taking the difference

between consecutive centroid positions to remove the noise. For the antiparallel circulation pattern

analysis, the circulation velocity and persistent time of the circulation direction were calculated at

each time point i (see Appendix C).

FIG. 1. Setup of the vertex model with a self-aligned tension gradient. (a) The system is represented by a polygonal tiling with periodic boundary conditions. Each polygon expresses the shape of an individual cell. Purple and gray polygons represent motile-type and nonmotile-type cells, respectively. Arrows indicate polarity vectors of the motile-type cells. (b) Schematic illustration of a T1 transition. When the edge length, l_ε, becomes shorter than Δl, the cell configuration is rearranged by reconnecting the network of vertices and edges. (c) Tension gradient generated within a motile-type cell at its boundary with adjacent cells. Relative to the polarity vector, negative tension at the leading edge causes the edge to extend, and positive tension causes the rear edge to contract. (d) Schematic illustration of self-alignment. Each polarity vector tends to align with the velocity of the cell centroid with an alignment rate, J, and undergoes rotational diffusion with noise intensity, D_r.

FIG. 2. Phase separation and the emergence of antiparallel cell circulation in the vertex model with a self-aligned tension gradient. (a) Time evolution of the system with an alignment rate of $J = 4.0$ and a noise intensity of $D_r = 0.0$ (also shown in Movies S1 and S2). Motile-type cells rotating clockwise (CW) around the center of circulation are colored red, and those rotating counterclockwise (CCW) are colored blue. (b) Time evolution of the separation degree, μ, for two parameter sets: $(J, D_r) = (6.0, 0.0)$ and $(J, D_r) = (2.0, 3.0)$. The line and band widths in the plot indicate the statistical average values and standard deviations, respectively. (c) Phase diagram of the separation degree, μ, as a function of J and D_r. (d) Separation degree, μ, as a function of J with $D_r = \{0.0, 1.0, 2.0, 3.0, 4.0, 5.0\}$. (e) Schematic illustration of antiparallel cell circulation within a motile-type cell domain. Red arrows represent CW trajectories, and blue arrows represent CCW ones. The circulations are concentrically aligned, with adjacent layers moving in antiparallel directions. (f) Distribution of cell migration directions (CW: red; CCW: blue) as a function of cell velocity angle with $(J, D_r) = (5.0, 0.01)$. The cell velocity angle is defined as the angle between the direction of cell migration and the vector from the domain center to the center of circulation. (g) Average cell speed as a function of cell velocity angle with $(J, D_r) = (5.0, 0.01)$. (h) Distribution of the circulation persistence time, τ_α, with J for $D_r = 0.01$ (blue) and $D_r = 1.00$ (orange). (i) Schematic definition of the geometric factor $\chi_{\alpha\alpha'}$ for calculating the pairwise order parameter S. If

the center of the α'-th cell, adjacent to the α-th cell, lies within the sector ranging from $\pi/3$

forward or $\pi/3$ backward relative to the polarity vector, $\boldsymbol{q}_\alpha$, of the α-th cell, then $\chi_{\alpha\alpha'} = 1$;

otherwise, $\chi_{\alpha\alpha'} = -1$. (j) Phase diagram of the pairwise order, S, as a function of J and D_r.

FIG. 3. Macroscopic behavior of motile-type cells and domains. (a) Mean-square displacement

(MSD) of motile-type cells as a function of the time shift, τ. (b) Time evolution of the domain

length, R, with $J = 6.0$ for $D_r = 0.25$ (blue) and $D_r = 1.0$ (orange). The line and band widths in

the plot indicate the statistical average values and standard deviations, respectively. (c) Growth

exponent, ν (black point), and separation degree, μ (blue line), as a function of noise intensity D_r

with $J = 2.0$ (left), $J = 6.0$ (middle), and $J = 10.0$ (right). The green dashed lines and red dash-

dotted lines indicate $\nu = 1/3$ and $\nu = 1/4$, respectively. (d) Spatial correlation function of cell

types, $C(r, t)$, as a function of the distance between cells, r, normalized by the domain length, R,

with $(J, D_r) = (6.0, 0.25)$. (e) Average speed of motile-type cells (orange line) and separation

degree, μ (blue line), as a function of time, t, with $(J, D_r) = (6.0, 0.25)$ (left) and $(J, D_r) =$

$(2.0, 3.0)$ (right).

FIG. 4. Translational motion of a pair of motile-type cells. (a) Snapshots of a motile-type cell pair in
translational motion (also shown in Movie S3). Motile-type and nonmotile-type cells are colored
purple and gray, respectively. Arrows indicate the polarity vectors of motile-type cells, denoted by
q_A and q_B. Edge colors indicate the intensity of edge tension. (b) Autocorrelation of the mean
velocity of the motile-type cells as a function of time shift, τ, with $D_r = 0.01$ (blue) and $D_r = 1.00$ (orange). (c) The statistical average of the interfacial tension between the motile-type cells as a
function of polarity alignment $q_A \cdot q_B$ (solid line) and the distribution of the polarity alignment
(dashed line), with $D_r = 0.01$ (blue) and $D_r = 1.00$ (orange).

FIG. 5. Stability of the circulation configuration in a two-rail model. (a) Setup of the two-rail model.
Each dashed line represents the cell array corresponding to cell circulation. The i-th array has a flow
velocity, v_i (black arrow), along the array and polarity, q_i (magenta arrow). The interaction
between cell arrays is characterized by an interaction index, β. (b) The vector field of the polarities,
q_1 and q_2, with $\beta = -1$ (left), $\beta = 0$ (middle), and $\beta = 1$ (right). The color indicates the value
of $v_1 v_2$. (c) Eigenvalues of the Jacobian matrix for fixed points corresponding to the steady parallel
arrangement (red) and the steady antiparallel arrangement (blue) as a function of β.

FIG. 6. Antiparallel circulation and translational motion of *Dictyostelium discoideum* cells. (a)

Experimental setup for the observation of *D. discoideum*. (b) A snapshot of a *D. discoideum*

aggregate in a 2D chamber. (c) Time evolution of the aggregate with circulation directions of cells

around the center (also shown in Movie S4). Cells exhibiting clockwise (CW) circulation are colored

red, and those with counterclockwise (CCW) circulation are colored blue. The yellow crosses mark

the center of circulation. (d) Distribution of cell migration directions (CW: red; CCW: blue) as a

function of cell velocity angle. (e) Average cell speed as a function of cell velocity angle. (f)

Distribution of the circulation persistence time, τ_α, for the raw data (blue) and randomized data

generated via bootstrap sampling (orange). (g) Snapshots of a pair of *D. discoideum* cells in

translational motion (also shown in Movie S5). (h) Distribution of the normalized velocity

alignment, $\widehat{\boldsymbol{v}}_A \cdot \widehat{\boldsymbol{v}}_B$, of a pair of *D. discoideum* cells.

FIG. 7. Jamming transition of the system. (a) Phase diagram of the effective diffusion coefficient,

D_{eff}, as a function of the target shape index, p_0, and strength of polarized tension, γ^{p}, with

$(J, D_{\mathrm{r}}) = (0.0, 1.0)$. (b) Phase diagram of the separation degree, μ, as a function of p_0 and J with

$\gamma^{\mathrm{p}} = 0.2$ and $D_{\mathrm{r}} = 1.0$. The white dotted line indicates the target shape index threshold, which

determines whether the tissue is in a solid-like state or fluid-like state. The yellow dotted line

indicates the alignment rate at which the value is equal to the noise.

FIG. 8. Dependence of phase separation on initial conditions. (a, b) Time evolution of the separation

degree, μ, with $(J, D_r) = (5.0, 2.5)$ (a) and $(J, D_r) = (1.0, 3.0)$ (b). In panels (a) and (b), the blue

and orange lines indicate the case with a homogeneous initial state and the case with a separated

initial state, respectively.