Natural variation in pigmentation caused by *cis*-regulatory changes affecting expression of the *ebony* gene in *Drosophila elegans* species subgroup

Anggun Sausan Firdaus[1], Atessha Negi[1], Kai-An You[1], Erick X. Bayala[2], Patricia J. Wittkopp[2,3], Ben-Yang Liao[4], and Shu-Dan Yeh[1]

[1]Department of Life Sciences, National Central University, Taoyuan, 32001, Taiwan.

[2]Department of Molecular, Cellular, and Developmental Biology, University of Michigan, Ann Arbor, MI, USA.

[3]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA.

[4]Institute of Population Health Sciences, National Health Research Institutes, Miaoli, Taiwan.

Correspondence: Shu-Dan Yeh[1]

e-mail: shudanyeh@ncu.edu.tw

Running Title: Pigmentation dimorphism and *ebony* regulation

Abstract

Repetitive evolution has offered valuable insights into how similar traits can arise across different

populations and species. The dimorphic body color morphs (brown and black) of geographically

isolated *Drosophila elegans*, along with its closely related species *D. gunungcola*, which exhibits black

morph, raise intriguing questions about whether these traits have emerged through independent

evolution, introgression, or shared ancestral variants. To investigate the genetic and evolutionary origins

of these differences, we perform quantitative trait locus (QTL) mapping on recombinant hybrids

generated from reciprocal crosses between brown and black morphs *D. elegans*. QTL mapping reveals

significant contributions from the X chromosome and Muller element E to intraspecific pigmentation

polymorphism, with *ebony* identified as one candidate gene. Analysis of *ebony* expression, including

allele-specific expression in F_1 hybrids, and sequence variation suggests that the intra- and interspecific

pigmentation differences within *D. elegans* species subgroup may have evolved independently from the

lineage-specific region upstream of *ebony*. By identifying a shared candidate regulatory region

associated with both intra- and interspecific variation, our findings suggest that recurrent use of the

same genetic region may contribute to the emergence of similar traits across populations and species.

Keywords: *cis*-regulatory divergence, *Drosophila elegans*, *ebony*, parallel evolution, pigmentation,

QTL mapping.

Introduction

Similar phenotypes within and between species often arise in response to environmental

challenges, indicating that ecological conditions often favor similar adaptive solutions. In many cases,

such similarities have been linked to divergence in the same genes or pathways, suggesting that certain

genetic architectures may be more prone to modification under selection (Martin and Orgogozo 2013;

Stern 2013; Massey and Wittkopp 2016; Cerca 2023; Bohutinska and Peichel 2024). This raises the

intriguing question of how similar phenotypes evolve across populations and species.

Among the phenotypes studied to address this question, pigmentation offers an ideal platform as

it is one of the most frequently studied morphological characteristics due to its visibility and ecological

relevance that often serve as targets for both natural and sexual selection across taxa (Cuthill et al. 2017;

Courtier-Orgogozo et al. 2020). In many *Drosophila* species, pigmentation polymorphisms often appear

as a continuous gradient (Brisson et al. 2005; Pool and Aquadro 2007; Massey and Wittkopp 2016).

These gradients are shaped by ecological pressures and commonly occur across populations living in

regions that are geographically proximate (Pool and Aquadro 2007; Parkash et al. 2008; Telonis-Scott

et al. 2011). While pigmentation in many *Drosophila* species follows a continuous gradient, there are

cases where geographically isolated populations exhibit distinct body color morphs.

The *Drosophila elegans* species subgroup is one particularly notable example of striking

dimorphic pigmentation between geographically isolated populations. *Drosophila elegans* populations

exhibit two distinct color morphs: brown morph distributed in regions closer to the equator, including

Indonesia, the Philippines, and Hong Kong, while a black morph distributed in higher-latitude regions

like Taiwan and Okinawa Island (Fig. 1A, B). The subgroup also consists of its sister species, *D.*

gunungcola (Fig. 1C). Notably, *D. gunungcola* is endemic to the southern hemisphere of Sumatra and

throughout the Java Islands (Hirai and Kimura, 1997). Despite having partial sympatry with the brown

morphs of *D. elegans* at higher elevation, *D. gunungcola* consistently exhibits black morph. This

intriguing pattern of pigmentation and distribution raises compelling questions about whether the

phenotypic similarities between the black morph of *D. elegans* and *D. gunungcola* are driven by shared

genetic variants or independent mutations.

Previous studies have shown that a small set of genes, including *yellow*, *tan*, and *ebony*, are

repeatedly implicated in both polymorphic and divergent pigmentation in *Drosophila* (Martin and

Orgogozo 2013; Johnson et al. 2015; Massey and Wittkopp 2016). For instance, pigmentation

polymorphism within *D. americana* and pigmentation divergence between *D. americana* and *D.*

novamexicana are both influenced by the genes *ebony* and *tan*, which likely originated from a shared

ancestral population (Wittkopp et al. 2009; Lamb et al. 2020). In the *D. elegans* species subgroup,

genetic loci linked to the *yellow* and *ebony* genes have been implicated in the pigmentation divergence

between the brown morph of *D. elegans* and *D. gunungcola* (Massey et al. 2021). Therefore, we

hypothesize that both intra- and interspecific pigmentation variations in the *D. elegans* are driven by

the same genes. However, whether the intraspecific dimorphism in *D. elegans* and the interspecific

divergence involving *D. gunungcola* share common genetic variants remains unknown.

58
59

Figure 1. (A) Brown morph *D. elegans*. **(B)** Black morph *D. elegans*. **(C)** *D. gunungcola*. **(D)** QTL map for female (red) and male (blue) thorax pigmentation intensity. The y-axis represents LOD score (the red and blue dashed line indicates the $p < 0.01$ threshold for female and male recombinants, respectively). The x-axis represents the physical map of Muller elements corresponding to *D. melanogaster* chromosome (X = X; B = 2L; C = 2R; D = 3L; E = 3R; and F = 4). Black tick marks on the x-axis represent individual markers along the chromosomes. **(E)** QTL mapping plots for the female abdominal pattern. **(F)** 16 recombinant females containing breakpoints near the thorax pigmentation QTL peaks were aligned to narrow down the candidate loci. Red, blue, and white bars represent homozygous genotype for brown allele, homozygous genotype for black allele, and heterozygous genotype, respectively.

In this study, we explored the genetic basis of dimorphic pigmentation within the *D. elegans*

species subgroup, combining QTL mapping, gene expression, and comparative sequence analyses. Our

findings identified that differential expression of the pigmentation gene *ebony* likely contributes to both

intra- and interspecific variation, potentially through variation in regulatory regions. In particular, a

divergent upstream sequence of *ebony* emerged as a candidate region for this dimorphic pigmentation,

although the precise causal variants remain unresolved. Together, these findings suggest the potential

role of regulatory divergence in shaping similar pigmentation phenotypes within and between species,

highlighting the *D. elegans* species subgroup as a valuable model for studying the evolution of

dimorphic traits across evolutionary scales.

Results

The X chromosome and Muller element E contribute to intraspecific pigmentation dimorphism

To examine the inheritance of intraspecific pigmentation variation, we performed reciprocal

crosses between two strains of *D. elegans*: HK (brown morph) and TPCC (black morph). The F1 flies

from both crosses exhibited intermediate body color (Supplemental Fig. S1A). We then generated

sibling crosses from the F1 hybrids for eight generations. We focused our analysis on the thorax

pigmentation polymorphism in both male and female recombinants. The resulting recombinants

displayed a spectrum of thorax pigmentation intensity.

To investigate the candidate loci linked to the thorax pigmentation variation, we performed QTL

mapping on 235 females and 212 males from F2 to F8 recombinants produced through reciprocal

crosses using individuals with the body color at two ends of the phenotypic distribution (Supplemental

Fig. S1B). There were no significant differences in thorax pigmentation intensity between male and

female recombinants within the brown morph or within the black morph, indicating that this trait is not

sexually dimorphic (Supplemental Fig. S2).

Our QTL mapping analyses revealed an X-linked QTL that was highly significant in the male

mapping dataset, but only marginally significant in the female mapping dataset (Fig. 1D; Table 1). This

difference in significance presumably results from differences in mapping power for X-linked loci in

the two sexes: males are hemizygous for X-linked genes, carrying only the black or brown morph allele,

whereas females include individuals that are homozygous for the black morph allele, homozygous for

the brown morph allele, or heterozygous for both. The entire Muller element E (corresponding to the

3R chromosome in *D. melanogaster*) exhibited strong associations with multiple peaks in both male

and female mapping datasets (Fig. 1D; Table 1). The broad QTL signals across Muller element E may

be attributed to: 1) multiple QTLs of large effect; 2) low resolution from the relatively small number of

recombinant hybrids; or 3) a low recombination rate on Muller element E. The latter scenario is less

likely as crossover frequencies were consistent among chromosomes (Supplemental Fig. S3).

Table 1. QTLs detected for thorax pigmentation intensity and abdominal banding pattern.

Traits	Sex	Chromosome	QTL interval (bp)[a]	QTL peak (bp)	LOD
Thorax pigmentation intensity	Female	X	7,445,655-18,244,170	15,476,093	3.71
Thorax pigmentation intensity	Male	X	7,955,254-14,476,391	12,486,125	12.72
Thorax pigmentation intensity	Female	E	8,622,548-14,156,589	14,038,360	35.38
Thorax pigmentation intensity	Male	E	8,422,980-8,972,186	8,487,591	29.01
A4-tergite dark band width	Female	X	7,955,254-15,699,310	14,725,205	6.77
A4-tergite dark band width	Female	E	600,044-2,080,945	1,069,099	47.11

109 [a]1.8-LOD drop support interval
110

We also observed variations in female abdominal banding patterns (Fig. 1E). To compare the

genetic basis of thorax pigmentation intensity and abdominal banding pattern variation, we measured

the width of dark bands on the 4th tergite of 235 female F2 to F8 recombinants. We detected a significant

positive correlation between thorax pigmentation intensity and the width of dark band on the A4

(Spearman's correlation: $R=0.76$, $P < 0.001$) (Supplemental Fig. S4), suggesting that darker thoraxes

are associated with wider abdominal dark bands. Our QTL mapping analyses identified a highly

significant QTL on Muller element E associated with variation in the abdominal banding pattern. A

QTL with lower significance on the X chromosome also showed genetic association with female

abdominal banding pattern (Fig. 1E; Table 1). This was unexpected, as thorax pigmentation intensity

and spatial banding pattern in the abdomen are typically regulated by different genetic mechanisms

(Rebeiz et al. 2009; Takahashi 2013; Massey and Wittkopp 2016). Our results suggest these traits may

share a genetic basis, potentially controlled by overlapping or closely linked regulatory elements.

Candidate loci within QTLs and genetic interaction between X chromosome and Muller element E

Muller element E contains multiple genes known to be associated with pigmentation in *Drosophila*,

including *ebony*, *Abdominal-B (abd-b)*, *Dop1R1*, *Dop1R2*, and *stripe (sr)*. The QTL signal spans a large

portion of Muller element E, suggesting that one or more known or previously uncharacterized genes

may contribute to pigmentation differences. To refine the candidate region on Muller element E, we

examined recombinant genotypes and phenotypes to identify the recombination breakpoints flanking

the thorax QTLs. Genotype data for each recombinant individual are provided in the Supplemental File.

This allowed us to narrow down only one candidate region to ~38 kb in females, containing 10 genes

in the region acting like a genetic switch controlling brown or black thorax pigmentation (Fig. 1F). In

males, we refined the region to ~891 kb, which includes the region identified in females (Supplemental

Fig. S5). Hybrids homozygous for the black loci in this region exhibited black thorax pigmentation,

whereas those with heterozygous or homozygous brown loci displayed brown thorax (Fig. 1F; Fig. 2A).

One gene in this region is *ebony* (*e*) which encodes the enzyme N-ß-alanyl-dopamine synthetase

required for the formation of yellow sclerotin (Wittkopp et al. 2002). *ebony* has been previously reported

to contribute in pigmentation variation both within and between *Drosophila* species (Wittkopp et al.

2009; Johnson et al. 2015; Massey and Wittkopp 2016), making it a strong candidate gene responsible

for pigmentation variation in *D. elegans*.

We also attempted to refine the candidate region on the X chromosome associated with thorax

pigmentation. However, we were unable to identify a specific region on the X chromosome, as

individuals carrying homozygous black, heterozygous, or homozygous brown loci across the entire X

chromosome could exhibit either brown or black thorax pigmentation (Supplemental Fig. S6). To test

for possible genetic interactions, we further performed a two-way ANOVA between the QTL identified

on the X chromosome (hereafter referred to as QTL-X) and the *ebony* locus on Muller element E. Our

analysis revealed tissue- and sex-specific genetic interactions between *ebony* and QTL-X, using the

marker with highest LOD score (Fig. 2B; Two-way ANOVA; Male thorax pigmentation intensity: $F_{2,206}$

$= 2.79$, $P = 0.064$; Female thorax pigmentation intensity: $F_{4,226} = 2.63$, $P = 0.035$; Female abdominal

dark band width: $F_{4,226} = 2.21$; $P = 0.069$). We observed that brown *ebony* allele was dominant to black

ebony allele regardless of QTL-X allele for thorax pigmentation. While male recombinants showed a

marginally significant interaction, female thorax pigmentation showed weak epistatic interaction. These

results suggest that while *ebony* is the primary driver of pigmentation, QTL-X likely contributes through

additive effects and weak epistatic interactions in a tissue- and sex-specific manner.

E chr: 1,582,436

Figure 2. (A) Effect plots of the genotype against QTL peak (marker: 1,582,436 near the *ebony* locus) for male thorax pigmentation intensity, female thorax pigmentation intensity, and female abdominal pattern. Dots represent individual replicates. **(B)** Genetic interactions between QTL on the X chromosome (QTL-X) and *ebony* on Muller element E. Error bars represent 95% confidence intervals for the phenotypic values of each genotype. Sample sizes (N) for flies with each genotype (blue for homozygous *D. elegans* black allele, purple for heterozygous allele, and red for homozygous *D. elegans* brown allele) are provided below each plot for the respective traits. Higher score means darker pigmentation intensity.

Variation in ebony expression is correlated with Drosophila elegans pigmentation differences

Given the pleiotropic role of *ebony* in *Drosophila* (Borycz et al. 2002; Takahashi 2013; Massey et al. 2019), we hypothesized that genetic variation in *ebony* might regulate its expression levels rather than altering its protein function, contributing to both intra- and interspecific pigmentation differences

in the *D. elegans* species subgroup. To test this, we performed RT-qPCR on brown morph *D. elegans*, black morph *D. elegans*, and black morph *D. gunungcola*. The inclusion of *D. gunungcola* allowed for a comparison between closely related species with the same body color morphs. We collected male and female samples within 0 to 2 hours post-eclosion, and focused on the head, thorax epidermis, and abdominal epidermis to capture tissue-specific expression patterns. We observed consistent *ebony* expression patterns across tissues and sexes. Generally, the brown morph showed significantly higher *ebony* expression than the black morphs (Fig. 3A). Notably, the thorax epidermis showed especially strong differences in *ebony* expression between brown and black morphs in both male and female datasets.

To test whether these *ebony* expression differences stem from *cis*-regulatory divergence, we analyzed expression of the *ebony*^{HK} (the brown morph allele) and *ebony*^{TP} (the black morph allele) in F_1 hybrids after eclosion. The *ebony*^{HK} consistently showed higher expression than the *ebony*^{TP} across all tissues (one-sample *t*-test, $P < 0.05$, Supplemental Table S1), indicating that *cis*-regulatory divergence does indeed contribute to the difference in expression level seen between these two morphs.

We further investigated the spatial distribution of *ebony* and *yellow*, a pigmentation gene within QTL-X peak, mRNA expression of black and brown morphs of *D. elegans* using the Hybridization Chain Reaction (HCR) method. The brown morph displayed stronger *ebony* expression than the black morph at 65-75 hours after puparium formation (hAPF) (Supplemental Fig. S7) and this difference become more pronounced at later developmental stages (75-85 hAPF, 85-95 hAPF, and post-eclosion), aligning with RT-qPCR findings (Fig. 3B, Supplemental Fig. S7). In contrast, *yellow* expression was detected in the epidermis in both morphs, though the signal appeared slightly stronger before eclosion in the black morph and after eclosion in the brown morph. These findings correlate with the effects observed for QTL-X and QTL-E, with larger differences in *ebony* expression than *yellow* expressing between the black and brown morphs. Together, our results from three kinds of gene expression analyses support that *cis*-regulatory differences in *ebony* underlie intraspecific pigmentation dimorphism in *D. elegans*.

A.

B.

Figure 3. (A) Relative *ebony* gene expression among different body morphs. Newly emerged flies were dissected into three parts: head, thorax epidermis, and dorsal abdominal epidermis. One-way ANOVA followed by *post hoc* Tukey HSD were used to assess significant differences among group. Significant *P value*: '*' ≤ 0.05; '**' ≤ 0.01; '***' ≤ 0.001. **(B)** Spatiotemporal expression of *ebony* and *yellow* in black and brown morphs of *D. elegans*. The epidermal tissues from brown morph (*D. elegans* HK) and black morph (*D. elegans* TP) were stained using the Hybridization Chain Reaction (HCR) at two developmental stages: (i; ii) 85–95 hours after puparium formation (hAPF); and (iii; iv) 0–2 hours post-eclosion. For each sample, three panels are shown: DNA (DAPI, blue), *ebony* mRNA (magenta), and *yellow* mRNA (green). All samples were processed and imaged under the same conditions; variation in DAPI signal may reflect staining efficiency or nuclear accessibility.

5' intergenic region of ebony displayed high nucleotide variation that correlates with dimorphic

pigmentation differences at both intra- and interspecific levels

To explore natural genetic variation within the *ebony* locus that may be associated with

pigmentation dimorphism, we sequenced a ~16 kb *ebony* region (including the 5' and 3' intergenic,

exons, and introns) from wild-caught *D. elegans* and *D. gunungcola* from Indonesia and Taiwan

(Supplemental Table S2). We first investigated protein sequence and found no fixed amino acid

differences between brown and black *D. elegans* morphs. In contrast, we identified seven amino acid

residues diverged between two species (Supplemental Table S3). Five of these residues are located

within the adenylation domain of the Ebony protein, hypothetically activate ß-alanine to

aminoacyladenylate for further conjugating on to dopamine (Richardt et al. 2003; Izore et al. 2019).

The functional significance of these five residues, however, have not been experimentally tested. The

remaining two sites are located in the C-terminal region (Izore et al. 2019). Based on the *D.*

melanogaster Ebony structure, none of these residues is at critical sites for substrate binding, orientation,

or catalysis, suggesting they are unlikely to influence Ebony's enzymatic function.

Instead of protein sequence differences, cis-regulatory changes in the *ebony* locus may result in

pigmentation differences. To explores lineage-specific versus shared genetic patterns linked to

pigmentation, we performed a sliding-window analysis of net divergence (*Da*) using the ~16 kb *ebony*

across three pairwise comparisons- between brown and black *D. elegans* as well as *D. gunungcola*.

Between-species comparisons exhibit higher nucleotide divergence than within-species comparisons

generally as expected, except in the region of 3635 bp to 4035 bp from the end of upstream gene coding

sequence. This 400-bp sequence in the 5' intergenic region of *ebony* shows high divergence between

brown and black morphs for both within and between species comparisons, but not between black

morph *D. elegans* vs. *D. gunungcola* (Fig. 4; Supplemental Fig. S8). The relatively low divergence

between black morph *D. elegans* and *D. gunungcola* implies that natural selection may have acted on

this region in the black morph to reflect similar evolutionary pressures in both species or shared

ancestral genetic variants.

Figure 4. Sliding window analysis of net divergence levels (DA) between *D. gunungcola* and *D.*
elegans. Three pairwise comparisons were performed: brown *D. elegans* vs. black *D. elegans* (blue),
brown *D. elegans* vs. *D. gunungcola* (red), and black *D. elegans* vs. *D. gunungcola* (black). Grey, black,
and red (400 bp) horizontal bars at the top of the figure indicate the transcribed regions of the *ebony*
gene, nearby gene, and *ebony* region with high divergence levels. Window size was set to 100 bp with
step size 25 bp.

Phylogenetic patterns of dimorphic pigmentation in D. elegans species subgroup

To investigate the evolutionary histories underlying dimorphic pigmentation in *D. elegans* species

subgroup, we reconstructed phylogenetic trees of wild-caught samples using three genomic regions: the

400-bp candidate region in the 5' intergenic region of *ebony*, the coding sequence (CDS) of *ebony*, and

mitochondrial DNA (*ND1*) (Supplemental Table S2). Examining these regions separately is important

as each may reflect distinct evolutionary trajectories and selective pressures.

The tree based on *ebony* coding sequences formed two major clades: *D. gunungcola* clade and *D.*

elegans clade, within which the black morph is nested in the brown morph clade (Fig. 5A). In contrast,

the tree based on the 400-bp 5' intergenic region of *ebony* formed three major clades: *D. gunungcola*

clade, brown morph *D. elegans* clade, and black morph *D. elegans* clade (Fig. 5B). On the other hand,

the mitochondrial DNA tree did not separate black and brown *D. elegans* into distinct clades (Fig. 5C).

While the phylogenetic tree based on the 400-bp candidate region revealed distinct clades for the brown

and black morphs of *D. elegans*, this pattern contradicts our hypothesis that the black morphs of *D.*

elegans and *D. gunungcola* share ancestral genetic variants for body melanization.

(A)   (B)

264   (C)

Figure 5. Phylogenetic trees reconstructed using Bayesian Inference based on (A) the *ebony* coding
sequence, (B) the 5' intergenic region of *ebony*, and (C) mitochondrial DNA (*ND1*). Numbers on the
branches represent posterior probabilities from Bayesian analysis, indicating statistical support for each
node.

If dimorphic pigmentation is resulted from shared genetic variants, we would expect a

substantially higher number of shared polymorphisms exclusive to black *D. elegans* and *D. gunungcola*.

We analyzed the sequence polymorphisms within the 400-bp candidate region (Supplemental Table S4)

and identified 18 diverged polymorphisms (16 nucleotide substitutions and 2 indels) between species,

12 shared nucleotide substitutions exclusive to black morph *D. elegans* and *D. gunungcola,* and 11

shared polymorphisms (10 nucleotide substitutions and 1 indel) exclusive to brown morph *D. elegans*

and *D. gunungcola*. It appears to be more diverged polymorphisms between species and similar shared

polymorphisms between body color morphs to species comparisons. Therefore, both phylogenetic and

polymorphism analyses do not support the evolutionary scenario of shared ancestral genetic variants.

Discussion

ebony: key driver of intraspecific and interspecific dimorphic pigmentation differences

Research on *Drosophila* pigmentation has revealed substantial variation in thorax, abdominal,

and wing pigmentation shaped by selection pressures, geographic variables, and genetic factors.

Notably, the sex chromosome and Muller element E (3R chromosome) harbor three key pigmentation

genes, *yellow*, *tan*, and *ebony*, that play a central role in pigmentation divergence across *Drosophila*

populations and species. Prior studies across *Drosophila* species have shown *ebony*'s key role in

pigmentation spectrum (Takahashi et al. 2007; Rebeiz et al. 2009; Wittkopp et al. 2009; Massey and

Wittkopp 2016). Our findings, along with those of Massey et al. (2021), examined divergence between

brown *D. elegans* and black *D. gunungcola*, highlighting *ebony* as a potential genetic hotspot underlying

dimorphic pigmentation traits within and between species.

In *D. elegans*, dimorphism in thorax pigmentation appears to result primarily from *cis*-regulatory

differences in *ebony*, rather than coding sequence mutations. The crucial role of *cis*-regulatory changes

in the *ebony* locus has also been well-documented in several *Drosophila* species (Takahashi et al. 2007;

Takahashi and Takano-Shimizu 2011; Telonis-Scott et al. 2011; Miyagi et al. 2015). One of the most

intriguing aspects of *ebony* is its pleiotropic nature, affecting not only pigmentation but also cuticular

hydrocarbon, vision, and behavioral rhythmicity (Borycz et al. 2002; Takahashi 2013; Massey et al.

2019). Evolution of the *cis*-regulatory region likely provides a more adaptable route in pleiotropic loci.

In this case, the long 5' intergenic region and intron 1 in the *ebony* locus offer opportunities to modulate

the control for gene expression pattern and adaptive responses among multiple biological functions,

thus making *ebony* an ideal target for selection.

In fact, the repetitive evolution has also been observed across various species, where the same

sets of genes and pathways are repeatedly utilized within and between organisms. This process can lead

to the emergence of novel traits or the development of similar traits across distinct species or

populations over time. One of the well-known examples in the stickleback fish, where recurrent

mutations of the *Eda* gene have independently led to the reduction of armor plates in multiple freshwater

populations (Colosimo et al. 2005; Archambeault et al. 2020). This demonstrating how the same genetic

changes can repeatedly drive similar adaptations in certain environments.

Our results suggest that repetitive evolution in dimorphic pigmentation differences within and

between species may be associated with regulatory genetic changes across lineages. Similar phenotypes

might evolve repeatedly in responding to parallel selective pressure in distinct populations or species.

Further studies are required to explore how other genetic and regulatory pathways interact with *ebony*,

which may uncover more profound insights into the mechanisms underlying such cases. Such research

will broaden our understanding of evolutionary biology within *Drosophila* and shed light on the

predictability of evolutionary changes across taxa.

Repetitive evolution of pigmentation linked to independent regulatory changes

Repetitive evolution can arise through shared genetic polymorphism (collateral evolution),

introgression, or independent genetic variation (parallel or convergent evolution), depending on

whether pre-existing alleles or new mutations are involved in the process (Maeso et al. 2012; Stern

2013; Cerca 2023). The phylogenetic trees constructed from the 400-bp sequence in 5' intergenic region

(5'IR), coding sequence (CDS), and mitochondrial DNA (mtDNA), offer new insights into the

evolutionary trajectories of dimorphic pigmentation within *D. elegans* species subgroup.

We use the mtDNA tree as a baseline to infer species-level divergence. The lack of distinct clades

for brown and black morphs implies that, there no clear lineage sorting for pigmentation traits at the

species level. However, the *ebony* CDS tree shows that black morphs nested within the brown morph

clade, implying that black morphs might be a more recent adaptation or a variation that evolved from

the brown morph *D. elegans* population. These contrasting patterns between mtDNA and *ebony* CDS

may suggest that the traits are diversifying under selection pressures, while the overall population

structure remains interconnected.

In contrast, the 5'IR tree shows two distinct clades of black and brown *D. elegans*, with *D.*

gunungcola as a sister species. This phylogenetic pattern corresponds to ASE analysis, which indicates

the link between *cis*-regulatory changes and the dimorphic pigmentation differences at both intra- and

interspecific levels. Although this 400-bp region is relatively conserved between *D. elegans* and *D.*

gunungcola, low sequence similarity of corresponding regions exists in other *Drosophila* species,

indicating potential lineage specificity. Notably, we observed a relatively even distribution and number

of shared genetic variants between black *D. elegans*- *D. gunungcola* contrast and brown *D. elegans*- *D.*

gunungcola contrast. This balanced pattern does not support a scenario of shared ancestral variant or

introgression, but instead suggests the involvement of independent evolutionary changes for the

dimorphic pigmentation in the *D. elegans* species subgroup.

The pleiotropic effect of *ebony* in cuticular hydrocarbon (CHC) composition may also

synergistically facilitate the evolutionary changes of *ebony* expression level in the epidermis. The CHC

composition plays a significant role in desiccation resistance (Wang et al. 2022) and mate recognition

(Ferveur 1997; Chung and Carroll 2015). The genetic ablation of *ebony* in *D. melanogaster, D.*

americana, and *D. novamexicana* results in qualitative changes of CHC profiles with increased

abundance of long chain CHCs (Massey et al. 2019; Lamb et al. 2020). In *D.* elegans, females of black

and brown morphs exhibit significant differences in CHC profiles, potentially influencing male mating

preference (Ishii et al. 2001). Thus, the interaction among desiccation resistance, mate recognition, and

body-color related adaptation may maintain a strong selection imposing lower expression level of *ebony*

in the epidermis of black morph *D. elegans* and *D. gunungcola.* Still, whether the same or overlapping

regulatory elements influencing the differences of body pigmentation and CHCs needs further

investigations.

Our results provide a valuable baseline for further studies into the genetic and ecological

underpinnings of pigmentation divergence. However, it remains unclear whether the divergent upstream

region of *ebony* contributes specifically to pigmentation or also influences other traits, such as CHC

composition. Moreover, the emergence of similar pigmentation in geographically distinct populations

and species suggests the potential for repetitive evolutionary changes, although the underlying

environmental or selective pressures require further investigations. Future research should aim to

disentangle these complex relationships and identify the regulatory elements and environmental drivers

that shape pigmentation and related traits in this system.

Conclusion

Our findings suggest that pigmentation differences at both intra- and interspecific levels may

have evolved independently through modifications in a lineage-specific sequence within the 5'

intergenic region of *ebony*. This highlights the predictability of adaptive evolution, where similar

phenotypes repeatedly arise via modifications in specific regulatory regions, potentially driven by

parallel selective pressures. Future research, including functional validation of this candidate region

will be essential to confirm the regulatory role and clarify its contribution to dimorphic pigmentation

differences within *D. elegans* species subgroup.

Materials and Methods

Fly strains

Drosophila elegans HK (brown morph, originated from Hong Kong, China) and *D. gunungcola*

SK (originated from Sumatra Island, Indonesia) strains were provided by John R. True (Stony Brook

University) who maintained the stocks established by M. T. Kimura. *Drosophila elegans* TPCC,

TPCC12, and TPCC13 strains (black morph) were established from several females collected in New

Taipei City, Taiwan. We maintained all strains on a standard cornmeal medium (food recipe is provided

by request) under constant light at 23ºC in the Department of Life Sciences, National Central University,

Taiwan.

The wild-caught flies used for sequencing the *ebony* and mitochondrial DNA were collected from

Sumatra and Java Island in 2018 and Taiwan in 2020-2021. The flies were preserved in 75% ethanol

upon field collection. Fly collection in Indonesia was carried out under a material transfer agreement

with Indonesian collaborators from the Brawijaya University.

Generating intraspecific recombinants for QTL mapping analysis

Virgin males and females from *D. elegans* HK (brown morph) and TPCC (black morph) strains were categorized 4-6 hours after eclosion and aged for 4-7 days to reach sexual maturity. We started the reciprocal crosses by setting up ten single-pair vials per direction, each containing a male from one strain and a virgin female from another. We allowed each parental pair to copulate and oviposit for 10-14 days, then collected the parental flies for genotyping. We transferred newly eclosed F_1 recombinants into new vials and allowed them to interbreed for 7-10 days to produce F_2 recombinants. We maintained the hybrids of each cross pair separately from those of other cross pairs. To promote further recombination, we continued random interbreeding using the same protocol for eight non-overlapping generations.

To map the genetic basis of thorax pigmentation, we phenotyped both male and female recombinants from the 2nd to 8th generations of each reciprocal cross. We visually inspected and classified flies at 7-10 days old. In the 2nd generation, we collected approximately 30 recombinants per color and sex. From the 3rd to 8th generations, we selected the darkest recombinants resembling the TPCC strain and the lightest ones resembling the HK strain (N=12-17 per color and sex).

Imaging and scoring pigmentation traits

We photographed individual flies to score the pigmentation traits before genotyping. To reduce light reflection interference, we immersed and positioned each fly in 70% ethanol. We captured fly images in RGB mode using a digital camera (MSHOT MS60) connected to a stereoscopic microscope (Leica MZ 125), applying identical optical parameters for all samples. We quantified thorax pigmentation intensity using ImageJ software (version 1.52a; Wayne Rasband, National Institutes of Health, USA; http://imagej.nih.gov/ij). We manually selected six regions of interest (ROIs), modified from (Miyagi et al. 2015), including the thoracic trident, thoracic scutellum, and areas outside the trident. For each ROI, we recorded the mean gray value. To align scores with a scale ranging from 0 (white) to 255 (black), we substracted each score from 255 (Gibert et al. 2016). We averaged the six scores to obtain

the final pigmentation intensity score for each sample. Intensity values are reported in arbitrary units

(A.U.).

For the female abdominal banding pattern, we measured the width of the dark band and the total

width of the 4th tergite. We calculated the percentage of coverage coverage by dividing the dark band

width by the total tergite width and multiplying the result by 100.

Library preparation, genome assembly, and marker generation

A total of 469 flies, including 226 male and 243 female recombinants, were individually

genotyped using Multiplexed Shotgun Genotyping (MSG) (Andolfatto et al. 2011) with 473 barcoded

adaptors, following the protocol described in (Cande et al. 2012). We extracted genomic DNA and

prepared sequencing libraries using the methods explained in (Massey et al. 2020). For read mapping,

we used the *D. elegans HK* reference genome (D.elegans_HK_3.0; accession number:

GCA_011057505.1). This genome reference has been assembled to chromosomal-level which

facilitates more precise mapping to specific chromosomal locations. We sequenced the TPCC strain

genome at 50x coverage allowing a confident variant calling between HK and TPCC strain. In total, we

identified 3198 informative markers between the two strains for QTL mapping. The markers, thorax

and female abdominal phenotypes, and script used for QTL mapping are available on Supplemental

Code.

QTL mapping analysis and gene annotation

We conducted QTL mapping analysis using the R/qtl package (Broman et al. 2003) in R for

Windows version 4.4.0 (R Core Team 2024). We inferred ancestry states across chromosomal regions,

recombinant breakpoints, and genotyping status using a Hidden Markov Model developed for MSG

data, and imported the results into the R/qtl package using a custom script

(https://github.com/dstern/read_cross_msg) (Andolfatto et al. 2011). We performed genome scans

using Haley-Knott regression and the "scanone" function of a single QTL model (Haley and Knott

1992). Due to the low sequencing resolution, we excluded 13 males and 8 females from the analysis,

resulting in a final dataset of 212 males and 235 females from the 2nd to 8th generations. We determined

the LOD significance threshold using 1,000 permutation replicates at a 1% significance level ($\alpha = 0.01$).

We extracted the chromosomal regions spanning the QTL interval (~38 kb on Muller element E) for

thorax and abdominal pattern from the assembled *D. elegans* HK genome (accession no: CM021661.1).

Gene annotation followed the method described in Massey et al. (2020).

Analyzing epistasis interactions between QTL-X and ebony

To investigate potential epistatic interactions between QTL on the X chromosome and Muller

element E, we compared thorax pigmentation intensity (in both males and females) and female

abdominal banding patterns across different genotypic combinations. The QTL-X genotypes were

identified at positions with the highest LOD scores from our QTL mapping (Table 1; Male thorax

pigmentation intensity: 12,486,125 bp; Female thorax pigmentation intensity: 15,476,093 bp; Female

abdominal dark band width: 14,725,205 bp). For Muller element E, we focused on a region previously

narrowed down (position: 1,582,436 bp, located near *ebony*).

RNA Sample Preparation and RT-qPCR

We assessed the relative gene expression of *ebony* between brown (HK strain) and black (TPCC12

strain) morphs of *D. elegans* using RT-qPCR. We used a different black morph strain collected from

the same locality as TPCC because the TPCC strain originally used for QTL mapping perished several

months after experiments. We also included *D. gunungcola* (SK strain) to test whether its *ebony*

expression was similar to black morph *D. elegans*, as both share similar body pigmentation.

We collected male and female flies within 0 to 2 hours after eclosion. After removing the wings,

we stored the bodies in RNAlater™ Solution (Thermo Fisher Scientific) for dissection. We dissected

flies into three body parts: head, thorax epidermis (including scutellum and pleura), and the dorsal side

of abdominal epidermis. For each body part, sex, and strain, we prepared three biological replicates

(~40 flies per replicate). We extracted Total RNA using the Quick-RNA™ Miniprep Kit (Zymo

Research, USA) following the manufacturer's instructions and quantified the RNA using Nanodrop

spectrophotometer. First-strand cDNA was synthesized from ~200 ng of total RNA using the MMLV

Reverse Transcription Kit (PROTECH).

We performed RT-qPCR using a Bio-Rad CFX96 system in 20-µl reaction containing SYBR

Green Supermix (BioRad, USA) and 1 µl of cDNA per reaction. Primer sequences used for *ebony*,

Actin42A, and *TBP* are listed in Supplemental Table S5. Relative expression of *ebony* was calculated

using the Pfaffl method (Pfaffl 2001), with *Actin42A* and *TBP* as reference genes. Whole-body RNA

from *D. elegans* HK used as the control sample. Cq values and calculation steps used for expression

normalization are provided in Supplemental Table S6.

RNA and DNA extraction for allele-specific expression (ASE) analysis

To investigate cis-regulatory divergence of the *ebony* gene between pigmentation morphs in *Drosophila*

elegans, we performed allele-specific expression (ASE) analysis using a Sanger sequencing. We used

genomic DNA from the parental strains, HK strain (brown morph) and TPCC13 strain (black morph),

as reference samples for normalization and baseline correction. We pooled five adult females and

extracted DNA using the GeneAid gSYNC DNA Extraction Kit. We constructed a standard curve for

normalization by mixing parental gDNA samples at the brown:black ratios of 1:10, 1:5, 1:4, 1:2, 1:1.5,

1:1, 1.5:1, 2:1, 4:1, 5:1.

We crossed 5 HK females with 5 TPCC13 males to obtain F_1 recombinants. To prepare the total

RNA from F_1 recombinants, we dissected females within 0-2 hours post-eclosion to isolate the head,

thorax epidermis, and abdominal epidermis. For each tissue type, we prepared three biological replicates,

each consisting of 20 pooled individuals. We extracted total RNA using the Gene-Spin Total RNA

Purification Kit (PROTECH), following the manufacturer's protocol. We reverse-transcribed

approximately 200 ng of RNA per replicate into cDNA using the MMLV Reverse Transcription Kit

(PROTECH).

We PCR-amplified the target region (~800 bp of *ebony* exon 2) from gDNA and cDNA samples

using e_F9 and e_R8 primers (Supplemental Table S5). PCR conditions were: initial denaturation at

95ºC for 2 min, followed by 35 cycles of 95ºC for 30 s, 59ºC for 1 min, and 72ºC for 20 s, with a final

extension at 72ºC for 5 min. We sent the samples for Sanger sequencing using the forward primer

(PB_e_F9).

 Allele-Specific Expression Quantification and Normalization

To quantify allele-specific expression, we extracted peak height values from sequencing chromatograms (.ab1) using QSVanalyser (Carr et al. 2009) for each F_1 biological replicate. We used gDNA from each parental strain (HK accession number: PV939603; TPCC13 accession number: PV939604) to identify informative reference SNPs that distinguish the brown and black alleles of *D. elegans*. We focused on a single reliable marker SNP (located at 489 bp), which was fixed between the brown and black parental strains and showed consistent allelic imbalance across all tissues and biological replicates. We excluded other candidate SNPs from the analysis because they were not fixed (i.e., both alleles were present in at least one parental strain, as indicated by their chromatograms).

We corrected the final peak heights using baseline signal values following the method of Carr et al. (2009). For each replicate, we calculated the allelic expression ratio by dividing the final peak height of the brown allele by the black allele. We normalized these ratios using the standard curve generated from mixed parental gDNA samples with known brown:black ratios. This normalization accounted for technical variation in peak heights and allowed us to more accurately estimate the relative expression of the two alleles.

Hybridization Chain Reaction (HCR)

We performed Hybridization Chain Reaction (HCR) *in situ* staining to detect mRNA expression of *ebony* and *yellow* in *Drosophila elegans* black (TP) and brown (HK) morphs, following the protocol described in (Bayala et al. 2024). DAPI was included to visualize DNA. The same custom hairpins targeting *yellow* and *ebony* described in this prior work were also used in the current study. To assess spatiotemporal gene expression, we stained whole-body samples at two developmental stages: 85-95 hours after puparium formation (hAPF) and 0-2 hours post eclosion. We did not select for specific sexes, as the objective was to capture gene expression at carefully matched developmental stages. To further examine spatiotemporal expression in the abdominal epidermis during pupal development, we additionally stained dissected pupal abdomens at 65-75 hAPF and 75-85 hAPF. After staining, tissues were mounted on slides and imaged using a Leica SP5 laser scanning confocal. Red, green, and blue channel levels were adjusted uniformly across all image panels.

Collection of wild D. elegans in Indonesia and Taiwan

We collected samples in Indonesia for over three months in 2018 across Sumatra and Java. In Taiwan, we conducted sample collection in 2020 and 2021. Flies were collected using an aspirator and transferred into vials containing standard fly media or Instant Drosophila medium (agar, glucose, cornmeal, and yeast extract). Some flies were immediately preserved in ethanol 70%, and the rest of the caught flies were maintained and reared in the laboratory to establish isofemale or wild-isolated strains. Fly collection in Indonesia was carried out under a material transfer agreement with Indonesian collaborators from the University of Brawijaya.

Haplotype sequence analysis of ebony region

We extracted genomic DNA from single adult male flies using Gsync™ DNA Extraction Kit, following the manufacturer's instruction. We analyzed a total of 20 individuals: 9 *D. elegans* and 6 *D. gunungcola* from Indonesia (Java and Sumatra), and 5 *D. elegans* from Taiwan (Supplemental Table S2). We used six pairs of primers targeting partially overlapping ~3 kb amplicons to amplify ~16 kb of the *ebony* locus (Supplemental Table S5). The amplified region spanned the entire *ebony* gene (including 5' and 3' intergenic region, introns, and exons).

We assessed PCR product quality using gel electrophoresis and checked the quantity using a Qubit® fluorometer (ThermoFisher Scientific) with High Sensitivity (HS) dsDNA assay kits and a NanoDrop spectrophotometer. We sent the amplicons for sequencing using the PacBio Single-Molecule Real-Time (SMRT). We evaluated the quality of the amplicon-specific consensus sequences using SequelTools program/SequelQC v1.3.0. (Hufnagel et al. 2020). For each sample, six read clusters corresponding to the six primer pairs were phased into haplotypes. We mapped the reads to the *D. elegans* reference genome (D.elegans_HK_3.0; accession number: GCA_011057505.1) using the pbmm2 program (https://github.com/PacificBiosciences/pbbioconda). We visualized variants from each cluster in IGV, carefully selected for each respective amplicon by aligning it to reference using NUCmer from the MUMmer suite version 3.1, and stitched into a complete haplotype (Kurtz et al. 2004).

We aligned DNA sequences using ClustalW in MEGA-X v10.2.5 (Kumar et al. 2018). To

examine patterns of genetic divergence, we performed a 100-bp sliding-window analysis of net

nucleotide divergence (*Da*) across the ~16 kb region spanning *ebony* regions and its neighboring genes

using DnaSP v6.12.03 (Rozas et al. 2017). Sample information, including sample ID, accession

numbers, and localities is described in Supplemental Table S2.

Phylogenetic tree reconstruction

We reconstructed phylogenetic trees using Bayesian Inference implemented in MrBayes v3.2.7a

(Ronquist et al. 2012). We analyzed three datasets separately: the complete coding region of *ebony*, 400

bp of the 5' intergenic region, and mitochondrial DNA (*ND1*: NADH dehydrogenase subunit 1). For

the mitochondrial dataset, we used a different set of samples due to limited availability. Genomic DNA

was extracted using Gsync^{TM} DNA Extraction Kit, following the manufacturer's protocol. We amplified

the *ND1* region using primers (mtF9 and mtR14) listed in Supplemental Table S5. The samples used

for each analysis are detailed in Supplemental Table S2.

For all datasets, we used *D. fuyamai* as an outgroup. For the *ebony* region, we extracted the coding

sequence along with upstream intergenic regions from the genome assembly of *D. fuyamai* (accession

number: JAECXW010000874.1). For the mitochondrial dataset, we used *ND1* sequence from *D.*

fuyamai (accession number: HQ631806). Due to poor sequence conservation, the alignment quality of

the 400-bp upstream region of *ebony* with the outgroup was not optimal. Despite this limitation, we

included it in the phylogenetic reconstruction to explore potential evolutionary relationships.

To determine the best-fitting nucleotide substitution model for each dataset, we used jModelTest

2.1.10 v20160303 (Darriba et al. 2012) and selected models based on the Bayesian Information

Criterion (BIC). Convergence was confirmed by examining the average standard deviation of split

frequencies (<0.01). We discarded the first 25% of sampled trees as burn-in and summarized the

remaining trees using a 50% majority-rule consensus to construct the final phylogeny.

Statistical analysis

We conducted all statistical analyses in R for Windows v4.4.0 (R Core Team 2024), unless otherwise

specified. Specific statistical analysis approaches varied by analysis and are described in detail in the

relevant method subsections (e.g., QTL mapping, epistasis analysis). For RT-qPCR, we used one-way

ANOVA, followed by Tukey's HSD test for post hoc pairwise comparisons. For allele-specific

expression, we applied one-sample t-tests on log-transformed allele ratios to evaluate significant

deviations from equal expression (1:1 ratio) between alleles. We used two-way ANOVAs for epistasis

analyses.

Data access

All raw sequencing data generated in this study have been submitted to the National Center for

Biotechnology Information (NCBI; https://www.ncbi.nlm.nih.gov/) under accession number listed in

Supplemental Table S2. All other data (including QTL genotype and phenotype matrices, RT-qPCR

results, and image files) are available in Supplemental Code, Supplemental File, and Supplemental

Table S6.

Competing interest statement

The authors have no conflicts of interest to declare.

Acknowledgments

We thank Jonathan H. Massey for assistance with library preparation and genome assembly for

QTL mapping, and for helpful feedback on the early version of the manuscript. We are also grateful to

Nia Kurniawan, Karuniawan Puji Wicaksono, and Hagus Tarno from the University of Brawijaya for

their support during field sampling in Indonesia, including assistance with research permits and the

material transfer agreement. This study was funded by National Science and Technology Council grant

MOST107-2311-B-008-002 and MOST110-2311-B-008-002 to S.-D.Y. The field trip of fly collection

was funded by Ministry of Education grant to S.-D. Y.

Author contributions: S.-D. Y., B.-Y. L., and P. J. W. designed the project. A. S. F. wrote the

manuscript with comments from all authors. A. S. F. analyzed the data, generated recombinants for

QTL mapping, performed phenotyping, RT-qPCR, and ASE analyses. A. S. F. and S.-D. Y. collected

wild flies. A. N. processed and phased long-read sequences to assemble haplotypes of the *ebony* region

from wild-caught samples. K.-A.Y. extracted genomic DNA and prepared PCR amplicons for

sequencing of the *ebony* region. E. X. B. performed HCR experiments and analyzed the resulting data.

All authors read and approved the final manuscript.

Reference

Andolfatto P, Davison D, Erezyilmaz D, Hu TT, Mast J, Sunayama-Morita T, Stern DL. 2011.

Multiplexed shotgun genotyping for rapid and efficient genetic mapping. *Genome Res* **21**: 610-

617.

Archambeault SL, Bartschi LR, Merminod AD, Peichel CL. 2020. Adaptation via pleiotropy and

linkage: Association mapping reveals a complex genetic architecture within the stickleback Eda

locus. *Evol Lett* **4**: 282-301.

Bayala EX, Sinha P, Wittkopp PJ. 2024. Protocol for dissecting Drosophila pupae and visualizing RNA

expression using hybridization chain reaction. *STAR Protoc* **5**: 103456.

Bohutinska M, Peichel CL. 2024. Divergence time shapes gene reuse during repeated adaptation.

Trends Ecol Evol **39**: 396-407.

Borycz J, Borycz JA, Loubani M, Meinertzhagen IA. 2002. tan and ebony genes regulate a novel

pathway for transmitter metabolism at fly photoreceptor terminals. *J Neurosci* **22**: 10549-10557.

Brisson JA, De Toni DC, Duncan I, Templeton AR. 2005. Abdominal pigmentation variation in

drosophila polymorpha: geographic variation in the trait, and underlying phylogeography.

Evolution **59**: 1046-1059.

Broman KW, Wu H, Sen S, Churchill GA. 2003. R/qtl: QTL mapping in experimental crosses.

Bioinformatics **19**: 889-890.

Cande J, Andolfatto P, Prud'homme B, Stern DL, Gompel N. 2012. Evolution of multiple additive loci

caused divergence between Drosophila yakuba and D. santomea in wing rowing during male

courtship. *PLoS One* **7**: e43888.

Carr IM, Robinson JI, Dimitriou R, Markham AF, Morgan AW, Bonthron DT. 2009. Inferring relative

proportions of DNA variants from sequencing electropherograms. *Bioinformatics* **25**: 3244-

3250.

Cerca J. 2023. Understanding natural selection and similarity: Convergent, parallel and repeated

evolution. *Mol Ecol* **32**: 5451-5462.

Chung H, Carroll SB. 2015. Wax, sex and the origin of species: Dual roles of insect cuticular

hydrocarbons in adaptation and mating. *Bioessays* **37**: 822-830.

Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G, Jr., Dickson M, Grimwood J, Schmutz J,

Myers RM, Schluter D, Kingsley DM. 2005. Widespread parallel evolution in sticklebacks by

repeated fixation of Ectodysplasin alleles. *Science* **307**: 1928-1933.

Courtier-Orgogozo V, Arnoult L, Prigent SR, Wiltgen S, Martin A. 2020. Gephebase, a database of

genotype-phenotype relationships for natural and domesticated variation in Eukaryotes. *Nucleic

Acids Res* **48**: D696-D703.

Cuthill IC, Allen WL, Arbuckle K, Caspers B, Chaplin G, Hauber ME, Hill GE, Jablonski NG, Jiggins

CD, Kelber A et al. 2017. The biology of color. *Science* **357**.

Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and

parallel computing. *Nat Methods* **9**: 772.

Ferveur JF. 1997. The pheromonal role of cuticular hydrocarbons in Drosophila melanogaster.

Bioessays **19**: 353-358.

Gibert JM, Mouchel-Vielh E, De Castro S, Peronnet F. 2016. Phenotypic Plasticity through

Transcriptional Regulation of the Evolutionary Hotspot Gene tan in Drosophila melanogaster.

PLoS Genet **12**: e1006218.

Haley CS, Knott SA. 1992. A simple regression method for mapping quantitative trait loci in line

crosses using flanking markers. *Heredity (Edinb)* **69**: 315-324.

Hufnagel DE, Hufford MB, Seetharam AS. 2020. SequelTools: a suite of tools for working with PacBio

Sequel raw sequence data. *BMC Bioinformatics* **21**: 429.

Ishii K, Hirai Y, Katagiri C, Kimura MT. 2001. Sexual isolation and cuticular hydrocarbons in

Drosophila elegans. *Heredity (Edinb)* **87**: 392-399.

Izore T, Tailhades J, Hansen MH, Kaczmarski JA, Jackson CJ, Cryle MJ. 2019. Drosophila

melanogaster nonribosomal peptide synthetase Ebony encodes an atypical condensation

domain. *Proc Natl Acad Sci U S A* **116**: 2913-2918.

Johnson WC, Ordway AJ, Watada M, Pruitt JN, Williams TM, Rebeiz M. 2015. Genetic Changes to a

Transcriptional Silencer Element Confers Phenotypic Diversity within and between Drosophila

Species. *PLoS Genet* **11**: e1005279.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary Genetics

Analysis across Computing Platforms. *Mol Biol Evol* **35**: 1547-1549.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile

and open software for comparing large genomes. *Genome Biol* **5**: R12.

Lamb AM, Wang Z, Simmer P, Chung H, Wittkopp PJ. 2020. ebony affects pigmentation divergence

and cuticular hydrocarbons in Drosophila americana and D. novamexicana. *Front Ecol Evol* **8**.

Maeso I, Roy SW, Irimia M. 2012. Widespread recurrent evolution of genomic features. *Genome Biol*

Evol **4**: 486-500.

Martin A, Orgogozo V. 2013. The Loci of repeated evolution: a catalog of genetic hotspots of

phenotypic variation. *Evolution* **67**: 1235-1250.

Massey JH, Akiyama N, Bien T, Dreisewerd K, Wittkopp PJ, Yew JY, Takahashi A. 2019. Pleiotropic

Effects of ebony and tan on Pigmentation and Cuticular Hydrocarbon Composition in

Drosophila melanogaster. *Front Physiol* **10**: 518.

Massey JH, Li J, Stern DL, Wittkopp PJ. 2021. Distinct genetic architectures underlie divergent thorax,

leg, and wing pigmentation between Drosophila elegans and D. gunungcola. *Heredity (Edinb)*

127: 467-474.

Massey JH, Rice GR, Firdaus AS, Chen CY, Yeh SD, Stern DL, Wittkopp PJ. 2020. Co-evolving wing

spots and mating displays are genetically separable traits in Drosophila. *Evolution* **74**: 1098-

1111.

Massey JH, Wittkopp PJ. 2016. The Genetic Basis of Pigmentation Differences Within and Between

Drosophila Species. *Curr Top Dev Biol* **119**: 27-61.

Miyagi R, Akiyama N, Osada N, Takahashi A. 2015. Complex patterns of cis-regulatory

polymorphisms in ebony underlie standing pigmentation variation in Drosophila melanogaster.

Mol Ecol **24**: 5829-5841.

Parkash R, Sharma V, Kalra B. 2008. Climatic adaptations of body melanisation in Drosophila

melanogaster from Western Himalayas. *Fly (Austin)* **2**: 111-117.

Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic*

Acids Res **29**: e45.

Pool JE, Aquadro CF. 2007. The genetic basis of adaptive pigmentation variation in Drosophila

melanogaster. *Mol Ecol* **16**: 2844-2851.

R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. R Foundation for

Statistical Computing, Vienna, Austria.

Rebeiz M, Pool JE, Kassner VA, Aquadro CF, Carroll SB. 2009. Stepwise modification of a modular

enhancer underlies adaptation in a Drosophila population. *Science* **326**: 1663-1667.

Richardt A, Kemme T, Wagner S, Schwarzer D, Marahiel MA, Hovemann BT. 2003. Ebony, a novel

nonribosomal peptide synthetase for beta-alanine conjugation with biogenic amines in

Drosophila. *J Biol Chem* **278**: 41160-41166.

Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B, Liu L, Suchard

MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model

choice across a large model space. *Syst Biol* **61**: 539-542.

Rozas J, Ferrer-Mata A, Sanchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sanchez-

Gracia A. 2017. DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data Sets. *Mol*

Biol Evol **34**: 3299-3302.

Stern DL. 2013. The genetic causes of convergent evolution. *Nat Rev Genet* **14**: 751-764.

Takahashi A. 2013. Pigmentation and behavior: potential association through pleiotropic genes in

Drosophila. *Genes Genet Syst* **88**: 165-174.

Takahashi A, Takahashi K, Ueda R, Takano-Shimizu T. 2007. Natural variation of ebony gene

controlling thoracic pigmentation in Drosophila melanogaster. *Genetics* **177**: 1233-1237.

Takahashi A, Takano-Shimizu T. 2011. Divergent enhancer haplotype of ebony on inversion

In(3R)Payne associated with pigmentation variation in a tropical population of Drosophila

melanogaster. *Mol Ecol* **20**: 4277-4287.

Telonis-Scott M, Hoffmann AA, Sgro CM. 2011. The molecular genetics of clinal variation: a case

study of ebony and thoracic trident pigmentation in Drosophila melanogaster from eastern

Australia. *Mol Ecol* **20**: 2100-2110.

Wang Z, Receveur JP, Pu J, Cong H, Richards C, Liang M, Chung H. 2022. Desiccation resistance

differences in Drosophila species can be largely explained by variations in cuticular

hydrocarbons. *Elife* **11**.

Wittkopp PJ, Stewart EE, Arnold LL, Neidert AH, Haerum BK, Thompson EM, Akhras S, Smith-

Winberry G, Shefner L. 2009. Intraspecific polymorphism to interspecific divergence: genetics

of pigmentation in Drosophila. *Science* **326**: 540-544.

Wittkopp PJ, True JR, Carroll SB. 2002. Reciprocal functions of the Drosophila yellow and ebony

proteins in the development and evolution of pigment patterns. *Development* **129**: 1849-1858.