RNA reinforces condensate nucleation on chromatin to amplify oncogenic transcription

Krista A. Budinich[1,2,3,4#], Xinyi Yao[5#], Chujie Gong[1,2,3,4#], Lele Song[1,2,4], Michelle C. Lee[1,2,4,6], Qinglan Li[1,2,4], Sylvia Tang[1,2,4], Kaeli M. Mathias[1,2,4,7,8], Yiman Liu[1,2,4], Son C. Nguyen[4,9], Eric F. Joyce[4,9], Yuanyuan Li[5*], Haitao Li[5,10,11*], and Liling Wan[1,2,4,12,13*]

[1]Department of Cancer Biology, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[2]Abramson Family Cancer Research Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[3]Cell and Molecular Biology Graduate Group, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.

[4]Penn Epigenetics Institute, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[5]State Key Laboratory of Molecular Oncology, MOE Key Laboratory of Protein Sciences, Beijing Frontier Research Center for Biological Structure, School of Basic Medical Sciences, Tsinghua University, Beijing 100084, China.

[6]School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[7]Biochemistry and Molecular Biophysics Graduate Group, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[8]Center for Computational and Genomic Medicine, The Children's Hospital of Philadelphia, Philadelphia, PA, 19104, USA.

[9]Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[10]Tsinghua-Peking Center for Life Sciences, Beijing 100084, China.

[11]SXMU-TM Collaborative Innovation Center for Frontier Medicine, Shanxi Medical University, Taiyuan 030001, China.

[12]Institute for Regenerative Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA.

[13]Lead contact

[#] These authors contributed equally.

[*]Correspondence should be addressed to L.W. (liling.wan@pennmedicine.upenn.edu), H.L.

(lht@tsinghua.edu.cn), Y. Li (liyuanyuan@tsinghua.edu.cn)

ABSTRACT

Aberrant chromatin-associated condensates have emerged as drivers of transcriptional dysregulation in

cancer, yet the mechanisms regulating their formation and function remain poorly understood. Gain-of-

function mutations in the histone acetylation reader ENL, found in leukemia and Wilms tumor, drive

oncogenesis by inducing condensate formation at highly selective genomic loci. Here, we uncover a

critical role for RNA in reinforcing the nucleation, chromatin engagement, and oncogenic activity of ENL

mutant condensates. Mutant ENL binds RNA via a conserved basic patch within its YEATS domain, and

this interaction enhances condensate formation both *in vitro* and across diverse cellular contexts. Using a

chemically inducible condensate displacement and re-nucleation system, we show that blocking ENL–

RNA interactions or transcription impairs condensate reformation at endogenous targets. RNA

interactions preferentially enhance ENL occupancy at condensate-permissive loci, leading to increased

transcriptional bursting and robust gene activation at the single-cell and single-allele level. In mouse

models, disrupting ENL–RNA interactions diminishes condensate formation and oncogenic transcription

in hematopoietic stem and progenitor cells, thereby suppressing ENL mutation-driven leukemogenesis.

Together, these findings establish RNA as an active cofactor that pathogenic condensates exploit to

reinforce locus-specific nucleation and amplify transcriptional programs that drive disease.

INTRODUCTION

An increasing number of chromatin-associated proteins have been shown to form nuclear condensates—high local-concentration assemblies often driven by dynamic multivalent interactions[1]. In some cases, these condensates enrich transcriptional machinery at specific genomic loci to regulate gene expression[2-8]. Dysregulation of condensate formation has been implicated in a range of diseases, notably cancer[9-11]. Genetic mutations or chromosomal translocations in regulatory proteins can induce aberrant condensates, which may be co-opted by cancer cells to activate and sustain oncogenic gene programs[9-11]. However, the molecular mechanisms that regulate the assembly and function of pathogenic condensates remain poorly understood.

RNA has long been recognized as a key component of various biomolecular condensates, including stress granules, the nucleolus, and nuclear speckles[12-14]. Emerging evidence suggests that RNA can also modulate condensates formed by transcriptional regulators. For instance, RNA has been proposed to regulate transcription through a feedback mechanism: low levels of RNA produced during transcription initiation promote condensate formation, while elevated RNA levels accumulated during elongation stimulate their dissolution[15]. In pluripotent stem cells, RNA transcribed from endogenous retroviral elements has been shown to facilitate the hijacking of super-enhancer-associated condensates[16]. Despite these advances, whether and how RNA regulates the formation and function of chromatin-associated pathogenic condensates remains largely unexplored.

To address this question, we leverage oncogenic ENL (Eleven-nineteen-leukemia) mutants found in pediatric acute myeloid leukemia (AML) and Wilms tumor[17-21]. In its wildtype form, ENL binds acetylated histones via its YEATS domain and promotes transcriptional elongation at target genes[22,23]. We previously demonstrated that cancer-associated hotspot mutations in the ENL YEATS domain induce aberrant condensate formation at selective target loci, such as the *HOXA* cluster. These condensates are required for ENL mutation-induced chromatin occupancy, hyperactivation of target genes, and tumorigenesis[24-26]. However, how these condensates are regulated remains unknown. Given the role of RNA in other nuclear condensates[12] and emerging evidence that many transcriptional regulators interact with RNA[27-32], we hypothesized that RNA may modulate ENL mutant condensates. The robust

condensate-forming ability, high locus specificity, and potent oncogenic activity of ENL mutants make

them an ideal model for dissecting the regulatory mechanisms of pathogenic condensates.

Here, we uncover a positive role for RNA in promoting ENL mutant condensate formation and

function. ENL mutants bind RNA via a conserved basic patch within the YEATS domain, and this

interaction enhances condensate nucleation and transcriptional bursting at permissive loci. Disrupting

ENL:RNA interactions impairs oncogenic gene activation and tumorigenesis in mouse models. These

findings reveal an RNA-dependent mechanism regulating pathogenic condensates and establish RNA as

a functional co-factor in condensate-mediated gene dysregulation in cancer.

RESULTS

ENL binds RNA via a conserved basic patch in the YEATS domain *in vitro* and in cells

A previous study showed that AF9, a paralog of ENL, can bind DNA in part through residues in its YEATS

domain[33]. To assess whether ENL shares this property, we performed electrophoretic mobility shift assay

(EMSA) using purified full-length wildtype ENL (ENL-WT) and the oncogenic ENL-T1 mutant, the most

common of eight patient-derived hotspot mutations found within ENL's YEATS domain (Fig. 1a). We

found that ENL-WT bound double-stranded DNA (dsDNA) in a concentration-dependent manner, and the

T1 mutation did not alter this capacity (Extended Data Fig. 1a, b).

To test whether ENL can also bind RNA, we performed EMSA with a single-stranded nascent

RNA (ssRNA) fragment derived from *HOXA11*—a known ENL-T1 target gene whose nascent transcripts

are detected within cellular ENL-T1 condensates[25]—and found that ENL-WT and T1 bound *HOXA11*

RNA with comparable affinities (Extended Data Fig. 1c,d). These results indicate that ENL harbors RNA-

binding capacity and this capability is preserved in cancer-associated ENL mutants. Given the strong

enrichment of nascent RNA in ENL-T1 condensates[25] and the reported role of RNA in modulating

transcriptional condensates[15], we focused our study on the role of ENL:RNA interactions in condensate

formation and function.

We next sought to identify the region within ENL that mediates RNA interaction. Given that AF9

binds DNA through its YEATS domain, we examined the electrostatic surface potential of ENL's YEATS

domain and identified four positively charged surface patches that could engage with negatively charged

RNA (Fig. 1b). To evaluate RNA-binding propensity, we modeled interactions between the ENL YEATS

domain and ~100 diverse RNA sequences using AlphaFold3 (Extended Data Fig. 1e and Supplementary

Table 1). This analysis highlighted the basic patch encompassing residues K61-K67 as the region most

likely to interact with RNA (Fig. 1c). Notably, several positively charged residues within this region align

with the DNA-binding residues identified in AF9 (Extended Data Fig. 1f). Structural inspection revealed

that among these residues, K61, R63, and K67 are surface-exposed and accessible, whereas R64 is

less exposed (Extended Data Fig. 1g).

To test the contribution of this region to RNA binding, we generated a charge-neutralizing triple

alanine mutant (K61A/R63A/K67A, hereafter '3A') predicted to preserve the overall YEATS domain

structure (Extended Data Fig. 1h). AlphaFold3 modeling showed that the T1-3A mutant exhibited

significantly fewer predicted contacts with RNA compared to T1, suggesting a potential role for these

residues in RNA interaction (Fig. 1d and Extended Data Fig. 1i). EMSA using full-length proteins and a

nascent *HOXA11* RNA fragment confirmed that T1-3A exhibited reduced RNA binding relative to T1,

while retaining partial interactions likely mediated by other basic regions outside of the YEATS domain

(Fig. 1e, f)**.** EMSA using purified YEATS domains alone showed that the 3A mutation markedly impaired

RNA binding (Extended Data Fig. 1j, k), validating this basic patch as the key RNA-binding surface within

the YEATS domain.

To determine whether ENL binds RNA in cells, we performed CLAP-seq (Covalent Linkage and

Affinity Purification followed by sequencing), an adaptation of CLIP-seq optimized to more directly detect

protein–RNA interactions under denaturing conditions[34]. We expressed Halo-tagged ENL-WT, T1, or T1-

3A at near endogenous levels in HEK293 cells (Extended Data Fig. 2a), UV-crosslinked protein–RNA

complexes, affinity-purified ENL and its associated RNAs using Halo ligand resin under denaturing

conditions, and sequenced the recovered RNAs (Fig. 1g). Comparison of CLAP pull-downs with matched

inputs identified ~1,600 RNA peaks across all three genotypes, with fewer than 10% unique to any single

condition (Extended Data Fig. 2b and Supplementary Table 2). Biological replicates were highly

correlated (Extended Data Fig. 2c). WT and T1 displayed similar CLAP signal intensities across the

merged peak set (Fig. 1h), consistent with their comparable RNA-binding capacity observed by EMSA

(Extended Data Fig. 1c, d). T1-3A bound largely to these same peak regions, but with reduced signal

(Fig. 1h, i). These results established that ENL binds RNA in cells and that this interaction is in part

mediated by the K61/R63/K67 basic patch.

Genomic analysis revealed that the majority of CLAP input peaks mapped to exonic regions,

reflecting the high abundance of spliced mRNA in the cell (Fig. 1j and Extended Data Fig. 2d). In

contrast, ENL CLAP peaks were strongly enriched in intronic regions (Fig. 1j and Extended Data Fig. 2e),

suggesting that ENL primarily binds nascent transcripts. Annotation of CLAP peaks to genes revealed

that ENL-bound RNAs were predominantly protein-coding (Fig. 1k and Extended Data Fig. 1f), consistent

with ENL's known localization to protein-coding genes[22,26] (Fig. 1k). When plotting CLAP signal across

the gene bodies of these targets, we found that ENL-WT and T1 exhibited similar levels of RNA

association, with T1 showing a modest increase. In contrast, T1-3A showed reduced RNA association

across these gene bodies (Fig. 1l-n, Extended Data Fig. 2g, and Supplementary Table 3).

Altogether, our *in vitro* and cellular findings demonstrate that ENL binds RNA and that this

interaction depends, at least in part, on a conserved basic patch within its YEATS domain. We therefore

adopted ENL-T1-3A as a molecular tool to probe how partially impaired ENL:RNA interactions affect the

formation and function of condensates driven by cancer-associated mutations.

RNA enhances droplet formation of oncogenic ENL mutant *in vitro*

To test whether RNA contributes to ENL condensate formation, we performed *in vitro* droplet

assays using purified full-length ENL and RNA, including total RNA extracted from whole cells or nuclear

fractions, as well as with chemically synthesized nascent RNA from *HOXA11*, a known ENL-T1 target

gene (Fig. 2a-c and Extended Data Fig. 3a-d). ENL-T1 formed droplets more readily than WT in the

absence of RNA, consistent with increased self-association conferred by the oncogenic mutation[25].

Addition of RNA enhanced droplet formation of both ENL-WT and T1 in a dose-dependent manner, with

T1 showing a more pronounced increase (Fig. 2b, c and Extended Data Fig. 3a-d).

To determine whether this RNA responsiveness depends on the basic patch in the YEATS

domain, we tested the T1-3A mutant. Notably, the 3A mutation did not impair the intrinsic droplet forming

abilities conferred by the T1 mutation, as observed in the absence of RNA (Fig. 2b, c and Extended Data

Fig. 3a-d). Rather, 3A specifically impaired RNA-dependent enhanced droplet formation (Fig. 2b, c and

Extended Data Fig. 3a-d). Fluorescent labeling further showed stronger incorporation of *HOXA11* RNA

into ENL-T1 droplets compared to T1-3A droplets (Fig. 2b, d), supporting that RNA-mediated droplet

enhancement is facilitated through interactions with the basic patch.

RNA binding promotes ENL mutant condensate formation at endogenous target loci

We previously showed that oncogenic ENL mutations induce condensate formation at highly specific

genomic targets in cells, driving aberrant gene activation and tumorigenesis[25,26]. These mutations thus

provide a tractable system to interrogate how specific features, such as RNA binding, affect condensate

formation and function in cells. We expressed doxycycline-inducible FLAG-tagged ENL variants at near-

endogenous levels in two human kidney-derived cell lines, HEK293 and WiT49 (Fig. 2e, f and Extended

Data Fig. 4a, b). We focused on two of the most common patient-derived mutants: ENL-T1 (insertion)

and ENL-T2 (deletion) (Fig. 1a), and generated RNA-binding-deficient T1-3A and T2-3A by introducing

the K61A/R63A/K67A triple mutation.

Consistent with previous findings, FLAG-ENL immunofluorescence (IF) revealed that ENL-T1 and

T2 formed discrete nuclear condensates not seen with ENL-WT (Fig. 2f-h and Extended Data Fig. 4b).

Notably, the 3A mutation significantly reduced the number of condensates per cell, indicating that RNA

binding promotes the ability of ENL-T1 and T2 to form condensates (Fig. 2f-h and Extended Data Fig.

4b). Cycloheximide chase assays showed comparable protein stability across all variants (Extended

Data Fig. 4c, d), indicating that the reduction in condensate formation is not simply due to differences in

protein turnover. These phenotypes were observed across additional cell lines, including A549 (lung

cancer) and MDA-MB-231 (breast cancer), where ENL-T1 and T2 again formed discrete nuclear

condensates while the 3A mutation reduced condensate number (Fig. 2i, j and Extended Data Fig. 4e-h).

To determine whether ENL:RNA interactions contribute to condensate formation at endogenous target genes, we performed concurrent FLAG-ENL IF and DNA-FISH using probes targeting the *HOXA* locus or *CBX3,* two key ENL-T1 and T2 target loci in kidney cell lines. In HEK293 cells expressing ENL-T1 or T2, the majority of *HOXA* and *CBX3* DNA alleles colocalized with ENL mutant puncta (Fig. 2k-r), indicating robust and frequent condensate formation at these loci. In contrast, colocalization was markedly reduced in cells expressing T1-3A or T2-3A variants (Fig. 2k-r), consistent with a lower overall number of nuclear condensates that likely reduces the frequency of condensate association at any given target locus.

Corroborating our imaging data, chromatin immunoprecipitation followed by quantitative PCR (ChIP–qPCR) showed that ENL-T1 and T2 exhibited markedly increased occupancy at *HOXA* and *CBX3* relative to ENL-WT, and this enrichment was significantly reduced in T1-3A and T2-3A (Fig. 2s, t). Importantly, isothermal titration calorimetry using purified YEATS domain proteins demonstrated that the 3A mutation does not impair histone acetylation binding affinity of the ENL-T1 or T2 YEATS domains (Extended Data Fig. 4i), ruling out loss of histone acetylation recognition as the main cause of reduced chromatin occupancy.

Together, these findings support a model in which ENL:RNA interactions, mediated in part through the K61/R63/K67 basic patch within the YEATS domain, facilitate condensate formation and chromatin engagement by oncogenic ENL mutants at endogenous target loci.

RNA binding promotes re-nucleation of ENL condensates at target loci following acute chemical displacement

As disruption of ENL:RNA interactions via the 3A mutation reduces condensate number (Fig. 2g-j), we hypothesized that RNA binding facilitates condensate nucleation. To directly test this, we established a chemical-induced condensate displacement and wash-off system to visualize and quantify condensate re-nucleation dynamics at endogenous targets (Fig. 3a). DNA-FISH combined with FLAG-ENL IF showed that ENL-T1 condensates frequently colocalize with the *HOXA* locus in HEK293 cells at basal conditions (Figs. 2k, l and 3a, left panel). To monitor condensate nucleation at this site, we first displaced

condensates using TDI-11055, a small-molecule inhibitor that competitively blocks the interaction between the ENL YEATS domain and acetylated histones[26,35] (Fig. 3a, middle panel). Following inhibitor wash-off, we tracked the re-formation of condensates at the *HOXA* locus over time in cells expressing either ENL-T1 or the RNA-binding-deficient T1-3A mutant (Fig. 3a, right panel, b).

In ENL-T1–expressing cells, TDI-11055 treatment markedly reduced the percentage of *HOXA* alleles associated with condensates compared to untreated controls. Upon wash-off, condensate occupancy at *HOXA* loci recovered over a 10-hour period, reaching levels comparable to untreated controls (Fig. 3c, d and Extended Data Fig. 5a). This was accompanied by restored expression of *HOXA11* and *HOXA13* (Fig. 3e and Extended Data Fig. 5b). In contrast, recovery was impaired in T1-3A–expressing cells, both in condensate nucleation and gene activation (Fig. 3c-e and Extended Data Fig. 5a, b).

To further probe of the role of nascent transcripts in condensate nucleation, we applied transcription inhibitors in the wash-off system (Fig. 3f, g). ENL-T1–expressing cells were pre-treated with either DMSO or a transcription inhibitor 1 hour prior to TDI-11055 wash-off. In DMSO-treated cells, condensates rapidly re-formed at the *HOXA* locus, with 50% of *HOXA* alleles showing condensate association within 6 hours post wash-off (Fig. 3h, j). This re-formation was accompanied by largely restored *HOXA11* and *HOXA13* expression (Fig. 3i, k and Extended Data Fig. 5c, d). In contrast, acute transcription inhibition using Actinomycin D (blocks initiation) or DRB (blocks elongation) significantly impaired condensate re-formation (Fig. 3h-k and Extended Data Fig. 5c, d).

Together, using a unique chemical strategy to acutely displace and monitor re-formation of condensates at defined genomic loci, we show that both RNA-binding capability and ongoing transcription are required for efficient nucleation of ENL-T1 condensates.

RNA binding preferentially enhances ENL mutant occupancy at condensate-permissive loci

Our DNA-FISH and ChIP-qPCR analyses showed that disrupting ENL:RNA interactions via the 3A mutation reduces chromatin occupancy of oncogenic ENL mutants at condensate-associated genomic loci (Fig. 2k-r). To assess whether RNA binding contributes to ENL mutant occupancy genome-

wide or preferentially at specific targets, we performed ChIP-seq for FLAG-ENL variants in HEK293 cells.

ENL-T1 and T2 largely occupied the same genomic regions as ENL-WT but with increased occupancy

(Extended Data Fig. 6a, b), consistent with increased self-association conferred by these oncogenic

mutations[25]. Ranking ENL-T1 or ENL-T2 peaks by their ChIP-seq signal revealed a highly asymmetric

distribution, with fewer than 1% of peaks showing disproportionately high occupancy (Fig. 4a and

Extended Data Fig. 6c). These top-ranked peaks included validated condensate-associated targets such

as the *HOXA* cluster and *CBX3* (Fig. 2k-r). Using the inflection point of the ranked plot, we classified

ENL-T1 and ENL-T2 peaks into top- and low-bound groups (Fig. 4a, Extended Data Fig. 6c, and

Supplementary Table 4). Top-bound peaks showed a greater fold increase in ENL-T1 or T2 occupancy

relative to ENL-WT (Fig. 4b and Extended Data Fig. 6d), suggesting that these target sites are most

likely to associate with condensates (Fig. 4c).

We next asked whether RNA interactions contribute differentially to ENL chromatin occupancy at

top- versus low-bound sites. At low-bound sites, both T1 and T1-3A showed similarly increased

occupancy relative to WT, with minimal differences between the two (Fig. 4d-f). In contrast, T1-3A

exhibited significantly reduced occupancy relative to T1 at top-bound sites, indicating that RNA binding

selectively reinforces ENL-T1 occupancy at condensate-associated loci. Similar results were observed

for T2 versus T2-3A comparisons (Extended Data Fig. 6e-g).

To test whether this selective requirement is conserved across cellular contexts with distinct

condensate-associated targets, we performed ChIP–seq in MDA-MB-231 breast cancer cells expressing

FLAG-ENL variants. As in HEK293, ENL-T1 and ENL-T2 exhibited an asymmetric distribution of ChIP–

seq signals. However, the top-bound peaks differed substantially between the two cell lines, reflecting

cell type-specific targeting (Fig. 4g, Extended Data Fig. 7a, b, and Supplementary Tables 5, 6).

Nevertheless, top-bound peaks in MDA-MB-231 also showed a greater increase in occupancy in T1 and

T2 relative to WT (Fig. 4h and Extended Data Fig. 7c). To test whether these top-bound sites are indeed

associated with condensates, we performed DNA-FISH for two top targets, *NR2F2* and *CHD2,* and found

that ENL-T1 and T2 condensates frequently colocalized with these targets (Fig. 4i-k and Extended Data

Fig. 7d-h). As in HEK293, the 3A mutation reduced ENL-T1 and T2 occupancy more strongly at top-

bound than at low-bound sites (Fig. 4l-n and Extended Data Fig. 7i-k), further supporting the conclusion

that RNA binding preferentially promotes ENL chromatin engagement at condensate-permissive loci.

Together, these findings support a model in which oncogenic mutations amplify ENL occupancy at

pre-existing target sites, but only a small, cell type–specific subset are permissive to condensate

formation. Although the molecular features of these sites remain to be defined, RNA interactions appear

to contribute more to ENL occupancy at these sites, likely by facilitating condensate nucleation than

general chromatin binding.

RNA binding enhances transcriptional bursting and gene activation at condensate-permissive

loci

Given that ENL mutant condensates drive hyperactivation of target genes[25,26], we next asked

whether RNA binding contributes to this function. We performed RNA-seq in HEK293 and MDA-MB-231

cells expressing ENL-T1, T2, or their RNA-binding-deficient 3A variants. Gene set enrichment analysis

(GSEA) using ChIP–seq–defined top-bound genes (Fig. 4a, g, Extended Data Fig. 6c, 7b, and

Supplementary Tables 7, 8) revealed upregulation of these targets in ENL-T1 and T2 expressing cells

compared to WT (Fig. 4o, p and Extended Data Fig. 8a, b). This enrichment was attenuated in T1-3A and

T2-3A mutants (Fig. 4o, p and Extended Data Fig. 8a, b), indicating that RNA binding contributes to the

hyperactivation of condensate-associated targets. RT–qPCR for nascent transcripts confirmed reduced

expression of key targets in T1-3A and T2-3A–expressing cells relative to T1 and T2 in both cell lines

(Fig. 4q and Extended Data Fig. 8c, d).

To examine transcriptional activity at single-cell and single-allele resolution, we performed RNA-

FISH targeting intronic *HOXA13* transcripts with Halo-tagged ENL staining in HEK293 cells. ENL-T1 and

T2 expression led to a significant increase in detected RNA-FISH foci per cell compared to WT (Fig. 4r, s

and Extended Data Fig. 8e, f), indicative of increased transcription bursting frequency at the *HOXA*

locus. Notably, nearly all FISH foci in T1 or T2–expressing cells colocalized with condensates,

highlighting a strong link between condensate formation and transcriptional bursting (Fig. 4r-t and

Extended Data Fig. 8e-g). The T1 and T2-induced increase in RNA-FISH foci was largely abrogated by

the 3A mutation (Fig. 4r-t and Extended Data Fig. 8e-g). Although T1-3A and T2-3A cells showed fewer

RNA-FISH foci, those detected still colocalized with condensates and exhibited comparable intensities to

those observed in T1 or T2 cells (Fig. 4t and Extended Data Fig. 8g, h), reinforcing the link between

condensates and gene activation. Together, these results suggest that RNA binding primarily promotes

the frequency of condensate formation on target genes, thereby increasing transcriptional bursting

frequency and enabling robust gene activation by oncogenic ENL mutants.

Disrupting RNA binding impairs ENL mutant–driven leukemogenesis

We previously demonstrated that cancer-associated ENL mutations are bona fide drivers of AML using

both conditional knock-in mouse models and transplantation models of ectopic expression in

hematopoietic stem/progenitor cells (HSPCs)[26]. Building on this, we sought to determine whether RNA-

binding is required for the oncogenic activity of ENL mutants. We isolated LSK (Lin⁻Sca1⁺cKit⁺) cells

from wildtype C57BL/6 mice and transduced them with FLAG-tagged ENL-WT, T1, or T1-3A constructs

(Fig. 5a), which were expressed at comparable RNA and protein levels (Extended Data Fig. 9a, b). FLAG

IF showed that ENL-T1 formed discrete condensates not seen with ENL-WT, while T1-3A exhibited a

marked reduction in condensate number (Fig. 5b, c). We next performed DNA-FISH targeting the *Hoxa*

cluster and *Meis1*, two key loci previously shown to colocalize with and are activated by ENL mutant

condensates in LSK cells[26]. Colocalization of these loci with ENL condensates was significantly more

frequent in T1 than in T1-3A–expressing cells (Fig. 5d-g), supporting that RNA binding promotes

condensate formation at critical leukemogenic gene loci in HSPCs.

To assess the transcriptional consequences of disrupting ENL:RNA interactions, we performed

RNA-seq in LSK cells expressing ENL-WT, T1, or T1-3A. Differential gene expression analysis (fold

change > 1.5; *padj* < 0.05) identified five gene clusters (Extended Data Fig. 9c and Supplementary Table

9). Consistent with ENL-T1 functioning as a transcriptional activator, most differentially expressed genes

(DEGs) were upregulated from WT to T1 (C3 and C5), while a smaller subset of DEGs were

downregulated (C1 and C2) (Extended Data Fig. 9c). Only a limited number of DEGs were uniquely

imparted by ENL-T1-3A (C4) (Extended Data Fig. 9c). Focusing on C3 and C5, which represent the

majority of DEGs and those upregulated by ENL-T1, we found that the 3A mutation impaired ENL-T1-

driven gene activation (Fig. 5h, i), including reduced expression of key condensate-associated targets,

the *Hoxa* cluster genes and *Meis1* (Fig. 5j).

Using our previously published ENL-T1 chromatin occupancy dataset generated in LSK cells[26],

we stratified ENL-T1-bound peaks into top- and low-bound groups (Extended Data Fig. 9d and

Supplementary Table 10). GSEA revealed that genes associated with top-bound peaks were significantly

upregulated in T1 versus WT and remained enriched in T1 compared to T1-3A (Fig. 5k and Extended

Fig. 9e). Moreover, genes associated with leukemic stem cells in AML patients were enriched in T1

relative to both WT and T1-3A (Fig. 5l and Extended Data Fig. 9f), indicating that RNA binding

contributes to ENL mutant-driven leukemogenic programs.

Finally, we assessed the impact of perturbing ENL:RNA interactions on leukemogenesis. We first

performed serial colony formation assays to evaluate self-renewal capacity. ENL-T1–expressing LSK

cells maintained robust colony-forming ability over multiple replatings, whereas WT and T1-3A cells

rapidly lost this potential (Fig. 5m), indicating that RNA binding is critical for sustaining the T1-induced

oncogenic state. We next transplanted LSK cells expressing ENL-WT, T1, or T1-3A into lethally irradiated

mice and monitored engraftment over time (Fig. 5n). All groups had similar initial engraftment at two

weeks post transplantation (Fig. 5o). However, at the terminal timepoint, defined by maximal disease

burden in ENL-T1 mice or one year post-transplantation for WT and T1-3A, ENL-T1 mice exhibited

significantly higher engraftment (Fig. 5o and Extended Data Fig. 9g) and markedly reduced survival

across two independent cohorts (Fig. 5p and Extended Data Fig. 9h). Postmortem analysis confirmed

hallmark AML features in ENL-T1 mice, including splenomegaly and leukemic blast infiltration into lung,

liver, and kidneys. In contrast, WT and T1-3A mice showed no signs of leukemia even one year post

transplantation (Fig. 5q, r). Together, these results demonstrate that ENL:RNA interactions are critical for

supporting the full oncogenic potential of cancer-associated ENL mutants in disease-relevant models.

DISCUSSION

Our findings reveal that RNA interactions actively promote the nucleation and oncogenic function

of chromatin-associated condensates. While ENL mutations are the primary trigger of condensate

formation[25], genetic and chemical perturbations demonstrate that RNA binding enhances condensate

nucleation efficiency at selective genomic loci. These interactions reinforce local enrichment of mutant

ENL—and likely associated co-factors—which amplifies transcriptional bursting and oncogenic activation

essential for tumorigenesis. This effect is particularly pronounced at condensate-permissive sites,

suggesting that condensates impose locus-specific transcriptional control through distinct, RNA-sensitive

mechanisms (Fig. 5s).

The positive role of RNA identified here in the context of pathogenic condensates contrasts with

models from normal cells, where accumulating RNA is thought to exert negative feedback on

transcription. For instance, while low levels of enhancer RNAs facilitate Mediator condensate formation

at super-enhancers in mouse embryonic stem cells, increasing levels of nascent RNA during

transcriptional bursts have been shown to disrupt condensates by altering the local electrostatic

balance[15,36]. Such feedback mechanisms likely serve to ensure proper gene expression during

development and to maintain homeostasis[36]. In contrast, cancer cells often depend on persistent

oncogene activation to maintain malignant states[37]. As exemplified by ENL mutations, one way to

achieve this is by co-opting condensates that both promote transcription and bypass RNA-mediated

negative feedback.

We propose that basal levels of nascent transcripts provide multivalent interactions that lower the

energetic barrier for condensate nucleation, enabling mutant ENL to efficiently concentrate at key

genomic targets and drive aberrant transcription. Elevated levels of locally produced RNA—rather than

triggering condensate dissolution—may be buffered through direct interactions with ENL and other

factors within the condensates. This buffering effect could further stabilize condensate association and

sustain high-level transcriptional output critical for tumorigenesis.

Beyond mutant ENL, other oncogenic drivers, such as NUP98-HOXA9, EWS-FLI1, and mutant

nucleophosmin (NPM1), also form condensates enriched in regulatory factors[6,38-42]. These proteins often

contain intrinsically disordered regions with charge blocks that may enable interactions with RNA[38,43,44].

Thus, we speculate that this RNA-dependent condensate regulation may represent a broader strategy

through which cancer cells achieve persistent pathogenic gene activation. While our study highlights how

RNA interactions promote condensate formation and function, condensates can also influence RNA fate.

For example, condensates formed by the m6A reader YTHDC1 were shown to protect *MYC* transcripts

from degradation in leukemia[45]. In other settings, high levels of nascent transcripts can suppress further

transcription from the same locus through autoregulatory mechanisms[46], independent of condensates. It

is therefore possible that pathogenic condensates might buffer these transcripts and bypass such

feedback, thereby enabling sustained transcription, a hypothesis warranting further investigation.

Together, our findings establish RNA as an active cofactor that pathogenic condensates can

exploit not only to promote their formation, but also to sustain aberrant transcriptional programs. This

feed-forward mechanism may represent a broadly used strategy in cancer and other diseases

characterized by persistent transcriptional dysregulation and could offer new avenues for mechanistic

understanding and therapeutic exploration.

DATA AVAILABILITY

All CLAP-seq, ChIP-seq, and RNA-seq raw and processed data have been deposited in the Gene

Expression Omnibus database under accession numbers GSE300418, GSE300420, GSE300421. All

other raw data generated or analyzed during this study are included in this paper, the Extended Data

Figures, and the Supplementary Information.

ACKNOWLEDGEMENTS

We thank members of the Wan Laboratory and the Li Laboratory for scientific input throughout the study;

the CDB Microscopy Core at the University of Pennsylvania; the Flow Cytometry and Cell Sorting Core at

the Children's Hospital of Pennsylvania; the Pathology Core at the Children's Hospital of Pennsylvania;

the AFCRI Research Facilities Team at the University of Pennsylvania, and the University Laboratory

Animal Resources team at the University of Pennsylvania. The research was supported by funding from

AUTHOR CONTRIBUTIONS

K.A.B, Y.Li, H.L., and L.W. conceived and designed the study. K.A.B performed the cellular, molecular,

genomics, and *in vivo* animal studies. X.Y. and Y.Li performed *in vitro* biochemical studies. C.G.

performed bioinformatic analysis of genomics data. Q.L. provided bioinformatics analysis pipelines.

M.C.L., S.T., Y.Liu, L.S., and K.M.M. provided technical assistance. S.C.N. and E.F.J. designed and

synthesized FISH probes. K.A.B. and L.W. wrote the manuscript with input from H.L., Y.Li, X.Y., L.S., and

C.G. H.L. and Y.Li jointly supervised the *in vitro* biochemical studies. L.W. supervised the overall study.

COMPETING INTERESTS

379   L.W. is a co-inventor on a patent filed (US No. 62/949,160) related to the inhibitor used in this

380   manuscript. All other authors declare no competing interests.

MATERIALS AND CORRESPONDENCE

383   Correspondence and requests for materials should be addressed to lead contact L.W.

384   (liling.wan@pennmedicine.upenn.edu).

REFERENCES

Banani, S. F., Lee, H. O., Hyman, A. A. & Rosen, M. K. Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285-298, doi:10.1038/nrm.2017.7 (2017).

Cho WK, S. J., Hecht M, Lee C, Li C, Grube V, Cisse II. Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* **361**, doi:10.1126/science.aar4199 (2018).

Mir, M. *et al.* Dynamic multifactor hubs interact transiently with sites of active transcription in Drosophila embryos. *Elife* **7**, doi:10.7554/eLife.40497 (2018).

Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. & Sharp, P. A. A Phase Separation Model for Transcriptional Control. *Cell* **169**, 13-23, doi:10.1016/j.cell.2017.02.007 (2017).

Bhat, P., Honson, D. & Guttman, M. Nuclear compartmentalization as a mechanism of quantitative control of gene expression. *Nat Rev Mol Cell Biol* **22**, 653-670, doi:10.1038/s41580-021-00387-1 (2021).

Chong, S. *et al.* Imaging dynamic and selective low-complexity domain interactions that control gene transcription. *Science* **361**, doi:10.1126/science.aar2555 (2018).

Sabari, B. R. *et al.* Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361**, doi:10.1126/science.aar3958 (2018).

Lu, H. *et al.* Phase-separation mechanism for C-terminal hyperphosphorylation of RNA polymerase II. *Nature* **558**, 318-323, doi:10.1038/s41586-018-0174-3 (2018).

Boija, A., Klein, I. A. & Young, R. A. Biomolecular Condensates and Cancer. *Cancer Cell* **39**, 174-192, doi:10.1016/j.ccell.2020.12.003 (2021).

Xu, C., Kim, A., Corbin, J. M. & Wang, G. G. Onco-condensates: formation, multi-component organization, and biological functions. *Trends Cancer* **9**, 738-751, doi:10.1016/j.trecan.2023.05.006 (2023).

Mathias, K. M., Liu, Y. & Wan, L. Dysregulation of transcriptional condensates in human disease: mechanisms, biological functions, and open questions. *Curr Opin Genet Dev* **86**, 102203, doi:10.1016/j.gde.2024.102203 (2024).

Han, T. W., Portz, B., Young, R. A., Boija, A. & Klein, I. A. RNA and condensates: Disease implications and therapeutic opportunities. *Cell Chem Biol* **31**, 1593-1609, doi:10.1016/j.chembiol.2024.08.009 (2024).

Wadsworth, G. M. *et al.* RNA-driven phase transitions in biomolecular condensates. *Mol Cell* **84**, 3692-3705, doi:10.1016/j.molcel.2024.09.005 (2024).

Roden, C. & Gladfelter, A. S. RNA contributions to the form and function of biomolecular condensates. *Nat Rev Mol Cell Biol* **22**, 183-195, doi:10.1038/s41580-020-0264-6 (2021).

Henninger, J. E. *et al.* RNA-Mediated Feedback Control of Transcriptional Condensates. *Cell* **184**, 207-225 e224, doi:10.1016/j.cell.2020.11.030 (2021).

Asimi, V. *et al.* Hijacking of transcriptional condensates by endogenous retroviruses. *Nat Genet* **54**, 1238-1247, doi:10.1038/s41588-022-01132-w (2022).

Perlman, E. J. *et al.* MLLT1 YEATS domain mutations in clinically distinctive Favourable Histology Wilms tumours. *Nat Commun* **6**, 10013, doi:10.1038/ncomms10013 (2015).

Hetzner, K., Garcia-Cuellar, M.P., Buttner, C., and Slany, R.K. The interaction of ENL with PAF1 mitigates polycomb silencing and facilitates murine leukemogenesis. *Blood* **131** (2018).

Gadd, S. *et al.* A Children's Oncology Group and TARGET initiative exploring the genetic landscape of Wilms tumor. *Nat Genet* **49**, 1487-1494, doi:10.1038/ng.3940 (2017).

Umeda, M. *et al.* A new genomic framework to categorize pediatric acute myeloid leukemia. *Nat Genet* **56**, 281-293, doi:10.1038/s41588-023-01640-3 (2024).

Bolouri, H. *et al.* The molecular landscape of pediatric acute myeloid leukemia reveals recurrent structural alterations and age-specific mutational interactions. *Nat Med* **24**, 103-112, doi:10.1038/nm.4439 (2018).

Wan, L. *et al.* ENL links histone acetylation to oncogenic gene expression in acute myeloid leukaemia. *Nature* **543**, 265-269, doi:10.1038/nature21687 (2017).

Erb, M. A. *et al.* Transcription control by the ENL YEATS domain in acute leukaemia. *Nature* **543**, 270-274, doi:10.1038/nature21688 (2017).

Wan, L. *et al.* Impaired cell fate through gain-of-function mutations in a chromatin reader. *Nature* **577**, 121-126, doi:10.1038/s41586-019-1842-7 (2020).

Song, L. *et al.* Hotspot mutations in the structured ENL YEATS domain link aberrant transcriptional condensates and cancer. *Mol Cell* **82**, 4080-4098 e4012, doi:10.1016/j.molcel.2022.09.034 (2022).

Liu, Y. *et al.* Condensate-Promoting ENL Mutation Drives Tumorigenesis In Vivo Through Dynamic Regulation of Histone Modifications and Gene Expression. *Cancer Discov* **14**, 1522-1546, doi:10.1158/2159-8290.CD-23-0876 (2024).

Oksuz, O. *et al.* Transcription factors interact with RNA to regulate genes. *Mol Cell*, doi:10.1016/j.molcel.2023.06.012 (2023).

Bose, D. A. *et al.* RNA Binding to CBP Stimulates Histone Acetylation and Transcription. *Cell* **168**, 135-149 e122, doi:10.1016/j.cell.2016.12.020 (2017).

Rahnamoun, H. *et al.* RNAs interact with BRD4 to promote enhanced chromatin engagement and transcription activation. *Nat Struct Mol Biol* **25**, 687-697, doi:10.1038/s41594-018-0102-0 (2018).

Holmes, Z. E. *et al.* The Sox2 transcription factor binds RNA. *Nat Commun* **11**, 1805, doi:10.1038/s41467-020-15571-8 (2020).

Sigova, A. A., Abraham, B.J., Ji, X., Molinie, B., Hannett, N.M., Guo, Y.E., Jangi, M., Giallourakis, C.C., Sharp, P.A., and Young, R.A. . Transcription factor trapping by RNA in gene regulatory elements. *Science* **350**, 978–981, doi:https://doi.org/10.1126/science.aad3346. (2015).

Jeon, Y. & Lee, J. T. YY1 tethers Xist RNA to the inactive X nucleation center. *Cell* **146**, 119-133, doi:10.1016/j.cell.2011.06.026 (2011).

Klein, B. J. *et al.* Structural insights into the pi-pi-pi stacking mechanism and DNA-binding activity of the YEATS domain. *Nat Commun* **9**, 4574, doi:10.1038/s41467-018-07072-6 (2018).

Guo, J. K. *et al.* Denaturing purifications demonstrate that PRC2 and other widely reported chromatin proteins do not appear to bind directly to RNA in vivo. *Mol Cell* **84**, 1271-1289 e1212, doi:10.1016/j.molcel.2024.01.026 (2024).

Liu, Y. *et al.* Small-Molecule Inhibition of the Acyl-Lysine Reader ENL as a Strategy against Acute Myeloid Leukemia. *Cancer Discov* **12**, 2684-2709, doi:10.1158/2159-8290.CD-21-1307 (2022).

Ogami, K. *et al.* RNA Dynamics Regulate Transcriptional Condensate Vivacity to Drive Gene Coordination. *bioRxiv*, doi:10.1101/2025.06.06.658161 (2025).

Bradner, J. E., Hnisz, D. & Young, R. A. Transcriptional Addiction in Cancer. *Cell* **168**, 629-643, doi:10.1016/j.cell.2016.12.013 (2017).

Chandra, B. *et al.* Phase Separation Mediates NUP98 Fusion Oncoprotein Leukemic Transformation. *Cancer Discov* **12**, 1152-1169, doi:10.1158/2159-8290.CD-21-0674 (2022).

Wang, X. Q. D. *et al.* Mutant NPM1 Hijacks Transcriptional Hubs to Maintain Pathogenic Gene Programs in Acute Myeloid Leukemia. *Cancer Discov* **13**, 724-745, doi:10.1158/2159-8290.CD-22-0424 (2023).

Ahn, J. H. *et al.* The phenylalanine-and-glycine repeats of NUP98 oncofusions form condensates that selectively partition transcriptional coactivators. *Mol Cell* **85**, 708-725 e709, doi:10.1016/j.molcel.2024.12.026 (2025).

Ahn, J. H. *et al.* Phase separation drives aberrant chromatin looping and cancer development. *Nature* **595**, 591-595, doi:10.1038/s41586-021-03662-5 (2021).

Terlecki-Zaniewicz, S. *et al.* Biomolecular condensation of NUP98 fusion proteins drives leukemogenic gene expression. *Nat Struct Mol Biol* **28**, 190-201, doi:10.1038/s41594-020-00550-w (2021).

Selig, E. E., Romero-Moreno, A. K., Akula, S., Xu, X. & Libich, D. S. The oncogenic fusion protein EWS-FLI1 promotes premature ageing of biomolecular condensates by catalyzing fibril formation. *bioRxiv*, doi:10.1101/2022.06.04.494830 (2022).

Ng KP, P. G., Savene RO, Denny CT, Uversky VN, Lee KA. Multiple aromatic side chains within a disordered structure are critical for transcription and transforming activity of EWS family oncoproteins. *Proc Natl Acad Sci U S A* **104**, 479-484, doi:10.1073/pnas.0607007104. (2007).

Cheng, Y. *et al.* N(6)-Methyladenosine on mRNA facilitates a phase-separated nuclear body that suppresses myeloid leukemic differentiation. *Cancer Cell* **39**, 958-972 e958, doi:10.1016/j.ccell.2021.04.017 (2021).

Kaikkonen, M. U. & Adelman, K. Emerging Roles of Non-Coding RNA Transcription. *Trends Biochem Sci* **43**, 654-667, doi:10.1016/j.tibs.2018.06.002 (2018).

FIGURES AND FIGURE LEGENDS

Fig. 1

Fig. 1. ENL binds RNA via a conserved basic patch in the YEATS domain. a, Schematic of ENL

wildtype protein sequence and its oncogenic hotspot mutations, T1-T8. Mutations are shown in red text.

YEATS, Yaf9, Enl, Af9, Taf14, Sas5; IDR, intrinsically disordered region; AHD, ANC1 homology domain.

b, Surface charge map of ENL-T1 YEATS domain (YD). Residues from positive patches are labeled. **c,**

Heatmap depicting confidence scores (PAE, predicted aligned error) of predicted RNA-interacting

regions in the ENL-T1 YEATS domain. Predictions generated by Alphafold3. **d,** Percentage of Alphafold3

models in which indicated ENL residue is predicted to be within 4Å of the RNA molecule. *P* values were

calculated using two-tailed Fisher's exact test. **e,** Representative image of EMSA performed with purified

full-length ENL protein and a chemically synthesized 15nt *HOXA11* nascent RNA fragment. **f,** EMSA

quantification of ENL:RNA binding. Data are mean ± s.d. *P* values were calculated using two-tailed

unpaired Student's t test. Representative data shown for *n* = 3 replicates. **g,** Schematic of CLAP-

sequencing workflow. **h,** Average (top) and heatmap (bottom) representations of ENL-bound RNA signal

on merged WT, T1, and T1-3A CLAP peaks. Peak signals are calculated by subtracting input reads per

million (RPM) from those of corresponding CLAP pull-downs. **i,** Box plots showing Log_2 fold changes of

RNA peak signal on merged CLAP peaks. Center line indicates median, and box limits are set to 25^{th} and

75^{th} percentiles. *P* values were calculated using two-tailed paired Student's t test. **j,** Distribution of

merged input or CLAP pull-down RNA peaks across genomic regions. TSS, transcription start site; TTS,

transcription termination site. **k,** Distribution of peak-associated genes from merged ENL CLAP pull-

downs or ENL ChIP-seq. ncRNA, noncoding RNA; snoRNA, small nucleolar RNA; rRNA, ribosomal RNA.

l, RNA signal across gene body regions. RPM, reads per million; TSS, transcription start site; TES,

transcription end site. **m,** Box plots showing RNA signal (CLAP pull-downs relative to inputs) of peak-

associated genes. Center line indicates median, and box limits are set to 25^{th} and 75^{th} percentiles. *P*

values were calculated using two-tailed paired Student's t test. **n,** Genome browser view of ENL-bound

RNA signal in inputs and CLAP pull-downs.

Fig. 2

Fig. 2. Interactions with RNA promote condensate formation by oncogenic ENL.

a, Schematic of *in vitro* droplet formation assay. **b,** *In vitro* droplet formation by purified recombinant GFP-tagged ENL protein with two representative concentrations of chemically synthesized and fluorescently labeled nascent *HOXA11* RNA. Yellow box indicates zoom-in region. **c,** Quantification of total droplet area per image field. Data are mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates. **d,** Total RNA signal (pixel intensity) per ENL droplet. a.u., arbitrary units. Data are mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates. **e,** Western blot showing similar expression of ENL transgenes to each other and to endogenous ENL in HEK293 cells. **f,** Representative IF staining of FLAG-ENL in HEK293 cells. Yellow dashed line outlines the nucleus. **g-j,** Quantification of puncta number per cell in HEK293 (**g**), WiT49 (**h**), A549 (**i**), and MDA-MB-231 (**j**) cells. Data are mean ± s.d. *P* values are calculated by two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates. **k, m,** Representative *HOXA* DNA-FISH and FLAG-ENL IF staining in HEK293 cells. Yellow dashed line outlines the nucleus **l, n,** Quantification of percentage of cells with indicated number of ENL puncta-associated *HOXA* alleles. *P* values were calculated by two-tailed Fisher's exact test. Representative data shown for *n* = 3 replicates. **o, q,** Representative *CBX3* DNA-FISH and FLAG-ENL IF staining in HEK293 cells. Yellow dashed line outlines the nucleus **p, r,** Quantification of percentage of cells with indicated number of ENL puncta-associated *CBX3* alleles. *P* values were calculated by two-tailed Fisher's exact test. Representative data shown for *n* = 3 replicates. **s, t,** ChIP-qPCR analysis of FLAG-ENL in HEK293 cells. Data are mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. Triplicate qPCR values are shown.

Fig. 3

Fig. 3. RNA binding facilitates dynamic re-nucleation of ENL condensates at key target loci after acute chemical displacement. a, Schematic depicting experimental design to measure condensate nucleation. Left panel shows ENL mutant condensates reliably occupying target genes. Middle panel shows ENL inhibitor TDI-11055 dislodging ENL condensates from target genes. Right panel shows TDI-11055 wash-off results in the reformation of ENL mutant condensates on target genes. **b,** Schematic of experimental workflow. **c,** Representative *HOXA* DNA-FISH and FLAG-ENL IF staining in HEK293 cells. Yellow dashed line outlines the nucleus. **d,** Quantification of percentage of total *HOXA* alleles associated with ENL mutant puncta in HEK293 cells. TDI, TDI-11055. *P* values were calculated by two-tailed Fisher's exact test. Representative data shown for *n* = 3 replicates. **e,** RT-qPCR showing nascent *HOXA13* expression relative to *GAPDH*. Data are mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. Triplicate qPCR values are shown. **f,** Schematic depicting experimental design to measure condensate nucleation under transcription inhibition. **g,** Schematic of experimental workflow. **h, j,** Quantification of percentage of total *HOXA* alleles associated with ENL mutant puncta in HEK293 cells. TDI, TDI-11055; ActD, actinomycin D; DRB, 5,6-Dichloro-1-β-D-ribofuranosylbenzimidazole. *P* values were calculated by two-tailed Fisher's exact test. Representative data shown for *n* = 3 replicates. **i, k,** RT-qPCR showing nascent *HOXA13* expression relative to *18s rRNA*. Data are mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates.

Fig. 4

Fig. 4. RNA binding preferentially promotes ENL mutant occupancy at condensate-permissive target loci and enhances transcriptional bursting and gene expression. a, ENL-T1 ChIP-seq peaks in HEK293 cells ranked by read count. Top-bound peaks fall to the right of the inflection point and are labeled in red text. Low-bound peaks fall to the left of the inflection point and are labeled in gray. Known condensate-associated targets, *HOXA* and *CBX3*, are labeled in red text. **b,** Box plot showing Log_2 fold change of T1/WT signal at top or low-bound sites in HEK293. Top, n = 58; Low, n = 5834. Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **c,** Schematic showing at that most ENL-bound sites (low-bound sites), ENL-T1 modestly gains occupancy compared to ENL-WT. At a very small subset of ENL bound sites (top-bound sites), ENL-T1 exhibits a large gain in chromatin occupancy, likely reflecting gain of condensate formation. **d,** Average chromatin occupancy of ENL at top or low-bound sites. RPM, reads per million. **e,** Box plot showing Log_2 fold change of T1-3A vs. T1 signal at top or low-bound sites. Top, n = 58; Low, n = 5834. Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **f,** Genome browser view of ENL ChIP signal at representative HEK293 top and low-bound genes. **g,** ENL-T1 ChIP-seq peaks in MDA-MB-231 cells ranked by normalized read count. Top-bound peaks fall to the right of the inflection point and are labeled in red text. Low-bound peaks fall to the left of the inflection point and are labeled in gray. **h,** Box plot showing Log_2 fold change of T1/WT signal at top or low-bound sites in MDA-MB-231. Top, n = 144; Low, n = 14244. Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **i,** Representative *NR2F2* DNA-FISH and FLAG-ENL IF staining in MDA-MB-231 cells. Yellow dashed line outlines the nucleus. **j, k,** Quantification of percentage of cells with indicated number of ENL puncta-associated *NR2F2* (**j**) or *CHD2* (**k**) alleles. *P* values were calculated by two-tailed Fisher's exact test. Representative data shown for n = 2 replicates. **l,** Average chromatin occupancy of ENL at top or low-bound sites in MDA-MB-231 cells. RPM, reads per million. **m,** Box plot showing Log_2 fold change of T1-3A/T1 signal at top or low-bound sites. Top, n = 144; Low, n = 14244. Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **n,** Genome browser view of ENL ChIP signal at

representative MDA-MB-231 top and low-bound genes. **o, p,** GSEA plots showing that top-bound genes

(determined by ChIP-seq) are upregulated in T1 compared to WT or to T1-3A in HEK293 (**o**) and MDA-

MB-231 (**p**) cells. NES, normalized enrichment score. **q,** RT-qPCR showing nascent *HOXA11* or

HOXA13 expression relative to *GAPDH*. Data are mean ± s.d. *P* values were calculated by two-tailed

unpaired Student's t test. Triplicate qPCR values are shown. **r,** Representative nascent RNA-FISH for the

HOXA13 locus and Halo staining for Halo-ENL in HEK293 cells. White arrows indicate RNA-FISH foci.

Yellow dashed line outlines the nucleus. **s,** Quantification of percentage of cells with indicated number of

nascent RNA-FISH foci. *P* values were calculated by two-tailed Fisher's exact test. Representative data

shown for *n* = 3 replicates. **t,** Quantification of percentage of total *HOXA13* FISH foci associated with

ENL mutant puncta. *P* values were calculated by two-tailed Fisher's exact test. Representative data

shown for *n* = 3 replicates.

Fig. 5

Fig. 5. RNA binding is required for ENL mutant–driven leukemogenesis. a, Experimental work flow

using hematopoietic stem cells (LSK). LSK, Lin$^-$Sca$^+$cKit$^+$. **b,** Representative FLAG-ENL IF staining in

LSK cells. Yellow dashed line outlines the nucleus. **c,** Quantification of puncta number per cell. Data are

mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. **d, f,** Representative *Hoxa*

(**d**) or *Meis1* (**f**) DNA-FISH and FLAG-ENL IF staining in LSK cells. Yellow dashed line outlines the

nucleus. **e, g,** Quantification of percentage of cells with indicated number of ENL puncta-associated

Hoxa (**e**) or *Meis1* (**g**) alleles. *P* values were calculated by two-tailed Fisher's exact test. **h,** Heatmap

depicting z-scores of genes upregulated in T1 vs. WT LSK cells. DEG, differentially expressed genes. **i,**

Log2 fold change in expression of T1 vs. WT upregulated genes. Center line indicates median and red or

blue dotted lines indicate interquartile range. *P* values were calculated by two-tailed paired Student's t

test. **j,** RT-qPCR showing gene expression of key target loci relative to *Gapdh*. Data are mean ± s.d. *P*

values were calculated by two-tailed unpaired Student's t test. Triplicate qPCR values are shown **k,**

GSEA plot showing that ENL-T1 top-bound genes (Extended Data Fig. 9d, *n* = 69) are upregulated in T1

compared to T1-3A in LSK. NES, normalized enrichment score. **l,** GSEA plot showing that genes

associated with AML patients (*n* = 47) are upregulated in T1 compared to T1-3A in LSK. NES, normalized

enrichment score. **m,** Quantification of colonies formed by LSK expressing indicated ENL variants. Data

are mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test. Representative data

shown for *n* = 3 replicates. **n,** Schematic depicting experimental workflow for bone marrow (BM)

transplantation. CD45.1, CD45.2, CD45.1/2 represent leukocyte markers. **o,** Percentage of peripheral

blood (PB) composed of cells derived from transplanted LSK at initial versus terminal timepoints. Each

dot represents *n* = 1 mouse. Horizontal line represents median. *P* values were calculated using two-tailed

unpaired Student's t test. **p,** Kaplan-Meier survival curves of mice receiving bone marrow transplant. *P*

value was calculated using Mantel-Cox log-rank test. WT, *n = 3*; T1, *n* = 4; T1-3A, *n = 4.* **q,**

Representative images (top) and quantification (bottom) of spleen masses harvested from transplanted

mice. Each dot represents *n* = 1 spleen. Horizontal line represents median. *P* values were calculated

using two-tailed unpaired Student's t test. **r,** H&E staining. Black dashed line indicates leukemic blast

infiltration border. **s,** Model (top) ENL:RNA interactions facilitate condensate formation at key target

genes, thus driving frequent transcriptional bursting and persistent activation at top-bound, condensate-

permissive loci, leading to potent leukemogenesis in mouse models. (Bottom) Disrupting these

interactions reduces condensate nucleation, transcriptional bursting, and gene activation, ultimately

impairing tumorigenesis.

Extended Data Fig. 1

Extended Data Fig. 1. ENL binds RNA via a conserved basic patch in the YEATS domain *in vitro*.

a, b, Representative image (**a**) and quantification (**b**) of EMSA performed with purified full-length ENL

protein and chemically synthesized 15nt *HOXA11* dsDNA fragment. Data are mean ± s.d. *P* values were

calculated using two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates. **c,**

d, Representative image (**c**) and quantification (**d**) of EMSA performed with purified full-length ENL

protein and chemically synthesized 15nt *HOXA11* ssRNA fragment. Data are mean ± s.d. *P* values were

calculated using two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates. **e,**

Distribution of nucleotides in 104 RNA sequences used for Alphafold3 modelling. **f,** Sequence alignment

of ENL and ENL paralog, AF9. AF9's DNA-binding residues are outlined in red. **g,** ENL-T1 structure

shows K61, R63, and K67 are surface-exposed, while R64 is less accessible. **h,** Overlay of resolved

ENL-T1 and predicted ENL-T1-3A YEATS domain structures. T1-3A structure predicted with Alphafold3.

i, Percentage of Alphafold3 models in which indicated ENL residue is predicted to be within 3Å or 5Å of

the RNA molecule. *P* values were calculated using two-tailed Fisher's exact test. **j, k,** Representative

image (**j**) and quantification (**k**) of EMSA performed with purified ENL YEATS domain protein and *in vitro*

transcribed 15nt *HOXA11* ssRNA fragment. Data are mean ± s.d. *P* values were calculated using two-

tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates.

Extended Data Fig. 2

Extended Data Fig. 2. ENL binds RNA via a conserved basic patch in the YEATS domain in cells.

a, Western blot showing comparable expression of ENL transgenes to each other and to endogenous

ENL in HEK293 cells. **b,** Fold change values of peak signal within each sample over the average signal

across WT, T1, and T1-3A conditions. **c,** Pearson correlation analysis of CLAP experimental replicates.

Each dot represents the read counts per 5kb genomic bin per chromosome. **d,** Distribution of merged

input RNA peaks across genomic regions. TSS, transcription start site; TTS, transcription termination

site. **e,** Distribution of merged CLAP pull-down RNA peaks across genomic regions. TSS, transcription

start site; TTS, transcription termination site. **f,** Distribution of genes associated with ENL CLAP pull-

down peaks. ncRNA, noncoding RNA; snoRNA, small nucleolar RNA; rRNA, ribosomal RNA.

Extended Data Fig. 3

Extended Data Fig. 3. RNA enhances droplet formation of purified ENL protein. a, *In vitro* droplet

formation by purified recombinant GFP-tagged ENL protein with RNA extracted from whole cell lysate. **b,**

Quantification of total droplet area per image field. Data are mean ± s.d. *P* values were calculated by

two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates. **c,** *In vitro* droplet

formation by purified recombinant GFP-tagged ENL protein with RNA extracted from cell nuclei. **d,**

Quantification of total droplet area per image field. Data are mean ± s.d. *P* values were calculated by

two-tailed unpaired Student's t test. Representative data shown for *n* = 3 replicates.

Extended Data Fig. 4

Extended Data Fig. 4. RNA binding enhances condensate formation by oncogenic ENL at

endogenous target loci. a, Western blot showing similar expression of ENL transgenes to each other

and to endogenous ENL in WiT49 cells. **b,** Representative IF staining of FLAG-ENL in WiT49 cells.

Yellow dashed line outlines the nucleus. **c,** Western blots of cycloheximide (CHX) pulse-chase assays. **d,**

Quantification of cycloheximide pulse chase assay. FLAG-ENL intensity at each time point is normalized

to matched vinculin intensity. **e, g,** Western blot showing similar expression of ENL transgenes to each

other and to endogenous ENL in A549 (**e**) and MDA-MB-231 (**g**) cells. **f, h,** Representative IF staining of

FLAG-ENL in A549 (**f**) and MDA-MB-231 (**h**) cells. Yellow dashed line outlines the nucleus. **i,** Isothermal

calorimetry of purified ENL protein and H3K27ac peptide.

Extended Data Fig. 5

a

b

c

d

Extended Data Fig. 5. RNA binding facilitates dynamic re-nucleation of ENL condensates at

endogenous target loci targets. a, Representative *HOXA* DNA-FISH and FLAG-ENL IF staining in

HEK293 cells. Yellow dashed line outlines the nucleus. **b,** RT-qPCR showing nascent *HOXA11*

expression relative to *GAPDH*. Data are mean ± s.d. *P* values were calculated by two-tailed unpaired

Student's t test. **c, d,** RT-qPCR showing nascent *HOXA11* expression relative to *18S rRNA*. Data are

mean ± s.d. *P* values were calculated by two-tailed unpaired Student's t test.

Extended Data Fig. 6

Extended Data Fig. 6. RNA binding enhances mutant ENL occupancy at condensate-permissive

loci in HEK293 cells. a, Average (top) and heatmap (bottom) representations of ENL ChIP signal on

merged ENL-WT/T1 peaks in HEK293 cells. RPM, reads per million. **b,** Average (top) and heatmap

(bottom) representations of ENL ChIP signal on merged ENL-WT/T2 peaks. RPM, reads per million. **c,**

ENL-T2 ChIP-seq peaks in HEK293 cells ranked by normalized read count. Top-bound peaks fall to the

right of the inflection point and are labeled in red text. Low-bound peaks fall to the left of the inflection

point and are labeled in gray. Known condensate-associated targets, *HOXA* and *CBX3*, are labeled in

red text. **d,** Box plot showing Log_2 fold change of T2/WT signal at top- or low-bound sites in HEK293.

Top, n = 67; Low, n = 10108. Center line indicates median and box limits are set to 25^{th} and 75^{th}

percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **e,** Average chromatin

occupancy of ENL at top or low-bound sites in HEK293 cells. RPM, reads per million. **f,** Box plot showing

Log_2 fold change of T2-3A/T2 signal at top or low-bound sites in HEK293. Top, n = 67; Low, n = 10108.

Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated

by two-tailed unpaired Student's t test. **g,** Genome browser view of ENL ChIP signal at representative

HEK293 top and low-bound genes.

Extended Data Fig. 7. RNA binding enhances mutant ENL occupancy at condensate-permissive target loci in MDA-MB-231 cells. a, Venn diagram showing shared or unique top-bound peaks in HEK293 and MDA-MB-231 cells. **b,** ENL-T2 ChIP-seq peaks in MDA-MB-231 cells ranked by normalized read count. Top-bound peaks fall to the right of the inflection point and are labeled in red text. Low-bound peaks fall to the left of the inflection point and are labeled in gray. **c,** Box plot showing Log_2 fold change of T2/WT signal at top or low-bound sites in MDA-MB-231. Top, n = 301; Low, n = 15,001. Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **d,** Representative *CHD2* DNA-FISH and FLAG-ENL IF staining in HEK293 cells. Yellow dashed line outlines the nucleus. **e, g,** Representative *NR2F2* (**e**) or *CHD2* (**g**) DNA-FISH and FLAG-ENL IF staining in MDA-MB-231 cells. Yellow dashed line outlines the nucleus. **f, h,** Quantification of percentage of cells with indicated number of ENL puncta-associated *NR2F2* (**f**) or *CHD2* (**h**) alleles. *P* values were calculated by two-tailed Fisher's exact test. **i,** Average chromatin occupancy of ENL at top or low-bound sites in MDA-MB-231 cells. RPM, reads per million. **j,** Box plot showing Log_2 fold change of T2-3A/T2 signal at top or low-bound sites in MDA-MB-231. Top, n = 301; Low, n = 15,001. Center line indicates median and box limits are set to 25^{th} and 75^{th} percentiles. *P* values were calculated by two-tailed unpaired Student's t test. **k,** Genome browser view of ENL ChIP signal at representative top- and low-bound genes in MDA-MB-231 cells.

Extended Data Fig. 8

Extended Data Fig. 8. RNA binding enhances transcriptional bursting and gene activation by

oncogenic ENL. a, b, GSEA plots showing that top-bound genes (determined by ChIP-seq) are

upregulated in T2 versus WT or to T2-3A in HEK293 (**a**) and MDA-MB-231 (**b**) cells. NES, normalized

enrichment score. **c,** RT-qPCR showing nascent *NR2F2* or *CHD2* expression relative to *GAPDH* in MDA-

MB-231 cells expressing T1 variants. Data are mean ± s.d. *P* values were calculated by two-tailed

unpaired Student's t test. **d,** RT-qPCR showing nascent top target gene expression relative to *GAPDH* in

HEK293 or MDA-MB-231 cells expressing T2 variants. Data are mean ± s.d. *P* values were calculated by

two-tailed unpaired Student's t test. **e,** Nascent RNA-FISH for the *HOXA13* locus and Halo staining for

Halo-ENL in HEK293 cells. White arrows indicate RNA-FISH foci. Yellow dashed line outlines the

nucleus. **f,** Quantification of percentage of cells with indicated number of nascent RNA-FISH foci. *P*

values were calculated by two-tailed Fisher's exact test. **g,** Quantification of percentage of total *HOXA13*

RNA-FISH foci associated with ENL mutant puncta. *P* values were calculated by two-tailed Fisher's exact

test. **h,** Quantification of RNA FISH foci intensity (normalized to nuclear intensity). a.u., arbitrary units.

Horizontal line represents median. *P* values were calculated by two-tailed unpaired Student's t test.

Extended Data Fig. 9

Extended Data Fig. 9. Disrupting RNA binding impairs ENL mutant–driven leukemogenesis. a, RT-qPCR showing total human (transgene) and murine (endogenous) *ENL* RNA expression in transduced LSK cells. **b,** Mean nuclear FLAG-ENL intensity in LSK cells represents relative protein expression. a.u., arbitrary units. Horizontal line represents median. *P* values were calculated using two-tailed unpaired Student's t test. **c,** Heatmap depicting z-scores of all differentially expressed genes (DEGs). **d,** ENL-T1 ChIP-seq peaks in LSK cells ranked by normalized read count. Top-bound peaks fall to the right of the inflection point and are labeled in red text. Low-bound peaks fall to the left of the inflection point and are labeled in gray. **e,** GSEA plot showing that ENL-T1 top-bound genes are upregulated in T1 versus WT in LSK cells. NES, normalized enrichment score. **f,** GSEA plot showing that genes associated with AML patients ($n = 47$) are upregulated in T1 versus WT LSK cells. NES, normalized enrichment score. **g,** Percentage of peripheral blood (PB) composed of cells derived from transplanted LSK. CD45.2, leukocyte marker of donor LSK. **h,** Kaplan-Meier survival curves of mice receiving bone marrow transplant. *P* value was calculated using Mantel-Cox log-rank test.

METHODS

Cell Lines

Reagent or Resource	Manufacturer/Source	Catalog Number
HEK293T	ATCC	CRL-3216
HEK293	ATCC	CRL-1573
WiT49	Laboratory of Herman Yeger	N/A
A549	Laboratory of Junwei Shi	N/A
MDA-MB-231	Laboratory of Yibin Kang	N/A

Antibodies

Reagent or Resource	Manufacturer	Catalog Number
Brilliant Violet 421™ anti-mouse CD3 Antibody	Biolegend	100227
Brilliant Violet 421™ anti-mouse CD19 Antibody	Biolegend	115549
Brilliant Violet 421™ anti-mouse/human CD11b Antibody	Biolegend	101251
Brilliant Violet 421™ anti-mouse/human CD45R/B220 Antibody	Biolegend	103240
Brilliant Violet 421™ anti-mouse TER-119/Erythroid Cells Antibody	Biolegend	116233
Brilliant Violet 421™ anti-mouse Ly-6G/Ly-6C (Gr-1) Antibody	Biolegend	108433
Brilliant Violet 421™ anti-mouse NK-1.1 Antibody	Biolegend	108741
Brilliant Violet 421™ anti-mouse CD11c Antibody	Biolegend	117343
Brilliant Violet 421™ anti-mouse TCR γ/δ Antibody	Biolegend	118119
Brilliant Violet 421™ anti-mouse TCR β chain Antibody	Biolegend	109229
PerCP/Cyanine5.5 anti-mouse Ly-6A/E (Sca-1)	Biolegend	108124
PE anti-mouse CD117 (c-Kit)	Biolegend	105808
CD45.1 mouse anti-mouse APC/Cy7	Fisher Scientific	BDB560579
CD45.2 mouse anti-mouse PE	Fisher Scientific	BDB560695
Flag	Sigma	F1804
MLLT1/ENL	Millipore Sigma	ABE2596
GAPDH	Cell Signaling Technology	2118
Anti-mouse IgG, HRP-linked Antibody	Cell Signaling Technology	7074S
Anti-rabbit IgG, HRP-linked Antibody	Cell Signaling Technology	7076S
Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Invitrogen	A32731
Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	Invitrogen	A32723
Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 568	Invitrogen	A-11036

Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 568	Invitrogen	A11031
Goat anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 647	Invitrogen	A-21245
Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor 647	Invitrogen	A-21236

Commercial Assays

Reagent or Resource	Manufacturer	Catalog Number
HisTrap HP His tag protein pur columns	Cytiva	29051021
Superdex 75 10/300 GL columns	Cytiva	28920504
Pierce microcentrifuge spin columns	Thermo Fisher	89879
RNA Clean & Concentrator Kit	Zymo Research	R1013
QIAquick PCR Purification Kit	Qiagen	28104
Vazyme Universal DNA Library Prep Kit for Illumina V4	Vazyme	ND610-04
Qubit 1X dsDNA High Sensitivity Kit	Invitrogen	Q33231
Agilent D1000 Screen Tape System	Agilent	N/A
RNeasy Mini Kit	Qiagen	74104
High-Capacity cDNA Reverse Transcription Kit	Applied Biosystems	4368814
NEBNext Poly(A) mRNA Magnetic Isolation Module	New England Biolabs	E7490L
NEBNext Ultra™ II RNA Library Prep Kit for Illumina	New England Biolabs	E7770L
NEBNext Multiplex Oligos for Illumina	New England Biolabs	E7335S
EasySep Mouse CD117 (cKIT) Positive Selection Kit	Stem Cell Technologies	18757

Media, chemicals, and other reagents

Reagent or Resource	Manufacturer	Catalog Number
DMEM with L-Glutamine, 4.5g/L Glucose and Sodium Pyruvate	Corning	MT10013CV
Eagle's Minimum Essential Medium (EMEM)	Corning	MT10009CV
DMEM/F-12	Sigma	D6421
IMDM + GlutaMAX	Gibco	31980030
Opti-MEM	Gibco	31985070
Hank's Balanced Salt Solution (HBSS)	Gibco	14-170-112
Fetal Bovine Serum	Cytiva	SH103IH2540
Penicillin/Streptomycin	Gibco	15070063
Cycloheximide	Cell Signaling Technology	2112S
MethoCult	Stem Cell Technologies	03234
Recombinant murine Flt3 ligand	PeproTech	250-31L
Recombinant murine SCF	PeproTech	250-03
Recombinant murine TPO	PeproTech	315-14
Recombinant murine IL6	PeproTech	216-16
Recombinant murine IL3	PeproTech	200-03
RBC Lysis Buffer (10x)	BioLegend	420301
Zombie Aqua Fixable Viability Kit	BioLegend	423101

SYTOX Blue Dead Cell Stain	Invitrogen	S34857
2-Mercaptoethanol	BioRad	1610710
L-glutamine	Gibco	A2916801
PEI MAX	Polysciences	24765
Polyethylene Glycol 8000 (PEG)	Fisher Scientific	BP233-100
Polybrene	Millipore Sigma	TR-1003-G
Doxycycline	Sigma	D9891-1G
RetroNectin Recombinant Human Fibronectin Fragment	Takara Bio	T100B
Bovine Serum Albumin	Sigma Aldrich	A7906-10G
16% Paraformaldehyde Aqueous	Electron Microscopy Sciences	15710
Tween-20	Fisher Scientific	BP337500
Triton-X-100	Fisher Scientific	BP151500
Goat serum	Cell Signalling Technology	5425s
Mounting Medium with DAPI - Aqueous, Fluoroshield	Abcam	ab104139
PowerUp™ SYBR™ Green Master Mix for qPCR	Applied Biosystems	A25778
Glycine	Fisher Scientific	BP3815
Tris Base	Fisher Scientific	BP152-1
Blotting grade blocker	BioRad	1706404
SDS	Thermo Fisher	BP8200500
Pierce ECL Western Blotting Substrate	Thermo Scientific	32106
Tris-HCl	Fisher Scientific	BP153-500
EDTA	Sigma Aldrich	E5134-500G
NaDOC	Sigma Aldrich	35915
NaCl	Fisher Scientific	BP358-1
Sarkosyl	Sigma Aldrich	61739
DTT	Sigma Aldrich	D9163-5G
cOmplete EDTA-free Protease Inhibitor Cocktail	Sigma-Aldrich	11836170001
Dynabeads Protein A for IP	Invitrogen	10001D
LiCl	Sigma-Aldrich	213233-500G
NP-40	BioVision	2111-100
RNase A	Millipore Sigma	9001-99-4
Proteinase K	Thermo Scientific	EO0491
Poly-L-lysine solution	Sigma Aldrich	P8920
20X SSC buffer	Corning	46-020-CM
Formamide	Fisher Scientific	BP227-500
Dextran Sulfate	Sigma Aldrich	D8906
PVSA solution	Sigma Aldrich	278424
Hoechst 33342 solution, 20 mM	Thermo Scientific	62249
SlowFade Gold Antifade Mountant	Invitrogen	S36936
Glycerol	Fisher Scientific	P2291
RNase inhibitor (Murine)	Beyotime	R0102
10X TBE RNase-free	MREDA	M212499
APS	Sigma Aldrich	7727-54-0
TEMED	Sigma Aldrich	110-18-9
30%Arc-Bis RNase-free	Macklin	A917692
Imidazole	Sigma Aldrich	12399
D-Desthiobiotin	IBA Life Sciences	2-1000-002

Isopropyl β-D-1-thiogalactopyranoside	Sigma Aldrich	10724815001
Kanamycin	Fisher Scientific	BP906-5
Ni IDA Beads 6FF	Smart-Lifesciences	SA00501L
Strep-4FF beads	Smart-Lifesciences	SA053100
Alexa 488 Fluor™ 488 5-TFP	Invitrogen	A30005
HEPES-Na	Sigma-Aldrich	H7006
H3$_{34-28}$K27ac	This paper	SciLight Biotechnology, LLC
5' CY5 HOXA11 ssRNA 15nt/30nt	This paper	hippobio
5' CY5 HOXA11 ssDNA 15nt/30nt	This paper	xianghongbio
DEPC	Sigma-Aldrich	1609-47-8
BL21 (DE3) competent cells	Millipore Sigma	69450-3
HaloLink resin	Promega	G1913
Bovine serum albumin (BSA) fraction V	Sigma-Aldrich	126609-5GM
50X protease inhibitor cocktail	Promega	G6521
TURBO DNase	Invitrogen	AM2238
RNase Inhibitor, murine	New England Biolabs	M0314S
Ambion RNase I	Invitrogen	AM2294
FastAP Thermosensitive Alkaline Phosphatase	Thermo Scientific	EF0651
T4 Polynucleotide Kinase (PNK)	New England Biolabs	M0201S
T4 RNA ligase high concentration, T4 RNA ligase reaction buffer, 50% PEG8000, ATP	New England Biolabs	M0204
Dynabeads MyOne Silane	Invitrogen	37002D
Maxima H Minus Reverse Transcriptase	Thermo Scientific	EP0752
ExoI	New England Biolabs	M0293S
Instant Sticky-end Ligase Master Mix	New England Biolabs	M0370S
AMPure XP beads	Beckman Coulter	A63881
Janelia Fluor 657 (JF657)	Laboratory of Luke Lavis	N/A
Ribonucleoside vanadyl complexes (RVC)	Sigma Aldrich	R3380-5ML
Actinomycin D	Millipore Sigma	A9415-2MG
5,6-Dichlorobenzimidazole 1-β-D-ribofuranoside (DRB)	Millipore Sigma	D1916
Triptolide	Sigma	T3652-1MG
Alpha-Amanitin	Sigma	A2263-1MG

Recombinant DNA

Reagent or Resource	Source	Catalog Number
VSVG	Addgene	8454
PAX2	Addgene	12260
pCL-ECO	Addgene	12371
pLenti-tetON-miniCMV-3xFLAG-2xHA-hsENL-WT-NLS	Song *et al., Mol Cell,* 2022	N/A
pLenti-tetON-miniCMV-3xFLAG-2xHA-hsENL-T1-NLS	Song *et al., Mol Cell,* 2022	N/A
pLenti-tetON-miniCMV-3xFLAG-2xHA-hsENL-T1(K61A/R63A/K67A)-NLS ('T1-3A')	This paper	N/A
pLenti-tetON-miniCMV-3xFLAG-2xHA-hsENL-T2-NLS	Song *et al., Mol Cell,* 2022	N/A

pLenti-tetON-miniCMV-3xFLAG-2xHA-hsENL-T2(K61A/R63A/K67A)-NLS ('T2-3A')	This paper	N/A
pLenti-tetON-miniCMV-Halo-3xFLAG-hsENL-WT-NLS	Song *et al., Mol Cell,* 2022	N/A
pLenti-tetON-miniCMV-Halo-3xFLAG-hsENL-T1-NLS	Song *et al., Mol Cell,* 2022	N/A
pLenti-tetON-miniCMV-Halo-3xFLAG-hsENL-T1(K61A/R63A/K67A)-NLS ('T1-3A')	This paper	N/A
pLenti-tetON-miniCMV-Halo-3xFLAG-hsENL-T2-NLS	Song *et al., Mol Cell,* 2022	N/A
pLenti-tetON-miniCMV-Halo-3xFLAG-hsENL-T2(K61A/R63A/K67A)-NLS ('T2-3A')	This paper	N/A
MSCV-3xFLAG-HA-hsENL-WT-PGK-puro-IRES-GFP	Liu *et al., Cancer Discov,* 2024	N/A
MSCV-3xFLAG-HA-hsENL-T1-PGK-puro-IRES-GFP	Liu *et al., Cancer Discov,* 2024	N/A
MSCV-3xFLAG-HA-hsENL-T1(K61A/R63A/K67A)-PGK-puro-IRES-GFP	This paper	N/A
pET28a-2*strep-ENL-WT-6*his	Song *et al., Mol Cell,* 2022	N/A
pET28a-2*strep-ENL-T1-6*his	Song *et al., Mol Cell,* 2022	N/A
pET28a-2*strep-ENL-T1(K61A/R63A/K67A)-6*his	This paper	N/A
pET28a-6*his-ENL-WT(1-148)	Song *et al., Mol Cell,* 2022	N/A
pET28a-6*his-ENL-T1(1-151)	Song *et al., Mol Cell,* 2022	N/A
pET28a-6*his-ENL-T1(K61A/R63A/K67A)(1-151)	This paper	N/A

Software and algorithms

Reagent or Resource	Manufacturer	Catalog Number
FIJI	Schindelin *et al.,* 2012	imagej.net/software/fiji/
GraphPad Prism	GraphPad	graphpad.com
Imaris 9.3.1	Oxford Instruments	imaris.oxinst.com
Origin 7.0	OriginLab	originlab.com
PyMOL 3.1	PyMOL	pymol.org
Alphafold3 Server	N/A	PMID: 38718835
FlowJo	BD Biosciences	flowjo.com
GSEA_4.3.2	N/A	PMID: 16199517; PMID: 12808457

Protein purification
For full-length ENL protein purification, N-terminal 2 X streptavidin tags and C-terminal 6 X histidine tags
were added to the pET28b expression vector containing the ENL-WT CDS (residues 1-559;
NP_005925.2). The T1 mutation (p. 117_118insNHL) and RNA-binding residue mutations (K61A, R63A,
and K67A) were subsequently added. Proteins were expressed in BL21(DE3) *E. coli* cells induced with
0.4mM IPTG (Isopropyl β-D-thiogalactopyranoside) for 16 h at 16°C. Cells were harvested and
resuspended in lysis buffer (25 mM Tris-HCl pH 7.5, 500 mM NaCl, 5% glycerol, 50 mM imidazole, 1 mM
PMSF). After lysis using an Emulsiflex C3 (Avestin) high-pressure homogenizer, crude lysate was clarified
by centrifugation. Clarified lysate was loaded onto a HisTrap column and washed with wash buffer (30 mM
Tris-HCl pH 7.5, 1 M NaCl, 5% glycerol). Protein was eluted using lysis buffer supplemented with 500 mM

Imidazole, concentrated, and buffer-exchanged to strep buffer (25 mM HEPES-Na pH 7.4, 500 mM NaCl, 5% glycerol, 1 mM EDTA, 2 mM β-mercaptoethanol) to decrease the imidazole concentration. After buffer exchange, the protein was incubated with strep-4FF beads, washed with strep buffer, and eluted using strep buffer supplemented with 5 mM desthiobiotin.

For purification of the ENL YEATS (residues 1-148) domain, N-terminal 6 X histidine tags were added to the pET28b vector containing the ENL-WT YEATS domain CDS (residues 1-148). The T1 mutation (p. 117_118insNHL) and RNA-binding residue mutations (K61A, R63A, and K67A) were subsequently added. Proteins were expressed in BL21(DE3) *E. coli* cells induced with 0.4mM IPTG for 16h at 16°C. Cells were harvested and resuspended in lysis buffer. After lysis using an Emulsiflex C3 high-pressure homogenizer, crude lysate was clarified by centrifugation. Clarified lysate was loaded onto a HisTrap column and washed with wash buffer (30 mM Tris-HCl pH 7.5, 1 M NaCl, 5% glycerol). Protein was eluted with a linear imidazole gradient from 20 mM to 500 mM. Protein was further purified and polished over a Superdex 75 10/300 gel filtration column using AKTA Purifier 10 systems (GE Healthcare). The monomer peak in elution buffer (500 mM NaCl, 25 mM Tris-HCl pH 7.5, and 2 mM β-mercaptoethanol) was pooled, concentrated to about 10 mg/mL, and stored at −80°C.

AlphaFold predictions and analysis

We used the Alphafold-Multimer (v3) to predict the structure of the ENL YEATS-RNA complex. The human ENL 1-142 T1 sequence was submitted as the protein sequence, and the first 15 nucleotides of 104 randomly selected transcripts were chosen as input RNA sequences (Supplementary Table 1). Each model was predicted using one ENL YEATS domain and one RNA fragment. For all predicted results, the top-scoring relaxed model was selected, with an average iPTM score > 0.6 being inferred as putative hits. Diagnostic plots (PAE plot, PLDDT plot, and sequence coverage) and generated structures were manually checked. The regions characterizing the interaction between ENL and RNA were extracted from the PAE matrix, and the minimum PAE corresponding to each ENL residue was calculated and normalized. The PAE values corresponding to each residue were summed and mapped onto the structure for visualization.

To examine the impact of the 3A mutation on ENL:RNA interactions, we performed the same Alphafold modelling with the T1-3A YEATS domain sequence and the 104 RNA sequences. For the 104 T1:RNA models and 104 T1-3A:RNA models, we measured the shortest possible distance between each of ENL's putative RNA binding residues (K61, R63, and K67) and the RNA molecule. We then plotted the percentage of models that show each individual residue as being within 3, 4, or 5 Å of the RNA molecule.

Electrophoretic mobility shift assay (EMSA)

The nucleic acids used in the experiment were all 5' cy5 labeled. ssRNA was first incubated at 95°C for 5 minutes and then quickly cooled on ice for 5 minutes. RNA refolding buffer (10 mM Bis-Tris pH 6.7, 50 mM KCl, 10 mM $MgCl_2$; 1X final) was then added to the RNA on ice. To refold, the RNA was transferred to a cold metal block and folded by warming to room temperature (~25°C) for 30 minutes. For dsDNA, the cy5 labeled DNA oligo was mixed with an equal amount of unlabeled complementary oligo in DNA annealing buffer (10 mM Tris-HCl pH 8.0, 50 mM NaCl, 1 mM EDTA,1 X final), heated at 90°C for 3 minutes, and then cooled slowly to room temperature. The prepared nucleic acids were incubated with purified ENL protein in RNase-free ultra-low adhesion PCR tubes. The reactions contained 1 X binding buffer (1x PBS with 5% glycerol, 2 mM β-ME, and 0.4 U/μL Murine RNase inhibitor (Beyotime)). The reactions were loaded on a 6% native polyacrylamide gel (29:1 Acrylamide: Bis-acrylamide) and electrophoresed at 100 V for 1 hour in 0.5x TBE. The gel was imaged and photographed using a Tanon gel imaging system with a 605 nm light source. The experiment was performed with at least three technical replicates and three biological replicates. All reagents and equipment used in the experiment were RNase-free.

***In vitro* droplet formation assay**

Droplet formation assay was performed in the droplet formation buffer (1X PBS with 5% glycerol and 2 mM β-ME). Proteins were fluorescently labeled for microscopy using Microscale Protein Labeling kits from Thermo Scientific: Alexa Fluor 488 (A30005). Synthetic nucleic acids labeled with 5' cy5 were used in the experiment. Labeled proteins were added to unlabeled proteins at a 1:10 molar ratio. Droplets were

assembled in 384 low-binding multi-well 0.17-mm microscopy plates (384-well microscopy plates) (In Vitro Scientific) and sealed with optically clear adhesive film. For the groups with RNase treatment, RNase A powder is pre-dissolved in droplet formation buffer. After ENL and RNA have been incubated in droplet formation buffer, RNase A is added to achieve a final concentration of 1 µg/mL and incubated at room temperature for 5 minutes. All reagents used in the experiment were RNase-free. After quick centrifugation, droplets were imaged under Olympus FV1200 using a 100 X oil objective. The results were processed by Imaris 9.3.1 (Oxford Instrument). Total droplet area was defined by and plotted as the total area of a field occupied by droplets. Total RNA signal per droplet is defined by the sum of RNA channel pixel intensities within a mask drawn over each individual droplet.

Isothermal titration calorimetry (ITC)
All ITC titrations were performed at 25°C using MicroCal iTC200 system (Malvern). Both synthetic histone H3 peptides (H3$_{24-28}$K27ac) and the recombinant ENL YEATS domain proteins were extensively dialyzed against ITC buffer (25 mM Tris-HCl pH 7.5, 500 mM NaCl, 5 % glycerol and 2 mM β-ME). Before titration, all samples were centrifuged at 13,000 rpm at 4°C for ~10-15 minutes to remove potential aggregates. ITC cells were rinsed with buffer for several times before protein solution was added to the cell. Cell contents were pipetted several times to mix with any trace buffer remaining in the cell. A small volume of protein solution was removed from the cell for concentration re-check using a NanoDrop. Each ITC titration consisted of 17 successive injections with 0.5 µL for the first and 2.4 µL for the rest. The intervals between injections were 150 seconds, and the stirring speed was 750 rpm. Histone H3 peptides (2.0-2.2 mM) were titrated into ENL YEATS domain proteins (0.15 mM) in all experiments. The resultant ITC curves were processed using the Origin (v.7.0) software (OriginLab) in accordance with the "One Set of Sites" fitting model. First injection data was always excluded from the analysis.

Human cell lines
All cell lines used in this study were cultured in a humidified incubator maintained at 37°C and 5% CO$_2$ and regularly tested mycoplasma negative. All cell lines not purchased from ATCC were subjected to STR authentication testing prior to any experimentation. HEK293T, A549, MDA-MB-231 were cultured in DMEM supplemented with 10% FBS and 100 U/mL penicillin/streptomycin. HEK293 cells were cultured in EMEM supplemented with 10% FBS and 100 U/mL penicillin/streptomycin. WiT49 cells were cultured in DMEM/F-12 supplemented with 10% FBS and 100 U/mL penicillin/streptomycin.

Lentivirus and retrovirus generation
HEK293T cells were used for all virus packaging. For lentivirus generation, cells were seeded 1 day prior to transfection. For transfection, lentiviral packaging plasmids VSVG and PAX2 were combined with the plasmid of interest and Polyethylenimine MAX (PEI MAX) in Opti-MEM at a ratio of 1:4, total DNA (ug): PEI (uL). These components were incubated to allow for complex formation and added to the HEK293T cells. ~6h after adding the mix, transfection media was removed and replaced with fresh culture media. The next day, virus was collected and fresh culture media was added back to the cells. A total of 5 virus collections occurred over the course of 72h post-transfection. For retrovirus generation, cells were seeded 1 day prior to transfection. For transfection, retroviral packaging plasmid pCL-ECO was combined with the plasmid of interest and Polyethylenimine MAX (PEI MAX) in Opti-MEM at a ratio of 1:3, total DNA (ug): PEI (uL). These components were incubated to allow for complex formation and added to the HEK293T cells. ~6h after adding the mix, transfection media was removed and replaced with fresh culture media. The next day, virus was collected and fresh culture media was added back to the cells. A total of 5 virus collections occurred over the course of 72h post-transfection. When necessary, virus was concentrated 100x with 5x PEG8000 solution.

Human cell line viral transduction and transgene expression
HEK293 and WiT49 stable cell lines were generated by incubating cells (~70% confluent) with concentrated lentivirus and 10 ug/mL polybrene overnight (~16h). A549 and MDA-231 stable cell lines were generated by adding concentrated lentivirus and 10 ug/mL polybrene to cells (~70% confluent), performing spinfection (2500 rpm for 1h), and incubating overnight. When applicable, selection regents were added 24-48 hours post-infection. Cells containing doxycycline-inducible transgenes (i.e., ENL) were

741 incubated for 48h with a pre-determined concentration of doxycycline such that exogenous ENL expression
742 was similar to that of endogenous ENL as measured by protein levels in Western blot.
743
CLAP pull-down and RNA isolation
CLAP-sequencing was conducted according to published protocols from the Dr. Mitchell Guttman lab[34]
with some modifications. For CLAP experiments, HEK293 cells stably expressing doxycycline-inducible
Halo-tagged ENL transgenes were used (~50-60M cells used per replicate per condition). Cells were
treated with pre-determined concentrations of doxycycline for 48h prior to experimentation. Plates of
adhered cells were UV crosslinked in PBS on ice at an energy setting of 400 mJ/cm^2. Immediately following
crosslinking, cells were harvested, washed with ice cold PBS, centrifuged, and stored as cell pellets at -
80°C.
752
Prior to performing pull-down, HaloLink resin was washed three times in PBST (PBS + 0.1% Tween-20)
and blocked in 1% nuclease-free BSA (fraction V) for 20 minutes at room temperature. Resins were then
washed three more times in PBST. Crosslinked cell pellets were incubated for 10 minutes on ice in lysis
buffer (RIPA buffer (1M HEPES pH 7.4, 5M NaCl, 10% deoxycholate, 20% SDS, 10% NP-40), 1X Promega
protease inhibitor cocktail, 1X TURBO DNase, 1x Mn^{2+}/Ca^{2+} mix (1M MnCl$_2$, 1M CaCl$_2$), and Murine RNase
inhibitor). Lysates were transferred to 1mL Bioruptor sonication tubes and sonicated on the "low setting"
for 5 minutes (30s on/30s off) at 4°C. Following sonication, TURBO DNase and diluted RNase I (1:25 in
PBS) were added to lysates and incubated at 37°C for 5 minutes with shaking at 1200 rpm. Immediately
then murine RNase inhibitor was added to quench the reaction and lysates were then centrifuged at 15,000
rcf for 10 minutes at 4°C. ~5% of the supernatant was reserved in -80°C to use later as an input control.
The remaining lysate was added to the prepared HaloLink resin and incubated at 4°C overnight with
rotation.
765
The next day, the following five CLAP wash buffers were prepared fresh and pre-warmed to 80°C prior to
use: CLAP wash buffer 1 (20% N-lauroylsarcosine, 500mM EDTA, PBS); CLAP wash buffer 2 (1M HEPES
pH 7.4, 5M NaCl, 10% NP-40, water); CLAP wash buffer 3 (1M HEPES pH7.4, 10% NP-40, 8M urea);
CLAP wash buffer 4 (1M HEPES pH 7.4, 10% NP-40, 10% Tween-20, water); CLAP wash buffer 5 (1M
HEPES pH 7.4, 10% NP-40, 500mM EDTA, water). Overnight captures were centrifuged and supernatants
were discarded. Resins were washed twice with RIPA buffer. Resins were then washed three times each
with each CLAP wash buffer in numerical order. For each wash, buffer was added to resins and incubated
at 90°C for 3 minutes with shaking at 1200 rpm before centrifugation and removal of supernatant. Final
resin pellets, as well as the reserved input samples, were then subjected to proteinase K elution by
incubating with CLAP wash buffer 1 and proteinase K at 50°C for 20 minutes with shaking at 1200 rpm.
The slurry was then transferred to Pierce micro-centrifuged columns to remove the resin. Supernatants
were then used for RNA extraction with the Zymo RNA clean and concentrator kit. Eluted RNA was used
for subsequent library preparation.
779
CLAP library preparation
Input samples were first incubated with 1x TURBO DNase buffer and TURBO DNase at 37°C for 15
minutes with shaking at 1200 rpm and subsequently subjected to RNA purification again using the Zymo
clean and concentrator kit. Eluted input RNA was then mixed with 1X FastAP buffer, fragmented at 91°C
for 90s, and incubated with FastAP enzyme and murine RNase inhibitor for 10 minutes at 37°C with
shaking at 1200rpm. At the same time, CLAP samples were mixed with 1X FastAP buffer, FastAP enzyme,
TURBO DNase, and murine RNase inhibitor, and incubated for 10 minutes at 37°C with shaking at
1200rpm. Then, for all input and CLAP samples, PNK mix (1X PNK buffer, PNK enzyme, TURBO DNase)
was added on top of the FastAP reaction and incubated for an additional 10 minutes at 37°C with shaking
at 1200rpm. RNA was then purified using the Zymo RNA clean and concentrator kit.
790
791 To each sample, we then added the RIL19 adaptor (5'–
792 /Phosphate/rArGrArUrCrGrGrArArGrArGrCrGrUrCrGrUrG/3ddC/ – 3') in DMSO and incubated at 65°C for
793 2 minutes. Then ligation mix (1X T4 RNA ligase buffer, ATP, PEG8000, murine RNase inhibitor, and T4

RNA ligase high concentration) was added and incubated at room temperature for 1 hour and 15 minutes with rotation. Adaptor-ligated RNA was then cleaned with SILANE beads.

Eluted RNA was then incubated with reverse transcription adaptor AR17 (5'-ACACGACGCTCTTCCGA-3') at 65°C for 2 minutes. Reverse transcription mix (1X Maxima RT buffer, dNTPs, murine RNase inhibitor, Maxima RT enzyme) was added to the adaptor ligated RNA and incubated at 50°C for 20 minutes. Samples were then incubated with ExoI at 37°C for 15 minutes and quenched with EDTA on ice. To fragment RNA, the samples were incubated with NaOH at 80°C for 6 minutes and neutralized with equal amount of HCl. Reverse transcription product was then cleaned with SILANE beads. The resulting cDNA was then mixed with a splint adaptor (detailed in the published protocol) and 1X Instant Sticky Master Mix and incubated for 4 hours at room temperature with shaking at 1000 rpm. Each reaction was then cleaned with SILANE beads.

With the eluted cDNA samples we then performed PCR with NEBNext Universal primers for Illumina, NEBNext Index primers for Illumina, and 1X Q5 master mix. The following PCR conditions were used: (1X) 98°C 1 minute; (4X) 98°C 15s, 69°C 15s, 72°C 90s; (4X for inputs, 10X for IP) 98°C 15s, 72°C 90s; (1X) 72°C 2 min). PCR products then underwent 1.2X SPRI bead size selection and elution. Concentrations of the final barcoded libraries were measured on the Qubit and size measurements were calculated using the Agilent Tapestation prior to sequencing on a NextSeq2000.

CLAP-seq data processing and analysis
The mapping of CLAP-sequencing was conducted according to published protocols from the Dr. Mitchell Guttman lab[34]. Adapters were trimmed from the paired end sequencing reads using cutadapt v4.9 from Trim Galore!. The RNA alignment was performed in two parts. First, the reads were aligned to hg38 genome reference containing both repetitive and structure RNAs using bowtie2. Second, the remaining reads were recycled and aligned to hg38 with STAR. After alignment, MarkDuplicates function from the Picard package was used to remove any PCR duplications. Finally, alignments from both mapping methods were finally merged into a single bam file for future analysis. CLAP peaks were identified by MACS2 peakcalling with IP and input from each condition using macs2 callpeak -t -c -f BAMPE -g hs -n --nomodel --broad --broad-cutoff 1e-10 -p 1e-10 --keep-dup all. Peak distribution and associated genes were identified using HOMER module annotatePeaks.pl v4.11. NGS plots were generated use ngs.plot.r v2.61.

Western blot
Cells were collected, washed once in PBS, and boiled (95°C) in 1X SDS loading buffer (5X SDS loading buffer: 0.25M Tris-HCl (PH6.8), 0.5M DTT, 10% SDS, 50% glycerol, 0.25% BPB, 1% b-mercaptoethanol (b-ME) for 20 minutes. Lysates were run on homemade 10% SDS-PAGE gels in homemade 1X running buffer (2.5mM Tris Base, 19.2 mM Glycine, 0.01% SDS in H2O, pH 8.3). Protein was transferred to PVDF membrane (Millipore, 0.45 um) using freshly made homemade transfer buffer (1.2 mM Tris Base, 9.6 mM Glycine in H2O, pH 8.3) + 20% methanol on ice. After transferring, membranes were blocked with 5% blocking buffer in PBST (PBS + 0.1% Tween) for 45 minutes at room temperature. Membranes were then incubated with primary antibody suspended in 1% blocking buffer overnight at 4°C with rocking. The next day, membranes were washed three times for 10 minutes each wash with PBST. Membranes were then incubated with secondary antibody suspended in 1% blocking buffer for 1 hour at room temperature and subsequently washed three more times in PBST as before. Blots were covered with Pierce ECL Western Blotting Substrate according to the manufacturer's instructions and imaged using film. Primary antibodies were diluted as follows: Flag (1:1000), MLLT1/ENL (1:1000), GAPDH (1:5000). Secondary antibodies were used as follows: Anti-mouse or Anti-rabbit IgG HRP-linked (1:2500).

Cycloheximide pulse chase
HEK293 cells expressing Flag-tagged ENL transgenes were incubated with doxycycline for 48h prior to experimentation. Cells were treated with 50 ug/mL cycloheximide for and collected at indicated timepoints. The intensity of each sample's ENL blot was normalized to the intensity of that sample's GAPDH blot. Intensities were measured in FIJI.

Immunofluorescence staining and confocal microscopy

Cells (150K/well) were seeded on 24-well plates containing coverslips and incubated 24-48h. Cells were fixed with 4% PFA in PBS for 10 min at room temperature, washed with PBST (PBS + 0.1% Tween-20) three times (5 min/wash), and then permeabilized with PBS containing 0.1% Triton-X 100 for 10 min, and washed three times with PBST as before. Subsequently, cells were blocked in PBS with 10% goat serum for 30 min at room temperature. Cells were then incubated overnight at 4°C with primary antibodies (1:300) in 1% BSA in PBS. The next day, cells were washed three times in PBST, incubated with secondary Alexa Fluor-conjugated antibodies (1:250) in 1% BSA for 1h at room temperature, and washed three more times in PBST. The coverslips were then mounted face down on slides in DAPI-containing mounting medium. Primary antibodies used: Flag (Sigma F1804) and MLLT1/ENL (Millipore ABE2596). Secondary antibodies used: goat anti-Rabbit or anti-Mouse IgG (H+L) conjugated with Alexa Fluor 488, 568, or 647 (Invitrogen). Images were captured on Zeiss LSM880 or Leica Stellaris 5 confocal microscopes.

DNA-FISH with immunofluorescence staining

Poly-L-lysine was first added to wells of rubber gaskets adhered to glass microscopy slides and removed/dried prior to cell plating. Cells were then plated in the gasket wells and allowed to adhere at 37°C for 1.5-2h (HEK293 and MDA-MB-231cells) or 20 minutes (LSK cells). Slides were then fixed in 4% PFA for 10 minutes at room temperature in coplin jars followed by three 10 minute washes in room temperature (RT) PBS. **DNA-FISH Part 1**: Slides were permeabilized in 0.5% Triton-X-100 in PBS for 15 minutes at RT and washed once in PBS for 5 minutes at RT. Next slides were incubated in 70% EtOH, 90% EtOH, and 100% EtOH for 2 minutes each time. Slides were then washed once in 2X SSCT (1X SSC + 0.1% Tween-20) for 5 minutes at RT and incubated once in 70% FMM buffer (70% formamide, 1X SSC, water) for 5 minutes at RT. Slides were then transferred to 70% FMM buffer pre-heated to 37°C for 1 h. Then primary FISH probe hybridization solution (50pmol primary probe, 100uM dNTP, 50% FMM, 1X dextran-sulfate PVSA hybridization mix, 10ug RNase A, water) was added to each slide, sealed with a coverslip and rubber cement, and denatured on a water-immersed heat block for 30 minutes at 83°C. Slides were then transferred to a humidity chamber and incubated overnight at 37°C. **DNA-FISH Part 2:** The next day, slides were washed once in 2X SSCT for 10 minutes at 60°C, once in 2X SSCT for 10 minutes at RT, and once in 0.2X SSC for 10 minutes at RT. Slides were then incubated with secondary FISH probe for 2 h at RT protected from light under coverslips sealed with rubber cement (secondary probe mix: 10pmol secondary probe, 10% FMM, 1X dextran-sulfate PVSA hybridization mix, water). Slides were then washed again in 2X SSCT 60°C, 2X SSCT RT, and 0.2X SSC as before. **IF Part 1:** Slides were next fixed again in 2% PFA for 10 minutes at RT, washed once in PBS for 5 minutes at RT, permeabilized with 0.1% Triton-X-100 for 15 minutes at RT, and washed three times in PBST (PBS + 0.1% Tween-20) for 10 minutes each at RT. Slides were then blocked in 1% BSA in PBST for 45 minutes at RT. Then primary antibody solution (1:300 primary antibody in 1% BSA) was added to each slide, sealed with a coverslip and rubber cement, and incubated overnight at 4°C in a humidity chamber. **IF Part 2:** The next day, slides were washed three times in PBST for 10 minutes each time. Then secondary antibody solution (1:250 secondary antibody in 1% BSA) was added to each slide, sealed with a coverslip and rubber cement, and incubated for 2 h at RT. Slides were washed again three times in PBST for 10 minutes each time. Finally, slides were incubated in DAPI staining buffer (2X SSC + 1:10,000 Hoescht stain) for 5 minutes at RT and washed once in PBS for 5 minutes at RT. Coverslips were added to each slide with anti-fade mounting media and sealed with clear nail polish.

Primary probes used: Probes were designed to cover ~100kb regions (4 40mer probes per kb) of the human *HOXA9-13* locus, human *CBX3* locus, human *NR2F2* locus, human *CHD2* locus, mouse *Hoxa9-10* locus, or mouse *Meis1* locus. Secondary probes were dye conjugated. Primary antibodies used: Flag for HEK293 and MDA-MB-231 cells (Sigma F1804) and MLLT1/ENL (Millipore ABE2596) for LSK cells. Secondary antibodies used: goat anti-Rabbit or anti-Mouse IgG (H+L) conjugated with Alexa Fluor 488, 568, or 647 (Invitrogen). Images were captured on Zeiss LSM880 or Leica Stellaris 5 confocal microscopes.

ChIP-qPCR and ChIP-sequencing

Between 20-25M cells were used per ChIP. Cells were collected from plates and fixed in freshly made 1% PFA in PBS (1mL 1% PFA per 10M cells). For Flag ChIP, fixation was performed for 10 minutes at room temperature with intermittent inversion. PFA was quenched with 125mM Glycine for 5 minutes at room temperature and the cell pellet was then washed three times with PBS. Cells were then resuspended in RIPA3.0 buffer (0.1% SDS, 1% Triton-X-100, 10mM Tris-HCl pH7.4, 1mM EDTA pH8, 0.1% NaDOC, 0.3M NaCl) freshly supplemented with 1/40 volume of 10% sarkosyl, 1mM DTT, and protease inhibitor. For MDA-MB-231 ChIPs, an extra 0.1% SDS (final 0.2% total) was added to the sonication buffer to improve sonication. Cells were sonicated on a Covaris Ultrasonicator. Concurrently, primary antibody was incubated with Protein A Dynabeads and then washed three times with PBS + 0.1% Tween-20. A small volume of sonicated lysate was reserved as an input control. Remaining lysate was rotated overnight at 4°C with the primary antibody-bound Dynabeads. For MDA-MB-231 ChIPs, sonicated lysates (which contain extra SDS in the sonication buffer) were diluted with non-SDS containing sonication buffer to a final concentration of 0.1% SDS before adding to the antibody-bound beads. While extra SDS improved sonication, this SDS can interfere with IP and must be re-diluted before IP. The next day, beads were washed three times with low salt buffer (150mM NaCl, 0.1% SDS, 1% Triton-X-100, 1mM EDTA pH8, 50mM Tris-HCl pH8), three times with high salt buffer (500mM NaCl, 0.1% SDS, 1% Triton-X-100, 1mM EDTA pH8, 50mM Tris-HCl pH8), and three times with LiCl buffer (150mM LiCl, 0.5% NaDOC, 0.1% SDS, 1% NP-40, 1mM EDTA pH8, 50mM Tris-HCl). Beads were then washed once with TE buffer (1mM EDTA, 10mM Tris-HCl pH8) + 50mM NaCl. Elution buffer (1% SDS, 50mM Tris-HCl ph8, 10mM EDTA, 200mM NaCl) was then added to the beads and incubated at 65°C with shaking for 0.5h. Beads were then centrifuged and supernatant was transferred to a new tube and incubated overnight at 65°C. At this point elution buffer was also added to reserved input lysate and incubated overnight at 65°C. The next day, the samples were first incubated with RNase A for 1h and 37°C and then with Proteinase K for 1h at 55°C. DNA was purified using the Qiagen PCR purification kit and eluted in PCR water.

For qPCR, ChIP DNA was diluted 50x in water and qPCR was performed using SYBR green master mix. Primers used are listed in Supplementary Table 11. For ChIP-sequencing, ChIP DNA was subjected to library preparation using the Vazyme Universal DNA Library Prep Kit for Illumina. Prior to sequencing, libraries were quantified and size confirmed using the Qubit and Tapestation according to the manufacturers' instructions. Final libraries were sequenced on an Illumina Nextseq 2000.

Reads were aligned to the human reference genome (hg19) by the package Bowtie2 v2.5.4 with the default parameters. The package samtools v1.9 was used for converting to BAM format, sort- ing, removing duplicates, and indexing. In ChIP-seq analysis, we utilized the corresponding input data as a control during peak calling. The chromatin distribution of the peaks was determined by HOMER module annotatePeaks.pl v 4.11. Peaks are listed in Supplementary Tables 2, 4). To make the heatmaps, the filtered, sorted, and indexed BAM files were converted to bigwig files using the bamCoverage from deeptools v3.5.6 and then normalized by counts per million (CPM). Heatmaps were generated using the module computeMatrix and plotHeatmap from deeptools v3.5.6.

RNA extraction, cDNA synthesis, and RT-qPCR
Total RNA was extracted using Qiagen RNeasy kits according to the manufacturer's instructions. Reverse transcription was performed using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to the manufacturer's protocol. Quantitative real-time PCR was performed using SYBR green master mix. Primers used are listed in Supplementary Table 11.

RNA-sequencing and Gene Set Enrichment Analysis (GSEA)
RNA was extracted as described above. Library preparation was performed with 500 ng total RNA beginning with the NEB polyA mRNA magnetic isolation module followed by NEB library preparation kit, both according to the manufacturer's instructions. Prior to sequencing, libraries were quantified and size confirmed using the Qubit and Tapestation according to the manufacturers' instructions. Final libraries were sequenced on an Illumina Nextseq 2000.

Reads were aligned to the mouse reference genome (mm10) using HISAT2 v 2.2.1 with default
parameters. The package featureCounts v2.0.2 was used for counting the mapped reads in each mouse
gene. Transcript per million (TPM) was used for normalizing the gene expression. Differentially expressed
genes between conditions were statistically determined by R package DESeq2 v1.38.3. Genes with
adjusted *P* value < 0.05 and fold change ≥ 1.5 were reported as differential genes. Differentially expressed
genes in LSK are listed in Supplementary Table 9.
960
961 For gene set enrichment analysis, custom gene signatures were made from our ChIP-sequencing datasets
962 (Supplementary Tables 5, 6, 8). Briefly, normalized read counts for ENL-T1 or -T2 peaks were ranked and
963 the inflection point determined each peak to be top or low-bound. Top-bound peaks were annotated with a
964 corresponding gene and this list of top-bound genes was used as the gene signature. DESeq2 files used
965 in the analysis were pre-ranked. GSEA was performed with the GSEA_4.3.2 desktop application.
966
RNA-FISH with Halo staining
HEK293 cells expressing Halo-tagged ENL transgenes were seeded on poly-L-lysine coated coverslips
and grown for 48h in the presence of doxycycline. Live cells were then incubated with 250 nM JF657 Halo
ligand for 30 minutes and washed five times in pre-warmed media (5 minutes per wash). Cells were then
washed once in PBS + 1% RVC (ribonucleoside vanadyl complex), fixed in 4% PFA + 1% RVC for 10
minutes at room temperature, and then washed twice more in PBS 1% RVC. The cells were then
permeabilized in 70% ethanol and 1% SDS overnight at 4°C. The next day, the ethanol/SDS solution was
removed and the cells were washed twice with 2xSSCT (2xSSC + 0.1% Tween-20) + 1% RVC at room
temperature for 5 minutes per wash. Primary probe hybridization mix (2x DSM (dextran sulfate PVSA),
50% formamide, 1% SDS, 30 nM primary probe, 1% RVC) was assembled and added dropwise onto a
parafilm sheet sitting on top of a damp paper towel in a petri dish. Coverslips were lifted and placed cell-
side down onto the hybridization mix drops and denatured at 70°C for 4 minutes. The apparatus was then
incubated at 42°C overnight. The following day, the coverslips were transferred back to 24-well plates and
washed twice with 2xSCCT + 1% RVC at room temperature, four times with 2xSCCT + 1% RVC at 65°C,
and twice more with 2xSCCT + 1% RVC at room temperature. Each wash was 5 minutes. Secondary
probe hybridization mix (2x DSM (dextran sulfate PVSA), 50% formamide, 1% SDS, 50 nM dye-conjugated
secondary probe, 1% RVC) was assembled and added dropwise onto a parafilm sheet sitting on top of a
damp paper towel in a petri dish. Coverslips were lifted and placed cell-side down onto the hybridization
mix drops and incubated for 1h at 37°C. Following secondary hybridization, the coverslips were transferred
back to 24-well plates and washed four times with 2xSCCT + 1% RVC at 37°C. The coverslips were finally
mounted cell side down on slides in DAPI-containing mounting medium. Images were captured on a Zeiss
LSM880 confocal microscope.
989
990 Primary probe used: Probes were designed to cover a ~750bp region (19 60mers) of the human *HOXA13*
991 intron. Secondary probes were dye conjugated. For RNA FISH foci intensity measurements, total RNA
992 pixel intensity was measured within each FISH foci boundary and normalized to the total RNA pixel intensity
993 of the corresponding nucleus.
994
Murine hematopoietic stem cell harvest and culture
Whole bone marrow was extracted from euthanized C57BL/6 mice using lower limb centrifugation and
subjected to RBC lysis. cKit+ cells were then isolated using the EasySep Mouse CD117 (cKIT) Positive
Selection Kit (StemCell). Cells were then stained with Zombie Aqua live/dead dye and cell surface markers
(hematopoietic lineage markers, Sca-1, cKit) prior to sorting out Lin⁻Sca-1⁺cKit⁺ cells on a BD FACSAria.
Lineage markers used are CD3, CD19, CD11b, B220, TER119, Gr-1, NK1.1, CD11c, TCRβ, and TCRγ/δ.
LSK cells were cultured for 48h prior to subsequent virus transduction. Cells are cultured in IMDM +
GlutaMAX supplemented with 10% FBS, 100 U/mL penicillin-streptomycin, 20 ng/mL mFlt3L, 20 ng/mL
mIL6, 100 ng/mL mSCF, 20 ng/mL mTPO, 1:1000 0.1M betamercaptoethanol, 1:100 200mM L-glutamine.
1004
1005 **Murine hematopoietic stem cell viral transduction**
1006 For LSK cell infection, the day prior to infection 420 uL of RetroNectin reagent (60 ug/mL) was added to
1007 untreated 24-well cell culture plates and incubated overnight at 4°C. The next day, RetroNectin was

removed and wells were blocked with PBS + 2% BSA for 30 minutes at room temperature. Wells were then washed once with Hank's balanced salt solution (HBSS). After removing the HBSS, desired volume of retrovirus was added to the wells. Plates were centrifuged at 4°C 3000 rpm for 1h. Next, retroviral supernatant was removed and cells were added to the virus-loaded wells with 10 ug/mL polybrene and spinfection was carried out at 37°C 1500rpm for 1.5h. Plates were then incubated overnight. Cell sorting was performed 48-72 hours post-infection.

Colony formation assay
6000 cells were resuspended in 600 uL of IMDM and added to 3mL of MethoCult methylcellulose media supplemented with 10 ng/mL mIL3, 10 ng/mL mIL6, and 20 ng/mL mSCF. 1 mL of the mixture was then added to each well of a 6-well plate (3 replicates total, 2000 cells per replicate). Plates were placed in the incubator and colonies were counted ~1-1.5 weeks after plating. Subsequent re-platings were carried out in the same way.

Bone marrow transplantation and engraftment measurements
LSK cells (CD45.2) were isolated and transduced with ENL transgenes as described above. Transduced LSK were mixed at a 1:3 ratio with RBC-lysed whole bone marrow from a CD45.1/2 mouse (100K LSK: 300K support cells). 400K cells resuspended in 100 uL PBS were injected retro-orbitally to each CD45.1 recipient mouse (B6.SJL-*Ptprc^a Pepc^b*/BoyJ, Jackson Labs #002014). The morning and afternoon prior to injection, CD45.1 recipient mice underwent 2 rounds of lethal irradiation (5.5 Gy per round, 11 Gy total).

Beginning 2 weeks after transplantation, engraftment was measured by peripheral blood staining. Peripheral blood (PB) was harvested by submandibular bleeding into EDTA coated tubes. ~15 uL PB was lysed in 1X RBC lysis buffer for 6 minutes on ice, quenched with FACS buffer (PBS + 2% FBS), and washed once in FACS buffer. Cells were stained with CD45.1 and CD45.2 antibodies (1:200) in FACS buffer for 30 minutes on ice protected from light. Cells were washed once more in FACS buffer and resuspended in FACS buffer + 1:2000 SYTOX blue dead cell stain. Samples were then run on an Attune NxT flow cytometer. Engraftment was measured approximately every 2 weeks for 1 year following transplantation.

Statistics and reproducibility
All experimental observations were confirmed by independent repeat experiments. Experimental data is presented as mean ± standard deviation unless stated otherwise. Statistical significance was calculated by two-tailed, unpaired or paired Student's t-test on two experimental conditions with $P < 0.05$ considered statistically significant unless stated other- wise. The number of replicates and the statistical test used were indicated in the corresponding figure legends. Statistics were performed using GraphPad Prism 10 and RStudio. No data were excluded from the analyses. No statistical methods were used to predetermine sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment.