Assessment of spring wheat responses to late season heat stress under Mediterranean conditions

Shaimaa Mahmoud Awad-Allah, Samer Mohamed Amer, Mahmoud Hassan Abdel-Moneim

1- Abstract

In Mediterranean climatic regions, wheat yield is significantly influenced by sowing date and cultivar selection, particularly where production could be limited by heat waves during critical growth stages. This investigation was undertaken to determine the effect of shifting sowing date on a cohort of six high yielding spring wheat cultivars (*Triticum aestivum* L.), by delaying sowing date 15 and 30 days in a two-year field trial, aiming to highlight agronomic traits contributing to sustained yield potential under heat stress induced by late sowing, with a focus on identifying tolerant cultivars.

Our findings showed that heat stress was detrimental when occurred during anthesis stage. Late sowing shifted the timing of heat stress, resulting in floral abortion and significantly reduced number of grains/spike. More importantly, some cultivars showed significant reduction following delaying sowing date, together with the strong correlation with yield especially at late sowing (r=0.74), imposing up to 42% yield penalty in the heat stressed late sowing conditions.

On the contrary, harvest index of sowing delayed by 30 days was significantly higher compared to both 15 days late sowing and the optimum sowing date, suggesting that decline in biological yield exceeded that of grain yield.

Upon shifting sowing date, the strength of traits association is altered, grain yield associations showed the strongest positive correlation in the 30 days late sowing with plant height (r=0.86), in the 15 days late sowing with number of total tillers/m^2 (r=0.63) and in optimum sowing date with biological yield (r= 0.62), suggesting that grain yield enhancement through selected traits with direct or indirect contributions should be evaluated independently across different growing conditions.

2- Introduction

Wheat (*Triticum aestivum* L.) is globally feeding 2.5 billion people in more than 85 countries and a source of at least 20% calories in the human diet in these countries [1], Moreover, it accounts for 40% of the protein and 33% of the calories in the Egyptian diet [2].

Wheat yield is a polygenic quantitative trait influenced by multiple components, e.g. number of spiked tillers, number of grains per spike, and grain weight. It is also highly sensitive to environmental conditions [3]. One critical agricultural practice for maximizing yield is identifying the optimal sowing date. Delayed sowing can negatively impact yield, its components, and overall wheat growth and development [4]. Specifically, it can disrupt key growth stages like heading time, grain filling rate and maturity[5], leading to reduced number of spikes per plant and number of grains per spike[6], ultimately causing yield penalty. However, the significance of these effects profoundly varies depending on the genotype [7].

Climate change poses significant threats to wheat production. It was reported that global temperatures are rising by 0.18°C per decade, with each degree of increase potentially reducing wheat production by 6% [8], with several empirical and simulated studies reporting 6.5%-27% yield reduction with above optimum temperature increase during various critical growth stages [9-11]. Therefore, a key focus for researchers is enhancing yield and stability while minimizing resource investment. Hütsch et al. [12] found that number of grains per spike was reduced under heat stress by 83% and 75% during grain filling and at maturity, respectively but decreased the individual grain weight only by 23% at maturity and that lead to a significant 36% reduction in harvest index.

Wheat is considered a heat sensitive crop [13] having optimal temperature of about 20°C for above ground growth and development [14]. Late sowing could put wheat under stress, specifically terminal heat stress which adds yield penalty because of reduction in grain yield key components *viz*., tillering potential, number of spikes/unit area, number of grains/spike and grain weight [12]. Heat stress during the terminal growth stage accelerates the assimilation of photosynthates due to early senescence, leading to smaller grains and reduced quality[15].

Number of grains per spike is established during and shortly after anthesis, mainly via the process of grain setting. A process affected by the maximum number of florets initiated at the double-ridge stage, which occurs simultaneously with the onset of tillering, and the proportion of these florets successfully developing into grains after pollination [16]. Consequently, this yield-determining factor is also partly shaped by events that take place during the vegetative growth phase. Heat stress can negatively impact the formation of double ridges on the shoot apex [12], impede fertilization due to pollen sterility [17] and cause the abortion of already fertilized grains because of inadequate assimilates supply.

The rate and duration of grain filling are the main drives of grain weight. Research investigating the effects of heat stress during this phase showed that while the grain filling rate increases, the duration is significantly shortened. Wheat plants exposed to heat stress reached maturity earlier than those grown under optimal conditions. However, the yield losses caused by the reduced grain filling period were not compensated for by the accelerated filling rate [18-20].

Thus, considering these constraints an experiment was conducted to investigate the effect of delayed sowing date and its possible consequent heat stress on wheat grain yield and yield components, using six spring wheat cultivars representing a range of attributes including yield potential and tolerance to biotic and abiotic stresses.

3- Materials and methods

3.1- Plant material

The used germplasm is a collection of six spring wheat cultivars, obtained from Ministry of Agriculture, Egypt, listed in Table 1.

Table 1. Particulars of Egyptian spring wheat cultivars

Cultivar	Attributes
Misr1	High yielding cultivar and resistant to rusts, especially stem rust UG99.
Giza171	Cultivated across Egypt, resistant to a wide range of rusts.
Gemeza11	Cultivated in the delta region, prone to late season shattering.
Giza168	Drought tolerant cultivar and resistant to yellow rust.
Sids12	Cultivated across Egypt, resistant to a wide range of rusts.
Shandwel	High tillering and high yielding cultivar.

3.2- Field trials

The genotypes panel was evaluated during 2018-2019 and 2019-2020 winter seasons at the experimental station of the Faculty of Agriculture, Alexandria University, Alexandria, Egypt (31º20' N, 30º E), where the soil is sandy loam (54% sand, 30% silt, and 16% clay), electrical conductivity of 1.30 dS m^{-1} and pH 8.05, average and maximum monthly temperature (°C) for the two seasons are illustrated in Figure 1.

Figure 1. Maximum and average temperature for the two growing seasons

A randomised complete block design was employed in a split plot, consisting of three blocks (replicates). The main plots were assigned for the three sowing dates, with 15 days interval (15th November, 30th November and 15th December, noting that optimum sowing date under such environment is mid-November, while the sub-plots were dedicated for the cultivars. Seeds were sown in 4 m^2 (2 x 2m) plots and seeding rate of 57 g/plot (143 kg/ha). Plots were maintained free of weeds and disease with the appropriate herbicides and fungicides and received standard irrigation and fertilization regime (Nitrogen fertilization as 476 kg/ha of urea 46.5% split over three doses, Phosphorus as 595 kg/ha of mono-superphosphate 15% added to the soil during seed bed preparation and Potassium as 119 kg/ha of potassium sulphate added 30 days after sowing.

3.3- Phenotyping

Preharvest traits; plant height (P.ht) was measured in cm using the rising plate [21] as the average of two measurements per plot at maturity, number of total tillers/m^2 (TT) measured as number of tillers in randomly picked 1m^2 within the plot and number of spiked tillers/m^2 (ST) measured as number of tillers bearing spikes in the same randomly picked 1m^2. Post harvest traits; spike length (SL), number of spikelets/spike (SS) and number of grains/spike (GS) measured as the average of ten spikes/plot randomly collected before harvesting, manually measuring the spike length and count number of spikelets/spike then threshed to estimate grain yield, biological yield (BY) by manually harvesting all above ground biomass/plot at maturity and grain yield (GY) quantified by mechanically threshing the harvested biological yield then weighing the grain yield, both estimated as kg/plot then transformed to ton/hectare, thousand grain weight in grams (TGW) measured as the average of two samples/plot collected from the harvested grain yield. Harvest index (HI) was estimated as grain yield/biological yield * 100.

3.4- Statistical analysis

R software R 3.3.4[22] was used to analyse experimental data. Means and standard errors were calculated using ANOVA to identify differences between treatments, R package 'Agricolae' [23] was used to perform split plot analysis for traits without homogeneity of error across years, 'lmerTest' R package[24] for combined analysis across years for traits with homogeneous error,' Corrplot' package[25] for correlation matrix and heat maps and R package 'ggplot2' [26] for graphical illustrations.

4- Results

The significance levels of measured traits responses under the effects of the two studied factors and their interactions are presented in Table 2 for the traits with heterogeneous

experimental error across years, where the main effect of sowing date, cultivars and their interaction varied among the three traits but showed the same pattern in TGW and GS while SS showed opposite pattern over the two years. TGW was not affected by sowing dates in both seasons but varied significantly among cultivars (Table 3), the significant interactions between sowing dates and cultivars (Table 4) shows some cultivars with increase in TGW upon delaying sowing, e.g. Giza 168 and Shandwel while the other cultivars showed varying degrees in TGW decrease with delayed sowing (Figure 2). Number of grains/spike was significantly affected by cultivars X sowing date interaction, in the first season D2:Giza168 was significantly superior with 58.67 grains/spike and D3:Gemeza11 was the lowest with 30.33 grains/spike, while in the second season D3:Giza168 had the highest grains/spike of 101.67 and D1:Sids12 had the lowest with 57.67 grains/spike (Figure 3).

In contrast to TGW, number of spikelets/spike in the first season significantly increased by delaying sowing date, as D2 and D3 yielded more SS than D1 by 2.3% and 10.6%, respectively, in the second season cultivars X sowing date interaction was significant, with delaying sowing Giza168 and Giza 171 SS increased, Misr1 and Shandwel SS decreased while Gemezaa11 and Sids12 SS peaked only in D2 (Figure 4).

Table 2. Levels of significance of thousand grain weight (TGW), number of spikelets/spike (SS) and number of grains/spike (GS) as affected by sowing dates and cultivars in 2018-2019 and 2019-2020 winter seasons.

Sources of variation	d.f.	TGW		SS		GS	
		2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020
Sowing Date (SD)	2	86.23ns	26.31ns	53.32*	129.43ns	931.70*	693.48*
Cultivar (C)	5	830.51***	481.26***	94.12***	233.35ns	1216.09***	2812.76***
SD X C	10	607.79**	351.50**	18.04ns	1127.71***	654.07***	1328.52*

ns: non-significant at $p \leq 0.05$, * significant at $p \leq 0.05$, ** significant at $p \leq 0.01$, *** significant at $p \leq 0.001$.

Table 3. Mean values for thousand grain weight (TGW), number of spikelets/spike (SS) and number of grains/spike (GS) as affected by sowing dates and cultivars in 2018-2019 and 2019-2020 winter seasons.

Trait	TGW		SS		GS	
	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020
Gemeza11	73.45 b	56.37 b	23.63 a	42.78 a	39.66 c	73.11 bcd
Giza168	68.36 d	52.47 d	23.65 a	40.41 ab	54.33 a	89.22 a

Giza171	68.74 cd	52.85 cd	22.04 b	37.01 b	47.55 b	69.44 cd
Misr1	78.54 a	60.25 a	22.60 ab	38.68 ab	42.55 c	78.89 b
Shandwel	67.00 d	51.45 d	23.46 a	41.90 a	49.66 b	75.44 bc
Sids12	72.55 bc	55.75 bc	19.91 c	42.31 a	47.66 b	67.11 d
LSD 0.05	3.98	3.03	1.08	4.65	3.62	7.58

D1	70.13	53.94	21.62 b	42.18	50.28 a	73.94 b
D2	73.15	55.01	22.11 b	40.97	49.39 a	80.50 a
D3	71.06	53.83	23.92 a	38.45	41.05 b	72.17 b
LSD 0.05	ns	ns	1.17	ns	6.33	4.46

144   Mean values followed by the same letter(s) are not significantly different.

145   **Table 4**. Mean values for thousand grain weight (TGW), number of spikelets/spike (SS) and

146 number of grains/spike (GS) as affected by sowing dates and cultivars in 2018-2019 and 2019-

147 2020 winter seasons.

	TGW		SS		GS	
	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020
D1:Gemeza11	71.07 cde	54.67 cdefg	21.63 gh	45.97 abc	44.00 fghij	72.67 cdef
D1:Giza168	58.93 f	45.33 h	23.53 bcdef	34.30 fg	54.33 abc	83.00 abc
D1:Giza171	71.07 cde	54.67 cdefg	20.80 hi	35.30 efg	50.33 cde	74.67 cde
D1:Misr1	80.60 a	62.00 a	22.07 efgh	44.87 abc	46.00 defghi	78bcde
D1:Shandwel	65.87 e	50.67 fg	22.73 cdefg	47.43 ab	57.67 ab	77.67 bcde
D1:Sids12	73.23 bcd	56.33 bcde	18.93 i	45.20 abc	49.33 cdefg	57.67 g
D2:Gemeza11	78.47 ab	59.00 abc	23.67 bcde	39.10 cdefg	44.67 efghij	77.33 bcde
D2:Giza168	70.93 cde	53.33 defg	22.33 defgh	40.63 bcdef	58.67 a	90.00 ab
D2:Giza171	68.77 cde	51.67 efg	22.73 cdefg	35.87 defg	43.67 ghij	68.33 efg
D2:Misr1	80.69 a	60.67 ab	21.67 fgh	36.63 defg	43.00 hij	82.33 abcd
D2:Shandwel	66.06 e	49.67 gh	23.13 cdefg	43.73 abcd	54.67 abc	82.00 abcd
D2:Sids12	74.04 abc	55.67 bcdef	19.13 i	49.53 a	51.67 bcd	83.00 abc
D3:Gemeza11	70.84 cde	53.67 defg	25.60 a	43.27 abcde	30.33 l	69.33 defg
D3:Giza168	75.24 abc	57.00 abcd	25.10 ab	46.30 abc	50.00 cdef	94.67 a
D3:Giza171	66.44 de	50.33 gh	22.60 defgh	39.87 bcdefg	48.67 cdefgh	65.33 efg
D3:Misr1	74.36 abc	56.33 bcde	24.07 abcd	34.53 fg	38.67 jk	76.33 cde
D3:Shandwel	69.08 cde	52.33 defg	24.53 abc	34.53 fg	36.67 k	66.67 efg
D3:Sids12	70.40 cde	53.33 defg	21.67 fgh	32.20 g	42.00 ijk	60.67 fg
L.S.D	6.89	5.24	1.87	8.05	6.26	13.13

148   Mean values followed by the same letter (s) are not significantly different.

Figure 2. Variations in thousand grain weight (TGW) as affected by the significant interaction between sowing dates and cultivars during (a) 2018-2019 and (b) 2019-2020 winter seasons.

Figure3. Variations in number of grains/spike (GS) as affected by the significant interaction between sowing dates and cultivars during (a) 2018-2019 and (b) 2019-2020 winter seasons.

Figure 4. Variations in number of spikelets/spike (SS) as affected by the significant interaction between sowing dates and cultivars during 2018-2019 and by significant main effect of sowing dates and cultivars during 2019-2020.

Table 5. Levels of significance of plant height (P.ht), biological yield (BY), grain yield (GY), harvest index (HI), number of total tillers/m^2 (TT), number of spiked tillers/m^2 (ST) and spike length (SL) as affected by year, sowing dates and cultivars in 2018-2019 and 2019-2020

Sources of variation	D.f	P.ht	BY	GY	HI	TT	ST	SL
year	1	1891.70*	9032.70**	131.71**	124.38*	52163ns	46927ns	99.57**
Sowing date (SD)	2	215.40ns	1039.70***	8.27**	166.35*	50398***	37041**	21.44*
Cultivar (C)	5	341.80**	85.70ns	9.68***	272.55***	77771***	72733***	41.91***
Year X SD	2	3373.60***	591.70***	12.35**	87.26ns	312ns	378ns	8.89ns
Year X C	5	421.00***	330.00***	13.55***	258.16***	6554*	7339*	24.00*
SD X C	10	98.10ns	760.60***	23.29***	531.77***	22153***	29960***	37.21*
Year X SD X C	10	212.80ns	396.10***	55.45***	697.35***	10998*	11810*	44.30**

ns: non-significant at p ≤ 0.05, * significant at p ≤ 0.05, ** significant at p ≤ 0.01, *** significant at p≤0.001.

Table 5 shows that year X cultivar effect was significant for all traits, the three way and SD X C were significant for all traits except plant height, while Year X SD was significant for plant height, biological and grain yield only. The main effect of cultivars was not significant for biological yield, while the main effect of sowing date was not significant for plant height and the year main effect was not significant for number of total tillers and spiked tillers/m^2.

Significance of plant height with Year X Cultivar interaction is mainly due to the stability of Sids12 height across the two seasons while all other cultivars hight was reduced significantly in the second season (Figure 5b). Sowing date effect varied significantly between the two years for D3 (decreased in the second year by 20.5%) while D1 and D2 effect was stable across years (Figure 5a).

The effect of sowing date X cultivar on biological yield varied significantly across years (Figure 7a), highlighting D2:Giza171 in the first year as the highest (42.11 t/ha) and D3:Giza168 in the second year as the lowest (5.30 t/ha), indicating a strong seasonal impact. Notably, all sowing date X cultivar combinations were significantly reduced for biological yield in the second season (except for D2:Shandwel) with various degrees ranging from 28.5% decrease in D1: Giza168 up to 85 % in D3: Giza168 (Table 7).

Table 6 shows significant 42% reduction in yield of D3 compared to D1 and D2 of the second season. The significant Year X SD X C interaction in grain yield showed the superiority of D2:Misr1in the first year (6.85 t/ha) (Table 7) and D3:Giza168 in the second year (same as BY) as the lowest (0.80 t/ha). In the second year, for each sowing date X cultivar combination grain yield was reduced by at least 40% and notably reduced by 83% in D3:Giza168, except for D2:Giza171 which didn't lose significant grain yield in the second year (Figure 7b). Table 7 illustrates that GY was reduced by delaying sowing date in most cultivars in each season, with few exceptions of increase in D2 or D3 compared to D1 for the same cultivars specially in 2018-2019 season, Moreover, Giza171 and Misr1 cultivars showed stability in the second season as yield reduction was not significant in D2 and D3 relative to D1.

A different pattern from biological and grain yield was observed in harvest index, although the Year X SD X C interaction was significant, Figure 7b shows that some sowing date X cultivar combinations significantly decreased, others increased, and some didn't change across years e.g. D3:Shandwel which had the highest harvest index across the two years (Table 7).

Figure 6a illustrates that most sowing date X cultivar combinations were significantly reduced for number of total tillers/m^2 in the second season, however D2: Sids12 and D3: Giza168 increased insignificantly as shown in Table 7, the significant decrease ranged from 9.8% in D3: Misr1 up to 28.8% in D2:Misr1, noting that the highest decrease for sowing date X cultivar combination between the two years was observed in D2 (20.6%-28.8%).

A similar pattern for the number of total tillers/m^2 was found in response of Number of spiked tillers/m^2 response to the three-way interaction (Figure 6b), with significant decrease in most sowing date X cultivar combinations in the second year ranging from 9.7% in D1:Giza171 up to 28.8% in D2:Misr1, with exceptional one significant increase in the second year for D2:Sids12 combination as 27.8% increase (Table 7).

The significant Year X SD X C interaction in terms of spike length showed D3: Giza168 in the second season with the highest value of 18.53 cm while the shortest spike length of 11.27 cm was found in D1:Sids12 in the first year (Table 7). For most sowing date X cultivar combinations, the spike length significantly increased in the second year from 1.67 cm in D2:Giza171 up to 4.36 cm in D1:Sids12. However, D3:Gemeza11 didn't follow the same trend and had a significantly decreased spike length in the second year (Figure 7).

Table 6. Mean values for plant height (P.ht), biological yield (BY), grain yield (GY), harvest index (HI), number of total tillers/m^2 (TT), number of spiked tillers/m^2 (ST) and spike length (SL)as affected by sowing dates and cultivars in 2018-2019 and 2019-2020.

Trait	P.ht		BY		GY		HI		TT		ST		SL	
	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020	2018-2019	2019-2020
Gemeza11	110.11 ab	101.23 cd	32.02 c	15.87 de	4.54 bc	1.92 f	14.24 def	12.61 ef	243.33 bc	192.22 gh	238.00 bc	183.11 g	15.10 abc	15.54ab
Giza168	109.06 ab	97.62 d	35.58 ab	15.25 e	5.06 a	1.88 f	14.30 def	13.87 def	217.11 def	170.33 hi	209.11 def	170.33 gh	14.25 cd	15.53 ab
Giza171	112.31 a	99.33 cd	37.73 a	14.91 e	4.37 c	2.28 ef	12.22 f	17.11 abc	199.55 fg	160.00 i	194.55 fg	156.66 h	12.75 ef	14.43 bcd
Misr1	110.84 ab	101.26 cd	35.91 ab	16.28 de	5.21 a	2.52 e	14.96 cde	16.21 bcd	254.55 b	197.55 fg	243.55 b	190.22 fg	12.27 f	15.23 abc
Shandwel	108.00 b	101.41 c	30.08 c	18.26 d	4.90 ab	3.27 d	17.68 ab	19.07 a	300.11 a	235.11 bcd	291.77 a	229.55 bcd	13.62 de	15.60 a
Sids12	101.40 c	100.64 cd	34.90 b	15.89 de	4.15 c	3.10 d	12.09 f	19.49 a	230.55 cde	213.77 ef	220.55 cde	207.11 ef	12.44 f	15.63 a
LSD 0.05	3.68		2.79		0.47		2.49		21.3		22.02		1.13	
D1	102.42 c	104.80 bc	35.99 a	20.00 c	4.89 a	2.91 b	14.18 b	14.87 b	250.39 a	204.00 b	246.56 a	199.00 b	12.53 d	15.23 a
D2	108.28 b	104.57 bc	33.83 ab	19.76 c	4.43 a	2.91 b	13.83 b	14.89 b	209.22 b	169.33 c	205.22 b	166.67 c	13.39 cd	15.13 ab
D3	115.17 a	91.38 d	33.30 b	8.48 d	4.79 a	1.67 c	14.73 b	19.41 a	263.00 a	219.33 b	247.00 a	209.33 b	14.30 bc	15.62 a
LSD 0.05	4.3		2.6		0.47		2.81		22.73		11.82		0.92	

Mean values followed by the same letter(s) are not significantly different.

Figure 5. Variations in plant height (P.ht) as affected by the significant effect of sowing dates (a) and interaction between year and cultivars during 2018-2019 and 2019-2020 seasons.

Figure 6. Variations in number total tillers/m² (TT) (a) and spiked tillers/m² (ST) (b) as affected by the significant interaction between sowing dates and cultivars during 2018-2019 and 2019-2020

Figure 7. Variations in biological yield (BY) (a), grain yield (GY) (b) harvest index (HI) (c) and (d) spike length (SL) as affected by the significant interaction between sowing dates and cultivars during 2018-2019 and 2019-2020.

Correlations (Pearson's Correlation Coefficient - PCC) between traits under different sowing dates (Figure 8) were calculated to assess the association among studied traits and the environmental dependences of these associations. Number of total tillers/m^2 and number of spiked tillers/m^2 had the highest correlation (r=0.98) over the three sowing dates.

TGW showed negative association with spike length, number of spikelets/spike and number of grains/spike with various degrees across sowing dates, its strongest (r= -0.8) with number of spikelets/spike and number of grains/spike at D2. On the other hand, number of spikelets /spike and number of grains/spike were both highly correlated and this association peaked (r=0.85) at D2. Figure 8 illustrates the positive correlation between spike length and SS and GS varied among sowing dates, with its highest at D1 (r=0.89 and r= 0.57, respectively). The sowing date effect on association between GS and SS is evidently shown in figure 8, with its lowest association in D1 (r=0.69) and its highest in D2 (r=0.85).

Grain yield associations varied with traits and within sowing dates, noting the strongest positive correlation in D3 with plant height (r=0.86) which fell in D2 (r=0.47) and vanished in D1, while GY positive correlation with ST varied slightly among sowing dates (r=0.38 to r=0.6). Correlations with post-harvest traits showed TGW to have positive

association with GY that increased with delaying sowing date, while SS and GS negatively
correlated with GY across the three sowing dates (Figure 8).

Harvest index had no significant correlation with GY in D3 but showed significant
positive association with GY (r= 0.43) and (r=0.67) in D1 and D2, respectively and showed
no significant associations with post-harvest traits across the sowing dates except with TGW
in D3 (r=-0.46).

Figure 8. Pearson's correlation coefficient for the measured traits. (a) D1, (b) D2 and (c) D3.

Table 7. Mean values for plant height (Pht), biological yield (BY), grain yield (GY), harvest index (HI), number of total tillers/m² (TT), number of spiked tillers/m² (ST) and spike length (SL) as affected by interaction between year, sowing dates and cultivars in 2018-2019 and 2019-2020 seasons.

Treatment	BY 2018-2019	BY 2019-2020	GY 2018-2019	GY 2019-2020	HI 2018-2019	HI 2019-2020	TT 2018-2019	TT 2019-2020	ST 2018-2019	ST 2019-2020	SL 2018-2019	SL 2019-2020
D1: Gemeza11	30.42 fgh	19.24 lmn	5.47 bcde	2.52 klm	18.18 cde	13.05 fghij	255.00 cde	216.00 fghij	252.00 bc	199.33 hijk	12.87 ijklm	16.43 bc
D1: Giza168	35.92 bcde	25.70 hij	5.58 bcd	2.61 jklm	15.75 defgh	10.60 ijkl	209.33 hijkl	176.67 jklm	209.33 fghij	176.67 jklm	14.20 defghi	14.20 defghi
D1: Giza171	39.33 abc	14.39 op	4.32 fgh	2.41 klm	10.99 ijkl	16.89 def	184.33 jklmno	164.67 mnop	177.33 jklm	160.00 lm	13.50 fghijkl	13.07 hijklm
D1: Misr1	36.92 bcd	20.48 klm	4.93 def	2.61 sjklm	14.42 efghi	13.11 fghij	294.33 ab	212.00 ghijk	289.33 ab	208.67 fghi	12.20 klm	15.77 bcd
D1: Shandwel	40.33 ab	24.79 jik	3.89 ghi	5.15 cde	9.65 jkl	21.21 abc	327.00 a	248.00 cdefg	326.00 a	247.33 cde	13.60 efghijkl	16.30 bc
D1: Sids12	33.00 def	15.43 nop	5.14 cde	2.17 klm	16.09 defgh	14.41 efghi	232.33 cdefgh	206.67 hijkl	225.33 cdefgh	202.00 ghij	11.27 m	15.63 bcd
D2: Gemeza11	33.83 def	21.24 jklm	4.74 ef	2.28 klm	13.79 fghij	10.87 ijkl	215.00 ghij	160.00 nop	212.00 efghi	158.00 lm	15.27 bcdef	15.20 cdefg
D2: Giza168	35.42 cde	14.76 nop	4.67 efgh	2.25 klm	13.19 defgh	15.67 defgh	212.33 ghijk	164.00 nop	212.33 defghi	164.00 klm	14.27 defghi	13.87 ghijk
D2: Giza171	42.11 a	21.95 jkl	2.84 jk	2.52 klm	6.79 l	11.58 hijk	193.00 jklmn	140.00 p	193.00 hijkl	140.00 m	13.40 fghijkl	15.07 cdefghijk
D2: Misr1	34.83 cdef	16.80 mno	6.85 a	2.73 jkl	19.63 abcd	16.34 defg	223.00 defghi	158.67 nop	220.00 cdefgh	156.67 lm	12.20 klm	14.733 cdefghi
D2: Shandwel	22.21 jkl	20.83 klm	4.93 def	2.55 klm	22.08 abc	12.30 ghij	258.67 bcd	205.33 hijkl	253.67 bc	202.00 ghij	12.40 jklm	15.53 bcde
D2: Sids12	34.58 cdef	22.98 jkl	2.61 jklm	5.18 bcde	7.56 kl	22.64 ab	153.33 op	188.00 jklmno	140.33 m	179.33 ijkl	12.80 ijklm	16.40 bc
D3: Gemeza11	31.83 efg	7.15 qr	3.41 ij	0.99 n	10.77 jkl	13.94 efghij	260.00 bc	200.67 hijklm	250.00 cd	192.00 hijkl	17.17 ab	15.00 cdefgh
D3: Giza168	35.42 cde	5.30 r	4.93 def	0.80 n	13.98 efghi	15.36 defgh	229.67 cdefgh	252.50 cdef	205.67 fghij	158.50 lm	15.00 cdefgh	18.53 a
D3: Giza171	31.75 efg	8.45 qr	5.97 b	1.90 m	18.87 bcd	22.87 ab	221.33 efghi	175.33 lmnop	213.33 defghi	170.00 jklm	13.07 hijklm	15.17 cdefg
D3: Misr1	35.98 bcde	11.59 pq	3.87 hi	2.24 klm	10.84 ijkl	19.21 bcd	246.33 cdefg	222.00 efghi	221.33 cdefgh	205.33 fghij	12.43 jklm	15.20 cdefg
D3: Shandwel	27.72 ghi	9.18 qr	5.90 bc	2.13 klm	21.32 abc	23.69 a	314.67 a	252.00 cdef	295.67 a	239.33 cdefg	14.87 cdefgh	14.97 cdefgh
D3: Sids12	37.13 bcd	9.28 qr	4.700 efg	1.95 lm	12.65 fghij	21.44 abc	306.00 a	246.67 cdefg	296.00 a	240.00 cdef	13.27 ghijkl	14.87 cdefgh
LSD$_{0.05}$	4.83		0.82		3.05		36.66		37.90		1.97	

Mean values followed by the same letter(s) are not significantly different.

5- Discussion

In a Mediterranean type environment the correct choice of sowing date and cultivar are critical determinants of yield, sowing date usually occurs within a sowing window starting with the first significant rainfall in autumn and closing when a sowing date is too late to achieve a reasonable yield [27].

Although sowing date didn't affect GY in the first season, we found a significant decline of 42% in the late sowing date D3 in the second season which coincided with above average temperature during late February and March (Figure 1), as early sowing dates increase the interception of solar radiation by the crop, allowing it to accumulate more dry matter and avoid terminal drought/heat at the end of the growing season or shorter harvest window in late season sowing [27, 28].

Number of spikelets/spike in D3 increased significantly in the first season, but didn't change significantly in the second year which can be explained by changes in spike length rather than effect of stress on fertility, as SL showed strong association with SS (Figure 8) and sowing date had the same effect on SL as SS (Tables 3 and 6). Conversely, number of grains/spike decreased significantly in D3, probably due to terminal heat stress during anthesis (Figure 1), which usually takes place at February/early March [29] suggesting that GY decrease in D3 is attributed to the poor seed set and decrease number of spikes/m^2 [12, 28, 30]. GS was reduced in D3 in both seasons by an average of 8.3 grains/spike, possibly due to incomplete fertilization, as heat elevation at anthesis adversely impacts pollens fertility [12, 17]. The number of grains is largely determined by the number of fertile florets, which in turn depends on the dynamics of floret generation and degeneration [31]. Floret degeneration or mortality typically occurs when the stem and spike are growing at their fastest rates [32]. It is believed that the variation in number of grains per spike is linked to changes in the availability of assimilates for spike growth during the period of rapid stem elongation, when floret abortion could occur due to adverse conditions [33]. With limited availability of assimilates, many floret primordia may degenerate or die, leaving less florets reaching the fertile stage at anthesis [32].

TGW didn't change following sowing date alteration in both years despite the decrease in grains number/spike in the second season which in normal conditions would have shown the trade-off between TGW and GS [34], possibly due to limited source capacity of biomass [35] which appeared as a positively strong correlation (r=0.89) between BY and TGW in Figure 8c.

Biological yield didn't change in the first season but showed significant 57% diminish in D3 of the second season, matching the pattern of grain yield, these results align with Aslani et al. [6] findings who reported negative impacts of delayed sowing on biomass of wheat. Under stress conditions, the plants adjust to the unfavourable conditions by limiting biomass growth, possibly facilitating increased transpiration while maintaining stable water consumption and water use efficiency [12].

Harvest index was stable in the first season across sowing dates following both BY and GY; Nevertheless, it showed an opposite pattern in the second season as HI in D3 was significantly higher than that of D1 and D2 contradicting several reports which found HI to follow BY and GY decline [11, 12], probably as decline in BY was greater than that of GY.

Plant height showed opposite performance across years, in D3 of the first year it was significantly higher than D1 and D2, an unusual finding reported in one of a two-years trial by Elbashier et al. [36]. In a different study, Sareen et al. [37] found no significant reduction under late sowing/heat stress, conversely, it was significantly lower in D3 of the second year as was most of the measured traits due to heat stress in February/March, agreeing with others who reported shorter plants under late sowing [38, 39].

TT and ST at D3 were equal to that of D1(optimal sowing date) in both years, probably because heat stress induced by late sowing in this experiment didn't coincide with tillering or booting , while others found it might even increase under late sowing [40, 41]

SL in D3 of the first year slightly surpassed D1, however, in the second change it showed stable performance under the three sowing dates, minimal or no impact of sowing date on spike length [40], on the contrary, SL get significantly reduced if the late sowing led to terminal heat stress [42, 43].

As the sowing date changed, the strength of traits association is altered, indicating that grain yield enhancement through prioritize traits with direct or indirect contributions evaluated independently across different growing conditions to account for environmental interactions [28]. This alteration was observed in several cases e.g. SL and GS association which shifted by delaying sowing as r= 0.57, 0.47 and 0.42 in D1, D2 and D3, respectively. Moreover, grain yield associations varied profoundly among sowing dates, noting the strongest positive correlation in D3 with plant height (r=0.86), (r=0.47) in D2 and vanished in D1(r= -0.05) (Figure 8).

The TGW negative association with spike length, number of spikelets/spike and number of grains/spike across sowing dates, this association aligns with the well-established trade-off between TGW, SS and GS [34, 44].

The genotypes showed wide range of responses both within growing seasons and across sowing dates, with significant interactions observed in most of the studied traits. Nevertheless, 'Giza171' and 'Misr1' demonstrated yield stability in the second year, aligning with their stability in number of grains/spike, highlighting the significance of such trait in sustaining yield potential under heat stress during anthesis, a result reported in investigations on other wheat germplasms [45, 46].

6- Conclusion

Heat stress induced by late sowing could significantly impair wheat grain yield, however, the yield components driving the yield penalty significantly vary depending on (1) differential genotypic responses and (2) sowing date-dependent heat stress across growth stages, fluctuating annually. Our results under mediterranean conditions illustrated that late sowing could impose heat stress during anthesis, resulting in significant floret abortion and consequent reduction in grain number/spike and ultimate grain yield, a consequence that could be alleviated by growing the cultivars with ability of sustaining yield potential under such stress.

7- Declarations:

7.1- Ethics approval and consent to participate:

Not applicable.

7.2- Consent for publication:

Not applicable.

7.3- Availability of data and materials:

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

7.4- Competing interests:

The authors declare that they have no competing interests.

7.5- Funding: No funding.

7.6- Authors' contributions:

The three authors contributed equally to designing, data collection of the experiment, data analysis and writing the manuscript.

8- References

1. Yashavanthakumar, K.J., et al., *Impact of heat and drought stress on phenological development and yield in bread wheat*. Plant Physiology Reports, 2021. **26**(2): p. 357-367.

2. Abdalla, A., M. Becker, and T. Stellmacher *The Contribution of Agronomic Management to Sustainably Intensify Egypt's Wheat Production*. Agriculture, 2023. **13**, DOI: 10.3390/agriculture13050978.

3. Pang, Y., et al., *High-Resolution Genome-wide Association Study Identifies Genomic Regions and Candidate Genes for Important Agronomic Traits in Wheat*. 2020(1752-9867 (Electronic)).

4. Andarzian, B., et al., *Determining optimum sowing date of wheat using CSM-CERES-Wheat model*. Journal of the Saudi Society of Agricultural Sciences, 2015. **14**(2): p. 189-199.

5. Yasmeen, A., et al., *PERFORMANCE OF LATE SOWN WHEAT IN RESPONSE TO FOLIAR APPLICATION OF Moringa oleifera Lam. LEAF EXTRACT*. Chilean journal of agricultural research, 2012. **72**: p. 92-97.

6. Aslani, F. and M.R. Mehrvar, *Responses of Wheat Genotypes as Affected by Different Sowing Dates*. Asian Journal of Agricultural Sciences, 2012. **4**: p. 72-74.

7. Ahamed, K.U., K. Nahar, and M. Fujita, *Sowing date mediated heat stress affects the leaf growth and dry matter partitioning in some spring wheat (Triticum aestivum L.) Cultivars*. IIOAB Journal, 2010. **1**: p. 8-16.

8. Wang, X., et al., *Genome-wide association study identifies QTL for thousand grain weight in winter wheat under normal- and late-sown stressed environments*. Theoretical and Applied Genetics, 2021. **134**(1): p. 143-157.

9. Ullah, A., et al., *Heat stress effects on the reproductive physiology and yield of wheat*. Journal of Agronomy and Crop Science, 2022. **208**(1): p. 1-17.

10. Barkley, A., et al., *The impact of climate, disease, and wheat breeding on wheat variety yields in Kansas, 1985–2011*. Bulletin, 2013. **665**: p. 1-31.

11. Bergkamp, B., et al., *Prominent winter wheat varieties response to post-flowering heat stress under controlled chambers and field based heat tents*. Field Crops Research, 2018. **222**: p. 143-152.

12. Hütsch, B.W., D. Jahn, and S. Schubert, *Grain yield of wheat (Triticum aestivum L.) under long-term heat stress is sink-limited with stronger inhibition of kernel setting than grain filling*. Journal of Agronomy and Crop Science, 2019. **205**(1): p. 22-32.

13. Wang, X., et al., *Physiological and proteome studies of responses to heat stress during grain filling in contrasting wheat cultivars*. Plant Science, 2015. **230**: p. 33-50.

14. Porter, J.R. and M. Gawith, *Temperatures and the growth and development of wheat: a review*. European Journal of Agronomy, 1999. **10**(1): p. 23-36.

15. Porter, J.R., *Rising temperatures are likely to reduce crop yields*. Nature, 2005. **436**(7048): p. 174-174.

16. Kirby, E.J.M., *Ear development in spring wheat*. The Journal of Agricultural Science, 1974. **82**(3): p. 437-447.

17. Prasad, P.V.V. and M. Djanaguiraman, *Response of floret fertility and individual grain weight of wheat to high temperature stress: sensitive stages and thresholds for temperature and duration*. 2014(1445-4416 (Electronic)).

18. Dias, A.S. and F.C. Lidon, *Evaluation of Grain Filling Rate and Duration in Bread and Durum Wheat, under Heat Stress after Anthesis*. Journal of Agronomy and Crop Science, 2009. **195**(2): p. 137-147.

19. Wardlaw, I.F., I. Sofield, and P. Cartwright, *Factors Limiting the Rate of Dry Matter Accumulation in the Grain of Wheat Grown at High Temperature*. Australian Journal of Plant Physiology, 1980. **7**: p. 387-400.

20. Wardlaw, I.F. and L. Moncur, *The Response of Wheat to High Temperature Following Anthesis. I. The Rate and Duration of Kernel Filling*. Australian Journal of Plant Physiology, 1995. **22**: p. 391-397.

471   21.   Sharrow, S.H., *A Simple Disc Meter for Measurement of Pasture Height and Forage Bulk.*
472   Journal of Range Management, 1984. **37**(1): p. 94-95.
473   22.   R development core team, R., *R development core team.* RA Lang Environ Stat Comput,
474   2013. **55**: p. 275-286.
475   23.   Mendiburu, F. and R. Simon, *agricolae.* 2009.
476   24.   Kuznetsova, A., P.B. Brockhoff, and R.H.B. Christensen, *lmerTest Package: Tests in*
477   *Linear Mixed Effects Models.* Journal of Statistical Software, 2017. **82**(13): p. 1 - 26.
478   25.   Simko, T.W.a.V. *R package 'corrplot': Visualization of a Correlation Matrix.* 2021; Available
479   from: https://github.com/taiyun/corrplot.
480   26.   Ginestet, C., *ggplot2: Elegant Graphics for Data Analysis.* Journal of the Royal Statistical
481   Society Series A: Statistics in Society, 2011. **174**(1): p. 245-246.
482   27.   Bassu, S., et al., *Optimising sowing date of durum wheat in a variable Mediterranean*
483   *environment.* Field Crops Research, 2009. **111**(1): p. 109-118.
484   28.   Ul-Allah, S., et al., *Phenotypic characterization of wheat germplasm for heritability and*
485   *dissection of association among post anthesis traits under variable sowing dates.*
486   Journal of King Saud University - Science, 2023. **35**(3): p. 102578.
487   29.   Wajid, A., et al. *Crop Models: Important Tools in Decision Support System to Manage*
488   *Wheat Production under Vulnerable Environments.* Agriculture, 2021. **11**,   DOI:
489   10.3390/agriculture11111166.
490   30.   Roy, D., et al., *Delayed sowing and its ramifications: biophysical, yield and quality*
491   *analysis of wheat cultivars in the northwest Indo-Gangetic plains.* Journal of the Science
492   of Food and Agriculture, 2024. **104**(11): p. 6831-6843.
493   31.   Prieto, P., et al., *Dynamics of floret initiation/death determining spike fertility in wheat as*
494   *affected by Ppd genes under field conditions.* Journal of Experimental Botany, 2018.
495   **69**(1460-2431 (Electronic)): p. 2633–2645.
496   32.   Youssefian, S., E.J.M. Kirby, and M.D. Gale, *Pleiotropic effects of the GA-insensitive Rht*
497   *dwarfing genes in wheat. 2. Effects on leaf, stem, ear and floret growth.* Field Crops
498   Research, 1992. **28**(3): p. 191-210.
499   33.   Reynolds, M., et al., *Achieving yield gains in wheat.* Plant, Cell and Environment, 2012.
500   **35,**(1365-3040 (Electronic)): p. 1799–1823.
501   34.   Griffiths, S., et al., *Genetic dissection of grain size and grain number trade-offs in*
502   *CIMMYT wheat germplasm.* Plose One, 2015. **16**(1932-6203 (Electronic)): p. 18.
503   35.   Zhang, B., et al., *Estimation of grain filling rate and thousand-grain weight of winter*
504   *wheat (Triticum aestivum L.) using UAV-based multispectral images.* European Journal
505   of Agronomy, 2024. **159**: p. 127258.
506   36.   Elbashier, E., Idris, S. , Tahir, I. , Elhanafi, S. , Saad, A. , Mustafa, H. , Idris, A. and
507   Tadesse, W., *Genome Wide Association Study of Yield and Yield-Related Traits in Elite*
508   *Spring Bread Wheat Genotypes Grown under High Temperature Environment in Sudan.*
509   American Journal of Plant Sciences, 2023. **14**: p. 18.
510   37.   Sareen, S., et al. *Resilience to Terminal Drought, Heat, and Their Combination Stress in*
511   *Wheat Genotypes.* Agronomy, 2023. **13**, DOI: 10.3390/agronomy13030891.
512   38.   Shaalan, A., M.A. Attia, and M.A. Hassaan, *Response of Some Wheat Cultivars to*
513   *Sowing Dates and Biofertilizers under North West Coast of Egypt.* Egyptian Journal of
514   Agronomy, 2019. **41**(3): p. 313-324.
515   39.   Basheir, S.M., et al. *Identification of Wheat Germplasm Resistance to Late Sowing.*
516   Agronomy, 2023. **13**, DOI: 10.3390/agronomy13041010.
517   40.   Kumar, S., et al., *Capturing agro-morphological variability for tolerance to terminal heat*
518   *and combined heat–drought stress in landraces and elite cultivar collection of wheat.*
519   Frontiers in Plant Science, 2023. **Volume 14 - 2023**.
520   41.   Rehman, H.U., et al. *Evaluation of Physiological and Morphological Traits for Improving*
521   *Spring Wheat Adaptation to Terminal Heat Stress.* Plants, 2021. **10**,   DOI:
522   10.3390/plants10030455.
523   42.   Gupta, V., et al., *AMMI and GGE biplot analysis of yield under terminal heat tolerance in*
524   *wheat.* (1573-4978 (Electronic)).

43.   Devi, K., et al., *Identification of Wheat Genotypes Resilient to Terminal Heat Stress Using*
GGE Biplot Analysis. Journal of Soil Science and Plant Nutrition, 2022. **22**(3): p. 3386-
3398.
44.   Vicentin, L., J. Canales, and D.F. Calderini, *The trade-off between grain weight and grain*
number in wheat is explained by the overlapping of the key phases determining these
major yield components. Frontiers in Plant Science, 2024. **Volume 15 - 2024**.
45.   Kim, J., R. Savin, and G.A. Slafer, *Quantifying pre- and post-anthesis heat waves on grain*
number and grain weight of contrasting wheat cultivars. Field Crops Research, 2024.
307: p. 109264.
46.   Qaseem, M.F., R. Qureshi, and H. Shaheen, *Effects of Pre-Anthesis Drought, Heat and*
Their Combination on the Growth, Yield and Physiology of diverse Wheat (Triticum
aestivum L.) Genotypes Varying in Sensitivity to Heat and drought stress. Scientific
Reports, 2019. **9**(1): p. 6955.