Title

Role of aridity in shaping adaptive genomic divergence and population
connectivity in a Southern African rodent

Abstract

Elucidating the drivers of evolution in dry environments is central to understanding how
organisms respond to climate change. While research on the genomics of adaptation is
growing, aridity-driven intraspecific divergence remains poorly quantified. Here, we
address this gap by using genomic data from 230 individuals of the arid-adapted four-
striped mouse *Rhabdomys bechuanae*, sampled across an aridity gradient in southern
Africa, a region facing increasing aridification. Combining these data with palaeoclimatic
reconstructions and present-day aridity indices, we investigate, from a spatio-temporal
perspective, how intraspecific genetic variation relates to aridity. Inference of past
effective population size revealed a sharp decline in the late Pleistocene, coinciding with
regional aridification and potentially reflecting changes in connectivity during dry
periods. Current population structure followed a pattern of isolation by distance and
mirrored the aridity gradient. Genotype-Environment Association analyses identified
SNPs and genes significantly associated with aridity and genetically differentiated
among populations, with functions related to water and energy conservation - as
expected under arid conditions - as well as neurotransmission. These findings highlight
the underappreciated role for neurological processes in coping with water and resource
scarcity. More broadly, our integrative genomics approach suggests that aridity shapes
population connectivity and adaptation, with implications for climate resilience.

Keywords

Introduction

Climate change has increasingly been associated with local extinctions [1], shifts in phenology [2] and alteration of species' geographic distributions [3]. These responses tend to be particularly pronounced in harsh environments, such as arid regions, where species already face several ecological and physiological constraints [4]. With projections indicating a rise in the intensity and frequency of drought events in the coming decades [5,6], it is essential to better understand how species respond to both temporal and spatial variation in aridity. By combining newly generated genomic data with available palaeoclimatic reconstructions and contemporary measures of aridity, this study aims to shed light on the role of aridification on the demographic history, spatial population structure and adaptive responses of a non-model rodent species. These insights are key for predicting resilience of natural populations under future climate scenarios and for guiding conservation strategies [7].

At the individual level, the primary challenges of life in arid environments include maintaining body temperature and conserving water [8]. These challenges are especially evident for species that rely on evaporative water loss mechanisms, such as sweating, to dissipate heat, particularly in mammals [9]. The physiological demands imposed by such environmental conditions, particularly when experienced over multiple generations, often exceed the limits of phenotypic plasticity in non-desert-adapted species. Consequently, it is assumed that strong selection has acted on complex physiological traits related to energy metabolism, water retention and thermoregulation in species inhabiting arid environments [8-10].

Genomic studies have identified recurrent genetic patterns among mammals living in deserts [10]. Evidence of adaptation to arid environments has included elevated allele frequency differentiation [11,12], reduced genetic diversity [11,13-15], increased non-synonymous-to-synonymous substitution rate ratios [16] and divergent gene expression patterns [17] in genomic regions involved in key functions for desert adaptation. These genomic regions often relate to key physiological traits relevant to the survival of both mammalian specialist and non-specialist species in deserts, including DNA repair, protein synthesis and degradation [15,16], thyroid hormone levels [11], water retention [18], lipid [16] and carbohydrate metabolism [11, 13, 19-21]. However, most of these studies have relied on qualitative comparisons between desert and non-desert species or populations, and relatively few have explicitly explored quantitative variation along continuous environmental gradients. Several other studies on the genomics of local adaptation have demonstrated the power of Genotype-Environment Association (GEA) analyses to identify polymorphisms associated with continuous environmental variation [22-24]. In this context, Rocha et al. [10] highlighted the use of the Aridity Index (AI), known to be a good predictor of certain aridity-adapted phenotypes, as a powerful, composite and quantitative environmental variable to inform genomic studies of adaptation to arid conditions.

In this study, we applied a GEA approach to investigate the genomic signatures of adaptation to arid conditions in *Rhabdomys bechuanae*, a four-striped mouse species

from Southern Africa. Its range overlaps with hot arid and hyper-arid summer rainfall regions of Southern Africa [25]. Divergence from its closest relatives within the *Rhabdomys* species complex occurred approximately 3.1-4.3 million years ago [26], and this diversification has been associated with environmental niche differentiation [27]. It likely reflects a long-term trend towards increasing aridity and seasonality in Southern Africa since the late Miocene, coupled with a shift from closed subtropical woodland to sparse, shrubby vegetation [28]. Currently, *R. bechuanae* thrives predominantly in sparsely vegetated areas and nests in bushes. Phenotypic traits suggest adaptation to aridity; these traits include morphological (longer tail length related to potential differences in locomotion and/or thermoregulation) and behavioural divergence (selection of different habitats) from more mesic congeners [23,29], as well as enhanced osmoregulatory ability, as suggested by differences in blood metabolite concentrations between populations of *R. bechuanae* and of its sister species *R. dilectus*, occurring syntopically in semi-arid environments [30]. Although *R. bechuanae* is considered one of the most arid-adapted species of its genus, it still occupies a wide environmental gradient, providing an ideal system to investigate whether the aridity gradient imposes divergent selection pressures on key physiological traits. We therefore hypothesised that this aridity gradient would impose divergent selective pressures on key physiological traits related to adaptation to aridity, leading to detectable intraspecific population differentiation and genetic variation at loci associated with aridity-related functions.

To investigate genomic signatures of evolution under arid conditions in *R. bechuanae*, we generated a genome-wide SNP dataset using restriction-site associated DNA sequencing (RAD-seq) [31]. Our overarching aim was to evaluate how the temporal and spatial variation in aridity has shaped the population history and structure of *R. bechuanae*, by integrating inferences of historical changes in effective population size with GEA analyses.

Using an extensive sample of individuals from multiple localities spanning the species' aridity gradient, our first objective was to reconstruct past changes in effective population size in relation to palaeoclimatically inferred shifts in aridity. We expected that historical episodes of increased aridity would be associated with population declines, either due to demographic bottlenecks [32,33], or via reductions in genetic diversity caused by habitat fragmentation and decreased population connectivity. Conversely, wetter periods were expected to correspond with population expansions or increased gene flow.

In addition to temporal variation, spatial variation in aridity may also influence current population structure and genetic diversity [12,34,35]. Therefore, our second objective was to characterise the current population structure within *R. bechuanae* and assess its relationship with spatial variation in aridity across the species' range. We predicted that population structure would correlate with the degree of aridity, reflecting its impact on gene flow and differentiation.

Our third objective was to assess the genomic basis of adaptation to aridity in *R. bechuanae* using a GEA approach with the Aridity Index as a quantitative predictor. This

approach enabled us to identify loci associated with spatial variation in arid conditions across the species' range and to identify potential biological functions that may underpin survival in arid environments. We hypothesised that site frequency variation would concentrate in genomic regions linked to key functions for survival in deserts such as DNA repair, protein modification, osmoregulation, and carbohydrate/fat metabolism [10,16].

Material & Methods

Samples, aridity index and RAD-seq data production

Individual samples of four-striped mice of the genus *Rhabdomys* were previously collected in their natural environment between 2007 and 2019 by us or collaborators [25]. Population-level samples from *R. bechuanae* were collected specifically for the present study in 2018 and 2019 (details for trapping procedures in [25]). Based on data obtained for *R. pumilio* indicating minimal relatedness at this distance [36], we aimed to avoid related individuals as much as possible, by selecting mice trapped at least 100 m apart, except for breeding pairs (a male and a female).

To study fine-scale intraspecific structure, as well as perform GEA and historical effective population size inferences in *R. bechuanae*, we selected a large number of individuals from this species, sampled in 11 different localities at least 20 km distant from each other and along the aridity gradient (*N*=230 individual samples in total, 19.82 +/- 4.72 individuals per locality, **Table S1**). We estimated the aridity level of each *R.bechuanae* sampling locality by computing the Aridity Index (AI) at each location ([38], see **Supplementary Material** for details). Species identification for all sampled individuals in the *Rhabdomys* species complex composed of cryptic species had previously been obtained via COI sequencing or RFLP genotyping ([25] and procedure therein; **Table S1**). DNA extractions for RAD-sequencing were performed from tail tissues preserved in 98% ethanol, using a 96-Well Plate Animal Genomic DNA Miniprep Kit (BioBasic Inc.). Since samples from other *Rhabdomys* species and subspecies had also been collected for a separate study, we proceeded with RAD-seq library preparation and sequencing for *R. bechuanae* samples analysed in this study concurrently with the other *Rhabdomys* samples (*N* = 411 in total). Library preparation followed the protocol of Etter et al. [39], with modifications as described in [40] (See **Supplementary Material** for details). Sequencing was performed using a 2x150 bp paired-end Illumina NovaSeq 6000 protocol at MGX-Montpellier GenomiX Facility (Montpellier, France).

RAD-seq raw data processing and filtering

Following standard Illumina processing and quality filtering, duplicate reads resulting from PCR amplification were discarded using the program *clone_filter* implemented in the *Stacks* v. 2.66 software [41]. The resulting paired-end reads were demultiplexed and quality filtered using the *process_radtags* pipeline, also implemented within *Stacks*. Sequence reads were aligned to the reference genome using *BOWTIE* (v. 2.4.5) [42]. The reference genome used was the one of the closely related species *R. pumilio* (2.3 Gb in length; GCA_030674055.1) [43]. Before mapping, putative repeated elements were hard-masked (i.e., repeated sequences were replaced by stretch of "N") from the reference genome using the mouse repeated element Repbase Update (v. 27.06) database using *Repeatmasker* v. 4.1.3. [44]. Reads mapping to more than one reference sequence were discarded, and the maximum number of mismatches allowed was three. The resulting files were converted to .bam files and sorted using *SAMtools* v. 1.18 [45].

The reference-based *Gstacks* pipeline was applied to cluster reads into RAD loci using the Marukilow model, minimum mapping quality of 40, and alpha thresholds (for mean and variance) of 0.05 for discovering single nucleotide polymorphisms (SNPs). Using the catalogue of loci generated with *Gstacks*, the *populations* program was then used to proceed to SNP and sample filtering steps and generate SNP datasets adapted to each type of downstream analyses (see **Supplementary Material** and **Table S2** for details).

Inferences of historical changes in effective population size

To evaluate historical events that could have significantly impacted current population genetic diversity and structure, and test whether historical variations in effective population size (N_e) coincide with known past variations in the degree of aridity, we estimated historical changes in N_e using *Stairway Plot 2* v.2.1.2, which conducts multi-epoch coalescent inference to assess changes in population size over time [46,47]. We ran this analysis at two different sample scales: 1) at the *R. bechuanae* species level, pooling individual samples from all localities, and 2) at the locality level: within *R. bechuanae*, analysing two localities representing the two extremes of the Aridity Index across the sampled distribution. At each scale, we applied a downsampling procedure and run the analysis with N_{max} = 19 per locality.

The Python program *easySFS* [48] was first used to generate the folded site frequency spectrum (SFS) formatted file (v0.0.1) [49]. Since demographic inference from the SFS is particularly dependent on the presence of rare alleles (i.e., singletons and doubletons) [50], we used the SNP data set obtained without the *-mac* filter. In all cases, the samples with the least missing data were included, and then projected down, to cover the optimal number of SNPs. For all analyses, the mutation rate was set to 5.7 × 10^{-9} per site per generation (based on the germline mutation rate found in *Mus musculus* in [51]), with a generation time of one year. We considered the entire allele

frequency spectrum with and without singletons (See **Supplementary Material** for details).

Population structure analyses

To describe the structure of genetic diversity within *R. bechuanae*, we first computed differentiation statistics (pairwise F_{ST}) [52] for all sampled localities with sufficient sample sizes using the *vcftools* program v.0.1.16 [53]. To test isolation-by-distance and isolation-by-environment within *R. bechuanae*, simple and partial Mantel tests (R *ncf* package v. 1.3-2) [54] were conducted to test for correlations between genetic differentiation among all *R. bechuanae* localities (using $F_{ST}/(1- F_{ST})$ following [55]) and log-transformed geographic distance, as well as log-transformed environmental distance (using the Aridity Index). To characterise population structure within *R. bechuanae*, we performed a principal component analysis (PCA) and a sparse non-negative matrix factorisation (*sNMF)* analysis using the R package *LEA* v.3.14 [56]. For *sNMF* analyses, we ran 1,500 repetitions for each value of *K* ranging from 1 to 8 and chose the best *K* by evaluating cross-entropy values. Additionally, a neighbour-joining tree was generated using SNP data of these same samples with the R package *ape* (v. 5.7-1) [57].

Genotype-Environment Association analyses

To explore signals of genetic adaptation to aridity, we conducted Genotype–Environment Association (GEA) analyses. Using extensions of F_{ST}-based models, the rationale was that environmental variables distinguishing differentiated populations should be associated with allele frequency differences at loci subjected to the selective constraints they impose [22]. To test for an association between allele frequency differences between all *R. bechuanae* localities and the environmental variable of interest, the Aridity Index, which was computed for all localities using [38], we used the program *BayPass* v.2.4, which implements a method accounting for the neutral correlation of allele frequencies across populations to detect loci associated with covariables [24]. We first pooled individual genotype information from each of the 11 *R. bechuanae* localities, before running *BayPass* under the standard covariate model, which uses an Importance Sampling (IS) approximation to identify SNPs associated with the Aridity Index, and corrects for population structure by using the scaled covariance matrix Ω of population allele frequencies. Three runs (with different starting seeds) were performed to test for convergence. To determine the level of association of each SNP with the Aridity Index and identify SNPs significantly associated, we computed the empirical Bayesian p-value (eBPis) and Bayes factors (*BF*) expressed in deciban units (dB). Three thresholds of *BF* (or degrees of association) were considered, following [58] : 10 dB < *BF* < 15 dB for "strong" evidence; 15 dB < *BF* < 20 dB for "very strong" evidence; *BF* > 20 dB for "decisive" evidence. Additionally, *XtX* statistics [22], which explicitly account for population structure through Ω, were outputted by the

BayPass core model to identify SNPs with significant genetic differentiation among *R.*
bechuanae populations (see **Supplementary Material** and **Figure S1** for details).

To construct an accurate gene annotation file, we mapped the *Mus musculus* GRCm39
assembly annotation [59] onto the *R. pumilio* genome using the default settings of *liftoff*
software version 1.6.3 [60]. SNPs with a *BF* greater than 10 dB, as well as those that
both had a *BF* greater than 10 dB and a significant *XtX*, were mapped onto the
annotation. We used this annotation to identify genes containing SNPs significantly
associated with the Aridity Index (outlier 'association' genes) and among those, genes
containing SNPs both significantly associated and differentiated among populations
(outlier 'association/differentiation' genes). For each gene, we first retrieved a full
description from the Alliance of Genome Resources [61], which was then summarised
to obtain a meaningful set of functional information. We performed functional
enrichment analyses on these two lists of outlier genes using *Metascape*, an online tool
for gene function annotation analysis. The enrichment was initially considered
statistically significant at $p < 0.05$. P-values were then corrected for false discovery
rates according to the Benjamini-Hochberg method. Only enrichments exhibiting a
corrected $p_{adj} < 0.05$ were considered significant.

Results

RAD-seq and genotype dataset

Across all 230 *R. bechuanae* samples analysed, an average of 2,316,057 sequence
reads per sample were mapped against the *R. pumilio* reference genome, after using
process_radtags to filter for quality and *clone-filter* to mask potential PCR duplicates
(see **Table S3** for details). The alignment rate to the repeat-masked *R. pumilio* genome
averaged 55.96 % across *R. bechuanae* samples. The *Gstacks* analysis resulted in an
average of 151,936 ± 1,779 loci per sample. RAD loci were genotyped with a mean per-
sample coverage of 10.94 ± 0.21 X and a mean of 227.88 ± 0.09 base pairs per locus.

The SNP and sample filtering procedures, used to generate the three different *R.*
bechuanae datasets for population structure analyses, inferences of past effective
population sizes and GEA analyses, resulted in 44,900 (population structure; GEA
datasets), and 147,761 (inferences of past effective population sizes dataset)) final sets
of SNPs (**Table S2**). After SNP and genotype quality filtering, missing data were on
average 9.56 ± 0.68 % per sample (**Figure S2**). A total of 154 samples were kept after
individual missing data filtering (< 50%) and downsampling for analysis of population
structure and historical changes in population effective size, while 218 samples were
kept after individual missing data filtering for the GEA (**Tables S2** and **S3**).

 ## Genetic diversity and population structure in time and space

Historical changes in effective population size in *R. bechuanae*

Aridity Index (AI) estimations validated our sampling of *R. bechuanae* along an aridity
gradient in South Africa: AI of sampled localities ranged from 0.0322 (Klein Pella,
classified as hyper-arid) to 0.2457 (Sandveld, classified as semi-arid) (**Table S4**).

We detected no major differences in the temporal patterns of N_e between two distinct
populations at the semi-arid (Sandveld) or hyper-arid (Klein Pella) extremes along the
aridity gradient, nor any evidence of recent population decline (**Figure 1**, green and
orange lines). The two populations showed stable and similiar N_e, from the LGM period
until present, detecting only a difference between the two populations during the LGM
period where Klein Pella population showed a reduction in N_e. When analysing historical
changes in N_e for *R. bechuanae* as a whole, pooling data from all sampled localities,
Stairway Plot 2 inferred a distinct historical trajectory of *Ne* compared with intra-
population inferences (**Figure 1**, red line). Similarly to population-specific inferences, a
steep increase in N_e beginning around 140 thousand years ago (kya) was detected,
followed by a stabilisation around 100 kya, with a median N_e estimate of 4,434,261
individuals reached around 60 kya. From ~30 kya - shortly before the Last Glacial
Maximum (LGM) - the species-level trajectory departed from population-specific
inference, with N_e declining gradually and then sharply from ~15 kya to the present. This
trajectory was also recovered in the analysis excluding singletons (**Figure S3**).
Comparison with palaeoclimatic reconstructions from the central summer rainfall zone
of Southern Africa [62], revealed that this period of sharp N_e decline coincides with
repeated and localised arid episodes occurring at 15 – 11, 7–5 and 1.5–0.5 kya (**Figure
1**).

Contemporary *R. bechuanae* population structure

The analysis of contemporary population structure on all retained *R. bechuanae*
individuals revealed some degree of genetic clustering despite low level of intraspecific
differentiation within this species (**Figure 2**, **Tables S5 and S6**). Pairwise F_{ST} estimates
among *R. bechuanae* sampling localities were relatively low, spanning from 0.010 to
0.058 (**Table S5**). Combined, the Principal Component Analysis (PCA) (**Figure S4**), the
sNMF analysis (**Figure 2A**) and the neighbour-joining tree (**Figure 2C**) did not indicate
a clear-cut separation of several sampling localities or groups of localities from each
other.

The *sNMF* analysis yielded similar cross-entropy values for K = 3 and 4 clusters (**Figure
S5**), suggesting that these two inferences of population structure are almost equally
likely (**Figure 2A**). At the lowest cross-entropy value (K=3; **Figure 2**, **Table S6**), a first
R. bechuanae genetic cluster grouped most localities from a region corresponding to
the Southern Kalahari (Lake Naute, Mariental, Tswalu, Molopo, Kolomela, Sandveld),
which ranged from a hyper-arid to a semi-arid climate according to the estimated Aridity

Index. Another cluster was more prevalent in the Gariep and Tussen die Riviere semi-arid localities, and a third cluster was mostly found in the hyper-arid Klein Pella locality (**Figure 2B**, **Table S6**). Genetic differentiation between these groups of localities was also visible on the neighbour-joining tree (**Figure 2C**). Most individuals from localities located centrally in our sampling scheme (Benfontein, Kolomela Mine) did not have ancestry coefficients > 85 % from one specific cluster and were instead composed of mixed continuous ancestries (**Figure 2C, Table S6**), as also suggested by the PCA plot (**Figure S4**).

Consistent with the gradual pattern of genetic ancestries observed across the sampled localities (**Figure 2**), simple Mantel tests using genetic distances based on F_{ST} values among *R. bechuanae* localities (**Table S5**) showed significant positive correlations between genetic distance and geographic distance (p = 0.038), indicating isolation by distance, as well as between genetic distance and the environmental variable of interest, the Aridity Index (p = 0.037), indicating isolation by environment (**Figure 3**). Since we found a very strong correlation between geographic and environmental distances (p <0.001) and no significant partial Mantel tests (**Table S7**), these results suggest that patterns of isolation by distance and by environment are closely intertwined, and that genetic differentiation may be shaped by the aridity gradient, which closely mirrors the geographic distance gradient.

Genetic basis of adaptation to aridity

Genotype-Environment Association analysis and genomic scan for differentiation

To take advantage of the full aridity gradient across *R. bechuanae*'s area of distribution, we performed the GEA analysis at the locality level using *BayPass*. Given the above correlations, we expected stringent conditions for the GEA analysis. Indeed, since *BayPass* explicitly standardises population allele frequencies for their correlation structure - here induced by the neutral population structure but also the isolation by environment signal - we expected a high rate of false negatives when testing for an association with the Aridity Index.

Overall, despite these stringent conditions, 108 SNPs were found to be significantly associated with the Aridity Index (significance threshold of 10 dB), distributed across all genomic scaffolds (**Figure 4A**, **Table S8**). Among these, 68 were mapped within an annotated coding region of the *R. pumilio* genome, corresponding to 63 outlier 'association' genes in total.

Across the three runs of the model, which gave similar results, a total of 1,682 variants were identified as *XtX* outliers (**Figure 4B**). Of these, 65 SNPs displayed "strong evidence" of association with aridity, as well as a significant *XtX* value. Among them, 45 were mapped within an annotated coding region of the *R. pumilio* genome, corresponding to 41 outlier 'association/differentiation' genes.

Functional analyses

All 63 genes associated with aridity, as well as the subset of 41 genes that were both associated with aridity and showed significant genetic differentiation across *R. bechuanae* populations, had identified homologues in the *M. musculus* genome, according to *Metascape* analyses (for a complete list of outlier genes with functional details, see **Table S8**).

The 41 'association/differentiation' outlier genes corresponded to twelve broad functional physiological pathways (**Table S8**), reflecting diverse metabolic functions. In **Figure 4**, we present one representative gene from each pathway, selected for showing the strongest signal of association with the Aridity Index. Among the most promising candidate genes were those involved in nervous system functions (e.g., *ADGRB1*, *WHAMM,* see **Table S8**). Notably, Gene Ontology (GO) terms related to neurotransmission showed the strongest degree of enrichment ($p_{adj} < 0.01$) among genes associated with the Aridity Index, with twelve GO terms significantly enriched overall **(Figure 5, Table S9)**.

Several genes were also linked to maintaining cellular integrity and stability of genetic or protein information. These included genes associated with osmoregulation (e.g., *EFCC1*), cellular response to stress (e.g., *SLC37A3*), and protein metabolism (e.g., *TMEM44*). Both gene sets – those associated with aridity and those both associated and differentiated – contained genes involved in the regulation of transcription (e.g., *ESRP2*, see **Table S8**). Relevant GO terms were significantly enriched among genes associated with aridity (**Figure 5**), and approached significance ($p_{adj} = 0.063$) among associated/differentiated genes **(Table S9)**. Furthermore, GO terms related to cytoskeleton regulation and GTPase activity were also near the threshold for enrichment significance in the associated/differentiated gene set ($p_{adj} = 0.063$) (**Table S9**), with genes such as *WHAMM* or *EEFSEC* among top candidates (**Figure 4, Table S8**).

Finally, several genes with potential relevance for energy conservation were identified. These included genes involved in lipid metabolism (e.g., *SDR9C7*), carbohydrate metabolism (e.g., *GNAQ*), immunity (e.g., *STING1*), and sweet taste perception (*TAS1R2*, see **Table S8**).

Discussion

In this study, we generated RAD-seq data to investigate patterns of intraspecific genetic diversity and responses to aridity in the African four-striped mouse, *Rhabdomys bechuanae*, a rodent species broadly distributed across an aridity gradient in Southern Africa. By developing an original integrative approach, combining genomic data with fine-scale contemporary and historical aridity data available for the Southern African region and leveraging a natural aridity gradient to investigate signatures of aridity-driven

intraspecific divergence using GEA analyses, our study provides significant insights into population-level responses to increased aridity in a homeothermic rodent species.

Our results suggest a recent decline in effective population size at the species level, coinciding with Holocene aridification. Additionally, we found that genetic diversity within the species is shaped by spatial structure of the environment, closely linked to variation in aridity across its distribution range. We detected allele frequency shifts associated with aridity levels, enabling identification of candidate genes and functions potentially involved in adaptation to arid environments. As predicted, we identified candidate genomic regions associated with classical desert survival phenotypes, including DNA repair/protein modification, osmoregulation and carbohydrate/fat metabolism [10,16]. However, remarkably, the strongest functional enrichment signal emerged for genes involved in neuronal transmission functions, bringing novel insights into the role of neurological processes in coping with osmotic stress and resource scarcity. Together, these findings support the hypothesis that genetic variation in *R. bechuanae* reflects both its demographic history, shaped by past climatic fluctuations, and the selective pressures imposed by spatial variation in aridity across its distribution range [63,64].

Population structure and historical changes in effective population size in response to aridification

Although genetic differentiation amongst *R. bechuanae* localities was relatively low overall, our analyses revealed genetic clustering (three clusters) with the *sNMF* approach, along with significant correlation between genetic and geographical distances. This intraspecific structure was also supported by both PCA and neighbour-joining analyses. The observed pattern of isolation by distance provided evidence of restricted gene flow across the species' distribution range, which aligns with the patchiness of favourable habitat available for *R. bechuanae* in the arid and semi-arid regions of Southern Africa [25,65]. Isolation by distance results from constraints on dispersal, through habitat patchiness for example, thereby promoting the population structure observed here despite the species' relatively continuous regional distribution. Our findings support this hypothesis through the detection of significant isolation by environment associated with the Aridity Index variable, although this variable showed strong correlation with geographic distance. The population structure analysis identified genetic clusters that correspond to populations at the opposing ends of the aridity gradient (Klein Pella representing the arid extreme, and Sandveld the semi-arid extreme), both of which have relatively peripheral positions in the species' range. This spatial arrangement suggests that the observed genetic structure could reflect adaptation to distinct environmental conditions along the aridity gradient, which may have imposed divergent selective pressures on *R. bechuanae* populations, supporting the hypothesis that aridity is a key driver of population structure in this species.

Historical demographic events, such as range expansions, population bottlenecks or divergence due to local adaptation or neutral processes, may have shaped patterns of population structure and diversity in *R. bechuanae* and may be partly related to past

climatic events. This motivated us to infer historical changes in effective population size
and correlate results with known past climatic events.

A marked decline in the hyper-arid locality Klein Pella around the Last Glacial Maximum
was observed, suggesting that biome changes induced by the last deglaciation in
Southern Africa [66] partly drove *R. bechuanae* population history. According to
palaeoclimatic reconstructions from central Southern Africa, aridification took place in
the *R. bechuanae* distribution range in the late Pleistocene [62]. During this time, the
Southern Kalahari experienced a gradual increase in temperature [67] that may have
contributed to N_e reduction in ancestral populations observed at the arid extreme,
upholding our prediction of N_e decline during dry periods. In more recent times, single-
locality effective population size (N_e) was largely stable and similar between 'arid' or
'semi-arid' localities. The similar effective population size trends inferred in relatively
peripheral localities at each environmental 'extreme' in recent times can be attributed to
low migrant rates [68].

Different trajectories between single-locality and species-level inferences were also
observed, as a decline in N_e was inferred at the species level from 15 thousand years
ago, but not in any of the localities on their own.

In this study, it seems unlikely that the recent reduction in N_e at the species level
reflects changes in effective population size : these patterns could instead
be consistent with a strong effect of structuration within *R. bechuanae*. Indeed, signals
of past changes in population size in such inferences can often reflect stable population
structure or variations in population connectivity [68-70]. However, this effect is likely a
complex mix of intra- and inter-deme effects and difficult to fully analyse with SFS-
based inference methods [68]. Given the coincidence between the period when a signal
of reduced N_e is detected and repeated and localised arid past episodes [62],
structuration and connectivity may have been influenced by fluctuation in aridity.
Specifically, arid conditions and their impact on the vegetation may alter population
dynamics of striped mice [71], as well as the distribution of nesting sites, concentrating
them in scattered patches of appropriate vegetation [65], and lack of favourable habitat
may create barriers that impede dispersal and gene flow between populations [72].

The patterns of genetic diversity emerging from aridification remain relatively
unexplored in the literature; previous studies on vertebrate models report a link between
past changes in aridity and genetic diversification [73], or changes in connectivity [74].
Overall, our results support the hypothesis that past climatic fluctuations in the Southern
Kalahari and central South Africa may have been important factors shaping effective
population size and population structure of *R. bechuanae,* although other neutral and
adaptive factors could also be involved. Historical periods of increased aridity could not
only have reduced population connectivity but may have also created opportunities for
local adaptation if geographically separated populations experienced different
environmental pressures along the aridity gradient. This dual effect of aridity, both
fragmenting populations and creating divergent selective environments, provides a
comprehensive explanation for the observed genetic structure in *R. bechuanae*.

This study enabled the identification of several SNPs associated with the Aridity Index
variable, despite the very stringent conditions of the *BayPass* analysis due to significant
correlations between genetic, geographic and environmental (Aridity Index) distances.
While these conditions reduced the overall power of the analysis, they also
strengthened the reliability of the identified SNPs. Additional approaches, such as
genome-wide association studies based on well characterised physiological phenotypes
[30], could complement our findings by revealing other genomic regions underlying key
traits for adaptation to aridity. It is unlikely our inferences were biased by small sample
sizes in some localities, as GEA studies using Bayes' factor are not particularly
sensitive to sample sizes [75]. Refining our candidate list of outlier SNPs by focusing on
SNPs located within coding regions and exhibiting significant genetic differentiation
among populations allowed us to identify the most promising outlier SNPs and
candidate genomic regions.

Our hypothesis of differential selective pressures acting on *R. bechuanae* populations
along the aridity gradient was supported by the identification of outlier SNPs located
within coding regions of genes associated with important biological functions for
adaptation to an arid or desert lifestyle. We identified genes involved in energetic
balance (e.g., lipid metabolism, immunity) and osmotic balance (e.g., osmoregulation,
neuronal processes, body fluids regulation), as well as cellular responses to dehydration
(e.g., protein metabolism, regulation of transcription and cytoskeleton). *Rhabdomys*
bechuanae face environmental constraints, which can involve high temperatures and
resource scarcity. Even in semi-arid conditions, these challenges cause a strain on
homeostasis, as revealed in *R. bechuanae* populations from semi-arid locations by
blood markers of osmoregulation, immunity and energy levels [30]. More arid conditions
could cause cellular stress, and exacerbate the effects of dehydration due to lack of
water and rapid water loss (e.g., respiratory water loss, evaporative cooling). Overall,
consistent with our predictions, our results suggest that various physiological
mechanisms may enable *R. bechuanae* to cope with harsh conditions and inhabit drier
environments. We synthesised previous findings from other species aligning with our
results in **Table S8**, and discuss in more detail the functions of interest for adaptation to
arid conditions in light of this literature review.

In numerous studies carried out on mammalian species, the same genes, or those
involved in similar functions as reported in our study, are highlighted as containing
SNPs varying with an aridity climatic index (e.g., Moisture index [76]), differentially
expressed in arid conditions (e.g., water restriction [20]), or showing signs of selection in
specialist organisms (e.g. [16]). Many of these gene lists identified from previous studies
and in our study intersect, suggesting a genetic basis for adaptation to aridity partly
shared across various mammals. Studies on the basis of adaptation to xeric
environments show convergent changes in tissue composition, gene expression as well
as coding sequences, with significant overlap mostly among related species [10,77].

Our findings align with key shared mechanisms of adaptation to low resource availability observed across mammals, especially those involved in maintaining energetic balance, such as lipid metabolism [10,13,19,78], and immune response [79,80]. Interestingly, acquiring energy in a food-scarce environment may also involve diet shifts in response to arid food sources, such as desert plants and insects. These shifts can be facilitated by mutations in taste receptor genes, such as the sweet taste receptor identified in our study or in the numbat *Myrmecobius fasciatus* [81], and bitter taste receptors in the *Peromyscus eremicus* cactus mouse [15].

At the cellular level, dehydration-induced changes, such as cytoskeletal rearrangements [82], may explain the observed association between aridity and genes regulating the cytoskeleton (e.g., GO terms: filopodium assembly), suggesting possible adaptations to hyperosmotic stress in the striped mouse. However, dehydration also affects cells through protein denaturation, transcription/translation suppression, altered adhesion, increased DNA breaks and protein oxidation, cell cycle arrest, and apoptosis [82-84]. Consistent with these responses, SNPs associated with aridity were found in genes involved in protein synthesis and modification, DNA repair, and cellular cycle regulation. Genes containing outlier SNPs were also enriched for GO terms related to transcription regulation. These findings align with previous results from selective sweeps studies in arid-adapted rodents (e.g., *Peromyscus eremicus* [15]), and meta-analyses of metazoan genomic and transcriptomic data showing selection on translation-related genes in harsh environments [85]. Indeed, cellular responses to hyperosmotic stress commonly involve regulating protein translation and degradation [84]. Overall, our results suggest strong selection on genes mitigating thermal or hyperosmotic stress and restoring cellular function after acute stress [83].

Arid environments impose strong selective pressures on osmoregulation mechanisms. In *R. bechuanae*, we identified genes containing significantly associated outlier SNPs involved in osmoregulation, many of which were also reported in other desert rodents (e.g., *Allactaga sibirica*, *Dipus sagitta*, *Meriones meridianus* and *Phodopus roborovskii* [16], **Table S8**). Water balance in mammals is regulated by brain-body communication mediated by central nervous system structures that exhibit marked plasticity in response to environmental cues in arid-adapted species [86,87]. This may help explain the strong enrichment of GO terms related to nervous system functions, such as nervous system development and neurotransmitter regulation, which emerged as the most significant signal in our study. Although neurological functions are occasionally noted in aridity adaptation studies (e.g., [13,75,88,89]), they are rarely emphasised as key survival factors under arid conditions. When mentioned, they are typically discussed in the context of selective brain cooling in large, heat-adapted mammals [90,91]. Neurotransmitters, essential for maintaining homeostasis [92], regulate blood flow – a critical function in arid environments where dehydration can restrict organ perfusion [93]. Our identification of outlier genes involved in heart rate regulation and vasculature development - also differentially expressed in other desert-adapted rodents such as *Abrothrix olivacea* [94], supports this. Together, these results support the involvement of a previously underappreciated functional category – neurological pathways - in the

547 genomic response to aridity, offering new directions for research in the evolutionary
548 physiology of aridity adaptation.

Conclusion

This study sheds new light on how historical climatic fluctuations and environmental
gradients shaped population structure and genetically based responses to aridity in
Rhabdomys bechuanae, a rodent species inhabiting some of the most arid regions of
Southern Africa. By integrating genomic data with fine-scale environmental information,
we uncovered signatures of population differentiation and polygenic adaptation along an
aridity gradient, involving functions linked to osmoregulation, energetic balance, and
cellular stress responses. The list of candidate genes and biological functions provides
a valuable foundation for further research on the molecular mechanisms of adaptation
to arid environments. While we focused on coding regions to identify genes and
functions associated with the Aridity Index, future work incorporating transcriptomic and
non-coding genomic data will be essential to assess the role of gene expression
regulation in this species' response to increasing aridity and to disentangle the relative
contributions of coding versus regulatory variation. Moreover, although this study
revealed genetically based signatures of adaptation, avoidance behaviours and plastic
physiological responses likely play a critical role in coping with environmental stress
imposed by seasonal variation in aridity [89]. Future studies manipulating water and
food availability, will be key to elucidating the interplay between phenotypic plasticity
and local adaptation in shaping species' responses to arid conditions.

568
569

References

1. Wiens JJ. Climate-related local extinctions are already widespread among plant and animal species. PLoS Biol. 2016 Dec;14(12):e2001104. (https://doi.org/10.1371/journal.pbio.2001104)

2. Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, et al. European phenological response to climate change matches the warming pattern. Glob Chang Biol. 2006 Oct;12(10):1969–76. (https://doi.org/10.1111/j.1365-2486.2006.01193.x)

3. Chen IC, Hill JK, Ohlemuller R, Roy DB, Thomas CD. Rapid Range Shifts of Species Associated with High Levels of Climate Warming. Science. 2011;333(6045):1024–6. (https://doi.org/10.1126/science.1206432)

4. Easterling DR, Meehl GA, Parmesan C, Changnon SA, Karl TR, Mearns LO. Climate extremes: Observations, modeling, and impacts. Science. 2000 Sept;289(5487):2068–74. (https://doi.org/10.1126/science.289.5487.2068)

5. Naumann G, Alfieri L, Wyser K, Mentaschi L, Betts RA, Carrao H, et al. Global changes in drought conditions under different levels of warming. Geophysical Research Letters. 2018;45(7):3285–96. (https://doi.org/10.1002/2017GL076521)

6. Engelbrecht FA, Steinkopf J, Padavatan J, Midgley GF. Projections of future climate change in southern Africa and the potential for regional tipping points. In: Ecological Studies. Cham: Springer International Publishing; 2024. p. 169–90. (Ecological studies: analysis and synthesis. Berlin, Heidelberg, New York NY).

7. Skelly DK, Joseph LN, Possingham HP, Freidenburg LK, Farrugia TJ, Kinnison MT, et al. Evolutionary responses to climate change. Conserv Biol. 2007 Oct;21(5):1353–5.

8. Willmer P, Stone G, Johnston I. Environmental Physiology of Animals. John Wiley & Sons; 2009.

9. Rymer TL, Pillay N, Schradin C. Resilience to droughts in mammals: A conceptual framework for estimating vulnerability of a single species. Q Rev Biol. 2016 June;91(2):133–76. (https://doi.org/10.1086/686810)

10. Rocha JL, Godinho R, Brito JC, Nielsen R. Life in deserts: The genetic basis of mammalian desert adaptation. Trends Ecol Evol. 2021 July;36(7):637–50. (https://doi.org/10.1016/j.tree.2021.03.007)

11. Malaspinas AS, Westaway MC, Muller C, Sousa VC, Lao O, Alves I, et al. A genomic history of Aboriginal Australia. Nature. 2016 Oct;538(7624):207–14. (https://doi.org/10.1038/nature18299)

12. Rocha JL, Silva P, Santos N, Nakamura M, Afonso S, Qninba A, et al. North African fox genomes show signatures of repeated introgression and adaptation to life in deserts. Nat Ecol Evol. 2023 Aug;7(8):1267–86. (https://doi.org/10.1038/s41559-023-02094-w)

13. Kim ES, Elbeltagy AR, Aboul-Naga AM, Rischkowsky B, Sayre B, Mwacharo JM, et al. Multiple genomic signatures of selection in goats and sheep indigenous to a hot arid environment. Heredity (Edinb). 2016 Mar;116(3):255–64. (https://doi.org/10.1038/hdy.2015.94)

14. Ababaikeri B, Abduriyim S, Tohetahong Y, Mamat T, Ahmat A, Halik M. Whole-Genome
Sequencing of Tarim Red Deer (Cervus Elaphus Yarkandensis) Reveals Demographic History
and Adaptations to an Arid-Desert Environment. Frontiers in Zoology. 2020 Oct;17(1):31.
(https://doi.org/10.1186/s12983-020-00379-5)

15. Tigano A, Colella JP, MacManes MD. Comparative and population genomics approaches
reveal the basis of adaptation to deserts in a small rodent. Mol Ecol. 2020 Apr;29(7):1300–14.
(https://doi.org/10.1111/mec.15401)

16. Cheng J, Peng X, Li H, Feijó A, Xia L, Shenbrot GI, et al. Similar adaptative mechanism but
divergent demographic history of four sympatric desert rodents in Eurasian inland. Commun
Biol. 2023 Jan;6(1):33. (https://doi.org/10.1038/s42003-023-04415-y)

17. Bittner N, Mack KL, Nachman MW. Gene expression plasticity and desert adaptation in
house mice. Evolution. 2021;75:1477–91. (https://doi.org/10.1111/evo.14172)

18. Bridges TE, James NV. THE HYPOTHALAMO☐NEUROHYPOPHYSIAL SYSTEM OF
NATIVE AUSTRALIAN DESERT RODENTS. THE VASOPRESSIN AND OXYTOCIN
CONTENTS OF HYPOTHALAMUS AND POSTERIOR PITUITARY OF NOTOMYS ALEXIS
AND PSEUDOMYS AUSTRALIS COMPARED WITH THOSE OF THE LABORATORY RAT
AND MOUSE IN DIFFERENT STATES OF WATER BALANCE. Australian Journal of
Experimental Biology and Medical Science. 1982;60(3):265-83.
(https://doi.org/10.1038/icb.1982.31)

19. Wu H, Guang X, Al-Fageeh MB, Cao J, Pan S, Zhou H, et al. Camelid genomes reveal
evolution and adaptation to desert environments. Nat Commun. 2014 Oct;5(1).
(https://doi.org/10.1038/ncomms6188)

20. MacManes MD, Eisen MB. Characterization of the transcriptome, nucleotide sequence
polymorphism, and natural selection in the desert adapted mousePeromyscus eremicus. PeerJ.
2014 Oct;2:e642. (https://doi.org/10.7717/peerj.642)

21. Marra NJ, Romero A, DeWoody JA. Natural selection and the genetic basis of
osmoregulation in heteromyid rodents as revealed byRNA☐seq. Mol Ecol. 2014
June;23(11):2699–711. (https://doi.org/10.1111/mec.12764)

22. Coop G, Witonsky D, Di Rienzo A, Pritchard JK. Using environmental correlations to identify
loci underlying local adaptation. Genetics. 2010 Aug;185(4):1411–23.
(https://doi.org/10.1534/genetics.110.114819)

23. Frichot E, Schoville SD, Bouchard G, François O. Testing for associations between loci and
environmental gradients using latent factor mixed models. Mol Biol Evol. 2013 July;30(7):1687–
99. (https://doi.org/10.1093/molbev/mst063)

24. Gautier M. Genome-wide scan for adaptive divergence and association with population-
specific covariates. Genetics. 2015 Dec;201(4):1555–79.
(https://doi.org/10.1534/genetics.115.181453)

25. Ganem G, Dufour CMS, Avenant NL, Caminade P, Eiseb SJ, Tougard C, et al. An update
on the distribution and diversification of Rhabdomys sp. (Muridae, Rodentia). J Vertebr Biol.
2020 July;69(2):1. (https://doi.org/10.25225/jvb.20013)

26. du Toit N, Jansen van Vuuren B, Matthee S, Matthee CA. Biome specificity of distinct
genetic lineages within the four-striped mouse Rhabdomys pumilio (Rodentia: Muridae) from
southern Africa with implications for taxonomy. Mol Phylogenet Evol. 2012 Oct;65(1):75–86.
(https://doi.org/10.1016/j.ympev.2012.05.036)

27. Meynard CN, Pillay N, Perrigault M, Caminade P, Ganem G. Evidence of environmental
niche differentiation in the striped mouse (Rhabdomyssp.): inference from its current distribution
in southern Africa. Ecol Evol. 2012 May;2(5):1008–23. (https://doi.org/10.1002/ece3.219)

28. Dupont LM, Rommerskirchen F, Mollenhauer G, Schefuß E. Miocene to Pliocene changes
in South African hydrology and vegetation in relation to the expansion of C4 plants. Earth and
Planetary Science Letters. 2013;375:408–17. (https://doi.org/10.1016/j.epsl.2013.06.005)

29. Dufour CMS, Pillay N, Avenant N, Watson J, Loire E, Ganem G. Habitat characteristics and
species interference influence space use and nest‐site occupancy: implications for social
variation in two sister species. Oikos. 2019 Apr;128(4):503–16.
(https://doi.org/10.1111/oik.05357)

30. Keilani H, Avenant N, Caminade P, Pillay N, Ganem G. Negative impact of mild arid
conditions in natural rodent populations revealed using markers of physiological condition in
natura. Peer Community Journal. 2025; (https://doi.org/10.24072/pcjournal.523)

31. Baird NA, Etter PD, Atwood TS, Currey MC, Shiver AL, Lewis ZA. Rapid SNP Discovery and
Genetic Mapping Using Sequenced RAD Markers. PLoS ONE. 2008;3(10).
(https://doi.org/10.1371/journal.pone.0003376)

32. Benning JW, Faulkner AE, Moeller DA. Rapid evolution during climate change: demographic
and genetic constraints on adaptation to severe drought. Proceedings of The Royal Society B:
Biological Sciences. 1998;290. (https://doi.org/10.1098/rspb.2023.0336)

33. Ventura-Rojas PD, González-Romero A, Moreno CE, Sosa VJ. Effect of rainfall,
temperature and climate change on the ecology of the rodents of arid zones: a review. Mamm
Rev. 2025 Apr;55(2). (https://doi.org/10.1111/mam.12372)

34. Gomulkiewicz R, Houle D. Demographic and genetic constraints on evolution. Am Nat. 2009
Dec;174(6):E218–29. (https://doi.org/10.1086/645086)

35. Sarabia C, vonHoldt B, Larrasoaña JC, Uríos V, Leonard JA. Pleistocene climate
fluctuations drove demographic history of African golden wolves (Canis lupaster). Mol Ecol.
2021 Dec;30(23):6101–20. (https://doi.org/10.1111/mec.15784)

36. Solmsen N, Johannesen J, Schradin C. Highly asymmetric fine-scale genetic structure
between sexes of African striped mice and indication for condition dependent alternative male
dispersal tactics. Mol Ecol. 2011 Apr;20(8):1624–34. (https://doi.org/10.1111/j.1365-
294X.2011.05042.x)

38. Zomer RJ, Xu J, Trabucco A. Version 3 of the Global Aridity Index and Potential
Evapotranspiration Database. Sci Data. 2022 July;9(1):409. (https://doi.org/10.1038/s41597-
022-01493-1)

39. Etter PD, Bassham S, Hohenlohe PA, Johnson EA, Cresko WA. SNP discovery and
genotyping for evolutionary genetics using RAD sequencing. In: Methods in Molecular Biology.

Totowa, NJ: Humana Press; 2012. p. 157–78. (Methods in molecular biology (Clifton, N.J.)).
(https://doi.org/10.1007/978-1-61779-228-1_9)

40. Cruaud A, Gautier M, Galan M, Foucaud J, Sauné L, Genson G, et al. Empirical assessment
of RAD sequencing for interspecific phylogeny. Mol Biol Evol. 2014 May;31(5):1272–4.
(https://doi.org/10.1093/molbev/msu063)

41. Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA. Stacks: an analysis tool set
for population genomics. Molecular Ecology. 2013;22(11):3124–40.
(https://doi.org/10.1111/mec.12354)

42. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. Genome Biol. 2009 Mar;10(3):R25.
(https://doi.org/10.1186/gb-2009-10-3-r25)

43. Richardson R, Feigin CY, Bano-Otalora B, Johnson MR, Allen AE, Park J, et al. The
genomic basis of temporal niche evolution in a diurnal rodent. Curr Biol. 2023 Aug;33(15):3289-
3298.e6. (https://doi.org/10.1016/j.cub.2023.06.068)

44. Smit AFA, Hubley R, Green P. RepeatMasker Open-4.0. 2013–2015. 2015

45. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. Genome Project Data
Processing Subgroup, The Sequence alignment/map (SAM) format and SAMtools.
Bioinformatics. 2009;(16):2078–9. (https://doi.org/10.1093/bioinformatics/btp352)

46. Liu X, Fu YX. Exploring population size changes using SNP frequency spectra. Nat Genet.
2015 May;47(5):555–9. (https://doi.org/10.1038/ng.3254)

47. Liu X, Fu YX. Stairway Plot 2: demographic history inference with folded SNP frequency
spectra. Genome Biol. 2020 Nov;21(1):280. (https://doi.org/10.1186/s13059-020-02196-9)

48. (https://github.com/isaacovercast/easySFS)

49. Coffman AJ, Hsieh PH, Gravel S, Gutenkunst RN. Computationally Efficient Composite
Likelihood Statistics for Demographic Inference. Molecular Biology and Evolution. 2016
Feb;33(2):591–93. (https://doi.org/10.1093/molbev/msv255)

50. Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency data. PLoS
Genet. 2009 Oct;5(10):e1000695. (https://doi.org/10.1371/journal.pgen.1000695)

51. Milholland B, Dong X, Zhang L, Hao X, Suh Y, Vijg J. Differences between germline and
somatic mutation rates in humans and mice. Nat Commun. 2017 May;8(1).
(https://doi.org/10.1038/ncomms15183)

52. Weir BS, Cockerham CC. Estimating F-statistics for the analysis of population structure.
evolution. 1984;1358-70. (https://doi.org/10.2307/2408641)

53. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, Depristo MA. & 1000 Genomes
Project Analysis Group. Bioinformatics. 2011;27(15):2156–8.

54. Bjornstad ON, Bjornstad MON. Package `ncf. Spatial nonparametric covariance functions.
2016;5:1–1.

55. Rousset F. Genetic differentiation and estimation of gene flow from F-statistics under
isolation by distance. Genetics. 1997 Apr;145(4):1219–28.
(https://doi.org/10.1093/genetics/145.4.1219)

56. Frichot E, François O. LEA: An R package for landscape and ecological association studies.
Methods in ecology and evolution. 2015;6(8):925–9. (https://doi.org/10.1111/2041-210X.12382)

57. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. Bioinformatics. 2019 Feb;35(3):526–28.
(https://doi.org/10.1093/bioinformatics/bty633)

58. Jeffreys H. Oxford Classic Texts in the Physical Sciences. In: Theory of Probability. Oxford:
Oxford Univ. Press; 1961.

59. Mouse Genome Sequencing Consortium, Waterston RH, Lindblad-Toh K, Birney E, Rogers
J, Abril JF, et al. Initial sequencing and comparative analysis of the mouse genome. Nature.
2002 Dec;420(6915):520–62.

60. Shumate A, Salzberg SL. Liftoff: accurate mapping of gene annotations. Bioinformatics.
2021 Jun;37(12):1639–1643. (https://doi.org/10.1093/bioinformatics/btaa1016)

61. Alliance of Genome Resources Consortium. Updates to the Alliance of Genome Resources
central infrastructure. Genetics. 2024 May;227(1). (https://doi.org/10.1093/genetics/iyae049)

62. Chevalier M, Chase BM. Determining the drivers of long-term aridity variability: a southern
African case study. Journal of Quaternary Science. 2016;31(2):143–51.
(https://doi.org/10.1002/jqs.2850)

63. Hereford J, Winn AA. Limits to local adaptation in six populations of the annual plantDiodia
teres. New Phytol. 2008 June;178(4):888–96. (https://doi.org/10.1111/j.1469-
8137.2008.02405.x)

64. Kolbe JJ, Leal M, Schoener TW, Spiller DA, Losos JB. Founder effects persist despite
adaptive differentiation: A field experiment with lizards. Science. 2012 Mar;335(6072):1086–9.
(https://doi.org/10.1126/science.1209566)

65. Dufour CMS, Meynard C, Watson J, Rioux C, Benhamou S, Perez J, et al. Space use
variation in co-occurring sister species: Response to environmental variation or competition?
PLoS One. 2015 Feb;10(2):e0117750. (https://doi.org/10.1371/journal.pone.0117750)

66. Zhao X, Lu X, Meadows M, Dupont L, Mao L, Xu Y, et al. Biome variability in southernmost
Africa since the last deglaciation recorded in marine sediments. Quaternary Science Reviews.
2024;343. (https://doi.org/10.1016/j.quascirev.2024.108912)

67. Lukich V, Ecker M. Pleistocene environments in the southern Kalahari of South Africa. Quat
Int. 2022 Mar;614:50–8. (https://doi.org/10.1016/j.quaint.2021.03.008)

68. Lesturgie P, Planes S, Mona S. Coalescence times, life history traits and conservation
concerns: An example from four coastal shark species from the Indo☐Pacific. Molecular
Ecology Resources 2022;22(2) :554-66. (https://doi.org/10.1111/1755-0998.13487)

69. Heller R, Chikhi L, Siegismund HR. The confounding effect of population structure on
Bayesian skyline plot inferences of demographic history. PloS one. 2013 8(5):e62992.
(https://doi.org/10.1371/journal.pone.0062992)

70. Mazet O, Rodríguez W, Grusea S, Boitard S, Chikhi L. On the importance of being
structured: instantaneous coalescence rates and human evolution—lessons for ancestral
population size inference?. Heredity. 2016 116(4) :362-71.

71. Nater CR, van Benthem KJ, Canale CI, Schradin C, Ozgul A. Density feedbacks mediate
effects of environmental change on population dynamics of a semidesert rodent. J Anim Ecol.
2018 Nov;87(6):1534–46. (https://doi.org/10.1111/1365-2656.12888)

72. Ota H. Geographic patterns of endemism and speciation in amphibians and reptiles of the
Ryukyu Archipelago, Japan, with special reference to their paleogeographical implications.
Researches on Population Ecology. 1998;40:189–204. (https://doi.org/10.1007/BF02763404)

73. Garcia JT, Alda F, Terraube J, Mougeot F, Sternalski A, Bretagnolle V, et al. Demographic
history, genetic structure and gene flow in a steppe-associated raptor species. BMC Evol Biol.
2011;11:333. (https://doi.org/10.1186/1471-2148-11-333)

74. Merchant HN, Ivanova A, Hart DW, García C, Bennett NC, Portugal SJ, et al. Patterns of
genetic diversity and gene flow associated with an aridity gradient in populations of common
mole-rats, Cryptomys hottentotus hottentotus. Genome Biology and Evolution. 2024;16(7).
(https://doi.org/10.1093/gbe/evae144)

75. Forester BR, Lasky JR, Wagner HH, Urban DL. Comparing methods for detecting multilocus
adaptation with multivariate genotype–environment associations. Mol Ecol. 2018
May;27(9):2215–33. (https://doi.org/10.1111/mec.14584)

76. Flori L, Moazami-Goudarzi K, Alary V, Araba A, Boujenane I, Boushaba N, et al. A genomic
map of climate adaptation in Mediterranean cattle breeds. Molecular ecology. 2019;28(5):1009–
29. (https://doi.org/10.1111/mec.15004)

77. Chalopin D, Rey C, Ganofsky J, Blin J, Chevret P. Convergent transcriptomic and genomic
adaptation in xeric rodents. Mouginot M, Boussau B, Pantalacci S, Sémon M, editors. 2025.
(https://doi.org/10.1101/2024.10.02.616319)

78. Sugden LA, Atkinson EG, Fischer AP, Rong S, Henn BM, Ramachandran S. Localization of
adaptive variants in human genomes using averaged one-dependence estimation. Nat
Commun. 2018 Feb;9(1). (https://doi.org/10.1038/s41467-018-03100-7)

79. McDade TW, Georgiev AV, Kuzawa CW. Trade-offs between acquired and innate immune
defenses in humans. Evol Med Public Health. 2016;2016(1):1–16.
(https://doi.org/10.1093/emph/eov033)

80. Brusch GA, Christian KA, Brown GP, Shine R, Denardo DF. Dehydration enhances innate
immunity in a semiaquatic snake from the wet-dry tropics. Journal of Experimental Zoology Part
A: Ecological and Integrative Physiology. 2019;331(4):245–52.
(https://doi.org/10.1002/jez.2260)

81. Peel E, Silver L, Brandies P, Hayakawa T, Belov K, Hogg CJ. Genome assembly of the
numbat (Myrmecobius fasciatus), the only termitivorous marsupial. Gigabyte. 2022.
(https://doi.org/10.46471/gigabyte.47)

82. Burg MB, Ferraris JD, Dmitrieva NI. Cellular Response to Hyperosmotic Stresses.
Physiological Reviews. 2007;87(4):1441–74. (https://doi.org/10.1152/physrev.00056.2006)

83. Kampinga HH. Thermotolerance in mammalian cells protein denaturation and aggregation,
and stress proteins. J Cell Sci. 1993 Jan;104(1):11–7. (https://doi.org/10.1242/jcs.104.1.11)

84. Lamitina T, Huang CG, Strange K. Genome-wide RNAi screening identifies protein damage
as a regulator of osmoprotective gene expression. Proc Natl Acad Sci U S A. 2006
Aug;103(32):12173–8. (https://doi.org/10.1073/pnas.0602987103)

85. Porcelli D, Butlin RK, Gaston KJ, Joly D, Snook RR. The environmental genomics of
metazoan thermal adaptation. Heredity (Edinb). 2015 May;114(5):502–14.
(https://doi.org/10.1038/hdy.2014.119)

86. Rabhi M, Ugrumov M, Polenova O, Bengelloun W, Calas A. Modifications
morphofonctionnelles du système hypothalamo-hypophysaire vasopressinergique après
déshydratation chronique chez la Mérione (Meriones shawi). Ann Endocrinol. 1993;54.

87. Djazouli Alim FZ, Romanova EV, Tay YL, Rahman AYBA, Chan KG, Hong KW, et al.
Seasonal adaptations of the hypothalamo-neurohypophyseal system of the dromedary camel.
PLoS One. 2019 June;14(6):e0216679. (https://doi.org/10.1371/journal.pone.0216679)

88. Yang J, Li WR, Lv FH, He SG, Tian SL, Peng WF, et al. Whole-Genome Sequencing of
Native Sheep Provides Insights into Rapid Adaptations to Extreme Environments. Molecular
Biology and Evolution. 2016 Oct;33(10):2576–92. (https://doi.org/10.1093/molbev/msw129)

89. Blumstein D, Macmanes M. The multi-tissue gene expression and physiological responses
of water deprived Peromyscus eremicus. BMC genomics. 2024;25(1).
(https://doi.org/10.1186/s12864-024-10629-z)

90. Hetem RS, Strauss WM, Fick LG, Maloney SK, Meyer LC, Fuller A, et al. Selective brain
cooling in Arabian oryx (Oryx leucoryx): a physiological mechanism for coping with aridity?
Journal of Experimental Biology. 2012;215(22):3917–24. (https://doi.org/10.1242/jeb.074666)

91. Strauss WM, Hetem RS, Mitchell D, Maloney SK, OaposBrien HD, Meyer LC, et al. Body
water conservation through selective brain cooling by the carotid rete: a physiological feature for
surviving climate change? Conservation Physiology. 2017;5.
(https://doi.org/10.1093/conphys/cow078)

92. Mittal R, Debs LH, Patel AP, Nguyen D, Patel K, OaposConnor G, et al. Neurotransmitters:
The critical modulators regulating gut-brain axis. Journal of cellular physiology.
2017;232(9):2359–72. (https://doi.org/10.1002/jcp.25518)

93. Trangmar SJ, González-Alonso J. New insights into the impact of dehydration on blood flow
and metabolism during exercise. Exercise and sport sciences reviews. 2017;45:146–53.
(https://doi.org/10.1249/JES.0000000000000109)

94. Giorello FM, Feijoo M, DaposElía G, Naya DE, Valdez L, Opazo JC, et al. An association
between differential expression and genetic divergence in the Patagonian olive mouse
(Abrothrix olivacea). Molecular Ecology. 2018;27(16):3274–86.
(https://doi.org/10.1111/mec.14778)

 Figure legends

Figure 1: Historical changes in effective population size in *R. bechuanae*.

Stairway Plot 2 analyses (with singletons) showing historical changes in effective population size (N_e) over the past 130,000 years. The red line indicates the estimated median N_e, for the whole species (N=129, 146 896 SNPs), the Klein Pella locality (hyper-arid climate, N=14, 146 534 SNPs) is represented in orange and the Sandveld locality (semi-arid climate, N=16, 145 767 SNPs) is represented in yellow-green. The lines represent the median, and the buffers represent the 95% confidence interval. The grey rectangle marks the period of the Last Glacial Maximum (LGM), while the yellow rectangles denote recent arid episodes in the Summer rainfall region of Southern Africa [67].

Figure 2: Population structure among *R. bechuanae* samples.

A. *sNMF* results from K=3 to K=4 for *R. bechuanae* samples in each locality. Abbreviations of sampling localities are : Klein Pella (KP); Kolomela Mine (Ko); Lake Naute (LN); Mariental (Ma); Molopo (Mo); Benfontein (Be); Tswalu (Ts); Sandveld (Sa); Soetdoring (So); TdR (Tussen die Riviere); Ga (Gariep Dam). **B.** Map showing the distributions and genetic compositions of *R. bechuanae* sampling localities. Pie diagrams indicate the frequencies of genetic clusters in each locality. Base map: World Topographic Map Esri Standard, Aridity Index layer was computed from a 0.5° global grid, using data from Version 3 of the Global Aridity Index and Potential Evapotranspiration Database [38]. **C.** Neighbour-joining tree built using *ape.* Individuals with an ancestry coefficient > 0.85 for one cluster are coloured according to the colour code of the corresponding genetic cluster used in panel A. *R. bechuanae* individuals without an ancestry coefficients > 0.85 for any genetic cluster are coloured in grey. All analyses were performed using *N*=154 individuals and 44,900 SNPs.

Figure 3: **Correlations between genetic distance and geographic/Aridity Index distances**.

Correlation plots with slopes and correlation coefficient R^2 estimates between genetic distance ($F_{ST}/(1-F_{ST})$) and log(geographic distance), and between environmental distance and geographic distance matrices (orange). Environmental distance is represented by Euclidean distance in terms of the Aridity Index.

Figure 4: **Genotype-Environment Associations and genetic differentiation in *R. bechuanae*.**

A. Manhattan plot of Bayes Factor (*BF*, in decibans) values from *BayPass* (standard
covariate model), showing associations between 44,900 SNPs and the Aridity Index
across 24 chromosome-level scaffolds (x-axis : genomic position). The dashed line
marks the *BF*=10 ("strong" level of association) significance threshold. Outlier SNPs are
in colour; others alternate black/white fill colour by scaffold). **B.** Manhattan plot of
genetic differentiation (*XtX* values) from *BayPass* (core model). The dashed line
indicates the 5% POD significance threshold (*XtX* = 14.4). Colour scheme as in A. **C.**
Joint plot of *BF* and *XtX* values. Thresholds are shown as dashed lines; SNPs
significant for both statistics are in orange. In all plots, we show the names of some
genes containing significantly associated and differentiated SNPs, involved in functions
relevant to the response to arid conditions (for each function, the gene containing the
SNP with the highest BF was highlighted; SNPs within these genes in red). All analyses
were performed using *N*=218 individuals and 44,900 SNPs.

Figure 5: Functional enrichment among genes significantly associated with the
Aridity Index.

Abbreviated description of significantly enriched GO terms ($p_{adj} < 0.05$) are presented
on the left, classified into more general physiological pathways, with level of significance
after Benjamini-Hochberg correction reported on the x-axis. Circle size represents the
number of outlier genes found to be associated with each GO term.

KP Ko LN Ma Mo Be Ts Sa So TdR Ga

K=3

K=4

A.

B.

Mariental

Lake Naute

Molopo

Tswalu

Sandveld

Klein Pella

Kolomela

Benfontein

Soetdoring

Gariep

Tussen die Riviere

Aridity Index

0.65

0 km 400 800 km

K1 K2 K3

K4

C.

LN Ko KP

Ma

Ts

Be

Mo

Sa

TdR

Ga So