Can foraging in pesticide-treated patches become an ecological trap for pollinators?

Modelling visitation patterns on a simulated landscape

Zoltán Tóth[1]

[1]Department of Zoology, Plant Protection Institute, HUN-REN Centre for Agricultural

Research, Budapest, Hungary

Corresponding author: Zoltán Tóth, e-mail: toth.zoltan@atk.hun-ren.hu; ORCID: 0000-0002-

2634-8393

Abstract

Understanding the risks of pesticide exposure in agricultural environments is crucial for maintaining essential agroecosystem services. Behaviour often plays a pivotal role in determining how well species cope with human-altered environments, but in pollinators, its significance in mitigating the exposure level to pesticides and, consequently, the accumulation of associated sub-lethal effects has received little attention. In this study, I extended the individual-based model implemented in the simulation tool BEE-STEWARD to investigate how previously overlooked behavioural processes, such as preference for mass-flowering crop, inadvertent social information (ISI) use, and a foraging bias induced by the pesticide applied on the crop may affect nectar and pollen visit rates in a common pollinator species, the buff-tailed bumblebee (*Bombus terrestris*), on far and closely located crop patches. All investigated behavioural processes increased foragers' visit rate on crop patches but their impact differed substantially in magnitude and context. As expected, preference for the crop was strongly positively associated with both visit rates. ISI use increased the visit rate during nectar foraging in interaction with preference on the close crop patches, and the modelled pesticide-induced foraging bias substantially enhanced pollen visits on both close and far crop patches. Simulation results indicate that if a pesticide applied to the crop causes a behavioural alteration that reduces foragers' ability to assess the level of patch depletion correctly, an ecological trap may emerge for pollinators in pesticide-treated agricultural landscapes.

Keywords: bumblebees; foraging; pesticide exposure; sub-lethal effects; individual-based modelling

Introduction

Pesticide use is an integrated element of intensive agriculture aiming for global food security. However, its environmental impact can threaten human health [1], biodiversity [2] and various ecosystem services [3]. Even though environmentally friendly agricultural approaches have been advocated by both governing and conservation organisations in the last decades, the majority of plant health programs are still based on chemical control [4], so pollinators and natural predators of pests are often at risk due to their frequent exposure to agrochemicals. As exposure risk can negatively impact population stability and species interactions at the ecosystem level [5,6], a better understanding of pesticides' adverse effects on non-target organisms and the way these effects are mitigated is essential to maintain key agroecosystem services and achieve sustainable food production.

The amount of pesticides found in agricultural landscapes is highly inconsistent, and it is not well understood if pollinators can influence their exposure risk when pesticide contamination varies spatially and temporally [7]. Evidence suggests that some pollinating insects, such as honeybees and bumblebees, cannot detect even the lethal concentrations of pesticides in nectar [8,9], so foraging individuals cannot behaviourally mitigate their pesticide exposure level by avoiding contaminated food sources (but see [7] for a lepidopteran counterexample). If so, exposure patterns are most likely determined by how pesticide-treated food patches are distributed within pollinators' foraging range. However, intrinsic and induced preference for specific food sources can modify such distribution-based exposure levels. Natural preference for certain types of flowers (e.g., [10,11]) may enhance exposure risk if those plant species are more often treated with pesticides than alternative food sources. Besides, mass-flowering crops can attract pollinators disproportionally to their patches during seasonal blooms [12]. Pesticide-induced preference may also emerge through physiological dependence, due to the addictive nature of some active ingredients or co-formulants. For instance, experimental findings proved that pollinators can develop a preference for food treated with neonicotinoid pesticides [8,13], and similarly induced preferences were also observed for other substances like alcohol and caffeine [14,15]. Finally, pesticide exposure may alter some specific aspects of foraging behaviour that lead to a shift in optimal food preferences. For instance, prolonged exposure to pesticides may hinder locating the floral scent source [16], or induce biases that result in visiting food sources located further apart [17] or foraging less persistently [18]. Pesticide exposure may also reduce activity [19] or motivation to forage [20,21], or directly alter pollinators' natural preference for the flower types they usually exploit [22,23]. Such

behavioural changes can also result in higher pesticide exposure if the more frequently visited

food patches are more likely to receive – directly or indirectly [24] – regular pesticide

treatments. Natural and pesticide-induced preferences are not mutually exclusive and may

dynamically shape actual exposure patterns together.

The ecological significance of inadvertent social information ('ISI' henceforward) use lies in

the possibility of inducing strong correlations in the behaviours and space use of many

individuals [25]. ISI use is a well-known phenomenon in pollinating insects [26], but its

contribution to exposure risk during foraging has remained unexplored. Bees often rely on

social cues when they do not have personal information about surrounding food sources or

finding them would be more costly [27,28], and so ISI use facilitates the localisation and

exploitation of new food sources [29,30] and helps to avoid predators [31]. The presence of

conspecifics can also provide a cue about the quantity of resources in the visited flowers [32].

However, relying on social information may also limit the time spent searching for alternative

food sources and making independent foraging decisions, resulting in sub-optimal foraging

[33]. For instance, attracting conspecifics to the same foraging patches could facilitate the

spread of foraging biases or induce preference among colony members.

Ecological traps are maladaptive consequences of naturally adaptive behaviour in human-

altered environments [34], and ISI use has previously been shown to be responsible for the

emergence of such traps in group-living mammals (e.g., [35,36]). In pesticide-treated

agricultural landscapes, the interaction between preference for pesticide-treated food patches

and ISI use could lead to elevated exposure patterns. Visiting pesticide-treated food patches

represents a sub-optimal foraging choice if the associated sub-lethal effects negatively affect

individual behaviour, physiology or colony growth. These adverse effects can be further

amplified if ISI use facilitates the exploitation of such patches and thus copying the patch

choice of conspecifics increases colony exposure to the pesticide's sub-lethal effects,

ultimately leading to severe reductions in fitness (e.g., [37,38]).

In this study, I conducted a theoretical investigation about how ISI use, preference for the

crop, and a recently demonstrated pesticide-induced foraging bias can influence the pesticide

exposure risk of pollinators on an agricultural landscape. I used the buff-tailed bumblebee

(*Bombus terrestris*) as the model species, an important pollinator of both crops and

wildflowers in temperate regions [39], and a common visitor of the used mass-flowering crop,

oilseed rape (*Brassica napus*; 'OSR' henceforward), the dominant oil crop in Europe [40]. I

included the above behavioural processes in the individual-based model implemented in BEE-

STEWARD, a research and decision-supporting simulation tool that enables researchers,

policymakers and practitioners to predict the impact of land-use characteristics and

interventions on colony and population dynamics in various bumblebee species [41]. I

examined how patch visitation – and thereby the level of exposure – change with the presence

of ISI use and different parameter values of preference for OSR in a landscape consisting of a

grassland (i.e., natural nesting and foraging habitat) and eight OSR (i.e., foraging habitat

during the crop's flowering period) patches. Moreover, a recent study showed that chronic

exposure to a formulation containing a neonicotinoid insecticide frequently applied in OSR

cultivations [42,43] reduced buff-tailed bumblebees' ability to orient toward the floral blend-

containing scent source [16]. By incorporating a similar effect into the model, I also explored

how a pesticide-induced behavioural alteration may further modulate foragers' pesticide

exposure on this simulated landscape. I assessed the magnitude of the effect of these three

parameters on the visit rate of OSR patches at two different distances (i.e., in close or far OSR

patches) during nectar and pollen foraging (i.e., proportion of nectar visits and proportion of

pollen visits, respectively). In addition, I evaluated the relationship between these visit rates

and proxies for pollination capacity and colony fitness under the default model settings.

Results

Nectar visits on OSR patches

Both preference for OSR and ISI use influenced which patches bees exploited during the OSR

flowering period, but differently. Preference for OSR substantially increased the proportion of

nectar visits to closely located OSR patches (Fig. 2a). In the baseline scenario (i.e., using the

original model setting), as well as in the 'only ISI use' and 'only foraging bias' scenarios, the

relationship between OSR preference and nectar visit proportion was best described by a

second-order polynomial approximation. However, when both ISI use and foraging bias were

present, this relationship was linear (Table S1). When the preference for OSR was set to zero,

the proportion of nectar visits (predicted value [95% CI]) ranged between 0.271 [0.26; 0.282]

and 0.376 [0.363; 0.388] across the four scenarios, with the lowest value observed in the

baseline scenario. At 100% OSR preference, this proportion increased to between 0.497

[0.486; 0.508] and 0.633 [0.62; 0.646]. ISI use also increased nectar visits to close OSR

patches, and the magnitude of its contribution linearly increased with preference; this trend

was observed both in the absence (*intercept* [95% CI]=0.11 [0.1; 0.12]; *β* [95% CI]=0.0003

[0.0001; 0.0005]) and presence of foraging bias (*intercept*=0.07 [0.05; 0.08]; *β*=0.0003

[0.0001; 0.0006]; Fig. 1a inset). In the absence of both OSR preference and ISI use, foraging

bias increased the proportion of nectar visits to a level equivalent to a scenario where 27.5%

of bees preferentially foraged on close OSR patches. This resulted in a modest increase in

nectar visits to these patches (from 0.271 [0.26; 0.282] to 0.311 [0.299; 0.322]). However,

when ISI use was present, foraging bias had no additional effect on visitation rates.

On far OSR patches, the proportion of nectar visits increased linearly with preference for OSR

(Fig. 2b; Table S1). The smallest increase occurred in the 'baseline' scenario (from 0.064

[0.06; 0.067] to 0.116 [0.112; 0.121] (back-transformed values)), while the largest increase

was observed in the 'both ISI use and foraging bias' scenario (from 0.082 [0.078; 0.086] to

0.137 [0.132; 0.142]). Overall, ISI use had a minor effect on the proportion of nectar visits

and did not increase with OSR preference in the absence of foraging bias (*intercept*=0.02

[0.01; 0.03]; *β*=0.0001 [-0.01; 0.01]). In the presence of foraging bias, its contribution was

marginal, showing a slight increase with preference (*intercept*=0.007 [0.0007; 0.0129];

β=0.0088 [-0.0015; 0.019]; Fig. 2b inset). When both OSR preference and ISI use were

absent, foraging bias led to only a slight increase in nectar visits to far OSR patches (0.064

[0.06; 0.067] vs. 0.075 [0.072; 0.077]), similar to a scenario where 23.8% of bees

preferentially visited OSR patches. However, when ISI use was included in the model,

foraging bias had no additional effect.

Pollen visits on OSR patches

Preference for OSR substaintily increased the proportion of pollen visits across all OSR

patches. On close OSR patches (Fig. 3a), this preference nearly quadrupled the proportion of

pollen visits in the 'baseline' scenario (from 0.111 [0.091; 0.131] at zero preference to 0.392

[0.372; 0.412] at 100% preference) and almost doubled it when both ISI use and foraging bias

were present (0.341 [0.327; 0.355] vs. 0.579 [0.564; 0.593]). In contrast, ISI use had little

impact on pollen visits to close OSR patches (Fig. 3a inset). In the absence of foraging bias,

its highest contributions were predicted when OSR preference was either zero (0.036 [0.008;

0.066]) or 100% (0.043 [0.014; 0.071]), with a minimum value of 0.013 [-0.004; 0.03] at

47.3% preference. When foraging bias was present, ISI use contributions did not change with

preference (*intercept* = 0.03 [0.01; 0.05]; *β* = 0.006 [-0.03; 0.04]). Most notably, foraging bias

increased pollen visits to close OSR patches to levels comparable to scenarios where 78.1%

(without ISI use) or 79.8% (with ISI use) of bees preferred OSR. As a result, the proportion of

pollen visits rose from 0.111 [0.091; 0.131] to 0.311 [0.297; 0.325] in the absence of ISI use

and from 0.148 [0.127; 0.168] to 0.341 [0.327; 0.355] in its presence.

The positive effect of OSR preference on pollen visits was also evident on far OSR patches

(Fig. 3b). In all scenarios, the proportion of pollen visits increased linearly with preference

(Table S1), leading to a substantial rise from 0.023 [0.02; 0.027] (back-transformed values) to

0.104 [0.097; 0.112] in the 'baseline' scenario and from 0.09 [0.083; 0.097] to 0.148 [0.139;

0.157] in the 'both ISI use and foraging bias' scenario. ISI use had a negligible effect on

pollen visits, regardless of foraging bias. In the absence of foraging bias, its maximum

predicted contribution was just 0.006 [-0.006; 0.018] at 0% preference, while in its presence,

the contribution remained minimal (*intercept*= 0.001 [-0.011; 0.013]; $\beta_{preference}$=0.002 [-0.018,

0.022]; Fig. 3b inset). Notably, foraging bias alone had a strong effect on pollen visits to far

OSR patches, even when preference for OSR was absent. This effect was comparable to

scenarios where 85.7% of foraging bees preferred OSR without ISI use (increasing from

0.023 [0.02; 0.027] to 0.089 [0.084; 0.095]) and 83.5% preferred OSR with ISI use

(increasing from 0.027 [0.024; 0.031] to 0.09 [0.083; 0.097]).

3.3. Relationship between the proportions of visits and the proxies for pollination capacity

and fitness

In the 'baseline' scenario, the proportion of pollen visits across all OSR patches was

positively related to the total number of adult workers (β=469.37 [79.35; 859.4]) and the

number of colonies (β=55.63 [16.77; 94.5]), but did not increase the total number of

hibernating queens (β=352.4 [-57.92; 762.7]; Fig. 4, Fig. S4). The proportion of nectar visits

across all OSR patches had no such effect on either measure (total number of adult workers:

β= 448.9 [-199.1; 1096.93]; number of colonies: β= 27.98 [-37.4; 93.36]; total number of

hibernating queens: β= 561.5 [-124.99; 1247.99]).

Discussion

The extent to which ecological traits can modify foragers' pesticide exposure and its

subsequent impacts is less known in non-*Apis* bees [44]. Simulation results indicated that all

investigated behavioural processes increased foragers' visit rate on OSR patches in the

simulated landscape where the buff-tailed bumblebee population existed in equilibrium. Still,

their effect differed substantially in magnitude and context. The original model predicted that,

on average, 33.9% of nectar and 14.2% of pollen visits occurred on the close and far OSR

patches. As expected, preference for the crop notably increased both visit rates, so when all

bees preferred collecting resources from OSR during its flowering period, 61.8% of nectar

and 49.8% of pollen visits occurred on these patches. The maximum exposure, however, was

observed when ISI use and the modelled foraging bias was also present, increasing the visit

rates further to 76% in nectar and 73.3% in pollen foraging, respectively. A decisive result is

that the foraging bias alone significantly increased the rate of pollen visits (from 13.4-17.4%

to 40-43.1%) on the crop patches, potentially amplifying this exposure pathway to the

pesticide's additional adverse effects. Under the original model settings, the proportion of

pollen visits on all OSR patches was positively associated with colony growth measures

representing the population's capacity for pollination but not with reproductive performance.

This observation corroborates with the findings of previous field studies that the presence of a

mass flowering crop within the colonies' foraging range does not necessarily translate into a

higher production of sexual offspring [45,46]. Because of that, a preference for mass-

flowering OSR crops is unlikely to be strongly favoured by natural selection. However, if the

pesticide applied to the crop induces a foraging bias – among other effects – that impairs

bees' ability to accurately assess patch depletion, an ecological trap may emerge in pesticide-

treated foraging patches, leading to more pronounced sub-lethal effects on colony growth.

This finding implies that even in the presence of alternative food sources during crop bloom

[47,48], bumblebees may collect resources disproportionally from contaminated resource

patches and the resulting risk of exposure can be prolonged if crops with different pesticide

application schedules (e.g., consecutive in time) are present within their foraging range.

Chronic pesticide exposure often leads to multiple sub-lethal effects [17,49,50]. Previous

studies showed that exposure to field-realistic concentrations of acetamiprid following

application (>3360 ppb; see in [51]) can result in various colony- and individual-level

symptoms in bumblebees including significant reductions in drone count [52,53], drone

weight [53,54], number of pupae [53], number of interactions in the colony, immune

response, worker body size (dependent also on food quality [55]), and biased foraging-related

parameters (some dependent also on weight [16,51]). The conducted simulations indicate that

even a single pesticide-induced foraging bias can change the patch exploitation pattern of

bumblebees in a way that makes other sub-lethal effects more likely to appear. There are

various forms of pesticide-induced biases in cognition or foraging behaviour, such as the

impairment of spatial memory [56], disruption of odour coding and discrimination ability

[57], foraging less persistently [18] or visiting food sources located further apart than

unexposed individuals [58], which might have similar exposure-enhancing effects in

bumblebees. This finding strongly suggests that induced foraging biases may have far-

reaching consequences in pollinating insects, not necessarily on overall consumption but on

the level of pesticide exposure and the likelihood of developing multiple sub-lethal effects.

Exposure to pesticides is known to disrupt within-nest social networks and social interactions

in bumblebees [38,55,59], but the extent to which social information use modulates patch

visitation patterns represents a crucial gap in knowledge (unlike in honeybees [60]). These

simulations provide the first evidence that ISI use increases visitation rates on specific

resource patches (predicted values ranging between 0.007 [0.001; 0.013] and 0.144 [0.132;

0.156]). In accordance with initial expectations and previous findings, ISI use enhanced the

impact of natural preference for the crop during nectar foraging, but its effect was mitigated

by the pesticide-induced foraging bias. In this study, ISI use was modelled as simply as

possible to obtain conservative estimates of its impact, i.e., patches visited by others were

preferred to unoccupied patches irrespective of the number, condition or foraging success of

those conspecifics. However, insects can follow various strategies regarding when to use

social information and whom they should copy [26], so incorporating more complex decision

rules in the model, if characteristic of buff-tailed bumblebees, could refine the obtained

predictions. Second, a few but large patches in the simulated landscape represented a coarse-

scale resolution of resources for the foraging bees. While such spatial distribution of crop

patches is not unrealistic in modern agricultural environments, the contribution of ISI use to

exploitation patterns could be considerably higher than the observed one at a fine-scale

resolution of small resource patches.

Simulations showed a substantial discrepancy in how the modelled foraging bias and ISI use

affected the visit rates on OSR patches during nectar and pollen foraging. Patch choice

decisions related to these two resources differ in the Bumble-BEEHAVE model incorporated

in BEE-STEWARD: the calculation of handling time for nectar foragers follows the model of

Harder [61] and the energetic efficiency of exploiting a patch (i.e., energy gained compared to

the energy needed to exploit given layer) plays a crucial role in determining the most

profitable food source, whereas pollen foraging, in the lack of detailed data and existing

models, is modelled to maximise foraging rate. In addition, the priority of nectar and pollen

foraging is also different in the model, and individuals always search for nectar if the stimulus

for this task in the bee's colony is higher than the bee's associated threshold, regardless of the

stimulus-threshold relationship of the other tasks (including pollen foraging). Since these

differences do not necessarily explain the observed asymmetry in the effect of the investigated

parameters, the possible mechanisms behind the pattern should be explored further, especially

as exposure routes via syrup and pollen may lead to distinct sub-lethal effects. For instance, in

a study where both routes of exposure to acetamiprid were examined within the same

experimental setup [53], exposure via nectar resulted in fewer drones in the examined

bumblebee colonies, while the pollen exposure route significantly reduced drone weight.

Besides, although bees may consume more syrup than pollen, pesticide intakes via the two

routes will depend on the pesticide concentrations in these two resources, thus their capacity

to exert sub-lethal effects may vary [62]. Nevertheless, Lonsdorf et al. [63] recently proved

that pesticide exposure via pollen foraging in bumblebees can be satisfyingly predicted by a

spatially explicit model characterised only by the pesticide level and a distance-based

foraging process from the colony locations. This finding implies that this exposure pathway

could be robustly estimated across heterogeneous agricultural landscapes, even if the

available information is limited.

This study provides *in silico* evidence that previously overlooked aspects of pollinator

behaviour do not mitigate but increase pesticide exposure risk and corroborates with field

observations that mass-flowering crops can competitively dilute pollinator densities in

agricultural landscapes [12,64]. In particular, simulation results imply that a pesticide-induced

foraging bias may lead to greater exploitation of crop patches, increasing bumblebees'

exposure level to the pesticide in pollen without changing overall pollen consumption.

Through positive feedback, such shifts in preference may facilitate the accumulation of sub-

lethal effects of pesticides in real-world scenarios. Future efforts should extend the BEE-

STEWARD model to more behavioural aspects of pollinator foraging that could potentially

influence visitation patterns to differently treated resource patches. In addition, if more sub-

lethal effects could be included in the model, it would allow an evaluation of the conditions

under which their accumulation is likely to occur and the magnitude of their overall impact on

colonies. In that way, we could have an even more powerful tool for predicting the impact of

pesticide exposure on bumblebee population dynamics and exploring intervention options to

mitigate the pesticide-associated risks for pollinators in agricultural landscapes.

Materials and methods

Description of the applied model

Simulations were run in BEE-STEWARD (www.beesteward.co.uk [41]), a software into

which the landscape exploration module of BEESCOUT [65] and the individual-based model

of Bumble-BEEHAVE [66] has been integrated. Pollinator behaviour is determined by stimuli

and thresholds that scale up to colony-level processes, enabling the simulation of population

survival and performance on the landscape level over multiple years. The underlying parts of

the software are parameterised with empirical data from the UK, describing colony dynamics

of six bumblebee species and plant species characteristics, including details on flower shape,

resource production, and flowering period. The model also incorporates energy budgets and

depletion of resources to facilitate the emergence of competition both within and between

pollinator species. With sensitivity analysis and validation, the Bumble-BEEHAVE model has

been demonstrated to make realistic predictions for buff-tailed bumblebee colonies within the

boundaries of independent empirical observations [66]. For a detailed description of the

underlying models and procedures, see Becher et al.'s [65,66] supplementary materials

(Appendices S02 and S05, respectively).

In short, the simulated landscape is 2-dimensional, consisting of 300×210 grid cells and

comprises patches that provide foraging and/or nesting opportunities. Patches are spatially

explicit and implemented as one or more food sources, each representing a single plant

species. If a patch contains multiple food sources, these can be layered, i.e. placed at the same

location and linked with each other. The simulations start at the beginning of the year with an

initial number of hibernating queens that, after emerging from hibernation, search for suitable

nest sites in semi-natural habitats (exact dates drawn from a characteristic distribution).

Queens that survived the searching process and successfully founded a colony must collect

resources to lay the first batch of eggs, then forage and care for the brood simultaneously until

the first adult workers emerge. After that, the queen's activity is limited to egg laying, while

the workers are responsible for nursing and foraging. Foragers' choices on which patch to

exploit are based on maximising the foraging rate for pollen and the energetic efficiency for

nectar, primarily determined by the distance from the colony, handling time, and the amount

of the given resource available in the patch [67]. Colony growth depends on resource

collection, whereas population dynamics result from the number of mated queens that leave

the colony and enter into hibernation until the following spring [66].

Model extensions

I modified the original code by incorporating the parameters 'ISI use', 'preference for OSR'

and 'foraging bias' into the model (Table 1). I also included additional functions in the

extended version to obtain patch-specific nectar and pollen visitation data. To reduce the

computational time of the simulations, the code of the original model was converted from

NetLogo 5.3.1 to 6.4.0 [68].

In the original foraging module, the model sorts the list of known foraging patches by their

distances to the bee's colony. In the presence of 'ISI use', foragers' choice of where to collect

nectar or pollen is influenced by the presence of conspecifics. Individuals' known patches are

first divided into two types: one type is where others are currently foraging, and the other type

is the unoccupied patches. Then, both lists are sorted by the distances of patches and the bee's

colony, and the lists are compiled so the occupied patches appear first in the compiled list (see

the 'Foraging_SortKnownPatchesListREP' function in the supplemented code). The modified

function is later called during both nectar and pollen foraging.

When foragers' activity is set to 'searching', unexperienced foragers and those unsatisfied

with their previous patch search for a new flower patch. In the original model, foragers'

chance of finding a patch is determined by a default detection probability function based on

the distance between a given food source and the colony's location ('DetectionProbREP'

[65]). If 'preference for OSR' is set to a non-zero value, the 'foraging_searchingProc'

procedure (which utilises the detection probability function) is modified in a way that a given

proportion of bees (represented by the provided non-zero value) search for a patch from the

list of available OSR patches during the OSR flowering period, while others bees' patch-

finding process is unaffected by the type of flower present in the patch.

The foraging bias included in the extended model is based on a recent finding of Kárpáti et al.

[16]. This study examined the peripheral olfactory detection of a floral blend and foraging

behaviour of buff-tailed bumblebee workers exposed to acetamiprid, a neonicotinoid

insecticide. One of the main results was that treated bumblebees had a decreased likelihood of

finding the floral blend source (that was associated with the presence of syrup during the

chronic pesticide treatment), resulting in a random choice between the control and blend scent

sources in the behavioural trials. This finding suggests that exposed individuals could not

correctly assess which presented scent source was rewarding. As scent is generally associated

with the presence of resources in flowers [69], I mimicked a similar effect by modifying the

'Foraging_bestLayerREP' function that reports the most profitable layer within the currently

used flower patch based on minimal handling time. In the original model, handling time

('handlingTimePerFlower_s' function) for nectar foraging is determined by the time needed

to travel to the next flower plus the time to test whether it contains nectar, divided by the

filling level of the flower, plus the time needed to collect the nectar once a filled flower is

found. During pollen foraging, handling time is calculated from the time needed to travel to

the next flower multiplied by the number of flowers needed to be visited, divided by the

filling level of the flower, plus the time needed to collect the pollen once a flower with pollen

is found. In the presence of 'foraging bias', filling levels are set to 1, so handling times are not

dependent on actual filling levels when bumblebees are exposed to a pesticide. Still, only

those patches are considered where the filling level > 0. For simplicity, I assumed that all bees

are affected by the pesticide application on OSR patches irrespective of the patch individuals

exploit (as the resources stored within the colonies can also be contaminated), but only during

the OSR flowering period (i.e., from day 114 to day 136).

Simulated agricultural landscape with foraging patches

The landscape consisted of one grassland and eight oilseed rape (*Brassica napus*) patches

(Fig. 1). The central grassland patch can provide nectar and pollen for the bumblebees year-

around and also serves as a nesting habitat where queens can establish new colonies after

emerging from winter hibernation. This patch is a good-quality habitat for bumblebees

containing 22 different flowering plant species [66]. The OSR patches represent habitats that

are unsuitable for nesting and solely consist of oilseed rape crop that flowers from day 114

until day 136 in each simulation year. All patches had a 200 m radius, resulting in a 125 663.7

m^2 patch area. Due to the spatial distribution of OSR patches around the grassland patch, four

OSR patches were closer to the central patch (distance between patch edges: 15 m) than the

diagonally located OSR patches (186.9 m). The former patches were termed 'close OSR

patches', whereas the latter 'far OSR patches'. The exploitation of close and far OSR patches

was examined separately.

Characteristics of the simulation runs

Simulations were run over 50 years in 50 iterations to determine if the default number of total

hibernating queens (set to 500) is appropriate for the model and from which year the

population can be assumed to be in equilibrium [70] (Fig. S1A). Based on the results, I took

year 10 as the year when the bumblebee population was likely to reach equilibrium in the

simulated landscape and thus subsequently analysed the data obtained in the 10th year. The

initial number of hibernating queens was also 500 in all iterations. I ran the simulations 50

times in all combinations of 'ISI use' (absent or present), preference for OSR (ranging from 0

to 1 by 0.1) and foraging bias (present or absent), totalling 2200 simulations. Models were run

individual-based under the 'intermediate foraging mortality' scenario. I also used the setting

where males were not unlimited.

I evaluated the model by its ability to reproduce the following patterns. First, as semi-natural

habitats support pollinators by providing nectar, pollen and nesting resources over the whole

season [46,71], the simulated bumblebee population reaches equilibrium if only the grassland

patch is present (Fig. S1B). Second, preference for OSR and ISI use do not substantially

affect the number of pollen and nectar visits, i.e., these parameters have little impact on bees'

general foraging activity (Fig. S2). Third, preference for OSR and ISI use do not considerably

alter the bumblebee population's pollination capacity and fitness either (Fig. S3).

Analysis of model outputs

All simulations and calculations were performed using R 4.3.2 [72] and RStudio

v2024.04.2.764 [73]. Instead of applying frequentist hypothesis testing during the analysis, I

evaluated the magnitude of differences between simulation runs with different parameter

settings and reported parameter estimates with corresponding uncertainty following the

recommendations of White et al. [74]. The extended model, the R script for the analysis of

model outputs and data supporting the results are archived and available at Figshare

(https://figshare.com/s/acee9613adfc7757267b).

From the simulation outputs, I assembled the following data from the 10th simulation year for

subsequent analysis: the number of visits associated with nectar foraging on each patch

between day 114 and day 136 (i.e., during the OSR flowering period), the number of visits

related to pollen foraging on each patch between day 114 and day 136, total number of adult

workers and number of colonies on day 149, and total number of hibernating queens on day

365. I calculated the proportions of nectar and pollen visits from the patch visits data for the

close and far OSR patches. Then, I used these variables to characterise the magnitude of

exploitation and, with it, the exposure level to the pesticide on these two types of OSR

patches. The sum of the proportions determined for the close and far OSR patches represented

the total exposure in the simulated landscape during nectar and pollen foraging. The total

number of adult workers and the number of colonies variables were used as proxies for the

pollination capacity of the simulated bumblebee population, and the total number of

hibernating queens was considered a proxy for fitness (as in [67]).

I fitted linear polynomial models on the proportion of nectar and pollen visits on the close and

far OSR patches with the preference for OSR as a predictor in four scenarios. In the 'baseline'

scenario, bumblebees' foraging decisions followed the original model settings (i.e., no ISI use

or foraging bias). In the 'only ISI use' scenario, conspecifics' presence in the patches could

affect bees' foraging decision on where to collect resources. In the 'only foraging bias'

scenario, bees suffered a foraging-related behavioural alteration due to the simulated exposure

to acetamiprid on the OSR patches. In the 'both ISI use and foraging bias' scenario, ISI use

and foraging bias was present and could influence bees' foraging behaviour. In one iteration,

the bumblebee population died out before day 136 of the 10^{th} year (in the 'only foraging bias'

scenario), and in an additional one, it went extinct before the end of the 10^{th} year (in the

'baseline' scenario); simulation results from these iterations were excluded from the

corresponding analyses. To better illustrate the role of ISI use, I also fitted linear polynomial

models on the contribution of ISI use to the proportion of nectar and pollen visits calculated

as mean differences between the two scenarios with the absence and presence of ISI use at

each preference value in far and close OSR patches. To obtain differences for these

calculations, I randomly paired corresponding iterations from the two scenarios. For all

models, the optimal number of degrees was determined by calculating the AICc value for

each candidate model (with k ranging from 1 to 3) using the 'AICcmodavg' R package [75].

Model assumptions were checked by plot diagnosis using the 'DHARMa' R package [76].

The final models were the most parsimonious from the candidate model set with appropriate

diagnostics and a good fit (i.e., $\Delta AICc < 2$). From these models, I evaluated the magnitude of

differences between simulation runs with different parameter settings and calculated

parameter estimates with 95% CI to characterise the effect of predictors on specific dependent

variables. In the case of second-order polynomial regressions, critical values were computed

with the help of the 'mosaic' R package [77]. Inverse estimations from the fitted models were

calculated using the 'investr' R package [78].

Acknowledgements

Z.T. was financially supported by the János Bolyai Research Scholarship of the Hungarian

Academy of Sciences (MTA, BO/00634/21/8).

Author contribution statement

The author confirms sole responsibility for the following: conceptualisation, methodology,

data curation, formal analysis, investigation, validation, visualisation, and writing.

Data availability statement

The extended model, the R script for the analysis of model outputs and data supporting the

results are archived and available at Figshare (https://figshare.com/s/acee9613adfc7757267b)

and are available from the corresponding author on reasonable request.

Competing interests statement

The author has no conflict of interest to declare.

Ethics approval

Not applicable.

References

1. Smagghe, G., Hayrabedya, S., Hayrabedyan, K., Topping, C., Kahru, A., Hardy, I. C. W.,
    ... & Norton, M. Neonicotinoids and their substitutes in sustainable pest control.
    EASAC policy report 45. Schaefer Druck und Verlag GmbH. (2023).

2. Sánchez-Bayo, F., & Wyckhuys, K. A. Worldwide decline of the entomofauna: A review of
    its drivers. *Biol. Conserv.* **232**, 8-27. (2019).

3. Rumschlag, S. L., Mahon, M. B., Hoverman, J. T., Raffel, T. R., Carrick, H. J., Hudson, P.
    J., & Rohr, J. R. Consistent effects of pesticides on community structure and ecosystem
    function in freshwater systems. *Nat. Commun.* **11**, 6333;
    https://doi.org/10.1038/s41467-020-20192-2 (2020).

4. Deguine, J. P., Aubertot, J. N., Flor, R. J., Lescourret, F., Wyckhuys, K. A., & Ratnadass,
    A. Integrated pest management: good intentions, hard realities. A review. *Agron.*
    Sustain. Dev. **41**, 38; https://doi.org/10.1007/s13593-021-00689-w (2021).

5. Richman, S. K., Maalouf, I. M., Smilanich, A. M., Marquez Sanchez, D., Miller, S. Z., &
    Leonard, A. S. A neonicotinoid pesticide alters how nectar chemistry affects bees.
    Funct. Ecol. **36**, 1063-1073. (2022).

6. Nicholson, C. C., Knapp, J., Kiljanek, T., Albrecht, M., Chauzat, M. P., Costa, C., ... & Rundlöf, M. Pesticide use negatively affects bumble bees across European landscapes. *Nature* **628**, 355-358. (2024).

7. Olaya‐Arenas, P., Scharf, M. E., & Kaplan, I. Do pollinators prefer pesticide‐free plants? An experimental test with monarchs and milkweeds. *J. Appl. Ecol.* **57**, 2019-2030. (2020).

8. Kessler, S. C., Tiedeken, E. J., Simcock, K. L., Derveau, S., Mitchell, J., Softley, S., ... & Wright, G. A. Bees prefer foods containing neonicotinoid pesticides. *Nature* **521**, 74-76. (2015).

9. Parkinson, R. H., Scott, J., Dorling, A. L., Jones, H., Haslam, M., McDermott-Roberts, A. E., & Wright, G. A. Mouthparts of the bumblebee (*Bombus terrestris*) exhibit poor acuity for the detection of pesticides in nectar. *eLife* **12**, RP89129; https://doi.org/10.7554/eLife.89129.2 (2023).

10. Lanterman Novotny, J., Lybbert, A., Reeher, P., Mitchell, R. J., & Goodell, K. Bumble bee banquet: Genus‐and species‐level floral selection by Midwestern Bombus. *Ecosphere* **14**, e4425; https://doi.org/10.1002/ecs2.4425 (2023).

11. Huang, W., Vallejo-Marín, M., Inouye, D. W., Yang, C. F., & Ye, Z. M. Bumblebees' flower preferences are associated with floral abundance and buzz frequency when buzz-pollinating co-flowering plants. *Entomol. Gen.* **44**, 133-141. (2024).

12. Riggi, L. G., Raderschall, C. A., Fijen, T. P. M., Scheper, J., Smith, H. G., Kleijn, D., ... & Lundin, O. Early‐season mass‐flowering crop cover dilutes wild bee abundance and species richness in temperate regions: A quantitative synthesis. *J. Appl. Ecol.* **61**, 452-464. (2024).

13. Arce, A. N., Ramos Rodrigues, A., Yu, J., Colgan, T. J., Wurm, Y., & Gill, R. J. Foraging bumblebees acquire a preference for neonicotinoid-treated food with prolonged exposure. *Proc. R. Soc. B* **285**, 20180655; https://doi.org/10.1098/rspb.2018.0655 (2018).

14. Arnold, S. E., Dudenhöffer, J. H., Fountain, M. T., James, K. L., Hall, D. R., Farman, D. I., ... & Stevenson, P. C. Bumble bees show an induced preference for flowers when primed with caffeinated nectar and a target floral odor. *Curr. Biol.* **31**, 4127-4131. (2021).

15. Ostap-Chec, M., Opalek, M., Stec, D., & Miler, K. Discontinued alcohol consumption elicits withdrawal symptoms in honeybees. *Biol. Lett.* **17**, 20210182; https://doi.org/10.1098/rsbl.2021.0182 (2021).

16. Kárpáti, Z., Szelényi, M. O., & Tóth, Z. Exposure to an insecticide formulation alters
chemosensory orientation, but not floral scent detection, in buff-tailed bumblebees
(*Bombus terrestris*). *Sci. Rep.* **14**, 14622; https://doi.org/10.1038/s41598-024-65388-4
(2024).

17. Siviter, H., Johnson, A. K., & Muth, F. Bumblebees exposed to a neonicotinoid pesticide
make suboptimal foraging decisions. *Environ. Entomol.* **50**, 1299-1303. (2021a).

18. Ohlinger, B. D., Schürch, R., Durzi, S., Kietzman, P. M., Silliman, M. R., & Couvillon,
M. J. Honey bees (Hymenoptera: Apidae) decrease foraging but not recruitment after
neonicotinoid exposure. *J. Insect Sci.* **22**, 16; https://doi.org/10.1093/jisesa/ieab095
(2022).

19. Baines, D., Wilton, E., Pawluk, A., de Gorter, M., & Chomistek, N. Neonicotinoids act
like endocrine disrupting chemicals in newly-emerged bees and winter bees. *Sci. Rep.* **7**,
10979; https://doi.org/10.1038/s41598-017-10489-6 (2017).

20. Lämsä, J., Kuusela, E., Tuomi, J., Juntunen, S., & Watts, P. C. Low dose of neonicotinoid
insecticide reduces foraging motivation of bumblebees. *Proc. R. Soc. B* **285**, 20180506;
https://doi.org/10.1098/rspb.2018.0506 (2018).

21. Muth, F., & Leonard, A. S. A neonicotinoid pesticide impairs foraging, but not learning,
in free-fying bumblebees. *Sci. Rep.* **9**, 4764; https://doi.org/10.1038/s41598-019-39701-
5 (2019).

22. Gill, R. J., & Raine, N. E. Chronic impairment of bumblebee natural foraging behaviour
induced by sublethal pesticide exposure. *Funct. Ecol.* **28**, 1459-1471. (2014).

23. Phelps, J. D., Strang, C. G., Gbylik-Sikorska, M., Sniegocki, T., Posyniak, A., & Sherry,
D. F. Imidacloprid slows the development of preference for rewarding food sources in
bumblebees (*Bombus impatiens*). *Ecotoxicology* **27**, 175-187. (2018).

24. Ward, L. T., Hladik, M. L., Guzman, A., Winsemius, S., Bautista, A., Kremen, C., &
Mills, N. J. Pesticide exposure of wild bees and honey bees foraging from field border
flowers in intensively managed agriculture areas. *Sci. Total Environ.* **831**, 154697;
https://doi.org/10.1016/j.scitotenv.2022.154697 (2022).

25. Gil, M. A., Hein, A. M., Spiegel, O., Baskett, M. L., & Sih, A. Social information links
individual behavior to population and community dynamics. *Trends Ecol. Evol.* **33**,
535-548. (2018).

26. Grüter, C., & Leadbeater, E. Insights from insects about adaptive social information use.
Trends Ecol. Evol. **29**, 177-184. (2014).

27. Dawson, E. H., Avarguès-Weber, A., Chittka, L., & Leadbeater, E. Learning by
observation emerges from simple associations in an insect model. *Curr. Biol.* **23**, 727-
730. (2013).

28. Jones, P. L., Ryan, M. J., & Chittka, L. The influence of past experience with flower
reward quality on social learning in bumblebees. *Anim. Behav.* **101**, 11-18. (2015).

29. Worden, B. D., & Papaj, D. R. Flower choice copying in bumblebees. *Biol. Lett.* **1**, 504-
507. (2005).

30. Avarguès-Weber, A., & Chittka, L. Local enhancement or stimulus enhancement?
Bumblebee social learning results in a specific pattern of flower preference. *Anim.*
Behav. **97**, 185-191. (2014).

31. Dawson, E. H., & Chittka, L. Bumblebees (*Bombus terrestris*) use social information as
an indicator of safety in dangerous environments. *Proc. R. Soc. B* **281**, 20133174;
https://doi.org/10.1098/rspb.2013.3174 (2014).

32. Leadbeater, E., & Chittka, L. Bumble-bees learn the value of social cues through
experience. *Biol. Lett.* **5**, 310-312. (2009).

33. Avarguès-Weber, A., Lachlan, R., & Chittka, L. Bumblebee social learning can lead to
suboptimal foraging choices. *Anim. Behav.* **135**, 209-214. (2018).

34. Robertson, B. A., & Hutto, R. L. A framework for understanding ecological traps and an
evaluation of existing evidence. *Ecology* **87**, 1075-1085. (2006).

35. Donaldson, R., Finn, H., Bejder, L., Lusseau, D., & Calver, M. The social side of human–
wildlife interaction: wildlife can learn harmful behaviours from each other. *Anim.*
Conserv. **15**, 427-435. (2012).

36. Sigaud, M., Merkle, J. A., Cherry, S. G., Fryxell, J. M., Berdahl, A., & Fortin, D.
Collective decision☐making promotes fitness loss in a fusion☐fission society. *Ecol.*
Lett. **20**, 33-40. (2017).

37. Baron, G. L., Jansen, V. A., Brown, M. J., & Raine, N. E. Pesticide reduces bumblebee
colony initiation and increases probability of population extinction. *Nat. Ecol. Evol.* **1**,
1308-1316. (2017).

38. Crall, J. D., De Bivort, B. L., Dey, B., & Ford Versypt, A. N. Social buffering of
pesticides in bumblebees: agent-based modeling of the effects of colony size and
neonicotinoid exposure on behavior within nests. *Front. Ecol. Evol.* **7**, 51;
https://doi.org/10.3389/fevo.2019.00051 (2019).

39. Hutchinson, L. A., Oliver, T. H., Breeze, T. D., Bailes, E. J., Brünjes, L., Campbell, A. J.,
... & Garratt, M. P. Using ecological and field survey data to establish a national list of

the wild bee pollinators of crops. *Agric. Ecosyst. Environ.* **315**, 107447; https://doi.org/10.1016/j.agee.2021.107447 (2021).

40. Perrot, T., Bretagnolle, V., Acar, N., Febvret, V., Matejicek, A., Grégoire, S., & Gaba, S. Bees improve oil quality of oilseed rape. *Basic Appl. Ecol.* **76**, 41-49. (2024).

41. Twiston□Davies, G., Becher, M. A., & Osborne, J. L. BEE□STEWARD: A research and decision□support software for effective land management to promote bumblebee populations. *Methods Ecol. Evol.* **12**, 1809-1815. (2021).

42. Kathage, J., Castañera, P., Alonso□Prados, J. L., Gómez□Barbero, M., & Rodríguez□Cerezo, E. The impact of restrictions on neonicotinoid and fipronil insecticides on pest management in maize, oilseed rape and sunflower in eight European Union regions. *Pest Manag. Sci.* **74**, 88-99. (2018).

43. Dworzańska, D., Moores, G., Zamojska, J., Strażyński, P., & Węgorek, P. The influence of acetamiprid and deltamethrin on the mortality and behaviour of honeybees (*Apis mellifera carnica* Pollman) in oilseed rape cultivations. *Apidologie* **51**, 1143-1154. (2020).

44. Raine, N.E., & Rundlöf, M. Pesticide exposure and effects on non-*Apis* bees. *Annu. Rev. Entomol.* **69**, 551-576. (2024).

45. Westphal, C., Steffan□Dewenter, I., & Tscharntke, T. Mass flowering oilseed rape improves early colony growth but not sexual reproduction of bumblebees. *J. Appl. Ecol.* **46**, 187-193. (2009).

46. Williams, N. M., Regetz, J., & Kremen, C. Landscape□scale resources promote colony growth but not reproductive performance of bumble bees. *Ecology* **93**, 1049-1058. (2012).

47. Rundlöf, M., Stuligross, C., Lindh, A., Malfi, R. L., Burns, K., Mola, J. M., ... & Williams, N. M. Flower plantings support wild bee reproduction and may also mitigate pesticide exposure effects. *J. Appl. Ecol.* **59**, 2117-2127. (2022).

48. Carturan, B. S., Siewe, N., Cobbold, C.A., & Tyson, R. C. Bumble bee pollination and the wildflower/crop trade-off: When do wildflower enhancements improve crop yield? *Ecol. Modell.* **484**, 110447; https://doi.org/10.1016/j.ecolmodel.2023.110447 (2023).

49. Desneux, N., Decourtye, A., & Delpuech, J. M. The sublethal effects of pesticides on beneficial arthropods. *Annu. Rev. Entomol.* **52**, 81-106. (2007).

50. Tosi, S., Sfeir, C., Carnesecchi, E., & Chauzat, M. P. Lethal, sublethal, and combined effects of pesticides on bees: A meta-analysis and new risk assessment tools. *Sci. Total Environ.* **844**, 156857; https://doi.org/10.1016/j.scitotenv.2022.156857 (2022).

51. Tóth, Z., & Kovács, Z. Chronic acetamiprid exposure moderately affects the foraging

behaviour of buff-tailed bumblebees (*Bombus terrestris*). *Ethology* **130**, e13425;

https://doi.org/10.1111/eth.13425 (2024).

52. Camp, A. A., Williams, W. C., Eitzer, B. D., Koethe, R. W., & Lehmann, D. M. Effects of

the neonicotinoid acetamiprid in syrup on Bombus impatiens (Hymenoptera: Apidae)

microcolony development. *PLoS ONE* **15**, e0241111;

https://doi.org/10.1371/journal.pone.0241111 (2020a).

53. Weitekamp, C. A., Koethe, R. W., & Lehmann, D. M. A comparison of pollen and syrup

exposure routes in Bombus impatiens (Hymenoptera: Apidae) microcolonies:

implications for pesticide risk assessment. *Environ. Entomol.* **51**, 613-620. (2022).

54. Camp, A. A., Batres, M. A., Williams, W. C., Koethe, R. W., Stoner, K. A., & Lehmann,

D. M. Effects of the neonicotinoid acetamiprid in pollen on Bombus impatiens

microcolony development. *Environ. Toxicol. Chem.* **39**, 2560-2569. (2020b).

55. Straub, F., Birkenbach, M., Leonhardt, S. D., Ruedenauer, F. A., Kuppler, J., Wilfert, L.,

& Ayasse, M. Land-use-associated stressors interact to reduce bumblebee health at the

individual and colony level. *Proc. R. Soc. B* **290**, 20231322;

https://doi.org/10.1098/rspb.2023.1322 (2023).

56. Samuelson, E.E., Chen-Wishart, Z. P., Gill, R. J., & Leadbeater, E. Effect of acute

pesticide exposure on bee spatial working memory using an analogue of the radial-arm

maze. *Sci. Rep.* **6**, 38957; https://doi.org/10.1038/srep38957 (2016).

57. Andrione, M., Vallortigara, G., Antolini, R., & Haase, A. Neonicotinoid-induced

impairment of odour coding in the honeybee. *Sci. Rep.* **6**, 38110;

https://doi.org/10.1038/srep38110 (2016).

58. Siviter, H., Richman, S. K., & Muth, F. Field-realistic neonicotinoid exposure has sub-

lethal effects on non-Apis bees: A meta-analysis. *Ecol. Lett.* **24**, 2586-2597. (2021b).

59. Crall, J. D., Switzer, C. M., Oppenheimer, R. L., Ford Versypt, A. N., Dey, B., Brown, A.,

... & de Bivort, B. L. Neonicotinoid exposure disrupts bumblebee nest behavior, social

networks, and thermoregulation. *Science* **362**, 683-686. (2018).

60. Hasenjager, M. J., Hoppitt, W., Cunningham☐Eurich, I., Franks, V. R., & Leadbeater, E.

Coupled information networks drive honeybee (*Apis mellifera*) collective foraging. *J.*

Anim. Ecol. **93**, 71-82. (2024).

61. Harder, L. D. Flower handling efficiency of bumble bees: morphological aspects of

probing time. *Oecologia* **57**, 274-280. (1983).

62. Azpiazu, C., Bosch, J., Viñuela, E., Medrzycki, P., Teper, D., & Sgolastra, F. Chronic oral
exposure to field-realistic pesticide combinations via pollen and nectar: effects on
feeding and thermal performance in a solitary bee. *Sci. Rep.* **9**, 13770;
https://doi.org/10.1038/s41598-019-50255-4 (2019).

63. Lonsdorf, E. V., Rundlöf, M., Nicholson, C. C., & Williams, N.M. A spatially explicit
model of landscape pesticide exposure to bees: Development, exploration, and
evaluation. *Sci. Total Environ.* **908**, 168146;
https://doi.org/10.1016/j.scitotenv.2023.168146 (2024).

64. Holzschuh, A., Dormann, C. F., Tscharntke, T., & Steffan-Dewenter, I. Expansion of
mass-flowering crops leads to transient pollinator dilution and reduced wild plant
pollination. *Proc. R. Soc. B* **278**, 3444-3451. (2011).

65. Becher, M. A., Grimm, V., Knapp, J., Horn, J., Twiston-Davies, G., & Osborne, J. L.
BEESCOUT: A model of bee scouting behaviour and a software tool for characterizing
nectar/pollen landscapes for BEEHAVE. *Ecol. Modell.* **340**, 126-133. (2016).

66. Becher, M. A., Twiston Davies, G., Penny, T. D., Goulson, D., Rotheray, E. L., &
Osborne, J. L. Bumble BEEHAVE: A systems model for exploring multifactorial
causes of bumblebee decline at individual, colony, population and community level. *J.*
Appl. Ecol. **55**, 2790-2801. (2018).

67. Knapp, J. L., Becher, M. A., Rankin, C. C., Twiston Davies, G., & Osborne, J. L.
Bombus terrestris in a mass flowering pollinator dependent crop: A mutualistic
relationship? *Ecol. Evol.* **9**, 609-618. (2019).

68. Wilensky, U. NetLogo. Center for Connected Learning and Computer-Based Modeling.
Northwestern University, Evanston, IL. http://ccl.northwestern.edu/netlogo (1999).

69. Wright, G. A., & Schiestl, F. P. The evolution of floral scent: the influence of olfactory
learning by insect pollinators on the honest signalling of floral rewards. *Funct. Ecol.* **23**,
841-851. (2009).

70. Hui, C. Carrying capacity, population equilibrium, and environment's maximal load. *Ecol.*
Modell. **192**, 317-320. (2006).

71. Rundlöf, M., Persson, A. S., Smith, H. G., & Bommarco, R. Late-season mass-flowering
red clover increases bumble bee queen and male densities. *Biol. Conserv.* **172**, 138-145.
(2014).

72. R Core Team R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/ (2023).

73. Posit Team RStudio: Integrated Development Environment for R. Posit Software, PBC,
Boston, MA. http://www.posit.co/ (2024).

74. White, J. W., Rassweiler, A., Samhouri, J. F., Stier, A. C., & White, C. Ecologists should
not use statistical significance tests to interpret simulation model results. *Oikos* **123**,
385-388. (2014).

75. Mazerolle, M. J. AICcmodavg: Model selection and multimodel inference based on
(Q)AIC(c). R package version 2.3.3, https://cran.r-project.org/package=AICcmodavg
(2023).

76. Hartig, F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)
Regression Models. R package version 0.4.6, https://CRAN.R-
project.org/package=DHARMa (2022).

77. Pruim, R., Kaplan, D. T., & Horton, N. J. The mosaic Package: Helping Students to
'Think with Data' Using R. *R J.* **9**, 77-102. (2017).

78. Greenwell, B. M., & Schubert Kabban, C. M. investr: An R Package for Inverse
Estimation. *R J.* **6**, 90-100. (2014).

Table 1. Additional parameters in the extended BEE-STEWARD model.

Parameter	Description	Range (unit)
Preference for OSR	When this parameter is set above zero, a corresponding proportion of bees search for a patch from the list of available OSR patches during the OSR flowering period.	0-1 (0.1)
ISI use	When using ISI, foragers collect nectar or pollen from patches where conspecifics are present, provided such foraging sites are known.	0-1 (1)[*]
Foraging bias	With foraging bias, filling levels (if >0) are set to 1, making handling times independent of actual filling levels when bumblebees are exposed to a pesticide during the OSR flowering period.	0-1 (1)[*]

[*]either absent (0) or present (1)

Figure legends

Figure 1. Schematic diagram of the simulated landscape consisting of a grassland (green) and

multiple oilseed rape (OSR) patches (yellow). All patches had a 200 m radius (r), resulting in

a 1256.64 m^2 patch area. Due to the spatial distribution of OSR patches around the grassland

patch, four OSR patches were closer to the central patch (d_1: 15 m) than the diagonally

located OSR patches (d_2: 186.9 m). The former patches were termed 'close OSR patches'

(with dotted pattern), whereas the latter 'far OSR patches' (with diagonal pattern). Distances

on the diagram are not on scale.

Figure 2. Relationship between the preference for OSR and the proportion of nectar visits on

the close (A) and far (B) OSR patches in the examined four scenarios ('Baseline': light blue;

'Only ISI use': dark blue; 'Only foraging bias': yellow; 'Both ISI use and foraging bias': dark

grey). Insets indicate the contributions of ISI use to this measure fitted against preference in

the absence (blue) and presence of foraging bias (brown). Symbols in the insets represent

mean differences (i.e., the mean difference in the proportion of nectar visits between the two

scenarios with the absence and presence of ISI use at each preference value), and vertical

lines show the corresponding SEs. All trend lines were fitted using polynomial

approximations with the degree of the corresponding final models. In the main panels, light

red arrows show at which preference value the predicted proportion of nectar visits in the

'baseline' scenario matches the value predicted in the 'only foraging bias' scenario at zero

preference (in the presence of ISI use, foraging bias did not increase the proportion of nectar

visits on either close or far OSR patches; Table S2). In panel b, the y-axis is on the square-

root scale because models were fitted to square-root transformed data.

Figure 3. Relationship between the preference for OSR and the proportion of pollen visits on

the close (A) and far (B) OSR patches in the examined four scenarios ('Baseline': light blue;

'Only ISI use': dark blue; 'Only foraging bias': yellow; 'Both ISI use and foraging bias': dark

grey). Insets indicate the contributions of ISI use to this measure fitted against preference in

the absence (blue) and presence of foraging bias (brown). Symbols in the insets represent

mean differences (i.e., the mean difference in the proportion of pollen visits between the two

scenarios with the absence and presence of ISI use at each preference value), and vertical

lines show the corresponding SEs. All trend lines were fitted using polynomial

approximations with the degree of the corresponding final models. In the main panels, red

arrows show at which preference value the predicted proportion of nectar visits in the

'baseline' scenario matches the value predicted in the 'only foraging bias' scenario at zero

preference (in the absence of ISI use: light red; in the presence of ISI use: dark red; Table S2).

In panel b, the y-axis is on the square-root scale because models were fitted to square-root

transformed data.

Figure 4. Forest plot for the standardised parameter estimates (with corresponding 95% CI) of

the proportion of nectar visits and the proportion of pollen visits on all OSR patches

calculated from the linear regressions fitted on the proxies for pollination capacity and fitness.

The total number of adult workers and the number of colonies were measured on day 149,

whereas the total number of hibernating queens was measured on day 365 of the 10^{th}

simulation year.