A slow unpicking of the landscape thread? Changes in tropical savanna birds

on a gradient of habitat modification over 12 years of monitoring

Alex S. Kutt [A,B,C]

A. School of Ecosystem and Forest Sciences, University of Melbourne, Victoria 3010,

AUSTRALIA

B. School of Natural Sciences, University of Tasmania, Hobart, Tasmania 7005,

AUSTRALIA

C. Research Institute for the Environment and Livelihoods, Casuarina campus, Charles

Darwin University, Darwin, Northern Territory, 0810 AUSTRALIA.

Email: alexanderkutt@gmail.com

ORCID: 0000-0001-9679-2206

Abstract
17

18   *Context*
19   In south-eastern Australia, land clearing has been extensive, resulting in an avifauna that
20   has been significantly depleted. In northern Australian tropical savannas, land clearing is
21   much less but increasing.
22
23   *Aims*
24   To examine the relative change in bird community composition of a Queensland tropical
25   savanna over a 12-year period where sites represent a gradient of habitat modification. I
26   investigate if these changes trigger avifauna disruption in the wider intact woodlands.
27
28   *Methods*
29   The data were collected from 60 sites sampled seven times between 2004 and 2016, using
30   an eight-count repeat census in each site, in each year of survey. The data was examined
31   via a range of non-parametric multivariate method.
32
33   *Key Results*
34   The analysis of bird community composition indicated that variation was not parallel across
35   the habitat modification gradient (i.e., cleared, thinned sites displayed higher temporal
36   change and instability). The strongest directional change and temporal turnover in β-diversity
37   was strongly linked to the gradient of modification from cleared to thinned, to intact.
38
39   *Conclusions*
40   In this study I demonstrated that, where tree clearing has commenced or trees are thinned
41   (even though it is only a small percentage of the total land cover), this initiates bird
42   community instability, though the surrounding intact vegetation seemed to remain unaffected
43   up until 2016.
44
45   *Implications*
46   Regional changes in bird populations can take many decades to appear after substantial
47   disturbance, and continued monitoring of landscapes being progressively cleared, should be
48   a priority to identify and prevent irreversible collapse of the avifauna, such has occurred in
49   south-eastern Australia.
50
51   **Keywords**: woodland birds, conservation, clearing and thinning, community dispersion,
52   temporal variability, beta-diversity, extinction debt, northern Australia

53
54

Introduction

Despite being the single most significant threat to biodiversity globally (Tilman *et al.* 2017), land clearing for agriculture continues at a depressingly high rate in Australia, and especially in the state of Queensland (Reside *et al.* 2017). This clearing is predominantly broad scale and often including pasture improvement using exotic perennial grasses, targeted clearing of productive areas such as alluvial floodplains for crops, or selective the removal of a proportion of the canopy or understory (i.e., thinning) to promote grass growth (Walker and Weston 1990). The extent of clearing at both a local scale and regional can have small to significant impacts on the fauna (Tilman *et al.* 2017) and in the case of birds - a functionally diverse component of woodlands and forests - the effects can be substantial (Hannah *et al.* 2007; Martin and McIntyre 2007). In southern and eastern Australia, land clearing has been extensive, and resulting in an avifauna that has been significantly depleted and therefore much less resilient to extended periods of climate extremes (Bennett *et al.* 2014). As a consequence, the subtropical and temperate woodland bird community in these areas, has been proposed for listing as a threatened ecological community under national legislation (Fraser *et al.* 2019).

The time lag to species decline and extinction after substantial landscape change, is long and can be commence many decades prior to the depletion of species becoming manifest (Liao *et al.* 2022). In Australia, the ongoing loss and fragmentation of woodlands and forests is linked to the regional loss of species (Ford *et al.* 2009) and the reconfiguration of both the bird community and vegetation has promoted a change in such as the dominance and despotism of hyper-aggressive colonising avifauna that continue to promote the reduction of functional groups such as small-bodied birds (Maron *et al.* 2013). In areas where the clearing for pastoralism is still limited, there is evidence that small scale disturbance can initiate changes in the avifauna (Kutt and Fisher 2011), that could signal the commencement of a longer term impact on birds (Hannah *et al.* 2007).

In this study, I examined the change in bird composition in a tropical savanna woodland in sites sampled over multiple years, and representing a gradient of habitat modification from clearing, thinning and intact vegetation. What is notable in this study is that the landscape is predominantly uncleared, unlike other parts of eastern Australia; however, clearing in Queensland continues to cumulatively increase. I investigated a conceptually straightforward question: what is the change in the bird community composition over the 12-year period of survey, after relative recent (1990-2000s) small scale habitat modification events, and does this trigger on-going community change in the wider landscape (i.e., in both the cleared and thinned vegetation and the adjacent intact vegetation). This paper is a companion piece to a study that modelled in more detail the relationship between individual species abundances

and remotely sensed vegetation metrics (Kutt *et al.* 2025c). Articulating how species and

communities change in the initial phase of habitat modification is important for planning,

managing, and mitigating the long-term negative effects of these changes, especially in the

context of overarching climate change.

Methods

Study Area

The study area is situated in the Desert Uplands bioregion in central-north Queensland (Fig.

1). The vegetation of the region consists of predominantly *Eucalyptus, Corymbia* and *Acacia*

woodlands, open grasslands and ephemeral lakes (DERM 2012). The region is within the

southern more arid area of Australia's tropical savannas, with a long term (1912-2023) mean

annual rainfall of 473 mm (Uanda Station, Bureau of Meteorology 2020). The dominant land

use is cattle grazing, and vegetation clearing in the Desert Uplands bioregion is estimated to

be 7.4%, largely within the Jericho, Bogie River Hills and Prairie – Torrens Creek sub-

regions (the latter sub-region being the location of this study) (Queensland Department of

Environment and Science 2022).

The study was undertaken across four properties (Kalleroo, Penrice, Timaru and Woura

Park stations) located within a 50 km radius of Torrens Creek (20°48'S, 145°00'E) (Fig. 1).

Sixty sites were selected for sampling reflecting three habitat modification types –

predominantly cleared of native vegetation, thinned native vegetation, and intact native

vegetation (hereafter cleared, thinned and intact). All sites sampled in this study were in a

single broad vegetation unit (1: 1,000,000) *Eucalyptus populnea* (poplar box) and *E.*

melanophloia (silver-leaved ironbark) or *E. whitei* (White's ironbark) dry woodlands to open

woodlands on sandplains or depositional plains.The three habitat modification types

including the thinning method is described in Tassicker *et al.* (2006). The details of the

survey periods, and treatment breakdown across the properties is provided (KUTT *et al.*

2025a; Kutt *et al.* 2025c).

Bird surveys

The bird surveys were undertaken seven times between 2004 and 2016, with no data

collected in 2007, 2009-2012 and 2015. All surveys were conducted in the dry season

between May and October, the exact timing dictated by access due to prevailing weather

and permission from property managers. Sixty sites were sampled in 2004 and 2005 before

Kalleroo managers revoked permission to access the property; therefore 50 sites were

sampled in the remaining years. One site at Kalleroo could not be accessed in 2005.

The bird surveys were undertaken within a 1 ha site. In each site, 8 x 10-minute meandering

diurnal bird census were conducted over a four-day period. Two counts were undertaken at

each site on each of the four days: one in the morning between dawn and the three hours

after dawn and the other after this period and before dusk. All birds seen and heard within

the site were recorded, though species flying over the site were not recorded unless they

were interacting with the site (i.e., hunting).

Analysis

For each site and for each survey, I was able to derive total species richness and total

abundance of all birds combined, and abundance of each species. Abundance data were

calculated as the total number of birds of each species recorded during each site surveys

(i.e., total of eight counts), rather than any calculation of absolute density. Though

unadjusted counts can have some biases, the census method used is well-established for

Australian tropical savanna bird communities (Woinarski and Tidemann 1991) and has

proven to be a reliable index for assessing bird community change in many locations in

northern Australia (Hannah *et al.* 2007; Kutt *et al.* 2012b; Kutt *et al.* 2016; Woinarski *et al.*

2012), as well as being a standard state-based approach to bird monitoring (Eyre *et al.*

2018). The scientific and common names, and taxonomic order for birds tabulated, follow the

BirdLife Australia working list of Australian birds version 4.1 (BirdLife 2022).

I examined the changes in bird composition across the habitat modification categories and

years via several multivariate methods. In the first instance I constructed a site by species

abundance array, fourth root transformed the data (to reduce the influence of taxa with high

abundance) and then created a Bray-Curtis resemblance (similarity) matrix. I then examined

the variation in bird composition across the survey years and habitat modification treatments

(and their interaction) via multivariate analysis using the PERMANOVA in PRIMER v7.0.13

(Anderson *et al.* 2008). PERMANOVA is a distance-based, non-parametric, multivariate

analysis of variance that calculates a pseudo *F*-statistic and associated *P*-value by means of

permutations, rather than relying on normal-theory tables (Anderson *et al.* 2008). In essence

this test determines if the centroids (the average point) of different groups (year, habitat

modification and the interaction) are significantly different.

Secondly, I examined the homogeneity of multivariate dispersions via PERMDISP (also in

PRIMER v7.0.13), which statistically assesses if the spread of data points (sites) is

significantly different among groups. I undertook three analyses: for year and habitat

modification alone (i.e., the stability of the entire bird community in each year and each

habitat category), and then for year and habitat modification categories combined (i.e., 2005

cleared, 2005 thinned, 2005 intact, etc). For visualization, mean distances to centroids were

plotted as boxplots for each treatment across all years (intact, thinned, cleared), with jittered

points showing site-level variability. The distances to group centroids were calculated using

the *betadisper* function in the vegan package (Oksanen *et al.* 2022) for each treatment in

each year, providing a measure of within-group multivariate spread. Greater dispersion

values indicate higher temporal or spatial heterogeneity in species composition within

treatments.

I also compiled the trajectory metrics in two dimensional NMDS to quantify the magnitude

and consistency of temporal change via path length (total change through time), net

displacement (first–last year distance), and tortuosity (i.e., path length / net displacement). A

greater path length and tortuosity indicate larger and more erratic compositional changes

(Collins *et al.* 2000). To help visualise these patterns, I created an ordination (non-metric

MDS) of the habitat modification and year centroids and overlaid a minimum spanning tree.

I examined temporal β-diversity turnover in bird assemblage composition within each

habitat-modification category in each year. For each habitat-modification category, the Bray–

Curtis dissimilarity centroid was calculated between the centroid of the baseline year (2004)

and the centroid for each subsequent year, providing a measure of compositional divergence

through time. The resulting temporal trajectories were plotted for each treatment to visualise

changes in assemblage similarity to the baseline. Linear models (*dissimilarity ~ year*) were

fitted separately for each treatment to estimate the rate of compositional change over time

(slope of dissimilarity per year), with steeper positive slopes indicating faster divergence and

greater temporal turnover. These analyses were undertaken using *vegan* (Oksanen *et al.*

2022) and R 4.4.3 (R Core Team 2024).

To visualise and test how bird assemblage composition in modified habitats changed

through time relative to reference conditions, I also used a Principal Response Curve

analysis (ter Braak and Šmilauer 2015) using the *prc* function in vegan (Oksanen *et al.* 2022)

and R 4.4.3 (R Core Team 2024). PRC is a constrained ordination technique derived from

redundancy analysis that extracts the main temporal pattern of treatment effects relative to a

specified reference level. In this case, the model structure was specified as *Bird Composition*

~ Habitat Modification × Year, using the intact sites as the reference treatment, against

which trajectories for the thinned and cleared treatments were expressed through time. The

PRC curves indicate the deviation of each modification category from the reference across

the years, and the proportion of total variance explained by the first PRC axis represents the

dominant trajectory of compositional change relative to the reference. PRC also supplies a

species weighting vector that indicates how strongly and in which direction individual species

contribute to those compositional deviations. High positive values denote species that

increased in relative abundance as habitats became more modified (i.e., cleared sites), while

negative values identify species that remained largely restricted to intact sites.

Finally for completeness, I examined the variation in species abundance across years in

each habitat modification class, using the Kruskal-Wallis non-parametric analysis of variance

in Statistica version 13.5.0.17 (TIBCO Software Inc 2018). I tabulated the average

abundance (and standard error) for all species for each year and habitat modification

category separately, and then indicated which species demonstrated significant variation

across each category.

All raw data is available via the scientific data repository *figshare* (Kutt *et al.* 2025b).

Results

This study recorded a total of 21,899 birds representing 116 species from the 389 sites over

the seven years of sampling. The most frequently recorded species were Weebill *Smicrornis*

brevirostris (n=341 sites), Singing Honeyeater *Gavicalis virescens* (n=296), Rufous Whistler

Pachycephala rufiventris (n=282), Striated Pardalote *Pardalotus striatus* (n=282) and Yellow-

throated Miner *Manorina flavigula* (n=265). The most abundantly recorded species were

Weebill (n=3263 records), Budgerigar *Melopsittacus undulatus* (n=1400), Variegated Fairy-

wren *Malurus lamberti* (n=1161), Zebra Finch *Taeniopygia castanotis* (n=1090) and Yellow-

throated Miner (n= 872). Twelve bird species were only recorded once, and 10 of these were

only via a single record. Considering the distribution of species across the habitat

modification categories, 23 species were only recorded in a single category, while 70

species were recorded in all three; for years, this breakdown was 15 in a single year and 48

in every year.

For all sites combined, the highest mean site species abundance was in 2005, and highest

mean site species richness was in 2014 (Table 1). Considering cleared sites alone, highest

abundance was in 2005, and highest species richness in 2013, for thinned sites alone,

highest abundance was in 2005, and highest species richness in 2014 and for intact sites,

highest abundance was also in 2005, and highest species richness in 2014 (Table 1).

Variation in individual species abundance for each habitat modification category across the

years is presented in Supplementary Table 1. As this is not the direct focus of the

manuscript, these results are not considered further, except to note that in cleared sites 41

species indicated significant variation in abundance across the years of sampling (from a

total of 101 species recorded), in thinned sites 22 species indicated significant variation in

abundance across the years of sampling (from a total of 85 species recorded), and in intact

sites 55 species indicated significant variation in abundance across the years of sampling

(from a total of 110 species recorded).

The PERMANOVA results indicated that there was a strong overall effect of year (df = 6,

Pseudo-F = 6.94, P(perm) = 0.001), habitat modification (df = 2, Pseudo-F = 25.4, P(perm) =

0.001) and the interaction (df = 20, Pseudo-F = 2.43, P(perm) = 0.001) on bird community

composition. The larger Pseudo-F (a measure of effect size) for habitat modification

suggests this is a stronger effect than year and the interaction. The PERMDISP indicated

that was a strong deviation from the centroid (i.e., evidence of dispersion) for habitat

modification, (F = 47.3, df1 = 2, df2 = 386, P(perm): 0.001) but less so for year (F = 10.0, df1

= 6, df2 = 382 P(perm) = 0.001) and the interaction (F = 11.3, df1 = 20, df2 = 368 P(perm) =

0.001). The boxplots of the multivariate dispersion provide further detail of the pattern (Fig.

2), namely, that intact sites consistently showed the lowest distances to group centroids,

indicating relatively homogeneous and stable assemblage composition within years,

whereas thinned sites exhibited intermediate levels of dispersion, with modest year-to-year

fluctuations and cleared sites displayed the greatest within-group dispersion across most

years, reflecting higher spatial or temporal heterogeneity in community composition.

In NMDS space, cleared sites exhibited the greatest total path length (1.64), followed by

thinned (1.35) and intact (0.70), indicating the largest cumulative change in cleared habitats.

However, tortuosity, reflecting irregular or non-directional change, was highest in Thinned

(38.4), compared with Cleared (8.52) and Intact (6.11). This pattern suggests that thinned

assemblages changed most erratically, whereas intact communities were the most stable

through time (Fig. 3).

Temporal β-diversity analyses revealed clear differences in the rate of compositional change

among habitat-modification treatments (Fig. 4). Bird assemblages in cleared sites exhibited

the greatest and most consistent increase in Bray–Curtis dissimilarity from baseline

conditions through time, indicating progressive and substantial community turnover. The

thinned sites showed intermediate levels of change, with moderate increases in dissimilarity

suggesting ongoing but less pronounced compositional shifts. Finally, the intact habitats

remained relatively stable across survey years, with only minor departures from baseline

assemblages. Linear regression slopes of temporal β-diversity against year confirmed this

pattern. The strongest positive slope occurred in cleared sites (Δ = 0.0487 yr^{-1}, p ≈ 0.06, R²

= 0.54), followed by thinned (Δ = 0.0322 yr^{-1}, p ≈ 0.07, R² = 0.51) and intact (Δ = 0.0370 yr^{-1},

p ≈ 0.27, R² = 0.24).

The overall PRC model was significant (F = 4.12, p = 0.001; 999 permutations), indicating

that bird community composition in modified treatments changed markedly and directionally

through time relative to intact habitats. The first PRC axis accounted for most of the habitat

modification related temporal variance, indicating a clear separation between the intact

reference data and the two modified habitats (Fig. 5). Bird community composition in cleared

habitats exhibited the greatest and most consistent deviation from intact bird community

baseline, reflecting progressive compositional change over the years. The thinned sites

habitats showed intermediate divergence, with less fluctuation and intact habitats remained

close to the baseline, indicating high temporal stability in community composition. The

species weights (greatest magnitude, either positive or negative) indicated species

characteristic of the gradient from the intact to modified (cleared and thinned sites) were

typical of good condition woodlands (e.g., Grey-crowned Babbler, Grey Shrike-thrush, Spiny-

cheeked Honeyeater, Brown Treecreeper, Striated Pardalote, Rufous Whistler) and

disturbed or cleared woodlands (e.g., Zebra Finch, Galah, Black-faced Woodswallow,

Australian Magpie, Brown Falcon, Crested Pigeon) (Fig. 6).

Discussion

The clearing of native vegetation in many parts of the world, including north-eastern and

south-eastern Australia, have caused a substantial past and continuing disarray of bird

communities and species (Berryman *et al.* 2024; Thomson *et al.* 2015). The primary cause

of bird community decline in Australia is land clearing associated with agriculture

(Lindenmayer 2022b). In this study I examined the pattern of bird species in a series of sites

(sampled over multiple years), representing the preliminary stages of more extensive land

clearing that has already occurred elsewhere in Australia. In general, there were well-defined

differences in species composition across the cleared and intact tropical savanna

woodlands, and more muted intermediate differences with the thinned vegetation. The

concept of land clearing thresholds for the prevention of species decline and local extirpation

have been well articulated (McAlpine *et al.* 2002), yet the attrition of native vegetation

continues in a dispiritingly substantial manner on many land tenures with little effective

legislative oversight (Ward *et al.* 2019). In this study I found that small scale clearing and

thinning triggers a degree of instability in the bird community in the cleared sites, which may

be the precursor of long-term more permanent depletion of the birds in the modified habitat,

even when the surrounding matrix is still intact. The data did not indicate a change in the

adjacent intact vegetation, but it is conceivable that over time (i.e., decades) this disruption

will influence the surrounding vegetation, for example via edge effects, exotic pasture

invasion or changes in stocking rates.

I identify a few limitations to the study and analysis. Firstly, as each site in each year, was

censused eight times, occupancy analysis may have provided more detailed and revealing

data regarding the changes in bird species over time in each modification category

(Montague-Drake *et al.* 2009). However, over the years of survey, the data was not

consistently structured in the data base (i.e., summaries versus each count separately) to

allow this analysis. Secondly, even though the study spanned 12 years, the sites were not

annually monitored, which provides some gaps the annual patterns, and any evidence of

unusual or predictable changes from year to year. Lastly, site based environmental factors,

including changes to the site management of these pastoral properties (i.e., stocking rates,

exotic pasture incursion, regrowth or re-clearing), which does not allow for a more nuanced

understanding of the mechanisms of change in the bird communities across the sites and

habitat modification categories. Any future work that may revisit these sites, should attempt

to address these issues. Regardless, this study provides a unique insight into the changes

over a 12-year period of a bird community at the initial phase of disturbance.

The analysis of bird community composition change indicated that the variation was not

parallel across the three habitat modification treatments (i.e., the significant interacting effect

in the PERMANOVA). Clearing and thinning have a strong effect on the avifauna in the study

area, but the temporal changes differed on the gradient of modification. Though

heterogenous landscapes can support multiple bird communities and increase the bird

diversity across the wider matrix, woodland and forest areas are numerically the most

significant for bird conservation (Lu *et al.* 2024). However the species in cleared areas are

not necessarily equivalent to natural grasslands, despite species overlap, and this may be in

part of changes in resources, such as available diet (Houston *et al.* 2023) and differences in

ground cover. In general, when land clearing continues, the bird depletion continues

(Lindenmayer 2022a) and only woodland restoration can reverse this effect (Bennett *et al.*

2022). In this study I found that disturbance increases community heterogeneity, but with

less stability in the modified sites. Evidence from the sites further south to the study area

with a greater extent of landscape modification and fragmentation (Hannah *et al.* 2007)

indicated that bird species richness in cleared sites were substantially reduced (8.1 species

per site compared to this study 14.4 [average over all years combined]) whereas species

richness in intact sites was remarkably coincident (19.9 species versus 19.8 in this

study[average over all years combined]).

The bird assemblages underwent the strongest directional change through time in the

cleared sites, with a slight trajectory towards the intact sites in 2008, which then reversed.

Thinned sites remained closer to the intact sites. This pattern is driven by disturbance-

tolerant taxa, or species that fluctuate with rainfall events (the study sites being on the edge

of more arid bioregion). Woodland bird assemblages in Australia can be highly variable from

year to year (Maron *et al.* 2005), and these patterns can be driven by climatic events

(Bennett *et al.* 2014), associated resource pulses or declines (Collett *et al.* 2022) as well as

seasonal regional or continental migration patterns (Chan 2001). The species that were

characteristic of, and influential in the temporal trajectories of cleared were granivores that

can demonstrate distinct pulses of abundance (i.e., Little Button-quail, Galah, Budgerigar,

Zebra Finch), species that typically occur in cleared and disturbed habitat (i.e., Crested

Pigeon, Australian Magpie), and species that occur in natural grasslands (i.e., Australian

Kestrel, Ground Cuckoo-shrike, Black-faced Woodswallow). Conversely the species in the

intact sites were characteristic of intact and good condition woodlands (Fraser *et al.* 2019).

This contrast is typical of those occurring across a treed to treeless gradient, such as

woodland dependent birds (i.e., species in Meliphagidae, Pachycephalidae, Acanthizidae)

and species associated with grasslands and bare ground (Barrett *et al.* 1994) and the

woodland-grassland nexus in tropical savannas (Kutt and Fisher 2011; Kutt *et al.* 2012a;

Woinarski *et al.* 2000; Woinarski and Tidemann 1991).

The temporal turnover in β-diversity in the study sites indicated that this pattern was strongly

linked to the gradient of modification from cleared to thinned, to intact. In addition, there are

large directional community shifts in the cleared sites (and a lesser degree in thinned sites)

compared to the intact sites which are relatively stable. Disturbance driven variability can be

an important precursor to more persistent ecological change (Fraterrigo and Rusak 2008),

and though variability can be natural, driven by, for example, climatic fluctuations (Recher

and Davis 2014), vegetation changes can have a stronger effect on bird composition

compared to weather patterns (Pavey and Nano 2009). The mechanisms for these species

decline in south-eastern Australia, where there is more substantial habitat loss and

fragmentation, include reduced dispersal and higher nest predation and then reduced

breeding leading to local extirpation (Ford *et al.* 2009). Though in this study I did not have

site based environmental data to demonstrate the oscillation in bird community is linked to

an equivalent fluctuation in, and possible reduction in, key resources for species, there is

ample evidence, albeit on a larger scale of clearing, that once clearing commences, this

unlocks a cascading influence on both resources, threats and habitat conditions, that create

an eventual species deficit (Ford *et al.* 2009).

Changes in the vertebrate fauna of tropical savannas in northern Australia are evident, which

mostly have become manifest in the mammal fauna (Fisher *et al.* 2014). The data for bird

assemblages seem to suggest more resilience poor land management (Woinarski *et al.*

2012), though there is likely a geographical effect. In some areas of north-eastern Australia

where overgrazing has significantly altered intact vegetation (i.e., permanently altered the

ground cover, and removed the shrub layer), the avifauna can also be depleted (Kutt and

Fisher 2011). In this study I have demonstrated that, where tree clearing has commenced or

trees are thinned (even though it is only a small percentage of the total land cover), this

seems to initiate instability in these modified habitats. During the timespan of this study, the

intact bird communities did not follow this pattern of high variability. Regional changes in bird

populations can take many decades to appear after substantial disturbance (i.e., extinction

debt Szabo *et al.* 2011), though at a landscape scale where change is multi-directional or

more limited, these can be both debits and credits (Haddou *et al.* 2022). The sites described

in this study were last sampled in 2016, and surveys have been discontinued. Long term

monitoring is a critical action for improved landscape management and conservation

(Lindenmayer *et al.* 2022). In this study I present data that hints at an unpicking of the

landscape condition for woodland birds; extension, enhancement or even recasting of long-

term research, such as those presented here, should be ongoing, and include more specific

focus on biological mechanisms for species variation, and if, over time, the adjacent intact

vegetation also succumbs to gradual disruption of the woodland bird community.

Acknowledgements

I am grateful for the help of landholders in granting us access to their properties at Woura

Park, Timaru, Penrice and Kalleroo, and the many people who assisted in the surveys, from

the Australian CSIRO and Queensland Department of Environment and Science. This

project was supported by the Queensland Government and the Australian Government

(CSIRO, National Heritage Trust, Tropical Savanna Cooperative Research Centre, TERN).

All surveys were conducted under the Queensland Government Scientific Purposes Permit

number WITK04645707.

References

Anderson MJ, Clarke KR, and Gorley RN (2008) 'PERMANOVA+ for Primer. Guide to
Software and Statistical Methods.' (University of Auckland and PRIMER-E Ltd: Plymouth
UK)

Barrett GW, Ford HA, and Recher HF (1994) Conservation of woodland birds in a
fragmented rural landscape. *Pacific Conservation Biology* **1**, 245-256. doi:
https://doi.org/10.1071/PC940245.

Bennett AF, Holland GJ, Haslem A, Stewart A, Radford JQ, and Clarke RH (2022)
Restoration promotes recovery of woodland birds in agricultural environments: A comparison
of 'revegetation' and 'remnant' landscapes. *Journal of Applied Ecology* **59**, 1334-1346. doi:
https://doi.org/10.1111/1365-2664.14148.

Bennett JM, Nimmo DG, Clarke RH, Thomson JR, Cheers G, Horrocks GFB, Hall M,
Radford JQ, Bennett AF, and Mac Nally R (2014) Resistance and resilience: can the abrupt
end of extreme drought reverse avifaunal collapse? *Diversity and Distributions* **20**, 1321-
1332. doi: https://doi.org/10.1111/ddi.12230.

Berryman AJ, Butchart S. H. M., Jackson M. V., Legge S. M., Olah G., Thomas J., Woinarski
J. C. Z., and Garnett ST (2024) Trends and patterns in the extinction risk of Australia's birds
over three decades. *Emu - Austral Ornithology* **124**, 55-67. doi:
10.1080/01584197.2023.2289999.

BirdLife (2022) The BirdLife Australia Working List of Australian Birds; Version 4.1. (Ed. B
Australia): Downloaded from http://www.birdlife.org.au/documents/BWL-
BirdLife_Australia_Working_List_v4.1.xlsx.)

Bureau of Meteorology (2020) 'Average annual, seasonal and monthly rainfall. Available at
http://www.bom.gov.au/climate/data/index.shtml [Verified 09 October 2020].'

Chan K (2001) Partial migration in Australian landbirds: a review. *Emu* **101**, 281-292.

Collett SJ, Crewe TL, Radford IJ, Garnett ST, and Campbell HA (2022) Sympatric finches
increase trophic and spatiotemporal partitioning to enable coexistence through periods of
low resource availability. *Functional Ecology* **36**, 2138-2151. doi:
https://doi.org/10.1111/1365-2435.14111.

Collins SL, Micheli F, and Hartt L (2000) A method to determine rates and patterns of
variability in ecological communities. *Oikos* **91**, 285-293. doi: https://doi.org/10.1034/j.1600-
0706.2000.910209.x.

DERM (2012) 'Biodiversity Planning Assessment, Desert Uplands Bioregion Landscape
Expert Panel Report, Central West Region.' (Department of Environment and Resource
Management, Queensland Government: Brisbane)

Eyre T, Ferguson D, Hourigan C, Smith G, Mathieson M, Kelly A, Venz M, Hogan L, and
Rowland J (2018) 'Terrestrial vertebrate fauna survey assessment guidelines for
Queensland (Version 3).' (Department of Environment and Science, Queensland
Government: Brisbane)

Fisher DO, .Johnson CN, Lawes MJ, Fritz SA, McCallum H, Blomberg SP, VanDerWal J,
Abbott B, Frank A, Legge S, Letnic M, Thomas CR, Fisher A, Gordon IJ, and Kutt AS (2014)
The current decline of tropical marsupials in Australia: is history repeating? *Global Ecology
and Biogeography* **23**, 181-190. doi: 10.1111/geb.12088.

Ford HA, Walters JR, Cooper CB, Debus SJS, and Doerr VAJ (2009) Extinction debt or
habitat change? – Ongoing losses of woodland birds in north-eastern New South Wales,
Australia. *Biological Conservation* **142**, 3182-3190. doi:
https://doi.org/10.1016/j.biocon.2009.08.022.
480
481 Fraser H, Simmonds JS, Kutt AS, and Maron M (2019) Systematic definition of threatened
482 fauna communities is critical to their conservation. *Diversity and Distributions* **25**, 462-477.
483 doi: 10.1111/ddi.12875.
484
485 Fraterrigo JM and Rusak JA (2008) Disturbance-driven changes in the variability of
486 ecological patterns and processes. *Ecology Letters* **11**, 756-770. doi:
487 https://doi.org/10.1111/j.1461-0248.2008.01191.x.
488
489 Haddou Y, Mancy R, Matthiopoulos J, Spatharis S, and Dominoni DM (2022) Widespread
490 extinction debts and colonization credits in United States breeding bird communities. *Nature
491 Ecology & Evolution* **6**, 324-331. doi: 10.1038/s41559-021-01653-3.
492
493 Hannah D, Woinarski JCZ, Catterall CP, McCosker JC, Thurgate NY, and Fensham RJ
494 (2007) Impacts of clearing, fragmentation and disturbance on the bird fauna of Eucalypt
495 savanna woodlands in central Queensland, Australia. *Austral Ecology* **32**, 261-276.
496
497 Houston WA, Black RL, and Wormington KR (2023) Grasslands of cleared woodlands have
498 lower invertebrate diversity and different assemblages to remnant woodlands in grazed
499 landscapes of eastern Australia. *Journal of Insect Conservation* **27**, 999-1011. doi:
500 10.1007/s10841-023-00515-6.
501
502 KUTT A, Healy A, and Hamer RP (2025a) Birdseye in the sky: the relationship between
503 fractional cover, rainfall and woodland birds in a partially modified tropical savanna - data.
504 (The University of Melbourne)
505
506 Kutt AS and Fisher A (2011) Increased grazing and dominance of an exotic pasture
507 (*Bothriochloa pertusa*) affects vertebrate fauna species composition, abundance and habitat
508 in savanna woodland. *The Rangeland Journal* **33**, 49-58. doi: doi:10.1071/RJ10065.
509
510 Kutt AS, Healy A, and Hamer RP (2025b) Birdseye in the sky: the relationship between
511 fractional cover, rainfall and woodland birds in a partially modified tropical savanna - data.
512 (The University of Melbourne)
513
514 Kutt AS, Healy AJ, and Hamer RP (2025c) Birdseye in the sky: the relationship between
515 fractional cover, rainfall and woodland birds in a partially modified tropical savanna. *The
516 Rangeland Journal* **47**, https://doi.org/10.1071/RJ25022. doi:
517 https://doi.org/10.1071/RJ25022.
518
519 Kutt AS, Perkins GC, Colman N, Vanderduys EP, and Perry JJ (2012a) Temporal variation
520 in a savanna bird assemblage: what changes over 5 years? *Emu* **112**, 32-38. doi:
521 http://dx.doi.org/10.1071/MU11054.
522
523 Kutt AS, Vanderduys EP, and O'Reagain P (2012b) Spatial and temporal effects of grazing
524 management and rainfall on the vertebrate fauna of a tropical savanna. *The Rangeland
525 Journal* **34**, 173-182. doi: http://dx.doi.org/10.1071/RJ11049.
526
527 Kutt AS, Vanderduys EP, Perry JJ, Mathieson MT, and Eyre TJ (2016) Yellow-throated
528 miners Manorina flavigula homogenize bird communities across intact and fragmented
529 landscapes. *Austral Ecology* **41**, 316-327. doi: http://10.1111/aec.12314.
530

Liao Z, Peng S, and Chen Y (2022) Half-millennium evidence suggests that extinction debts
of global vertebrates started in the Second Industrial Revolution. *Communications Biology* **5**,
1311. doi: 10.1038/s42003-022-04277-w.

Lindenmayer D (2022a) Birds on farms: a review of factors influencing bird occurrence in the
temperate woodlands of south-eastern Australia. *Emu - Austral Ornithology* **122**, 238-254.
doi: 10.1080/01584197.2022.2106875.

Lindenmayer D (2022b) Birds on farms: a review of factors influencing bird occurrence in the
temperate woodlands of south-eastern Australia. *Emu - Austral Ornithology* **122**, 238-254.
doi: 10.1080/01584197.2022.2106875.

Lindenmayer DB, Lavery T, and Scheele BC (2022) Why We Need to Invest in Large-Scale,
Long-Term Monitoring Programs in Landscape Ecology and Conservation Biology. *Current
Landscape Ecology Reports* **7**, 137-146. doi: 10.1007/s40823-022-00079-2.

Lu X, Jia Y, and Wang Y (2024) The effect of landscape composition, complexity, and
heterogeneity on bird richness: a systematic review and meta-analysis on a global scale.
Landscape Ecology **39**, 132. doi: 10.1007/s10980-024-01933-w.

Maron M, Grey MJ, Catterall CP, Major RE, Oliver DL, Clarke MF, Loyn RH, Mac Nally R,
Davidson I, and Thomson JR (2013) Avifaunal disarray due to a single despotic species.
Diversity and Distributions **19**, 1468-1479. doi: 10.1111/ddi.12128.

Maron M, Lill A, Watson DM, and Mac Nally R (2005) Temporal variation in bird
assemblages: How representative is a one-year snapshot? *Austral Ecology* **30**, 383-394. doi:
https://doi.org/10.1111/j.1442-9993.2005.01480.x.

Martin TG and McIntyre S (2007) Impacts of Livestock Grazing and Tree Clearing on Birds
of Woodland and Riparian Habitats. *Conservation Biology* **21**, 504-514. doi:
https://doi.org/10.1111/j.1523-1739.2006.00624.x.

McAlpine CA, Fensham RJ, and Temple-Smith DE (2002) Biodiversity conservation and
vegetation clearing in Queensland: principles and thresholds. *The Rangeland Journal* **24**,
36-55. doi: https://doi.org/10.1071/RJ02002.

Montague-Drake RM, Lindenmayer DB, and Cunningham RB (2009) Factors affecting site
occupancy by woodland bird species of conservation concern. *Biological Conservation* **142**,
2896-2903. doi: https://doi.org/10.1016/j.biocon.2009.07.009.

Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'hara R,
Solymos P, Stevens M, and Szoecs E (2022) vegan: Community ecology package (2.6-4).
CRAN.

Pavey CR and Nano CEM (2009) Bird assemblages of arid Australia: Vegetation patterns
have a greater effect than disturbance and resource pulses. *Journal of Arid Environments*
73, 634-642.

Queensland Department of Environment and Science (2022) Statewide Landcover and
Trees Study – Methodology Overview v1.1. Queensland Government. (Brisbane)

R Core Team (2024) R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Recher H and Davis J, William (2014) Response of birds to episodic summer rainfall in the
Great Western Woodlands, Western Australia. *Australian Zoology* **37**, 206-224. doi:
10.7882/az.2014.011.

588
589 Reside AE, Beher J, Cosgrove AJ, Evans MC, Seabrook L, Silcock JL, Wenger AS, and
590 Maron M (2017) Ecological consequences of land clearing and policy reform in Queensland.
591 *Pacific Conservation Biology* **23**, 219-230. doi: https://doi.org/10.1071/PC17001.
592
593 Szabo JK, Vesk PA, Baxter PWJ, and Possingham HP (2011) Paying the extinction debt:
594 woodland birds in the Mount Lofty Ranges, South Australia. *Emu - Austral Ornithology* **111**,
595 59-70. doi: 10.1071/MU09114.
596
597 Tassicker AL, Kutt AS, Vanderduys E, and Mangru S (2006) The effects of vegetation
598 structure on the birds in a tropical savanna woodland in north-eastern Australia. *Rangeland
599 Journal* **28**, 139-152.
600
601 ter Braak CJ and Šmilauer P (2015) Topics in constrained and unconstrained ordination.
602 *Plant Ecology* **216**, 683-696.
603
604 Thomson JR, Maron M, Grey MJ, Catterall CP, Major RE, Oliver DL, Clarke MF, Loyn RH,
605 Davidson I, Ingwersen D, Robinson D, Kutt A, MacDonald MA, and Mac Nally R (2015)
606 Avifaunal disarray: quantifying models of the occurrence and ecological effects of a despotic
607 bird species. *Diversity and Distributions* **21**, 451-464. doi: https://doi.org/10.1111/ddi.12294.
608
609 TIBCO Software Inc (2018) Statistica (data analysis software system), version 13.5.0.17
610 Available from: http://statistica.io. (TIBCO Software Inc: Palo Alto, CA, USA;)
611
612 Tilman D, Clark M, Williams DR, Kimmel K, Polasky S, and Packer C (2017) Future threats
613 to biodiversity and pathways to their prevention. *Nature* **546**, 73-81. doi:
614 10.1038/nature22900.
615
616 Walker B and Weston E (1990) Pasture development in Queensland–a success story.
617 *Tropical Grasslands* **24**, 257-268.
618
619 Ward MS, Simmonds JS, Reside AE, Watson JEM, Rhodes JR, Possingham HP, Trezise J,
620 Fletcher R, File L, and Taylor M (2019) Lots of loss with little scrutiny: The attrition of habitat
621 critical for threatened species in Australia. *Conservation Science and Practice* **1**, e117. doi:
622 https://doi.org/10.1111/csp2.117.
623
624 Woinarski JC, Connors G, and Franklin DC (2000) Thinking honeyeater: nectar maps for the
625 Northern Territory, Australia. *Pacific Conservation Biology* **6**, 61-80.
626
627 Woinarski JCZ, Fisher A, Armstrong M, Brennan K, Griffiths AD, Hill B, Choy JL, Milne D,
628 Stewart A, Young S, Ward S, Winderlich S, and Ziembicki M (2012) Monitoring indicates
629 greater resilience for birds than for mammals in Kakadu National Park, northern Australia.
630 *Wildlife Research* **39**, 397-407. doi: https://doi.org/10.1071/WR11213.
631
632 Woinarski JCZ and Tidemann SC (1991) The Bird Fauna of a Deciduous Woodland in the
633 Wet-Dry Tropics of Northern Australia. *Wildlife Research* **18**, 479-500.
634
635
636

637 **Figure 1.** The location of the properties where the bird surveys were undertaken in this
638 study. The grey shaded are indicates intact vegetation and the white areas, cleared. Areas
639 that are thinned are included within the grey shaded areas, as they still represent intact
640 vegetation in the mapping product used for this figure.

641
642
643

644   **Figure 2.** Boxplots of multivariate dispersion (Bray–Curtis distance to the group centroid) for
645   bird assemblages under three habitat-modification treatments (intact, thinned, and cleared)
646   across survey years. Points denote individual site distances to the treatment centroid.
647

648
649

Figure 3. Ordination via non-metric multidimensional scaling indicating the centroids for bird
composition for each habitat modification category by year, joined by a minimum spanning
tree. Cleared sites are the open circles (o), thinned sites are the crosses (+) and intact sites
are solid circles (●).

Figure 4. Temporal trajectories of bird assemblage change, expressed as mean Bray–Curtis
dissimilarity from the baseline year for each habitat-modification treatment (intact, thinned,
cleared). Lines represent treatment-specific community turnover through time, with higher
values indicating greater compositional divergence from baseline assemblages. Points show
annual mean dissimilarities, and the horizontal line at zero indicates no change relative to
baseline.
662

663
664
665

Fig. 5. Principal Response Curves (PRC) showing treatment-specific temporal trajectories of bird assemblage composition relative to intact reference habitats. The horizontal zero line represents no difference from the reference condition, with greater deviations indicating stronger compositional divergence.

Figure 6. Species weights from the Principal Response Curve analysis showing the taxa
most strongly contributing to compositional differences among habitat-modification
treatments relative to the intact baseline. The top 15 species characteristic of intact sites and
the top 15 species characteristic of the most modified sites (cleared) are presented. Larger
absolute weights indicate taxa exerting greater influence on treatment separation and
temporal community trajectories. Positive values denote species that increased in relative
abundance as habitats became more modified, while negative values identify species that
declined or remained largely restricted to intact conditions.
681

682
683

684 **Table 1.** The mean (and standard error) of bird abundance and species richness recorded across all surveys for each of the years. Data
685 presented is all sites combined and then for each habitat modification category.
686

Sites	Group	2004	2005	2006	2008	2013	2014	2016	H	p
All sites	Abundance	56.1 (4.1)	**85.8 (12.1)**	49.8 (3.0)	38.7 (2.9)	60.4 (4.4)	54.7 (4.0)	48.0 (3.1)	14.1	<0.05
	Richness	19.2 (0.8)	17.2 (0.7)	17.0 (0.7)	13.8 (0.8)	19.1 (0.7)	**21.0 (0.7)**	19.1 (0.6)	25.1	<0.001
Cleared	Abundance	34.1 (6.1)	**147.1 (58.1)**	36.6 (6.2)	16.6 (3.1)	57.7 (7.7)	47.2 (8.9)	40.0 (6.0)	35.7	<0.001
	Richness	13.4 (1.4)	13.7 (1.3)	12.0 (1.2)	8.9 (1.1)	**18.7 (1.1)**	17.6 (1.1)	15.8 (1.1)	36.7	<0.001
Thinned	Abundance	55.3 (12.0)	73.9 (26.0)	64.4 (5.3)	45.4 (6.8)	36.0 (4.3)	62.2 (3.4)	68.6 (10.2)	10.8	ns
	Richness	14.7 (1.1)	15.2 (1.6)	20.2 (1.1)	14.4 (1.5)	18.6 (2.2)	**26.4 (1.4)**	20.8 (1.8)	23.5	<0.001
Intact	Abundance	61.9 (4.8)	**73.2 (8.0)**	52.7 (3.4)	49.1 (3.2)	66.8 (6.1)	58.3 (4.4)	49.6 (3.4)	13.5	<0.05
	Richness	21.7 (1.0)	18.6 (0.9)	18.8 (0.8)	16.4 (1.0)	19.4 (1.0)	**22.4 (0.8)**	20.9 (0.6)	25.9	<0.001