Research Article

Repeated sleep deprivation selectively reactivates hippocampal CA1

pyramidal neuron

Yutong Wang[1,2], Emily N. Walsh[2,3,5], Junko Kasuya[1], Colton Remedies[4], Jon Resch[1,2],

Lisa C. Lyons[4], and Ted Abel[1,2]

[1]Department of Neuroscience and Pharmacology, Carver College of Medicine, 51 Newton Road,

2-417B Bowen Science Building, University of Iowa, Iowa City, IA 52242, USA.

[2]Iowa Neuroscience Institute, Carver College of Medicine, 169 Newton Road, 2312 Pappajohn

Biomedical Discovery Building, University of Iowa, Iowa City, IA 52242, USA.

[3]Interdisciplinary Graduate Program in Neuroscience, University of Iowa, 356 Medical Research

Center, Iowa City, IA 52242, USA.

[4]Program in Neuroscience, Department of Biological Science, Florida State University, 319

Stadium Drive, Tallahassee, FL32306, USA.

[5]Department of Neurology, Washington University School of Medicine, 660 South Euclid Avenue,

St Louis, MO 63110, USA

Address for Correspondence:

Ted Abel, Ph.D.

Department of Neuroscience and Pharmacology, Iowa Neuroscience Institute, Carver College of

Medicine, University of Iowa, 169 Newton Road, 2314 Pappajohn Biomedical Discovery Building,

Iowa City, IA 52242, United States.

E-mail: ted-abel@uiowa.edu

Author Email Addresses:

Yutong Wang: yutong-wang@uiowa.edu

Emily N. Walsh: enwalsh@wustl.edu

Junko Kasuya: junko-kasuya@uiowa.edu

Colton Remedies: colton.remedies@med.fsu.edu

Jon Resch: jon-resch@uiowa.edu

Lisa C. Lyons: lyons@bio.fsu.edu

Ted Abel: ted-abel@uiowa.edu

Author's ORCID numbers:

Yutong Wang: 0000-0003-3625-0936

Emily N. Walsh: 0000-0003-3638-7238

Junko Kasuya: 0009-0002-9266-2679

Colton Remedies: 0009-0008-6264-3382

Jon Resch: 0000-0002-5409-7771

Lisa C. Lyons: 0000-0002-3294-6744

Ted Abel: 0000-0003-2423-4592

ABSTRACT

Sleep supports a variety of physiological processes, ranging from metabolic to immune system homeostasis, and plays a critical role in cognition and memory. A brief period of sleep loss impairs memory, particularly hippocampus-dependent memories, and alters molecular signaling and synaptic plasticity in the hippocampus. Studies have shown that sleep deprivation (SD), alters neuronal activation as indicated by broad changes in gene expression signatures and by the altered expression of c-Fos, an immediate early gene that functions as a molecular marker of neuronal activity. In the present study, we examined hippocampal subregion-specific c-Fos induction patterns via immunohistochemical staining. We find that CA1 pyramidal neurons exhibit the most robust c-Fos induction after SD. Using an activity-driven ribosomal tagging system and a repeated SD model, we labeled sleep deprivation activated CA1 neurons and observed a population of excitatory neurons in area CA1 that are reactivated by repeated SD. Using the c-Fos-RiboTag system that enables the isolation of ribosomes attached mRNA from labeled neurons, we performed fosTRAP-seq and identified activity-dependent gene expression changes in c-Fos+ CA1 neurons. Our results revealed that synapse organization, protein dephosphorylation, cellular response to endogenous stimulus (such as insulin) are upregulated, whereas mRNA processing and splicing being downregulated. In summary, our study provides a detailed view of the activation of hippocampal neurons after SD, revealing a subset of CA1 pyramidal neurons that have higher sensitivity to the effect of sleep loss, shown as reactivation during repeated SD, allows investigation of molecular changes in neurons specifically impacted by repeated sleep loss. Our work uncovers a population of CA1 pyramidal neurons that are sensitive to repeated sleep loss and sheds light on a possible connection between acute and chronic sleep loss at the cellular and molecular levels.

Keywords: sleep deprivation; hippocampus; CA1; pyramidal neurons; c-Fos; RiboTag; fosTRAP-seq

64   **Graphical Abstract**

INTRODUCTION

Sleep facilitates numerous physiological processes including cognition, learning, and memory consolidation (1). Over 30% of adults in the United States routinely experience insufficient sleep, leading to subsequent health problems (2, 3). Sleep loss has pronounced impacts on cognitive function, with hippocampus-dependent memories being especially vulnerable (4). The hippocampus plays a critical role in consolidating spatial, episodic, and contextual memories in rodents (5) and declarative memory in humans (6). It is comprised of subfields with specialized roles in memory: the Dentate Gyrus (DG) enables pattern separation (7), Cornu Ammonis 3 (CA3) integrates separation and completion (8), and Cornu Ammonis 1 (CA1) recodes CA3 outputs while linking to the neocortex for memory retrieval (9). Brief periods of sleep deprivation (SD) impact hippocampal function via impairments in the cAMP signaling and related long-lasting forms of synaptic plasticity (10-12), suppressed protein synthesis (13), and altered synaptic connectivity (14-16).

In humans, sleep loss can often span several nights or even weeks. Chronic insomnia is related to accelerated cognitive decline and is associated with increased risk of dementia in humans (17, 18). Work using rodents to model chronic sleep restriction (CSR) has demonstrated impairment of hippocampal synaptic plasticity and severe spatial memory deficits that are resistant to recovery sleep (19, 20). CSR also promotes tau hyperphosphorylation and amyloid-β accumulation in rodent models of Alzheimer's Disease (21, 22). However, how repeated bouts of sleep loss affect neuronal activity in the hippocampus and how acute SD-induced impairments progress to CSR-promoted neurodegeneration remains not known.

A widely recognized marker of neuronal activity is the immediate early gene (IEG) c-Fos. The c-Fos protein is rapidly activated during synaptic plasticity associated with learning and memory (23), and its downstream effector genes, such as *Timp1* and *Mmp9*, regulate the neuronal structural and functional plasticity required for memory formation (23). c-Fos expression has

been used as a marker to locate and identify the neuronal populations activated by sleep and

extended wakefulness (24-27). While sleep is typically linked to low c-Fos expression and

neuronal activity across most brain regions (24, 25), several hours of SD significantly increases

c-Fos expression, particularly in the cortex, medial preoptic area, posterior hypothalamus, and

the hippocampus (24-27). This brain-wide activation resembles the increased c-Fos expression

pattern during spontaneous wakefulness. Given the anatomical and functional complexity of the

hippocampus, this study aims to precisely map c-Fos expression patterns within hippocampal

subregions after single and repeated SD exposures, and identify activity-driven gene expression

changes within SD-activated neurons.

Our results suggest that neuronal activation, as measured by c-Fos expression, varies across

the hippocampus following both single and repeated SD. Specifically, CA1 subregion showed

the most pronounced increase in activation, while DG exhibited a decrease after both single and

repeated SD. Applying the Target Recombination in Active Populations (fosTRAP) method in

conjunction with the ribosome tagging strategy (RiboTag) (28, 29), we devised an active neuron

labeling system that was amenable to analysis of individual gene expression. Through this c-

Fos-RiboTag approach, we confirmed that a substantial proportion of CA1 pyramidal neurons

were reactivated by repeated bouts of SD. As prior genomic studies identified translation-level

changes in brain regions exhibiting increased c-Fos expression after SD, including the

hippocampus (30-33), we further combining the c-Fos-RiboTag labeling system with Translating

Ribosome Affinity Purification sequencing (TRAPseq) technique and mapped activity-dependent

translatome changes in c-Fos+ CA1 neurons.

METHODS & MATERIALS

Animals

All experimental animals were 3- to 5-month-old male C57BL/6J mice obtained from the Jackson Laboratories (Catalog# 000664, Bar Harbor, ME). Mice were group housed (5 per cage) prior to the start of the experiment in standard ventilated cages with access to food (NIH-31 irradiated modified mouse diet #7913) and water *ad libitum.* Animals were maintained in the animal care facility at the University of Iowa on a 12h light/dark cycle, in a temperature- and humidity-controlled environment (21-22°C and 60-70%, respectively). All experiments were conducted in accordance with the standards established by the US National Institutes of Health Guidelines for Animal Care and use was approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Iowa.

Sleep deprivation

Total SD was performed acutely, for 5h beginning at Zeitgeber time (ZT) 0 using the gentle handling method, which involved lightly tapping and shaking the cage to keep the animal awake (12). Mice were single housed in cages with corncob bedding and a handful of soft bedding for nest building for 7 days prior to SD or non-sleep deprivation (NSD). Mice had *ad libitum* access to food and water in these housing conditions and during the sleep deprivation experiments. Mice were habituated to the experimenter and gentle handling stimuli, which was done by the experimenter holding each mouse in the palm for 2min and cage tapping for 2min for 4 consecutive days prior to the 1st sleep deprivation bout. After 5 hours, mice were returned to the original standard housing room where NSD mice were kept undisturbed throughout the experimental time window (ZT0-ZT5). The second SD or NSD bout was performed a week later using the same gentle handling method and timeline. After the second bout of SD/NSD, mice went through perfusion fixation for immunohistochemistry validation of c-Fos/RiboTag expression or cervical dislocation to collect tissue for RNA extraction from the hippocampal CA1.

For active neuron tagging experiments: injection habituation occurred for 2 consecutive days prior to the first sleep deprivation experiment in which mice received an i.p. injection of saline

after the 2min of handling and before being placed back in the cage for 2min of light cage

tapping. All mice received an i.p. injection of 15mg/kg or 50mg/kg 4-hydroxytamoxifen (4-OHT)

at ZT2 on the day of first SD/NSD.

Adeno-associated virus (AAV) constructs and stereotactic surgeries

Surgeries were performed under isoflurane anesthesia (5% induction, 2% maintenance)

following administration of Meloxicam (5mg/kg subcutaneous, s.c.). Animal health was

monitored for 5 days following surgery along with administration of Meloxicam (5mg/kg, s.c.) for

the first 2 days post-surgery. To label the SD activated hippocampal neurons, 10-12 week old

mice were injected intrahippocampally with a cocktail of AAV8-c-Fos-ERT2-Cre-ERT2-PEST-

WPRE (titer- 1.57E+13 GC/mL, generated and packaged by Stanford University Gene Vector,

catalog# GVVC-AAV-139) and AAV9-EF1a.DIO.HA-mRpl22.IRES2.eYFP.WPRE.hGH (titer-

1.14E+13 GC/mL, generated and packaged by the University of Pennsylvania Viral Vector Core).

For labeling cells activated by SD throughout the hippocampal subregions, the AAV cocktail was

injected into the dorsal hippocampus bilaterally (1000nL at a rate of 200nL/min) using the

following coordinates relative to bregma: anteroposterior (AP) -1.9mm, mediolateral (ML)

±1.5mm, dorsoventral (DV) -1.5mm from the surface of the brain. For labeling cells activated by

SD specifically in area CA1, the AAV cocktail was injected into dorsal CA1 bilaterally (300nL at a

rate of 50nL/min), the following coordinates were used upon validation: AP -1.8mm, ML

±1.45mm, DV -1.65 from the surface of the skull. Experiments were performed after 3 to 4

weeks to allow recovery from surgery and for spread of the viral particles in the targeted brain

regions.

Drug preparation

4-OHT solution was prepared fresh on the day of experiment as previously described (34). 4-

OHT (Catalog #H6278, Sigma-Aldrich, MO, USA) was dissolved in 100% ethanol at 37°C for

15min with constant rotation to prepare a 20mg/mL stock solution. Stock solution was then

diluted to 10mg/mL in 100% corn oil followed by another 15min incubation at 37°C with constant

rotation. The solution was vacuum centrifuged for 15min to evaporate the ethanol, yielding an

injectable solution of 10mg/mL 4-OHT in corn oil, and was administered intraperitoneally (i.p.) at

a dose of 15mg/kg or 50mg/kg body weight.

Immunohistochemistry (IHC)

Perfusion fixed brains (4% v/v paraformaldehyde in 1X phosphate buffer saline (PBS), Catalog

#15710, Electron Microscopy Sciences, PA, USA) were equilibrated in sucrose solution

(30%w/v in 1X PBS) for 48h. The brains were then sliced into 30µm sections using a Leica

cryostat (Leica CM3050S, IL, USA) at -20°C and stored in cryoprotectant solution (30% w/v

sucrose, 30% v/v ethylene glycol, and 0.01% Sodium Azide in 1X PBS). Sections containing the

dorsal hippocampus were rinsed three times for 5min each in 1X PBS , followed by a 1h

incubation in blocking solution (3% normal Donkey/Goat serum in 0.4% triton-X100/PBS buffer)

and an overnight incubation at room temperature (RT) with primary antibodies: rabbit anti-c-Fos

(1:3000, #226008, Synaptic Systems, Goettingen, Germany), guinea pig anti-c-Fos (1:3000,

#226308, Synaptic Systems), rabbit anti-HAtag (1:2000, #3724, Cell Signaling, MA, USA),

mouse anti-NeuN (1:1500, #ab104224, Abcam, MA, USA), rabbit anti-Sox9 (1:1500, #ab185966,

Abcam), mouse anti-GAD67 (1:1000, #MAB5406, Sigma-Aldrich). Following overnight

incubation with primary antibodies, slices were rinsed three times for 5 min in 1X PBS and

incubated with secondary antibodies: Alexa Fluor 488 goat anti-rabbit (1:1000, #A11008,

Invitrogen, CA, USA), Alexa Fluor 647 donkey anti-rabbit (1:1000, #A32795, Invitrogen), Alexa

Fluor 555 goat anti-guinea pig (1:1000, #150186, Abcam), Alexa Fluor 647 goat anti-guinea pig

(1:1000, #A21450, Invitrogen), and Alexa Fluor 488 donkey anti-mouse (1:1000, #A21202,

Invitrogen) for 2h at RT followed by three rinses in 1X PBS for 5min each. Slices were mounted

onto Superfrost Plus (Fisherbrand) slides with Prolonged Diamond Antifade mountant with DAPI

(P36962, Thermo Fisher Scientific, MA, USA) to stain cell nuclei.

Confocal Microscopy and Image Analysis

Following IHC, images of the dorsal hippocampus were acquired using a Leica SPE Confocal

microscope equipped with lasers at 405nm, 488nm, 561nm, and 635nm. All images (8-bit) were

obtained with identical settings for laser power, detector gain, and pinhole diameter (1.0AU)

using a 20X oil immersion objective at 1024×1024-pixel resolution and 1.5X optical zoom.

The Adult Mouse Allen Brain Reference Atlas was used as a reference for the hippocampus and

hippocampal subregions. A total of 2-3 sections (2 hippocampi per section) were imaged per

mouse and averaged get the number of c-Fos puncta per animal. Images were processed using

the ImageJ software. Background fluorescent signals generated from AAV infusion surgery was

subtracted from c-Fos and HAtag channels using the Image Calculator function in ImageJ.

Images were converted to 16-bit for c-Fos+ cell quantification. The c-Fos+ cells within the cell

body layers of each subregion were then plotted and counted automatically using the Analyze

Particle function in ImageJ. The HAtag+ cells, DAPI+ nuclei, and double-labeled with

HAtag/NeuN/Sox9 cells were counted manually. For c-Fos and GAD67 overlapping comparison,

GAD67 images were first merged with the DAPI channel and overlapping cells within cell layers

were identified as inhibitory interneurons. Then, interneurons overlapping with c-Fos labeling

were quantified as activated inhibitory interneurons in each subregion. The percentage of

activated excitatory neuron in each subregion was calculated by subtracting the inhibitory

(GAD67+) neuronal fraction from the neuronal (NeuN+) fraction and normalizing to the fraction

of NeuN+ neurons.

Immunoprecipitation and mRNA Extraction

For translatome sequencing, mice were sacrificed at ZT5 by cervical dislocation after the 2nd

SD/NSD. Hippocampi were rapidly dissected into ice-cold 1X PBS followed by microdissection

of CA1 on ice under the Leica S4E stereo microscope as previously described (35). CA1

tissues were collected in sterile, DNase and RNase-free tubes chilled on dry ice and

immediately stored at -80ºC. Ribosome-associated mRNA was immunoprecipitated as

previously described (29). Both CA1 tissues were binned from 5 mice in each group were

homogenized in 1.2mL homogenization buffer (50mM Tris pH 7.4, 100mM KCL, 12mM MgCl$_2$,

10% Nonidet P-40 (NP40), 1mM DTT, 200U/mL Promega RNasin, 1mg/mL heparin, 100μg/mL

cycloheximide, 1X protease inhibitor (#P8340, Sigma, MA, USA). The homogenates were

centrifuged at 10,000xg for 10min at 4˚C. The supernatant was incubated with 5μL Anti-HA.11

Epitope Tag Antibody (#16B12, Biolegend, CA, USA) for 4h at 4 ˚C with rotation, followed by

overnight incubation with Pierce Protein A/G magnetic beads (#88803, Pierce, MA, USA) at 4 ˚C

with rotation. Supernatant was removed from the beads using a magnetic rack and the beads

were washed 3 times with a high salt buffer (50mM Tris pH 7.4, 300mM KCL, 12mM MgCl$_2$, 10%

NP40, 0.5mM DTT, 100μg/mL cycloheximide). mRNA bound to the beads were eluted at 4˚C in

350μL of RLT buffer from the RNeasy Micro Kit (#74004, Qiagen, Venlo, Netherland)

supplemented with β-mercaptoethanol mRNA extraction at 10μL/mL. DNAse digestion of IP-

eluted mRNA were performed following Qiagen's RNeasy extraction kit instructions. RNA

concentration and quality was assessed using the Qubit^{TM} RNA High Sensitivity Assay Kit

(#Q32852, Thermo Fisher Scientific, MA, USA).

Library Preparation

RNA quality was assessed using an Agilent Bioanalyzer. RNA library preparation from samples

of SD and NSD mice were prepared at the Iowa Institute of Human Genetics (IIHG), Genomics

Division, using Illumina Stranded Total RNA Prep Ligation with Ribo-Zero Plus v01 sample

preparation kit (Illumina, CA, USA). Pooled library was sequenced on the Element Aviti24

sequencer with 150-bp paired-end chemistry at the IIHG core.

fosTRAP-seq Data Analysis

Data analysis was performed using Partek Flow interface (Partek, Illumina, CA). Following

QA/QC, raw reads were subjected to the base trimming to remove poorly scored 3'-bases and

aligned using STAR aligner version 2.7.8a (36). Aligned reads were quantified by Partek E/M

model and annotated to mouse reference genome mm39 Ensemble release 110. The resulting

mRNA counts were filtered to exclude mRNA with minimum count $\leq$ 5 (1%) and normalized

using median ratio method. Differential gene expression analysis was performed with DESeq2

(37). Differentially Expressed Genes (DEGs) were identified by the standard: fold change >1.2

or <-1.2, false discovery rate (FDR) value <0.05.

Enrichment of upregulated/downregulated DEG-associated pathways from the fosTRAP-seq

data was performed with a combination of Gene Ontology (GO-Biological Process-EBI-UniProt-

GOA-ACAP-ARAP) and Kyoto Encyclopedia of Genes and Genomes (KEGG) database. The

GO-BP analyses were performed with the clusterProfiler package in R (38) and the Cytoscape

(version 3.10.4 https://cytoscape.org/) (39) plug-in ClueGo (version 2.5.10) (40). Only the

common GO-BP pathways between the two methods with an adjusted p value <0.05 and gene

counts $\geq$ 3 were considered as significant. Volcano plot of all DEGs and Cnet plots representing

enriched GO-BP pathways with their associated DEGs were generated using SRplot website

(41) (Fig. 6). KEGG analyses were performed using the Cytoscape plug-in ClueGo, pathways

were considered significantly enriched with an adjusted p value < 0.05 and gene counts $\geq$ 3. To

connect terms in the network, ClueGO utilizes kappa statistics in which here was set as $\geq$ 0.4

and generated the map figure of enriched functions (additional file1: Fig. S3, 4). For comparison

between our fosTRAP-seq and previous excitatory TRAPseq data published by Lisa Lyons et al.,

(31) (Fig. 6), we first ran the same DEG identification protocol as described above on excitatory

TRAPseq raw data and compared common DEGs between the two datasets. The quadrant

diagram illustrating common DEGs (Fig. 6) were generated using GraphPad Prism v10. GO-BP

analysis was performed to identify enrichment biological processes associated with common

DEGs using clusterProfiler package in R (38) and the pathways with their associated genes

were presented using a Sankey diagram generated using SRplot (41) as well. Biological

functions associates with DEGs specific for each database were performed using KEGG

analysis (additional file1: Fig. S4) with Cytoscape as described.

Statistical analysis

All statistical analyses were performed using GraphPad Prism v10 with $\alpha = 0.05$ for analyses.

Imaging data comparing c-Fos and RiboTag expression and neuronal reactivation were

analyzed using unpaired two-tailed t-tests to compare the NSD and SD groups (Fig. 1, 3C, 4, 5)

and the distal and proximal CA1 sections (additional file1: Fig. S2). Two-way analysis of

variance (ANOVA) comparisons were used to determine the interaction between c-Fos

activation and number of sleep deprivation epochs (Fig. 3D, E). Data is presented as Mean ±

SEM and *n* represents the number of animals in all experiments, IHC data is presented as an

average of 3-4 sections per animal per condition. In figures, ns refers to non-significant, * refers

to a *p* value < 0.05, ** refers to $p < 0.01$, *** refers to $p<0.001$, and **** refers to $p<0.0001$.

RESULTS

Sleep deprivation causes non-uniform activation of hippocampal subregions as

measured by c-Fos expression

Induction of IEGs, such as *Fos, Arc, and Egr1*, mark recent neuronal activation, and previous

studies have investigated the impact of sleep and sleep loss on neuronal activity across rodent

brains by measuring mRNA or protein levels of IEGs (24-27, 30, 32, 42). To investigate the

impact of acute SD on neuronal activation within hippocampal subregions, we sleep deprived

mice for 5h by the gentle handling method. This method was chosen to avoid confounds with

methods that may cause novelty-induced hippocampal activation. Following our 5h SD, we

mapped c-Fos protein expression, a well characterized marker of neuronal activity, within the

hippocampus as an indicator of neuronal activation. Quantification of c-Fos protein abundance

by immunohistochemistry in SD and NSD mice found increased c-Fos in SD samples

suggesting an overall increase in hippocampal activity (Fig. 1A and B_1) after 5h of SD. As the

hippocampal subregions have distinct functions, we quantified c-Fos expression within each

hippocampal subregion finding that sleep deprivation for 5h induced c-Fos expression in

pyramidal cell layers in CA1 and CA3 subfields. However, we observed decreased c-Fos

expression in the superior (DGsup) and inferior blades (DGinf) of the DG (Fig. $1B_2$).

Furthermore, although CA1 and CA3 both showed an increase in c-Fos expression after 5h of

SD, the induction of c-Fos expression was more prominent in CA1 (8-fold induction compared to

NSD) compared to CA3 (1.6-fold induction compared to NSD).

We then asked whether a specific cell type within each subregion is selectively activated by

sleep deprivation. To answer that question, we performed co-staining with c-Fos and cell-type

specific markers (NeuN for neurons and Sox9 for astrocytes). Our results suggest that neurons

(NeuN+) represent the majority of activated cells in all the subregions while a minor fraction of

those c-Fos+ cells in each subregion are astrocytes (Sox9+) (Fig. 2A-B). In addition, depending

on the subregion, a variable fraction of activated cells (c-Fos+) was classified as other cell types

(e.g. microglia, oligodendrocytes) because they were negative for NeuN and Sox9. To further

identify whether those c-Fos+ neurons are excitatory or inhibitory, we performed double labeling

against c-Fos and GAD67, results indicated that sleep deprivation significantly activates

excitatory pyramidal neurons (about 95% of all activated neurons) rather than inhibitory

interneurons in all hippocampal subregions (Fig. 2C-D).

Repeated SD has similar effects on c-Fos expression

In humans, SD is not usually a singular event, but frequently occurs on multiple nights or a weekly basis. However, how hippocampal neuronal activity is affected by repeated sleep loss remains poorly understood. To address this gap and to determine whether the prominent c-Fos induction in CA1 was due to stochastic neuronal activity or an inherent bias that makes a subset of neurons susceptible to sleep loss, we subjected animals to a repeated SD paradigm (Fig. 3A). In this paradigm animals were subject to SD twice with a 1 week-long recovery between the first and the second SD. This allowed us to determine how repeated SD alters c-Fos expression across hippocampal subregions. A second SD period revealed a similar pattern of c-Fos expression in the hippocampus as a single SD with induction in CA1 and CA3, and a decreasing trend in DG inferior blade (Fig. 3 B, C). The repeated protocol itself does not have a significant effect on c-Fos abundance within subregions. We observed similar c-Fos expression levels between single and repeated NSD and SD groups (1xNSD vs 2xNSD, and 1xSD vs 2xSD) within subregions, although the DG region showed decreased c-Fos after repeated NSD (Fig. 3D, E & additional file2: Table S1). These results indicate consistent levels of neuronal activation after each individual SD session. More importantly, the c-Fos induction patterns evident with a single SD were conserved after the second SD: high c-Fos induction in CA1 (7-fold) and a smaller induction in CA3 (3.5-fold).

Together, the results suggest that SD consistently induces region-specific changes in c-Fos expression and neuronal activation within the hippocampus, with the CA1 and CA3 regions showing increased c-Fos and the DG consistently showing decreased c-Fos expression.

c-Fos driven RiboTag labels hippocampal neurons activated by sleep deprivation

Although there is a significant induction of c-Fos expression in CA1 pyramidal neurons after both single and repeated SD compared to NSD, the number of c-Fos+ neurons represent a

small portion of the total number of CA1 pyramidal neurons. While the results from the repeated

SD paradigm suggest that CA1 neurons are activated by SD, it remains unclear if the neurons

within CA1 that show c-Fos induction are randomly recruited by individual SD sessions or

whether neurons activated by an initial SD period are then more likely to be activated by

additional periods of SD. To further characterize the neurons activated by repeated bouts of SD,

we took advantage of the fosTRAP system, which can label neurons activated by defined stimuli

(28).

We employed an AAV-based neuron labeling system in which animals were injected

intrahippocampally with a cocktail of a c-Fos driven Tamoxifen-dependent Cre vector (AAV8-

Fos-ERT2-Cre-ERT2) and a Cre-dependent RiboTag vector (AAV9-EF1a-DIO-HA-mRpl22-

IRES- eYFP). Administration of 4-OHT facilitates expression of the tamoxifen-inducible c-Fos-

driven Cre recombinase (Cre^{ERT2}) upon neuronal activation, which subsequently drives

expression of mRpl22-HA (RiboTag) (28, 29) (Fig. 4A) to label CA1 pyramidal neurons. Using

this strategy, we first tested CA1 neuronal labeling efficacy, indicated by RiboTag (HAtag)

expression, using different labeling durations, 4-OHT doses, and AAV infusion methods

(additional file1: Fig. S1 and Fig. 4).

Unlike c-Fos protein itself, which is expressed rapidly following neuronal activation, expression

of the c-Fos activation-promoted RiboTag depends on Cre^{ERT2} mediated recombination of loxP

sites and consequent expression of mRpl22-HA (RiboTag). We first assessed the amount of

time needed to get robust expression of HA following 4-OHT delivery and SD or NSD. In a 5h

labeling test (additional file1: Fig. S1A-C), where 4-OHT was administered at ZT0 and tissue

was collected after SD or NSD (ZT0-5), we observed only weak expression of the HAtag, and

there were no differences between the SD and NSD groups, confirming that the RiboTag marker

takes longer to develop. To provide a longer expression window, we waited one week based on

previous results showing that repeated sleep deprivation leads to c-Fos activation in CA1 and

CA3 that are comparable to a single SD (Fig. 3). Consequently, this approach requires the use

of c-Fos-driven RiboTag labeling for neurons activated by an initial period of SD and subsequent

c-Fos immunohistochemistry to identify neurons activated by a second period of sleep

deprivation. We divided mice into repeated sleep deprivation (SD-SD) or repeated sleep (NSD-

NSD) groups. All mice received an injection of 4-OHT at ZT2 of the 1st SD/NSD (34) to drive

RiboTag expression in neurons activated during the 5h of SD/NSD. We also tested two

concentrations (15mg/kg and 50mg/kg) of tamoxifen and found that a higher dose of 4-OHT

(50mg/kg) was necessary to reflect the proportion of cells activated by SD (additional file1: Fig.

S1D-F).

To further optimize RiboTag expression, we targeted the AAV cocktail infusion to hippocampal

area CA1 bilaterally. Specific CA1 expression revealed a 3-fold induction of RiboTag expression

after SD compared to the NSD group (Fig. 4). The expression level of RiboTag using this CA1

targeting strategy was higher than the normalized number of cells showing c-Fos expression

that we observed in both NSD and SD for the single or repeated SD groups (Fig. 1 and 3),

particularly for the NSD groups. This could be due to the short half-life (~2h) of c-Fos (43), and

consequently the c-Fos protein that was induced in the early stages of the light cycle may not be

reliably detectable at ZT5 using IHC in both the NSD and SD conditions, whereas the RiboTag

will persist upon neuronal activation. Another factor that is likely to increase RiboTag expression

in the NSD group is the unavoidable wakefulness and subsequent c-Fos expression induced by

i.p. injection of 4-OHT. Together, these optimization strategies established a method for activity-

driven RiboTagging that successfully labels CA1 pyramidal neurons that are activated by SD.

CA1 pyramidal neurons are reactivated by repeated SD

To answer the previous question that if subsets of neurons reactivate during repeated SD in the

CA1 or whether hippocampal CA1 pyramidal neurons are randomly activated by repeated sleep

loss, we performed RiboTagging combined with c-Fos immunolabeling. Immunofluorescence

was used to identify the cells positive for RiboTag (the neurons that were activated during first

SD or NSD) and c-Fos (the neurons that were activated during the second SD or NSD). Cells

that are positive for both c-Fos and the HA tag are neurons that undergo reactivation during the

second SD period (Fig. 5A). Our analysis showed that about 30% of CA1 neurons activated by

the first SD (HAtag positive) were reactivated by the second SD (c-Fos positive; Fig. 5B).

Furthermore, of the CA1 pyramidal neurons activated by the second SD (c-Fos+), 40% were

previously activated by the first SD (Fig. 5C). We examined HAtag and c-Fos immunolabeling

across CA1 and found higher HAtag and c-Fos expression in proximal CA1, where pyramidal

neurons are mainly driven by medial entorhinal cortex projections and involved in spatial

information processing (44, 45), with similar reactivation rate between proximal and distal CA1

(additional file1: Fig. S2), which aligns with impaired spatial memory.

Importantly, overlap in c-Fos and HAtag was minimal in the NSD-NSD group: about 5% of

RiboTagged cells during 1^{st} NSD were activated by the 2^{nd} NSD (Fig. 5B), and double-labeled

cells make up approximately 15% of the limited population of c-Fos+ cells after the second NSD

(Fig. 5C). This overlap is likely due to background RiboTag expression (additional file1: Fig.

S1G). This neuronal reactivation confirms that repeated SD stimulates a similar population of

pyramidal neurons in hippocampal CA1 rather than a random activation, which suggests that a

subset of neurons is more likely to be activated during single and repeated SD.

Activity-dependent translatome changes in SD-sensitive CA1 neurons

In addition to tagging activated neurons, the RiboTag approach we applied also allows analyzing

gene expression and mRNA translation in specific cell types (29, 31), in our case the c-Fos+

neurons, by performing Translating Ribosome Affinity Purification with RNA sequencing

(fosTRAP-seq).   We tagged SD-sensitive CA1 neurons by subjecting all mice to sleep

deprivation on day 0, followed by division into NSD and SD groups on day 7 to induce changes

in activity and gene expression (Fig. 6A). Immediately after that SD/NSD session, actively

translating mRNA from CA1 RiboTag-labeled neurons were pooled using immunoprecipitation

with antibodies for the HAtag. Following TRAP RNA extraction (5 animals pooled per sample),

we used an unbiased RNA sequencing approach, aligned raw reads using STAR aligner (36),

and identified differentially expressed genes (DEGs) by the criterion of: FDR < 0.05, and fold

change >1.2 or <-1.2 to avoid false positives. From our fosTRAP-seq data, we identified 631

DEGs being upregulated while 221 DEGs being downregulated in RiboTag labeled CA1

neurons (Fig. 6B, additional file2: Table S2). As expected, the IEGs (such as *Fosb, Arc, Egr1,*

and Homer1) that are bound for translation are upregulated by sleep deprivation indicating

reactivation of certain SD-sensitive CA1 neurons during the repeated sleep loss. We also

observed decreased ribosome associated transcripts of *Cirbp, Rbm3*, *Srsf5* which have been

reported by previous bulk transcription analysis (33, 46).

We then separately analyzed the upregulated and downregulated transcripts for the enriched

biological processes and functions that were affected by SD against the GO: biological process

(GO-BP) and KEGG databases. The analyses on the upregulated DEGs revealed strong

enrichment in biological processes of protein dephosphorylation (e.g. *Dusp1/4/5/19/26,*

Ppm1a/1f, Ptpn1/4/11), negative regulation of kinase activity (e.g. *Adarb1, Dnajc3, Pkia,*

Dusp1/19), response to insulin (e.g. *Akt2, Foxo4, Gsk3b, Irs1/2, Capn10, Egr1, Tsc1, Vgf,*

Ptpn1/11), dendrite development (e.g. *Arc, Cobl, Mack6, Mecp2, Ngfr, Ppp1r9b*), regulation of

neurotransmitter secretion (e.g. *Adra2, Cplx2, Kcnc4, Lrrk2, Rim3/4*), cognition (e.g. *Arc, Egr1,*

Ccnd2, Nr4a2, Uba6) having higher enrichment scores among all altered pathways (Fig. 6C,

additional file2: Table S3-1), and the biological functions of longevity regulation and insulin

signaling pathways (additional file1, 2: Fig. S3A, Table S4-1).   The downregulated DEGs are

mainly associated with regulation of mRNA processing and splicing processes (e.g. *Cirbp,*

Rbm3/x, Srsf5, Luc7l2, Cdc40, and Sfswap), as well as negative regulation of angiogenesis

(*Adgrb3, Flt1, Krit1, Ptn, Tek, and Tie1*) (Fig. 6D, additional file2: Table S3-2), and the biological

functions of ECM-receptor interaction and spliceosome (additional file1,2: Fig. S3B, Table S4-2).

We also compared these data to our previous work performing TRAPseq (31) on bulk

hippocampal excitatory neurons from the hippocampus after 5h of sleep deprivation. There are

62 DEGs overlapped between our fosTRAP-seq and the excitatory TRAPseq and all DEGs

share the same direction of alteration (Fig. 6E-F, additional file2: Table S5). Among the common

DEGs, IEGs (including *Fosb, Arc, Homer1, Nr4a1*), *Upf* 2 (regulator of nonsense mediated

mRNA decay), and *Spred1* (negative regulation of kinase activity/protein phosphorylation) are

upregulated in both datasets. The *Cirbp* and *Rbm3* genes, involved in RNA splicing and mRNA

processing regulation, are downregulated from the two datasets. GO-BP analysis indicates that

protein phosphorylation, mRNA splicing and processing processes are shared in both datasets

(Fig. 6G, additional file2: Table S6). We also compared distinct DEGs between the two datasets

and KEGG analysis revealed that biological functions of insulin-signaling pathway, longevity

regulation pathway, and focal adhesion being enriched specifically in our fosTRAP-seq, while

melanogenesis and apoptosis are altered only in excitatory TRAPseq (additional file1,2: Fig. S4,

Table S7). The shared and unique DEGs between the two datasets further validate the reliability

and robustness of our fosTRAP-seq method.

Taken together, our fosTRAP-seq identified activity-dependent translatome alteration in SD-

vulnerable CA1 neurons with a higher specificity and resolution revealing post-transcriptomic-

and translational-modification, synapse connection, neuron development, insulin signaling being

vulnerable functions affected by sleep loss.

DISCUSSION

The hippocampus is a structurally distinct and functionally diverse brain region, comprised of

interconnected subregions that contribute uniquely to memory encoding processes (7-9). To

better understand how sleep deprivation affects hippocampal function, we investigated c-Fos

expression patterns across different hippocampal subregions. In this study, we sleep-deprived

animals from ZT0-ZT5 by gentle handling, a widely accepted approach that minimizes stress

(47) while avoiding the confounding effects of spatial or object novelty on neuronal activation in

the hippocampus (47, 48), and examined hippocampal subregional c-Fos expression after

single and repeated SD. Results indicated that c-Fos expression is altered differently among

hippocampal subregions: CA1, particularly proximal CA1, shows the most robust c-Fos

induction; CA3 shows a moderate activity increase; and the DG shows reduced c-Fos

expression. (Fig. 1-3). Using the c-Fos driven RiboTag labeling system, we observed 30% to 40%

overlap ratio between neurons activated during the first and second SD and confirmed that CA1

pyramidal neurons are reactivated by repeated sleep loss (Fig. 4-5). We then performed

fosTRAP-seq and revealed translatome changes in RiboTag labeled CA1 neurons that are

sensitive to sleep deprivation (Fig. 6).

In the present study, we observed distinct patterns of neuronal activation among hippocampal

subregions caused by sleep deprivation. Hippocampal neuronal activation is regulated by

coordinated interactions among glutamatergic, GABAergic, serotonergic, and noradrenergic

systems, and regional differences in receptor expression may account for distinct c-Fos

activation patterns. During sleep deprivation, there are a multitude of changes in

neurotransmitter systems and signaling that differ between areas of the cornu ammonis and

dentate gyrus (49, 50). Of note, locus coeruleus–derived norepinephrine release is elevated

during SD and is known to induce c-Fos expression and enhance CA1 pyramidal neuron

excitability via β-adrenergic signaling (51-53). Consistent with this, our fosTRAP-seq data show

that CA1 neurons are enriched for *Adrb1* relative to *Adrb2* and *Adrb3*, suggesting elevated β_1-

adrenergic receptor expression enhances sensitivity to norepinephrine and promotes c-Fos

activation via cAMP-CREB signaling. This differential pattern of neuronal activation is consistent

with previous spatial transcriptomic studies, which identified unique gene expression changes

and enriched molecular functions specific to hippocampal subregions after SD (49, 54) and

adds to a growing literature emphasizing the heterogeneity of SD effects in various hippocampal

subregions. However, our findings differ from a recent study by Sarma and colleagues (2025),

which reported that 3h, but not 6h, of gentle handling sleep deprivation alters c-Fos expression

across hippocampal subregions (55). In contrast, our 5 h sleep deprivation paradigm produced

a more robust increase in c-Fos, particularly in CA1, which may reflect the higher sensitivity of

immunofluorescence compared with DAB-based immunohistochemistry. Notably, our c-Fos

expression pattern is consistent with prior genomic, translational, and behavioral studies

showing upregulation of c-Fos and other IEGs after 5 h of sleep deprivation (31, 33, 46). This

same 5 h gentle-handling protocol reliably induces hippocampus-dependent memory deficits

and synaptic plasticity alterations (12, 14) without elevating corticosterone or confounding

behavioral stress effects (47, 48). Together, these findings suggest that sleep-loss duration is a

critical determinant of hippocampal c-Fos responses, with 5 h representing a key time point at

which molecular, electrophysiological, and behavioral consequences of sleep deprivation

converge.

Utilizing a viral c-Fos-RiboTag active neuron tagging approach (28, 29), we successfully

captured the robust induction of CA1 neuronal activation and identified a specific population of

CA1 pyramidal neurons that is repeatedly activated by sleep deprivation. Our approach labeled

a substantial number of activated neurons compared to a previous study that reported minimal

labeling of CA1 neurons following sleep deprivation using a TRAP transgenic mouse model (56).

Our results also demonstrate over 30% overlap between c-Fos+ and RiboTag-labeled neurons

across repeated sleep deprivation events (Fig. 5). This represents a significant overlap of

neuronal activation after repeated SD comparable to the hippocampal engram reactivation rate

observed using IEG-dependent tagging methods (e.g. TRAP2 and Tet-tag) during memory

consolidation and retrieval, which typically ranges from 5-20% (49, 57-59). This prominent reactivation in our dataset provides compelling evidence supporting the hypothesis that a specific subset of CA1 pyramidal neurons have distinct molecular or electrophysiological characteristics (60, 61) that confer heightened sensitivity to repeated and chronic sleep loss. Notably, c-Fos expression is widely used as indicator of engram activation and reactivation in hippocampus-dependent memory formation and consolidation. In CA1, higher neuronal place field preference, place field activity, stability of spatial maps, and more precise spatial information decoding is found in the neurons that were c-Fos positive during spatial learning tasks (62). Thus, their strong activation during sleep deprivation –despite the absence of learning– raises a mechanistic question regarding the functional significance of CA1 activation under sleep loss. Consistent with this idea, several hours of sleep deprivation following learning similarly induces CA1 c-Fos expression yet impairs memory consolidation (49, 55), implying that sleep loss may either recruit neuronal populations distinct from memory engrams or aberrantly activate learning-related ensembles while disrupting gene expression, protein synthesis, or sharp-wave ripple–mediated replay (63-65). Future experiments comparing whether learning and sleep deprivation activates a similar population of CA1 pyramidal neurons could be helpful answer the question. Together, these observations highlight a sleep deprivation-sensitive CA1 neuronal subpopulation and provide a framework for dissecting how aberrant neuronal activation and reactivation contribute to hippocampal dysfunction and memory impairment under chronic sleep loss.

Taking advantage of the c-Fos-RiboTag system combined with repeated SD paradigm, we isolated activity-dependent translatome changes in CA1 neurons that are selectively sensitive to sleep loss (c-Fos+). Our fosTRAP-seq analysis identified DEGs that have been reported in previous gene expression studies after acute sleep deprivation (31, 33, 46, 54) supporting the robustness of our repeated SD and neuronal labeling strategy. We reported upregulated

translation of transcripts involved in synaptic vesicle exocytosis (e.g. *Cplx2, Rims4, Slc4a8*),

regulation of synaptic plasticity (e.g., *Arc, Egr1, Adcy1/8*), and dendrite development (e.g. *Arc,*

Cobl, Mack6, Mecp2) all of which are essential for memory consolidation. In contrast, transcripts

associated with mRNA splicing and processing (e.g., *Rbm3/25, Rbmx, Cirbp, Srsf5, Luc7l2,*

Tia1, Snrnp70, *Fus*) were markedly downregulated by sleep deprivation. In the context of

memory impairment by SD, this pattern suggests that although CA1 neurons may initiate

synaptic remodeling during SD, disruptions in RNA processing and translational control could

ultimately impair effective memory consolidation.

Beyond previously reported DEGs identified from excitatory neurons from the entire

hippocampus (31), we further detected (Fig. 6) increased translation of genes involved in insulin

signaling and insulin resistance pathways, including *Akt2, Gsk3b, Irs1/2, Pik3r3, Ppp1r3c,*

Prkab2, Ptpn1/11, Ptprf, and Rps6ka2. Brain insulin signaling regulates multiple cellular

processes including glucose metabolism, cell survival, and synaptic plasticity through PI3K-Akt

signaling and downstream phosphorylation of targets such as glycogen synthase kinase 3 beta

(GSK-3β) and mTOR (66). Disruption of hippocampal insulin signaling such as developing

insulin resistance impairs neuroplasticity and spatial memory in rats (67) and overactivation of

GSK-3β has been linked to inhibition of hippocampal long-term potentiation (68). Consistent

with this framework, our fosTRAP-seq data indicate that insulin response pathways, particularly

those relate to insulin resistance, are substantially altered in SD-sensitive CA1 neurons. These

disruptions may serve as a central mechanism linking SD to interrupted synaptic plasticity,

impaired signal transmission and reduced memory performance. Importantly, GSK-3β also

participates in different aspects and pathways relevant to the onset and development of

Alzheimer's Disease neuropathology via directly promoting tau hyper-phosphorylation, amyloid

production and accumulation, and inflammatory molecules (69). Additionally, we observed

increased translation of *Ttbk1/2* (tau tubulin kinase 1 and 2), kinases that phosphorylate tau at

multiple sites and contribute to tauopathy in neurodegenerative disorders (70). Given that

chronic sleep restriction causes hippocampus-dependent memory deficits (71) and promotes

tau hyperphosphorylation in both human and rodents (18, 21), our findings suggest a

mechanistic continuum whereby repeated acute SD initiates insulin signaling dysfunction and

upregulation of tau-related kinases (GSK-3 β , TTBK1/2), which, when sustained, may

accelerate tau pathology and neurodegeneration in CA1. Collectively, these results highlight

insulin signaling dysfunction as a key molecular feature of SD-induced hippocampal impairment

and propose a trajectory linking acute sleep loss to chronic sleep restriction-associated

tauopathy and increased vulnerability to neurodegenerative disease.

For future studies, a direct comparison of gene expression profiles between CA1 pyramidal

neurons that are sleep loss sensitive (c-Fos+) and insensitive (c-Fos-) during SD will be

important for delineating molecular features that confer selective vulnerability to sleep loss. In

parallel, the fosTRAP system will enable the manipulation of these SD-sensitive neurons for

causal tests of their functional significance. Specifically, determining whether selective activation

of this neuronal population is sufficient to induce memory impairments, or whether their

inhibition can mitigate SD-induced deficits, will provide critical insight into the circuit

mechanisms underlying sleep loss-associated cognitive dysfunction.

In conclusion, this study reveals the subregional impact of sleep deprivation on hippocampal

neuronal activation, which highlights the importance of spatially segregated analysis to

understand the response to sleep loss. The reactivation of CA1 pyramidal neurons after

repeated SD supports the premise that certain excitatory neuron populations are particularly

susceptible to repeated sleep loss. Using the c-Fos-RiboTag system, we identified activity-

driven gene expression alteration at a high resolution in SD-sensitive CA1 neurons and

expanded the understanding of how SD could impair memory consolidation and progress

toward chronic sleep restriction induced damages. Our research provides a detailed insight into

hippocampal neurons responsive to SD and establishes a foundation for future molecular and

functional investigations of neuron subsets activated by sleep loss. Identifying these subsets will

help inform strategies to mitigate cognitive deficits associated with SD.

List of Abbreviation

SD: sleep deprivation

NSD: non-sleep deprivation

CSR: chronic sleep restriction

CA1: Cornus Ammonis 1

CA3: Cornus Ammonis 3

DG: Dentate Gyrus

IEG: immediate early gene

ZT: zeitgeber time

RT: room temperature

h: hour

min: minute

AAV: adeno-associated virus

4-OHT: 4-hydroxytamoxifen

i.p.: intraperitoneally

IHC: immunohistochemistry

PBS: phosphate buffer saline

NP40: nonidet P-40

TRAP: target recombination in active populations

RiboTag: ribosome tagging strategy

TRAPseq: translating ribosome affinity purification sequencing

ANOVA: analysis of variance

DEG: differentially expressed gene

FDR: false discovery rate

GO-BP: gene ontology – biological process

KEGG: Kyoto encyclopedia of genes and genomes

SEM: standard error of the mean

DECLARATIONS

Ethics approval and consent to participate

All experiments were conducted in accordance with the standards established by the US

National Institutes of Health Guidelines for Animal Care and use was approved by the

Institutional Animal Care and Use Committee (IACUC) at the University of Iowa.

Consent for publication

Not Applicable

Availability of data and materials

The datasets generated and/or analyzed during the current study are available in the NCBI's

Gene Expression Omnibus repository, GEO Series accession GSE316227,

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE316227, and DEG accession

GSE156925, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE156925.

Any additional information underlying this article will be shared on reasonable request to the

corresponding author.

Competing Interests

The authors declare that they have no competing interest.

Funding

This work was supported by grant funding from the National Institutes on Aging 5R01AG062398

to T.A. and L.C.L. T. A. is supported by a Roy J. Carver Directorship.

Authors' contribution

Y.W. and T.A. designed the study with input from E.N.W., J.R., and L.C.L.. Y.W. and E.N.W.

performed sleep deprivation and tissue collection. Y.W. performed protein expression

experiments and data analysis. Y.W. performed mRNA extraction for fosTRAP-seq and the

sequencing data analysis by J.K., C.R., and Y.W.. Y.W. wrote the manuscript with input from all

other authors.

Acknowledgements

The sequencing was performed by Genomics Division in the Iowa Institute of Human Genetics

which is supported, in part, by the University of Iowa Carver College of Medicine. We thank Dr.

Yong-Seok Lee for his guidance on the CA1 microdissection technique. We also thank Dr. Satya

Tadinada and Pravda Quinones-Labernik for their valuable feedback on this manuscript.

Figure Legends

Figure 1. cFos expression is differently altered within hippocampal subregions after

acute sleep deprivation. A) Immunofluorescent images of c-Fos expression in NSD and SD

hippocampus. **B**) Comparison of area normalized cFos expression between NSD (n=5) and SD

(n=6) groups from the entire hippocampus (1) and within different subregions (2): CA1, CA3,

and superior and inferior blades of Dentate Gyrus (DGsup and DGinf). Data are shown as mean

$\pm$ standard error of the mean (SEM). Unpaired two-tailed t test for B_1: NSD=180.64±11.53,

SD=408.39±14.27, t(9)=12.05, p<0.0001. Multiple unpaired t tests for B_2: CA1:

NSD=161.50±17.15, SD=1072.02±26.98, t(9)=27.10, p<0.00001; CA3: NSD=142.53±6.97,

SD=218.99±14.11, t(9)=4.54, p=0.0014; DGsup: NSD=176.18±11.26, SD=131.76±6.58,

t(9)=3.56, p=0.0061; DGinf: NSD=289.11±51.80, SD=99.07±6.38, t(9)=4.02, p=0.0030.

Figure 2. **Excitatory neurons are selectively activated by SD**. **A**) Representative images of

c-Fos double labeling with NeuN and Sox9 after SD. **B**) Percentage of c-Fos+NeuN+ and c-

Fos+Sox9+ double labeled cells to the total number of c-Fos+ cells within each subregion in SD

(n=3) hippocampus. **C**) Representative images of c-Fos double labeling with GAD67 (inhibitory

neuronal marker) after SD. **D**) Percentage of excitatory and inhibitory neurons to the neuronal

population within cell layers of each subregion in SD (n=3) hippocampus.

Figure 3. **Repeated sleep deprivation induces similar pattern of cFos expression in the**

hippocampus. A) Schematic of repeated sleep deprivation. **B**) Immunofluorescent images of c-

Fos expression after repeated NSD (NSD-NSD) and repeated SD (SD-SD) in the hippocampus.

C) Comparison of area normalized c-Fos expression between NSD-NSD (n=6) and SD-SD (n=6)

whole hippocampus. Unpaired two-tailed t-test: NSD=113.13±15.84, SD=336.38±36.42,

t(10)=5.62, p=0.0002. **D, E**) Comparison of area normalized cFos expression between single

(1x NSD/SD) and repeated sleep deprived (2x NSD/SD) hippocampus (D) and hippocampal

subregions (E). Data are shown as mean ± SEM. D: Two-way ANOVA shows a significant effect

of sleep deprivation ($F_{(1,19)}$=98.43, *p*<0.0001) and a significant effect of repeat manipulation

($F_{(1,19)}$=9.42, *p*=0.0063), but no significant effect of interaction ($F_{(1,19)}$=0.01, *p*=0.9224). E:

Two-way ANOVA for each subregion: E1: effect of SD ($F_{(1,19)}$=158.5, p<0.0001), ns for effect

of repeat manipulation and interaction; E2: effect of SD ($F_{(1,19)}$=29.29, p<0.0001), ns for effect

of repeat manipulation and effect of interaction; E3: ns for effect of SD, effect of repeat

manipulation ($F_{(1,19)}$=14.23, p=0.0013), effect of interaction ($F_{(1,19)}$=5.57, p=0.0291); E4:

effect of SD ($F_{(1,19)}$=19.45, p=0.0003), ns for effect of repeat manipulation, effect of interaction

($F_{(1,19)}$=4.81, p=0.041). Bonferroni post hoc tests were performed for panel D & E to compare

single vs repeated exposure under SD & NSD conditions, as well as repeated NSD vs SD c-Fos

expression. Detailed 2-way ANOVA analyses results see additional file2: Table S1.

Figure 4. cFos-RiboTag strategy to label activated cells in area CA1. A) Diagram of the

activity (cFos)-driven ribosomal tagging (RiboTag) strategy**. B**) Diagram of CA1 neuron labeling

using cFos-RiboTag strategy with repeat sleep deprivation. **C**) Representative images of neuron

labeling shown as RiboTag (HAtag) expression in the NSD and SD hippocampus. **D**) Neuronal

labeling efficacy in hippocampal CA1: area normalized HAtag+ cells in NSD (n=5) and SD (n=4)

groups.   Unpaired   two-tailed   t-test   performed,   NSD=476.91±02.01,   SD=1408.06±186.79,

t(7)=4.79, p=0.0020. Data shown as mean ± SEM.

Figure 5. CA1 pyramidal neurons are reactivated by repeated SD. A) Immunofluorescence

images of cFos and HAtag double labeling in area CA1 in NSD-NSD and SD-SD groups. **B, C**)

Neuronal reactivation: percentage (%) of double labeled cells / HAtag+ cells (B) and double

labeled cells / cFos+ cells (C) overlap in NSD-NSD (n=5) and SD-SD (n=4) CA1. Unpaired two-

tailed t-test. Data shown as mean ± SEM. B: NSD-NSD=8.82±2.71, SD-SD=30.97±5.56,

t(7)=5.44, p=0.0064. C: NSD-NSD=36.02±7.95, SD-SD=38.96±6.03, t(7)=3.26, p=0.7866.

Figure 6. fosTRAP-seq reveals activity-dependent gene expression alteration in SD-

sensitive hippocampal CA1 neurons after sleep deprivation. A) Diagram of translating

mRNA extraction from RiboTag labeled CA1 neurons. All mice were sleep deprived on day 0

then randomly divided into NSD and SD groups on day 7. (n=10 mice/group, CA1 from 5 mice

were pooled together for further RNA extraction.) **B**) Volcano plot of all DEGs (FDR<0.05,

|FC|>1.2) identified in response to sleep deprivation in hippocampal CA1 SD-sensitive neurons.

C, D) Cnet plot showing GO biological process (BP) analyzed pathway enrichment (adjusted

Pvalue<0.05, gene≥3) associated with DEGs significantly upregulated (C) or downregulated (D)

after sleep deprivation in SD-sensitive CA1 neurons. Full details of C, D see additional file2:

Table S3. **E**) Venn diagram representing the number of DEGs overlapped between our

fosTRAP-seq results and our previously published (Lyons et al., 2020) excitatory TRAPseq after

SD identified 62 common genes, 790 only in fosTRAP-seq and 286 only in excitatory TRAPseq.

F) Quadrant diagram showing comparison of common DEGs between fosTRAP-seq and

excitatory TRAPseq. **G**) Sankey diagram (left) shows association of overlapped DEGs to the

identified GO biological processes (p<0.05), along with pathway enrichment values as a Dot plot

(right). Full details of G see additional file2: Table S6.

ADDITIONAL FILES

Additional file1.pdf:

Figure S1. Supplement data for RiboTag expression following single and repeated SD. **A**)

Schematic of the c-Fos promoted RiboTag expression following 5h of SD. **B**) Representative

image of RiboTag (HAtag) expression in NSD/SD hippocampus. **C**) Comparison of area

normalized HAtag+ expression between NSD (n=3) and SD (n=4) groups within area CA1.

Unpaired t-test, NSD=48.28±6.60, SD=77.92±19.24, t(6)=0.7073, p>0.05. **D**) Timeline of the

CA1 neuron labeling using the cFos-RiboTag strategy (whole hippocampus infusion) with

repeated sleep deprivation. **E**) Representative images of neuron labeling shown as RiboTag

(HAtag) expression in the NSD and SD hippocampus injected with 15mg/kg or 50mg/kg 4-OHT.

F) Comparison of neuronal labeling efficacy in hippocampal CA1: area normalized HAtag+ cells

in NSD and SD groups injected with 15mg/kg and 50mg/kg of 4-OHT, n= 4-7 mice in each group.

Two-way ANOVA test performed. Data shown as mean ± SEM. D: No significant effect of

repeated sleep loss ($F_{(1,18)}$=3.815, p=0.0665) but a significant effect of 4-OHT dose

($F_{(1,18)}$=13.21, p=0.0019) and a significant interaction of repeat manipulation and 4-OHT dose

($F_{(1,18)}$=4.049, p=0.0594). Bonferroni post hoc: 15mg/kg: NSD=1145.54±168.88,

SD=1209.69±135.90, t(18)=0.0384, p>0.9999; 50mg/kg: NSD=1420.40±139.43,

SD=2092.38±328.09, t(18)=3.106, p=0.0366. 15mg/kg vs 50mg/kg: p(NSD)>0.9999,

p(SD)=0.0044. **G**) Representative image of background RiboTag expression (~200/mm2) in

negative control mice injected with vehicle (corn oil) rather than 4-OHT.

Figure S2. Proximal CA1 shows greater cFos expression than distal CA1 after SD. **A**)

Representative image of DAPI, cFos+ and HAtag+ cells in distal vs proximal CA1 in SD-SD

group. **B**) Comparison of area normalized DAPI nuclei (1), cFos+ (2), HAtag+ (3), and double

labeled (4) cells between distal vs proximal CA1 in SD-SD mice (n=6). Unpaired t-tests. Data

shown as mean ± SEM. B1) dCA1=7827.39±190.20, pCA1=7732.98±227.45, t(10)=0.32,

p=0.7576. B2) dCA1= 776.04±154.53, pCA1=1243.15±140.26, t(10)=2.24, *p*=0.0491. B3)

dCA1=1287.53±65.68, pCA1=1993.05±142.29, t(10)=4.50, *p*=0.0011. B4) dCA1=305.80±78.63,

pCA1=452.91±41.19, t(10)=1.66, *p*=0.1285. **C**) Comparison of neuronal reactivation rate:

percentage (%) of double labeled cells / HAtag+ cells (1) and double labeled cells / cFos+ cells

(2) between distal vs proximal CA1 in SD-SD mice. Unpaired t-tests, data shown as mean ±

SEM. C1 (%): dCA1=21.97±3.71, pCA1=23.99±1.46, t(10)=0.51, p=0.6230. C2 (%):

dCA1=38.43±5.24, pCA1=39.62±3.73, t(10)=0.18, *p*=0.8572.

Figure S3. fosTRAP-seq biological function enrichment analysis. A, B) Map of KEGG

pathways associated with upregulated DEGs (A) and downregulated DEGs (B) after sleep

deprivation in SD-sensitive hippocampal CA1 neurons (adjusted p<0.05). Full details of KEGG

pathways see additional file2: Table S4.

Figure S4. Comparison between fosTRAP-seq and excitatory TRAPseq. A) Venn diagram

representing DEGs specific to fosTRAP-seq analysis. **B**) KEGG pathways (adjusted p<0.05)

associated with fosTRAP-seq unique DEGs. **C**) Venn diagram representing DEGs specific to

excitatory TRAP-seq analysis. **D**) KEGG pathways (adjusted p<0.05) associated with excitatory

TRAPseq unique DEGs. Full details see additional file2: Table S7

Additional file2.xlsx:

Table S1. Supplement statistical results of 2-way ANOVA analyses of c-Fos expression after

single and repeated SD. **Table S2.** List of DEGs revealed by fosTRAP-seq (FDR<0.05,

|FC|>1.2). **Table S3.** Enriched GO-BP processes associated with fosTRAPseq DEGs (common

pathways enriched in both clusterProfiler and ClueGO analysis). **Table S4.** KEGG biological

functions associated with fosTRAP-seq DEGs. **Table S5**. List of common DEGs between

fosTRAP-seq and excitatory TRAPseq. **Table S6.** Enriched GO-BP processes associated with

common DEGs shared by fosTRAP-seq and excitatory TRAPseq. **Table S7**. Enriched KEGG

biological functions of fosTRAP-seq and excitatory TRAPseq unique DEGs.

REFERENCES

1.	Zielinski MR, McKenna JT, McCarley RW. Functions and Mechanisms of Sleep. AIMS Neurosci. 2016;3(1):67-104.

2.	Adjaye-Gbewonyo D, Ng AE, Black LI. Sleep difficulties in adults: United States, 2020. 2022.

3.	Altevogt BM, Colten HR. Sleep disorders and sleep deprivation: an unmet public health problem. 2006.

4.	Killgore WD. Effects of sleep deprivation on cognition. Prog Brain Res. 2010;185:105-29.

5.	Zemla R, Basu J. Hippocampal function in rodents. Current opinion in neurobiology. 2017;43:187-97.

6.	Grunwald T, Kurthen M. Novelty detection and encoding for declarative memory within the human hippocampus. Clin EEG Neurosci. 2006;37(4):309-14.

7.	Yassa MA, Stark CE. Pattern separation in the hippocampus. Trends Neurosci. 2011;34(10):515-25.

8.	Deuker L, Doeller CF, Fell J, Axmacher N. Human neuroimaging studies on the hippocampal CA3 region - integrating evidence for pattern separation and completion. Front Cell Neurosci. 2014;8:64.

9.	Kesner RP, Rolls ET. A computational theory of hippocampal function, and tests of the theory: new developments. Neurosci Biobehav Rev. 2015;48:92-147.

10.	Vecsey CG, Baillie GS, Jaganath D, Havekes R, Daniels A, Wimmer M, et al. Sleep deprivation impairs cAMP signalling in the hippocampus. Nature. 2009;461(7267):1122-5.

11.	Prince TM, Wimmer M, Choi J, Havekes R, Aton S, Abel T. Sleep deprivation during a specific 3-hour time window post-training impairs hippocampal synaptic plasticity and memory. Neurobiol Learn Mem. 2014;109:122-30.

12.	Walsh EN, Shetty MS, Diba K, Abel T. Chemogenetic Enhancement of cAMP Signaling Renders Hippocampal Synaptic Plasticity Resilient to the Impact of Acute Sleep Deprivation. eNeuro. 2023;10(1).

13.	Tudor JC, Davis EJ, Peixoto L, Wimmer ME, van Tilborg E, Park AJ, et al. Sleep deprivation impairs memory by attenuating mTORC1-dependent protein synthesis. Sci Signal. 2016;9(425):ra41.

14.	Havekes R, Park AJ, Tudor JC, Luczak VG, Hansen RT, Ferri SL, et al. Sleep deprivation causes memory deficits by negatively impacting neuronal connectivity in hippocampal area CA1. Elife. 2016;5.

15.	Raven F, Meerlo P, Van der Zee EA, Abel T, Havekes R. A brief period of sleep deprivation causes spine loss in the dentate gyrus of mice. Neurobiol Learn Mem. 2019;160:83-90.

16.	Cirelli C, Tononi G. Effects of sleep and waking on the synaptic ultrastructure. Philos Trans R Soc Lond B Biol Sci. 2020;375(1799):20190235.

17.	de Almondes KM, Costa MV, Malloy-Diniz LF, Diniz BS. Insomnia and risk of dementia in older adults: Systematic review and meta-analysis. J Psychiatr Res. 2016;77:109-15.

18.	Simmonds E, Levine KS, Han J, Iwaki H, Koretsky MJ, Kuznetsov N, et al. Sleep disturbances as risk factors for neurodegeneration later in life. npj Dementia. 2025;1(1):6.

19.   Konakanchi S, Raavi V, MI HK, Shankar Ms V. Effect of chronic sleep deprivation and
sleep recovery on hippocampal CA3 neurons, spatial memory and anxiety-like behavior in rats.
Neurobiol Learn Mem. 2022;187:107559.
20.   Giri S, Ranjan A, Kumar A, Amar M, Mallick BN. Rapid eye movement sleep deprivation
impairs neuronal plasticity and reduces hippocampal neuronal arborization in male albino rats:
Noradrenaline is involved in the process. J Neurosci Res. 2021;99(7):1815-34.
21.   Zhang F, Niu L, Zhong R, Li S, Le W. Chronic Sleep Disturbances Alters Sleep Structure
and Tau Phosphorylation in AβPP/PS1 AD Mice and Their Wild-Type Littermates. J Alzheimers
Dis. 2023;92(4):1341-55.
22.   Zhao B, Liu P, Wei M, Li Y, Liu J, Ma L, et al. Chronic Sleep Restriction Induces Aβ
Accumulation by Disrupting the Balance of Aβ Production and Clearance in Rats. Neurochem
Res. 2019;44(4):859-73.
23.   Jaworski J, Kalita K, Knapska E. c-Fos and neuronal plasticity: the aftermath of
Kaczmarek's theory. Acta Neurobiol Exp (Wars). 2018;78(4):287-96.
24.   Pompeiano M, Cirelli C, Tononi G. Immediate-early genes in spontaneous wakefulness
and sleep: expression of c-fos and NGFI-A mRNA and protein. J Sleep Res. 1994;3(2):80-96.
25.   Sherin JE, Shiromani PJ, McCarley RW, Saper CB. Activation of ventrolateral preoptic
neurons during sleep. Science. 1996;271(5246):216-9.
26.   Cirelli C, Pompeiano M, Tononi G. Sleep deprivation and c-fos expression in the rat brain.
J Sleep Res. 1995;4(2):92-106.
27.   Terao A, Greco MA, Davis RW, Heller HC, Kilduff TS. Region-specific changes in
immediate early gene expression in response to sleep deprivation and recovery sleep in the
mouse brain. Neuroscience. 2003;120(4):1115-24.
28.   Guenthner CJ, Miyamichi K, Yang HH, Heller HC, Luo L. Permanent genetic access to
transiently active neurons via TRAP: targeted recombination in active populations. Neuron.
2013;78(5):773-84.
29.   Sanz E, Bean JC, Carey DP, Quintana A, McKnight GS. RiboTag: Ribosomal Tagging
Strategy to Analyze Cell-Type-Specific mRNA Expression In Vivo. Curr Protoc Neurosci.
2019;88(1):e77.
30.   Thompson CL, Wisor JP, Lee CK, Pathak SD, Gerashchenko D, Smith KA, et al.
Molecular and anatomical signatures of sleep deprivation in the mouse brain. Front Neurosci.
2010;4:165.
31.   Lyons LC, Chatterjee S, Vanrobaeys Y, Gaine ME, Abel T. Translational changes induced
by acute sleep deprivation uncovered by TRAP-Seq. Mol Brain. 2020;13(1):165.
32.   Delorme J, Wang L, Kuhn FR, Kodoth V, Ma J, Martinez JD, et al. Sleep loss drives
acetylcholine- and somatostatin interneuron-mediated gating of hippocampal activity to inhibit
memory consolidation. Proc Natl Acad Sci U S A. 2021;118(32).
33.   Vecsey CG, Peixoto L, Choi JH, Wimmer M, Jaganath D, Hernandez PJ, et al. Genomic
analysis of sleep deprivation reveals translational regulation in the hippocampus. Physiol
Genomics. 2012;44(20):981-91.
34.   Maciel R, Yamazaki R, Wang D, De Laet A, Cabrera S, Agnorelli C, et al. Is REM sleep a
paradoxical state?: Different neurons are activated in the cingulate cortices and the claustrum
during wakefulness and paradoxical sleep hypersomnia. Biochem Pharmacol. 2021;191:114514.

35.   Lein ES, Zhao X, Gage FH. Defining a molecular atlas of the hippocampus using DNA
microarrays and high-throughput in situ hybridization. J Neurosci. 2004;24(15):3879-89.
36.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
37.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.
38.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing biological
themes among gene clusters. Omics. 2012;16(5):284-7.
39.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks. Genome Res.
2003;13(11):2498-504.
40.   Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, et al. ClueGO: a
Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation
networks. Bioinformatics. 2009;25(8):1091-3.
41.   Tang D, Chen M, Huang X, Zhang G, Zeng L, Zhang G, et al. SRplot: A free online
platform for data visualization and graphing. PLoS One. 2023;18(11):e0294236.
42.   Delorme JE, Kodoth V, Aton SJ. Sleep loss disrupts Arc expression in dentate gyrus
neurons. Neurobiol Learn Mem. 2019;160:73-82.
43.   Kruijer W, Cooper JA, Hunter T, Verma IM. Platelet-derived growth factor induces rapid
but transient expression of the c-fos gene and protein. Nature. 1984;312(5996):711-6.
44.   Lee H, GoodSmith D, Knierim JJ. Parallel processing streams in the hippocampus. Curr
Opin Neurobiol. 2020;64:127-34.
45.   Neunuebel JP, Yoganarasimha D, Rao G, Knierim JJ. Conflicts between local and global
spatial frameworks dissociate neural representations of the lateral and medial entorhinal cortex.
J Neurosci. 2013;33(22):9246-58.
46.   Gaine ME, Bahl E, Chatterjee S, Michaelson JJ, Abel T, Lyons LC. Altered hippocampal
transcriptome dynamics following sleep deprivation. Mol Brain. 2021;14(1):125.
47.   Vecsey CG, Wimmer ME, Havekes R, Park AJ, Perron IJ, Meerlo P, et al. Daily
acclimation handling does not affect hippocampal long-term potentiation or cause chronic sleep
deprivation in mice. Sleep. 2013;36(4):601-7.
48.   Raven F, Heckman PRA, Havekes R, Meerlo P. Sleep deprivation-induced impairment of
memory consolidation is not mediated by glucocorticoid stress hormones. J Sleep Res.
2020;29(5):e12972.
49.   Wang L, Park L, Wu W, King D, Vega-Medina A, Raven F, et al. Sleep-dependent
engram reactivation during hippocampal memory consolidation associated with subregion-
specific biosynthetic changes. iScience. 2024;27(4):109408.
50.   Lopez-Rodriguez F, Wilson CL, Maidment NT, Poland RE, Engel J. Total sleep
deprivation increases extracellular serotonin in the rat hippocampus. Neuroscience.
2003;121(2):523-30.
51.   Cirelli C, Pompeiano M, Tononi G. Neuronal gene expression in the waking state: a role
for the locus coeruleus. Science. 1996;274(5290):1211-5.
52.   Bacon TJ, Pickering AE, Mellor JR. Noradrenaline Release from Locus Coeruleus
Terminals in the Hippocampus Enhances Excitation-Spike Coupling in CA1 Pyramidal Neurons
Via β-Adrenoceptors. Cereb Cortex. 2020;30(12):6135-51.

53. Liu Y, Cui L, Schwarz MK, Dong Y, Schlüter OM. Adrenergic Gate Release for Spike
Timing-Dependent Synaptic Potentiation. Neuron. 2017;93(2):394-408.

54. Vanrobaeys Y, Peterson ZJ, Walsh EN, Chatterjee S, Lin LC, Lyons LC, et al. Spatial
transcriptomics reveals unique gene expression changes in different brain regions after sleep
deprivation. Nat Commun. 2023;14(1):7095.

55. Sarma A, Ronde M, Smit S, Meerlo P, Havekes R. Does It Matter What Keeps You
Awake? Effects of Two Different Sleep Deprivation Methods on Object-Location Memory and
Hippocampal c-Fos Expression in Mice. J Sleep Res. 2025:e70079.

56. Yamazaki R, Wang D, De Laet A, Maciel R, Agnorelli C, Cabrera S, et al. Granule cells in
the infrapyramidal blade of the dentate gyrus are activated during paradoxical (REM) sleep
hypersomnia but not during wakefulness: a study using TRAP mice. Sleep. 2021;44(12).

57. Tayler KK, Tanaka KZ, Reijmers LG, Wiltgen BJ. Reactivation of neural ensembles
during the retrieval of recent and remote memory. Curr Biol. 2013;23(2):99-106.

58. Li J, Jiang RY, Arendt KL, Hsu YT, Zhai SR, Chen L. Defective memory engram
reactivation underlies impaired fear memory recall in Fragile X syndrome. Elife. 2020;9.

59. Gulmez Karaca K, Brito DVC, Kupke J, Zeuch B, Oliveira AMM. Engram reactivation
during memory retrieval predicts long-term memory performance in aged mice. Neurobiol Aging.
2021;101:256-61.

60. Mozzachiodi R, Byrne JH. More than synaptic plasticity: role of nonsynaptic plasticity in
learning and memory. Trends Neurosci. 2010;33(1):17-26.

61. Parsons RG. Behavioral and neural mechanisms by which prior experience impacts
subsequent learning. Neurobiol Learn Mem. 2018;154:22-9.

62. Pettit NL, Yap EL, Greenberg ME, Harvey CD. Fos ensembles encode and shape stable
spatial maps in the hippocampus. Nature. 2022;609(7926):327-34.

63. Sadowski JH, Jones MW, Mellor JR. Sharp-wave ripples orchestrate the induction of
synaptic plasticity during reactivation of place cell firing patterns in the hippocampus. Cell
reports. 2016;14(8):1916-29.

64. Girardeau G, Benchenane K, Wiener SI, Buzsáki G, Zugaro MB. Selective suppression
of hippocampal ripples impairs spatial memory. Nature neuroscience. 2009;12(10):1222-3.

65. Giri B, Kinsky N, Kaya U, Maboudi K, Abel T, Diba K. Sleep loss diminishes hippocampal
reactivation and replay. Nature. 2024;630(8018):935-42.

66. He Y, Sun M, Qu M, Lu Y, Yang H, Wang R, et al. Brain Insulin Signaling Pathway
Regulation of Hippocampal Neuroplasticity in Neurocognitive Disorders: Mechanisms and
Therapeutic Implications. J Integr Neurosci. 2025;24(8):39446.

67. Grillo CA, Piroli GG, Lawrence RC, Wrighten SA, Green AJ, Wilson SP, et al.
Hippocampal Insulin Resistance Impairs Spatial Learning and Synaptic Plasticity. Diabetes.
2015;64(11):3927-36.

68. Giese KP. GSK-3: a key player in neurodegeneration and memory. IUBMB Life.
2009;61(5):516-21.

69. Lauretti E, Dincer O, Praticò D. Glycogen synthase kinase-3 signaling in Alzheimer's
disease. Biochim Biophys Acta Mol Cell Res. 2020;1867(5):118664.

70. Nozal V, Martinez A. Tau Tubulin Kinase 1 (TTBK1), a new player in the fight against
neurodegenerative diseases. Eur J Med Chem. 2019;161:39-47.

71.   Wang X, Chen H, Tang T, Zhan X, Qin S, Hang T, et al. Chronic Sleep Deprivation
Altered the Expression of Memory-Related Genes and Caused Cognitive Memory Dysfunction
in Mice. Int J Mol Sci. 2024;25(21).

Figure 1

A

NSD   SD

c-Fos

CA1

Superior DG   CA3

Inferior DG

500 μm   500 μm

B₁   **Whole Hippocampus**

B₂   **Subregions**

no. of c-Fos+ cells/mm²

NSD   SD

CA1   CA3   DGsup   DGinf

○ NSD (n=5)
○ SD (n=6)

Figure 2

A

| SD-CA1 | NeuN | Sox9 |
| c-Fos | c-Fos/NeuN | c-Fos/Sox9 |

CA1

500 µm

100 µm

B

CA1

1.2%

96.7%

CA3

0.9%

87.2%

DGsup

2.7%

70.5%

DGinf

3.0%

68.7%

SD

- NeuN
- Sox9
- Others

C

SD-CA1

c-Fos

GAD67

Merge

100 µm

D

CA1

96.4%

CA3

97.5%

DGsup

94.9%

DGinf

94.3%

SD

- Excitatory
- Inhibitory

Figure 3

Figure 4

A

cFos
cFos promoter   Cre^ERT2   →   Cre^ERT2

+ 4-OHT

eYFP   IRES   Rpl22-HAtag

B

-3 weeks   Day 0
                        4-OHT i.p.   Day 7

CA1 local infusion:
AAV8-cFos-Cre^ERT2
AAV9-DIO-HA-mRpl22(RiboTag)-EYFP

NSD-NSD

SD-SD

C

HAtag

NSD-NSD   SD-SD

CA1   CA1

500 μm   500 μm

D

CA1

no. of HAtag+ cells / mm²

**

NSD-NSD   SD-SD

Figure 5

Figure 6