Title: Inferences from epigenetic information in an ecological context: a case study of early-life environmental effects on DNA methylation in zebra finches

Authors : Marianthi Tangili[1*], Per J. Palsbøll[1,2], Simon Verhulst[1*]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, The Netherlands

[2] Center for Coastal Studies, Provincetown, Massachusetts, USA

*Corresponding authors: tangilim@gmail.com, s.verhulst@rug.nl

Marianthi Tangili : 0000-0002-2871-5265

Per J. Palsbøll : 0000-0002-4198-7599

Simon Verhulst : 0000-0002-1143-6868

Abstract

DNA methylation (DNAm) plays a key role in mediating phenotypic responses to environmental variation. Various approaches exist to link DNAm variation to phenotypes, ranging from single nucleotide resolution to the development of composite indexes ('epigenetic clocks'). We here discuss the conceptual differences between these approaches through a case study using whole genome, longitudinal DNAm data from adult zebra finches raised in experimentally created small or large broods. Specifically, we (i) identified sex and age-specific CpG sites where DNAm was affected by brood size, and (ii) developed two DNAm-based indexes of the early-life environment (DMSi: based on differentially methylated sites with respect to rearing brood size; SMLmi: supervised machine learning-derived index optimized to predict brood size). We also compared results obtained by either merging or analyzing longitudinal DNAm samples separately and discussed how the permanence or transience of DNAm changes shapes responses to environmental variation through life. Our findings confirm that early-life environment leaves lasting DNAm signatures detectable in adulthood, and this effect is stronger later in life. Importantly, both indexes predicted early-life growth rates, demonstrating that DNAm-based indexes can be leveraged to retroactively quantify aspects of early-life conditions, providing a powerful novel tool in the study of wild populations.

Keywords: DNA methylation, brood size, early-life stress, epigenetic data, case study

1. Introduction

Evolutionary theory has long considered genetic variation to be the main driver of phenotypic plasticity. However, mounting evidence reveals the insufficiency of genetic variation in explaining the rapid phenotypic responses of wild animals to environmental changes (Charmantier et al., 2008; Réale et al., 2003). Epigenetic modifications are known to be sensitive to the environment (Angers et al., 2010; Li et al., 2021), modulate gene expression (Gibney & Nolan, 2010), and generate phenotypic variation without altering the primary nucleotide sequence with direct effects on evolution and adaptation (Kronholm & Collins, 2016). Emerging research also recognizes epigenetic mechanisms such as chromatin modifications and DNA methylation (DNAm) as possible regulators that modulate phenotypic plasticity and thereby link environmental variation to phenotypic outcomes. Epigenetic modifications are especially relevant in contexts when genetic variation is limited, potentially generating heritable phenotypic diversity that can be subject to natural selection (Lind & Spagopoulou, 2018).

Since methylation of different parts of the genome has different effects on gene expression (Moore et al., 2013), the design and resolution of DNAm studies can greatly impact the insights that can be drawn. All levels of analysis complement each other, providing a comprehensive understanding of the DNAm landscape and how that influences gene expression, cellular function and ultimately individual phenotype across different biological contexts. Global DNAm, which provides a single overall measure of DNAm across the entire genome of an individual (e.g. with gel-based techniques or ELISA kits) is a simple, fast and cost-effective way of studying DNAm. It can provide a broad overview of the epigenetic state of individuals, cells or tissues, which can be indicative of overall genomic stability and health (reviewed by Li & Tollefsbol, 2021). However, effects of age and environmental factors can result in increased DNAm at some CpG sites, while simultaneously resulting in DNAm decreases at other sites. As a consequence, whole genome DNAm levels are much coarser and less informative as they are skewed by repetitive regions and cannot be directly connected to regulatory pathways as evidenced by previous research in mammals and birds (Laubach et al., 2021; Meyer et al., 2023).

Higher-resolution approaches assess DNAm at the level of individual CpG sites (positions in the genome where a 5' cytosine is followed by a guanine, Moore et al., 2013) providing valuable insights, especially when directly linked to gene expression of a gene with known effects on the phenotype or life history of the organism. In both biomedical and ecological contexts, DNAm patterns of specific CpG sites have been associated with numerous traits. In clinical research, CpG-specific DNAm patterns have been linked to specific diseases (Hedman et al., 2017) and neuropsychiatric disorders (Zheleznyakova et al., 2016). In evolutionary ecology, the methylation of specific CpG sites, or entire genes has been studied in relation to a plethora of ecologically relevant traits across a wide variety of species. In birds, methylation has been associated with brood size of rearing (Jimeno et al., 2019; Sepers et al., 2021), food deprivation (Xiao et al., 2020), and urbanization (Caizergues et al., 2022). In wild baboons, methylation was linked to resource availability (Lea et al., 2016), in lizards to habitat quality (Hu et al., 2019), and in crustaceans to the exposure to pollution (Harney et al., 2022). Collectively, these findings confirm that DNAm is sensitive to environmental perturbations and a putative mechanism for generating phenotypic variation.

A fundamentally different approach to make inferences based on DNAm, compared to the focus on individual CpG sites, is the development of methylation-based indexes. Utilizing the methylation of selected positions in the genome, an algorithm can be trained and later used as a

predictive tool using e.g. supervised machine learning. Such indexes have proven to be valuable
for diagnostic and therapeutic purposes, for example for non-invasive cancer detection and
identifying cancer types (Draškovič & Hauptman, 2024; Wei et al., 2021), prediction of patient
response to chemotherapy (Wick et al., 2014), and assessment of the impact of lifestyle and
therapeutic interventions (Keller et al., 2020; Lu et al., 2023). This approach has in particular
been applied to develop epigenetic 'clocks' of chronological age, algorithms that use DNAm
patterns to predict chronological age for humans, model, and non-model animals (Horvath,
2013; Petkovich et al., 2017; Tangili et al., 2023).

Despite the growing use of DNAm based indexes, it is still debated whether the CpG sites
underlying such indexes are functionally involved in the targeted process, or if the level of
methylation at the identified sites correlates with those processes in absence of any biological
function. The functional role of the DNAm of CpG sites outside of regulatory regions remains
poorly understood (Moore et al., 2013) and confounded by the limited overlap between the CpG
sites identified across different studies aimed at the same biological process within the same
species (Galkin et al., 2020; Liu et al., 2020). Part of the explanation might be that phenotypic
variation rarely correlates strongly with single genes and hence methylation-based indexes likely
capture CpG sites across gene networks scattered across the genome. In this sense, an index
comprising DNAm at many CpG sites may better match the causes of the phenotypic variation
under study, compared to the DNAm of single sites. Nevertheless, whether these sites are
causally linked to the observed phenotypes remains an open question, demanding more research
across diverse species to elucidate the underlying mechanisms.

We here present a case study to explore different approaches to utilize DNAm data to gain
information on the role of DNAm in mediating the effects of the environment on the phenotype.
We have utilized longitudinal whole genome methylation data from fifty adult zebra finches
(*Taeniopygia castanotis*) of both sexes that were reared in experimentally created small or large
broods. This manipulation is of interest, firstly because it is reasonable to assume that parents
and offspring have evolved responses to variation in brood size because it shows strong natural
variation, and hence we can reasonably assume that phenotypic effects are not artefacts.
Secondly, previous studies from the present study population and others, have documented
multiple long-term effects of the nutritional stress caused by this manipulation; for example,
individuals reared in large broods have shorter lifespans (Briga et al., 2017) and lower sexual
attractiveness (Holveck & Riebel, 2010).

Longitudinal data, collected from the same individuals at multiple time points, are essential
when studying processes such as development, aging and adaptation and allow for the
continuous monitoring of the molecular state of an individual. However, such data is scarce and
most epigenetic studies are cross-sectional. Knowing how to analyze and interpret longitudinal
epigenetic data is crucial for understanding complex phenotypic traits, but there is as still little
experience with utilizing longitudinal samples in epigenetic studies. In our case study we
sampled all individuals twice, once in early adulthood, and a second time closer to the end of
their lives, and explored different ways to combine the information of longitudinal samples.

In our case study, we first identified CpG sites with differential methylation between treatments,
for the sexes pooled but also for the sexes separately, to identify sex-specific effects of
developmental conditions. Secondly, to translate methylation information into interpretable
metrics, we developed two methylation-based indexes: an epigenetic early life quality index
(DMSi) calculated from methylation data of differentially methylated sites between individuals

raised in small and large broods and a predicted brood size index (SMLmi) calculated using a
supervised machine learning algorithm with the goal to predict brood size of rearing. We
hypothesized that these indexes could serve to quantify effects of early life conditions on the
methylome within the treatment categories, and as such provide a more accurate reflection of an
individual's phenotypic state than the coarse binary trait of manipulated brood size. If
successful, an estimate could in this way be obtained of the early life conditions experienced by
individuals first encountered in adulthood, providing a new tool for studies in the wild where a
substantial part of the population can consist of immigrants with unknown history.

2. Case study

2.1 Case study: methods

2.1.1 Subjects and samples

The present data set is based on samples taken from birds in the hard foraging conditions in the
experiment reported in (Briga et al., 2017), where methods are described in detail. In brief, adult
zebra finches were randomly matched and provided with nest boxes and nesting material.
Fortified canary food was supplied to the parents in addition to tropical seed mixture and water
until the hatching of the first chick in the nest but not thereafter. Nestboxes were checked daily
for the presence of eggs or chicks. Chicks were cross-fostered at age five days or before to broods
of either two or six chicks, which is the range of brood sizes observed in the wild (Zann, 1996).
On the day of cross-fostering, all chicks were weighed and individually marked, and weighed
again when 15 days old (or as near to that as practically possible). Siblings were assigned to
different broods, and parents never reared their own offspring. When reaching adulthood (≥3
months), birds were moved to outdoor aviaries each containing single sex groups of 18–24
adults. We here used blood samples collected from the brachial vein and stored in -80℃ in
glycerol buffer from 50 individuals distributed equally over the small and large broods (n=25 x
2) and almost equally over the two sexes (26 males, 24 females). Given all other selection
criteria, we could not make within natal family comparisons with respect to the effect of rearing
brood size, although this would have further increased statistical power. Each bird was sampled
every six months after entering the experiment, and DNAm data were extracted for two samples
per individual collected early and late in their life, with an average sampling interval of 995
(S.D.=742) days - see Fig.S1 for the age distribution at sampling.

2.1.2 Enzymatic Methyl-seq data

We extracted DNA according to the manufacturer's protocol using innuPREP™ DNA Mini Kit
(Analytik Jena GmBH) from 3 uL of nucleated red blood cells. Next-generation sequencing was
conducted by the Hospital for Sick Children (Toronto, Canada). The amount of DNA was
quantified using a Qubit High Sensitivity Assay (Thermo Fisher Scientific). Two hundred ng of
DNA was used as input material for library preparation using the Next Enzymatic Methyl-
seq™ Kit (New England Biolabs Inc.) following the manufacturer's instructions. Briefly, DNA was
fragmented by sonication to an average length of ~500 base pairs (bp) using a Covaris LE220
(Covaris Inc.). Fragmented DNA was end-repaired and ligated to Illumina sequencing adapters.
The 5-methylcytosines and 5-hydroxymethylcytosines were oxidized by TET2 and cytosines
were deaminated by APOBEC. Methyl-seq libraries were subjected to five PCR amplification
cycles. Libraries assessed on a Bioanalyzer™ DNA High Sensitivity chip (Agilent Inc.). The

amount of DNA was quantified by quantitative PCR using the Kapa Library Quantification
Illumina/ABI Prism™ kit according to the manufacturer's instructions (La Roche Ltd.). Libraries
were pooled in equimolar amounts and sequenced to a minimum of ~100M paired-end reads
(2x150bp) per sample on a 10B flow cell using a NovaSeqX™ (Illumina Inc.) platform.
174
Sequences were trimmed using Trim Galore! v0.6.10 (Krueger et al., 2023) in paired-end mode.
The data were assessed before and after trimming using FastQC v. 011.9 (Andrews, 2010) and
MultiQC v. 11.14 (Ewels et al., 2016). Trimmed reads were aligned against the *in silico* bisulfite
converted zebra finch (*Taeniopygia castanotis*) reference genome (GCA_003957565.4, Rhie et al.,
2021) using Bismark v. 0.14.433 (Krueger & Andrews, 2011) using the Bowtie 2 v. 2.4.5
alignment algorithm (Langmead & Salzberg, 2012). The average mapping efficiency was at
65.14% (SD: 2.79). Mean autosomal DNAm did not appear to be affected by the storage time
(range 8-16 years) of the samples (estimate= 0.03 ± 0.04, t = 0.85, p = 0.39).

2.1.3 Differentially methylated site (DMS) identification

We performed multiple epigenome wide association studies (EWASs) in order to identify
differentially methylated sites (DMSs) between zebra finches raised in small and large broods
using *Methylkit* v. 1.18.0 (Akalin et al., 2015).
For all EWAS analyses the following steps were followed:
1. CpG sites were filtered for coverage, retaining those with a minimum coverage of 5 and
maximum coverage the 99.9[th] percentile coverage in order to avoid PCR bias (Wreczycka
et al., 2017). For the analysis where the two samples from the same individual were
merged, a minimum coverage filter of 12 was applied.
2. Read coverage distributions across all samples were normalized using the
normalizeCoverage function of the *Methylkit* package.
3. Only CpG sites shared by all samples were retained using the *unite* function in the
Methylkit package.
4. Invariable CpG sites with a methylation of zero or one hundred percent in all samples
were excluded.
5. Logistic regression models with overdispersion correction following the McCullagh and
Nelder method ("MN", McCullagh & Nelder, 1989) were used to test for significant
methylation differences between birds raised in small and large broods, accounting for
extra-binomial variation in read counts. Resulting p-values were adjusted for multiple
testing using a Bonferroni correction (Dunn, 1961).
We considered as DMSs all sites with a significant (≤0.05) Bonferroni-corrected p-value.
204
205 In order to explore the long-term effects of brood size on DNAm per CpG site, data of the two
206 observations per individual were merged by adding the amount of methylated counts and
207 dividing by the combined coverage of the two observations. Merging the DNAm data per
208 individual increases coverage and therefore presumably precision.
209
210 When analyzing individual samples, filtering retained 1,114,704 CpG sites, whereas merging two
211 samples per individual increased the number to 2,085,308 sites. Running the analysis by sex
212 resulted in 1,896,932 and 1,787,898 CpG sites retained for females and males, respectively. After
213 filtering, 60,046 CpG sites remained in the analysis for the first sample of all individuals (young
214 age), and 82,273 sites in the second sample of all individuals (old age).
215

216 To assess whether age at sampling of the zebra finches influenced the differential methylation
217 calculation, we adding age as a covariate (in all analyses using *Methylkit* except from the one
218 where samples from the same individual were merged). The inclusion of age at collection as a
219 covariate did not affect the results (methylation difference, significance) so it was omitted from
220 subsequent analyses.

2.1.4 Epigenetic early-life quality index (DMSi) calculation

222 We calculated an epigenetic early life quality index, which we termed DMSi, as follows: DNAm of
223 all identified DMSs for the analysis per sample as mean centered, and for sites where DNAm was
224 on average higher in birds from enlarged broods, mean centered DNAm was multiplied by -1.
225 DNAm was then averaged per sample or individual over all DMSs. To facilitate comparisons with
226 future studies we z-scored the index prior to analyses.

2.1.5 Supervised Machine Learning methylation index (SMLmi) calculation

As a second method to develop a DNAm index of early life quality, we performed an elastic net
regression using DNAm at variant sites (i.e. DNAm not of zero or one hundred in all samples)
with a coverage between ten and the 99.9^{th} percentile of coverage. We used DNAm per CpG site
per sample as the predictor variable and brood size of rearing as continuous dependent variable
(coded as -0.5 and +0.5 for small and large rearing brood size respectively). The elastic net
mixing parameter was set to $\alpha = 0.1$, as a titration procedure showed this value to yield the most
accurate model (i.e. most samples assigned to the correct brood size). To determine the optimal
regularization parameter λ, we first used the function *cv.glmnet* from package *glmnet* v. 4.1-8
(Friedman et al., 2010) with a 10-fold cross-validation. The selected λ was the applied in a leave-
one-out cross-validation (LOOCV) framework, in which one sample was omitted each time and
the model was re-fitted on the remaining data, and the brood size of the omitted sample was
predicted. Lastly, a final elastic net model was fit using all samples and the optimized
hyperparameters. CpG sites with non-zero regression coefficients were extracted as features
retained by the final model.

2.1.6 Statistical analyses

We implemented linear mixed effects models using package *lme4* v. 1.1.35.1 (Bates et al., 2015)
in R v. 4.1.2 (R Core Team, 2023) in order to explore how the DMSi or SMLmi differed between
the sexes and individuals raised in small and large broods. We employed Akaike's An Information
Criterion (AIC) values (Akaike, 1974) to compare the predictive performance of models. Partial
correlations were calculated using the *pcor* function from the *ppcor* v. 1.1 package (Kim, 2015) in
R.

2.2 Case study: differentially methylated sites (DMSs)

When analyzing longitudinal DNAm data to extract differential methylation information, several
analytical strategies can be applied. We identified DMSs between birds raised in small and large
broods (i) separately for the two samples from each individual, (ii) merged per individual, (iii)
separately for the first and second samples per individual (young and old) and (iv) separately
for males and females. In this way, we can evaluate the effect of pooling samples taken at
different ages on the detection of DMSs, on the consistency of DMSs over life, and on the age-
dependence of frequency and magnitude of DMSs. Moreover, we can get insights into sex-specific
effects of developmental conditions on DNAm.

2.2.1 DMSs per sample vs. individual

When considering individual samples, the range of percent methylation difference between zebra finches raised in small and large broods was from -34.1 to 34.1% (Fig.1a). We identified 27 DMSs (Fig.1a), 20 of which were located in chromosome 3, two in chromosome 1A, two in chromosome 2 and one each in chromosomes 1, 10 and 7.

We extended the above analysis by combining the methylation data from the two samples collected early and late in life for each individual. The range of percent methylation difference between zebra finches raised in small and large broods was -34.5 to 34.2% (Fig.1b). However, this approach did not yield any DMSs between individuals raised in small and large broods (Fig.1b).

When comparing the absolute difference in methylation of all CpG sites in the analysis per sample vs. the merged samples using a Wilcoxon rank-sum test, this revealed a small but highly significant difference between groups (Fig.S2; $W = 1.15 \times 10^{12}$, $p<0.001$), with higher absolute differences between birds raised in small and large broods observed in the merged dataset. This is somewhat paradoxical, given that there are fewer DMSs identified in the pooled samples, for which we have no clear explanation. Importantly, these results indicate that effect-sizes are sensitive to how repeated measurements are treated, with the apparent gain in precision due to a higher coverage potentially outweighed by changes over life overshadowed by pooling samples.

2.2.2 DMSs in young vs. old cohorts

The range of percent methylation difference between zebra finches raised in small and large broods was from -19.1 and 21.1% between young (Fig. 2a) and -25.8 and 28.41% between old birds (Fig.2b). After filtering, 49,538 sites (of the 60,046 in the young and 82,273 in the old cohort) were shared between the young and old cohort data sets, Across all 49,538 CpG sites, methylation differences between zebra finches raised in small and large broods were positively among age cohorts (Fig.S3, r= 0.29, $p < 0.001$), indicating a significant repeatability over life of the brood size effect on DNAm. However, we identified zero brood size-related DMSs when individuals were respectively young or old (Fig.2). CpG sites with the largest methylation differences between small and large broods were not shared by the young and old datasets. In samples taken early in life, there were more methylation differences around zero, while the most extreme methylation differences are found in the old cohort (Fig.S4). Indeed, when comparing the absolute methylation difference over all CpG sites using a Wilcoxon rank-sum test, this revealed a small but highly significant difference between samples taken early and late in life (Fig.S5; $W = 2.16 \times 10^9$, $p < 0.001$). Thus, the magnitude of the effect of early-life conditions on DNAm differed significantly within individuals over life, with the changes being more pronounced later in life, but, at the same time, no significant DMSs were observed when considering samples taken early and late in life separately.

2.2.3 DMSs in females vs. males

The analysis for identification of DMSs was run separately for the two sexes. The range of percent methylation difference between zebra finches raised in small and large broods was from -48.6 and 47.4% in females (Fig.3a) and from -43.5 to 45.3% for males (Fig.3b), i.e. considerably larger then when observed when the sexes were pooled (section 2.2.1). After filtering, 1,108,263 CpG sites were shared between males and females. Across all 1,108,263 CpG sites, there was a

weak but significant negative correlation between brood size effects on DNAm in females and males (Fig.S6, r = -0.04, p<0.001).

We identified 34 DMSs between females and 38 DMSs between males raised in small and large broods (Fig.3). The majority of female DMSs (N=16) were located in chromosome 4 while the majority of male DMSs (N=19) were located in chromosome 3. Moreover, in females, the majority of DMSs exhibited negative methylation differences (7 positive vs. 27 negative) whereas in males, the majority were positive (30 positive vs. 8 negative). This difference in distribution was highly significant (χ^2 = 22.18, df = 1, p < 0.001). These findings indicate a sex-specific pattern in methylation changes between zebra finches raised in small and large broods.

In addition to the sex difference in average direction of DMSs, the absolute effect of brood size on all CpG sites was also sex-dependent. When comparing the absolute difference in methylation between birds raised in small and large broods across all CpG sites that passed the filters for females (N = 1,896,932) and males (N = 1,787,898) using a Wilcoxon rank-sum test, this revealed a small but highly significant difference between groups (Fig.S7; W = 1.64 × 10^{12}, p<0.001), with females having larger absolute differences in methylation (Fig.S7). The above observations indicate sex-specific epigenetic responses to early-life environment.

The identification of CpG sites across the genome with significant methylation differences between adult zebra finches raised in small and large indicates that early-life conditions are associated with persistent, site-specific variation in DNAm across the genome. Previous work on nestling zebra finches has also shown consistent effects of brood size on DNAm (Jimeno et al., 2019; Sheldon et al., 2018). These changes in DNAm between individuals raised in different brood sizes might underlie the phenotypic differences observed between treatment groups such as the significantly shorter lifespan of zebra finches raised in large broods we have previously identified (Briga et al., 2017). Using longitudinal samples, that are therefore immune to effects of selective disappearance, we further extend these findings, by revealing that early-life environment effects on DNAm are stronger later in life. This raises the question whether early-life environmental effects on the phenotypic level also increased with age, but this remains to be investigated. In any case, our findings highlight the potential role of DNAm in mediating the enduring impact of early-life environmental conditions on adult phenotypes (Dettmer & Chusyd, 2023).

2.3 Case study: Differentially Methylated Site index (DMSi)

Methylation-based, predictive indexes are widely used in clinical research and diagnostics (Xu et al., 2024; Yousefi et al., 2022; Zhang et al., 2025), but have so far only been used to predict chronological age in an ecological context (Tangili et al., 2023). Developing methylation-based biomarkers could allow quantification of other traits that are difficult to estimate directly in individuals not previously encountered, e.g. early-life stress and other significant life events. The development of such biomarkers holds promise especially when studying wild populations, where detailed, individual life-history information might be labor-intensive to obtain and therefore scarce. The development of targeted, cost-effective methods to measure DNAm of selected CpG sites (e.g. Morselli et al., 2021), makes feasible the collection of population data with unprecedented accuracy.

We constructed an epigenetic index of early developmental conditions (i.e. brood size) using the methylation of the DMSs identified between zebra finches raised in small and large broods

(DMSi) using individual samples (N=27). Because birds raised in large broods achieved a shorter lifespan (Briga et al 2017), we hypothesized that this index would serve as a quantitative measure of the overall phenotypic quality of each individual.

DMSi calculated over samples collected early and late in life on the same individuals was highly correlated (Fig 4a, r = 0.99, $p < 0.001$), indicating consistence of the developmental effects on DNAm over life. The DMSi differed strongly between birds reared in small and large broods, as expected (Fig. 4b-d). Interestingly, there was also a large difference in the DMSi variance between birds reared in small (0.04) and large (1.08) broods (F = 0.038, 95% C.I. 0.021 - 0.066, d.f. = 49, $p < 0.001$). We infer this to reflect that development in small broods is less variable compared to large broods, due to nutritional stress being more likely to occur in large broods, with larger variation between individuals in how they are impacted due to sibling competition (Tangili et al., 2025). In our study population, females have a shorter lifespan than males due to higher actuarial senescence (Briga et al., 2017), and we therefore anticipated the DMSi to be lower in females. However, independent of age, the DMSi in females was only marginally lower compared to males (Table 1a; Fig.4b). Because birds from large broods achieved a shorter lifespan (Briga et al., 2017), we speculated that the DMSi would decline with age when being old would imply having a phenotype that more resembles having grown up in a large brood. However, no significant effects of brood size and age on DMSi were detected (Fig. 4c,d), and this did not change when treating age as a numerical variable instead of a category. Additionally, neither sex nor interactions between age cohort, sex, or brood size significantly influenced DMSi values (Table 1a), suggesting that the effect of brood size on DMSi is consistent across ages and sexes.

2.4 Case study: Supervised Machine Learning methylation index (SMLmi)

When predicting brood size using the merged samples per individual, a ridge regression ($\alpha=0$, using the methylation of all CpG sites to predict brood size with optimized weights) yielded the best result. However, the predictive accuracy was only 42%, i.e. slightly worse than random assignment. This indicates that no individual CpG or small subset of CpGs in the pooled samples was strongly predictive of rearing brood size, which is in line with the identification of DMSs in this data set (Fig. 1). A predictive modeling approach on all individual samples using elastic net regression with leave-one-out cross-validation yielded a different result, predicting rearing brood size with an accuracy of 89% (Fig.5). When inspecting the 11 wrongly assigned samples, it became apparent that eight of these samples were from four individuals of which both samples were wrongly assigned (Table S1). The elastic net model SMLmi utilized the methylation of 455 CpG sites (0.04% of all CpG sites that passed filtering, Fig.S8) spread over 30 autosomal chromosomes and the Z chromosome (Fig.S8b), where it appears that there were more included CpG sites than expected by chance on chromosome 3, and fewer than expected by chance on 1 (Fig.S8a). Five SMLmi CpG sites were shared with the DMSs per sample and three with the male-specific DMSs (Fig.S4). As expected, the SMLmi differed strongly between birds reared in small and large broods but the difference in the SMLmi variance between birds reared in small (0.020) and large (0.023) broods did not approach significance (F = 0.86, 95% C.I. 0.49 - 1.52 , d.f. = 49, $p = 0.61$), in contrast to the finding with the DMSi. Age, either as category (young / old) or as continuous variable, and sex did not explain significant variation in the SMLmi (Table 1b).

2.5 Case study: Comparing indexes

Across samples, we identified a strong negative correlation between DMSi and SMLmi (Fig.6, r = -0.73, p < 0.001). This pattern suggests that both indices capture biologically meaningful epigenetic information about the early-life environment. There was no support for a non-linear relationship, as the inclusion of SMLmi squared or otherwise transformed did not improve the model. When examining this relationship within brood size treatments, the negative association was apparent in both groups (small broods: r = -0.5, p < 0.001, large broods: r = -0.56, p < 0.001). A z-test revealed that the difference in slopes between small and large broods was statistically significant (z=3.7, p < 0.001) suggesting that the relationship between DMSi and SMLmi differed depending on brood size. This finding is in agreement with the larger variance in DMSi among individuals reared in large broods. However, a mixed-effects model including an interaction between SMLmi and brood size did not detect a significant difference in the strength of association between brood sizes (brood size * SMLmi: β = 0.19 ± 0.27, t = 0.7, p = 0.49), which we attribute to the different position of the two data sets on the X-axis (Fig.6).

For most female-specific DMSs the methylation difference was negative and for male-specific DMSs the difference was positive (Fig.S10), meaning that birds in large broods tended to have lower methylation at female specific DMSs while the opposite pattern occurred in male-specific DMSs. The CpGs selected for the SMLmi showed intermediate methylation differences compared to the DMSs (Fig.S10). This distinction reflects the different goals of the two analyses; identifying the CpG sites with the largest methylation differences versus accurately predicting brood size. The CpGs selected to increase SMLmi performance do not necessarily represent the largest differences in methylation between treatments. This is to be expected, with machine learning algorithms selecting CpG sites that add the largest amount of information given the other selected CpG sites. CpG sites where DNAm responds strongly to treatment (i.e. brood size in our study) in most cases add relatively little information because DNAm on these sites will be strongly correlated with DNAm on selected sites where DNAm responded to treatment.

2.6 Case study: Indexes and early-life growth rate

To assess to what extent the two methylation-based indexes relate to individual variation in early-life conditions on a phenotypic level instead of brood size treatment, we correlated DMSi and SMLmi to pre-fledging growth rates. Early-life growth rate is affected by brood size (Briga et. al 2017) and a robust predictor of fitness prospects across a wide range of species (Douhard et al., 2014; Drake et al., 2025), including birds (Brown et al., 2022). Both indices significantly predicted individual variation in growth rate (Fig.8; Tables S2,3). The partial correlation coefficients for mass gain per day were 0.5 for the DMSi and -0.48 for the SMLmi, and comparison of the AIC values of the predictive models revealed that the DMSi was a slightly better predictor of early-life growth rate than SMLmi (AIC = -47.7 and -46.8, respectively). However, when adding rearing brood size to the models in tables S2 and S3, this caused the index effects to be no longer significant (DMSi: β = 0.02 ± 0.02, t = 1.05, p = 0.3; SMLmi: β = 0.03 ± 0.05, t = 0.53, p = 0.6). Thus, both indexes can be predictive of early-life growth, but are not an improvement over knowing the actual early-life conditions, indicating also that growth does not fully characterize the phenotypic consequences of developmental conditions.

3. Discussion

Epigenetic information can be utilized in different ways to understand how environmental factors modulate phenotypic traits in the short and long-term, affecting organismal sensitivity, resilience and adaptability. Studying DNAm at the level of each individual CpG site offers the highest resolution for understanding how DNAm directly affects phenotypes depending on its location, because the methylation of different parts of the genome has different effects on gene expression (Moore et al., 2013). However, even CpG site-level analyses can differ in resolution. Depending on the approach employed to quantify methylation, and the resulting resolution of the data, studies examined methylation in CpG sites across the entire genome, focus on CpG islands, or target specific genomic regions or functionally relevant loci. However, focusing exclusively on genes or genomic regions predicted to be affected by a treatment introduces significant limitations. Novel effects outside of the predefined regions may be overlooked, leading to a possibly misleading understanding of the treatment's effects on the epigenome. Furthermore, this approach limits the opportunity to use the data in ways not envisaged when designing the study. Taking into consideration all the above, we consider comprehensive, high resolution, genome-wide analyses essential to capture the full spectrum of epigenetic differences between phenotypes that can be leveraged to get insights into the mechanistic underpinnings of phenotypic differences between individuals.

Leveraging whole-genome methylation data from adult zebra finches raised in either small or large broods, we presented a case study to compare different ways to quantify the long-term consequences of early-life developmental conditions on the methylome using longitudinal samples. When identifying DMSs, longitudinal samples, taken at different times from an individual, can either be merged, increasing coverage per site and therefore potentially precision, or analyzed separately. We have here presented analyses on merged samples, separately for the two samples from each individual, taken early and late in life in each individual and separately for males and females.

Judging by the number of DMSs identified, using individual samples was the more powerful approach, and we consider these findings the best estimate of the (long-term) effects of rearing brood size on CpG site specific DNAm. Merging the samples concealed the interesting finding that DNAm differences due to rearing brood size were larger later in life, raising the question whether phenotypic differences due to rearing brood size were also larger later in life. We therefore conclude (not surprisingly), that whether it is recommended to merge longitudinal samples depends on the details of the question that is being addressed, with merging being useful to increase precision when estimating long-term effects, while addressing age-related variation requires samples taken at different times to be analyzed separately.

Statistically, higher coverage at a given CpG site reduces stochastic noise and improves reliability of the methylation estimate but systematic biological differences between the two samples may be averaged out, potentially masking temporal or condition-dependent changes. Whether longitudinal samples were merged or not also affected the development of indexes of early-life conditions. For example, we observed a strong reduction in power of the SMLmi to predict brood size from the merged samples compared to a similar analysis based on the individual samples. This finding indicates that using aggregated longitudinal data might not be suitable for trait prediction and offers further support to the idea that the difference in methylation between individuals raised in small versus large broods varies across different life stages as previously demonstrated in great tits (*Parus major*, Sepers et al., 2023, 2024). We therefore tend towards

recommending the calculation of indexes using individual samples, but acknowledge that more
comparisons between indexes from merged versus single samples will be necessary before
general conclusions can be reached with confidence.

When studying natural populations, developmental conditions of a proportion of individuals will
usually not be known, and this proportion can be very high. Our findings suggest that the indexes
we calculated based on the effects of manipulated brood size on DNAm offer the opportunity to
obtain information on the developmental conditions of individuals first encountered in
adulthood. The high predictive accuracy of brood size by SMLmi is encouraging in this respect, as
was the strong association between the indexes and early-life growth rate. The actual brood size
of rearing was still a better predictor of growth rate than either index, but when direct
information about developmental conditions is absent, indexes as developed in this study are
possibly the best available alternative, besides traits that can be measured on adults (e.g.
morphological traits that are fixed early in life). We recognize, however, that such indexes
probably have to be developed and validated for each species or population, and also within
populations it remains to be investigated as how stable such indexes are in space and time.
Moreover, it remains to be demonstrated that our approach yields similar results in the wild,
where environmental heterogeneity is likely to be larger, and survival selection on early life
development is stronger. Nonetheless, our findings suggest that at least within the context of an
ongoing study where developmental conditions are known for part of the population, DNAm
based indexes potentially provide a unique tool to assess the developmental conditions of
individuals first encountered as adults.

Indexes integrating DNAm information from many CpG sites are likely to have greater
explanatory power with respect to phenotypic variation, when compared to individual CpG sites.
Nevertheless, we acknowledge that in order to fully understand the mechanisms through which
DNAm mediates phenotypic variation, it is ultimately necessary to investigate the functional
roles of individual CpG sites. The SMLmi may then be a better starting point than the DMSi, as
that CpG selection method includes validation on independent samples (through the leave-one-
out protocol), but for either indexes it would be advisable to test its repeatability using an
independent but fully comparable data set. While both indexes we calculated (DMSi & SMLmi)
were highly correlated with each other, the two indexes draw information from sets of CpG sites
that showed limited overlap (Fig.S9). Instead of considering this a weak point, we see it as
broadening of the opportunity to explore the mechanisms underlying phenotypic variation. This
divergence generates the possibility that the two indexes capture multiple and possibly partially
independent epigenetic signatures that can contribute to the same phenotypic outcomes. Thus,
we see the development of DNAm based indexes of phenotypic traits as a stepping stone towards
the identification of (networks of) CpG sites that together play an important role in shaping
phenotypic variation.

As illustrated by our findings when comparing results based on either single samples or merged
longitudinal samples per individual, DNAm is a dynamic and modifiable process, not only during
early life and development (Watson et al., 2019) but throughout an individual's entire lifespan.
Information on the dynamics of DNAm at individual CpG sites is limited, in particular in non-
model systems, but whether DNAm responses to environmental changes are transient or
permanent has large consequences for the evolution of such responses (Fig.8). Environmental
changes that induce a phenotypic response through DNAm can impact fitness in different ways,
both before and during adulthood, depending on whether the change in DNAm is transient or

persistent. In either case, it can be assumed that an environmentally-induced change in methylation provides fitness benefits in the short term when it is an evolved response. However, different scenarios can be envisaged for long-term consequences (Fig.8). In one scenario, the induced DNAm causes a permanent (favorable) change in the phenotype, in which case the fitness benefits are permanent, regardless of whether the change in DNAm is maintained, making this scenario less relevant in the context of this discussion. Alternatively, methylation patterns can be stably maintained when the different DNAm state is also beneficial in adulthood, maintaining also associated fitness benefits. When the effect on DNAm is transient, and no longer beneficial, the DNAm change may diminish as the individual matures, becoming neutral with respect to fitness. Lastly, an antagonistic pleiotropic effect can arise, where due to a constraint on phenotypic flexibility a change in DNAm that is beneficial early in life is permanent and becomes deleterious in later life. The above scenarios showcase how early-life methylation modifications resulting from early-life environmental exposures can be reinforced, attenuated or altered by later-life exposures, which can lead to complex, lifelong phenotypic effects. A better understanding of the extent to which DNAm dynamics are transient or permanent will be essential for our understanding of the impact of DNAm on the relationship between environment, aging, phenotype and fitness.

3.1 Limitations

While present findings reveal a clear association between site-specific DNAm and brood size of rearing, the mechanistic pathways through which early-life conditions affect DNAm remain unclear. In avian species, a harsh development can have profound, direct effects on Darwinian fitness, influencing traits such as growth (de Kogel, 1997; Nilsson & Svensson, 2001), recruitment (Schwagmeyer & Mock, 2008), nestling survival (Mock et al., 2009), foraging behavior (Andrews et al., 2015) and (lifetime) reproductive success (Eastwood et al., 2019; Spagopoulou et al., 2020). More specifically, in zebra finches being raised in a large has been found to affect nestling behavior (Tangili et al., 2025), social integration (Gerritsma et al., 2025), mate preference (Holveck & Riebel, 2010) and lifespan (Briga et al., 2017). Future research efforts should aim to disentangle whether these behavioral and physiological consequences arise through direct effects of the development on the methylome or that they reflect secondary pathways mediated by altered physiological and social dynamics. A relevant concern in this context is that DNAm effects of developmental conditions may arise indirectly, in response to phenotypic changes triggered by the early-life environment (Höglund et al., 2020) , instead of underlying the observed phenotypic changes.

The extent to which our findings can be generalized remains an open question, given that the identification of differential methylation can be highly study and/or method-dependent, in that for example data type, analysis tools, filtering choices, statistical framework, and significance thresholds can strongly influence the results. For example, in great tits, initial approaches using pooled data and relatively underpowered methods suggested significant effects of brood size on nestling DNAm (Sepers et al., 2021), but a subsequent study using more refined methods did not find such an effect (Sepers et al., 2023).

Although our study in effect controlled for genetic background by randomly cross-fostering chicks from different genetic backgrounds to treatments, differences in methylation identified in this study should be interpreted with the caveat that some may reflect underlying genetic differences rather than causal effects of the brood size manipulation. Lastly, the modest sample size prevented subgroup analyses, such as age comparisons separately for the two sexes.

3.2 Conclusions

This study demonstrates effects of (manipulated) brood size in which an individual bird is raised on site-specific DNAm in adulthood in a sex-specific manner. Moreover, we have here illustrated how methodological choices can have profound effects on the identification of differential methylation, underscoring the need to align analytical pipelines to the specific study aims. It also suggests there is a need for simulation studies to investigate power and biases inherent in the different approaches to the analysis of DNAm. Lastly, the present study highlights the need to understand the causal pathways by which early-life environments affect the phenotype through methylation, including when the methylation marks are established, how stable they are across tissues and life stages, and under which conditions they translate into lasting physiological and behavioral changes.

Acknowledgements

We thank the Center for Information Technology at the University of Groningen for support and access to Hábrók high-performance computing cluster. We express our gratitude to Ellis Mulder for her help in the laboratory, to Michael Briga and Blanca Jimeno for running the long-term experiment and to the animal caretakers at the University of Groningen as well as numerous students whose invaluable help made this project possible.

References

Akaike, H. (1974). A New Look at the Statistical Model Identification. In E. Parzen, K. Tanabe, & G. Kitagawa (Eds.), *(eds) Selected Papers of Hirotugu Akaike*. Springer Series in Statistics. https://doi.org/https://doi.org/10.1007/978-1-4612-1694-0_16

Akalin, A., Franke, V., Vlahovic, K., Mason, C. E., & Schu, D. (2015). Genome analysis genomation: a toolkit to summarize , annotate and visualize genomic intervals. *Bioinformatics*, *31*(November 2014), 1127–1129. https://doi.org/10.1093/bioinformatics/btu775

Andrews, C., Viviani, J., Egan, E., Bedford, T., Brilot, B., Nettle, D., & Bateson, M. (2015). Early life adversity increases foraging and information gathering in European starlings , Sturnus vulgaris. *Animal Behaviour*, *109*(109), 123–132. https://doi.org/10.1016/j.anbehav.2015.08.009

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data.* https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Angers, B., Castonguay, E., & Massicotte, R. (2010). Environmentally induced phenotypes and DNA methylation: How to deal with unpredictable conditions until the next generation and after. *Molecular Ecology*, *19*(7), 1283–1295. https://doi.org/10.1111/j.1365-294X.2010.04580.x

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using {lme4}. *Journal of Statistical Software*, *67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Briga, M., Koetsier, E., Boonekamp, J. J., Jimeno, B., & Verhulst, S. (2017). Food availability affects adult survival trajectories depending on early developmental conditions. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1846), 20162287. https://doi.org/http://dx.doi.org/10.1098/rspb.2016.2287

Brown, T. J., Dugdale, H. L., Hammers, M., Komdeur, J., & Richardson, D. S. (2022). Seychelles warblers with silver spoons: Juvenile body mass is a lifelong predictor of annual survival , but not annual reproduction or senescence. *Ecology and Evolution*, *12*(7), 1–16.

605   https://doi.org/10.1002/ece3.9049

606 Caizergues, A. E., Le, J., Arnaud, L., Marta, G., Senar, J.-C., Charmantier, A., & Perrier, C. (2022).
607   Epigenetics and the city⬚: Non- - parallel DNA methylation modifications across pairs of
608   urban- - forest Great tit populations. *Evolutionary Applications*, *15*, 149–165.
609   https://doi.org/10.1111/eva.13334

610 Charmantier, A., McCleery, R. H., Cole, L. R., Perrins, C., Kruuk, L. E. B., & Sheldon, B. C. (2008).
611   Adaptive phenotypic plasticity in response to climate change in a wild bird population.
612   *Science*, *320*(5877), 800–803. https://doi.org/10.1126/science.1157174

613 de Kogel, C. H. (1997). Long-Term Effects of Brood Size Manipulation on Morphological
614   Development and Sex-Specific Mortality of Offspring. *The Journal of Animal Ecology*, *66*(2),
615   167. https://doi.org/10.2307/6019

616 Dettmer, A. M., & Chusyd, D. E. (2023). Early life adversities and lifelong health outcomes: A
617   review of the literature on large, social, long-lived nonhuman mammals. *Neuroscience and*
618   *Biobehavioral Reviews*, *152*(105297), 1–29.
619   https://doi.org/10.1016/j.neubiorev.2023.105297

620 Douhard, M., Plard, F., Gaillard, J., Capron, G., Delorme, D., Duncan, P., Loe, L. E., Bonenfant, C., &
621   Lyon, D. (2014). Fitness consequences of environmental conditions at different life stages
622   in a long-lived vertebrate. *Proceedings of the Royal Society B: Biological Sciences*,
623   *281*(20140276), 1–8. https://doi.org/http://dx.doi.org/10.1098/rspb.2014.0276

624 Drake, E. D., Ravindran, S., Bal, X., Pemberton, J. M., Pilkington, J. G., Nussey, D. H., Froy, H., &
625   Drake, E. D. (2025). Sex-specific effects of early-life adversity on adult fitness in a wild
626   mammal. *Proceedings of the Royal Society B: Biological Sciences*, *292*(20250192), 1–12.
627   https://doi.org/https://doi.org/10.1098/rspb.2025.0192

628 Draškovič, T., & Hauptman, N. (2024). Discovery of novel DNA methylation biomarker panels for
629   the diagnosis and differentiation between common adenocarcinomas and their liver
630   metastases. *Scientific Reports*, *14*(1), 1–25. https://doi.org/10.1038/s41598-024-53754-1

631 Dunn, O. J. (1961). Multiple Comparisons among Means. *Journal of the American Statistical*
632   *Association*, *56*(293), 52–64.
633   https://doi.org/http://dx.doi.org/10.1080/01621459.1961.10482090

634 Eastwood, J. R., Peters, A., Hall, M. L., Teunissen, N., Kingma, S. A., Hidalgo, N., Marie, A., Michael,
635   F., & Simon, R. (2019). Early - life telomere length predicts lifespan and lifetime
636   reproductive success in a wild bird. *Molecular Ecology*, *28*, 1127–1137.
637   https://doi.org/10.1111/mec.15002

638 Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for
639   multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048.
640   https://doi.org/10.1093/bioinformatics/btw354

641 Friedman, J., Hastie, T., & Tibshirani, R. (2010). Regularization Paths for Generalized Linear
642   Models via Coordinate Descent. *Journal of Statistical Software*, *33*(1), 1–22.
643   https://doi.org/https://doi.org/10.18637/jss.v033.i01

644 Galkin, F., Mamoshina, P., Aliper, A., de Magalhães, J. P., Gladyshev, V. N., & Zhavoronkov, A.
645   (2020). Biohorology and biomarkers of aging: Current state-of-the-art, challenges and
646   opportunities. *Ageing Research Reviews*, *60*. https://doi.org/10.1016/j.arr.2020.101050

647 Gerritsma, Y. H., Lamarque, F., Driessen-Roelfszema, M. M., & Verhulst, S. (2025). Growing up
648   with nutritional stress leads to peripheral social network positions, independent of
649   'personality.' *Animal Behaviour*, *224*, 1–11.

Gibney, E. R., & Nolan, C. M. (2010). Epigenetics and gene expression. *Heredity*, *105*, 4–13.
https://doi.org/10.1038/hdy.2010.54

Harney, E., Paterson, S., Collin, H., Chan, B. H. K., Bennett, D., & Plaistow, S. J. (2022). Pollution
induces epigenetic effects that are stably transmitted across multiple generations. *Evolution*
Letters, *6*(2), 118–135. https://doi.org/10.1002/evl3.273

Hedman, Å. K., Mendelson, M. M., Marioni, R. E., Gustafsson, S., Joehanes, R., Irvin, M. R., Zhi, D.,
Sandling, J. K., Yao, C., Liu, C., Liang, L., Huan, T., McRae, A. F., Demissie, S., Shah, S., Starr, J.
M., Cupples, L. A., Deloukas, P., Spector, T. D., … Ingelsson, E. (2017). Epigenetic Patterns in
Blood Associated with Lipid Traits Predict Incident Coronary Heart Disease Events and Are
Enriched for Results from Genome-Wide Association Studies. *Circulation: Cardiovascular*
Genetics, *10*(1), 1–13. https://doi.org/10.1161/CIRCGENETICS.116.001487

Höglund, A., Henriksen, R., Fogelholm, J., Churcher, A. M., Guerrero-bosagna, C. M., Martinez-
barrio, A., Johnsson, M., Jensen, P., & Wright, D. (2020). The methylation landscape and its
role in domestication and gene regulation in the chicken. *Nature Ecology and Evolution*, *4*,
1713–1724. https://doi.org/10.1038/s41559-020-01310-1

Holveck, M. J., & Riebel, K. (2010). Low-quality females prefer low-quality males when choosing
a mate. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1678), 153–160.
https://doi.org/10.1098/rspb.2009.1222

Horvath, S. (2013). DNA methylation age of human tissues and cell types. *Genome Biology*, *16*(1).
https://doi.org/10.1186/s13059-015-0649-6

Hu, J., Askary, A. M., Thurman, T. J., Spiller, D. A., Palmer, T. M., Pringle, R. M., Barrett, R. D. H., &
Pal, C. (2019). The Epigenetic Signature of Colonizing New Environments in Anolis Lizards.
Molecular Biology and Evolution, *36*(10), 2165–2170.
https://doi.org/10.1093/molbev/msz133

Jimeno, B., Hau, M., Gómez-Díaz, E., & Verhulst, S. (2019). Developmental conditions modulate
DNA methylation at the glucocorticoid receptor gene with cascading effects on expression
and corticosterone levels in zebra finches. *Scientific Reports*, *9*(1), 1–11.
https://doi.org/10.1038/s41598-019-52203-8

Keller, M., Yaskolka Meir, A., Bernhart, S. H., Gepner, Y., Shelef, I., Schwarzfuchs, D., Tsaban, G.,
Zelicha, H., Hopp, L., Müller, L., Rohde, K., Böttcher, Y., Stadler, P. F., Stumvoll, M., Blüher, M.,
Kovacs, P., & Shai, I. (2020). DNA methylation signature in blood mirrors successful weight-
loss during lifestyle interventions: the CENTRAL trial. *Genome Medicine*, *12*(1), 1–18.
https://doi.org/10.1186/s13073-020-00794-7

Kim, S. (2015). ppcor: An R Package for a Fast Calculation to Semi-partial Correlation
Coefficients. *Communications for Statistical Applications and Methods*, *22*(6), 665–674.
https://doi.org/10.5351/CSAM.2015.22.6.665.ppcor

Kronholm, I., & Collins, S. (2016). Epigenetic mutations can both help and hinder adaptive
evolution. *Molecular Ecology*, *25*(8), 1856–1868. https://doi.org/10.1111/mec.13296

Krueger, F., & Andrews, S. R. (2011). Bismark: A flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics*, *27*(11), 1571–1572.
https://doi.org/10.1093/bioinformatics/btr167

Krueger, F., James, F., Ewels, P., Afyounian, E., Weinstein, M., Schuster-Boeckler, B., Hulselmans,
G., & sclamons. (2023). *FelixKrueger/TrimGalore: v0.6.10*. Zenodo.
https://doi.org/10.5281/zenodo.7598955

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*
Methods, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Laubach, Z. M., Greenberg, J. R., Turner, J. W., Montgomery, T. M., Pioon, M. O., Sawdy, M. A.,
Smale, L., Cavalcante, R. G., Padmanabhan, K. R., Lalancette, C., vonHoldt, B., Faulk, C. D.,
Dolinoy, D. C., Holekamp, K. E., & Perng, W. (2021). Early-life social experience affects
offspring DNA methylation and later life stress phenotype. *Nature Communications*, *12*(1),
1–15. https://doi.org/10.1038/s41467-021-24583-x

Lea, A. J., Altmann, J., Alberts, S. C., & Tung, J. (2016). Resource base influences genome-wide DNA
methylation levels in wild baboons (Papio cynocephalus). *Molecular Ecology*, *25*(8), 1681–
1696. https://doi.org/10.1111/mec.13436

Li, D., Yang, Y., Li, Y., Zhu, X., & Li, Z. (2021). Epigenetic regulation of gene expression in response
to environmental exposures: From bench to model. *Science of the Total Environment*,
776(1), 1–12. https://doi.org/10.1016/j.scitotenv.2021.145998

Li, S., & Tollefsbol, T. O. (2021). DNA methylation methods: global DNA methylation and
methylomic analyses. Shizhao. *Methods*, *187*, 28–43.
https://doi.org/10.1016/j.ymeth.2020.10.002.DNA

Lind, M. I., & Spagopoulou, F. (2018). Evolutionary consequences of epigenetic inheritance.
Heredity, *121*, 205–209. https://doi.org/10.1038/s41437-018-0113-y

Liu, Z., Leung, D., Thrush, K., Zhao, W., Ratliff, S., Tanaka, T., Schmitz, L. L., Smith, J. A., Ferrucci, L.,
& Levine, M. E. (2020). Underlying features of epigenetic aging clocks in vivo and in vitro.
Aging Cell, *19*(10), 1–11. https://doi.org/10.1111/acel.13229

Lu, C. Y., Hsiao, C. Y., Peng, P. J., Huang, S. C., Chuang, M. R., Su, H. J., & Huang, K. W. (2023). DNA
Methylation Biomarkers as Prediction Tools for Therapeutic Response and Prognosis in
Intermediate-Stage Hepatocellular Carcinoma. *Cancers*, *15*(18), 1–16.
https://doi.org/10.3390/cancers15184465

McCullagh, P., & Nelder, J. A. (1989). *Generalized Linear Models*. Chapman and Hall.

Meyer, B. S., Moiron, M., Caswara, C., Chow, W., Fedrigo, O., Formenti, G., Haase, B., Howe, K.,
Mountcastle, J., Uliano-Silva, M., Wood, J., Jarvis, E. D., Liedvogel, M., & Bouwhuis, S. (2023).
Sex-specific changes in autosomal methylation rate in ageing common terns. *Frontiers in
Ecology and Evolution*, *11*(January), 1–9. https://doi.org/10.3389/fevo.2023.982443

Mock, D. W., Schwagmeyer, P. L., & Dugas, M. B. (2009). Parental provisioning and nestling
mortality in house sparrows. *Animal Behaviour*, *78*(3), 677–684.
https://doi.org/10.1016/j.anbehav.2009.05.032

Moore, L. D., Le, T., & Fan, G. (2013). DNA methylation and its basic function.
Neuropsychopharmacology, *38*(1), 23–38. https://doi.org/10.1038/npp.2012.112

Morselli, M., Farrell, C., Rubbi, L., Fehling, H. L., & Pellegrini, M. (2021). Targeted Bisulfite
Sequencing for Biomarker Discovery. *Methods*, *March*(187), 13–27.
https://doi.org/10.1016/j.ymeth.2020.07.006.

Nilsson, J. Å., & Svensson, M. (2001). Sibling competition affects individual growth strategies in
marsh tit , Parus palustris , nestlings. *Animal Behaviour*, *61*, 357–365.
https://doi.org/10.1006/anbe.2000.1602

Petkovich, D. A., Podolskiy, D. I., Lobanov, A. V, Lee, S., Miller, R. A., Gladyshev, V. N., Petkovich, D.
A., Podolskiy, D. I., Lobanov, A. V, Lee, S., & Miller, R. A. (2017). Using DNA Methylation
Profiling to Evaluate Biological Age and Longevity Interventions Short Article Using DNA
Methylation Profiling to Evaluate Biological Age and Longevity Interventions. *Cell
Metabolism*, *25*(4), 954–960. https://doi.org/10.1016/j.cmet.2017.03.016

R Core Team. (2023). *R: A language and environment for statistical computing.* (RStudio

741   2021.09.0+351 "Ghost Orchid" Release). https://www.r-project.org/

742 Réale, D., McAdam, A. G., Boutin, S., & Berteaux, D. (2003). Genetic and plastic responses of a
743   northern mammal to climate change. *Proceedings of the Royal Society B: Biological Sciences*,
744   *270*(1515), 591–596. https://doi.org/10.1098/rspb.2002.2224

745 Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W.,
746   Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-
747   Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D. (2021). Towards complete and
748   error-free genome assemblies of all vertebrate species. *Nature*, *592*(7856), 737–746.
749   https://doi.org/10.1038/s41586-021-03451-0

750 Schwagmeyer, P., & Mock, D. (2008). Parental provisioning and offspring fitness⬚: size matters.
751   *Animal Behaviour*, *75*, 291–298. https://doi.org/10.1016/j.anbehav.2007.05.023

752 Sepers, B., Erven, J. A. M., Gawehns, F., Laine, V. N., & van Oers, K. (2021). Epigenetics and Early
753   Life Stress: Experimental Brood Size Affects DNA Methylation in Great Tits (Parus major).
754   *Frontiers in Ecology and Evolution*, *9*(March), 1–15.
755   https://doi.org/10.3389/fevo.2021.609061

756 Sepers, B., Mateman, A. C., Gawehns, F., Verhoeven, K. J. F., & Oers, K. Van. (2023). Developmental
757   stress does not induce genome- - wide DNA methylation changes in wild great tit (Parus
758   major) nestlings. *Molecular Ecology*, *32*(March), 3960–3974.
759   https://doi.org/10.1111/mec.16973

760 Sepers, B., Verhoeven, K. J. F., & Oers, K. Van. (2024). Early developmental carry- - over effects on
761   exploratory behaviour and DNA methylation in wild great tits (Parus major). *Evolutionary*
762   *Applications*, *December 2023*, 1–14. https://doi.org/10.1111/eva.13664

763 Sheldon, E. L., Schrey, A. W., Ragsdale, A. K., & Griffith, S. C. (2018). Brood size influences patterns
764   of DNA methylation in wild Zebra Finches (Taeniopygia guttata). *Auk*, *135*(4), 1113–1122.
765   https://doi.org/10.1642/AUK-18-61.1

766 Spagopoulou, F., Teplitsky, C., Lind, M. I., Chantepie, S., Gustafsson, L., & Maklakov, A. A. (2020).
767   Silver-spoon upbringing improves early-life fitness but promotes reproductive ageing in a
768   wild bird. *Ecology Letters*, *23*(6), 994–1002. https://doi.org/10.1111/ele.13501

769 Tangili, M., Briga, M., & Verhulst, S. (2025). Begging costs rather than food received cause brood
770   size effect on growth in zebra finches. *BioRxiv*, 1–16.
771   https://doi.org/https://doi.org/10.1101/2025.01.29.635459

772 Tangili, M., Slettenhaar, A. J., Sudyka, J., Dugdale, H. L., Pen, I., Palsbøll, P. J., & Verhulst, S. (2023).
773   DNA methylation markers of age(ing) in non-model animals. *Molecular Ecology*, *32*(17),
774   4725–4741. https://doi.org/10.1111/mec.17065

775 Watson, H., Salmón, P., & Isaksson, C. (2019). Dynamic changes in DNA methylation during
776   embryonic and postnatal development of an altricial wild bird. *Ecology and Evolution*, *9*,
777   9580–9585. https://doi.org/10.1002/ece3.5480

778 Wei, B., Wu, F., Xing, W., Sun, H., Yan, C., Zhao, C., Wang, D., Chen, X., Chen, Y., Li, M., & Ma, J.
779   (2021). A panel of DNA methylation biomarkers for detection and improving diagnostic
780   efficiency of lung cancer. *Scientific Reports*, *11*(1), 1–10. https://doi.org/10.1038/s41598-
781   021-96242-6

782 Wick, W., Weller, M., Van Den Bent, M., Sanson, M., Weiler, M., Von Deimling, A., Plass, C., Hegi, M.,
783   Platten, M., & Reifenberger, G. (2014). MGMT testing - The challenges for biomarker-based
784   glioma treatment. *Nature Reviews Neurology*, *10*(7), 372–385.
785   https://doi.org/10.1038/nrneurol.2014.100

Wreczycka, K., Gosdschan, A., Yusuf, D., Grüning, B., Assenov, Y., & Akalin, A. (2017). *Strategies*
for analyzing bisulfite sequencing data. 261, 105–115.
https://doi.org/10.1016/j.jbiotec.2017.08.007

Xiao, Y., Wang, J., Siegel, P. B., Cline, M. A., & Gilbert, E. R. (2020). Early-life stress induced
epigenetic changes of corticotropin-releasing factor gene in anorexic low body weight–
selected chicks. *Life*, *10*(5). https://doi.org/10.3390/life10050051

Xu, X., Guo, Y., Liu, M., Hu, Y., & Li, S. (2024). Advancements in the clinical application of gene
methylation for early cancer detection. *Frontiers in Epigenetics and Epigenomics*, *2*(July), 1–
22. https://doi.org/10.3389/freae.2024.1430294

Yousefi, P. D., Suderman, M., Langdon, R., Whitehurst, O., Davey Smith, G., & Relton, C. L. (2022).
DNA methylation-based predictors of health: applications and statistical considerations.
Nature Reviews Genetics, *23*(6), 369–383. https://doi.org/10.1038/s41576-022-00465-w

Zann, R. A. (1996). *The zebra finch. A synthesis of field and laboratory studies.* Oxford University
Press.

Zhang, W., Lukacsovich, D., Young, J. I., Gomez, L., Schmidt, M. A., Martin, E. R., Kunkle, B. W.,
Chen, X. S., O'Shea, D. M., Galvin, J. E., & Wang, L. (2025). DNA methylation signature of a
lifestyle-based resilience index for cognitive health. *Alzheimer's Research and Therapy*,
17(88), 1–17. https://doi.org/10.1186/s13195-025-01733-7

Zheleznyakova, G. Y., Cao, H., & Schiöth, H. B. (2016). BDNF DNA methylation changes as a
biomarker of psychiatric disorders: Literature review and open access database analysis.
Behavioral and Brain Functions, *12*(1), 1–14. https://doi.org/10.1186/s12993-016-0101-4

Ethics

810 All methods and experiments detailed in this manuscript were performed under the approval of
811 the Central Committee for Animal Experiments (Centrale Commissie Dierproeven) of the
812 Netherlands, under license AVD1050020174344.

Data accessibility

Raw reads of the whole methylomes of all samples are available at the National Center for
Biotechnology Information under BioProject ID PRJNA1108628. The bioinformatics code is
available in https://datadryad.org/dataset/doi:10.5061/dryad.wm37pvmw8, the R code and
intermediate data files used in this study are available in
http://datadryad.org/share/LINK_NOT_FOR_PUBLICATION/ZxtGRgc_KmEIC-
S2ZKYyAj0cLw_CdjcNT1UmfT4XeT8 (reviewer URL).

Author's contributions

821 MT and SV designed the study. MT performed the analyses. MT and SV wrote the manuscript
822 with input from all authors. All authors approved the final version of the manuscript.

Competing interests

We have no competing interests.

Funding

Contributions by PJP were supported by the European Union's Horizon 2020 Research and
Innovation Programme under the Marie Skłodowska-Curie grant agreement no. 813383, and the
University of Groningen.

Figures

Figure 1. Volcano plot of Bonferroni-corrected p-values as a function of % DNAm differences
between individuals raised in small and large broods for (a) the analysis performed per sample
(N=1,114,704) and (b) for individuals merged together (N=2,085,308). Each point represents a CpG
site, with colored points indicating statistically significant differences in % DNAm between
individuals raised in small and large broods. The dashed horizontal lines mark the genome wide
significance thresholds.
838
839

841 *Figure 2. Volcano plot of Bonferroni-corrected p-values as a function of % DNAm differences*
842 *between individuals raised in small and large broods for (a) the analysis performed only for the*
843 *young cohort (N= 60,046) and (b) only for the old cohort (N=82,273). Each point represents a CpG*
844 *site. The dashed horizontal lines mark the genome wide significance thresholds. For information*
845 *about the age distribution of samples in each age cohort see Fig.S1.*

Figure 3. Volcano plot of Bonferroni-corrected p-values as a function of % DNAm differences between individuals raised in small and large broods for (a) the analysis performed only for females (N= 1,896,932) and (b) only for males (N=1,787,898). Each point represents a CpG site, with colored points indicating statistically significant differences in % DNAm between individuals raised in small and large broods.

Figure 4. (a) Within individual changes in DMSi (z-scores, r = 0.99, p < 0.001). The dashed line indicates equal values (y=x), so no within-individual change in DMSi with age. Levels of DMSi (z-scores) in zebra finches raised in small and large broods divided by sex in the merged samples (b), and divided by age cohort in females (c) and males (d).

Figure 5. Predicted brood size (small brood: -0.5, large brood: 0.5) from the DNA methylation of 455 CpG sites selected by the final model trained on all samples using the optimized hyperparameters α and λ. The model has a predictive accuracy of 89% (89/100 samples correctly assigned to their brood size). The brood size of the points above the dashed line on the left and under the dashed line on the right was wrongly assigned.

Figure 6. Correlation between DMSi and predicted brood size values (r = -0.73, p < 0.001). For

individuals raised in small broods r=-0.05 (p < 0.001) and for individuals raised in large broods r=-

0.56 (p < 0.001). Colors denote the brood size of rearing. Solid line represents the best-fit regression

through the data and the colored area is 95% CI.

Figure 7. Correlations of DMSi (a) and SMLmi (b) with mass gain per day (gr) from day five to day
15 (N=42). Solid line represents the best-fit regression through the data and the colored area is
95% CI. Growth rate data were corrected for age at measurement using the estimates in Tables S2
and S3.

Figure 8. Short and long-term effects of early-life environmental change on fitness through
methylation. Solid lines show the net fitness effects of methylation changes caused either by
environmental change (green star and vertical arrow) or the onset of adulthood (vertical dotted
line). Black solid lines represent the presence and grey solid lines the absence of methylation effects
on fitness. For simplicity, the effects of methylation change are here considered either fully
permanent or fully transient and the net fitness changes are only due to changes in methylation.

Tables

Table 1. The DMSi (a) and the SMLmi (b) in relation to brood size and sex. Variables were mean
centered with females, small broods and young individuals coded as -0.5 and males, large broods
and old individuals coded as +0.5. Bird identity was included as a random effect.

(a)	DMSi			
Predictors	*Estimates (±SE)*	*df*	*t-value*	*Pr(>|t|)*
Intercept	-0.004(±0.1)	46	-0.33	0.97
brood size	-1.31(±0.21)	46	-6.12	<0.001
age cohort	-0.005(±0.02)	46	-0.24	0.81
sex	0.28(±0.21)	46	1.29	0.21
brood size* age cohort	-0.02(±0.04)	46	-0.36	0.72
brood size* sex	0.1(±0.43)	46	0.23	0.82
age cohort * sex	-0.03(±0.04)	46	-0.63	0.53
brood size* age cohort *sex	0.001(±0.09)	46	0.01	0.99
Random Effects				
BirdID	0.56			
residual	0.12			
(b)	SMLmi			
Predictors	*Estimates (±SE)*	*df*	*t-value*	*Pr(>|t|)*
Intercept	6.13e-04 (±0.019)	47	0.03	0.98
brood size	0.295 (±0.038)	47	7.72	<0.001
age cohort	0.017 (±0.018)	47	0.94	0.35
sex	-0.032 (±0.038)	47	-0.83	0.41
brood size* age cohort	0.011 (±0.037)	47	0.32	0.76
sex * age cohort	1.34e-03 (±0.037)	47	0.04	0.97
Random Effects				
BirdID	0.01			
residual	0.008			

908 **Supplementary Information**
909

910 Inferences from epigenetic information in an ecological context: a case study of early-life
911 environmental effects on DNA methylation in zebra finches

912
913
914 Marianthi Tangili, Per J. Palsbøll, Simon Verhulst
915
916

Figure S1. Age at sampling (years) distribution of samples included in the analyses. Color denotes relative time of sampling (young=first sample from an individual, old= second sample of an individual).

Fig.S2. Distribution of CpG-site specific absolute methylation differences between zebra finches
raised in small and large broods for using individual samples (N=1,114,704) and for individuals
merged together (N=2,085,308). Violin plot widths represent the data density and boxplots show
the median and quartiles. The y-axis values were pseudo-log transformed.

Figure S3. Percent DNA methylation difference between individuals raised in small vs. large broods
in old against young individuals. Each data point represents one CpG site. Solid line represents the
best-fit regression through the data.

Figure S4. Density plot of percent methylation difference between individuals raised in small vs.
large broods in young and old birds.

Figure S5. Distribution of CpG-site specific absolute methylation differences between zebra finches
raised in small and large broods for the young (N= 60,046) and for the old cohort (N= 82,273).
Violin plot widths represent the data density and boxplots show the median and quartiles. The y-
axis values were pseudo-log transformed.

Figure S6. Relationship between CpG methylation differences (N = 1,108,263) observed in male and female zebra finches raised in small and large broods. Solid line represents the best-fit regression through the data.

955
956
957

Figure S7. Distribution of CpG-site specific absolute methylation differences between zebra finches raised in small and large broods for females (N= 1,896,932) and only for males (N=1,787,898). Violin plot widths represent the data density and boxplots show the median and quartiles. The y-axis values were pseudo-log transformed.

962
963
964
965

979   *Table S1. Sample information and predicted brood size. Small brood size is coded as -0.5 and large*
980   *brood size as +0.5. Accuracy denotes whether or not the sample was assigned to the correct brood*
981   *size. Predictions were considered correct if they matched the direction of the actual brood size*
982   *deviation (i.e., both positive or both negative).*

Sample	BirdID	Brood size	Predicted brood size	Accuracy
1	1810	-0.5	0.190	Incorrect
2	1825	-0.5	-0.030	Correct
3	1904	0.5	-0.106	Incorrect
4	1907	-0.5	-0.046	Correct
5	1920	0.5	0.068	Correct
6	1944	-0.5	-0.114	Correct
7	1953	-0.5	-0.085	Correct
8	1983	0.5	0.103	Correct
9	2045	-0.5	0.003	Incorrect
10	2053	-0.5	-0.059	Correct
11	2058	-0.5	-0.078	Correct
12	2069	0.5	0.010	Correct
13	2087	-0.5	-0.078	Correct
14	2108	0.5	-0.030	Incorrect
15	2124	-0.5	-0.060	Correct
16	2142	-0.5	-0.086	Correct
17	2153	0.5	-0.093	Incorrect
18	2155	-0.5	-0.190	Correct
19	2163	0.5	0.141	Correct
20	2175	-0.5	-0.055	Correct
21	2229	0.5	0.086	Correct
22	2260	0.5	0.125	Correct
23	2291	0.5	0.084	Correct
24	2236	0.5	0.195	Correct
25	2259	0.5	0.086	Correct
26	1983	0.5	0.028	Correct
27	2429	0.5	0.111	Correct
28	2473	-0.5	-0.342	Correct
29	1923	0.5	0.115	Correct
30	1942	0.5	0.001	Correct
31	2058	-0.5	-0.175	Correct
32	2269	-0.5	-0.078	Correct
33	2087	-0.5	-0.179	Correct
34	2172	-0.5	-0.079	Correct
35	2288	-0.5	-0.028	Correct
36	2423	0.5	0.268	Correct
37	2487	-0.5	-0.024	Correct
38	2045	-0.5	-0.108	Correct
39	2257	-0.5	-0.049	Correct

40	2277	0.5	-0.060	Incorrect
41	2291	0.5	0.050	Correct
42	2418	0.5	0.065	Correct
43	2390	-0.5	-0.055	Correct
44	2511	-0.5	-0.205	Correct
45	1953	-0.5	-0.019	Correct
46	2163	0.5	0.092	Correct
47	2247	0.5	0.141	Correct
48	2260	0.5	0.200	Correct
49	2463	0.5	-0.028	Incorrect
50	1920	0.5	0.097	Correct
51	2155	-0.5	-0.043	Correct
52	1810	-0.5	0.148	Incorrect
53	1825	-0.5	-0.060	Correct
54	2053	-0.5	0.089	Incorrect
55	2069	0.5	0.079	Correct
56	2153	0.5	-0.116	Incorrect
57	2508	0.5	0.024	Correct
58	2508	0.5	0.102	Correct
59	2116	0.5	-0.003	Incorrect
60	2259	0.5	0.147	Correct
61	2423	0.5	0.293	Correct
62	2257	-0.5	-0.023	Correct
63	1897	0.5	0.091	Correct
64	2463	0.5	-0.035	Incorrect
65	2038	0.5	-0.046	Incorrect
66	2229	0.5	0.044	Correct
67	2277	0.5	0.055	Correct
68	2354	-0.5	-0.127	Correct
69	2124	-0.5	-0.109	Correct
70	1907	-0.5	0.012	Incorrect
71	2511	-0.5	-0.296	Correct
72	2429	0.5	0.280	Correct
73	2473	-0.5	-0.081	Correct
74	2035	0.5	0.069	Correct
75	1944	-0.5	0.071	Incorrect
76	1897	0.5	0.084	Correct
77	2108	0.5	0.014	Correct
78	1923	0.5	0.166	Correct
79	2172	-0.5	-0.057	Correct
80	2415	-0.5	-0.141	Correct
81	2392	-0.5	-0.195	Correct
82	2038	0.5	0.144	Correct
83	1904	0.5	-0.096	Incorrect
84	1942	0.5	0.094	Correct
85	2470	-0.5	-0.120	Correct

86	2142	-0.5	-0.044	Correct
87	2487	-0.5	0.034	Incorrect
88	2035	0.5	0.110	Correct
89	2116	0.5	0.129	Correct
90	2175	-0.5	-0.148	Correct
91	2470	-0.5	-0.235	Correct
92	2247	0.5	0.076	Correct
93	2415	-0.5	0.047	Incorrect
94	2269	-0.5	-0.045	Correct
95	2390	-0.5	-0.184	Correct
96	2288	-0.5	-0.042	Correct
97	2236	0.5	0.192	Correct
98	2354	-0.5	-0.138	Correct
99	2418	0.5	0.097	Correct
100	2392	-0.5	-0.263	Correct

Figure S8. (a) CpG sites in the final brood size prediction model (SMLmi) against total CpG sites per chromosome. The dotted line represents the expected number of CpG sites per chromosome if the CpGs included in the final brood size prediction model were distributed proportionally to the total number of CpG sites in each chromosome. (b) CpG sites in the final brood size prediction model (SMLmi) per chromosome.

994   *Figure S9. Shared CpG sites between DMSs in the analysis per sample (light blue), in the analysis*
995   *only for females (pink), in the analysis only for males (blue), and the CpG sites used to predict brood*
996   *size (i.e. included in the SMLmi, yellow).*

Figure S10. Density plots of methylation differences between individuals raised in small and large broods. Light blue plot: DMSs identified in the analysis per sample (N=27). Pink plot: DMSs identified in the analysis only for females (N=34). Blue plot: DMSs identified in the analysis only for males (N=38). Yellow plot: CpGs selected for the SMLmi (N=455).

998
999
1000
1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

 Table S2. Results of linear model exploring the effects of DMSi and age at measurement on mass
 gain per day.

	Mass gain per day (gr)		
Predictor	*Estimate (±SE)*	*t-value*	*Pr(>|t|)*
Intercept	1.4 (±0.16)	8.89	<0.001
DMSi	0.07 (±0.02)	3.54	0.001
age at first mass measurement	0.04 (±0.01)	2.82	0.007
age at day 15 measurement	-0.06 (±0.01)	-5.166	<0.001

 Table S3. Results of linear model exploring the effects of SMLmi and age at measurement on mass
 gain per day.

	Mass gain per day (gr)		
Predictor	*Estimate (±SE)*	*t-value*	*Pr(>|t|)*
Intercept	1.43 (±0.16)	8.82	<0.001
SMLmi	-0.34 (±0.11)	-3.09	0.004
age at first mass measurement	0.04 (±0.02)	2.63	0.01
age at day 15 measurement	-0.06 (±0.01)	-5.11	<0.001

1023
1024