Elevational Shifts in Tropical Tree Leaf Traits: Interactions Between Soil, Climate, Light, and Phylogeny

Jiří Doležal[1,2,] Kirill Korznikov[1], Vaclav Bazant[3], Adam Taylor Ruka[1,2], Jorge Gago[4]

1 Institute of Botany, The Czech Academy of Sciences, Dukelská 135, Třeboň, 379 01, Czech Republic.
2 Department of Botany, Faculty of Science, University of South Bohemia, Branišovská 1760, 370 05, České Budějovice, Czech Republic.
3 Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Kamýcká 129, CZ-165 21, Prague 6 - Suchdol, Czech Republic
4 Research Group on Plant Biology under Mediterranean Conditions, Universitat de les Illes Balears (UIB) – Instituto de Investigaciones Agroambientales y de Economía del Agua (INAGEA), Palma, Illes Balears, Spain

ORCID:
Jiri Dolezal (0000-0002-5829-4051)
Adam Taylor Ruka (0000-0001-9255-5086)
Kirill Korznikov (0000-0003-2850-1483)
Jorge Gago (0000-0002-4293-0105)

Corresponding author: Jiri Dolezal, jiriddolezal@gmail.com, +420774545707
Institute of Botany, The Czech Academy of Sciences, Dukelská 135, Třeboň, 379 01, Czech Republic.

Abstract

Understanding how tropical trees respond to complex environmental gradients is essential for predicting forest resilience under climate change. We examined variation in key leaf traits—specific leaf area (SLA), foliar nitrogen (N) and phosphorus (P), C:N and N:P ratios, and stable isotope composition ($\delta^{13}C$, $\delta^{15}N$)—in 160 tree species along a 3,200-m elevational transect on Mount Cameroon. This gradient spans hyper-humid coastal rainforests to arid Afroalpine savannas, capturing sharp transitions in climate, soils, and forest structure. Leaf traits shifted nonlinearly with elevation, from acquisitive strategies at mid-elevations to conservative syndromes in lowlands and highlands. Mid-elevation forests (~1,000–1,500 m), characterized by moderate climate and canopy disturbance by elephants, supported nutrient-rich, high-SLA foliage. In contrast, high-elevation forests (>2,000 m) exhibited low SLA, high C:N, and enriched $\delta^{13}C$, consistent with stress tolerance under cold, dry, and fire-prone conditions. The strongest phosphorus limitation occurred in hyper-humid lowlands, where extreme rainfall (>12,000 mm/year) drives leaching losses. Foliar $\delta^{15}N$ declined markedly with elevation (from +5‰ to −5‰), indicating a shift from mineral N uptake and N-fixation in lowland Fabaceae to ecto- and ericoid mycorrhizal associations in montane Ericaceae. A bimodal $\delta^{15}N$ pattern—enrichment in both lowland and upper montane forests—reflects N-fixation under leaching and fire-driven N scarcity, respectively. Phylogenetic analyses showed that climate, soils, forest structure, and lineage jointly shaped trait–environment relationships. Traits related to $\delta^{13}C$, C:N, and $\delta^{15}N$ exhibited strong phylogenetic signal, highlighting evolutionary constraints. These findings underscore the value of integrating functional traits, isotopes, and phylogeny to predict tropical forest responses to global change.

Keywords: Leaf functional traits, Stable isotopes (δ^{13}C, δ^{1}N), Tropical montane forests, Elevational gradient, Nitrogen cycling, Mycorrhizal associations, Phylogenetic trait conservatism, Plant ecophysiology, Trait–environment relationships, Climate change adaptation

1. Introduction

Understanding how plant functional traits vary across environmental gradients is central to ecology, biogeography, and global change biology (Wright et al., 2004; Reich, 2014; Garnier et al., 2015). Leaf traits such as specific leaf area (SLA), leaf nitrogen (N) and phosphorus (P) content, and leaf N:P ratios are widely used as proxies for resource acquisition strategies, photosynthetic capacity, and growth responses (Wright et al., 2004; Reich, 2014). These traits are shaped by multiple environmental filters—including nutrient availability, moisture regimes, temperature, and light conditions—that often covary along elevation gradients (Körner, 2021). However, because these drivers are interdependent, it can be challenging to isolate their respective contributions to trait variation, which limits the ecological interpretation and predictive utility of trait-based approaches (Fyllas et al., 2017; Oliveras et al., 2020). Despite the rise of global trait databases, tropical mountain systems—particularly those in Africa—remain underrepresented in trait-based ecology, despite sustaining high biodiversity and regulating major global carbon and water fluxes.

Elevational gradients provide natural laboratories for examining trait–environment relationships, encompassing steep and relatively continuous transitions in climate, edaphic conditions, and forest structure over short geographic distances (Körner, 2007; Girardin et al., 2014). While such gradients have been widely used in temperate and boreal regions, studies spanning tropical rainforest to high-montane savanna remain scarce. The elevational transect on Mount Cameroon—a geologically young, exceptionally steep volcano rising from sea level to over 4,000 m—is one of the most dramatic tropical elevation gradients globally (Hořák et al., 2019; Maicher et al., 2020; Doležal et al., 2023). Yet, functional trait data from this region are virtually absent from global syntheses. This setting presents a rare opportunity to investigate how plant strategies transition from humid lowland rainforest to montane and Afroalpine ecosystems, thereby filling a critical empirical gap in trait-based biogeography and ecosystem modeling.

In tropical forest systems, lowland forests are generally characterized by high temperatures, relatively stable moisture conditions, and high radiation, while montane forests experience cooler temperatures, frequent cloud immersion, and often lower light availability (Kitayama & Aiba, 2002; Girardin et al., 2016). Soil fertility may decline or shift in nutrient composition with elevation, depending on parent material, erosion, and leaching processes (Vitousek & Sanford, 1986; Soethe et al., 2008). Leaf trait variation along these gradients— such as declining SLA or leaf N content with elevation—is often interpreted as an adaptive response to colder or nutrient-limited conditions. Yet such patterns may also reflect light limitation under dense canopies or nutrient reallocation in response to drought or other environmental stressors.

Different tropical forests are constrained by distinct environmental factors, and these constraints are reflected in characteristic leaf trait syndromes (Pringle et al., 2011; Méndez-Alonzo et al., 2012). In seasonally dry tropical forests or savanna ecotones, where water availability is the principal limiting factor, plants tend to exhibit conservative leaf traits—low SLA, and low leaf N and P concentrations—associated with drought tolerance and slow resource turnover (Markesteijn & Poorter, 2009; Wright et al., 2001). By contrast, in moist lowland rainforests where light is the main limiting resource beneath a closed canopy, species often display higher SLA and elevated leaf nutrient content, reflecting acquisitive strategies optimized for high light capture and fast growth in gaps or upper canopy strata (Díaz et al.,

2016; Asner et al., 2018). In tropical montane or cloud forests, low temperatures and
persistent cloud cover can limit both photosynthesis and transpiration, leading to thick, tough
leaves (low SLA, high LDMC), relatively low nutrient concentrations, and in some cases,
decoupling of leaf traits from typical stoichiometric expectations due to constrained nutrient
uptake or translocation (Vitousek et al., 1992; Moser et al., 2007). In nutrient-poor forests,
such as those developed on old or highly weathered tropical soils, species often exhibit low
leaf nutrient concentrations and high N:P ratios, indicating phosphorus limitation (Townsend
et al., 2007; Porder & Hilley, 2011).

Leaf traits can thus serve as ecological indicators of the dominant resource constraints
in a given habitat. High SLA and high leaf N are commonly associated with nutrient-rich and
high-light environments, whereas low SLA, high LDMC, and low leaf nutrient content are
indicative of stress tolerance under drought, cold, or nutrient limitation (Wright et al., 2004;
Laughlin, 2014). Leaf N:P ratios provide further insights into the type of nutrient limitation:
low N:P (<10–14) suggests nitrogen limitation, high N:P (>16–20) indicates phosphorus
limitation, and intermediate values suggest co-limitation or balanced nutrition (Koerselman &
Meuleman, 1996; Güsewell, 2004). These trait–environment relationships are increasingly
being used to infer ecosystem-level nutrient cycling, plant resource strategies, and responses
to environmental change.

Stable isotope ratios of carbon ($\delta^{13}C$) and nitrogen ($\delta^{15}N$) in leaves provide an
additional layer of inference, reflecting the integration of physiological and ecological
processes (Givnish, 2002; Santiago et al., 2017). Leaf $\delta^{13}C$ is widely used as a proxy for
intrinsic water-use efficiency (WUEi), as it reflects the ratio of carbon assimilation to
stomatal conductance over time (Farquhar et al., 1989). Higher $\delta^{13}C$ values indicate greater
WUE and are often associated with plants experiencing drought stress or growing in arid
microclimates (Cernusak et al., 2007; Cornwell et al., 2018). In elevational gradients, $\delta^{13}C$
typically increases (becomes less negative) with altitude by about 1.1‰ per kilometer on
average, due to reduced atmospheric pressure and CO_2 partial pressure (Körner et al., 1991).
However, it can also vary with canopy position and light exposure, as well as increasing
evaporative demand (Holtum & Winter, 2005; Doležal et al., 2021). As leaves become thicker
and denser with elevation—reflected in decreasing SLA—a corresponding decline in leaf
nutrient content and increased $\delta^{13}C$ may suggest a trade-off between photosynthetic nitrogen
efficiency and water-use efficiency (Wright et al., 2001; Cornwell et al., 2018). Thicker
leaves, while more durable and better protected against abiotic stressors, often limit CO_2
diffusion (mesophyll conductance), resulting in lower photosynthetic capacity per unit N and
higher $\delta^{13}C$ values. Thus, a negative correlation between leaf N and $\delta^{13}C$ across the gradient
could be expected, and may reflect this fundamental trade-off in resource allocation in
response to increasingly harsh abiotic conditions and reduced competitive pressure at the
treeline.

In contrast, $\delta^{15}N$ values may exhibit a complex pattern, diverging from $\delta^{13}C$ trends and
necessitating a more nuanced interpretation. $\delta^{15}N$ is influenced by N cycling processes,
symbiotic associations, and rooting depth (Craine et al., 2015; Yang et al., 2015). Mycorrhizal
associations, especially with ectomycorrhizal or ericoid fungi, tend to result in lower $\delta^{15}N$
values due to fractionation during uptake (Högberg, 1997; Mayor et al., 2014). Conversely,
Fabaceae often show higher or near-zero values, depending on their N source (e.g., symbiotic
fixation, nitrate vs. ammonium uptake). $\delta^{15}N$ values may thus reflect both ecological
processes (e.g., nutrient access, waterlogging, or periodic anoxia at low elevation) and
phylogenetic filtering along the gradient, underscoring the need for careful disentangling of
abiotic drivers from lineage-specific trait syndromes (Cornwell & Ackerly, 2009; Laughlin &
Messier, 2015).

In this study, we investigate elevational trends in key leaf traits across 160 tropical tree
species sampled along a nearly 3,200 m elevational gradient on Mount Cameroon, spanning
from sea-level rainforests to upper montane and Afroalpine vegetation. For each species, we
measured SLA, leaf N and P content, N:P ratios, and stable isotope composition ($\delta^{13}C$ and
$\delta^1\square N$), alongside structural traits (tree height, stem diameter) and detailed environmental
data, including soil chemistry, microclimate (temperature, humidity), and light availability
(from hemispherical photography). We hypothesized that trait variation across the gradient
reflects a combination of environmental filtering and evolutionary conservatism, with
acquisitive strategies (e.g., high SLA, high leaf N and P, low $\delta^{13}C$) favored in warm, fertile
lowlands, and conservative traits (e.g., low SLA, high C:N, elevated $\delta^{13}C$) emerging in colder,
drier, and nutrient-limited high elevations. These predictions are grounded in the leaf
economic spectrum (LES) framework, which posits that plant traits are structured by trade-
offs between resource acquisition and conservation. However, we also expected deviations
from LES predictions in areas where fire, hydrology, forest elephant disturbances, or nutrient
decoupling disrupt classical trait–soil patterns. By testing these hypotheses, we aim to identify
the dominant environmental constraints underlying observed trait variation and to advance
trait-based ecological frameworks for Afrotropical forests—one of the world's most under-
sampled and climatically dynamic ecosystems.
167
168

169
170 **Figure 1.** *A study area on the southwestern slopes of Mount Cameroon, West Africa. (a) The*
171 *map shows forest cover (%) with red transects indicating sampling locations. (b) Photos on*
172 *the right illustrate vegetation types and leaf morphology along the gradient.*
173

2.1. Study Area
This study was conducted along a continuous elevational gradient on the southwestern slopes
of Mount Cameroon, located in the Republic of Cameroon (Figure 1). Data were collected
from 176 permanent monitoring plots spanning a near-pristine rainforest-to-alpine gradient,
ranging from coastal lowland forests to the treeline at approximately 3200 m a.s.l.—a unique
ecological continuum in West Africa, situated within Mount Cameroon National Park
(Maicher et al. 2020). The plots encompassed all major vegetation zones, including forested
areas from sea level to montane cloud forests at 2400 m, as well as isolated tree stands and
solitary trees extending into the afromontane savanna up to 3200 m (Dolezal et al., 2022). At
each elevation level, 16 plots were systematically established approximately 150 meters apart

along the contour to capture variability in vegetation structure, canopy light conditions, leaf functional traits, and soil chemistry. Because the southwestern transect of the mountain is bordered by agricultural land and coastal settlements below 300 m a.s.l., the lowland rainforest was instead sampled within the nearby Bimbia Forest Reserve. This protected coastal forest harbors high tree species richness, with over 100 species per hectare, and provides a representative baseline for the coastal rainforest zone (Ferenc et al., 2018; Maicher et al., 2020).

Mount Cameroon lies within a perhumid tropical climate zone influenced by seasonal shifts between the moist southwest monsoon and the dry northeast harmattan winds. The region experiences a distinct rainy season from June to September and a dry season from late December to February, with brief transitional periods in between. Annual precipitation in the coastal lowlands exceeds 11,500 mm (Rehakova et al., 2022), making this one of the wettest locations on the Planet. Rainfall decreases steadily with increasing elevation and distance from the ocean, creating a pronounced dry belt between 2300 and 3200 m. This rain shadow effect facilitates the development of a treeless afromontane savanna above the forest limit (Dolezal et al. 2022).

The flora of Mount Cameroon is exceptionally diverse, with more than 2300 vascular plant species recorded, encompassing over 800 genera and 210 families (Cheek et al. 1996; Cable and Cheek 1998). The mountain's forest vegetation is stratified into distinct zones along the elevational gradient (Hall, 1973; Payton, 1993; Proctor et al., 2007). Coastal rainforest (0–250 m a.s.l.) is dominated by tall *Desbordesia glaucescens*, along with *Ceiba pentandra*, *Coelocaryon preussii*, *Pycnanthus angolensis*, *Scyphocephalium mannii*, and *Lophira alata*. A closed canopy and sparse understory characterize this zone. Lowland rainforest (300–1000 m a.s.l.) harbors species such as Pycnanthus angolensis, Coelocaryon preussii, Crudia gabonensis, Berlinia bracteosa, Anthonotha fragrans, and Baikiaea insignis, characterized by a structurally intact canopy. Mid-elevation forest (1000–1600 m a.s.l.) forms a transitional belt marked by reduced canopy closure due to disturbance by forest elephants (*Loxodonta cyclotis*), which maintain a mosaic of open clearings and closed forest. Dominant species include *Anthonotha noldeae*, *Schefflera mannii*, *Ficus chlamydocarpa*, and *Entandrophragma utile*. Montane forest (1600–2300 m a.s.l.) occurs under persistent cloud cover and mist, with a denser understory and canopy dominated by *Ilex mitis*, *Schefflera mannii*, *S. abyssinica*, *Syzygium staudtii*, and *Prunus africana*. Afromontane vegetation (2400–2300 m a.s.l.) is dominated by C4 grasses (*Loudetia simplex*, *Pennisetum monostigma*) and forbs (*Trifolium simense*, *Satureja robusta*), with sparse stands of fire-tolerant trees such as *Myrica arborea* and *Agauria salicifolia*, limited by volcanic activity, porous substrate, and low precipitation. Although the climatic tree line would occur around 3200–3500 m based on temperature, actual tree growth ceases at much lower elevations. At the highest elevations, from approximately 3300 m to the summit at 4000 m, the Afroalpine zone is characterized by cold, windy conditions and sparse plant cover, composed of C3 grasses, herbs, and low-growing shrubs (Dolezal et al., 2022).

2.2. Leaf trait sampling

We analyzed leaf functional trait data from 412 individual trees collected across a network of 176 monitoring plots, representing 164 species, 122 genera, and 53 families (Fig. 2), along an elevational gradient on Mount Cameroon. We used a single rope technique and 3-m-long pruning shears to collect 15–20 leaves from the upper, well-lit crown regions. Fresh leaf blades were photographed in the field to assess leaf area and SLA. For each individual, we measured eight leaf traits: leaf area (LA), specific leaf area (SLA), leaf nitrogen (LNC), leaf phosphorus (LPC), leaf carbon (LCC) content, carbon-to-nitrogen ratio (C:N), nitrogen-to-phosphorus ratio (N:P), and stable isotope ratios of carbon (δ^{13}C) and nitrogen (δ^{1}N). Total

carbon and nitrogen content, along with δ¹³C and δ¹⁵N, were measured at the Stable Isotope Facility at UC Davis, USA, using an elemental analyzer-isotope ratio mass spectrometer (EA-IRMS). Phosphorus concentrations were determined via spectrophotometry following HClO₄ digestion using a SHIMADZU UV-1650PC spectrophotometer.

2.3. Vegetation Structure and Light Environment

To assess the relationship between leaf traits, forest structure, and species composition, we conducted a detailed inventory of all trees with a diameter at breast height (DBH) exceeding 10 cm in each plot. Each tree was georeferenced, identified to the species level, and measured for DBH and total height. In total, we recorded over 10,000 individual trees belonging to more than 250 species. From these measurements, we derived several structural metrics for each forest plot, including stand basal area, total volume of live and dead trees, mean and maximum stem diameter (DBH) and tree height, the coefficients of variation (CV) for DBH and height (to capture vertical stratification), and the stem slenderness index (SSI). Canopy light conditions were quantified using hemispherical photography. In each forest plot, five hemispherical images were captured at a height of 1.8 meters aboveground—one at the plot center and four at points 10 meters away in the cardinal directions—resulting in a total of 480 photographs. These images were analyzed with WinSCANOPY software to estimate canopy openness, leaf area index, and both direct and diffuse solar radiation penetration.

2.4. Climate Variables

Due to the absence of meteorological stations above 1000 m a.s.l. on Mount Cameroon, we relied on high-resolution gridded climate data from the TerraClimate database (Abatzoglou et al., 2018) to characterize the climatic conditions across our study sites. This dataset provides monthly records of multiple climate variables, including mean air temperature, total precipitation, vapor pressure deficit (VPD), and the Palmer Drought Severity Index (PDSI), at a spatial resolution of 1/24 degrees. For each monitoring plot, we extracted the corresponding grid cell data and calculated the long-term mean annual values for the period from 2000 to 2014. To ground-truth the accuracy of these gridded estimates and ensure they reflect the microclimatic realities of the elevational gradient on Mount Cameroon, we conducted extensive field-based climate monitoring. Approximately 160 temperature dataloggers (EMS Brno and TOMST Praha, Czechia) were deployed at hourly resolution along the full elevational transect. Additionally, we installed 15 manual rain gauges (EMS Brno, Czechia) at key sites to capture site-specific precipitation data. These gauges were regularly maintained to remove accumulated vegetation, leaf litter, and other debris, and were protected from damage, particularly from seasonal fires in the savanna zone. Comparative analyses revealed strong correlations between the gridded climate data and the in situ measurements, with Pearson correlation coefficients ranging from 0.7 to 0.9 (P < 0.01). This high degree of agreement validates the use of the TerraClimate dataset as a reliable representation of local climate conditions along Mount Cameroon's complex elevational gradient.

2.5. Soil characteristics

To assess soil fertility, approximately 150 g of soil was collected from each plot. Samples were oven-dried at 100 °C, manually ground using a mortar, and then sieved through a 2 mm mesh after removing coarse roots and organic debris (Rehakova et al., 2022). Key macronutrients—including calcium (Ca²⁺), magnesium (Mg²⁺), potassium (K⁺), sodium (Na⁺), nitrogen, and phosphorus—were analyzed for each sample (see Supplementary Table S1). Cation concentrations were determined using atomic absorption spectroscopy (SpectrAA 640, Varian Techtron). Nitrogen forms (ammonium, nitrate, and total nitrogen) were measured colorimetrically following Kjeldahl digestion, using a FIAstar 5010 flow injection analyzer

284 (Tecator). Phosphorus was quantified colorimetrically after perchloric acid (HClO₄)
285 digestion using a SHIMADZU UV-1650PC spectrophotometer. In addition to nutrient
286 concentrations, we assessed a suite of basic soil properties, including pH, gravimetric water
287 content, soil carbon content (SOC), and soil texture, expressed as the proportion of coarse
288 particles (diameter>0.5 mm) (Fig. 3).
289
2.6. Data analysis
We used generalized additive models (GAM) to determine changes in environmental
parameters (climate, soil, forest structure) and leaf traits along the elevational gradient studied
(Fig. 3). The GAM framework helps discover elevation-specific patterns, does not rely on any
predefined relationships, and provides easy-to-interpret visualizations (McCullagh and Nelder
1989). Because many leaf traits were intercorrelated, we conducted a principal component
analysis (PCA) to summarize the major axes of variation in these traits (Fig. 4). Leaf trait
GAMs were fitted for all individuals and separately for small-statured species (typically
subcanopy trees <15 m tall) and large-statured species (canopy and emergent trees) (Fig. 5).
To assess how environmental factors, encompassing climate, soil, light, and forest structure,
shape functional trait variation across tropical montane tree species, we quantified
phylogenetically-informed standardized effect sizes (SES) for associations between nine key
leaf traits and 30 environmental variables encompassing climate, soil, and forest structure. To
evaluate the extent to which trait–environment relationships reflect shared evolutionary
history, we quantified phylogenetic signal using standardized effect sizes (SES) derived from
a phylogenetic permutation framework. For each trait–environment pair, we compared the
observed correlation coefficient to a null distribution generated by 999 random tip-shufflings
of the phylogeny, which preserves trait and environmental values but disrupts evolutionary
structure. Significant positive SES values ($|SES| > 2$) indicate that closely related species tend
to respond similarly to a given environmental gradient, consistent with phylogenetic
clustering and trait conservatism. For example, a strong positive association between foliar
$\delta^{15}N$ and mean annual temperature (MAT) would suggest that lineages adapted to warmer
environments—such as Fabaceae or Euphorbiaceae—consistently exhibit higher $\delta^{15}N$
values. In contrast, significant negative SES values indicate weaker-than-expected trait–
environment coupling under phylogenetic constraint, suggesting that phylogenetic
overdispersion may arise from convergent evolution or ecological equivalence across
distantly related taxa. The phylogenetic distance matrix was constructed from a species-level
megaphylogeny using patristic distances. Leaf trait data (e.g., specific leaf area, nitrogen,
$\delta^{13}C$) were averaged across individuals per species. Environmental variables were extracted
from plot-level soil and light measurements or interpolated climate surfaces and averaged by
species occurrence. Analyses were conducted in Python using numpy, scipy, pandas, and
matplotlib. Phylogenetic permutations and SES computation were implemented with custom
scripts. We reconstructed the phylogeny of the studied species using MrBayes 3.1.2, based on
nucleotide sequences obtained from GenBank and those generated from field-collected
material (Plavcova *et al.*, 2024).
325

Figure 2. *Variation in leaf traits across plant families along an elevational gradient on the SW slopes of Mount Cameroon. Bars represent family means for elevation, SLA, LNC, foliar C:N ratio, stable carbon isotope ratio (δ¹³C), and nitrogen isotope ratio (δ¹⁵N). Families are ordered by mean elevation, from high-elevation taxa (e.g., Ericaceae, Primulaceae, Myricaceae) to low-elevation taxa (e.g., Combretaceae, Clusiaceae), highlighting coordinated trait shifts from stress tolerance to resource acquisitiveness.*

3. Results
3.1. Environmental and Elevational Gradients

Our study encompassed a 3,200-m elevational gradient along the southwestern flank of Mount Cameroon (Fig. 1a), capturing a broad range of forest types, microclimates, and edaphic conditions (Figs. 2–3). Elevation was strongly and negatively associated with mean annual temperature ($R^2 = 0.95$, $P < 0.001$; Fig. 3a), which declined from >25 °C in the lowlands to ~8 °C at the uppermost elevations with sparse tree cover. Soil moisture displayed a pronounced mid-elevation peak ($R^2 = 0.56$, $P < 0.001$; Fig. 3b), mirroring patterns observed for soil organic carbon ($R^2 = 0.29$, $P < 0.001$; Fig. 3c), calcium ($R^2 = 0.14$, $P < 0.001$; Fig. 3d), and nitrogen ($R^2 = 0.37$, $P < 0.001$; Fig. 3e).

Soil phosphorus exhibited a unimodal distribution ($R^2 = 0.21$, $P < 0.001$; Fig. 3f), with lowest values from low to mid elevations and a distinct peak around 2,200 m. Stoichiometric ratios varied with elevation: soil C:N declined with altitude up to the montane forest limit (~2,200 m), followed by an upswing toward the treeline (~3,200 m; $R^2 = 0.24$, $P < 0.001$; Fig. 3g). Soil N:P ratio also followed a unimodal pattern, peaking between 1,500 and 1,600 m ($R^2 = 0.41$, $P < 0.001$; Fig. 3h), indicating sharp shifts in nutrient dynamics across ecotones.

Figure 3. *Environmental variation along the elevational gradient. Relationships between (a) elevation and temperature, (b) soil moisture, and soil chemical properties, including (c) organic carbon, (d) calcium, (e) nitrogen, (f) phosphorus, (g) C:N ratio, and (h) N:P ratio. Blue lines indicate fitted GAM (generalized additive model) curves; shaded areas show 95% confidence intervals. Boxplots represent binned elevation intervals. All relationships are statistically significant (P < 0.001).*

3.2. Shifts in Leaf Functional Traits

PCA of leaf traits revealed strong trait–environment coordination and phylogenetic clustering along elevation (Fig. 4a). The first two PCA axes captured the dominant patterns across the nine traits, explaining 60.88% of the total variance (Fig. 4a). This axis reflected a gradient from acquisitive strategies—characterized by high leaf area (LA), specific leaf area (SLA), and foliar nitrogen (LNC) and phosphorus (LPC), typically found in lowland rainforest species—to conservative strategies with low SLA, low leaf N and P, and elevated C:N ratios and $\delta^{13}C$ values, common among high-elevation taxa. Trait means (standard deviation) were as follows: LA = 295.7   180.8 cm², SLA = 17.8   9.3 cm²/g, LNC = 2.38   0.79 %, LPC = 0.17   0.08 %, LCC = 45.2   2.98 %, C:N = 21.9   9.97, N:P = 15.0   5.05, $\delta^{13}C$ = –32.5   2.15 ‰, and $\delta^{1}N$ = 0.32   2.63 ‰. High-elevation families—such as Ericaceae, Stilbaceae, and Myricaceae—clustered around high $\delta^{13}C$, high C:N, and low SLA and nutrient content. In contrast, lowland families (e.g., Irvinigiaceae, Clusiaceae) exhibited higher SLA, LA, and nutrient concentrations, consistent with fast-growing, resource-acquisitive strategies. Intermediate families, such as Fabaceae and Rosaceae, occupied a broader, ecologically versatile trait space.

3.3. Trait Coordination and Stoichiometric Trade-offs Across the Elevational Gradient

Trait correlations mirrored the elevational and environmental gradients, revealing strong coordination among functional, stoichiometric, and isotopic leaf properties (Fig. 4b). Leaf carbon content (LCC) and C:N ratio were positively associated with $\delta^{13}C$ (r = 0.18 and r = 0.35, P < 0.01), consistent with increased intrinsic water-use efficiency (WUE) under drier or cooler microclimates. This supports the hypothesis that thicker or denser leaves, with higher structural carbon, may exhibit reduced stomatal conductance and thus higher $\delta^{13}C$ values under stressful conditions. Leaf nitrogen (LNC) and phosphorus content (LPC) were tightly coupled (r = 0.60, P < 0.001), reflecting co-regulation of foliar nutrients involved

in photosynthesis and protein synthesis. Both nutrients were strongly and negatively correlated with the C:N ratio (LNC vs. C:N: r = −0.88, P < 0.001; LPC vs. C:N: r = −0.55, P < 0.001), confirming that nutrient-rich leaves exhibit lower stoichiometric C:N ratios and align with acquisitive strategies.

The leaf N:P ratio was negatively associated with LPC (r = −0.56, P < 0.001) and weakly with LNC (r = 0.22, P < 0.05), suggesting that phosphorus availability exerts a stronger control on stoichiometric ratios in this system than nitrogen. $\delta^1 N$ exhibited a moderately strong positive correlation with LNC (r = 0.56, P < 0.001) and a weaker correlation with LPC (r = 0.30, P < 0.01), possibly reflecting shifts in nitrogen sources or turnover rates with foliar nutrient availability. However, $\delta^1 N$ declined with C:N (r = −0.58, P < 0.001) and elevation (r = −0.69, P < 0.001), supporting the idea that higher-altitude species exhibit more conservative nitrogen cycling or increased mycorrhizal associations that reduce foliar $\delta^1 N$ values.

SLA was positively correlated with LNC (r = 0.50, P < 0.001) and LPC (r = 0.49, P < 0.001), and negatively with elevation (r = −0.40, P < 0.001), in line with expectations from the leaf economics spectrum (LES). Similarly, LA showed a positive relationship with SLA (r = 0.42, P < 0.001) and LNC (r = 0.38, P < 0.01), suggesting that large, thin leaves dominate in nutrient-rich lowland forests, while high-elevation communities are characterized by smaller, thicker, nutrient-conservative foliage. Importantly, we observed a negative correlation between LNC and $\delta^{13}C$ across the gradient (r = −0.29, P < 0.01), consistent with a trade-off between photosynthetic nitrogen efficiency and water-use efficiency in increasingly stressful environments. This pattern reflects the inherent physiological constraint whereby nitrogen-rich, high-photosynthesis leaves often require open stomata and high transpiration rates, resulting in lower WUE and more negative $\delta^{13}C$ values.

Figure 4. (a) *PCA biplot illustrates the relationships between leaf traits and plant families, revealing trait–environment associations and compositional changes with elevation. (b) A correlation matrix showing that elevation is negatively associated with leaf area, nitrogen, phosphorus, and $\delta^1 N$, and positively associated with $\delta^{13}C$ and the C:N ratio, reflecting shifts in plant functional strategies.*

3.4. Elevational Trends in Leaf Functional Traits

Leaf traits varied nonlinearly along the 3,000-meter elevational gradient, reflecting shifts in
resource-use strategies, physiological constraints, and forest structure (Fig. 5). Specific leaf
area (SLA), leaf area (LA), and foliar nitrogen (LNC) and phosphorus (LPC) all showed
hump-shaped responses to elevation, peaking at mid-elevations (~1,000–1,500 m) and
declining sharply at higher altitudes. These mid-elevation forests—dominated by tall canopy
trees and disturbed by elephants—likely provide optimal conditions for rapid growth and
nutrient uptake, combining moderate temperatures, relatively high nutrient availability, and
dynamic canopy openings. In contrast, high-elevation forests (>2,000 m) were characterized
by smaller, thicker leaves (lower SLA and LA), lower foliar nutrient concentrations, and
elevated C:N ratios, consistent with more conservative strategies under colder, windier, and
nutrient-constrained conditions. Elevation was negatively correlated with LA (r = –0.43,
P < 0.001, Fig. 5a) and SLA (r = –0.40, P < 0.001, Fig. 5b), reflecting anatomical
investment in structural leaf tissue at altitude. Stoichiometric ratios showed contrasting
responses: the C:N ratio increased steeply with elevation ($R^2 = 0.46$, $P < 0.001$, Fig. 5e),
indicating a shift toward greater structural investment and lower nutrient allocation. The N:P
ratio displayed a shallow U-shaped pattern, with the lowest values at mid-elevation, where
both N and P peaked—again suggesting favorable conditions for balanced nutrient
acquisition. Stable isotope signatures further illustrated functional responses to elevation. $\delta^{13}C$
values increased markedly above 1,000 m ($R^2 = 0.35$, $P < 0.001$, Fig. 5g), suggesting
greater intrinsic water-use efficiency (WUE) at higher elevations, where evaporative demand
may be high despite lower temperatures. This pattern likely reflects reduced stomatal
conductance and thicker leaves that limit CO_2 diffusion but conserve water. In contrast,
$\delta^{15}N$ declined sharply with elevation ($R^2 = 0.57$, $P < 0.001$, Fig. 5h), indicating either
increasing reliance on mycorrhizal nitrogen uptake, lower nitrogen turnover rates, or reduced
access to mineral N in cooler soils.
Canopy trees showed higher leaf sizes (LA and SLA) than smaller subcanopy trees,
particularly at low to mid elevations. This suggests a more acquisitive trait strategy in canopy
dominants, likely reflecting their exposure to higher light and a greater competitive advantage
for resource uptake at mid-elevations. However, these differences reverse toward higher
elevations, where taller trees showed reduced SLA and LA due to environmental constraints
(e.g., temperature, wind, and limited nutrient supply).

454
455 **Figure 5.** *Figure 5. Elevational patterns in leaf functional traits of tropical trees on Mount*
456 *Cameroon. Relationships between elevation and (a) leaf area, (b) specific leaf area (SLA), (c)*
457 *leaf nitrogen concentration, (d) leaf phosphorus concentration, (e) leaf carbon-to-nitrogen*

ratio (C:N), (f) leaf nitrogen-to-phosphorus ratio (N:P), (g) leaf carbon isotope composition (δ¹³C), and (h) leaf nitrogen isotope composition (δ¹⁵N). Generalized additive model (GAM) fits are shown for all individuals (purple line), small-statured species (red line), and large-statured species (green line). Boxplots represent trait distributions within elevation bands. All relationships are statistically significant (P < 0.001), except for N:P ratio (P = 0.004). Coefficients of determination (R²) indicate model fit.

3.5. Phylogenetically informed trait–environment relationships

To quantify how environmental factors shape trait variation while accounting for shared evolutionary history, we calculated standardized effect sizes (SES) for 261 trait–environment combinations based on 999 phylogenetically constrained permutations. Of these, 92 associations were statistically significant (|SES| > 2), indicating that observed trait–environment correlations deviated substantially from null expectations under a model of random tip shuffling (Figs. 6 and 7). Traits related to nutrient acquisition and water-use efficiency exhibited the strongest phylogenetic structuring. δ¹⁵N showed the highest number of significant associations (n = 18), followed by δ¹³C (n = 15), C:N (n = 14), and leaf N (LNC, n = 12). In contrast, traits such as leaf P concentration (LPC) showed little or no phylogenetically structured environmental response. Among the 92 significant associations, 50 were positive (SES > 2) and 42 were negative (SES < –2), suggesting both overdispersion and clustering relative to evolutionary expectation.

Climatic variables emerged as the most pervasive environmental filters. δ¹³C declined sharply with rainfall (e.g., SES = –5.9 for total annual rainfall; –5.6 for wet-season rainfall), and increased significantly with wind speed (+3.0), vapor pressure deficit (+3.9), and soil calcium (+3.3)—consistent with altered water-use efficiency and stomatal regulation strategies across phylogenetic lineages. δ¹⁵N increased strongly with mean annual temperature (+7.3), dry-season temperature (+8.3), and wet-season temperature (+8.0), and also responded positively to forest structure metrics, including leaf area index (+5.1) and stem slenderness index (+3.8), while decreasing with wind speed (–7.6) and soil pH (–7.3). These patterns suggest δ¹⁵N is tightly linked to both thermal regimes and forest physiognomy, potentially reflecting shifts in nitrogen sources and mycorrhizal associations.

The C:N ratio exhibited negative associations with rainfall (e.g., SES = –2.3 with dry-season rainfall) and mean annual temperature (–2.7), and positive associations with several edaphic variables, including soil pH (+3.6), calcium (+2.9), magnesium (+2.5), and soil organic carbon (+3.5), suggesting enhanced structural investment and lower foliar nitrogen in nutrient-poor, acidic environments. N:P ratios also reflected phylogenetically structured environmental filtering, increasing in response to mean annual temperature (+2.3) and dry-season temperature (+2.3), while declining with wet-season rainfall (–5.6) and total rainfall (–2.8).

Soil variables exerted strong but trait-specific filtering. Both δ¹³C and δ¹⁵N were significantly influenced by soil calcium, magnesium, and pH, but LNC and LPC showed weaker correlations with measured soil nutrients. For example, leaf N declined with soil magnesium (–3.0), while leaf P showed virtually no significant associations, indicating a weak correspondence between soil nutrient availability and foliar N or P concentrations. This suggests that nutrient uptake strategies are highly species-specific or mediated by unmeasured traits such as root architecture or mycorrhizal type.

Forest structure further modulated trait–environment relationships. SLA declined with wind speed (SES = –3.4), possibly reflecting investment in thicker, more durable leaves under mechanical stress. δ¹⁵N decreased under open canopies (–5.4) and increased with leaf area index (+5.1) and stem basal area (+3.1), indicating that nutrient cycling and nitrogen sources are closely tied to forest canopy complexity and vertical structure. Similarly, δ¹³C increased

with canopy openness (+2.2) and decreased with wet-season rainfall (–5.6), further underscoring its role as a sensitive indicator of water-use strategy modulated by both environmental and phylogenetic constraints.

Figure 6. Phylogenetically-structured trait–environment associations across tropical montane tree species. The heatmap shows standardized effect sizes (SES) for the relationships between nine key leaf traits and 29 environmental variables. SES values reflect the deviation from null expectations under phylogenetic constraints. Environmental variables are grouped by tree size variables, climate, forest structure and soil chemistry. Warm colors indicate strong positive associations; cool colors indicate negative associations.

Figure 7. Trait–environment relationships for key foliar traits across the elevational gradient. Panels show associations between (a) leaf area, (b–c) specific leaf area, (d) foliar C:N ratio, (e) δ¹³C, and (f) δ¹⬜N with selected climatic, edaphic, and structural variables. Lines represent fitted third-order polynomial regressions ±95% confidence intervals.

4. Discussion
4.1. Environmental drivers of functional trait coordination along a tropical mountain gradient

Our findings reveal a robust and ecologically coherent shift in leaf functional traits across a 3,200-meter elevational transect on Mount Cameroon—an Afrotropical biodiversity hotspot—highlighting how climate, soils, forest structure, and evolutionary history jointly shape plant strategies in tropical mountain systems. Trait–environment relationships tracked a continuum from acquisitive to conservative leaf syndromes, structured by elevation-dependent changes in temperature, moisture regimes, nutrient availability, and disturbance. Mid-elevation forests (~1,000–1,500⬜m)—characterized by high productivity, dynamic canopy gaps from elephant disturbance, and moderate climate—harbored species with high SLA, leaf N and P, and low C:N ratios, consistent with fast-growth strategies. In contrast, both lowland rainforests (500–1000⬜m) and upper montane habitats (>2,000⬜m) supported more stress-tolerant species with smaller, tougher leaves, lower nutrient content, and higher δ¹³C and C:N values, shaped respectively by chronic waterlogging or cold and fire-driven nutrient stress. These nonlinear patterns reinforce recent calls to move beyond linear or monotonic expectations in trait–environment relationships across tropical elevation gradients (Fyllas et al., 2017; Oliveras et al., 2020).

Stable isotopes (δ¹³C, δ¹⬜N) and stoichiometric traits (C:N, N:P) emerged as particularly sensitive indicators of environmental filtering. Their variation captured integrated responses to microclimate, moisture, soil nutrient supply, and mycorrhizal strategies, and their strong phylogenetic signal underscores the role of evolutionary history in shaping trait–environment associations. The combined influence of climate, fire, hydrology, and canopy architecture on trait distributions—modulated by taxonomic lineage—supports a model of multidimensional environmental filtering where biotic and abiotic constraints act in concert. Notably, our results diverge from global expectations linking leaf nutrient content tightly to soil fertility (e.g., Ordoñez et al., 2009), instead emphasizing the importance of hydrological

context, leaching, and disturbance in decoupling foliar stoichiometry from total soil nutrient pools in hyper-humid tropical systems.

4.2. Trait shifts along elevation: a move toward conservative strategies

The observed decline in specific leaf area (SLA), leaf area (LA), and nutrient concentrations (N and P), along with increasing carbon-to-nitrogen (C:N) ratios and $\delta^{13}C$ values with elevation, indicates a shift toward more conservative resource-use strategies under cooler, nutrient-poor, and potentially water-limited conditions. These patterns are consistent with findings from Andean and Asian montane systems (Homeier et al., 2021; Helsen et al., 2023), where declining temperatures, nutrient availability, and increasing evaporative demand jointly select for thicker, denser leaves with longer lifespans. Oliveras et al. (2020) similarly reported consistent elevation-related trait shifts across three tropical environmental gradients in South America and Africa, where leaf structure and chemistry co-varied predictably with climate and soil properties. Canopy trees, particularly in lowland forests, consistently exhibited higher SLA than subcanopy individuals, likely reflecting access to deeper nutrient pools and a stronger competitive advantage for light. However, these differences diminished at higher elevations, where temperature and nutrient constraints override vertical stratification, echoing patterns reported from other tropical mountains (Kitayama & Aiba, 2002; Moser et al., 2007).

Foliar $\delta^{13}C$ values increased markedly above 1000 m, suggesting enhanced intrinsic water-use efficiency (WUE), likely due to reduced stomatal conductance (Cernusak et al., 2007; Cornwell et al., 2018). The parallel increase in foliar carbon content and C:N ratio further supports a shift toward structural investment over nutrient acquisition. As leaves become thicker and denser with elevation—reflected in declining specific leaf area (SLA)— they tend to exhibit reduced nutrient concentrations and higher $\delta^{13}C$ values. This pattern likely reflects a shift in resource allocation strategies under cold, dry, and irradiated conditions. Thicker leaves, while conferring greater durability and resistance to desiccation or photodamage, often reduce mesophyll conductance, thereby limiting CO_2 diffusion to the chloroplasts and lowering photosynthetic capacity per unit nitrogen (Wright et al., 2001; Cornwell et al., 2018). The associated increase in $\delta^{13}C$ reflects lower intercellular CO_2 concentrations, thereby enhancing WUE. Accordingly, we observed a negative correlation between leaf nitrogen content and $\delta^{13}C$ across the gradient, consistent with a trade-off between photosynthetic nitrogen efficiency and water-use efficiency in increasingly stressful environments. The inverse relationship between LNC and $\delta^{13}C$ thus highlights the coordinated adjustment of structural, nutritional, and physiological traits in response to the shifting balance between carbon acquisition and stress tolerance along the elevation gradient. Similar trait shifts have been documented in montane forests of Taiwan (Helsen et al., 2023) and in global meta-analyses across elevational gradients (Fyllas et al., 2017)

4.3. Stable Carbon Isotopes Trace Plant Responses to Moisture and Elevational Stress

Stable isotope data provided further insight into plant ecophysiology. Foliar $\delta^{13}C$ values increased with elevation, indicating greater WUE or diffusion limitation of CO_2 under cold, low-pressure conditions—consistent with denser leaf structures and lower SLA (Wright et al., 2001; Cornwell et al., 2018). The steep decline in $\delta^{13}C$ values observed along the elevational gradient from approximately 1000 to 3000 m deviates markedly from the global average of ~1‰ per kilometer typically reported for humid, well-ventilated sites (Körner et al., 1991). Instead, the slope in our dataset is substantially steeper, likely reflecting the combined influence of changing moisture regimes, forest structure, and atmospheric conditions. Notably, total annual precipitation drops sharply from 10,000–12,000 mm at low elevations (300–1100 m) to less than 5000 mm between 1100 and 1600 m, and further declines to below

500 mm in the afromontane zone above 2400 m, where *Agarista salicifolia* and *Myrica*
arborea trees occur in scattered savanna formations (Dolezal et al., 2022). Above 1600 m,
montane forests experience a pronounced dry season from December to March, which likely
contributes to reduced stomatal conductance and elevated $\delta^{13}C$ values at higher elevations. In
contrast, the very negative $\delta^{13}C$ values recorded at lower elevations—approaching –32‰—
may result from consistently high humidity and low light availability under a near-permanent
cloud layer in the lowland rainforest, as well as the influence of soil-respired CO_2 enriched
in ^{12}C. Such conditions elevate internal CO_2 concentrations, enhancing discrimination
against ^{13}C. Collectively, these findings indicate a non-linear, moisture-sensitive response of
foliar $\delta^{13}C$ to elevation, characterized by a transition from saturated, shaded lowland forests to
seasonally dry montane zones and arid Afroalpine environments.
615
4.4. Waterlogging Effects on Foliar $\delta^{13}C$ in Lowland Rainforests
The very negative foliar $\delta^{13}C$ values observed below 1000 m may, in part, reflect the
physiological effects of waterlogging associated with extremely high annual rainfall,
exceeding 10,000–12,000 mm, in these lowland rainforests. Prolonged saturation of soils can
lead to anoxic conditions, impairing root function and reducing both water and nutrient
uptake. This in turn can constrain photosynthetic capacity more strongly than stomatal
conductance, resulting in elevated intercellular CO_2 concentrations (Ci) and increased
discrimination against ^{13}C. The consequence is a shift toward more negative $\delta^{13}C$ values in
leaf tissues. Additionally, waterlogged soils often emit respired CO_2 derived from anaerobic
decomposition, which carries a depleted ^{13}C signature. In humid, poorly ventilated forest
understoreys, re-fixation of this CO_2 may further reduce foliar $\delta^{13}C$. These processes—
combined with low vapor pressure deficit (VPD) and deep canopy shading—mirror the
physiological effects of cloud immersion and reinforce the observed $\delta^{13}C$ depletion at low
elevations. Thus, the extremely moist conditions of the lowland zone likely contribute
significantly to the lower $\delta^{13}C$ values, independent of elevation per se, and should be
considered when interpreting carbon isotope patterns across tropical mountain gradients.
632
4.5. Nitrogen Isotope Signatures Trace Environmental Filtering and Symbiotic
Strategies
Foliar $\delta^{15}N$ values declined markedly with elevation across the Mount Cameroon gradient,
ranging from approximately +5‰ in lowland forests to –5‰ at higher elevations. This trend
reflects a shift from mineral nitrogen acquisition toward organic and mycorrhiza-mediated
uptake in cooler, less biologically active environments, in line with global patterns (Hobbie
and Högberg, 2012; Craine et al., 2015). According to known isotopic signatures, $\delta^{15}N$
values around 0‰ typically indicate nitrogen fixation, values near –2‰ reflect arbuscular
mycorrhizal (AM) associations, and more negative values (–4‰ to –6‰) are characteristic of
ectomycorrhizal (EM) and ericoid mycorrhizal (ERM) strategies (Hobbie and Högberg,
2012). Our dataset supports these distinctions: species from Fabaceae showed relatively
enriched $\delta^{15}N$ values in lowland sites, consistent with nitrogen-fixation and AM-mediated
uptake under high rainfall and leaching. In contrast, at the upper elevational limit, *Agarista*
salicifolia (Ericaceae) exhibited a notably depleted $\delta^{15}N$ value of –3.2‰, suggesting ERM-
mediated nitrogen acquisition in highly organic, N-poor soils typical of high montane
ecosystems. This agrees with the observation by Hobbie and Högberg (2012) that ERM plants
acquire nitrogen from recalcitrant organic sources, leading to strong isotopic discrimination
and depleted foliar $\delta^{15}N$.
Additionally, a bimodal distribution in $\delta^{15}N$ values was observed. The first enrichment
peak (0–1000 m) coincides with intense precipitation and nitrogen leaching on the
southwestern slope, which favors nitrogen-fixing taxa, such as Fabaceae. The second peak

occurred in upper montane forests (2,200–2,500 m), where frequent dry-season fires likely
induce nitrogen losses through volatilization (Doležal et al., 2022). Here, the dominance of
Myrica arborea (Myricaceae), another nitrogen-fixing species, underscores the role of
disturbance-driven nutrient scarcity in shaping nitrogen acquisition strategies. In both cases,
enriched $\delta^{15}N$ signals correspond to disturbance-related nitrogen scarcity and nitrogen
fixation capability. Together, these findings suggest that $\delta^{15}N$ variation in tropical trees is
jointly driven by environmental filtering (rainfall, temperature, fire) and phylogenetically
constrained symbiotic strategies (e.g., N-fixation, AM, EM, ERM), echoing patterns reported
by Hobbie and Högberg (2012) and reinforcing the functional link between nitrogen isotopes,
mycorrhizal type, and plant adaptation across elevation.

Foliar $\delta^{15}N$ showed a strong positive correlation with dry-season rainfall
($R^2 = 0.42$, $P < 0.001$; Fig. 7), suggesting that wetter dry seasons promote nitrogen
losses via leaching or denitrification, enriching the $\delta^{15}N$ of residual soil nitrogen. This
isotopic signal corresponds spatially with the lowland rainforest zone on Mt. Cameroon,
where tree growth is paradoxically constrained during the peak wet season due to anoxia and
low light conditions under high cloud cover (Plavcová et al., 2024). In these forests, tree
radial growth often peaks during the dry season, when soils are aerated but remain sufficiently
moist to permit active xylem expansion. In contrast, upper montane trees experience growth
limitation during the dry season due to water scarcity. The elevated $\delta^{15}N$ values in wetter
lowland forests may thus reflect both increased nitrogen loss and the prevalence of nitrogen-
fixing species that dominate under such leached, anaerobic conditions. This co-occurrence of
enriched $\delta^{15}N$ and growth suppression during the wettest periods highlights the complex
coupling between nitrogen cycling, water availability, and growth phenology in Afrotropical
forests.

4.6. Divergence from Global Patterns: Decoupling of Leaf Nutrients from Soil
Fertility
Our results contrast markedly with global-scale findings by Ordoñez et al. (2009), who
reported that SLA, LNC, and LPC generally increased with soil fertility across ecosystems
worldwide. In our Afrotropical elevational gradient on Mt. Cameroon, we found no consistent
relationship between foliar N or P concentrations and corresponding soil nutrient pools. This
decoupling likely reflects the overriding influence of extreme rainfall and leaching in low-
elevation tropical rainforests (650 m), where soils are heavily weathered, organic matter is
low, and nutrient retention is poor—conditions that suppress mineralization and reduce
nutrient bioavailability despite high productivity and biomass.

Indeed, the lowest soil and foliar N and P values were observed in these hyper-humid
lowland forests, which receive up to 10,000 mm of rainfall annually. This stands in contrast
to global patterns where soil nutrient levels explain substantial variation in foliar traits (cf.
Ordoñez et al., 2009; Wright et al., 2004). Only in mid-elevation forests between 1500 and
1800 m—subject to frequent disturbance by forest elephants—did we observe concurrent
peaks in both soil and leaf nutrient concentrations. These disturbed zones appear to maintain
more dynamic nutrient turnover through bioturbation, canopy gap formation, and enhanced
microbial activity, reinforcing localized nutrient availability and uptake.

This pattern highlights how edaphic control of leaf functional traits may be contingent
on hydrological context and disturbance regimes. In hyper-humid systems, leaching and redox
constraints decouple foliar stoichiometry from soil total nutrient pools (e.g., Vitousek et al.,
2010; Townsend et al., 2007), while in montane systems, moderate disturbance and improved
drainage can enhance nutrient accessibility despite lower temperatures. These findings
underscore the limitations of applying generalized global trait–soil models to hyperdiverse,
moisture-saturated ecosystems and support calls to incorporate regional climatic and

ecological dynamics into predictive trait-based models (e.g., Fyllas et al., 2017; Oliveras et
al., 2020).
706
4.7. Phosphorus Availability is Shaped by Hydrology and Fire, Not Elevation Alone
Contrary to expectations based on the leaf economic spectrum (LES), our results reveal that
phosphorus limitation is most pronounced at lower elevations, rather than at high-elevation
sites, where colder temperatures and slower nutrient cycling are typically thought to constrain
P availability (Vitousek et al., 2010 and references therein). Both foliar and soil phosphorus
concentrations were lowest between 500 and 1000 m elevation—within the hyper-humid
rainforest belt receiving up to 10,000 mm of annual rainfall (Řeháková et al., 2022). These
lowland soils are often organic-poor, poorly drained, and subject to temporal waterlogging,
creating anoxic conditions that suppress mineralization and promote phosphorus fixation
through redox-sensitive sorption processes. In contrast, mid- to high-elevation soils (2200–
2400 m) exhibited notably higher available phosphorus concentrations, coinciding with
montane forest and Afroalpine savanna zones that experience regular dry-season fires during
January (Doležal et al., 2022). These fire events can enhance soil phosphorus availability
through combustion of organic material, ash deposition, and altered microbial turnover,
contributing to transient but ecologically significant peaks in soil P at higher elevations
(Hogue & Inglett, 2012; Veldhuis et al., 2016).
Interestingly, while soil P peaks at higher elevations, foliar P content displays the
opposite pattern—declining consistently with elevation in line with LES predictions. This
divergence suggests that low leaf P at high elevation is not driven by reduced soil P
availability, but rather by physiological or morphological constraints on nutrient uptake,
transport, or demand. Cold temperatures may reduce root activity and phosphorus mobility,
while shorter growing seasons and conservative life-history strategies may reduce foliar
nutrient investment despite adequate supply in the soil. Conversely, in the lowlands, the
combination of high foliar P demand and low P bioavailability in saturated, leached, and
redox-active soils likely intensifies actual phosphorus limitation. Together, these findings
reveal that soil and plant phosphorus dynamics are decoupled along elevation gradients and
shaped by the interplay of hydrological context, disturbance regimes, and plant functional
strategies.
735
4.8. Stoichiometric Shifts and Nutrient Limitation Along Elevation
Elevational patterns in both soil and foliar nutrient stoichiometry revealed significant
nonlinear shifts in ecosystem nutrient dynamics along the Mount Cameroon gradient. The soil
nitrogen-to-phosphorus (N:P) ratio exhibited a pronounced hump-shaped response to
elevation ($R^2 = 0.41$, $P < 0.001$), peaking in mid-elevation forests (~1500 m) that are
regularly disturbed by forest elephants. These open-canopy sites likely support enhanced
microbial activity, nutrient cycling, and mineralization, driven by increased light availability,
accelerated litter decomposition, and nutrient redistribution through dung deposition and
trampling. In contrast, foliar N:P ratios showed a much weaker elevational trend ($R^2 = 0.03$,
$P = 0.004$), though they too peaked modestly in the same mid-elevation zone. This partial
decoupling between soil and foliar stoichiometry highlights the influence of additional
factors, including species identity, functional strategy, and mycorrhizal associations, on leaf-
level nutrient ratios.
Our observed foliar N:P ratios largely fell within the intermediate range (10–20),
suggesting co-limitation by N and P. However, higher values at mid-elevation may indicate
increasing P limitation in sites with relatively high N availability. These patterns are broadly
consistent with the global synthesis by Güsewell (2004), who demonstrated that foliar N:P
ratios reflect both nutrient supply and a species' ecological strategies, with higher N:P values

being more common in stress-tolerant or graminoid species. They also align with findings from Townsend et al. (2007), who emphasized that while tropical forests are often considered N-rich and P-limited, foliar N:P ratios are not tightly constrained across tropical ecosystems, largely due to species-level variability and local environmental heterogeneity. Similarly, Soethe et al. (2008) found elevated foliar N:P at mid-elevations in Ecuatorian montane forests, where litter decomposition was more active and nutrient availability more favorable than at higher altitudes.

4.9. Phylogenetically structured environmental filtering

Our results indicate that environmental filtering of functional traits in tropical forest communities is profoundly shaped by phylogenetic history (Cavender-Bares et al., 2009; Swenson and Enquist, 2009). Traits related to nutrient acquisition and physiological regulation—particularly δ^{15}N, δ^{13}C, and C:N—demonstrated the strongest departures from phylogenetically randomized null expectations, indicating that both climate and soil gradients select for conserved trait syndromes within evolutionary lineages (Cornwell & Ackerly, 2009; Laughlin & Messier, 2015). Climatic factors were key drivers of trait differentiation. The strong negative SES values for δ^{13}C in wetter environments and positive values with wind stress suggest that lineages differ in intrinsic stomatal behavior and water-use efficiency, shaped by both habitat moisture and mechanical exposure (Farquhar et al., 1989; Cernusak et al., 2007; Cornwell et al., 2018). Similarly, the consistent positive associations of δ^{15}N with temperature and LAI may reflect evolutionary shifts in nitrogen source pools, uptake mechanisms, or mycorrhizal symbioses in warmer and less disturbed environments (Högberg, 1997; Craine et al., 2015; Mayor et al., 2014).

Soil properties exerted clear but trait-specific constraints. The positive SES values of C:N with base saturation and pH imply that nutrient allocation strategies are tightly linked to edaphic fertility and acidity, with conservative nutrient use prevailing in lineages associated with richer soils (Townsend et al., 2007; Porder & Hilley, 2011). Increasing base saturation and reduced nitrogen availability at higher pH may constrain nitrogen uptake relative to carbon assimilation. As soil pH rises, ammonium is increasingly converted to nitrate, which is more mobile and susceptible to leaching in humid tropical systems (Vitousek & Sanford, 1986). Additionally, enhanced microbial activity under near-neutral pH can promote nitrogen immobilization in microbial biomass, further reducing nitrogen availability to plants (Güsewell, 2004). Base-rich soils may also induce nutrient imbalances or competition between cations (e.g. Ca^{2+}, Mg^{2+}, K^+) and ammonium, limiting nitrogen uptake (Townsend et al., 2007). The resulting increase in foliar C:N reflects a shift toward nutrient-conservative strategies, emphasizing structural over metabolic investment under nutrient-limiting conditions. Similar patterns have been reported from base-rich montane forests and fire-affected soils (Oliveras et al., 2020; Santiago et al., 2017), supporting the interpretation that foliar stoichiometry integrates both soil chemistry and functional adaptation to resource constraints.

By contrast, traits such as P showed weak or absent phylogenetic signal, suggesting higher plasticity or lower evolutionary constraint (Poorter et al., 2009; Baraloto et al., 2010). Forest structure also shaped trait patterns in phylogenetically non-random ways. Increases in C:N and δ^{15}N with openness, and declines in SLA with wind exposure, point to the role of vertical canopy architecture and microclimatic variation in shaping resource-use and structural traits (Kitajima & Poorter, 2010; Markesteijn et al., 2011). These relationships reflect how mechanical stress and light availability interact with evolutionary legacy to constrain functional adaptation (Asner et al., 2018; Givnish, 2002).

Functional differentiation and community assembly patterns revealed further insight into evolutionary constraints. PCA showed distinct trait syndromes and strong family-level

clustering. High-elevation lineages—such as Ericaceae, Myrtaceae, and Myricaceae—were associated with conservative strategies including high $\delta^{13}C$ and C:N, and low SLA and foliar nutrient content, reflecting adaptations to cold, nutrient-poor, or fire-prone environments. In contrast, lowland families such as Irvingiaceae and Clusiaceae exhibited acquisitive traits associated with rapid growth under warm, moist, and fertile conditions—aligning with global LES expectations (Wright et al., 2004; Díaz et al., 2016). Some families, such as Fabaceae and Rosaceae, occupied broader functional spaces. Fabaceae in particular maintained elevated $\delta^1\square N$ values across the gradient, consistent with their N-fixing capacity and resilience under both leaching and fire-prone regimes. This taxonomic control over trait expression reinforces findings by Oliveras et al. (2020), who demonstrated that family identity often explains more variance in plant traits than site-level environmental variation.

Taken together, these findings support a model of multidimensional environmental filtering where functional traits are shaped not only by climate and soils but also by lineage-specific evolutionary history and disturbance regimes. Acquisitive trait syndromes (e.g., high SLA, N, P) are favored in warm, nutrient-rich, open forests, while conservative syndromes (e.g., high C:N, $\delta^{13}C$) dominate in cooler, less fertile, or more structurally complex habitats. The strength and direction of SES patterns, combined with phylogenetic clustering in multivariate trait space, highlight the importance of integrating taxonomic and evolutionary information when interpreting trait–environment relationships and forecasting tropical forest responses to global change (Fyllas et al., 2017; Swenson & Enquist, 2009).

5. Synthesis and implications for tropical trait-based ecology

In sum, the Mount Cameroon elevational gradient reveals strong and coordinated shifts in plant functional traits that reflect both adaptive responses to local environmental stressors and deep phylogenetic legacies. Trait–environment relationships—especially those involving SLA, C:N, $\delta^{13}C$, and $\delta^1\square N$—aligned with resource-use trade-offs that were modulated by temperature, precipitation, fire regimes, and soil chemistry, but also structured by lineage-specific constraints and strategies. Mid-elevation forests emerged as functional hotspots, combining high productivity, canopy disturbance, and nutrient availability to support acquisitive leaf traits. In contrast, lowland and highland forests converged on conservative strategies through different mechanisms—waterlogging and nutrient leaching in the former, cold stress and fire-driven nutrient losses in the latter.

These results reinforce the global utility of trait-based approaches in tropical forest ecology, especially when combined with stable isotope and stoichiometric markers that integrate physiological function and environmental history. As climate change accelerates, altering temperature regimes, rainfall seasonality, and fire frequency across tropical mountains (Fyllas et al., 2017; Helsen et al., 2023), the trait strategies documented here can serve as predictive tools for understanding shifts in community composition, productivity, and biogeochemical cycling. Our findings underscore the need for regionally grounded, mechanistically informed trait-based models that account for hydrology, disturbance, and evolutionary constraints when forecasting forest responses to global environmental change.

Author contributions
JD conceived the ideas and designed the methodology; JD, VB and KK collected the data; JD and KK analysed the data; JD, AR and JG led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Funding
The project was supported by the Czech Science Foundation (GACR 24-11954S), the Czech Academy of Sciences (RVO 67985939), and the Ministry of Education, Youth and Sport of

854   the Czech Republic (MSMT) (#VES24, INTER-ACTION, LUAUS24258). JG wants to thank
855   the national project PID2022-139455NB-C31 funded by MCIN/AEI/
856   10.13039/501100011033 and by the "European Union NextGenerationEU/PRTR".

Data Availability Statement

Data available from the Zenodo repository doi: XXX (Dolezal, 2025).

Conflicts of Interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Abatzoglou, J.T., Dobrowski, S.Z., Parks, S.A., and Hegewisch, K.C. (2018). TerraClimate, a high-resolution global dataset of monthly climate and climatic water balance from 1958 to 2015. Sci. Data 5: 170191.

2. Ågren, G.I. (2004). The C:N:P stoichiometry of autotrophs – theory and observations. Ecol. Lett. 7: 185–191.

3. Asner, G.P., Brodrick, P.G., Philipson, C., Vaughn, N.R., Martin, R.E., Knapp, D.E., Heckler, J., Evans, L.J., Jucker, T., Goossens, B., Stark, D.J., Reynolds, G., Ong, R., Renneboog, N., Kugan, F., and Coomes, D.A. (2018). Mapped aboveground carbon stocks to advance forest conservation and recovery in Malaysian Borneo. Biol. Conserv. 217: 289–310.

4. Cable, S., and Cheek, M. (1998). The plants of Mt Cameroon: a conservation checklist. Royal Botanic Gardens, Kew.

5. Cavender-Bares, J., Kozak, K.H., Fine, P.V.A., and Kembel, S.W. (2009). The merging of community ecology and phylogenetic biology. Ecol. Lett. 12: 693–715.

6. Cernusak, L.A., Winter, K., Turner, B.L. (2007). Leaf nitrogen to phosphorus ratios of tropical trees: experimental assessment of physiological and environmental controls. New Phytol. 172: 170–180.

7. Cheek, M., Cable, S., Hepper, F.N., Ndam, N., and Watts, J. (1996). Mapping plant biodiversity on Mount Cameroon. In: van der Maesen, L.J.G., van der Burgt, X.M., and van Medenbach de Rooy, J.M. (eds), The biodiversity of African plants. Springer, pp. 110–120.

8. Cornwell, W.K., and Ackerly, D.D. (2009). Community assembly and shifts in plant trait distributions across an environmental gradient in coastal California. Ecol. Monogr. 79: 109–126.

9. Cornwell, W.K., Wright, I.J., Turner, J., Maire, V., Barbour, M.M., Cernusak, L.A., Dawson, T.E., Ellsworth, D.S., Farquhar, G.D., and Griffiths, H. (2018). Quantifying the advantage of thinner leaves for carbon uptake across environmental gradients. New Phytol. 219: 87–96.

10. Craine, J.M., Brookshire, E.N.J., Cramer, M.D., Hasselquist, N.J., Koba, K., Marin-Spiotta, E., and Wang, L. (2015). Ecological interpretations of nitrogen isotope ratios of terrestrial plants and soils. Plant Soil 396: 1–26.

11. Díaz, S., Kattge, J., Cornelissen, J.H.C., et al. (2016). The global spectrum of plant form and function. Nature, 529: 167–171.

12. Doležal, J., Dančák, M., Kučera, J., Majeský, Ľ., Altman, J., Řeháková, K., Čapková, K., Vondrák, J., Fibich, P., and Liancourt, P. (2022). Fire, climate and biotic interactions shape diversity patterns along an Afrotropical elevation gradient. J. Biogeogr. 49: 1248–1259.

13. Doležal, J., Korznikov, K., Altman, J., Fibich, P., Řeháková, K., Lltonga, E.N., Majeský, Ľ., and Dančák, M. (2023). Ecological niches of epiphyllous bryophytes along Afrotropical elevation gradient. Oikos 2023: e09772. https://doi.org/10.1111/oik.09772

14. Farquhar, G.D., Ehleringer, J.R., and Hubick, K.T. (1989). Carbon isotope discrimination and photosynthesis. Annu. Rev. Plant Biol. 40: 503–537.

15. Ferenc M, Sedláček O, Tropek R, et al. Something is missing at the bottom: Importance of coastal rainforests for conservation of trees, birds and butterflies in the Mount Cameroon area. Afr J Ecol. 2018; 56: 679–683. https://doi.org/10.1111/aje.12506

16. Fyllas, N.M., Bentley, L.P., Shenkin, A., Asner, G.P., Atkin, O.K., Díaz, S., Enquist, B.J., Farfan-Rios, W., Gloor, E., Guerrieri, R., et al. (2017). Solar radiation and functional traits explain the decline of forest primary productivity along a tropical elevation gradient. Ecol. Lett. 20: 730–740. https://doi.org/10.1111/ele.12771

17. Garnier, E., Navas, M.-L., and Grigulis, K. (2015). Plant Functional Diversity: Organism Traits, Community Structure, and Ecosystem Properties. Oxford University Press.

18. Girardin, C.A.J., Malhi, Y., Aragao, L.E.O.C., et al. (2014). Seasonality of above-ground net primary productivity along an Andean altitudinal transect in Peru. J. Trop. Ecol. 30: 1–17. https://doi.org/10.1017/S0266467414000443

19. Girardin, C.A.J., Malhi, Y., Aragao, L.E.O.C., et al. (2016). Seasonal trends of Amazonian rainforest phenology, net primary productivity, and carbon allocation. Glob. Biogeochem. Cycles 30: 700–715. https://doi.org/10.1002/2015GB005270

20. Givnish, T.J. (2002). Adaptive significance of evergreen vs deciduous leaves: solving the triple paradox. Silva Fenn. 36: 703–743.

21. Goldsmith, G.R., Bentley, L.P., Shenkin, A., Salinas, N., Blonder, B., Martin, R.E., et al. (2017). Variation in leaf wettability traits along a tropical montane elevation gradient. New Phytol. 214: 989–1001. https://doi.org/10.1111/nph.14121

22. Güsewell, S. (2004). N:P ratios in terrestrial plants: variation and functional significance. New Phytol. 164: 243–266. https://doi.org/10.1111/j.1469-8137.2004.01192.x

23. Helsen, K., Viana, J.L., Lin, T.-Y., Kuo, L.-Y., and Zelený, D. (2023). Functional-trait contrasts between terrestrial and epiphytic ferns in Taiwanese subtropical cloud forests. J. Veg. Sci. 34: e13220. https://doi.org/10.1111/jvs.13220

24. Hobbie, E.A., and Högberg, P. (2012). Nitrogen isotopes link mycorrhizal fungi and plants to nitrogen dynamics. New Phytol. 196: 367–382. https://doi.org/10.1111/j.1469-8137.2012.04292.x

25. Högberg, P. (1997). Tansley Review No. 95: 15N natural abundance in soil-plant systems. New Phytol. 137: 179–203. https://doi.org/10.1046/j.1469-8137.1997.00808.x

26. Hogue, B. A., & Inglett, P. W. (2012). Nutrient release from combustion residues of two contrasting herbaceous vegetation types. Science of the Total Environment, 431, 9–19.

27. Holtum, J.A.M., Winter, K. Carbon isotope composition of canopy leaves in a tropical forest in Panama throughout a seasonal cycle. Trees 19, 545–551 (2005). https://doi.org/10.1007/s00468-005-0413-8

28. Homeier, J., Seeler, T., Pierick, K. et al. Leaf trait variation in species-rich tropical Andean forests. Sci Rep 11, 9993 (2021). https://doi.org/10.1038/s41598-021-89190-8

29. Hořák, D., Ferenc, M., Sedláček, O., Motombi, F. N., Svoboda,M., Altman, J., Albrecht, T., Djomo Nana, E., Janeček, Š.,Dančák, M., Majeský, Ľ., Lltonga, E. N.

and Doležal, J. 2019.Forest structure determines spatial changes in avian communityalong an elevational gradient in tropical Africa. – J. Biogeogr.46: 2466–2478.

30. Kitayama, K. and Aiba, S.-I. (2002), Ecosystem structure and productivity of tropical rain forests along altitudinal gradients with contrasting soil phosphorus pools on Mount Kinabalu, Borneo. Journal of Ecology, 90: 37-51. https://doi.org/10.1046/j.0022-0477.2001.00634.x

31. Koerselman, W. & Meuleman, A.F.M. (1996). The vegetation N:P ratio: a new tool to detect the nature of nutrient limitation. Journal of Applied Ecology, 33(6), 1441–1450.

32. Körner, C. (2021). Alpine plant life: Functional plant ecology of high mountain ecosystems. Springer Nature.

33. Körner, C., Farquhar, G.D., & Roksandic, Z. (1991). A global survey of carbon isotope discrimination in plants from high elevation. Oecologia, 88(1), 30–40.

34. Laughlin, D. C., & Messier, J. (2015). Fitness of multidimensional phenotypes in dynamic adaptive landscapes. Trends in Ecology & Evolution, 30(8), 487–496. https://doi.org/10.1016/j.tree.2015.06.003

35. Laughlin, D.C. (2014), The intrinsic dimensionality of plant traits and its relevance to community assembly. J Ecol, 102: 186-193. https://doi.org/10.1111/1365-2745.12187

36. Laughlin, D.C. (2014). The intrinsic dimensionality of plant traits and its relevance to community assembly. Journal of Ecology, 102(1), 186–193.

37. Maicher, V., Delabye, S., Murkwe, M., Doležal, J., Altman, J.,Kobe, I. N., Desmist, J., Fokam, E. B., Pyrcz, T. and Tropek,R. 2020b. Effects of disturbances by forest elephants on diver-sity of trees and insects in tropical rainforests on Mount Cam-eroon. – Sci. Rep. 10: 21618

38. Maicher, V., Sáfián, S., Murkwe, M., Delabye, S., Przybyłowicz, Ł.,Potocký, P., Kobe, I. N., Janeček, Š., Mertens, J. E. J., Fokam,E. B., Pyrcz, T., Doležal, J., Altman, J., Hořák, D., Fiedler, K.and Tropek, R. 2020. Seasonal shifts of biodiversity patternsand species elevation ranges of butterflies and moths along acomplete rainforest elevational gradient on Mount Cameroon.– J. Biogeogr. 47: 342–354.

39. Markesteijn, L. and Poorter, L. (2009), Seedling root morphology and biomass allocation of 62 tropical tree species in relation to drought- and shade-tolerance. Journal of Ecology, 97: 311-325. https://doi.org/10.1111/j.1365-2745.2008.01466.x

40. Mayor, J.R. et al. (2014). Leaf nitrogen and phosphorus drawdown during leaf development reflects changes in cell structure and metabolism. New Phytologist, 203(4), 1260–1270.

41. McCullagh, P., and Nelder, J.A. (1989). Generalized Linear Models (2nd ed.). Chapman and Hall, London. https://doi.org/10.1007/978-1-4899-3242-6

42. Méndez-Alonzo, R., Paz, H., Zuluaga, R.C., Rosell, J.A. and Olson, M.E. (2012), Coordinated evolution of leaf and stem economics in tropical dry forest trees. Ecology, 93: 2397-2406. https://doi.org/10.1890/11-1213.1

43. Moser, G., Hertel, D. & Leuschner, C. 2007. Altitudinal change in LAI and stand leaf biomass in tropical montane forests: a transect study in Ecuador and a pan-tropical meta-analysis. Ecosystems 10:924-935.

44. Oliveras, I., Bentley, L., Fyllas, N.M., Gvozdevaite, A., Shenkin, A.F., Peprah, T., Morandi, P., Peixoto, K.S., Boakye, M., Adu-Bredu, S., Schwantes Marimon, B., Marimon Junior, B.H., Salinas, N., Martin, R., Asner, G., Díaz, S., Enquist, B.J., and Malhi, Y. (2020). The influence of taxonomy and environment on leaf trait variation along tropical abiotic gradients. Front. For. Glob. Change 3: 18. https://doi.org/10.3389/ffgc.2020.00018

45. Ordoñez, J.C., Van Bodegom, P.M., Witte, J.-P.M., Wright, I.J., Reich, P.B., and Aerts, R. (2009). A global study of relationships between leaf traits, climate and soil measures of nutrient fertility. Glob. Ecol. Biogeogr. 18: 137–149. https://doi.org/10.1111/j.1466-8238.2008.00441.x

46. Payton, R.W. (1993). Final project report R4600: Ecology, altitudinal zonation and conservation of tropical rain forests of Mount Cameroon. Soil Survey and Land Research Centre.

47. Plavcová, L., Jandová, V., Altman, J., Liancourt, P., Korznikov, K., and Doležal, J. (2024). Variations in wood anatomy in Afrotropical trees with a particular emphasis on radial and axial parenchyma. Ann. Bot. 134: 151–162.

48. Porder, S., and Ramachandran, S. (2013). The phosphorus concentration of common rocks—a potential driver of ecosystem P status. Plant Soil 367: 41–55.

49. Pringle, E.G., Adams, R.I., Broadbent, E., Busby, P.E., Donatti, C.I., Kurten, E.L., Renton, K., and Dirzo, R. (2011). Distinct leaf-trait syndromes of evergreen and deciduous trees in a seasonally dry tropical forest. Biotropica 43: 299–308.

50. Proctor, J., Edwards, I.D., Payton, R.W., and Nagy, L. (2007). Zonation of forest vegetation and soils of Mount Cameroon, West Africa. Plant Ecol. 192: 251–269.

51. Řeháková, K., Čapková, K., Altman, J., Dančák, M., Majeský, Ľ., and Doležal, J. (2022). Contrasting patterns of soil chemistry and vegetation cover determine diversity changes of soil phototrophs along an Afrotropical elevation gradient. Ecosystems 25: 1020–1036.

52. Santiago, L., Silvera, K., Andrade, J., and Dawson, T. (2017). Functional strategies of tropical dry forest plants in relation to growth form and isotopic composition. Environ. Res. Lett. 12: 1088–9326.

53. Soethe, N., Lehmann, J., and Engels, C. (2008). Nutrient availability at different altitudes in a tropical montane forest in Ecuador. J. Trop. Ecol. 24: 397–406.

54. Swenson, N.G., and Enquist, B.J. (2009). Opposing assembly mechanisms in a Neotropical dry forest: Implications for phylogenetic and functional community ecology. Ecology 90: 2161–2170.

55. ter Braak, C.J.F., and Smilauer, P. (2012). Canoco reference manual and user's guide: software for ordination, version 5.0. Microcomputer Power.

56. Townsend, A.R., Cleveland, C.C., Asner, G.P., and Bustamante, M.M.C. (2007). Controls over foliar N:P ratios in tropical rain forests. Ecology 88: 107–118. https://doi.org/10.1890/0012-9658(2007)88[107:COFNRI]2.0.CO;2

57. Veldhuis, M.P., Hulshof, A., Fokkema, W., Berg, M.P., and Olff, H. (2016). Understanding nutrient dynamics in an African savanna: Local biotic interactions outweigh a major regional rainfall gradient. J. Ecol. 104: 913–923.

58. Vitousek, P.M., and Sanford, R.L. (1986). Nutrient cycling in moist tropical forest. Annu. Rev. Ecol. Syst. 17: 137–167.

59. Vitousek, P.M., Porder, S., Houlton, B.Z., and Chadwick, O.A. (2010). Terrestrial phosphorus limitation: Mechanisms, implications, and nitrogen–phosphorus interactions. Ecol. Appl. 20: 5–15.

60. Wright, I.J., Reich, P.B., Westoby, M., et al. (2004). The worldwide leaf economics spectrum. Nature 428: 821–827. https://doi.org/10.1038/nature02403

61. Wright, I.J., Reich, P.B., Westoby, M., et al. (2004). The worldwide leaf economics spectrum. Nature 428: 821–827.

62. Yang, Y., Fang, J., Ji, C., and Han, W. (2010). Above- and belowground biomass allocation in Tibetan grasslands. J. Veg. Sci. 21: 493–500.

1051   63. Yang, Y., Siegwolf, R.T.W., and Körner, C. (2015). Species-specific and environment-
1052   induced variation of δ13C and δ15N in alpine plants. Front. Plant Sci. 6: 423.
1053   https://doi.org/10.3389/fpls.2015.00423