Enhanced muscle uptake of chemically optimized miR-23b antisense oligonucleotides as lead compounds for Myotonic Dystrophy type 1

Irene González-Martínez,[1,2,3,#] Estefanía Cerro-Herreros,[1,2,#†] Marc Carrascosa-Sàez,[4,‡] Andrea García-Rey,[1,2] Diego Piqueras Losilla,[4] Anna Colom-Rodrigo,[4,§] Nerea Moreno,[1,2,3] Mouli Chakraborty,[4] Aline Huguet-Lachon,[5] Anchel González-Barriga,[5] Neia Naldaiz-Gastesi,[6,7,8] Martxel Dehesa,[6,7,8] Ana Díaz-Maqueda,[4] Nuria Barquero,[4] Miguel A Varela,[9,10] Adolfo López de Munain,[6,7,8,11,12] Ramon Eritja,[13,14] Geneviève Gourdon,[5] Arturo López-Castel,[1,2,3] Manuel Pérez-Alonso,[1,2,3] Beatriz Llamusi,[4] and Rubén Artero[1,2,3*]

[1] Human Translational Genomics Group, University Institute for Biotechnology and Biomedicine (BIOTECMED), University of Valencia; Av. Dr. Moliner 50, 46100 Valencia, Spain.

[2] INCLIVA Biomedical Research Institute; Av. Menendez Pelayo 4, 46010 Valencia, Spain.

[3] CIBERER, IISCIII; Av. Monforte de Lemos 3-5, 28029 Madrid, Spain.

[4] ARTHEx Biotech, Centro de Investigación Príncipe Felipe (CIPF); Valencia, Spain.

[5] Sorbonne Université, Inserm, Institut de Myologie, Centre de Recherche en Myologie; Paris, France.

[6] Biodonostia-BioGipuzkoa, Neurosciences Area, Neuromuscular Diseases Laboratory; San Sebastian, Spain.

[7] Department of Neurology, Hospital Universitario Donostia. OSAKIDETZA, San Sebastian, Spain.

[8] CIBERNED, ISCIII; Madrid, Spain.

[9] Department of Paediatrics, Institute of Developmental and Regenerative Medicine (IDRM), University of Oxford; Oxford, UK.

[10] MDUK Oxford Neuromuscular Centre, University of Oxford; Oxford, UK.

[11] Department of Neurosciences, University of the Basque Country, San Sebastian, Spain.

[12] Faculty of Health Sciences, University of Deusto, Bilbao, Spain.

[13] Nucleic Acids Chemistry Group, Institute for Advanced Chemistry of Catalonia (IQAC-CSIC),

Barcelona, Spain.

[14] Networking Center on Bioengineering, Biomaterials and Nanomedicine (CIBER-BBN), Barcelona,

Spain.

# Contributed equally.

*Corresponding author. Email: ruben.artero@uv.es

† Present address: ARTHEx Biotech, Centro de Investigación Príncipe Felipe (CIPF); Valencia, Spain.

‡ Present address: Institute for Integrative Systems Biology (I2SysBio), Consejo Superior de

Investigaciones Científicas, Universitat de València; Paterna, Valencia, Spain.

§ Present address: Human Translational Genomics Group, University Institute of Biotechnology

and Biomedicine (BIOTECMED), Universitat de València; Burjasot, Spain.

Short title: Oleic Conjugation Enhances antimiR-23b Therapy

Abstract:

Myotonic dystrophy type 1 (DM1) is a multisystemic disorder caused by CTG repeat expansions in *DMPK*. Mutant transcripts containing expanded CUG repeats form ribonuclear foci that sequester muscleblind-like splicing regulator (MBNL) proteins, key regulators of RNA splicing and metabolism. This functional depletion leads to widespread mis-splicing and persistence of fetal transcript profiles, which underlie muscle weakness, myotonia, and muscle atrophy. In addition, miR-23b is upregulated in DM1 muscle and further represses MBNL1 translation, amplifying molecular defects. We developed chemically optimized miRNA-targeting antisense oligonucleotides (antimiRs) to inhibit miR-23b and restore functional MBNL1 levels. Using a multi-step screening process, we evaluated antimiRs with varying sequences, lengths, chemical modifications, and lipid conjugations. A key optimization was a 3'-oleic acid conjugation combined with specific chemical modifications, which enhanced muscle uptake and efficacy. Lead candidates showed strong activity in preclinical models (HSA^{LR} and DMSXL mice, and human myoblasts), increasing MBNL1 levels, correcting mis-splicing, improving muscle strength, and reducing myotonia. They also exhibited efficient biodistribution to skeletal muscle, a critical DM1-affected tissue. *In vitro* toxicology indicated a favorable safety profile with minimal immune or renal toxicity. The antimiR mechanism was conserved in rat and pig fibroblasts. Overall, two lead antimiRs emerged as promising therapeutic candidates for DM1, with improved pharmacokinetics, tissue targeting, and safety, supporting the potential of microRNA-based approaches to correct key molecular defects in this disorder.

Main Text:

INTRODUCTION

Myotonic dystrophy type 1 (DM1, OMIM#160900) is a rare genetic disorder with an overall prevalence of one in 3,000–8,000 individuals worldwide. [1] DM1 is characterized by expansion of an unstable CTG microsatellite repeat in the 3'-untranslated region (UTR) of *DM1 protein kinase* (*DMPK*, OMIM*605377). Mutant *DMPK* transcripts accumulate in the nucleus and induce toxicity by affecting RNA processing mechanisms, disturbing downstream regulation of gene expression.[2] One key molecular event contributing to DM1 pathogenesis involves sequestration of the muscleblind-like (MBNL1 and 2) splicing regulator family of proteins by CUG expansions, particularly MBNL1 in skeletal muscles, which forms ribonuclear aggregates (foci).[3] This anomalous functional depletion of MBNLs severely affects the available levels of MBNL factors, impairing their critical regulatory role in alternative pre-mRNA splicing, microRNA biogenesis, and polyadenylation.[4–6] Additional stress responses caused by toxic *DMPK* mRNA accumulation leads to the activation of MBNL1 antagonists such as CELF1.[7] Together, MBNL1 and CELF1 proteins function as developmental switches, and their imbalance in DM1 causes abnormal persistence of fetal patterns in processed mRNAs, resulting in inadequate protein isoforms or altered gene expression levels.[8] This pathological mechanism affects hundreds of genes in various tissues and organs, many of which have been directly linked to the specific symptoms of DM1. [9–12] However, several reports support that lack of MBNL1 function contributes the most to muscular and cognitive DM1 phenotypes. [13–19] This is particularly favorable from a therapeutic perspective because the limited availability of MBNL1 and MBNL2 proteins, which are partially functionally redundant in the muscles,[20] can be compensated by upregulating *MBNL1* (OMIM*606516) and/or *MBNL2* (OMIM*607327), which remain normal at the genomic level in patients. Indeed, MBNL1

overexpression effectively rescues DM1 phenotypes in preclinical models[21–23] and is well tolerated

in transgenic mice.[17]

Recent findings indicate that miR-218 and miR-23b (translational repressors of MBNL1 and 2)

[24,25] are significantly overexpressed in diseased muscle,[25,26] exacerbating functional protein

depletion by sequestration. The observed downregulation of MBNL1 in DM1 cell models,

including primary myoblasts differentiated in vitro, can be explained by elevated miRNA levels [27]

and antagonizing these miRNAs with antimiRs increased MBNL1 protein expression and reduced

DMPK in human DM1 cells.[27] These observations further support miRNA inhibition as a strategy

to restore MBNL1 function.

Beyond these molecular mechanisms, DM1 exhibits a highly variable clinical presentation

influenced by repeat instability and multisystem involvement. [28] Due to errors in DNA replication

and repair mechanisms, these repeats can expand during genetic transmissions, a phenomenon

known as genetic anticipation, leading to earlier onset and more severe symptoms in successive

generations. The length of the CTG repeat broadly correlates with disease severity, but somatic

and germline instability, as well as non-CTG interruptions in expanded alleles, contribute to the

variable clinical presentation of DM1.[29] Skeletal muscle alterations, including muscle atrophy and

myotonia, cause disabling symptoms. Muscle atrophy leads to progressive weakness, especially in

distal muscles, impairing mobility and dexterity. Myotonia, characterized by delayed muscle

relaxation, causes stiffness and difficulty in gripping or walking, further limiting physical function

and independence.[1] DM1 patients suffer from clinically relevant multisystem manifestations. In

the cardiac muscles, arrhythmias and conduction defects may lead to sudden cardiac death.

Progressive heart muscle degeneration, marked by fibrosis and fatty infiltration, impairs systolic

and diastolic function.[30] The smooth muscles in the gastrointestinal tract are also impacted, leading

to symptoms such as dysphagia, constipation, and other digestive issues.[31] Additionally,

respiratory muscles are weakened, which contributes to breathing difficulties and increases the

risk of respiratory failure.[32] Central nervous system effects cause features such as hypersomnia,

behavioral autism spectrum disorder, and cognitive deficits.[33,34] Despite several ongoing drug

development programs, [35] as yet no specific drug has obtained regulatory approval for DM1

patients.

AntimiRs are synthetic nucleic acid strands designed to bind to specific miRNA molecules,

thereby modulating downstream gene expression with precision. Although many closely related

splice-switching oligonucleotides and gapmers have reached clinical trials,[36] no antimiRs are

currently used in medical practice. Research in the medicinal chemistry of nucleic acids has

experienced substantial and rapid advances aimed at improving their pharmacokinetic (PK) and

pharmacodynamic (PD) properties. [37] For instance, nuclease resistance and enhanced plasma

protein binding (leading to reduced clearance) are markedly improved by substituting the natural

phosphodiester inter-nucleotide bonds with phosphorothioate (PS) linkages or incorporating

locked nucleic acids (LNAs).[38,39] Some of these features can also be achieved using nucleotides

with pentose modifications, including various 2'-modified sugars such as 2'-O-methyl (2'-O-Me),

2'-O-methoxyethyl (2'-MOE), and 2'-fluoro (2'-F) modifications.[37,40] However, incorporating

some of these modifications into candidate antisense oligonucleotide (ON) drugs can result in toxic

accumulation in the liver and kidneys due to nonspecific interactions of the ON backbone with

proteins, contributing to some degree of hepatic and renal damage.[41–43] Moreover, antisense ON

uptake by several tissues and cell types is poor, one of the most challenging being skeletal muscles,

as shown in patients with the first ON that reached clinical trials in DM1.[44,45] One active research

approach to solve these issues is to conjugate the ON to an antibody[46] or fragment of antibody for

cell receptor recognition or to conjugate to small biomolecules that can influence the

biodistribution of the molecule or its half-life *in vivo*.[47–49] Therapeutic ONs represent a rapidly

developing and expanding therapeutic platform for treating neuromuscular and other disorders,

resulting in nearly twenty approved ON-based therapies[37,50,51] for a variety of conditions, though

none yet for DM1.

We report a streamlined approach to developing antimiR-23b hits[24,52,53] into lead drugs for human

use, by addressing chemical optimization and ON conjugation in parallel and thus significantly

saving time and resources. Using the therapeutic index (T_{index}) to prioritize candidates with the

greatest therapeutic window, alongside established knowledge on nucleic acid medicine,[54] top *in*

vitro candidates underwent *in vivo* evaluation for dose-responsiveness, duration of activity,

toxicity, immune activation, and organ biodistribution, and the mechanism of action was

confirmed in rat and pig fibroblasts.

MATERIAL AND METHODS
144
145   **Study design**
146
147   The design encompasses antimiR-23b molecule optimization and testing both *in vitro* and

148 *in vivo* to assess efficacy, toxicity, and immune activation potential, to select a lead compound

149 with the highest potential for subsequent promotion to human use. For *in vitro* studies, antimiR

150 candidates were screened using various chemical modifications and conjugates. The screening

151 evaluated their EC_{50}, TC_{50}, and T_{index} in TDMs. The *in vivo* experiments used DMSXL and *HSA^{LR}*

mouse models. AntimiR candidates were administered to HSA^{LR} mice in different doses (3, 6, 100 mg/kg), and at 25 mg/kg to DMSXL mice, after which relative muscle strength and myotonia were evaluated. Molecularly, key readouts included quantification of miRNA and MBNL1 protein and transcript levels, and alternative exon inclusion levels in skeletal muscles. Toxicity and immune activation assays were also conducted using multiple antimiR doses. RPTEC-TERT1 cells were used to assess potential toxicities and cell proliferation, while human peripheral blood mononuclear cells (PBMCs) were utilized to measure cytokine release and complement activation. Additionally, transcriptome-wide RNA-seq and Sylamer analyses were performed to assess the global impact of miR-23b inhibition on gene expression and splicing biomarkers in patient-derived muscle cells.

Oligonucleotide synthesis

ON were provided by AXOLABS and resuspended in 1x PBS to prepare 5 mM stock solutions, stored at -20°C. 3´-oleyl-antimiRs were synthesized according to [55]. Tables S1–S3 include the sequence, modifications, length, concentrations tested, and TC_{50}, EC_{50}, E_{max} and T_{index} parameters of each tested ON.

Cell culture experimentation

Immortalized MyoD-inducible (doxycycline) DM1 and control fibroblasts were kindly provided by Dr. Furling (Institute of Myology, Paris) and were grown in Dulbecco's Modified Eagle Medium (DMEM) with 4.5 g/l glucose, 1% penicillin/streptomycin, and 10% FBS (Sigma, Saint Louis, Missouri).[56] Fibroblast trans-differentiation into myotubes was performed according to [24]. Trans-differentiation was induced at day 0, and test compounds were added to the cell medium at the different concentrations indicated 24 h later by lipofection with X-tremeGENE™ HP (Roche,

Basel, Switzerland) and were replaced with fresh differentiation medium 4 h afterward. Cells were

collected on day 4 in the differentiation medium and processed for protein extraction.

RPTEC/TERT1 cells (CHT-003-0002, EVERCYTE GmbH) were cultured in phenol red-free

DMEM/F12 medium (Gibco™ 11039021) supplemented with 1% penicillin/streptomycin, 3.5

µg/ml L-ascorbic acid (A4544), 25 ng/ml prostaglandin E1 (P5515), 8.65 ng/ml sodium selenite

(S5261), 25 ng/ml hydrocortisone (H0396), 10 ng/ml human EGF (E9644), 5 µg/ml human insulin

(I9278), 5 pM 3,3´,5-triiodo-L-thyronine (T6397) [all from Sigma-Aldrich], 100 µg/ml geneticin

(InvivoGen, ant-gn-5), 5 µg/ml human transferrin (Merck-Millipore, 616424), and 1% fetal bovine

serum (Gibco™ 10270106). Cells were maintained in 25 cm² flasks and passaged 1–2 times per

week using a 1:2 to 1:3 subculturing ratio, following the manufacturer's guidelines.

For ON toxicity assessment, RPTEC/TERT1 cells were seeded into 96-well plates (Greiner Bio-

One™, 655180) at 20,000 cells/well in the specified medium and grown to confluence. On day 0,

cells were treated with antimiRs prepared in PBS at final concentrations of 1, 10, 30, and 100 µM

in 50 µl volumes. A toxic ON, SGLT2-MOE-ON (66), served as a positive control at 20, 300, 400,

and 500 µM in 50 µl volumes. Final ON solutions were added to 150 µl of complete medium, with

PBS as a vehicle control. On day 6, the medium was collected, centrifuged at 10,000 rpm to remove

sediment, and the supernatants were stored at -80°C for EGF evaluation.

Primary myoblasts were grown on 0.1% gelatin-coated flasks in a proliferation medium containing

DMEM supplemented with 10% FBS, 22% M-199, PSF 1x, insulin 1.74 µM, L-glutamine 2 mM,

FGF 1.39 nM and EGF 0.135 mM. When cells reached 80–90% confluence, the medium was

substituted by differentiation media containing DMEM supplemented with 2% FBS, 22% M-199,

PSF 1x, insulin 1.74 μM and L-glutamine 2 mM. At this point, cells were transfected with X82107

at 100 nM using Ribocellin (BioCellChallenge, #RC1000) as a transfection reagent following the

manufacturer's instructions. Cell pellets were collected for DNA, RNA, and protein analyses 5

days after differentiation started when treatments were also transfected.

Rat embryo (*Rat2*, ATCC CRL-1764) and pig testis (ST, ATCC CRL-1746) fibroblasts were

grown in a complete medium (the same as for human fibroblasts) and seeded at 10^5 cells/ml in

petri plates. These cells were treated with X82108 at 50 nM and transfected with the same

procedure as DM1 and control fibroblasts.

RNA extraction, semi-quantitative RT-PCR and RT-qPCR

Total RNA from murine gt and qd muscle was isolated using the miRNeasy Mini Kit (Qiagen,

Hilden, Germany) according to the manufacturer's instructions. One microgram of RNA was

digested with DNase I and reverse-transcribed with SuperScript II (Invitrogen, Carlsbad,

California) using random hexanucleotides. For subsequent PCR reactions, 20 ng of cDNA was

used with GoTaq polymerase (Promega, Madison, Wisconsin). Specific primers were used to

analyze the alternative splicing of *Atp2a1* (OMIM*108730), *Nfix* (OMIM*164005), *Mbnl1*, and

Clcn1 (OMIM*118425) in mouse samples (both muscles). *Gapdh* (OMIM*138400) levels

established the endogenous reference using 0.2 ng of cDNA. PCR products were separated on a

2% agarose gel and quantified using ImageJ software (NIH, Bethesda, Maryland). The PSR index

per alternative exon was defined as the value: %EI minus $\bar{X}$%DEI, divided by $\bar{X}$%DEI minus

$\bar{X}$%HEI (EI: exon inclusion of each sample; DEI: disease exon inclusion; HEI: healthy exon

inclusion). Overall PSR was the average of the above PSR for the four quantified alternative

splicing events and the two muscles analyzed. The primer sequences and exons analyzed are

available in [24].

We used 1 ng of mouse tissue cDNA as a template for multiplex RT-qPCR using the QuantiFast

Probe PCR Kit reagent. Commercial TaqMan probes (Qiagen, Hilden, Germany) were used for

mouse (*Mbnl1* and *Mbnl2*; FAM-labeled probes) and reference (*Gapdh* MAX-labeled probe)

genes. Results were normalized to *Gapdh* endogenous ~~gene~~ expression. The sequences of the

probes and primers are included in Table S4.

MiRNA expression in muscle tissues was quantified using specific miRCURY™-locked nucleic

acid microRNA PCR primers (Qiagen, Hilden, Germany) according to the manufacturer's

instructions. Relative gene expression was normalized to U1 (YP00203909) and U6

(YP00203907) snRNAs. Expression levels were measured using a QuantStudio 5 Real-Time PCR

System (Applied Biosystems, Foster City, California). Expression relative to the endogenous gene

and control group was calculated using the $2^{-\Delta\Delta Ct}$ method.

Animal experimentation and antimiR administration

Mouse handling and experimental procedures for HSA^{LR} and FVB animals followed the European

law regarding laboratory animal care and experimentation (2003/65/C.E.) and were approved by

the regional Ministry for Agriculture (Conselleria de Agricultura, Generalitat Valenciana).

Homozygous transgenic HSA^{LR} (line 20 b) mice[57] were provided by Prof. C. Thornton (University

of Rochester Medical Center, Rochester, NY, USA). Experimental groups were FVB as wild-type

control, HSA^{LR} treated with PBS as negative control, and HSA^{LR} mice treated with each

experimental antimiR. The sample size was four mice per treatment group, twelve mice for PBS,

and eighteen mice for the FVB group. All groups were injected intravenously (tail vein) with 1×

PBS (vehicle) or the specific antimiR with a single dose of 3 or 6 mg/kg, except for the 100 mg/kg group, which received a single subcutaneous injection. At 5, 7 or 15 days after injection, the mice were sacrificed, and the tissues of interest were frozen in liquid nitrogen for molecular assays. Additionally, DMSXL mice (>99% C57BL/6 background) provided by Prof. G Gourdon (Sorbonne University, Paris, France), carrying over 1500 CTG repeats within the *DMPK* transgene,[58,59] were used to study the effects of antimiR treatment (authorization APAFIS#30275-2021030414389550). These mice were 3 weeks old at the start of the experiment and received subcutaneous injections of antimiR X821 at a dose of 25 mg/kg, administered twice per week for 4 weeks. Age-matched wild-type littermate mice served as controls. Following the treatment regimen, the gastrocnemius and tibialis muscles were collected, rapidly frozen in liquid nitrogen, and stored at -80°C for subsequent molecular analyses.

Electromyography studies

Electromyography was performed before treatment and at the time of sacrifice under general anesthesia, as previously described.[22] Myotonia was assessed from a total of 10 needle insertions (5 per quadriceps). The overall myotonia score obtained from these insertions was then normalized to a 5-point scale, defined as follows: 0, no myotonia; 1, occasional myotonic discharges present in ≤50% of the insertions; 2, myotonic discharges present in >50% of the insertions; 3, myotonic discharges in nearly all insertions; and 4, myotonic discharges in all insertions.

Forelimb grip strength test

Forelimb grip strength was measured using a grip strength meter (BIO-GS3; Bioseb, Pinellas Park, Florida). Peak pull strength (g) was recorded on a digital strength transducer as the mouse released its grip. The transducer was reset to 0 g after each measurement, and three measurements were

taken at 30 s intervals. Body weight was recorded simultaneously, and normalized grip strength

was calculated by dividing average strength by body weight. Experiments were conducted with

coded animals to prevent bias.

ELISA determinations

Custom hybridization-based ELISA determinations of antimiR conjugate concentrations in gt and

qd from treated HSA^{LR} mice were performed as described in [60] using the following PS probe: miR-

23b (5'->3') [BIO] ATCACATT$_o$G$_o$C$_o$C$_o$A$_o$G$_o$GGATTACC [DIG] double-labeled with

digoxigenin (DIG) and biotin (BIO). A control sample (HSA^{LR} treated with PBS) was used to

subtract the background.

Extracellular EGF assay

To determine extracellular EGF, Human EGF ELISA Kits (Invitrogen™ KHG0062) were used,

and EGF concentration was determined according to manufacturer's instructions. The

experimental protocol was adapted from [61].

***In vitro* toxicity assay**

The French Blood Donors Organization (EFS, Etablissement Français du Sang) provided human

serum from healthy volunteers. Complement activation, coagulation, and platelet activation were

performed by SQY Therapeutics (France) in human blood samples. Test compounds were

incubated with human serum, and complement activation was evaluated using commercial ELISA

measurement of human C3a. The extrinsic and intrinsic coagulation pathways were evaluated in

human plasma incubated with test compounds. PT and aPTT were measured on a semi-automated

START max coagulometer (Stago). Flow cytometry assays evaluated human platelet activation by

test compounds by detecting surface antigens (usually glycoproteins) expressed during platelet

activation. Assays with PBMCs were performed by Axolabs GmbH (Germany).

RNAseq assay

Total RNA was extracted from DM1 and control immortalized MyoD-converted

fibroblasts [56] 72 h after transfection with 200 nM X82108 using X-tremeGENE HP (Roche)

following culture in DMEM GlutaMAX with 10% FBS. RNA was purified with the RNeasy Mini

Kit (QIAGEN). RNA sequencing was performed by Novogene using Illumina platforms. Poly(A)+

RNA was isolated using oligo-dT magnetic beads, fragmented, and converted to first- and second-

strand cDNA. Libraries were prepared following standard end repair, A-tailing, adaptor ligation,

size selection, and PCR amplification, and were assessed by Qubit, qPCR, and Bioanalyzer before

pooling and sequencing. Paired-end reads were aligned to the human reference genome

(GRCh38.p14) using STAR (v2.7.11b) with splicing-aware parameters and GeneCounts enabled.

BAM files were processed with SAMtools, and gene-level quantification was performed with

featureCounts.

Sylamer (v18-131) was used to assess enrichment of the miR-23b seed motif (AATGTGA)

across differentially expressed genes (adjusted $p < 0.05$), ranked by $\log_2$ fold change.

Corresponding 3′UTR sequences were retrieved from hg38 using

TxDb.Hsapiens.UCSC.hg38.knownGene and BSgenome.Hsapiens.UCSC.hg38, ordered

according to gene ranking - with positive values corresponding to upregulated genes and negative

values to downregulated genes-, and analyzed using a window growth parameter of 10. Output

tables were used to visualize motif representation across bins and to overlay statistical thresholds

(nominal $\pm p = 0.05$ and $\pm E = 0.01$).

Alternative splicing was quantified using rMATS (v4.1.2) with 150-nt unstranded read

settings. Analyses were restricted to a validated panel of 22 DM1-relevant splicing events

described in Provenzano *et al.*[62] Percent Spliced-In (PSI) values were calculated as the fraction of

exon inclusion isoform reads relative to all exclusion and inclusion isoform counts. Percent

Splicing Rescue (PSR) for each replicate x and event y was computed as: (PSIx, y – PSIMedian

Control, y)/(PSIMedian DM1, y – PSIMedian Control, y).

RNA-seq data are publicly available with accession number S-BSST2306 in the BioStudies

database.

Statistical analyses

TC_{50} values were calculated using non-linear least-squares regression. For molecular and

functional parameters, one-way ANOVA was performed where applicable, preceded by the

Shapiro-Wilk normality test. Normal samples were Brown-Forsythe corrected if required, while

non-normal samples were analyzed using Kruskal-Wallis without Dunn correction. Differences

relative to control samples were assessed with two-tailed Student's t-test ($\alpha = 0.05$), applying

Welch's correction as needed. Sample sizes (n) are indicated in each figure.

The values obtained for the HSA^{LR} mice are represented (Figure 4C) as the recovery index (RI),

which measures the proximity of the different parameter values obtained with treated HSA^{LR} mice

to those of FVB controls. This RI was obtained for the different parameters (MBNL1 protein,

Mbnl1 and Mbnl2 and miR-23b or -218 expression level, PSR, *Mbnl1* ex5 inclusion recovery, and

functional recovery) of each mouse after treatment according to the following formula: value %

MT minus X% MNT, divided by X% MH minus X% MNT [where MT is the value of each mice

treated (PBS or oligonucleotide), MNT is HSA^{LR} mice treated with PBS, and MH is healthy mice

value (FVB)]. These values range from 0 to 100, where 0 are untreated mice (H*SA*^{LR}-PBS) and

100 are healthy mice (FVB).

327
328 **RESULTS**
329

In vitro **screening of antimiR-23b ON**

In our search for the most effective and safest antimiR-23b, we performed *in vitro* screenings in

DM1 fibroblasts transdifferentiated into myotubes (TDMs). The first screening addressed specific

miR-23b-targeting sequences, length, and chemical modifications. In TDMs transfected with each

candidate antimiR variant, we established the T_{index} as E_{max}* (TC_{50}/EC_{50}), which integrates the

maximum activity (E_{max}), median toxic concentration (TC_{50}), and median effective concentration

(EC_{50}) into a single metric. EC_{50} and E_{max} values were obtained through a Quantitative Dot Blot

assay[63] for MBNL1 protein quantification to measure antimiR activity, and TC_{50} was determined

using an MTT/MTS toxicity assay. In all cases, we included for comparison purposes the antimiR-

23b molecule used in our previous proof-of-concept studies, here called X820,[26,52] with the

sequence 5'-GGUAAUCCCUGGCAAUGUGAU-3', 2'-OMe chemistry, PS linkages at end-

proximal nucleotides, and conjugated to cholesterol at the 3' end, and its unconjugated version

(X82--).

A panel of 21 unconjugated antimiR-23b molecules containing distinct sequence and chemistry

variants were synthesized and tested for their T_{index} *in vitro* (Figure 1A,B and ~~Supp~~ Table S1). The

results identified X821 as the most promising antimiR-23b candidate, with a T_{index} of 4018.3. This

ON demonstrated a 423-fold improvement compared to X82-- (T_{index}: 9.5), our unconjugated

benchmark, and a 5.9-fold improvement over the second-ranked X822 (T_{index}: 686.7). X821

exhibited the best toxicity profile *in vitro*, requiring around 16 times higher concentration than

X820 to reach TC_{50} (Figure 1B and Table S1). Comparing unconjugated versions among

themselves, 18 of the 21 antimiRs evaluated had a T_{index} higher than X82--, while three worsened

the parameter.

352   Effects of X821 on MBNL1 levels, alternative splicing, and functional muscle defects in two

353   DM1 mouse models

Prospective lead candidate X821 was evaluated *in vivo* subcutaneously to assess its therapeutic

activity in a well-established adult-onset DM1 murine model, the HSA^{LR} mouse, which expresses

expanded 250 CUG repeat tracts in human skeletal actin (HSA) transcripts. [57] Additionally, we

examined the effectiveness of the antimiR approach in DMSXL mice, which carry a complete

DMPK with more than 1,500 CTG repeats, providing a contrast to the muscle-specific HSA^{LR}

strain and enabling evaluation of antimiR activity in the context of a full-length human transgene.

[58] These complementary models allow for a comprehensive evaluation of the therapeutic efficacy

of X821.

In the HSA^{LR} mouse, treatment with 100 mg/kg of X821 significantly increased MBNL1 protein

levels in muscle without altering blood parameters (Figure S1A,B). Regarding C.P.K. levels,

although they appear higher in the treated group, the difference compared to non-treated animals

is not statistically significant. It also rescued *Nfix* ex 7, *Atp2a1* ex 22, and *Clcn1* ex 7a mis-splicing

in both gastrocnemius (gt) and quadriceps (qd) muscles (Figure 1C,D, S1C,D), leading ultimately

to a significantly reduced myotonia grade and increased grip strength in treated mice (Figure 1E).

In the DMSXL mouse, X821 also increased MBNL1 protein levels in the tibialis anterior (TA)

and gt muscles and rescued *Numa1* (OMIM*164009) exon 16 inclusion[64] in TA (Figure 1F,G).

Molecular rescues were sufficient to produce a significant functional improvement in grip strength

in the rear legs 2 weeks after the last injection (Figure 1H). Taken together, these findings

demonstrate that X821 effectively addressed key molecular defects, such as MBNL1 protein

upregulation and alternative splicing correction in both mice, while also improving functional

outcomes related to myotonia and muscle strength. Even though X821 effectively targeted miR-

23b, the effects observed on muscle function did not surpass those achieved with X820 at a lower

concentration.[52] This lack of correlation between *in vitro* and *in vivo* activity underscored the need

for subsequent optimization of X821 to enhance its therapeutic potential.

***In vitro* screening of lipidic conjugates to improve the therapeutic index**

A second *in vitro* screening addressed conjugation to various moieties to improve potency and PK

properties.[65,66] Conjugating hydrophobic molecules to antisense ON and employing various

strategies to transport them from blood to tissue might improve their transport to muscle tissues,

leading to enhanced biodistribution and functional uptake in muscle cells. [67,68] Fatty acid

conjugation, as yet not attempted with anti-miR-type molecules, was tested here to assess the

potential effects of various ligands. Additional antimiR molecules were generated by synthetically

conjugating different hydrophobic moieties (Figure 2A,B, and Table S2). To isolate the impact of

the moiety, the nucleotide sequence was kept identical to the original X82--, and experiments

included this unconjugated version. In this screening evaluating conjugates in the 5´ end of the

ONs, oleic acid (oleyl moiety; Ol) conjugation ranked first (X8200; T_{index}= 1101.4), followed by

the linoleyl-antimiR (X8201; T_{index}= 400.9). These two conjugates had the greatest ability to

reduce cell toxicity (Figure 2B), respectively showing a 16.10- and 5.84-fold superiority to the

original X820 hit, and 155.13- and 56.35-fold enhancement compared to the naked version of the

antimiR. Other lipid conjugations also improved T_{index} compared to the unconjugated benchmark

X82-- (tocopherol, X8202; cholesterol, X8203), while conjugation to palmitic, elaidic, and stearic

acids was similar or lower in terms of T_{index}.

In vivo screening of antimiR lead candidates

We hypothesized that the positive impact of Ol on T_{index} could enhance the high performance of

antimiR X821, aiming for synergistic or additive effects. Two X821 variants (X821-7 and X821-

8) were designed, retaining the same sequence but including known chemical modifications to

improve *in vivo* druggability, such as methylated cytosines designed to enhance stability (Table

S3), improve RNA target interaction, and reduce immunogenicity *in vivo* [69]. Both X821 variants

differed only slightly in terms of LNA/MOE content, with X821-8 having a higher Tm than X821-

7. X821-7 served as a benchmark to evaluate the effect of Ol conjugation on *in vivo* rescue

performance in *HSA^{LR}* mice. Therapeutic potential was assessed by comparing the rescue of key

DM1 molecular and functional parameters after a single low-dose (3 mg/kg) IV injection of

unconjugated X821-7 or ON with Ol at the 5´ (5´X82107) or 3´ (X82107) ends (Figure 2C).

At 5 days after injection, muscle tissue analysis revealed that both X82107 conjugate variants

consistently increased MBNL1 protein levels by 1.5-fold compared to untreated animals, unlike

unconjugated X821-7 (Figure 2D). For X82107, this translated into a significant rise in the overall

percentage splicing recovery (PSR; Figure 2E and S2A), generated from a selected panel of four

mis-spliced genes in this model (*Atp2a1, Clcn1, Nfix,* and *Mbnl1*). Increases at the *Mbnl1* and

Mbnl2 mRNA level were significant and similar for all three antimiR variants, whereas the

reduction in detected miR-23b was robust for X82107 (Figure S2B-D). ELISA determinations revealed that the oleic-conjugated antimiR X82107 achieved higher concentrations in the skeletal muscles, which are tissues relatively resistant to ON uptake, compared to the unconjugated version (Figure 2F). Also, the levels detected in the kidney and liver were higher with X82107 compared to the naked version, as was expected for this type of compound, as normally they are excreted by these organs. Functionally, normalized muscle strength improved significantly and similarly for all three antimiR variants, while myotonia reduction was higher for X82107 (Figure S2E-G). Taken together, the Ol-conjugated versions at either end of the molecule were more effective than the unconjugated version.

Similar to X821-7, we generated unconjugated X821-8, featuring a different set of chemical modifications (Table S3) and contrasted the effects of Ol (X82108) and palmitoyl [Pal] (X82148)-conjugation at the 3′-end of the molecule (Figure 3A), using Pal as a benchmark conjugate for its reported action as an effective enhancer of ON effects in muscle.[67] AntimiRs were administered as before, and in a similar way to X821-7, both X82108 and X82148 conjugates outperformed the unconjugated version, and in comparison to Pal, Ol moiety notably increased MBNL1 protein levels, significantly improving overall PSR in skeletal muscles (Figure 3B,C; S3A). The increase at the *Mbnl1* mRNA level and reduction in detected miR-23b were significant and similar for all three X82108 variants, whereas *Mbnl2* transcript levels significantly increased only after X82108 treatment (Figure S3B-D). Accordingly, ELISA determinations revealed that X82108 achieved higher concentrations in the skeletal muscle than the unconjugated version (Figure 3D). Functionally, X82108-treated mice showed a significant increase in normalized strength and a robust reduction in myotonia grade, which dropped from 3 to less than 1 (Figure S3E–G). Overall,

435   from these *in vivo* experiments, X82108 and X82107 were selected as new prospective lead

436   candidates for further assessment of preclinical features.

Boosting PD and therapeutic potential of antimiRs via oleic acid conjugation

Previous results support that Ol conjugation improves the PD and therapeutic impact of the

antimiR candidates X82107 and X82108 *in vivo*. We hypothesized that these benefits could be

sequence independent and would also apply to other antimiRs. To test this hypothesis we designed

miR-218 antimiRs, as this miRNA had previously been shown to upregulate MBNL1 in human

cells and mouse models and could also have therapeutic potential in DM.[24,26,27] This assessment

involved a 5-day treatment of HSA^{LR} mice with 3 mg/kg of an Ol-conjugated variant (X83104)

compared to the unconjugated antimiR-218 control (X831-4) with an identical chemical structure

(Figure 4A,B). The spider graphs in Figure 4C illustrate that Ol-conjugated antimiRs against miR-

23b or miR-218 consistently outperformed their unconjugated counterparts across all evaluated

parameters (data from FVB control mice are included in dataset S1). These conjugates showed

greater efficiency in reducing target miRNA levels, indicating enhanced molecular targeting.

Additionally, treatment with the conjugated antimiRs resulted in a significant increase in MBNL1

protein levels and *Mbnl1* and *Mbnl2* transcript expression, underscoring their ability to restore

critical molecular markers associated with DM1. Splicing recovery, including improved *Mbnl1*

exon 5 inclusion, was notably higher in mice treated with the conjugates, and conjugated variants

also provided the most robust recovery of functional parameters, suggesting that the Ol

modification enhances therapeutic potential. The overall performance of conjugated antimiRs

highlights the advantages of Ol conjugation for improving molecular and functional outcomes in

this disease model.

Sustained therapeutic effects of conjugated oligos

To build on the results obtained for prospective lead candidates X82107 and X82108 from the 5-day single-dose experiments, we tested the activity duration of these 3´Ol-conjugated molecules after the same 3 mg/kg IV dose by evaluating tissue response at 5, 7, and 15 days post-injection (Figure 5A, dataset S1). Throughout the study period, both antimiRs exhibited a sustained increase in Mbnl1 protein levels compared to phosphate-buffered saline (PBS)-injected controls, with the peak effect observed at 5 days post-treatment (Figure 5B). Notably, X82108 consistently outperformed X82107 in this parameter at each time point. These findings are consistent with a significant overall PSR on days 5 and 7, during which protein levels were still substantially higher than in disease controls, and X82108 provided notably better splicing rescue than X82107 (Figure 5C, dataset S1). Both antimiRs demonstrated a comparable peak in normalized grip strength at 5 and 7 days post-injection, nearing normal values before declining to PBS-injected levels by day 15. In the context of myotonia, X82108 achieved a greater reduction than X82107 at day 5, with both converging at later time points (Figure 5D,E, dataset S1). Taken together, these results suggest that the activity duration for both lead candidates ranged between 7 and 15 days, with very similar outcomes observed for the two compounds.

Dose-response efficacy of X82107 and X82108

Upon demonstrating the promising effects of Ol conjugation on X82107 and X82108 variants, we further investigated the efficacy of these molecules at a higher dose (6 mg/Kg), 15 days after injection, when the effects of the 3 mg/Kg injection were already substantially reduced. A single intravenous injection of 6 mg/kg of each variant was administered to HSA^{LR} mice (Figure 6A,B,

dataset S1), and several parameters were evaluated. Specifically, both antimiRs led to significantly

higher levels of MBNL protein in the skeletal muscles qd and gt compared to PBS-treated animals.

However, the increase observed with X82108 was notably higher than with X82107 (Figure 6C,

dataset S1). This trend was also reflected in the reduction of miR-23b levels, which decreased by

up to 80% in X82108-treated animals (dataset S1), and in a higher PSR achieved by this antimiR

(Figure 6D, dataset S1). These molecular changes translated into functional improvements, with

treated mice displaying similarly increased front leg grip strength, normalized to body weight

(Figure 6E, dataset S1). Both candidates demonstrated a significant increase in PSR and

normalized strength 15 days after injection at 6 mg/kg (Figure 6F), confirming prolonged activity

compared to results at the lower dose of 3 mg/kg conjugated with oleic acid or cholesterol.[52]

Overall, both molecules exhibited a dose-dependent duration of effect, further supporting their

potential as promising therapeutic candidates.

***In vitro* assessment of renal and immune activation profiles of X82107 and X82108**

Potential renal toxicity was assessed using an *in vitro* model with RPTEC/TERT1 cells, human

renal proximal tubular epithelial cells modified with telomerase reverse transcriptase for long-term

studies. RPTEC/TERT1 cells are ideal for nephrotoxicity, renal metabolism, and drug absorption

studies, as they retain primary renal cell functions, including electrolyte transport, toxin response,

and epithelial barrier integrity;[70] furthermore, EGFR activation is a pivotal mediator for renal

fibrosis and has a major role in activating pathways that mediate podocyte injury and loss in

diabetic nephropathy.[71] After treatment with antimiRs, extracellular epidermal growth factor

(EGF) levels rose significantly at the highest dose of 100 μM for antimiR X82108 compared to

PBS-treated controls (Figure 7A), while a moderate but non-significant increase was observed for

X82107 at the same dose (Figure 7B). No substantial changes in EGF levels were detected at 1,

10, or 30 µM for either compound. Parallel MTS assays revealed significant cytotoxicity at 100

µM for both compounds, with cell viability inhibition exceeding 80% (Figure 7C,D). This aligns

with the accumulation of extracellular EGF, as non-viable cells have reduced EGF uptake capacity.

In summary, significant effects on extracellular EGF levels and cell viability were observed only

at 100 µM, a concentration typically associated with toxicity in these assays. At lower doses, both

antimiRs displayed a favorable safety profile.

We also conducted *in vitro* tests in human blood to assess the potential immunogenic modulation

and safety of lead candidates X82107 and X82108. Neither compound activated the complement

system, as evidenced by the unchanged C3a levels observed in serum and plasma samples, even

at a high dose of 300 µM (Figure 7E). Given that different chemical modifications in

oligonucleotide-based drugs can significantly alter the pharmacokinetic and toxicological profiles

of antisense ON, including potential effects on coagulation times,[72] we measured these parameters

in normal human plasma. In coagulation assays, both X82107 and X82108 produced a significantly

prolonged activated partial thromboplastin time (aPTT, which measures the intrinsic coagulation

pathway) at 30 µM and higher doses in normal human plasma, while the prothrombin time (PT,

extrinsic pathway) remained unchanged for both molecules compared to the PBS control (Figure

7F). Platelet count and activation levels remained similar to control across all doses for both

compounds (Figure 7G). An additional assessment of cytokine release in human peripheral blood

mononuclear cells (PBMCs) showed no significant cytokine increase following X82108 treatment

(Figure 7H). Overall, our preclinical data support a favorable toxicological profile for lead

compounds X82107 and X82108, with no significant safety-related red flags.

AntimiR-23b-mediated MBNL upregulation is conserved in different mammal species

Human, rat (*Rattus norvegicus*) and pig (*Sus scrofa*) mature miR-23b sequences are identical.

Given this conservation, we tested the hypothesis that our lead antimiRs could similarly antagonize

miR-23b function upon transfection and upregulate MBNL1 protein expression in cell lines from

these species. Transfection of X82108 in fibroblasts of these two animals led to a significant

increase in *Mbnl1* levels and a dramatic reduction in miR-23b detection (Figure 8A,B).

Importantly, these results confirm that miR-23b inhibition produces *Mbnl1* upregulation in

different relevant preclinical mammalian species.

To rule out cell line bias, X82107 and X82108 were tested on patient-derived primary myoblasts

after 10 days of differentiation. This has already been reported for X82108 [27] and is shown here

for X82107 in two cell lines from that same study (Figure 8C–E). Briefly, treatment of primary

human myoblast cell lines MP09-57 (carrying 702 CTG repeats in *DMPK*) and MP10-31 (carrying

117 repeats) with X82107 resulted in a marked reduction of miR-23b detection (Figure 8C).

Overexpression of miR-23b and reduction in the presence of lead antimiRs correlated well with a

significant increase of MBNL1 protein levels that approached or surpassed those observed in

healthy control cells (MP-49F; Figure 8D,E). Results from Figure 8 and [27] demonstrate that lead

antimiRs can significantly upregulate MBNL1 levels in diverse genetic backgrounds and CTG

repeat expansion sizes. Additionally, treatment with antimiR-23b significantly reduced the number

of foci in these cell lines at different time points, as well as increased MBNL1 protein in a diffused,

free-form state in the nucleus. [27]

Transcriptomic analysis of miR-23b inhibition: miR-23b targets analysis and splicing

correction

To further characterize the molecular consequences of miR-23b inhibition, we evaluated

transcriptomic alterations in immortalized DM1 myotubes, where miR-23b is markedly

upregulated (Figure S4), consistent with observations in HSA^{LR} mice [25] and primary patient-

derived myotubes (Figure 8C and [27]). Myotubes were treated with 200 nM antimiR X82108 for 3

days and subjected to total RNA-seq. Sylamer analysis [73] was performed to determine whether

miR-23b seed–complementary sequences were enriched or depleted among differentially

expressed genes (DEGs), highlighting the miRNA-driven regulatory effects. Moreover, we

additionally evaluated the impact of miR-23b inhibition on the transcriptomic biomarker

Composite Alternative Splicing Index (CASI-22), a robust clinical biomarker consisting of 22

DM1-specific splicing events strongly correlated with muscle function and widely used to evaluate

disease-modifying treatments [62].

Genome-wide Sylamer analysis revealed a significant overrepresentation of the miR-23b-3p seed

motif (AATGTGA) among the downregulated mRNAs in DM1 compared with controls (Figure

S5A and Table S5), consistent with the repressive activity of elevated miR-23b levels. Following

antimiR treatment, this enrichment was found only among upregulated genes (Figure S5B),

indicating selective derepression of endogenous miR-23b targets. These results confirm that miR-

23b targets are repressed in the disease condition and antimiR X82108 specifically blocks miR-

23b activity, leading to increased expression of its repressed targets in the DM1 cellular context.

Given that RNA mis-splicing is the core molecular hallmark of DM1, we assessed whether miR-

23b inhibition could ameliorate splicing defects using the CASI-22 biomarker panel. Of the 17

CASI events detectable in the cells, 11 were significantly altered, 3 exhibited borderline significant

changes, and 3 remained unchanged relative to controls. Notably, antimiR X82108 rescued the

overall PSR up to 35% and significantly improved 80% of the splicing events altered, with 20%

of them fully rescued (Figure S5C, Figure 8F). Many of the corrected events correspond to splicing

alterations closely associated with muscle-strength decline in DM1—most notably *ATP2A1*,

DMD, *NFIX*, *OPA1*, and the autoregulatory exon of *MBNL1*—underscoring their functional and

clinical relevance. [62]

Collectively, these transcriptomic and splicing analyses demonstrate that antimiR X82108 exerts

a precise and therapeutically meaningful molecular effect, derepressing miR-23b targets, and

improving key DM1 biomarkers implicated in muscle dysfunction.

DISCUSSION

576
We aimed to enhance the therapeutic potential of miR-23b inhibition in DM1 by developing

optimized antimiRs. Inhibiting miR-23b has been shown to boost MBNL1 levels and counteract

critical DM1 molecular defects. Given its elevated expression in DM1,[27] we hypothesized that

reducing miR-23b activity could mitigate disease effects. Starting with the non-optimized antimiR-

23b molecule X820 (antagomiR-23b),[52] we employed a two-step *in vitro* and *in vivo* screening

strategy to refine miR-23b-targeting sequences, chemical modifications, and lipid conjugations,

optimizing activity and toxicity in DM1 muscle cells. This approach combined the best features

from each screening, yielding advanced candidates. Animal experiments confirmed the target

engagement of these optimized molecules, validating the use of cell-based assays to streamline *in*

vivo testing.[74]

The PK/PD characteristics of ON drugs are strongly influenced by their chemical modifications

and specific properties of any associated conjugates. [67,75] To be effective in DM1, therapy must

reach skeletal muscles, which can constitute up to approximately 40% of the total body mass in

average healthy adults. However, the first clinical trial evaluating a therapeutic ON targeting

DMPK transcripts in DM1 was discontinued due to insufficient drug levels in muscle tissue, even

at the highest doses tested.[44] This underscores the inherent delivery challenges faced in treating

genetic myopathies. Most ONs approved for human use are administered systemically by

intravenous or subcutaneous injections to achieve broad distribution throughout the body. ON

plasma concentration peaks immediately after administration, followed by rapid distribution to

major body tissues within hours, except for the central nervous system.[47]

Receptor-specific delivery has been addressed by conjugating *DMPK*-degrading ONs to

monoclonal antibodies or Fab fragments recognizing the transferrin receptor, aimed at enhancing

muscle tissue uptake.[35] The mechanism by which Ol conjugation may confer desirable properties

to an ON has not been formally elucidated, but an intriguing hypothesis is that lipids can facilitate

either fusion with or disruption of the endosomal membrane, allowing the antisense ON to escape

into the cytoplasm.[76,77] ONs conjugated with fatty acids longer than C16 are known to form self-

assembled vesicular structures, which may further enhance their intracellular delivery. The

interaction of these fatty acid chains with mouse and human serum albumin (MSA/HSA) results

in the formation of stable adducts, with a near-linear correlation between fatty acid-ON

hydrophobicity and binding strength to albumin.[67,68] This enhanced albumin binding likely

prolongs circulation time, improves systemic stability, and facilitates efficient transport across

endothelial barriers to reach target tissues, including skeletal muscle. These properties may

collectively contribute to the superior therapeutic potential observed with Ol-conjugated antimiRs.

Importantly, while we compared unconjugated and conjugated ONs at the same dose in mg/kg, the

higher molecular weight of the conjugated ONs means that fewer molecules were administered

relative to their unconjugated counterparts. As a result, the observed increase in potency may be

slightly underestimated, further underscoring the enhanced efficacy of the Ol-conjugated ONs.

Safety is the primary concern in early drug development for human use, so we prioritized ON

candidates that demonstrated the greatest therapeutic window. This prioritization strategy enabled

efficient progression from initial hits to lead candidates. Interestingly, significant toxicity was

observed only at very high doses of antimiRs *in vitro* (e.g., 30–100 µM) for most of the compounds

tested. This yielded remarkable (over 100-fold) differences between the TC_{50} and the EC_{50}

concentrations.

Our results indicate that lead compounds X82107 and X82108 do not activate human platelets or

increase C3a in serum, with X82108 showing a slight, non-clinically meaningful prolongation in

coagulation time at high doses. Overall, the compounds exhibit a safe toxicological profile, without

adverse effects on coagulation or platelet activation. Although microRNAs regulate numerous

biological pathways, their inhibition does not necessarily lead to widespread systemic effects.

Functional knockout and CRISPR/Cas9 studies have shown that deletion of miR-23b or the miR-

23b/27b/24-1 cluster in mice produces only mild phenotypes, such as a minor increase in pain

sensitivity in neuropathic models or altered glucose homeostasis under a high-fat diet, without

major developmental or systemic defects. [78,79] These findings, together with reports identifying

miR-23b inhibition as therapeutically beneficial in cancer, support a favorable safety profile for

miR-23b inhibition. [80]

Since miRNAs are short and highly conserved regulatory molecules, the same antimiR molecule

can usually be tested in several mammalian animal models during regulatory preclinical steps.[81]

In this case, after obtaining solid results in mice, we moved on to test our leading candidates in

normal rat and pig fibroblasts. This paves the way for future use of minipigs as a large animal

model in preclinical toxicology testing of antimiRs, given the numerous physiological similarities

with humans, especially in cardiac studies.[82] Notably, these experiments confirmed that the

antimiR mechanism of action was conserved across mammalian species, even without expressing

toxic CTG repeat expansions. Finally, the robust increase in MBNL1 levels in primary myoblast

derived from two DM1 patients upon X82107 or X82108[27] treatment demonstrated the

effectiveness of these antimiRs across different human genetic backgrounds affected by DM1.

Moreover, the new transcriptomic analyses further support the specificity and therapeutic

relevance of miR-23b inhibition, showing selective derepression of endogenous targets and partial

correction of key DM1-associated splicing defects, thereby complementing our previously

published transcriptomic study in eight primary DM1 myotube lines treated with antimiR-23b [27].

Studies with the original antagomiR-23b molecule determined an optimal dose of 12.5 mg/kg to

significantly increase MBNL1 levels *in vivo.*[24] In comparison, the new lead candidates exhibited

lower toxicity and higher potency, achieving the same effect on MBNL1 levels 5 days after

administration with an approximately fourfold lower dose (3 mg/kg) and sustaining this level of

MBNL1 upregulation for at least 15 days with approximately half the dose (6 mg/kg). Notably,

this enhanced potency also translated into improved alternative splicing correction, as reflected by

a significantly higher overall PSR at both 3 mg/kg and 6 mg/kg doses, with the strongest splicing

rescue observed at the peak of MBNL1 upregulation (Figure 5B,C).

While our study demonstrates the efficacy and safety of X82107 and X82108 in preclinical models,

several limitations must be considered. First, although we observed robust MBNL1 upregulation

and splicing correction, further studies are needed to evaluate whether these molecular

improvements translate into sustained functional benefits beyond muscle strength and myotonia

reduction. Additionally, while *in vivo* biodistribution analyses confirmed efficient skeletal muscle

uptake, the extent of off-target effects in other tissues was not fully characterized by these

experiments. Given the high conservation of miR-23b across species, the potential for unintended

modulation of physiological pathways outside DM1-affected tissues was carefully examined in

additional studies. Another key limitation is the absence of long-term toxicity data from these

experiments, particularly regarding immune activation and hepatic or renal clearance after

repeated dosing, which require further investigation to dose humans on a chronic basis.

To bridge the gap toward clinical application, the next steps should be to include large-animal

studies to confirm the efficacy and pharmacokinetics of these lead compounds in models with

greater physiological similarity to humans, such as minipigs and non-human primates.

Additionally, regulatory preclinical studies require detailed dose-escalation experiments to

establish the maximum tolerated dose and long-term safety, particularly in the context of systemic

administration. As ON-based therapies for neuromuscular diseases continue to evolve,

complementary formulation strategies may further enhance the bioavailability and T_{index} of these

compounds. Ultimately, clinical trial design will incorporate biomarker-driven endpoints,

including MBNL1 target engagement and splicing correction, to facilitate a rigorous evaluation of

therapeutic efficacy in DM1 patients.

In conclusion, our approach combining *in vitro* screening, rational design, and *in vivo* validation

efficiently identifies potent and safe candidates. This strategy enhances therapeutic development

in the antimiR field and advances DM1 treatments by optimizing delivery and reducing toxicity.

Acknowledgments:

We thank Inmaculada Noguera for veterinary assistance at the University of Valencia SCSIE

animal facility. Part of the equipment employed in this work has been funded by Generalitat

Valenciana and co-financed with ERDF funds (OP ERDF of Comunitat Valenciana 2014-2020).

Antibody MB1a (4A8) was provided by the MDA Monoclonal Antibody Resources. This work

was supported by "la Caixa" Banking Foundation grant HR17-00268 (RA, AL-M, GG),

Generalitat Valenciana grants PROMETEO/2020/081 and CIPROM/2023/22 (RA), Instituto de

Salud Carlos III grant DTS19/0128 (RA), Generalitat Valenciana predoctoral grant

FDEGENT/2020/011 (IG-M), Torres Quevedo post-doctoral fellowship PTQ2020-011110 (EC-

H), CDTI NEOTEC grant SNEO-20201136 (BL), GVA-IVACE grant IMIDTA/2021/65 (BL),

Talent Promotion Program-Line 3 of GVA-AVI grant INNTA3/2023/16 (DP-L), and Instituto de

Salud Carlos III grant PI21/00557 (NN-G and AL-M). Illustrations were created with BioRender.

Author contributions:

693
694   Conceptualization: RA, EC-H, BL.

695   Methodology: IG-M, MAV, NM, MC-S, NB, AD-M.

696   Validation: EC-H, AG-R, DP-L, NB.

697   Formal analysis: IG-M, EC-H, MC-S, AG-R, DP-L, AC-R, MCH, AD-M.

698   Investigation: IG-M, EC-H, MC-S, AG-R, DP-L, AC-R, MCH, AH-L, AG-B, NM,

699   MD.

700   Resources: RA, NN-G, AL-M, GG.

701   Visualization: IG-M, EC-H, AD-M.

702   Funding acquisition: RA, BL, AL-M, GG.

703   Project administration: RA, BL, AL-C.

704   Supervision: RA, EC-H, GG, AL-M.

705   Writing – original draft: RA, AL-C.

706   Writing – review & editing: RA, IG-M, EC-H, AL-C.

708 **Declaration of interests:**

709
Beatriz Llamusi and Ruben Artero are founders and shareholders, and CSO and scientific advisor

of Arthex Biotech, respectively. B.LL and E.C-H are employees of Arthex Biotech. B.LL, E.C-H

and R.A, are co-inventors in patents PCT/EP2017/073685 (Modulation of microRNAs against

myotonic dystrophy type 1 and antagonists of microRNAs therefor) and EP22382493

(Oligonucleotides conjugated to oleic acid and uses thereof), and IG-M also in the last one,

currently licensed to Arthex Biotech. The remaining authors declare no competing interests.

Data and code availability:

All materials used in this study are available upon reasonable request. AntimiRs will

require a material transfer agreement. All data are available in the main text or the supplementary

materials. RNA-seq data are publicly available with accession number S-BSST2306 in the

BioStudies database.

Keywords

Antisense oligonucleotides; MBNL proteins; myotonic dystrophy; oleic acid-conjugate; muscle

uptake.

References

1. Ashizawa, T., Gagnon, C., Groh, W.J., Gutmann, L., Johnson, N.E., Meola, G., Moxley, R., Pandya, S., Rogers,
M.T., Simpson, E., et al. (2018). Consensus-based care recommendations for adults with myotonic dystrophy
type 1. Neurol Clin Pract *8*, 507–520. https://doi.org/10.1212/cpj.0000000000000531.

2. Ozimski, L.L., Sabater-Arcis, M., Bargiela, A., and Artero, R. (2021). The hallmarks of myotonic dystrophy
type 1 muscle dysfunction. Biol Rev Camb Philos Soc *96*, 716–730. https://doi.org/10.1111/brv.12674.

3. Holt, I., Mittal, S., Furling, D., Butler-Browne, G.S., Brook, J.D., and Morris, G.E. (2007). Defective mRNA in
myotonic dystrophy accumulates at the periphery of nuclear splicing speckles. Genes Cells *12*, 1035–1048.
https://doi.org/10.1111/j.1365-2443.2007.01112.x.

4. Goers, E.S., Purcell, J., Voelker, R.B., Gates, D.P., and Berglund, J.A. (2010). MBNL1 binds GC motifs
embedded in pyrimidines to regulate alternative splicing. Nucleic Acids Res *38*, 2467–2484.
https://doi.org/10.1093/nar/gkp1209.

5. Batra, R., Charizanis, K., Manchanda, M., Mohan, A., Li, M., Finn, D.J., Goodwin, M., Zhang, C., Sobczak, K.,
Thornton, C.A., et al. (2014). Loss of MBNL leads to disruption of developmentally regulated alternative
polyadenylation in RNA-mediated disease. Mol Cell *56*, 311–322.
https://doi.org/10.1016/j.molcel.2014.08.027.

6.   Lin, X., Miller, J.W., Mankodi, A., Kanadia, R.N., Yuan, Y., Moxley, R.T., Swanson, M.S., and Thornton,
C.A. (2006). Failure of MBNL1-dependent post-natal splicing transitions in myotonic dystrophy. Hum Mol
Genet *15*, 2087–2097. https://doi.org/10.1093/hmg/ddl132.

7.   Kuyumcu-Martinez, N.M., Wang, G.S., and Cooper, T.A. (2007). Increased steady-state levels of CUGBP1 in
myotonic dystrophy 1 are due to PKC-mediated hyperphosphorylation. Mol Cell *28*, 68–78.
https://doi.org/10.1016/j.molcel.2007.07.027.

8.   Wang, E.T., Ward, A.J., Cherone, J.M., Giudice, J., Wang, T.T., Treacy, D.J., Lambert, N.J., Freese, P.,
Saxena, T., Cooper, T.A., et al. (2015). Antagonistic regulation of mRNA expression and splicing by CELF and
MBNL proteins. Genome Res *25*, 858–871. https://doi.org/10.1101/gr.184390.114.

9.   Freyermuth, F., Rau, F., Kokunai, Y., Linke, T., Sellier, C., Nakamori, M., Kino, Y., Arandel, L., Jollet, A.,
Thibault, C., et al. (2016). Splicing misregulation of SCN5A contributes to cardiac-conduction delay and heart
arrhythmia in myotonic dystrophy. Nat Commun *7*, 11067. https://doi.org/10.1038/ncomms11067.

10. Wheeler, T.M., Lueck, J.D., Swanson, M.S., Dirksen, R.T., and Thornton, C.A. (2007). Correction of ClC-1
splicing eliminates chloride channelopathy and myotonia in mouse models of myotonic dystrophy. J Clin Invest
117, 3952–3957. https://doi.org/10.1172/jci33355.

11. Savkur, R.S., Philips, A.V., and Cooper, T.A. (2001). Aberrant regulation of insulin receptor alternative
splicing is associated with insulin resistance in myotonic dystrophy. Nat Genet *29*, 40–47.
https://doi.org/10.1038/ng704.

12. Cisco, L.A., Sipple, M.T., Edwards, K.M., Thornton, C.A., and Lueck, J.D. (2024). Verapamil mitigates
chloride and calcium bi-channelopathy in a myotonic dystrophy mouse model. J Clin Invest *134*, e173576.
https://doi.org/10.1172/JCI173576.

13. Braz, S.O., Acquaire, J., Gourdon, G., and Gomes-Pereira, M. (2018). Of Mice and Men: Advances in the
Understanding of Neuromuscular Aspects of Myotonic Dystrophy. Front Neurol *9*, 519.
https://doi.org/10.3389/fneur.2018.00519.

14. Thomas, J.D., Oliveira, R., Sznajder Ł, J., and Swanson, M.S. (2018). Myotonic Dystrophy and Developmental
Regulation of RNA Processing. Compr Physiol *8*, 509–553. https://doi.org/10.1002/cphy.c170002.

15. André, L.M., van Cruchten, R.T.P., Willemse, M., and Wansink, D.G. (2019). (CTG)n repeat-mediated
dysregulation of MBNL1 and MBNL2 expression during myogenesis in DM1 occurs already at the myoblast
stage. PLoS One *14*, e0217317. https://doi.org/10.1371/journal.pone.0217317.

16. Manta, A., Stouth, D.W., Xhuti, D., Chi, L., Rebalka, I.A., Kalmar, J.M., Hawke, T.J., and Ljubicic, V. (2019).
Chronic exercise mitigates disease mechanisms and improves muscle function in myotonic dystrophy type 1
mice. J Physiol *597*, 1361–1381. https://doi.org/10.1113/JP277123.

17. Chamberlain, C.M., and Ranum, L.P. (2012). Mouse model of muscleblind-like 1 overexpression: skeletal
muscle effects and therapeutic promise. Hum Mol Genet *21*, 4645–4654. https://doi.org/10.1093/hmg/dds306.

18. Choi, J., Personius, K.E., DiFranco, M., Dansithong, W., Yu, C., Srivastava, S., Dixon, D.M., Bhatt, D.B.,
Comai, L., Vergara, J.L., et al. (2015). Muscleblind-Like 1 and Muscleblind-Like 3 Depletion Synergistically
Enhances Myotonia by Altering Clc-1 RNA Translation. EBioMedicine *2*, 1034–1047.
https://doi.org/10.1016/j.ebiom.2015.07.028.

19. Lee, K.-Y., Chang, H.-C., Seah, C., and Lee, L.-J. (2019). Deprivation of Muscleblind-Like Proteins Causes
Deficits in Cortical Neuron Distribution and Morphological Changes in Dendritic Spines and Postsynaptic
Densities. Front Neuroanat *13*, 75. https://doi.org/10.3389/fnana.2019.00075.

20. Wang, E.T., Cody, N.A.L., Jog, S., Biancolella, M., Wang, T.T., Treacy, D.J., Luo, S., Schroth, G.P., Housman,
D.E., Reddy, S., et al. (2012). Transcriptome-wide regulation of pre-mRNA splicing and mRNA localization by
muscleblind proteins. Cell *150*, 710–724. https://doi.org/10.1016/j.cell.2012.06.041.

21. Sellier, C., Cerro-Herreros, E., Blatter, M., Freyermuth, F., Gaucherot, A., Ruffenach, F., Sarkar, P., Puymirat,
J., Udd, B., Day, J.W., et al. (2018). rbFOX1/MBNL1 competition for CCUG RNA repeats binding contributes
to myotonic dystrophy type 1/type 2 differences. Nat Commun *9*, 2009. https://doi.org/10.1038/s41467-018-
04370-x.

22. Kanadia, R.N., Shin, J., Yuan, Y., Beattie, S.G., Wheeler, T.M., Thornton, C.A., and Swanson, M.S. (2006).
Reversal of RNA missplicing and myotonia after muscleblind overexpression in a mouse poly(CUG) model for
myotonic dystrophy. Proc Natl Acad Sci U S A *103*, 11748–11753. https://doi.org/10.1073/pnas.0604970103.

23. Gladman, J.T., Mandal, M., Srinivasan, V., and Mahadevan, M.S. (2013). Age of Onset of RNA Toxicity
Influences Phenotypic Severity: Evidence from an Inducible Mouse Model of Myotonic Dystrophy (DM1).
PLOS ONE *8*, e72907. https://doi.org/10.1371/journal.pone.0072907.

24. Cerro-Herreros, E., Sabater-Arcis, M., Fernandez-Costa, J.M., Moreno, N., Perez-Alonso, M., Llamusi, B., and
Artero, R. (2018). miR-23b and miR-218 silencing increase Muscleblind-like expression and alleviate myotonic
dystrophy phenotypes in mammalian models. Nat Commun *9*, 2482. https://doi.org/10.1038/s41467-018-
04892-4.

25. Piasecka, A., Szcześniak, M.W., Sekrecki, M., Kajdasz, A., Sznajder, Ł.J., Baud, A., and Sobczak, K. (2024).
MBNL splicing factors regulate the microtranscriptome of skeletal muscles. Nucleic Acids Res *52*, 12055–
12073. https://doi.org/10.1093/nar/gkae774.

26. Cerro-Herreros, E., González-Martínez, I., Moreno, N., Espinosa-Espinosa, J., Fernández-Costa, J.M., Colom-
Rodrigo, A., Overby, S.J., Seoane-Miraz, D., Poyatos-García, J., Vilchez, J.J., et al. (2021). Preclinical
characterization of antagomiR-218 as a potential treatment for myotonic dystrophy. Mol Ther Nucleic Acids
26, 174–191. https://doi.org/10.1016/j.omtn.2021.07.017.

27. Cerro-Herreros, E., Núñez-Manchón, J., Naldaiz-Gastesi, N., Carrascosa-Sàez, M., García-Rey, A., Losilla,
D.P., González-Martínez, I., Espinosa-Espinosa, J., Moreno, K., Poyatos-García, J., et al. (2024). AntimiR
treatment corrects myotonic dystrophy primary cell defects across several CTG repeat expansions with a dual
mechanism of action. Sci Adv *10*, eadn6525. https://doi.org/10.1126/sciadv.adn6525.

28. Brook, J.D., McCurrach, M.E., Harley, H.G., Buckler, A.J., Church, D., Aburatani, H., Hunter, K., Stanton,
V.P., Thirion, J.P., and Hudson, T. (1992). Molecular basis of myotonic dystrophy: expansion of a trinucleotide
(CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. Cell *68*, 799–808.
https://doi.org/10.1016/0092-8674(92)90154-5.

29. Savić Pavićević, D., Miladinović, J., Brkušanin, M., Šviković, S., Djurica, S., Brajušković, G., and Romac, S.
(2013). Molecular genetics and genetic testing in myotonic dystrophy type 1. Biomed Res Int *2013*, 391821.
https://doi.org/10.1155/2013/391821.

30. Mahadevan, M.S., Yadava, R.S., and Mandal, M. (2021). Cardiac Pathology in Myotonic Dystrophy Type 1.
Int J Mol Sci *22*, 11874. https://doi.org/10.3390/ijms222111874.

31. Hilbert, J.E., Barohn, R.J., Clemens, P.R., Luebbe, E.A., Martens, W.B., McDermott, M.P., Parkhill, A.L.,
Tawil, R., Thornton, C.A., Moxley, R.T., et al. (2017). High frequency of gastrointestinal manifestations in
myotonic dystrophy type 1 and type 2. Neurology *89*, 1348–1354.
https://doi.org/10.1212/WNL.0000000000004420.

32. Peterson, J.A.M., and Cooper, T.A. (2022). Clinical and Molecular Insights into Gastrointestinal Dysfunction in
Myotonic Dystrophy Types 1 & 2. Int J Mol Sci *23*, 14779. https://doi.org/10.3390/ijms232314779.

33. Okkersen, K., Buskes, M., Groenewoud, J., Kessels, R.P.C., Knoop, H., van Engelen, B., and Raaphorst, J.
(2017). The cognitive profile of myotonic dystrophy type 1: A systematic review and meta-analysis. Cortex *95*,
143–155. https://doi.org/10.1016/j.cortex.2017.08.008.

34. van der Velden, B.G., Okkersen, K., Kessels, R.P., Groenewoud, J., van Engelen, B., Knoop, H., and
Raaphorst, J. (2019). Affective symptoms and apathy in myotonic dystrophy type 1 a systematic review and
meta-analysis. J Affect Disord *250*, 260–269. https://doi.org/10.1016/j.jad.2019.03.036.

35. Pascual-Gilabert, M., Artero, R., and López-Castel, A. (2023). The myotonic dystrophy type 1 drug
development pipeline: 2022 edition. Drug Discov Today *28*, 103489.
https://doi.org/10.1016/j.drudis.2023.103489.

36. van Roon-Mom, W., Ferguson, C., and Aartsma-Rus, A. (2023). From Failure to Meet the Clinical Endpoint to
U.S. Food and Drug Administration Approval: 15th Antisense Oligonucleotide Therapy Approved Qalsody
(Tofersen) for Treatment of SOD1 Mutated Amyotrophic Lateral Sclerosis. Nucleic Acid Ther *33*, 234–237.
https://doi.org/10.1089/nat.2023.0027.

37. Egli, M., and Manoharan, M. (2023). Chemistry, structure and function of approved oligonucleotide
therapeutics. Nucleic Acids Res *51*, 2529–2573. https://doi.org/10.1093/nar/gkad067.

38. Khvorova, A., and Watts, J.K. (2017). The chemical evolution of oligonucleotide therapies of clinical utility.
Nat Biotechnol *35*, 238–248. https://doi.org/10.1038/nbt.3765.

39. Eckstein, F. (2014). Phosphorothioates, essential components of therapeutic oligonucleotides. Nucleic Acid
Ther *24*, 374–387. https://doi.org/10.1089/nat.2014.0506.

40. De Serres-Bérard, T., Ait Benichou, S., Jauvin, D., Boutjdir, M., Puymirat, J., and Chahine, M. (2022). Recent
Progress and Challenges in the Development of Antisense Therapies for Myotonic Dystrophy Type 1. Int J Mol
Sci *23*. https://doi.org/10.3390/ijms232113359.

41. Frazier, K.S. (2015). Antisense oligonucleotide therapies: the promise and the challenges from a toxicologic
pathologist's perspective. Toxicol Pathol *43*, 78–89. https://doi.org/10.1177/0192623314551840.

42. Hagedorn, P.H., Yakimov, V., Ottosen, S., Kammler, S., Nielsen, N.F., Høg, A.M., Hedtjärn, M., Meldgaard,
M., Møller, M.R., Orum, H., et al. (2013). Hepatotoxic potential of therapeutic oligonucleotides can be
predicted from their sequence and modification pattern. Nucleic Acid Ther *23*, 302–310.
https://doi.org/10.1089/nat.2013.0436.

43. Alhamadani, F., Zhang, K., Parikh, R., Wu, H., Rasmussen, T.P., Bahal, R., Zhong, X.B., and Manautou, J.E.
(2022). Adverse Drug Reactions and Toxicity of the Food and Drug Administration-Approved Antisense
Oligonucleotide Drugs. Drug Metab Dispos *50*, 879–887. https://doi.org/10.1124/dmd.121.000418.

44. Ionis Pharmaceuticals Industry Updates On Drug Development (2018 Myotonic Dystrophy Foundation annual
conference). Available online: https://www.myotonic.org/digital-academy/ionis-pharmaceuticals-industry-
updates-drug-development-2018-mdf-annual-conference [accessed on 1 December 2022].

45. Hammond, S.M., Aartsma-Rus, A., Alves, S., Borgos, S.E., Buijsen, R.A.M., Collin, R.W.J., Covello, G.,
Denti, M.A., Desviat, L.R., Echevarría, L., et al. (2021). Delivery of oligonucleotide-based therapeutics:
challenges and opportunities. EMBO Mol Med *13*, e13243. https://doi.org/10.15252/emmm.202013243.

46.   Brown, K.M., Nair, J.K., Janas, M.M., Anglero-Rodriguez, Y.I., Dang, L.T.H., Peng, H., Theile, C.S.,
Castellanos-Rizaldos, E., Brown, C., Foster, D., et al. (2022). Expanding RNAi therapeutics to extrahepatic
tissues with lipophilic conjugates. Nat Biotechnol *40*, 1500–1508. https://doi.org/10.1038/s41587-022-01334-x.

47.   Geary, R.S., Norris, D., Yu, R., and Bennett, C.F. (2015). Pharmacokinetics, biodistribution and cell uptake of
antisense oligonucleotides. Adv Drug Deliv Rev *87*, 46–51. https://doi.org/10.1016/j.addr.2015.01.008.

48.   Malecova, B., Burke, R.S., Cochran, M., Hood, M.D., Johns, R., Kovach, P.R., Doppalapudi, V.R., Erdogan,
G., Arias, J.D., Darimont, B., et al. (2023). Targeted tissue delivery of RNA therapeutics using antibody-
oligonucleotide conjugates (AOCs). Nucleic Acids Res *51*, 5901–5910. https://doi.org/10.1093/nar/gkad415.

49.   Nakagawa, O., Ming, X., Huang, L., and Juliano, R.L. (2010). Targeted intracellular delivery of antisense
oligonucleotides via conjugation with small-molecule ligands. J Am Chem Soc *132*, 8848–8849.
https://doi.org/10.1021/ja102635c.

50.   Aartsma-Rus, A., and Corey, D.R. (2020). The 10th Oligonucleotide Therapy Approved: Golodirsen for
Duchenne Muscular Dystrophy. Nucleic Acid Ther *30*, 67–70. https://doi.org/10.1089/nat.2020.0845.

51.   Luo, T., Huo, C., Zhou, T., and Xie, S. (2023). Progress on RNA-based therapeutics for genetic diseases.
Zhejiang Da Xue Xue Bao Yi Xue Ban *52*, 406–416. https://doi.org/10.3724/zdxbyxb-2023-0190.

52.   Cerro-Herreros, E., González-Martínez, I., Moreno-Cervera, N., Overby, S., Pérez-Alonso, M., Llamusí, B.,
and Artero, R. (2020). Therapeutic Potential of AntagomiR-23b for Treating Myotonic Dystrophy. Molecular
Therapy - Nucleic Acids *21*, 837–849. https://doi.org/10.1016/j.omtn.2020.07.021.

53.   Fernández-Garibay, X., Ortega, M.A., Cerro-Herreros, E., Comelles, J., Martínez, E., Artero, R., Fernández-
Costa, J.M., and Ramón-Azcón, J. (2021). Bioengineeredin vitro3D model of myotonic dystrophy type 1
human skeletal muscle. Biofabrication *13*. https://doi.org/10.1088/1758-5090/abf6ae.

54.   Muller, P.Y., and Milton, M.N. (2012). The determination and interpretation of the therapeutic index in drug
development. Nat Rev Drug Discov *11*, 751–761. https://doi.org/10.1038/nrd3801.

55.   Navarro, N., Serantes, S., Aviñó, A., Fàbrega, C., and Eritja, R. (2023). Synthesis and Evaluation of 3′-Oleyl–
Oligonucleotide Conjugates as Potential Cellular Uptake Enhancers. Synlett *35*, 721–727.
https://doi.org/10.1055/s-0042-1751528.

56.   Arandel, L., Polay Espinoza, M., Matloka, M., Bazinet, A., De Dea Diniz, D., Naouar, N., Rau, F., Jollet, A.,
Edom-Vovard, F., Mamchaoui, K., et al. (2017). Immortalized human myotonic dystrophy muscle cell lines to
assess therapeutic compounds. Dis Model Mech *10*, 487–497. https://doi.org/10.1242/dmm.027367.

57.   Mankodi, A., Logigian, E., Callahan, L., McClain, C., White, R., Henderson, D., Krym, M., and Thornton, C.A.
(2000). Myotonic dystrophy in transgenic mice expressing an expanded CUG repeat. Science *289*, 1769–1773.
https://doi.org/10.1126/science.289.5485.1769.

58.   Seznec, H., Lia-Baldini, A.S., Duros, C., Fouquet, C., Lacroix, C., Hofmann-Radvanyi, H., Junien, C., and
Gourdon, G. (2000). Transgenic mice carrying large human genomic sequences with expanded CTG repeat
mimic closely the DM CTG repeat intergenerational and somatic instability. Hum Mol Genet *9*, 1185–1194.
https://doi.org/10.1093/hmg/9.8.1185.

59.   Gudde, A.E., González-Barriga, A., van den Broek, W.J., Wieringa, B., and Wansink, D.G. (2016). A low
absolute number of expanded transcripts is involved in myotonic dystrophy type 1 manifestation in muscle.
Hum Mol Genet *25*, 1648–1662. https://doi.org/10.1093/hmg/ddw042.

60. Burki, U., and Straub, V. (2017). Ultrasensitive Hybridization-Based ELISA Method for the Determination of
Phosphorodiamidate Morpholino Oligonucleotides in Biological samples. Methods Mol Biol *1565*, 265–277.
https://doi.org/10.1007/978-1-4939-6817-6_22.

61. Moisan, A., Gubler, M., Zhang, J.D., Tessier, Y., Dumong Erichsen, K., Sewing, S., Gérard, R., Avignon, B.,
Huber, S., Benmansour, F., et al. (2017). Inhibition of EGF Uptake by Nephrotoxic Antisense Drugs In Vitro
and Implications for Preclinical Safety Profiling. Mol Ther Nucleic Acids *6*, 89–105.
https://doi.org/10.1016/j.omtn.2016.11.006.

62. Provenzano, M., Ikegami, K., Bates, K., Gaynor, A., Hartman, J.M., Jones, A., Butler, A., Berggren, K.N.,
Dekdebrun, J., Hung, M., et al. (2025). The Splice Index as a prognostic biomarker of strength and function in
myotonic dystrophy type 1. J Clin Invest *135*, e185426. https://doi.org/10.1172/JCI185426.

63. Moreno, N., González-Martínez, I., Artero, R., and Cerro-Herreros, E. (2022). Rapid Determination of MBNL1
Protein Levels by Quantitative Dot Blot for the Evaluation of Antisense Oligonucleotides in Myotonic
Dystrophy Myoblasts. Methods Mol Biol *2434*, 207–215. https://doi.org/10.1007/978-1-0716-2010-6_13.

64. Dincã, D.M., Lallemant, L., González-Barriga, A., Cresto, N., Braz, S.O., Sicot, G., Pillet, L.-E., Polvèche, H.,
Magneron, P., Huguet-Lachon, A., et al. (2022). Myotonic dystrophy RNA toxicity alters morphology,
adhesion and migration of mouse and human astrocytes. Nat Commun *13*, 3841.
https://doi.org/10.1038/s41467-022-31594-9.

65. Østergaard, M.E., Jackson, M., Low, A., E. Chappell, A., G. Lee, R., Peralta, R.Q., Yu, J., Kinberger, G.A.,
Dan, A., Carty, R., et al. (2019). Conjugation of hydrophobic moieties enhances potency of antisense
oligonucleotides in the muscle of rodents and non-human primates. Nucleic Acids Research *47*, 6045–6058.
https://doi.org/10.1093/nar/gkz360.

66. Tran, P., Weldemichael, T., Liu, Z., and Li, H. (2022). Delivery of Oligonucleotides: Efficiency with Lipid
Conjugation and Clinical Outcome. Pharmaceutics *14*, 342. https://doi.org/10.3390/pharmaceutics14020342.

67. Prakash, T.P., Mullick, A.E., Lee, R.G., Yu, J., Yeh, S.T., Low, A., Chappell, A.E., Østergaard, M.E., Murray,
S., Gaus, H.J., et al. (2019). Fatty acid conjugation enhances potency of antisense oligonucleotides in muscle.
Nucleic Acids Res *47*, 6029–6044. https://doi.org/10.1093/nar/gkz354.

68. Kusznir, E.-A., Hau, J.-C., Portmann, M., Reinhart, A.-G., Falivene, F., Bastien, J., Worm, J., Ross, A., Lauer,
M., Ringler, P., et al. (2023). Propensities of Fatty Acid-Modified ASOs: Self-Assembly vs Albumin Binding.
Bioconjug Chem *34*, 866–879. https://doi.org/10.1021/acs.bioconjchem.3c00085.

69. Henry, S., Stecker, K., Brooks, D., Monteith, D., Conklin, B., and Bennett, C.F. (2000). Chemically modified
oligonucleotides exhibit decreased immune stimulation in mice. J Pharmacol Exp Ther *292*, 468–479.

70. Simon-Friedt, B.R., Wilson, M.J., Blake, D.A., Yu, H., Eriksson, Y., and Wickliffe, J.K. (2015). The
RPTEC/TERT1 Cell Line as an Improved Tool for In Vitro Nephrotoxicity Assessments. Biol Trace Elem Res
166, 66–71. https://doi.org/10.1007/s12011-015-0339-y.

71. Izzedine, H., and Perazella, M.A. (2017). Adverse kidney effects of epidermal growth factor receptor inhibitors.
Nephrology Dialysis Transplantation *32*, 1089–1097. https://doi.org/10.1093/ndt/gfw467.

72. Dirin, M., and Winkler, J. (2013). Influence of diverse chemical modifications on the ADME characteristics
and toxicology of antisense oligonucleotides. Expert Opin Biol Ther *13*, 875–888.
https://doi.org/10.1517/14712598.2013.774366.

73. van Dongen, S., Abreu-Goodger, C., and Enright, A.J. (2008). Detecting microRNA binding and siRNA off-
target effects from expression data. Nat Methods *5*, 1023–1025. https://doi.org/10.1038/nmeth.1267.

74. González-Martínez, I., Cerro-Herreros, E., Moreno, N., García-Rey, A., Espinosa-Espinosa, J., Carrascosa-
Sàez, M., Piqueras-Losilla, D., Arzumanov, A., Seoane-Miraz, D., Jad, Y., et al. (2023). Peptide-conjugated
antimiRs improve myotonic dystrophy type 1 phenotypes by promoting endogenous MBNL1 expression.
Molecular Therapy - Nucleic Acids *34*, 102024. https://doi.org/10.1016/j.omtn.2023.09.001.

75. Juliano, R.L. (2016). The delivery of therapeutic oligonucleotides. Nucleic Acids Res *44*, 6518–6548.
https://doi.org/10.1093/nar/gkw236.

76. González-Barriga, A., Nillessen, B., Kranzen, J., van Kessel, I.D.G., Croes, H.J.E., Aguilera, B., de Visser,
P.C., Datson, N.A., Mulders, S.A.M., van Deutekom, J.C.T., et al. (2017). Intracellular Distribution and
Nuclear Activity of Antisense Oligonucleotides After Unassisted Uptake in Myoblasts and Differentiated
Myotubes In Vitro. Nucleic Acid Ther *27*, 144–158. https://doi.org/10.1089/nat.2016.0641.

77. Leonetti, J.P., Mechti, N., Degols, G., Gagnor, C., and Lebleu, B. (1991). Intracellular distribution of
microinjected antisense oligonucleotides. Proc Natl Acad Sci U S A *88*, 2702–2706.
https://doi.org/10.1073/pnas.88.7.2702.

78. Jiang, Y.-H., Man, Y.-Y., Liu, Y., Yin, C.-J., Li, J.-L., Shi, H.-C., Zhao, H., and Zhao, S.-G. (2021). Loss of
miR-23b/27b/24-1 Cluster Impairs Glucose Tolerance via Glycolysis Pathway in Mice. Int J Mol Sci *22*, 550.
https://doi.org/10.3390/ijms22020550.

79. Hao, S.-J., Zhao, Y.-Y., Wu, Z.-J., An, M.-Y., Song, C.-C., and Li, J. (2025). MiR-23b regulation of
metabolites in neuropathic pain: A novel approach. Behavioural Brain Research *495*, 115768.
https://doi.org/10.1016/j.bbr.2025.115768.

80. Hannafon, B.N., Cai, A., Calloway, C.L., Xu, Y.-F., Zhang, R., Fung, K.-M., and Ding, W.-Q. (2019). miR-23b
and miR-27b are oncogenic microRNAs in breast cancer: evidence from a CRISPR/Cas9 deletion study. BMC
Cancer *19*, 642. https://doi.org/10.1186/s12885-019-5839-2.

81. de Rie, D., Abugessaisa, I., Alam, T., Arner, E., Arner, P., Ashoor, H., Åström, G., Babina, M., Bertin, N.,
Burroughs, A.M., et al. (2017). An integrated expression atlas of miRNAs and their promoters in human and
mouse. Nat Biotechnol *35*, 872–878. https://doi.org/10.1038/nbt.3947.

82. Huang, C.K., Kafert-Kasting, S., and Thum, T. (2020). Preclinical and Clinical Development of Noncoding
RNA Therapeutics for Cardiovascular Disease. Circ Res *126*, 663–678.
https://doi.org/10.1161/circresaha.119.315856.

     Tables and figures

Figure 1. *In vitro* screening of antimiR-23b variants and assessment of safety and efficacy of a prospective lead candidate in two mouse models.

978
979

Figure 1. *In vitro* **screening of antimiR-23b variants and assessment of safety and efficacy of a prospective lead candidate in two mouse models. (A)** Schematic representation of the antimiRs tested in the screening. Modifications 1, 2, and 3 refer to 2'-O-Methyl RNA nucleotides,

2'-O-MOE RNA nucleotides, and LNA nucleotides. Backbones 1 and 2 represent a mix of PS

(phosphorothioate) and PO (phosphodiester) linkages. **(B)** Representative graphs illustrating the

ranking of the compounds tested in the screening, based on their EC_{50} (µM) and TC_{50} (µM) values.

The graph also includes a table displaying the T_{index} of each antimiR. In both graph and table, the

benchmark compounds are indicated in dark gray, the antimiR hit of the screening in greenish-

blue, and the remaining antimiRs in gray. The parameters measured in the two mouse models

(HSA^{LR} **(C,D,E)** and DMSXL **(F,G,H)**) treated subcutaneously with X821 at SD 100 mg/kg and

MD 25 mg/kg, respectively, were: **(C,F)** MBNL1 levels of treated mice relative to saline controls

in skeletal muscles affected in DM1; **(D,G)** the percentage of exon inclusion (Ψ) of the indicated

exons and muscles. **(E,H)** Functional rescues after treatment: **(E)** relative force and myotonia grade

change compared to controls; **(H)** change in relative muscle strength of antimiR-treated mice

compared to saline controls at the indicated days after treatment. Error bars indicate mean ± SEM.

*: $p<0.05$; **: $p<0.01$; ***: $p<0.001$. The data were analyzed using unpaired Student's t-test,

compared with untreated DM1 mice (phosphate-buffered saline, PBS). AI: after injection; SD:

single dose, MD: multiple dose. HSA^{LR}: PBS n=7, X821 n=3; DMSXL: PBS n=6, X821 n=6.

1000

1001

1002

Figure 2. *In vitro* screening of antimiRs-23b with different conjugates and assessment of safety and efficacy of a group of lead candidates in *HSA^{LR}* mice. (**A**) Schematic representation of the antimiRs tested in the screening. Modifications 1, 2, and 3 refer to 2'-O-Methyl RNA nucleotides, 2'-O-MOE RNA nucleotides, and LNA nucleotides. Backbones 1 and 2 represent a mix of PS (phosphorothioate) and PO (phosphodiester) linkages. (**B**) Representative graph illustrating the ranking of the compounds tested based on their EC$_{50}$ (µM) and TC$_{50}$ (µM) values. The graph also includes a table displaying the T$_{index}$ of each antimiR. In both graph and table, the benchmark compounds are indicated in dark gray, the antimiR hit of the screening in greenish-blue, and the remaining antimiRs in gray. (**C-F**) Selected compounds were tested *in vivo* using a single 3 mg/kg IV injection in *HSA^{LR}* mice. After 5 days, several parameters were measured, and skeletal muscle tissues were dissected for analysis. (**C**) Schematic representation of the antimiRs tested *in vivo*. (**D**) MBNL1 levels of treated mice relative to saline controls in skeletal muscle 5

days after injection (mean value of gt and qd). **(E)** Overall percentage splice recovery (PSR) of leading antimiR candidates in both muscles. **(F)** AntimiR X82107 vs X821-7 levels in muscle tissues, kidney and liver detected by ELISA. The multiplier value indicates the fold-change between the different versions tested. Error bars indicate mean ± SEM. *: $p < 0.05$; ***: $p < 0.001$. The data were analyzed by one-way ANOVA or Kruskal-Wallis test if required, compared with untreated HSA^{LR} mice (PBS). Individual values are indicated as data points. PBS n=3, X821-7 n=4, 5'X82107 n=3, X82107 n=4.

1025

Figure 3. Single-dose experiment with leading antimiR derivatives from the previous experiment. (A) Schematic representation of the antimiRs tested. Modifications 1, 2, and 3 refer to 2'-O-Methyl RNA nucleotides, 2'-O-MOE RNA nucleotides, and LNA nucleotides. Backbones 1 and 2 represent a mix of PS (phosphorothioate) and PO (phosphodiester) linkages. Selected compounds were tested *in vivo* using a single 3 mg/kg IV injection in *HSA^{LR}* mice. **(B)** MBNL1 levels of treated mice relative to saline controls in skeletal muscle (mean value of gt and qd). **(C)** Overall PSR of leading antimiR candidates in both muscles. **(D)** AntimiR X82108 and X821-8 levels in muscle tissues (mean value of qd and gt) detected by ELISA. The multiplier value indicates the fold-change between the different versions tested. Error bars indicate mean ± SEM. *: $p<0.05$; **: $p<0.01$; ****: $p<0.0001$. The data were analyzed by one-way ANOVA or Kruskal-Wallis test if required, compared with untreated *HSA^{LR}* mice (PBS). Individual values are indicated as data points. PBS n=3, X821-8 n=3, X82108 n=4, X82148 n=4.

1038

1039
1040

1041

1042

Figure 4. Effect of Oleyl conjugate in different antimiRs. (A) Schematic representation of the

experimental design using a single 3 mg/kg IV injection in HSA^{LR} mice. Modifications 1, 2, and 3

refer to 2'-O-Methyl RNA nucleotides, 2'-O-MOE RNA nucleotides, and LNA nucleotides.

Backbones 1 and 2 represent a mix of PS (phosphorothioate) and PO (phosphodiester) linkages.

(B) Schematic representation of the antimiRs tested. **(C)** Spider graphs depicting the readouts

obtained in the HSA^{LR} mouse model with leading antimiR candidates. Functional recovery includes

strength and myotonia measurements. PSR includes rescues of *Atp2a1* ex22, *Clcn1* ex7a, *Nfix* ex7

and *Mbnl1* ex5 in qd and gt. FVB n=5, PBS n=3, X821-7 n=4, X82107 n=3, X821-8 n=3, X82108

n=3, X831-4 n=4, X83104 n=4.

1052

1053

1054

1055

1056

1057
Figure 5. Selected antimiRs effects until day 15. (A) Schematic representation of the
experimental design using a single 3 mg/kg IV injection in *HSA*^{*LR*} mice. During 15 days after
injection, several parameters were measured, and skeletal muscle tissues were collected for
analysis: **(B)** MBNL1 levels of treated mice relative to saline controls and **(C)** overall PSR score
in qd and gt muscles, **(D)** strength/body weight (g) and **(E)** myotonia grade at different time points
after injection (AI). *: p<0.05; **: p<0.01; ***: p<0.001; ****: p<0.0001. The data were analyzed
by one-way ANOVA or Kruskal-Wallis test if required, compared with untreated *HSA*^{*LR*} mice
(PBS). The shaded area indicates the error in each case.

A

B

1066
1067

Figure 6. *In vivo* preclinical analysis with leading antimiR candidates. (A) Schematic representation of the experimental design using a single 6 mg/kg IV injection in *HSA^{LR}* mice. **(B)** Schematic representation of the antimiRs tested. Modifications 1, 2, and 3 refer to 2'-O-Methyl RNA nucleotides, 2'-O-MOE RNA nucleotides, and LNA nucleotides. Backbones 1 and 2 represent a mix of PS (phosphorothioate) and PO (phosphodiester) linkages. **(C)** MBNL1 levels of treated mice relative to saline controls in both muscles (mean value of qd and gt). **(D)** Overall PSR scores. **(E)** Front leg grip strength normalized to body weight. **(F)** Percentage of PSR and MSR (muscle strength rescue) rescues. Blue line indicates wild-type levels. Error bars indicate mean ± SEM. *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$; ****: $p<0.0001$. The data were analyzed by ANOVA one-way test or Kruskal-Wallis test, when necessary, compared with untreated *HSA^{LR}* mice (PBS). Color legend: PBS: gray; X82107: blue; X82108: orange.

1079

Figure 7. Toxicity and immune activation experiments with ascending doses of X82107 and X82108. EGF measurement **(A, B)** and cell proliferation inhibition assay **(C, D)** in RPTEC/TERT1 cells upon antimiR treatment. **(E-H)** Immune activation analysis in blood samples from three different donors: **(E)** activated complement (C3a) measurement, **(F)** coagulation assay (prothrombin and activated partial thromboplastin times), and **(G)** total and activated platelet count as measured by several cell surface markers. **(H)** Cytokine release in human peripheral blood mononuclear cells (PBMCs) upon X82108 treatment. *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$; ****: $p<0.0001$ according to one-way ANOVA or Kruskal-Wallis test if required, compared with negative control. aPPT: activated partial thromboplastin time; CD61: integrin beta-3; CD62P: platelet surface P-selectin; IFN: interferon; IL: interleukin; MCP1: monocyte chemoattractant protein-1; PAC1: activated GP IIb/IIIa; PBS: phosphate buffered saline; PT: prothrombin time; TNF: tumor necrosis factor.

1093

Figure 8. Confirmation of conserved antimiR-23b regulation activity on MBNL levels in various cell types. (A) MBNL1 levels and **(B)** miR-23b relative levels in Rat2 and ST cells upon treatment with X82108. *: $p<0.05$, ****: $p<0.0001$ according to Student's t-test compared with non-treated cells; ST: *Sus scrofa* (testis). **(C)** miR-23b relative levels and **(D,E)** MBNL1 levels in primary myoblasts derived from DM1 patients upon treatment. MP-49F: control cell line; MP-09-57: DM1 cell line carrying 702 CTG repeats in *DMPK*; MP-10-31: DM1 cell line carrying 117

repeats. **(F)** Overall Percent Splicing Rescue (PSR) values for the adapted biomarker panel based on the Composite Alternative Splicing Index (CASI-22) in immortalized DM1 myotubes treated with 200 nM antimiR X82108; each colored dot represents an individual splicing event. *ANK2*-green; *ATP2A1*-fuchsia; *OPA1*-bright green; *BIN1*-light blue, *DMD*-yellow; *GFPT1*-red; *GOLGA4*- Blue; *MBNL1*-Orange, *INSR*-light violet; *NFIX*- mustard yellow and *SOS1*- magenta. -: untreated; +: treated with X82107; *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$. The data were analyzed by one-way ANOVA or Kruskal-Wallis test if required, compared with untreated control cell line (MP-49F). ST: *Sus scrofa* (testis).