Genus-wide homologous recombination of tail fibers maintains tailocin diversity in *Pectobacterium*

Lakhansing A. Pardeshi[1,2,*], Anne Kupczok[1], Dick de Ridder[1], Sandra Smit[1], and Theo A. J. van der Lee[2,*]

[1]Bioinformatics Group, Wageningen University, Droevendaalsesteeg 1, 6708PB, Wageningen, The Netherlands

[2]Biointeractions and Plant Health, Wageningen Plant Research, Droevendaalsesteeg 1, 6708PB, Wageningen, The Netherlands

[*]Corresponding authors: lakhansing.pardeshi@wur.nl, theo.vanderlee@wur.nl

Abstract

Due to their ability to kill closely related strains, phage tail-like bacteriocins, also called tailocins, play an important role in shaping bacterial communities. One such tailocin, called carotovoricin, is also known to be present in the *Pectobacterium* species. However, little is known about its evolutionary dynamics and the scope of impact on species interactions in this genus. To investigate the diversity and evolution of carotovoricin, we performed a genus-wide, phylogenetically-structured pangenome study. This analysis inferred that the gene cluster responsible for carotovoricin biosynthesis is conserved across the genus and is located in the same gene neighborhood in all the species. Within the carotovoricin cluster, the tail fiber genes, which determine the host range specificity, exhibit high variability and discordance with the species phylogeny. We show evidence for an evolutionary mechanism involving recombination-mediated exchange of these tail fiber loci across the entire *Pectobacterium* genus, which complements the previously known mechanism for DNA sequence inversion to maintain tailocin polymorphism at the population level. In addition, the ability to exchange tail-fiber loci in a highly targeted and genus-wide manner could influence the community dynamics in nutrient rich environments such as infected plant tissues. In conclusion, the strong signal for carotovoricin retention and ability to exchange tail fibers indicates that it significantly contributes to the community interactions of the *Pectobacterium* phytopathogens.

Significance Statement

A widespread presence of tailocins among various gram-negative bacteria and maintenance of their tail fiber diversity underscore their role in inter-bacterial interactions. A tailocin is also found to be conserved in *Pectobacterium*, a pathogen causing soft rot. However, the mechanism maintaining the diversity of the tailocin tail fibers, which enable recognition of the target bacteria, is not yet completely understood. Here, we characterized the genomic diversity of this tailocin and discovered that the diversity is maintained through the exchange of the tail fiber locus DNA across the genus.

Keywords: Tailocins, Horizontal gene transfer, *Pectobacterium*, Pangenomics

1 Introduction

Bacterial genomes harbor integrated temperate bacteriophages (called prophages) (*1–4*), as well as defec-
tive prophages that degrade over time (*5*). Evidence suggests that some defective prophages, lacking capsid
and replication genes, were repurposed as phage tail-like bacteriocins, also known as tailocins (*6–9*). Struc-
turally, tailocins resemble "headless phages" with tail and tail fibers forming a functional unit to kill target
bacteria by puncturing the cell membrane (*10*). These tailocins are widespread throughout gram-negative
bacteria (*9*, *11–14*) and are divided into two classes, based on having a rigid (R-type) or flexible (F-type)
core structure. The F-type tailocins are noncontractile and flexuous. In contrast, R-type tailocins have a
rigid contractile tail, formed of a hollow tube encapsulated by a contractile sheath, attached to a baseplate
along with tail fibers (*15*, *16*). In general, the tailocin tail fibers recognize and bind the target bacterial
cell surface receptors, followed by contraction of the sheath and extension of the core tube.

As the tail fibers enable the recognition of specific target cell membrane receptors, tailocins offer an effective
and targeted mechanism of killing co-occurring bacteria (*17*). Normally, the target range is very narrow
and limited to strains of the same species or phylogenetically close relatives (*9*). Increasing the variability
of tail fiber genes extend the recognition spectrum. Bacteria possess various mechanisms to increase the
tailocin target range, such as recombination of tail fiber genes to produce new sequence combinations (*18*),
the presence of multiple tail fiber genes within tailocin loci (*19*), and tail-fiber locus DNA inversion by
invertase (*20*). These variations in the tail fibers yield a diverse pool of tailocins that are maintained in a
species (*21*).

A tailocin was discovered in the global phytopathogen *Pectobacterium carotovorum*, hence named caro-
tovoricin (*14*, *22*). Subsequently, it was shown to be present across the entire *Pectobacterium* genus (*23–
25*). Carotovoricin is produced at the basal level by most of the *Pectobacterium* species, which results
in inter-species competitive interactions (*25*). Although maintenance of diversity in tailocin tail fibers is
known in other species, such as *Pseudomonas* (*26*), the existence and scope of such diversity are not yet
known for the *Pectobacterium* genus. *Pectobacterium* pathogens are known to cause multi-species infections
(*27–29*); they also show strain-specific competitive interactions, where certain strains can inhibit others,
but none can inhibit the entire species (*30*). A systematic study of tailocin diversity at the genus level
would allow understanding the broader inter-strain and inter-species interactions. It can further help to
understand how carotovoricin shapes the community dynamics of co-inhabiting *Pectobacterium* strains and
species.

We recently applied a comparative pangenomics approach to identify orthologous prophages and their
contribution in the *Pectobacterium* genus pangenome dynamics (*24*). Here, we elaborate our approach
to systematically study the extent of conservation of the carotovoricin-biosynthesis gene cluster in the
Pectobacterium genus. Furthermore, we study local variation within the carotovoricin cluster to understand
the polymorphic nature of tailocin tail fibers. Subsequently, we evaluate the evolution of the genes within
the carotovoricin cluster to help understand the underlying mechanism generating polymorphisms. Finally,

we discuss the potential role of carotovoricin in the intra- and inter-species community dynamics observed
in *Pectobacterium* species and the potential applications of leveraging tailocins as biocontrol strategy.

Results

Carotovoricin is conserved in the *Pectobacterium* genus

We recently constructed a *Pectobacterium* genus pangenome with PanTools (*31*, *32*) and discovered the
contribution of prophages to pangenome growth and the emergence of new lineages (*24*). This pangenome
comprises 454 *Pectobacterium* genomes (g_1 to g_454 in Supplementary Table S1), where 1,977,865 protein-
coding genes were grouped into 30,156 homology groups. To study the dynamics of 1,319 putative prophages
identified in these genomes, we represented prophages as a sequence of homology groups and traced these
homology group signatures across the pangenome. We identified orthologous prophages by quantifying
syntenic conservation of homology group signatures between prophages and dereplicated the prophages
to 436 clusters (see Methods, Supplementary Tables S2, S3). The largest of these clusters, containing
428 orthologous prophages, was identified as the phage tail-like bacteriocin carotovoricin, based on the
functional annotation of homology groups. Each carotovoricin region contained genes encoding the core
structural proteins necessary for tailocin production: holin, lytic transglycosylase, tail sheath protein, tail
tube protein, tail assembly protein, tail tape measure protein, baseplate proteins, tail spike protein, and
tail fiber proteins (Figure 1a).

We observed a single copy of the carotovoricin biosynthesis gene cluster in 428 *Pectobacterium* genomes
(Supplementary Table S4). Of these, 365 were present on a single contig and the remaining were split over
two contigs due to fragmented genome assemblies. Because of the presence of such fragmented assemblies
in the public repository, a 5kb region flanking the carotovoricin was used to determine the flanking gene
structural conservation rather than a fixed number of neighboring genes. This gene neighborhood flanking
carotovoricin cluster, comprising three to four genes, is syntenically conserved across the entire *Pectobac-*
terium genus (Figure 1b), suggesting that a common ancestor acquired the carotovoricin cluster. Genes
tolC_2 and *ybiB* mark the upstream and downstream boundary of the carotovoricin biosynthesis gene
clusters in all *Pectobacterium* species.

Complete carotovoricins encoding all core structural genes were found to be present in all but three *Pec-*
tobacterium species (Figure 1b), detailed below. In addition to the three species-wide losses, there were
seven strains from various *Pectobacterium* species that showed disruption of the carotovoricin cluster (Sup-
plementary Results). Species with a complete carotovoricin contained 18 to 27 genes. Most of these genes
were conserved, as evidenced by a single homology group per gene across all species. A partial carotovoricin
cluster was observed in the *P. betavasculorum* and *P. atrosepticum* genomes, where only the first five and
seven genes from the 5' end were found to be present, respectively. The genes in these partial carotovoricins
are part of the first transcriptional unit out of the four identified in the carotovoricin cluster (*7*). Publicly
available transcriptomics data (*33*) suggests that the genes within this partial carotovoricin cluster are
expressed and regulated in *P. atrosepticum* (see Methods); this could not be verified for *P. betavasculorum*,

due to a lack of data. Based on a phylogeny built on 1,949 core genes in the pangenome (Supplementary Figure S1), *P. betavasculorum* and *P. atrosepticum* share a common ancestor with *P. peruviense* and *P. zantedeschiae* respectively, both with intact carotovoricins. Given the difference in the deleted carotovoricin cluster and the phylogenetic distance to the closest carotovoricin-carrying species, it seems that carotovoricin cluster was degraded independently in *P. betavasculorum* and *P. atrosepticum*. The third species without carotovoricin is *P. cacticida*, which lacks carotovoricin completely and shows no remnants. A recent phylogenomic analysis proposed a reclassification of *P. cacticida* into a new genus, *Alcorniella comb. nov* (*34*), which likely explains the lack of carotovoricin in this genome. Overall, carotovoricin in conserved and transmitting vertically in the *Pectobacterium* genus.

Tail fiber genes show higher variation than other carotovoricin genes

To understand the inter-species variation in carotovoricin, we investigated if it has further evolved independently in each species. To avoid analysis artifacts due to assembly quality, we selected carotovoricin cluster from 365 genomes in which it was present on a single contig together with the flanking genes. The optimal homology grouping that accounted for sequence diversity (see Methods) resulted in a single homology group for each of the carotovoricin core structural genes across the *Pectobacterium* genus. However, we found a total of 130 homology groups mapped to carotovoricin regions across the *Pectobacterium* genus, in contrast to the anticipated small set of carotovoricin-related homology groups. Closer inspection of the carotovoricin cluster across genomes showed a conserved region where the genes were represented by a single homology group, and a variable region where the genes were represented by multiple homology groups (Figure 1b, Supplementary Table S5).

First, we observed presence-absence variation of the nitric oxide deoxygenase gene (Supplementary Table S5) (present in 88/365 the genomes) (Figure 2a). The gene encoding the tail tape measure protein, located between the tail assembly chaperon and P2 GpU family protein, also showed variation across the *Pectobacterium* genus and was represented by 9 homology groups. Finally, the 3' region flanked by the tail protein and the *ybiB* gene was represented by multiple homology group combinations across the pangenome annotation graph (Figure 2a). Interestingly, the invertase *ein*, responsible for the inversion of tail fiber locus DNA, also showed presence-absence variation: it was found in 47% (172/365) of the carotovoricins. A similar degree of diversity in carotovoricin cluster was also observed within a species, as exemplified by *P. brasiliense* (Supplementary Figure S4) and *P. versatile* species (Supplementary Figure S5). In summary, the carotovoricin cluster seems to be broadly organized into two differentially conserved loci, both at the genus and species levels.

As illustrated in Figure 2a, we define the carotovoricin cluster region spanning from the holin to the tail protein encoding gene as the conserved locus, and the remaining region as the tail fiber locus (TFL), for subsequent analysis. We further used the uniquely ordered combinations of homology groups in the TFL to define TFL haplotypes. In total, there were 93 TFL haplotypes in the *Pectobacterium* genus (Supplementary Table S6). The species *P. brasiliense* and *P. versatile* showed the most haplotypes, 21 and 27 respectively (Figure 2b). Five species had 8 or more TFL haplotypes, and the remaining species

had less than 5 haplotypes, which might be because of the lower number of genome assemblies for these species. Interestingly, intra-species average nucleotide identity (ANI), is not related to the number of TFL haplotypes. For example, *P. brasiliense* is more diverse than *P. versatile* (based on broader intra-species ANI in *P. brasiliense* (Supplementary Figure S6)), however, we observed more TFL haplotypes in the ecologically and geographically broadly distributed *P. versatile* (*35*) compared to *P. brasiliense*.

The same tail fiber loci are observed across different species

We observed that genomes from two or more *Pectobacterium* species can have the same TFLs (Figure 2b, Supplementary Table S6). For example, among the samples collected at the watershed niche along the Durance River in the southeast of France (*36*), *P. carotovorum* (g_106, g_57) had the same TFL as *P. versatile* (g_53). In the Netherlands, we found that *P. brasiliense* (g_145, g_194, g_429) and *P. aroidearum* (g_421, g_442) have the same TFL haplotype. *P. versatile* (g_150, g_447) and *P. odoriferum* (g_434) from the Netherlands likewise shared the same TFL haplotype. In contrast, some species did not share TFL haplotypes, for example *P. parmentieri* and *P. brasiliense*. To further understand the complete extent of TFL haplotype sharing, we mapped all TFL haplotypes on the *Pectobacterium* genus phylogeny (Figure 3a-c). We observed that these haplotypes do not follow the phylogeny and found widespread evidence of shared TFL haplotypes, both intra- and inter-species.

To study the intra- and inter-species variation of conserved locus and the TFL (as defined in Figure 2a) at a higher resolution, we aimed to compare the DNA sequences between these two loci across the *Pectobacterium* genus. However, the presence of multiple structural variants in the TFL region, including inversions, insertions, and deletions, impedes the reconstruction of a reliable multiple sequence alignment, which would be required for phylogeny reconstruction. Therefore, we used a k-mer approach to calculate pairwise distance between these loci. Pairwise species distance was calculated as the cophenetic distance in the core-SNP phylogeny derived from the pangenome.

Under the assumption of uniform mutation rate across the genome, we expect the locus distance to correlate with the species distance. In line with this, pairwise locus distances for the conserved locus showed a strong positive correlation with species distance ($R = 0.86$, $p < 2.2 \times 10^{-16}$) (Figure 4a). Additionally, the intra-species pairwise locus distances for the conserved locus were smaller than the inter-species distances, as might be expected. For the TFL, we also observed a positive, albeit weaker, correlation between locus distance and species distance ($R = 0.55$, $p < 2.2 \times 10^{-16}$) (Figure 4b). However, the locus distance range was significantly broader for both intra- and inter-species TFL compared to the conserved locus (Levene's test, $p < 2.2 \times 10^{-16}$). This overall broad range of pairwise distances for TFLs underlines the sequence level diversity of the tail fiber pool in the *Pectobacterium* genus. As expected, TFL pairs with the same haplotypes exhibited lower locus distances, regardless of whether they were intra- or inter-species comparisons. Strikingly, however, some of the inter-species TFL pairs had a very small locus distance (sometimes even zero), indicating (nearly) identical TFLs in different *Pectobacterium* species (Figure 4b).

Genus-wide homologous recombination-mediated exchange of tail fiber loci

Given the presence of identical TFLs across different species, suggesting a possible horizontal gene transfer (HGT), we further analyzed phylogenetic incongruence between the conserved locus and the TFL using 16 genomes from 10 *Pectobacterium* species. These genomes encompassed three different TFL haplotypes that lacked inversions, thus allowing us to reconstruct a reliable multiple sequence alignment and phylogenetic tree. In the absence of HGT, the expectation is that both phylogenetic trees should show a consensus and follow the *Pectobacterium* species tree. However, while the conserved locus phylogeny showed strains clustering by *Pectobacterium* species, the tree derived from the variable TFL region alignment did not follow the *Pectobacterium* species phylogeny (Figure 5a-b). This therefore supports that the TFLs are exchanged among *Pectobacterium* species, independent of the rest of the carotovoricin locus.

A horizontal transfer of such loci between two species and strains can be mediated by homologous recombination. A key requirement for successful DNA recombination is the presence of a homologous region. To assess the existence of such compositional similarity at the sequence level in the regions flanking the TFLs, we identified locally collinear blocks of DNA sequences in selected TFLs using Mauve (*37*, *38*).

First, the intra-species comparison of *P. brasiliense* TFLs with *ein* showed inversions in this region across several genomes, flanked by positionally conserved locally collinear blocks with high conservation across the genomes (Figure 5c). These positionally conserved collinear blocks imply a common mechanism of DNA inversion, most likely Ein-mediated. The upstream conserved collinear block in TFL overlaps with the first half of the tail fiber gene, implying a conserved N-terminal domain of the protein encoded by this gene. This observation extends the previously reported modular structure of tail fibers in *P. carotovorum* (*7*) to the other *Pectobacterium* species. The remaining region of the TFL shows complex structural rearrangements and results in variable C-terminal tail fiber domains.

Another comparison involving TFLs from multiple *Pectobacterium* species also showed a complex reordering and inversion of locally collinear blocks (Figure 5d). The flanking collinear blocks in TFLs without *ein* also showed positional conservation (Figure 5d), although it was less pronounced. The overall conservation of positions of the collinear blocks flanking the TFLs in both sets supports our hypothesis of recombination-mediated horizontal gene transfer.

Discussion

Carotovoricin, a tailocin, is produced constitutively as well as under stress by *Pectobacterium* species during host infection, both on symptomatic and surrounding healthy tissue (*25*). Previous studies have indicated that the tailocins target the strains of producer species or closely related species. Therefore, the target spectrum diversity of these tailocins plays a critical role in shaping the bacterial community (*39*). The comparative pangenomics approach used in this study captured the diversity of carotovoricin in the *Pectobacterium* genus. We identified a variable degree of conservation within the carotovorocin cluster and a genus-wide horizontal gene transfer of tail fiber loci. Here we discuss the observed evolutionary dynamics and suggest new research directions based on this work.

Evolutionary dynamics

In order to understand the scope of tailocin-mediated interactions in *Pectobacterium*, we conducted a genus-wide analysis of their organization and evolution. Here we discuss the possible implications of the observed evolutionary dynamics in *Pectobacterium* and compare them with other tailocin-producing species.

Conservation of both carotovoricin core functional genes and their chromosomal location across the *Pectobacterium* genus, shown previously ([23–25](#)) and at a higher resolution in this study, underscores the important role of this tailocin in competitive interactions. The competitive interactions experienced by the bacteria may act as a negative selection pressure to preserve the tailocin functionality, evidenced by carotovoricin loss in less competitive and host-deprived niches (Supplementary Results and Discussion). This selection pressure is further emphasized by the observation that carotovoricin was found to be present in a single copy and no other tailocin was identified in *Pectobacterium* species genomes. This is different from *Pseudomonas* species, where multiple tailocins exist with various integration sites, evolutionary history and regulatory genes ([40](#), [41](#)). We observed species-wide partial degradation of carotovoricin cluster in *P. betavasculorum* and *P. atrosepticum*. Although we found transcriptional evidence for genes in the partially deleted carotovoricin cluster of *P. atrosepticum*, the function of such partial tailocin biosynthesis gene clusters remains unclear, as they do not code for a complete functional unit. Nonetheless, this indicate that the phage tail-like bacteriocins are not immune to dynamic degradation, a phenomenon observed in their prophage ancestors where a rapid initial deactivation is followed by gradual decay ([5](#)). We hypothesize that their role in inter-bacterial interactions likely imposes a purifying selection pressure that protects tailocins from rapid initial deactivation, resulting in different domestication dynamics of tailocins compared to prophages.

Although carotovoricin is broadly conserved in the *Pectobacterium* genus, we observed varying degrees of conservation for multiple loci within the carotovoricin cluster, namely the nitric oxide deoxygenase gene, the tape measure gene and the tail fiber locus. We focused primarily on the TFLs because variation in their number and composition is a generic property of tailocins that directly determines the target specificity ([42](#)). Multiple sequence alignment has been used to estimate TFL haplotypes for *Pseudomonas* species tailocins ([21](#)). However, generating a multiple sequence alignment for genomic regions with complex structural variation, such as the TFLs, is challenging. Our pangenome approach to identify TFL haplotypes using homology group signatures overcomes this complexity.

We hypothesize that the *Pectobacterium* community in a plant host niche maintains a pool of carotovoricin tail fiber diversity to resist invasion by other closely related *Pectobacterium* species pathogens. Indeed, such a competitive exclusion mechanism is noted for R-tailocin producing *Pseudomonas chlororaphis* in mixed-species rhizosphere communities ([19](#), [43](#)). Additionally, the maintenance of TFL diversity in *Pectobacterium* might be over both space and time, as reported for another phytopathogen, *Pseudomonas* ([21](#)). Such a mechanism could be beneficial for the established colonizers in case of multi-species soft-rot infections, which are commonly observed in *Pectobacterium* species pathogens ([24](#), [28](#), [29](#)).

Unexpectedly, the known mechanism to achieve TFL diversity, Ein-mediated inversion ([20](#)), was absent

in approximately half of the carotovoricins within the *Pectobacterium* genus. However, we showed the existence of an additional mechanism to acquire the TFL diversity: horizontal transfer of TFLs across the *Pectobacterium* genus. Phylogenetic incongruence showed that HGT occurs in the region encoding the tail fiber of the carotovoricin and is likely mediated by site-specific recombination. Although the recombinational exchange of TFLs is known in other tailocin-producing bacteria (*18*), a combination of the two mechanisms, complementing each other, to maintain and circulate a pool of TFLs make *Pectobacterium* phytopathogens a unique case, to our knowledge. Additionally, it would be intriguing to test whether the HGT mechanism we found in the *Pectobacterium* genus can also be found in other genera involved in soft-rot plant diseases, for example, the *Dickeya* species.

Given the role played by carotovoricin in competitive interactions, it is puzzling why *P. atrosepticum* and *P. betavasculorum* never reacquired a functional copy of carotovoricin by HGT from other *Pectobacterium* species. One explanation could be that HGT of a complete carotovoricin locus is not possible due to its size or required homology, as we only observed transfer of TFLs in the *Pectobacterium* genus. Another possibility is that these species do not co-exist with other *Pectobacterium* species, because of niche adaptation or other inter-species antagonistic interactions (*29*). Co-existence is essential for any successful HGT between two bacteria. Multiple *Pectobacterium* species share ecological niches and even co-occur on both symptomatic and asymptomatic plant material (*27–29*). In line with this, common TFL haplotypes are observed between such co-existing species. In contrast, species that do not co-exist, specifically *P. brasiliense* and *P. parmentieri* (*29*), do not share any TFL haplotypes.

The success and stability of heterogeneous community depend on having the means to overcome competition from other closely related strains or species. Thus, the ability to exchange TFLs adds a new dimension to the carotovoricin-mediated community dynamics involving multiple *Pectobacterium* species and strains. Based on the observed TFL diversity and the sharing of TFL haplotypes across species, innovations in terms of functional diversity appear to be shared within the *Pectobacterium* genus.

New research directions

Our new insights on carotovoricin evolution in the *Pectobacterium* genus opens up various alleys for future data-driven research to study bacterial interactions. Here we discuss two promising directions.

First, our work provides a foundation for data-driven experimental design of tailocin competition assays to discover new interaction rules between multiple species. Recently, inter-genus killing activity of tailocin was shown in *Dickeya* species, which could target several *Pectobacterium* species (*25*). Therefore, a data-first approach could be used to select the candidate strains across multiple species and spanning different TFL haplotypes to perform tailocin interaction experiments. Such an experimental design would facilitate the development of tailocin interaction prediction models, which can further resolve the tailocin-mediated interactions in bacteria. Due to carotovoricin cluster conservation, presence of a single copy per strain and localization at the same gene neighborhood across the genus, we propose to use *Pectobacterium* as a model to study inter-species tailocin mediated interactions.

Second, a fundamental understanding of tailocin diversity is essential to design novel tailocin-based appli-

cations targeting a specific group of pathogens. Recently, several of these applications have been developed, ranging from phytopathogen control (*13*, *44*, *45*), engineering broad-range tailocins from phages (*46*, *47*) as well as designing broad-range chimeric phages based on the tailocins (*48*). The hyper-variable regions, such as the tail fiber loci, are difficult to compare at a large scale using a traditional comparative genomics approach. The pangenome approach taken here allows a detailed characterization of these TFLs and their contribution to tailocin evolution, which is essential for the development of tailocin-based biocontrol applications. With a growing number of high quality genomes in public databases, our comparative pangenomics approach provides a template to explore this as well as discover whether similar or novel mechanisms exist in other bacterial species.

Materials and Methods

Genome collection and pangenome construction

A *Pectobacterium* genus pangenome comprising 454 genomes was recently constructed by us (*24*). Briefly, we used PanTools (v4.1.1) (*31*, *32*) to build a De Bruijn graph pangenome, including gene annotations, functional annotations using InterProScan (v5.56-89.0) (*49*, *50*) and EggNOG-mapper (v2.1.10) (*51*, *52*). We optimized the sequence identity threshold at which genes are grouped together in homology groups. For example, a strict threshold can result in orthologous genes being distributed into multiple homology groups, consequently inflating the total number of homology groups in the pangenome. Conversely, a relaxed threshold would yield a smaller number of homology groups by incorrectly aggregating diverse genes into single group. To study the carotovoricin evolutionary dynamics, an optimal sequence identity threshold for the whole pangenome was estimated by evaluating various settings based on BUSCO gene sets. The PanTools homology grouping setting of 4 (equivalent to >65% sequence identity, 0.05 intersection rate for the k-mer set comparison, a 7.2 inflation rate for Markov clustering and a contrast factor of 5) optimally captures the sequence diversity of BUSCO gene sets across the *Pectobacterium* genus.

Phylogenetic analysis

Multiple sequence alignment of genes in each of the 1,949 core homology groups was performed using the PanTools *msa* function. These alignments were used to construct a maximum likelihood phylogeny using the *core_phylogeny* function (*53*), which uses a concatenation-based phylogenomic approach. Phylogenetic trees were processed using R (v4.2.1) packages Ape (v5.7.1) (*54*, *55*) and treeio (v1.22.0) (*56*) and visualized using ggtree (v3.6.2) (*57*) and ComplexHeatmap (v2.15.1) (*58*).

Carotovoricin identification

Prophages were predicted using the geNomad (v1.5.0) pipeline (*59*) and completeness was assessed using checkV (v1.0.1) (*60*). Any bacterial genomic regions at the prophage borders were identified using CheckV and subsequently trimmed. Putative prophages without any gene of viral origin were not considered. Each prophage was represented as a sequence of pangenomic homology groups, referred to as a prophage signature. When multiple prophage signatures from a genome could map in tandem to a larger prophage signature in another genome with ≥ 80% coverage, prophages from the first genome were considered as fragmented. Such

fragmented prophages were merged into a single one and labeled as fragmented (110 fragmented prophages were merged into 50). All putative prophages with length < 5kb were further excluded, resulting in a final collection of 1,369 prophages.

The pairwise similarity between prophage homology group signatures was quantified as the syntenic conservation of homology groups: the syntenic Jaccard index. To calculate the syntenic Jaccard index, dynamic programming was used to align two prophage signatures, with a score of +5 for a match and −2 for a mismatch of homology groups between two prophage signatures. The longest common subsequence between two prophage signatures, with at most two gaps and at least five matching homology groups, was considered a valid alignment of homology groups. The syntenic matching homology groups from such valid alignments were used to determined the intersection size while calculating the Jaccard index. The syntenic Jaccard index matrix for all-vs-all comparisons was clustered using complete linkage hierarchical clustering with the 'hclust' function in R. Next to various small clusters, this yielded one large group of 428 orthologous prophages. Among these, prophages from *P. carotovorum* matched the phage tail-like bacteriocin encoding region, called "Carotovoricin Er". All other prophages belonging to this cluster showed similar gene structures and hence were denoted as carotovoricin.

Transcriptomics evidence for carotovoricin in *P. atrosepticum*

Normalized count data for wild-type and polyphenol treated RNAseq samples for *P. atrosepticum* was downloaded from the GEO record GSE196675 (*33*). Genes from the partial carotovoricin locus were located in the region NC_004547.2:3113731-3119520. These genes were found to be differentially expressed upon polyphenol treatment.

Carotovoricin region synteny visualization

A 5kb region flanking carotovoricin cluster was included to provide broader context to understand integration at same or different site in the genome. An R script was written to extract the homology groups for carotovoricin regions and the data was stored in the JSON format required by clustermap.js (https://github.com/gamcil/clustermap.js). For species wise visualization for *P. brasiliense* and *P. versatile*, carotovoricins for the respective species were clustered based on the homology group present-absence matrix and the dendrogram was cut at the appropriate to select representatives for clinker plotting.

Comparative phylogenetics analysis of tail fiber locus

The carotovoricin region from tail protein (hg_22427604) to *ybiB* (hg_22427603) was defined as the tail fiber locus (TFL) and the remaining upstream region as the conserved locus. Haplotypes for the tail fiber locus were determined based on the unique ordering of homology groups in this region across all *Pectobacterium* genomes. Locus distance, estimated as a sequence-based distance between carotovoricin pairs, was calculated using the MinHash approach in Mash (v2.3) (*61*). Pairwise locus distance was calculated separately for the conserved locus and tail fiber locus. Pairwise species distances were calculated as the cophenetic distance using 'cophenetic.phylo()' function from the R package Ape (*54*, *55*). Cophenetic distances are calculated as pairwise distances between the pairs of tips of a phylogenetic tree using its branch lengths.

Carotovoricins from 16 genomes spanning three TFL haplotypes and representing 10 *Pectobacterium* species were selected for performing comparative phylogenetics analysis. The DNA sequences for conserved loci and TFLs for these carotovoricin regions were independently aligned using MAFFT (v7.508) (*62*). Maximum-likelihood phylogenetic trees for these two loci were estimated using IQ-TREE2 (2.2.0.3) (*53*). The trees were visualized using R (v4.2.1) package ggtree (v3.6.2) (*57*).

Visualization of structural variation in tail fiber locus

DNA sequences of carotovoricin regions from tail protein (hg_22427604) to *ybiB* (hg_22427603) were extracted as TFL sequences. The homology group level annotations from the PanTools' pangenome and DNA sequences were combined into a GenBank file format using the *seqret* tool from the EMBOSS software suite (v6.6.0.0) (*63*). These GenBank format sequences were aligned and visualized using the 'progressiveMauve' command of the Mauve v(2.4.0) program (*37, 38*).

Acknowledgements

We would like to acknowledge Inge van Duivenbode from Dutch General Inspection Service for agricultural seeds and seed potatoes (NAK) and Michiel J. C. Pel from the Netherlands Institute for Vectors, Invasive plants and Plant health (NIVIP) for providing *Pectobacterium* isolates and their valuable discussion in this research.

Funding

This research was funded by the Dutch Ministry of Economic Affairs in the Topsector Program "Horticulture and Starting Materials" under the theme "Plant Health" (project number: LWV20.235) and its partners (NVWA, NAK, Naktuinbouw and BKD).

Authors' contributions

LP, SS, TL, DR: Conceptualization. LP: Data curation, Formal analysis, Investigation, Software, Writing – original draft. SS, TL, DR: Supervision, Funding acquisition, Project administration. LP, SS, TL, DR, AK: Writing - review and editing. All authors reviewed the manuscript.

Conflicts of interest

The authors declare no competing interests.

Supplementary data

Supplementary tables: Supplementary_Tables.xlsx

Supplementary results and figures: Supplementary_Material.pdf

Data Availability

Genome assemblies used in this study were downloaded from the NCBI database. Assembly accessions and metadata for 454 genomes are listed in Supplementary Table S1. All scripts used for preprocessing, pangenome construction and analysis are available in the GitHub repository `https://lakhanp1.github.io/Pectobacterium_pangenome`. Notebooks to reproduce the analysis, tables and figures can be accessed at `https://lakhanp1.github.io/Pectobacterium_pangenome/scripts/notebooks/carotovoricin_analysis.html`. GitHub repository is archived in Zenodo and can be accessed using DOI: `https://doi.org/10.5281/zenodo.16585541`.

References

(*1*) Casjens, S. (2003). Prophages and bacterial genomics: what have we learned so far? *Molecular Microbiology 49*, 277–300.

(*2*) Varani, A. M., Monteiro-Vitorello, C. B., Nakaya, H. I., and Van Sluys, M.-A. (2013). The Role of Prophage in Plant-Pathogenic Bacteria. *Annual Review of Phytopathology 51*, 429–451.

(*3*) Sharma, V., Hünnefeld, M., Luthe, T., and Frunzke, J. (2023). Systematic analysis of prophage elements in actinobacterial genomes reveals a remarkable phylogenetic diversity. *Scientific Reports 13*, 4410.

(*4*) Costa, A. R., Monteiro, R., and Azeredo, J. (2018). Genomic analysis of Acinetobacter baumannii prophages reveals remarkable diversity and suggests profound impact on bacterial virulence and fitness. *Scientific Reports 8*, 15346.

(*5*) Bobay, L.-M., Touchon, M., and Rocha, E. P. C. (2014). Pervasive domestication of defective prophages by bacteria. *Proceedings of the National Academy of Sciences 111*, 12127–12132.

(*6*) Nakayama, K., Takashima, K., Ishihara, H., Shinomiya, T., Kageyama, M., Kanaya, S., Ohnishi, M., Murata, T., Mori, H., and Hayashi, T. (2000). The R-type pyocin of Pseudomonas aeruginosa is related to P2 phage, and the F-type is related to lambda phage. *Molecular Microbiology 38*, 213–231.

(*7*) YAMADA, K., HIROTA, M., NIIMI, Y., NGUYEN, H. A., TAKAHARA, Y., KAMIO, Y., and KANEKO, J. (2006). Nucleotide Sequences and Organization of the Genes for Carotovoricin (Ctv) from Erwinia carotovora Indicate That Ctv Evolved from the Same Ancestor as Salmonella typhi Prophage. *Bioscience, Biotechnology, and Biochemistry 70*, 2236–2247.

(*8*) Stice, S. P., Jan, H.-H., Chen, H.-C., Nwosu, L., Shin, G. Y., Weaver, S., Coutinho, T., Kvitko, B. H., and Baltrus, D. A. (2023). Pantailocins: phage-derived bacteriocins from *Pantoea ananatis* and *Pantoea stewartii* subsp. *indologenes*. *Applied and Environmental Microbiology 89*, ed. by Alexandre, G., e00929–23.

(*9*) Borowicz, M., Krzyżanowska, D. M., Narajczyk, M., Sobolewska, M., Rajewska, M., Czaplewska, P., Węgrzyn, K., and Czajkowski, R. (2023). Soft rot pathogen Dickeya dadantii 3937 produces tailocins resembling the tails of Peduovirus P2. *Frontiers in Microbiology 14*, DOI: `10.3389/fmicb.2023.1307349`.

(*10*) Scholl, D. (2017). Phage Tail–Like Bacteriocins. *Annual Review of Virology 4*, 453–467.

(11) Ghequire, M. G., and De Mot, R. (2014). Ribosomally encoded antibacterial proteins and peptides from Pseudomonas. *FEMS Microbiology Reviews 38*, 523–568.

(12) Yao, G. W., Duarte, I., Le, T. T., Carmody, L., LiPuma, J. J., Young, R., and Gonzalez, C. F. (2017). A Broad-Host-Range Tailocin from Burkholderia cenocepacia. *Applied and Environmental Microbiology 83*, e03414–16.

(13) Ishii, T. et al. (2024). Rhizoviticin is an alphaproteobacterial tailocin that mediates biocontrol of grapevine crown gall disease. *The ISME Journal 18*, wrad003.

(14) Kamimiya, S., Izaki, K., and Takahashi, H. (1977). Bacteriocins in *Erwinia aroideae* with Tail Like Structure of Bacteriophages. *Agricultural and Biological Chemistry 41*, 911–912.

(15) Ge, P., Scholl, D., Leiman, P. G., Yu, X., Miller, J. F., and Zhou, Z. H. (2015). Atomic structures of a bactericidal contractile nanotube in its pre- and postcontraction states. *Nature Structural & Molecular Biology 22*, 377–382.

(16) Desfosses, A., Venugopal, H., Joshi, T., Felix, J., Jessop, M., Jeong, H., Hyun, J., Heymann, J. B., Hurst, M. R. H., Gutsche, I., and Mitra, A. K. (2019). Atomic structures of an entire contractile injection system in both the extended and contracted states. *Nature Microbiology 4*, 1885–1894.

(17) Bhattacharjee, R., Nandi, A., Sinha, A., Kumar, H., Mitra, D., Mojumdar, A., Patel, P., Jha, E., Mishra, S., Rout, P. K., Panda, P. K., Suar, M., and Verma, S. K. (2022). Phage-tail-like bacteriocins as a biomedical platform to counter anti-microbial resistant pathogens. *Biomedicine & Pharmacotherapy 155*, 113720.

(18) Baltrus, D. A., Clark, M., Smith, C., and Hockett, K. L. (2019). Localized recombination drives diversification of killing spectra for phage-derived syringacins. *The ISME journal 13*, 237–249.

(19) Dorosky, R. J., Pierson, L. S., and Pierson, E. A. (2018). Pseudomonas chlororaphis Produces Multiple R-Tailocin Particles That Broaden the Killing Spectrum and Contribute to Persistence in Rhizosphere Communities. *Applied and Environmental Microbiology 84*, e01230–18.

(20) Nguyen, H. A., Tomita, T., Hirota, M., Kaneko, J., Hayashi, T., and Kamio, Y. (2001). DNA Inversion in the Tail Fiber Gene Alters the Host Range Specificity of Carotovoricin Er, a Phage-Tail-Like Bacteriocin of Phytopathogenic Erwinia carotovora subsp. carotovora Er. *Journal of Bacteriology 183*, 6274–6281.

(21) Backman, T. et al. (2024). A phage tail-like bacteriocin suppresses competitors in metapopulations of pathogenic bacteria. *Science 384*, eado0713.

(22) Itoh, Y., Izaki, K., and Takahashi, H. (1978). PURIFICATION AND CHARACTERIZATION OF A BACTERIOCIN FROM *ERWINIA CAROTOVORA*. *The Journal of General and Applied Microbiology 24*, 27–39.

(23) Arizala, D., and Arif, M. (2019). Genome-Wide Analyses Revealed Remarkable Heterogeneity in Pathogenicity Determinants, Antimicrobial Compounds, and CRISPR-Cas Systems of Complex Phytopathogenic Genus Pectobacterium. *Pathogens (Basel, Switzerland) 8*, E247.

(24) Pardeshi, L. A., van Duivenbode, I., Pel, M. J., Jonkheer, E. M., Kupczok, A., de Ridder, D., Smit, S., and van der Lee, T. A. J. (2025). Pangenomics to understand prophage dynamics in the Pectobacterium genus and the radiating lineages of Pectobacterium brasiliense. *Microbial Genomics 11*, 001392.

(25) Borowicz, M., Krzyżanowska, D. M., Sobolewska, M., Narajczyk, M., Mruk, I., Czaplewska, P., Pédron, J., Barny, M.-A., Canto, P. Y., Dziadkowiec, J., and Czajkowski, R. (2025). Tailocin-Mediated Interactions Among Soft Rot Pectobacteriaceae. *Molecular Ecology 34*, e17728.

(26) Backman, T., Burbano, H. A., and Karasov, T. L. (2024). Tradeoffs and constraints on the evolution of tailocins. *Trends in Microbiology 0*, DOI: `10.1016/j.tim.2024.04.001`.

(27) Motyka-Pomagruk, A., Zoledowska, S., Sledz, W., and Lojkowska, E. (2021). The occurrence of bacteria from different species of Pectobacteriaceae on seed potato plantations in Poland. *European Journal of Plant Pathology 159*, 309–325.

(28) Zhou, J., Hu, M., Hu, A., Li, C., Ren, X., Tao, M., Xue, Y., Chen, S., Tang, C., Xu, Y., Zhang, L., and Zhou, X. (2022). Isolation and Genome Analysis of Pectobacterium colocasium sp. nov. and Pectobacterium aroidearum, Two New Pathogens of Taro. *Frontiers in Plant Science 13*.

(29) Smoktunowicz, M., Jonca, J., Stachowska, A., May, M., Waleron, M. M., Waleron, M., and Waleron, K. (2022). The International Trade of Ware Vegetables and Orna-Mental Plants—An Underestimated Risk of Accelerated Spreading of Phytopathogenic Bacteria in the Era of Globalisation and Ongoing Climatic Changes. *Pathogens 11*, 728.

(30) Barny, M.-A., Thieffry, S., Gomes De Faria, C., Thebault, E., and Pédron, J. (2024). Bacterial pathogens dynamic during multi-species infections. *Peer Community Journal 4*, e49.

(31) Sheikhizadeh, S., Schranz, M. E., Akdel, M., de Ridder, D., and Smit, S. (2016). PanTools: representation, storage and exploration of pan-genomic data. *Bioinformatics (Oxford, England) 32*, i487–i493.

(32) Jonkheer, E. M., van Workum, D.-J. M., Sheikhizadeh Anari, S., Brankovics, B., de Haan, J. R., Berke, L., van der Lee, T. A. J., de Ridder, D., and Smit, S. (2022). PanTools v3: functional annotation, classification and phylogenomics. *Bioinformatics 38*, 4403–4405.

(33) Kang, J. E., Hwang, S., Yoo, N., Kim, B. S., and Chung, E.-H. (2022). A resveratrol oligomer, hopeaphenol suppresses virulence activity of Pectobacterium atrosepticum via the modulation of the master regulator, FlhDC. *Frontiers in Microbiology 13*, 999522.

(34) Jonca, J., Pirhonen, M., Waleron, M. M., Gawor, J., Mrozik, A., Smoktunowicz, M., Waleron, K., and Waleron, M. (2024). Comprehensive phenomic and genomic studies of the species, Pectobacterium cacticida and proposal for reclassification as Alcorniella cacticida comb. nov. *Frontiers in Plant Science 15*, DOI: `10.3389/fpls.2024.1323790`.

(35) Portier, P., Pédron, J., Taghouti, G., Dutrieux, C., and Barny, M.-A. (2020). Updated Taxonomy of Pectobacterium Genus in the CIRM-CFBP Bacterial Collection: When Newly Described Species Reveal "Old" Endemic Population. *Microorganisms 8*, 1441.

(36) Ben Moussa, H., Bertrand, C., Rochelle-Newall, E., Fiorini, S., Pédron, J., and Barny, M.-A. (2022). The diversity of soft rot Pectobacteriaceae along the Durance River stream in the south-east of France revealed by multiple seasonal surveys. *Phytopathology®*, DOI: `10.1094/PHYTO-12-21-0515-R`.

(37) Darling, A. C., Mau, B., Blattner, F. R., and Perna, N. T. (2004). Mauve: Multiple Alignment of Conserved Genomic Sequence With Rearrangements. *Genome Research 14*, 1394–1403.

(38) Darling, A. E., Mau, B., and Perna, N. T. (2010). progressiveMauve: Multiple Genome Alignment with Gene Gain, Loss and Rearrangement. *PLoS ONE 5*, e11147.

(39) Heiman, C. M., Vacheron, J., and Keel, C. (2023). Evolutionary and ecological role of extracellular contractile injection systems: from threat to weapon. *Frontiers in Microbiology 14*, 1264877.

(40) Ghequire, M. G., Dillen, Y., Lambrichts, I., Proost, P., Wattiez, R., and De Mot, R. (2015). Different Ancestries of R Tailocins in Rhizospheric Pseudomonas Isolates. *Genome Biology and Evolution 7*, 2810–2828.

(41) Hockett, K. L., Renner, T., and Baltrus, D. A. (2015). Independent Co-Option of a Tailed Bacteriophage into a Killing Complex in Pseudomonas. *mBio 6*, 10.1128/mbio.00452–15.

(42) Patz, S., Becker, Y., Richert-Pöggeler, K. R., Berger, B., Ruppel, S., Huson, D. H., and Becker, M. (2019). Phage tail-like particles are versatile bacterial nanomachines - A mini-review. *Journal of Advanced Research 19*, 75–84.

(43) Dorosky, R. J., Yu, J. M., Pierson, L. S., and Pierson, E. A. (2017). Pseudomonas chlororaphis Produces Two Distinct R-Tailocins That Contribute to Bacterial Competition in Biofilms and on Roots. *Applied and Environmental Microbiology 83*, e00706–17.

(44) Baltrus, D. A., Clark, M., Hockett, K. L., Mollico, M., Smith, C., and Weaver, S. (2022). Prophylactic Application of Tailocins Prevents Infection by Pseudomonas syringae. *Phytopathology® 112*, 561–566.

(45) Kouzai, Y., Sagehashi, Y., Watanabe, R., Kajiwara, H., Suzuki, N., Ono, H., Naito, K., and Akimoto-Tomiyama, C. (2024). BglaTNB6, a tailocin produced by a plant-associated nonpathogenic bacterium, prevents rice seed-borne bacterial diseases. *PLOS Pathogens 20*, e1012645.

(46) Williams, S., Gebhart, D., Martin, D., and Scholl, D. (2008). Retargeting R-type pyocins to generate novel bactericidal protein complexes. *Applied and Environmental Microbiology 74*, 3868–3876.

(47) Woudstra, C., Sørensen, A. N., Sørensen, M. C. H., and Brøndsted, L. (2024). Strategies for developing phages into novel antimicrobial tailocins. *Trends in Microbiology*, DOI: 10.1016/j.tim.2024.03.003.

(48) Yao, G., Le, T., Korn, A. M., Peterson, H. N., Liu, M., Gonzalez, C. F., and Gill, J. J. (2023). Phage Milagro: a platform for engineering a broad host range virulent phage for Burkholderia. *Journal of Virology 97*, e0085023.

(49) Jones, P. et al. (2014). InterProScan 5: genome-scale protein function classification. *Bioinformatics 30*, 1236–1240.

(50) Blum, M. et al. (2021). The InterPro protein families and domains database: 20 years on. *Nucleic Acids Research 49*, D344–D354.

(51) Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., and Huerta-Cepas, J. (2021). eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Molecular Biology and Evolution 38*, 5825–5829.

(52) Hernández-Plaza, A., Szklarczyk, D., Botas, J., Cantalapiedra, C. P., Giner-Lamia, J., Mende, D. R., Kirsch, R., Rattei, T., Letunic, I., Jensen, L. J., Bork, P., von Mering, C., and Huerta-Cepas, J. (2023). eggNOG 6.0: enabling comparative genomics across 12 535 organisms. *Nucleic Acids Research 51*, D389–D394.

(53) Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution 37*, 1530–1534.

(54) Paradis, E., Claude, J., and Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R language. *Bioinformatics 20*, 289–290.

(55) Paradis, E., and Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics 35*, 526–528.

(56) Wang, L.-G., Lam, T. T.-Y., Xu, S., Dai, Z., Zhou, L., Feng, T., Guo, P., Dunn, C. W., Jones, B. R., Bradley, T., Zhu, H., Guan, Y., Jiang, Y., and Yu, G. (2020). Treeio: An R Package for Phylogenetic Tree Input and Output with Richly Annotated and Associated Data. *Molecular Biology and Evolution 37*, 599–603.

(57) Xu, S., Li, L., Luo, X., Chen, M., Tang, W., Zhan, L., Dai, Z., Lam, T. T., Guan, Y., and Yu, G. (2022). Ggtree: A serialized data object for visualization of a phylogenetic tree and annotation data. *iMeta 1*, e56.

(58) Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics 32*, 2847–2849.

(59) Camargo, A. P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain, P. S. G., Nayfach, S., and Kyrpides, N. C. (2023). Identification of mobile genetic elements with geNomad. *Nature Biotechnology*, 1–10.

(60) Nayfach, S., Camargo, A. P., Schulz, F., Eloe-Fadrosh, E., Roux, S., and Kyrpides, N. C. (2021). CheckV assesses the quality and completeness of metagenome-assembled viral genomes. *Nature Biotechnology 39*, 578–585.

(61) Ondov, B. D., Treangen, T. J., Melsted, P., Mallonee, A. B., Bergman, N. H., Koren, S., and Phillippy, A. M. (2016). Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biology 17*, 132.

(62) Katoh, K., Rozewicki, J., and Yamada, K. D. (2019). MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics 20*, 1160–1166.

(63) Rice, P., Longden, I., and Bleasby, A. (2000). EMBOSS: the European Molecular Biology Open Software Suite. *Trends in genetics: TIG 16*, 276–277.

Figure 1: Carotovoricin cluster across the *Pectobacterium* genus. (a) An illustration of a carotovoricin particle where structural units are colored to match the carotovoricin biosynthesis cluster alignment shown in panel b. (b) Carotovoricin biosynthesis gene clusters are ordered along the phylogeny of the *Pectobacterium* genus. Genes in 5kb regions flanking the carotovoricin clusters are shown in white. Genes belonging to the same homology group are connected with vertical black lines.

Figure 2: Carotovoricin locus variation. (a) An illustration of gene-level variation in carotovoricin cluster, with gene presence-absence variation shown as loops. Flanking genes are represented as white arrows. The conserved locus of carotovoricin is defined as genes encompassing holin and tail protein. The variable tail fiber locus is defined as the genes between the tail protein and *ybiB*. The unique homology group combinations in the TFLs across the *Pectobacterium* genus are used to define haplotypes. Pangenome homology group identifiers for the relevant genes are mentioned in parenthesis. (b) The UpSet plot shows the number of *Pectobacterium* species that share a TFL haplotype. Species are shown along the rows of the UpSet plot and the black dotted lines show the intersection of TFL haplotypes. The top bar chart shows number of unique TFL haplotypes shared in the respective intersection. The two bar charts on the right show total number of TFL haplotypes per species and number of genomes for each species in the pangenome, respectively.

Figure 3: Carotovoricin polymorphism due to tail fiber locus diversity. (a) A core-genome phylogenetic tree is shown for the 365 *Pectobacterium* genomes that contain the carotovoricin cluster in a single contig. (b) A species key marking species from left to right: *P. brasiliense*, *P. polaris*, *P. parvum*, *P. aquaticum*, *P. quasiaquaticum*, *P. versatile*, *P. carotovorum*, *P. odoriferum*, *P. actinidiae*, *P. parmentieri*, *P. wasabiae*, *P. punjabense*, *P. polonicum*, *P. zantedeschiae*, *P. peruviense*, *P. aroidearum*, *Pectobacterium sp. CFBP8739*, *P. colocasium*, *P. fontis* and *P. cacticida*. (c) The presence of 52 TFL haplotypes is shown as yellow bands. Only the haplotypes that are present in two or more genomes (52 out of 93) are shown.

Figure 4: Sequence diversity of the tail fiber locus. Pairwise locus distances for carotovoricin regions, estimated using Mash, were compared against the species evolutionary distance, estimated from the *Pecto-bacterium* core-SNP phylogeny derived from the pangenome. These comparisons are shown for two loci in the carotovoricin: (a) the conserved locus, and (b) the variable TFL. The comparisons in which both loci belong to the same TFL haplotype are colored in yellow in both scatter plots for reference. Such matching haplotypes show smaller locus distances, also at higher species distances, for inter-species comparisons of TFLs (b). This trend is not observed for the conserved loci in (a).

Figure 5: Horizontal gene transfer of tail fiber loci. (a) An unrooted maximum-likelihood phylogenetic tree for the conserved locus of carotovoricin from 16 genomes. (b) An unrooted maximum-likelihood phylogenetic tree generated from the TFLs from 16 genomes comprising 3 different haplotypes. For both the trees, bootstrap values are shown next to internal nodes and tips are colored as per the TFL haplotype. (c) Complex structural rearrangements of tail fiber locus (TFL) haplotypes are shown in the form of Mauve collinear blocks for carotovoricin loci in *P. brasiliense* accessions with the invertase gene *ein*. (d) Mauve collinear blocks in TFLs of various *Pectobacterium* species are shown and labels for TFLs that have invertase *ein* are colored green. Locally collinear blocks are shown as colored boxes above (forward) and below (reverse) the black horizontal line that represents the TFL haplotype region in a genome. Conserved blocks across the genomes are linked by vertical lines. Inside each block, a similarity profile is drawn which represents the average level of conservation across the TFL haplotypes. White areas represent sequences specific to a TFL and are not aligned.