A high-throughput heterologous expression platform for plant synthetic biology based on Arabidopsis suspension cells

Lukas Meile[1,2,*], Germán Alonso-Tolo[1], Zeltia Ferreiro-Eiras[1], Xi Jiang[1], Stefan Burén[1], and Luis M. Rubio[1,*]

[1] Centro de Biotecnología y Genómica de Plantas, Universidad Politécnica de Madrid e Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria/Consejo Superior de Investigaciones Científicas, Madrid 28223, Spain.

[2] Present address, Institute of Microbiology, ETH Zurich, 8093 Zurich, Switzerland.

*To whom correspondence should be addressed: Luis M. Rubio: +34 910679189 – luis.rubio@csic.es; Lukas Meile lmeile@ethz.ch.

Other e-mail addresses: Germán Alonso-Tolo (german.atolo@upm.es), Zeltia Ferreiro-Eiras (zeltiaferreiroeiras@gmail.com), Xi Jiang (xi.jiang@upm.es), and Stefan Burén (stefan.buren@upm.es).

Word count: 3082 words and 6 Figures. Supplementary data: 6 figures and 4 supplementary tables.

Running title: High-Throughput Heterologous Expression in Arabidopsis Cells

Highlight (30 words max.)

We developed a fast and scalable expression platform in Arabidopsis suspension
cells, enabling transient and stable multigene expression for applications in plant
synthetic biology such as nitrogenase engineering.

Abstract (200 words max, graphical abstract possible)

Efficient heterologous expression platforms are essential for plant synthetic biology,
particularly for engineering complex multigene pathways. Here, we establish a high-
throughput system for both transient and stable transformation of *Arabidopsis*
thaliana suspension cells using plant cell pack infiltration. This method requires no
specialized equipment or consumables and is compatible with several cell lines. It
enables rapid generation of 100 g of transgenic cells within two weeks and allows
expression of at least 6 stacked genes from a single construct. We characterized
constitutive promoters for gene expression in Arabidopsis cells and validated plastid
targeting peptides. A library of NifB homologs was screened for expression and
solubility and several archaeal variants suitable for plant expression were identified.
We further engineered stable cell lines expressing up to six genes, encoding the NifB
module components NifU, NifS, FdxN, and NifB, demonstrating that the newly
developed platform integrates into an established workflow for nitrogenase
engineering. The platform accelerates design–build–test cycles and facilitates the
production of delicate proteins that require large amounts of transgenic biomass. It
thus represents a versatile and scalable tool for advancing synthetic biology and for
tackling major biotechnological challenges, such as biological nitrogen fixation.

Keywords (6-9, alphabetical order)

Arabidopsis thaliana transformation

Heterologous expression

Nitrogen fixation protein engineering

Plant biotechnology

Plant cell packs

Plant suspension cultures

Plant synthetic biology

Plastid protein expression

Protein solubility

Abbreviations

54	2,4-D	2,4-dichlorophenoxyacetic acid
55	BNF	biological nitrogen fixation
56	CTP	chloroplast targeting peptide
57	MS	Murashige and Skoog
58	NAA	1-naphthaleneacetic acid
59	PCP	plant cell pack
60	TS	Twin-Strep

62 Introduction

Plant cell suspension cultures provide a valuable system for heterologous protein
production with several advantages over whole plants (Fischer *et al.*, 1999; Menges
and Murray, 2004). They offer faster growth rates, simpler cultivation protocols, and
reduced biological complexity due to a more uniform cell population. The
homogeneous nature of suspension cultures facilitates reproducibility and scalability,
which are essential for synthetic biology applications that require precise control over
gene expression. Furthermore, the direct contact of all cells with the growth medium
enables rapid and uniform exposure to external compounds such as expression
inducers, signaling molecules, or nutrients. Together, these features make plant
suspension cultures a powerful platform for accelerating design–build–test cycles in
plant synthetic biology.

Several plant species have been employed for cell suspension-based expression,
with tobacco (*Nicotiana tabacum*), rice (*Oryza sativa*), and carrot (*Daucus carota*)
being the most widely used ones (Santos *et al.*, 2016). *Arabidopsis thaliana*, although
used less frequently, offers some advantages that are yet to be fully exploited. As the
most extensively studied model plant, Arabidopsis provides many genetic resources,
including mutant collections and a well-annotated genome (Provart *et al.*, 2016). In
addition, certain Arabidopsis cell lines retain the ability to form functional chloroplasts,
allowing photomixotrophic growth (McCarthy *et al.*, 2016; Sello *et al.*, 2017) or even
photoautotrophy when gradually reducing the external carbon source (Hampp *et al.*,
2012). Chloroplast development in suspension cells provides opportunities to study
and engineer plastid-targeted proteins and chloroplast-related processes (Li *et al.*,
2021; McCarthy *et al.*, 2016; Sello *et al.*, 2016). Although Arabidopsis suspension
cells have been used in various studies, and protocols for stable transformation and
transient expression have been reported (Burén *et al.*, 2011; García-León *et al.*,
2018; Menges and Murray, 2004; Van Leene *et al.*, 2011), throughput and fast
scalability remain limited. Furthermore, as for other plant systems, integration of
multi-gene modules is still not a routine application (Rizzo *et al.*, 2023).

Engineering plants to fix atmospheric dinitrogen through direct transfer of prokaryotic
nitrogen fixation (*nif*) genes is a key challenge of plant biotechnology (Curatti and
Rubio, 2014; Oldroyd and Dixon, 2014). Nitrogenase is the prokaryotic multimeric
metalloenzyme responsible for biological nitrogen fixation (BNF). Its function requires
the coordinated expression of more than 10 Nif proteins, many of which are involved

in the biosynthesis of nitrogenase-specific essential Fe-S cofactors (Burén *et al.*,
2020). In addition to the high number of required Nif proteins, oxygen-sensitivity and
low solubility of nitrogenase and other Nif proteins are major obstacles for
engineering a functional plant nitrogenase (Burén and Rubio, 2018). Current
strategies to engineer Nif proteins for eukaryotic expression have exploited the
natural diversity of Nif proteins by systematically testing natural Nif protein variants
for expression and solubility (Burén *et al.*, 2019; Jiang *et al.*, 2022; Jiang *et al.*,
2021).

The nitrogen fixation pathway can be divided into functional modules, each mediating
an independent step of nitrogenase assembly. These modules can be optimized
independently, reducing complexity and providing a tractable route toward full
pathway reconstruction. Beyond optimizing protein expression, successful
nitrogenase engineering is also thought to require subcellular targeting, with
mitochondria and plastids being the two main candidate compartments. Mitochondria
offer a favorable redox environment in which oxygen is consumed by respiration, but
necessarily require protein import, which might be a limiting step for some Nif
proteins (Oldroyd and Dixon, 2014). Plastids, by contrast, are genetically more
accessible and support high expression levels (Boehm and Bock, 2019; Scotti *et al.*,
2012). Furthermore, reducing power is abundant in chloroplasts and plastid electron
transport components are compatible with nitrogenase (Yang *et al.*, 2017). Although
oxygen production during photosynthesis poses a major obstacle (Aznar-Moreno *et*
al., 2021; Eseverri *et al.*, 2020b; Ivleva *et al.*, 2016), strategies such as targeting non-
photosynthetic plastids or temporally separating nitrogenase expression from
photosynthesis have been proposed (Curatti and Rubio, 2014; Dixon *et al.*, 1997). In
this context, Arabidopsis suspension cultures represent a particularly versatile model.
They can be cultured under heterotrophic or photomixotrophic conditions, enabling
studies on how oxygen production in chloroplasts affects Nif protein stability.

Biotechnological challenges such as nitrogenase engineering highlight the need for
efficient platforms that allow iterative and flexible testing of complex biosynthetic
pathways in plants. Here, we describe a platform for both transient and stable
transformation of Arabidopsis suspension cells and demonstrate its utility by
integrating it into an established workflow for nitrogenase engineering. We
characterized a suite of constitutive promoters and performed expression and
solubility screenings of a NifB library, providing new promising NifB candidates
suitable for plant expression. We furthermore showed that the platform supports the
stacking of at least six transgenes and allows quick scalability for protein purification.
By enabling rapid protein production and synthetic pathway assembly in plant cells,
this simple and low-cost platform contributes to advancing experimental approaches
in plant synthetic biology.

136 **Materials and methods**

Plant cell lines and bacterial strains:

The Arabidopsis cell lines YG-1 and T87 were provided by the Riken Plant Cell Bank
(Yokohama, Japan). PSB-L cells were a gift from David Alabadí (IBMCP, Valencia,
Spain) and Col-0 and MM1 cells were a gift from Laszlo Bako (Umeå University,
Sweden). The *Escherichia coli* strain TOP10 was used for all cloning steps. The
Agrobacterium tumefaciens strain GV3101::pMP90 was a gift from Vicente Rubio and
used for PCP infiltration if not stated otherwise.

Arabidopsis cell suspension cultures:

Col-0 cells were routinely cultured in Murashige and Skoog (MS) medium including
vitamins with 3% sucrose, pH 5.7, supplemented with 2,4-dichlorophenoxyacetic acid
(2,4-D, 0.24 µg/ml) and kinetin (0.014 µg/ml) with weekly passages of 7 ml culture
into 40 ml of fresh medium in 250-ml flasks. MM1 and PSB-L cells were routinely
cultured in 1x MS Basal Salts medium with 3% sucrose, pH 5.7, supplemented with
thiamine (0.4 µg/ml), myo-inositol (0.1 mg/ml), 1-naphthaleneacetic acid (NAA, 0.5
µg/ml) and kinetin (0.05 µg/ml) with weekly passages of 2 ml culture into 48 ml of
fresh medium and 10 ml culture into 40 ml of fresh medium, respectively. YG-1 cells
were cultured in 1x MS Basal Salts medium with 200 mg/L KH_2PO_4, 3% sucrose, pH
5.8, supplemented with thiamine (0.4 µg/ml), myo-inositol (0.1 mg/ml), and 2,4-D (0.2
µg/ml) with weekly passages of 5 ml sedimented cells into 40 ml of fresh medium.
T87 cells were cultured in MS medium including vitamins with 1.5% sucrose, pH 5.7,
supplemented with NAA (0.224 µg/ml) with biweekly passages of 1.2 ml of culture
into 40 ml fresh medium. Col-0 and YG-1 cells were routinely cultured at 25°C
without light and MM1, PSB-L, and T87 cells were cultured at 22°C under a 16-h
photoperiod.

Transient and stable plant cell transformation:

The methods for transient and stable transformation of Arabidopsis suspension cells
are summarized in Fig. 1. Precultures for transient expression assays were prepared
as described above and grown for 3-7 days before transformation. Transient
expression assays were based on the plant cell pack (PCP) infiltration method
(Rademacher *et al.*, 2019). For PCP infiltration of Col-0, the medium was first
removed by pressing the tip of a serological pipette against the flask or tube bottom
followed by slow aspiration, similarly as described (Cortese *et al.*, 2021). The
resulting compacted cells were then resuspended in Paul's Medium (Buschmann,
2016) to a density that barely allowed pipetting of the suspension with wide-bore tips.
For small PCPs of approximately 200 mg (fresh weight), 500 µl of cell suspension
was loaded onto the top part of 1000-µl filter pipette tips. The excess liquid was
removed by vacuum for approximately 10 s, resulting in PCP formation. *A.*
tumefaciens suspensions were prepared from 1-day-old precultures as described
before (Gengenbach *et al.*, 2020), but without acetosyringone, at OD_{600nm} 0.4 if not
stated otherwise. PCPs were infiltrated with 300 µl of *A. tumefaciens* suspension and
incubated at 25°C for 1 h before removing the excess liquid by vacuum for
approximately 30 s. The infiltrated PCPs were incubated inside a tip box equipped

with sterile wet paper to maintain high humidity for 2-4 days at 25°C in the dark,
unless indicated otherwise.

For small-scale stable transformations of around 200 mg of starting biomass, PCPs
were prepared the same way as for transient expression assays. For larger PCPs of
approximately 8 g, cell suspensions were prepared similarly but 20 ml was loaded
onto a Zymo-Spin V-PS midiprep column assembly (with the DNA-binding matrix
removed) with a 15-ml Reservoir-X (Zymo Research, CA, USA). The excess medium
was removed by gravity and 20 ml of *A. tumefaciens* suspension was passed through
the packed cells by gravity. The excess liquid was removed by centrifugation in a
swing-bucket rotor at 200 g for 2 min. The compacted, semi-dry PCPs were then
transferred to a 50-ml centrifugation tube with a partially closed lid, sealed with
Micropore tape (3M Health Care, USA) and incubated in a moist box. After 2-4 days,
small PCPs were gently resuspended in 4 ml of cultivation medium described above
containing carbenicillin (250 µg/ml) and vancomycin (165 µg/ml) to select against *A.*
tumefaciens. For cell lines Col-0 and YG1, selection with hygromycin was applied at
this point (20 µg/ml), whereas for MM1 hygromycin was only applied 2-3 days after
resuspension. For large PCPs, cells were resuspended in 40 ml of growth medium,
agitated on an orbital shaker at 100 rpm for 30 min before replacing the medium and
supplementing with antibiotics as for small PCPs.

198 Plasmids, molecular cloning, gene library design and codon optimization:

Molecular cloning was performed using the Modular Cloning (MoClo) standard
(Weber *et al.*, 2011; Werner *et al.*, 2012) as described before (Eseverri *et al.*,
2020b). For domestication of Level 0 parts, the Phusion Hot Start II DNA Polymerase
(Thermo Fisher) was used with the primers and templates listed in Table S1. The
sequence of all generated constructs used in this study was verified using a Sanger
sequencing service (Macrogen, Madrid, Spain) or an Oxford Nanopore whole-
plasmid sequencing service (Plasmidsaurus, Inc., Eugene, USA). All constructs and
the individual parts used for their assembly are listed in Table S2. The MoClo Plant
Parts Kit and pUAP1 were a gift from Nicola Patron (Addgene kit #1000000047, and
plasmid #63674, respectively) and the MoClo Toolkit was a gift from Sylvestre
Marillonnet (Addgene kit #1000000044). 35S:RUBY was a gift from Yunde Zhao
(Addgene plasmid #160908) and pTHsp18.2 (GB0035) was a gift from Diego Orzaez
(Addgene plasmid #68186). The *nifB* library was designed by keyword search in the
Uniprot database (Bateman *et al.*, 2021). Archaeal NifB homologs were identified by
discarding protein sequences lacking the six described NifB characteristic motifs
(Arragain *et al.*, 2017), resulting in 321 candidate archaeal NifB homologs. Because
Nif proteins from thermophilic organisms have been successfully expressed in
eukaryotes before (Burén *et al.*, 2017; Jiang *et al.*, 2022), NifB homologs from
thermophilic sources were considered as high-priority candidates. To identify them,
NCBI taxon IDs associated with the Uniprot search results were used to obtain
information on growth temperature and habitats in the Bac*Dive*, JGI IMG/D and
GOLD databases (Chen *et al.*, 2021; Mukherjee *et al.*, 2025; Schober *et al.*, 2025).
Furthermore, homologs from known diazotrophs were considered good candidates.

Diverse source organisms were also considered by not including more than two
homologs from the same taxonomic order. The workflow for the library design is
summarized in Fig. S1. To improve expression in Arabidopsis, all *nifB* sequences as
well as the sequences of *nifU*, *nifS*, *fdxN*, and the TS-tag were codon optimized using
the DNA Chisel tool (Zulkower and Rosser, 2020). The optimization constraints were
high GC and GC3 contents of at least 50 and 69%, respectively and avoiding BpiI,
BsaI, and BsmBI restriction sites. The optimization objectives included codon
optimization for Arabidopsis, avoiding rare codons (minimum frequency = 0.25),
avoiding hairpins (stem size = 15, window = 200) as well as avoiding the instability
motifs considered by previous work (Eseverri *et al.*, 2020b). The optimized
sequences flanked by BsaI sites for cloning (Table S3) were ordered from GenScript
(USA).

Luciferase reporter assay:

Promoter activities were measured as previously reported (Li *et al.*, 2017) with
modifications. PCPs were lysed in a buffer containing 100 mM Tris-HCl (pH 8.6), 200
mM NaCl, 10% glycerol, 0.1% protease inhibitor cocktail (P9599, Sigma-Aldrich) with
4:1 (v/w) buffer/biomass ratio by metal bead (7 mm) beating in a Qiagen Retsch
MM300 TissueLyser for 60 s. Ten µl cell lysate was mixed with 100 µl luciferase
buffer containing 25 mM glycyl-glycine (pH 8), 14 mM K_2HPO_4-KH_2PO_4 (pH 8), 4 mM
EGTA, 2 mM ATP, 1 mM DTT, 15 mM $MgSO_4$, 150 µg/ml D-luciferin in a white 96-
well plate. Luminescence was measured with an integration time of 2 s in a
Varioskan LUX plate reader (Thermo Scientific). For normalization of luciferase
levels, every construct contained the same GUS reporter cassette. GUS activity was
measured in the same device by mixing 10 µl cell lysate with 50 µl buffer containing
10 mM Tris-HCl (pH 8.0), 1 mM 4-methylumbelliferyl-β-D-glucuronide, and 2 mM
$MgCl_2$ in a black 96-well plate. The reaction was incubated at 37°C for 30 min with
100-ms measurements in 15-s intervals with an excitation wavelength of 365 nm with
a bandwidth of 12 nm and an emission wavelength of 455 nm. The GUS activity was
determined by calculating the change in fluorescence over time in the linear phase.
The promoter activity was quantified by dividing the luminescence intensity by the
GUS activity and normalized to the strongest promoter of each run.

Confocal microscopy:

PCPs of the cell line Col-0 were infiltrated with *A. tumefaciens* harbouring the
plasmids 1608-1614 (Table S2) as described above and harvested after 2 days by
resuspension in growth medium. Confocal laser scanning microscopy was performed
with a Zeiss LSM 880 microscope equipped with a Plan-Apochromat 40X/1.2 water-
immersion objective and 488-nm and 561-nm lasers. The detection ranges were 493-
574 nm for GFP and 578-696 nm for mCherry. Images were acquired using the ZEN
2.6 Black software (Zeiss) and processed with the Fiji software (Schindelin *et al.*,
2012); modifications included cropping, maximum-intensity projections, and
adjustments of brightness and contrast.

Protein extraction, immunoblotting and purification:

Protein extraction for immunoblotting was performed similarly as described (Burén *et al.*, 2017) with modifications, if not stated otherwise. PCPs were added to a 2-ml Eppendorf tube and subjected to 4 freeze-thaw cycles with liquid N_2. Lysis buffer, containing 100 mM Tris-HCl (pH 8), 150 mM NaCl, 10 mM $MgCl_2$, 0.2% NP-40, 5% glycerol, 5 mM β-mercaptoethanol, 5 mM EDTA, and 0.5% protease inhibitor cocktail (Sigma P9599; added immediately before use), was added at a ratio of 2:1 (v/w) with a 7-mm diameter steel ball and the cells were homogenized in a Qiagen Retsch MM300 TissueLyser for 60 s, followed by a 15-min incubation at 4°C in a tube rotator. The resulting homogenate was mixed at a ratio of 1:1 with 2x Laemmli buffer (125 mM Tris-HCl pH 6.8, 4% SDS, 20% (v/v) glycerol, 10% β-mercaptoethanol, 0.005% bromophenol blue) and heated for 10 min at 98°C, yielding the total cell extract. For the soluble extract, the cell homogenate was centrifuged at 20,000 g for 15 min and the supernatant containing the soluble proteins was treated with 2x Laemmli buffer as described for the total extract. The heat-denatured protein samples were centrifuged at 20,000 for 10 min before SDS-PAGE electrophoresis and immunoblotting. After blotting, membranes were stained with a Ponceau S solution (0.2% in 5% acetic acid) as transfer control. The Twin-Strep (TS) tag was detected with a monoclonal mouse StrepII antibody (GenScript A01732, 1:5,000), NifU and NifS with rabbit polyclonal anti-NifU and anti-NifS antibodies (Rubio lab, 1:20,000), the HA-tag with a rat monoclonal antibody (Roche 12013819001, 1:5,000), GFP with a mouse monoclonal antibody (Santa Cruz sc-9996, 1:2,000), and RbcL with a rabbit polyclonal antibody (Agrisera AS03 037, 1:5,000). Secondary HRP-conjugated antibodies were used in 2% milk powder in TBS-T diluted 1:20,000 (anti-mouse and anti-rabbit) or 1:5,000 (anti-rat). NifB purifications were performed from Col-0 cells grown in the dark. Cells were transformed with plasmid pN2LM282 as described above and scaled up to four 250-ml shake flasks with 60 ml culture each. To collect cells, the grown transformant cultures were bubbled with pure N_2 and then moved to an anaerobic chamber, where the following steps were performed. Cells were harvested by filtration through a ZymoPURE Syringe Filter-X (Zymo Research, CA, USA) followed by drying with tissue paper and snap-freezing and storage in N_2. Cells were homogenized in a blender (Oster Classic 4655) for 5 min at maximum speed with lysis buffer (100 mM Tris-HCl [pH 8.6], 200 mM NaCl, 10% glycerol, 5 mM β-mercaptoethanol, 2 mM sodium dithionite, 1 mM phenylmethylsulfonyl fluoride, 1 mg/ml leupeptin, 5 mg/ml DNase I) with a buffer/biomass ratio of 2:1. The homogenate was then filtered through 2 layers of miracloth. Lysate clearing, NifB purification and concentration was performed as described (Jiang *et al.*, 2022), but with a 5-ml Strep-Tactin®XT 4Flow® column and a peristaltic pump setup. Protein purity was assessed by SDS-PAGE analysis followed by peptide mass fingerprinting of individual bands excised from the SDS-PAGE gel. Mass spectrometry analysis was performed by a service of the Proteomic Unit at Universidad Complutense de Madrid.

Results

A transient expression system for Arabidopsis suspension cells.

We first sought to improve the throughput of the *A. tumefaciens*-mediated transient expression workflow for Arabidopsis suspension cells. The method was based on a recently developed approach of plant cell pack (PCP) infiltration, which was originally established for tobacco BY-2 cells (Rademacher *et al.*, 2019). For easy and individual handling of PCPs, they were cast into the top part of filter pipette tips and the RUBY (He *et al.*, 2020) and GFP reporters were used to assess successful transformation. We first assessed four different *A. tumefaciens* strains for their suitability for PCP infiltration with different Arabidopsis cell lines and found that the strain GV3101::pMP90 led to RUBY expression in the cell line Col-0 (Fig. 2A) and YG-1 (Fig. S2A), but not in the cell lines MM1, PSB-L, and T87 (Fig. 2A). Strain AGL-1 led to RUBY expression in Col-0 but not in the other tested cell lines MM1, PSB-L, and T87 (Fig. 2A). Similarly, strain EHA105 led to RUBY expression only in Col-0 but not in the other tested cells MM1 and PSB-L (Fig. S2A). Finally, PCPs infiltrated with the strain LBA4404 never produced visible RUBY signal (Fig. 2A). The culture method of *A. tumefaciens* did not affect infiltration success, as cultures grown in liquid or solid media both led to RUBY expression (Fig. S2B). Since both Col-0 and YG-1 were routinely cultured in darkness and the other cell lines under a 16-h photoperiod, we hypothesized that cultivation under light might hamper transformation. Col-0 and MM1 cells grown either in darkness or under a 16-h photoperiod were then compared for their receptiveness to PCP infiltration. MM1 cells grown under darkness were more amenable to PCP infiltration; however, only small PCPs cast from 150 µl cell suspension but not from 500 µl consistently yielded high transformation efficiency (Fig. 2A & D). A similar effect of light conditions on transformation efficiency was observed for Col-0 cells, for which transformation was abolished when cultured under a 16-h photoperiod (Fig. 2C).

Next, we tested whether Arabidopsis PCP infiltration can be improved by the co-expression of the inhibitor of silencing p19. Co-expression of GFP and p19 in Col-0 PCPs was performed, either using a single *A. tumefaciens* strain harbouring both genes on the same plasmid or using two co-infiltrated strains, each harbouring one of the genes. To avoid the detection of GFP expressed in *A. tumefaciens* cells, which has been described before (Buschmann, 2016), a *gfp* version containing the *DEM1* intron (Christie *et al.*, 2011) was deployed. The p19 improved GFP expression irrespectively of whether *gfp* and *p19* were present on the same plasmid or co-infiltrated (Fig. 2E).

PCP infiltration is a fast method for generating stably transformed plant cells.

We hypothesized that infiltrated PCPs could serve as an efficient source for stably transformed cell cultures. This hypothesis was tested with the *A. tumefaciens* strain GV3101::pMP90 because this strain has been successfully used for stable transformation of Arabidopsis suspension cells (Burén *et al.*, 2011; García-León *et al.*, 2018). When resuspended in fresh culture medium amended with carbenicillin

and vancomycin to eliminate *A. tumefaciens*, PCPs carrying a hygromycin selection
cassette quickly formed a transgenic suspension culture under hygromycin selection,
as demonstrated by expression of the RUBY reporter (Fig. S2C-E). For the cell lines
Col-0 and MM1, transformation was possible with small PCPs of approximately 200
mg and 60 mg, respectively, as well as with large PCPs of approximately 8 g. The
transformation of large PCPs followed by hygromycin selection and scaling up the
culture allowed the production of nearly 100 g (fresh weight) of transformed Col-0
cells within 13 days after PCP infiltration (Fig. S2D). Similarly to what was observed
for transient expression, the formation of transgenic lines from PCPs prepared from
the cell line MM1 was only consistently possible if the cells were grown under
darkness before infiltration (Fig. S2E). Furthermore, 2 days of co-culture with *A.*
tumefaciens yielded more consistent transformations than 3 days and an optical
density of the applied *A. tumefaciens* of 0.1 favoured transformation compared to a
higher density of 0.4 (Fig. S2E).

363 Characterization of a promoter set for heterologous expression in plant cells.

To expand the molecular toolkit for synthetic biology applications including
heterologous expression from multigene constructs, we characterized 12 promoters
for constitutive gene expression in Arabidopsis suspension cells. A luciferase-based
reporter assay was used to measure promoter activity in infiltrated PCPs. The
cassava vein mosaic virus (CsVMV) promoter led to the highest luciferase expression
and the activity of the other promoters is therefore shown as percentage of the
CsVMV activity (Fig. 3). The widely used cauliflower mosaic virus *35S* (CMV35S)
promoter exhibited similar activity (96±16%), followed by the Arabidopsis ubitquitin-
10 (*AtUbq10)* promoter (73±3%) and the *A. tumefaciens* mannopine synthase
(*AtuMas*) promoter (30±12%). The tomato histone H4 (*SlH4*) promoter and the *A.*
tumefaciens nopaline synthase (*AtuNos*) promoters were considerably weaker (3±1%
and 3±3%, respectively; Fig. 3). The activities of the 6 weakest promoters were
between 0.6±0.2% and 0.3±0.2%. These findings suggest that the CsVMV, the
CVM35S, the *AtUbq10*, and the *AtuMas* promoters are most suitable for
heterologous expression of proteins to be purified and biochemically characterized in
Arabidopsis suspension cells. The weaker promoters can be used to drive the
expression of accessory proteins that are required at lower quantities.

381 Solubility screening of plastid targeted proteins.

We next sought to apply the newly established method for transient and stable
transformation of Arabidopsis suspension cells and the promoters validated for this
system to the engineering of BNF through heterologous expression of prokaryotic
nitrogen fixation (Nif) proteins. We aimed to identify new NifB protein variants that
can be used to engineer BNF in plants by performing a solubility screen of naturally
occurring NifB variants. Since NifB function and potentially also solubility depend on
the accessory proteins NifU, NifS and FdxN, we first generated a stable cell line
expressing those proteins to be used as a background line for the NifB solubility
screening. We therefore first tested a subset of the previously characterized

promoters and previously described chloroplast targeting peptides (CTPs) for
expression and plastid localization of NifU, NifS, and FdxN. NifX was co-expressed
from the same construct to test its solubility due to its function in nitrogenase co-
factor biosynthesis downstream of NifB. We found that the Arabidopsis *ubiquitin-10*
promoter, the tomato histone *H4* promoter and the Cassava Vein Mosaic Virus
(CsVMV) promoter led to the accumulation of NifU, NifS, and FdxN, respectively, in
the soluble fraction of protein extracts (Fig. 4A), whereas NifX was not soluble (Fig.
S3). Since the latter protein is not needed for NifB activity, *nifX* was excluded from
later constructs. The Arabidopsis CTPs from the TOCC, CAB6 and GLTB2 proteins
allowed plastid localization, as assessed by the presence of a single band in
immunoblots, suggesting the absence of full-length proteins with uncleaved CTPs
(Fig. 4A), and by the co-localization of GFP-fusion proteins with an established
plastid marker protein, as determined by confocal microscopy (Fig. 4B and Fig. S4).
In a cell line that stably expressed NifU, NifS, and FdxN, a library of 15 different
BCCP1::TS::nifB homologs was tested in transient expression assays (Fig. 5A),
employing a TS-tag sequence that contains an intron to avoid false positive detection
resulting from misexpression in *A. tumefaciens*. Since archaeal Nif variants have
been successfully expressed in eukaryotes before, we tested 13 *nifB* homologs from
archaea and only two from bacteria. Two archaeal homologs originated from
uncultivated samples of extreme environments, the others are from thermophilic
and/or known diazotrophic archaea (Fig. 5B, Fig. S1, Table S4). Comparing total and
soluble extracts of *nifB*-infiltrated PCPs led to the accumulation of soluble NifB
protein for 11 out of 15 tested NifB homologs (Fig. 5C, Fig. S5). None of the two
tested bacterial NifB homologs were soluble in plant cells and only two of the tested
archaeal homologs were insoluble.

Gene stacking and protein purification.

To test the capacity of this method for stable expression of complex multi-gene
constructs, we generated Arabidopsis suspension cell lines carrying a single T-DNA
with six genes: a hygromycin resistance marker (*hygR*) for selection, *gfp* as a
transformation control, and the four gene components of the NifB module (*nifU*, *nifS*,
fdxN, and *nifB*). The construct design is shown in Fig. 6A. All four proteins were
detected in the soluble fraction of protein extracts from stably transformed cells by
immunoblotting (Fig. 6B), confirming successful expression. To further validate
expression and protein integrity, we performed affinity purification using a TS-tag
fused to NifB. Coomassie-stained gels of purification fractions revealed bands at the
expected size alongside some nonspecific background (Fig. 6C), and a Western blot
confirmed that NifB was specifically enriched during the purification (Fig. 6D).
Additional immunoblots for GFP and the plastid-localized RbcL protein confirmed the
release of both cytosolic and plastid contents during tissue homogenization. Mass
spectrometry confirmed the identity of the purified protein of the expected mass (39.5
kDa) as TS::NifB (Fig. S6). Together, these results demonstrate that our platform
supports large multi-gene constructs and simultaneous expression of several plastid-
targeted proteins in plant cells.

Discussion

In this study, we established and validated a rapid and efficient workflow for transient and stable expression of multigene constructs in *A. thaliana* suspension cultures, with the aim of enabling complex engineering tasks such as the reconstruction of the nitrogen fixation pathway in plant cells. Our approach is based on the plant cell pack infiltration method (Rademacher *et al.*, 2019) and adapts it for Arabidopsis, demonstrating robust transformation efficiency, compatibility with several cell lines, and rapid generation of stably transformed cultures. By applying this system to the expression of plastid-targeted Nif proteins, we showcase its utility for synthetic biology applications. A key innovation of this work is the combination of transient and stable transformation workflows into a single streamlined platform. Unlike previous methods, which require more than 40 days to yield 200 ml of transgenic cell cultures (García-León *et al.*, 2018) or 29 days to yield 42 ml of culture (Van Leene *et al.*, 2011), the workflow presented here can generate 300 ml transgenic culture containing 100 g of cells in less than 2 weeks. This acceleration is probably due to a combination of a higher amount of starting material and a higher transformation efficiency. The workflow relies on basic low-cost equipment found in most biology laboratories. It significantly increases experimental throughput, which is especially important for iterative testing of multi-component systems like nitrogenase biosynthesis. Furthermore, transformed cultures could be propagated without the need for selection, opening the way for simple marker-less transformation applications.

While these features represent major advantages for plant synthetic biology, there are also limitations to consider. The current workflow yields heterogeneous cell populations rather than clonal lines, leading to varying T-DNA integration sites and expression profiles. While this population-based approach suffices for solubility screening and expression optimization, applications requiring precise genotype-phenotype correlations may benefit from purified transgenic lines. However, isolating and propagating individual Arabidopsis suspension cells is technically challenging due to their requirement for high cell densities (Muir *et al.*, 1954). This constraint can be addressed through the use "feeder cells" that maintain the growth of isolated transformant cells or through microdroplet-based approaches, in which single cells are cultured in small volumes (Schweiger *et al.*, 1987; Spangenberg and Koop, 1992). The ability to isolate purified cell lines will also be critical for forward-genetic screens, which are another possible future application of the presented platform thanks to the high transformation efficiency combined with the possibility to generate stable transformants. For example, bulk-transformations of fully assembled *nif* libraries could be employed to select for nitrogenase activity using fluorescent H_2 or ethylene biosensors in combination with fluorescence activated cell sorting.

The NifB solubility screen yielded results consistent with previous observations in
Nicotiana benthamiana leaves, supporting the use of Arabidopsis suspension
cultures as a relevant model system for plastid engineering. Specifically, the only
three NifB homologs that had been shown to be soluble in *N. benthamiana* leaves
when targeted to chloroplasts and/or mitochondria were also included in the library of
the present study and all of them were soluble in our Arabidopsis system. Notably,
none of the 26 previously tested bacterial NifB homologs were soluble in *N.*
benthamiana (Jiang *et al.*, 2022). Similarly, in the present study, none of the two
tested bacterial NifB homologs were soluble in Arabidopsis cells as opposed to 11
out of 13 tested archaeal homologs that exhibited some level of solubility. This
provides further evidence that archaeal Nif variants might generally be superior for
engineering BNF in eukaryotes.

This work lays the foundation for implementing multi-component pathways in plastids
and for exploring the effects of subcellular localization and nutrient availability on
nitrogenase stability and activity. Some Arabidopsis suspension cultures can be
readily shifted between heterotrophic and photomixotrophic conditions, enabling
controlled studies on the impact of photosynthetically produced oxygen on Nif protein
accumulation and function. However, not all cell lines are capable of consistently
developing chloroplasts even under light conditions, as shown for Col-0 (Fig. 2C);
therefore, the cell line MM1 would be more suitable to address research questions
related to chloroplast dynamics. This flexibility, in combination with high
transformation speed, makes the system highly suitable for prototyping nitrogenase
expression strategies and other complex biosynthetic pathways.

498 **Supplementary data**

Figure S1: *nifB* library design and homolog selection criteria.

Figure S2: Assessment of *A. tumefaciens* strain EHA105 and cell line YG1 for PCP
infiltration as well as examples of stable transformation of different cell lines.

Figure S3: NifX solubility testing.

Figure S4: Localization controls for confocal microscopy.

Figure S5: Additional replicate of NifB solubility screening.

Figure S6: NifB protein confirmation by mass spectrometry.

Table S1: Primers used in this study.

Table S2: Plasmids used in this study.

Table S3: Synthetic sequences ordered for this study.

Table S4: NifB library and corresponding source organisms.

Acknowledgements

513   We thank Álvaro Eseverri for help with MoClo assemblies, David Alabadí for
514   providing the cell line PSB-L, and Laszlo Bako for providing the cell lines Col-0 and
515   MM1.

Author contributions

518   L.M., G.A.T., Z.F.E., X.J., and S.B. performed the experimental work, L.M., G.A.T.,
519   and L.M.R. designed experiments, analyzed data, and wrote the manuscript.

Conflict of interest

521   The authors declare no conflicts of interest.

Data availability

523   All data supporting the findings of this study are available within the paper and its
524   Supplementary Information. A preprint version of the manuscript is available at
525   bioRxiv under CC BY 4.0 license. The Supplementary Information is available at
526   Dryad.

Funding

This work was supported, in whole or in part, by the Gates Foundation grants INV-
005889 and INV-067006. Under the grant conditions of the Foundation, a Creative
Commons Attribution 4.0 License has already been assigned to the Author Accepted
Manuscript version that might arise from this submission. L.M. was supported by the
Swiss National Science Foundation grant P2EZP3_199998. G.A.T. was supported by
Programa Propio UPM de I+D+i 2022.

References

Arragain S, Jiménez-Vicente E, Scandurra AA, Burén S, Rubio LM, Echavarri-Erasun C.
2017. Diversity and functional analysis of the FeMo-cofactor maturase NifB. Frontiers in
Plant Science **8**.

Aznar-Moreno JA, Jiang X, Burén S, Rubio LM. 2021. Analysis of nitrogenase Fe protein
activity in transplastomic tobacco. Frontiers in Agronomy **3**.

**Bateman A, Martin MJ, Orchard S*, et al.* 2021. UniProt: the universal protein
knowledgebase in 2021. Nucleic Acids Research **49**, D480–D489.

Boehm CR, Bock R. 2019. Recent advances and current challenges in synthetic biology of
the plastid genetic system and metabolism. Plant Physiology **179**, 794–802.

Burén S, Jiang X, López-Torrejón G, Echavarri-Erasun C, Rubio LM. 2017. Purification
and *in vitro* activity of mitochondria Targeted nitrogenase cofactor maturase NifB. Frontiers in
Plant Science **8**.

Burén S, Jiménez-Vicente E, Echavarri-Erasun C, Rubio LM. 2020. Biosynthesis of nitrogenase cofactors. Chemical Reviews **120**, 4921–4968.

Burén S, Ortega-Villasante C, Blanco-Rivero A, Martínez-Bernardini A, Shutova T, Shevela D, Messinger J, Bako L, Villarejo A, Samuelsson G. 2011. Importance of post-translational modifications for functionality of a chloroplast-localized carbonic anhydrase (CAH1) in *Arabidopsis thaliana*. PLoS ONE **6**, e21021.

Burén S, Pratt K, Jiang X*, et al.* 2019. Biosynthesis of the nitrogenase active-site cofactor precursor NifB-co in *Saccharomyces cerevisiae*. Proceedings of the National Academy of Sciences of the United States of America **116, 25078–25086.

Burén S, Rubio LM. 2018. State of the art in eukaryotic nitrogenase engineering. FEMS Microbiology Letters **365**, 274.

Buschmann H. 2016. Plant cell division analyzed by transient *Agrobacterium*-mediated transformation of tobacco BY-2 cells. Methods in molecular biology **1370**, 17–25.

Chen IMA, Chu K, Palaniappan K*, et al.* 2021. The IMG/M data management and analysis system v.6.0: new tools and advanced capabilities. Nucleic Acids Research **49, D751–D763.

Christie M, Croft LJ, Carroll BJ. 2011. Intron splicing suppresses RNA silencing in Arabidopsis. The Plant Journal **68**, 159–167.

Cortese E, Carraretto L, Baldan B, Navazio L. 2021. Arabidopsis photosynthetic and heterotrophic cell suspension cultures. *Arabidopsis Protocols*, Vol. 2200: Springer US, 167–185.

Curatti L, Rubio LM. 2014. Challenges to develop nitrogen-fixing cereals by direct *nif*-gene transfer. Plant Science **225**, 130–137.

Dixon R, Cheng Q, Shen G-F, Day A, Dowson-Day M. 1997. *Nif* gene transfer and expression in chloroplasts: Prospects and problems. Plant and Soil **194**, 193–203.

Eseverri Á, Baysal C, Medina V, Capell T, Christou P, Rubio LM, Caro E. 2020a. Transit peptides from photosynthesis-related proteins mediate import of a marker protein into different plastid types and within different species. Frontiers in Plant Science **11**, 1.

Eseverri Á, López-Torrejón G, Jiang X, Burén S, Rubio LM, Caro E. 2020b. Use of synthetic biology tools to optimize the production of active nitrogenase Fe protein in chloroplasts of tobacco leaf cells. Plant Biotechnology Journal **18**, 1882–1896.

Fischer R, Emans N, Schuster F, Hellwig S, Drossard J. 1999. Towards molecular farming in the future: using plant-cell-suspension cultures as bioreactors. Biotechnology and Applied Biochemistry **30**, 109–112.

García-León M, Iniesto E, Rubio V. 2018. Tandem affinity purification of protein complexes from Arabidopsis cell cultures. *Two-Hybrid Systems*, Vol. 1794: Springer New York, 297–309.

Gengenbach BB, Opdensteinen P, Buyel JF. 2020. Robot cookies – plant cell packs as an automated high-throughput screening platform based on transient expression. Frontiers in Bioengineering and Biotechnology **8**.

Hampp C, Richter A, Osorio S, Zellnig G, Sinha AK, Jammer A, Fernie AR, Grimm B, Roitsch T. 2012. Establishment of a photoautotrophic cell suspension culture of *Arabidopsis thaliana* for photosynthetic, metabolic, and signaling studies. Molecular Plant **5**, 524–527.

He Y, Zhang T, Sun H, Zhan H, Zhao Y. 2020. A reporter for noninvasively monitoring gene expression and plant transformation. Horticulture Research **7**, 1–6.

Ivleva NB, Groat J, Staub JM, Stephens M. 2016. Expression of active subunit of nitrogenase via integration into plant organelle genome. PLoS ONE **11**, e0160951.

Jiang X, Coroian D, Barahona E, Echavarri-Erasun C, Castellanos-Rueda R, Eseverri Á, Aznar-Moreno JA, Burén S, Rubio LM. 2022. Functional nitrogenase cofactor maturase NifB in mitochondria and chloroplasts of *Nicotiana benthamiana*. mBio **13**.

Jiang X, Payá-Tormo L, Coroian D, García-Rubio I, Castellanos-Rueda R, Eseverri Á, López-Torrejón G, Burén S, Rubio LM. 2021. Exploiting genetic diversity and gene synthesis to identify superior nitrogenase NifH protein variants to engineer N_2-fixation in plants. Communications Biology **4**, 1–11.

Li C, Liu Y, Liu X*, et al.* 2021. Chloroplast thylakoid ascorbate peroxidase PtotAPX plays a key role in chloroplast development by decreasing hydrogen peroxide in *Populus tomentosa*. Journal of Experimental Botany **72, 4333–4354.

Li Z, Zhang D, Xiong X, Yan B, Xie W, Sheen J, Li JF. 2017. A potent Cas9-derived gene activator for plant and mammalian cells. Nature Plants 2017 3:12 **3**, 930–936.

McCarthy A, Chung M, Ivanov AG, Krol M, Inman M, Maxwell DP, Hüner NPA. 2016. An established *Arabidopsis thaliana* var. Landsberg *erecta* cell suspension culture accumulates chlorophyll and exhibits a stay-green phenotype in response to high external sucrose concentrations. Journal of Plant Physiology **199**, 40–51.

Menges M, Murray JAH. 2004. Cryopreservation of transformed and wild-type *Arabidopsis* and tobacco cell suspension cultures. The Plant Journal **37**, 635–644.

Muir WH, Hildebrandt AC, Riker AJ. 1954. Plant tissue cultures produced from single isolated cells. Science **119**, 877–878.

Mukherjee S, Stamatis D, Cindy*, et al.* 2025. Genomes OnLine Database (GOLD) v.10: new features and updates. Nucleic Acids Research **53, D989–D997.

Oldroyd GE, Dixon R. 2014. Biotechnological solutions to the nitrogen problem. Current Opinion in Biotechnology **26**, 19–24.

Provart NJ, Alonso J, Assmann SM*, et al.* 2016. 50 years of Arabidopsis research: highlights and future directions. New Phytologist **209, 921–944.

Rademacher T, Sack M, Blessing D, Fischer R, Holland T, Buyel J. 2019. Plant cell packs: a scalable platform for recombinant protein production and metabolic engineering. Plant Biotechnology Journal **17**, 1560.

Rizzo P, Chavez BG, Leite Dias S, D'Auria JC. 2023. Plant synthetic biology: from inspiration to augmentation. Current Opinion in Biotechnology **79**, 102857.

Santos RB, Abranches R, Fischer R, Sack M, Holland T. 2016. Putting the spotlight back on plant suspension cultures. Frontiers in Plant Science **7**.

Schindelin J, Arganda-Carreras I, Frise E*, et al.* 2012. Fiji: an open-source platform for biological-image analysis. Nature Methods **9, 676–682.

Schober I, Koblitz J, Joaquim*, et al.* 2025. Bac*Dive* in 2025: the core database for prokaryotic strain data. Nucleic Acids Research **53, D748–D756.

Schweiger HG, Dirk J, Koop HU, Kranz E, Neuhaus G, Spangenberg G, Wolff D. 1987. Individual selection, culture and manipulation of higher plant cells. Theoretical and Applied Genetics **73**, 769–783.

Scotti N, Rigano MM, Cardi T. 2012. Production of foreign proteins using plastid transformation. Biotechnology Advances **30**, 387–397.

Sello S, Moscatiello R, La Rocca N, Baldan B, Navazio L. 2017. A rapid and efficient method to obtain photosynthetic cell suspension cultures of *Arabidopsis thaliana*. Frontiers in Plant Science **8**, 1–8.

Sello S, Perotto J, Carraretto L, Szabò I, Vothknecht UC, Navazio L. 2016. Dissecting stimulus-specific Ca^{2+} signals in amyloplasts and chloroplasts of *Arabidopsis thaliana* cell suspension cultures. Journal of Experimental Botany **67**, 3965–3974.

Spangenberg G, Koop HU. 1992. Low density cultures: microdroplets and single cell nurse cultures. *Plant Tissue Culture Manual*: Kluwer Academic Publishers, 51–78.

Van Leene J, Eeckhout D, Persiau G, Van De Slijke E, Geerinck J, Van Isterdael G, Witters E, De Jaeger G. 2011. Isolation of transcription factor complexes from *Arabidopsis* cell suspension cultures by tandem affinity purification. *Plant transcription factors*: Humana Press, 195–218.

Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S. 2011. A modular cloning system for standardized assembly of multigene constructs. PLoS ONE **6**, 16765.

Werner S, Engler C, Weber E, Gruetzner R, Marillonnet S. 2012. Fast track assembly of multigene constructs using golden gate cloning and the MoClo system. Bioengineered Bugs **3**, 38–43.

Yang J, Xie X, Yang M, Dixon R, Wang Y-P. 2017. Modular electron-transport chains from eukaryotic organelles function to support nitrogenase activity. Proceedings of the National Academy of Sciences **114**, E2460–E2465.

Zulkower V, Rosser S. 2020. DNA Chisel, a versatile sequence optimizer. Bioinformatics **36**, 4508–4509.

Figure legends

Fig 1: Workflow and applications of plant cell pack (PCP) infiltration in Arabidopsis suspension cells for stable and transient transformation at different scales. (A) Plant cells are maintained by weekly or monthly passaging in liquid or solid MS medium, respectively. Washed plant cells grown in liquid medium under darkness serve as starting material for both transient and stable transformations. 200-mg PCPs allow high throughput, whereas 8-g PCPs allow the production of up to 100 g (fresh weight) of transgenic cells in 2 weeks. (B) For high-throughput applications, a dense suspension of washed cells is loaded on top of a pipette filter tip. The excess liquid is then removed by vacuum and the semi-dry PCP is infiltrated with an *A. tumefaciens* suspension. After 60 min, the excess liquid is removed and the PCPs show transgene expression after 2-3 days. Individual PCPs can be conveniently transferred to microcentrifuge tubes downstream applications such as protein extraction, microscopy, or the generation of stably transformed cells. (C) For high-volume applications such as the purification of heterologously produced proteins, washed suspension cells are loaded onto midi-prep columns and excess liquid is removed by gravity or gentle vacuum. *A. tumefaciens* is loaded by gravity and, after 60 min, excess liquid and bacteria are removed by gentle centrifugation and PCPs are transferred to 50-ml centrifugation tubes. After 2-3 days, transformed PCPs are resuspended in fresh medium to start a liquid culture. Created in https://BioRender.com.

Fig 2: Optimization of transient expression in Arabidopsis suspension cells by plant cell pack (PCP) infiltration. (A) PCPs of four different Arabidopsis cell lines infiltrated with three different *A. tumefaciens* strains harboring a plasmid with the RUBY reporter. Pictures were taken 2 and 3 days post infiltration (dpi). PCPs treated with the infiltration buffer only were included as negative control. (B) PCPs of the Arabidopsis cell line Col-0 infiltrated with the *A. tumefaciens* strain GV3101::pMP90 carrying the RUBY cassette prepared from liquid or solid culture medium. (C) PCPs from the cell line Col-0 grown either under constant darkness or a 16-h photoperiod and infiltrated with *RUBY*-carrying *A. tumefaciens* strain GV3101::pMP90 (+) or with infiltration buffer only (-). (D) PCP infiltration of cell line MM1 grown either under darkness or a 16-h photoperiod with RUBY-carrying *A. tumefaciens* strain GV3101::pMP90 prepared at different optical densities (+) or with infiltration buffer only (-). (E) Co-expression of the suppressor of silencing P19 driven by the *Nos* promoter and GFP driven by the *35S* promoter in PCPs of cell line Col-0. The *p19* and *gfp* genes were either present on the same or on different plasmids. Three technical replicates are shown.

Fig 3: Various constitutive promoters of different strength allow transgene expression in Arabidopsis suspension cells.

Relative promoter activity as determined by LUC reporter assays. Different constitutive promoters driving expression the firefly luciferase (*fluc*) gene were co-transformed with a GUS reporter cassette (*p35S::GUS::t35S*) for normalization and the suppressor of silencing *p19* from a single T-DNA (grey box; not to scale). Promoter activity was defined as luminescence divided by GUS activity. Bars show the average of 3 biological replicates. The y-axis is segmented with two different scales. Individual replicates are shown as black dots and error bars represent the standard deviation. Letters represent statistical groups according to one-way ANOVA and Tukey's test. LB: left T-DNA border; RB: right T-DNA border.

Fig 4: Soluble proteins of the NifB functional module can be expressed and targeted to plastids in Arabidopsis suspension cells. (A) Western blots of total and soluble plant cell extracts transformed with *nifU*, *nifS*, and *fdxN* driven by the Arabidopsis *ubiquitin-10* promoter, the tomato histone H4 promoter, and the Cassava Vein Mosaic Virus (CsVMV) promoter, respectively. The proteins were fused to minimal versions of the chloroplast-targeting peptides (CTPs) of the Arabidopsis proteins TOCC, CAB6, and GLTB2, respectively (Eseverri *et al.*, 2020a). FdxN was fused to an HA-tag for detection, whereas NifU and NifS were untagged. Col-0 cells were transformed with plasmid pN2LM165 (rep1) and plasmid pN2LM166 (rep2) and proteins extracted with a buffer containing 100 mM Tris-HCl (pH8), 150mM NaCl, 10% glycerol and 0.5% protease inhibitor cocktail. (B) Detection of GFP signal in Arabidopsis cells transformed with *nifU*, *nifS* and *fdxN* fused to *gfp* and the TOCC, CAB6 and GLTB2 CTPs, respectively, by confocal laser scanning microscopy. The mCherry reporter fused to the RecA CTP was used as a control for plastid localization. Merged pictures of the mCherry and GFP channels are also shown. Scale bars: 10 μm.

Fig 5: Solubility screening identified new archaeal NifB protein variants suitable for plant expression. (A) Construct and expression design for the NifB solubility screening. A stably transformed background line expressing *nifU*, *nifS*, and *fdxN::HA* was generated first. In this background line, the different *nifB* homologs fused to the BCCP1 chloroplast targeting peptide (Eseverri *et al.*, 2020b) were transiently co-expressed with *gfp* as a transformation control and the suppressor of silencing p19. (B) Workflow overview for the *nifB* library and synthetic gene design. NifB variants annotated by Uniprot were filtered for known NifB protein motifs (Arragain *et al.*, 2017) and for archaeal origin. Public databases such as Bac*Dive* (Schober *et al.*, 2025) and information on isolation sources were used to select 8 NifB homologs from thermophilic organisms. Additionally, four homologs from known diazotrophic archaea, one from an uncultured archaeon, and two from extremophilic bacteria were included. The synthetic gene design was performed using the DNA Chisel program (Zulkower and Rosser, 2020), focusing on matching the codon usage to Arabidopsis and obtaining high GC and GC3 content as well as avoiding known

mRNA instability motifs. (C) Western blots of total (T) and soluble (S) plant cell extracts. NifB variants were fused to the Twin-Strep (TS) tag for detection with a Strep monoclonal antibody (StrepMAB). A control with the same expression vector but without *nifB* was included (-NifB). LB: left T-DNA border; RB: right T-DNA border; CTP: chloroplast targeting peptide; *hygR*: hygromycin selection marker.

Fig 6: Multi-gene stacking of the NifB module in stably transformed plant cells. (A) Construct design for stable transformation for expression plastid-targeted NifU, NifS, FdxN::HA, Twin-Strep::NifB, and cytosolic GFP. CTP: chloroplast targeting peptide (B) Western blots and ponceau-stained membranes of total and soluble plant extracts from MM1 plant cells transformed with the construct depicted in (A) with *nifB* from *Methanobacterium bryantii*, using α-NifU, α-NifS, α-HA, α-StrepII and α-GFP antibodies. (C) Left panel: Coomassie-stained SDS-PAGE gel loaded with different fractions of the Strep-Tactin affinity chromatographic purification of Col-0 cells transformed with the construct depicted in (A) with *nifB* from *Methanosarcina acetivorans.* Asterisk indicates the expected mass of TS-NifB, 39.5 kDa. Right panel: concentrated elution. Bands labelled with red numbers were analyzed by mass spectrometry. Band 3 was identified as TS::NifB, whereas bands 1, 2, and 4 corresponded to co-purified Arabidopsis proteins methylcrotonoyl-CoA carboxylase subunits alpha (Q42523) and beta (Q9LDD8), and the biotin carboxyl carrier protein of acetyl-CoA carboxylase (F4KE21), respectively. Band 5 was identified as a mix of the Arabidopsis proteins A0A384KVD0 and Q9SSR9 of unknown functions. (D) Western blots of the purification steps shown in (C) with the α-Strep (StrepMAB), α-RbcL, α-GFP antibodies.

Fig S1: *nifB* library design. Archaeal NifB proteins present in the Uniprot database (Bateman *et al.,* 2021) were identified by first obtaining a list of all entries with the keyword "nitrogenase", then filtering this dataset by presence of the word "cofactor" in the protein name, by archaeal origin, and by the presence of six NifB protein motifs (Arragain *et al.,* 2017). The dataset obtained from Uniprot contained information on the taxonomy of source organisms, which was used to select a diverse selection of NifB sequences, as well as the NCBI Taxon ID, which was used to obtain information on growth temperature, habitats, and metabolism. Main selection criteria were taxonomic diversity of the library, high growth or isolation temperature, extreme habitats, and reported diazotrophy.

Fig S2: Plant cell pack (PCP) infiltration is a simple and fast method for transient and stable transformation of Arabidopsis suspension cells. (A) PCPs cast from 500 µl dense cell suspension and infiltrated with the *A. tumefaciens* strains EHA105 and GV3101::pMP90 harboring a *35S::RUBY* reporter cassette (+) or with infiltration buffer only (-). (B) A 1000-µl filter pipette tip and a midi-prep column with a 15-ml reservoir loaded with PCPs before infiltration. (C) 500-µl PCPs of cell line Col-0 infiltrated with the *35S:RUBY* reporter cassette 2 days after infiltration and after resuspension in growth medium and a culture passage with and without hygromycin selection. (D) Different stages of biomass production of stably transformed suspension cells derived from an 8-g PCP. Shown are a PCP in a 50-ml

centrifugation tube, cells after resuspension, after a 3-day selection with hygromycin, and after a passage from one to four flasks. (E) Recovery of stable transformants of MM1 cells after treatment of 2 or 3 days with an *A. tumefaciens* strain carrying *35S:RUBY* at different optical densities and selected with hygromycin for 1 week. Pictures show cultures after transferring to a spectrophotometry cuvette and sedimentation. Before PCP infiltration, suspension cultures were grown either under a 16-h photoperiod or without light. Shown are 3 biological replicates. OD values refer to the *A. tumefaciens* suspensions used for PCP infiltration. dpi: days post infiltration.

Fig S3: NifX is expressed but insoluble in plant cell packs. Western blots and ponceau-stained membranes of total and soluble protein extracts of plant cell packs transformed with *nifX* driven by the Arabidopsis *act2* or *TCTP* promoters. The protein was fused to the minimal version of the Arabidopsis RCA chloroplast targeting peptide (Eseverri *et al.*, 2020a), to the TS-tag for detection, and to the TEV protease cleavage site.

Fig S4: Subcellular localization of plastid-targeted Nif proteins. Detection of GFP signal in Arabidopsis cells transformed with *nifU*, *nifS* and *fdxN* fused to *gfp* and the TOCC, CAB6 and GLTB2 CTPs, respectively, by confocal laser scanning microscopy (rows 1-3). Arabidopsis suspension cells expressing localization control constructs are also shown. Row 4: mCherry fused to the Arabidopsis RecA chloroplast targeting peptide. Row 5: GFP lacking a chloroplast targeting peptide, showing diffuse cytosolic localization. Row 6: Co-expression of both controls from a single plasmid. For each construct, representative wide-field images and individual GFP and mCherry channels, as well as merged fluorescence channels are shown. Scale bars: 10 µm.

Fig S5: Solubility screening identified new archaeal NifB protein variants suitable for plant expression. Replicate of experiment shown in Fig. 5. Western blots and ponceau-stained membranes of total (T) and soluble (S) plant cell extracts. NifB variants were fused to the Twin-Strep tag for detection with a Strep monoclonal antibody (StrepMAB). A control with the same expression vector but without *nifB* was included (-NifB). Asterisks mark the positions of full-length NifB proteins.

Fig S6: Peptide mass fingerprinting of purified TS::NifB protein. Sequence of BCCP1$_{CTP}$::TS::NifB of *Methanosarcina acetivorans*. Blue residues show the part of the BCCP1 chloroplast targeting peptide predicted to be cleaved during transfer to the plastid (Eseverri *et al.*, 2020b). Red bold residues show the peptides identified by mass spectrometry.

A

monthly passage → weekly passage

harvest, wash & concentrate

wash buffer
plant cells

high throughput, low volume
200 mg cells

high volume, low throughput
8 g cells

transient expression
2-3 days

stable transformation
up to 100 g biomass in 14 days

B High throughput, low volume:

plant cells
filter

filter tip

Dry by vacuum

Infiltrate PCPs with *A. tumefaciens*

dry PCPs by vacuum

2-3 days

harvest PCPs

resuspension

protein extraction

protein detection   microscopy

Stable transformation:

propagation → scale-up

C High volume, low throughput:

gravity-pack plant cells

15-ml reservoir

modified midiprep column

load *A. tumefaciens* by gravity

dry by centrifugation

transfer to 50-ml tube
3 days

resuspension, selection, propagation

Stable transformation:

protein purification