Title: Impacts of changing ocean circulation, temperature, and food supply on larval

recruitment of purple sea urchins in Southern California: A biophysical modeling study

Authors: Rachel D. Simons[1*] and Daniel K. Okamoto[2]

[1*] Earth Research Institute, University of California, Santa Barbara, CA, USA

[2] Department of Integrative Biology, University of California, Berkeley, CA, USA

[*] Corresponding Author. Address: Email address: simons@eri.ucsb.edu

Abstract:

Changes in ocean conditions among years and across decades can alter populations in marine

ecosystems. This study evaluates the impact of changing ocean conditions on larval recruitment

of purple sea urchins in the Southern California Bight using 3D biophysical modeling, surface

chlorophyll, and recruitment data. The influence of circulation, temperature, and food supply on

larval recruitment is quantified using five modeled variables, larval dispersal distance, larval

source location, larval food supply, and temperature exposure for larvae and adults, which are

derived for 18 years and five larval recruitment sites. Sensitivity testing of the variables to

different plankton larval durations (PLDs), larval behaviors, and nearshore retention is

performed. All variables are found to be relatively insensitive to changes in PLD greater than 26

days. Larval dispersal distance and source location, representing changes in circulation, are

found to be more sensitive to larval behavior and nearshore retention than larval food supply

and temperature exposure for larvae and adults. All variables are statistically compared to

recruitment field data. Temperature exposure for adults during the fall reproductive season is

found to be a strong driver of larval recruitment while temperature exposure for larvae during

the spring recruitment season is not. Food supply is not found to be a driver of larval

recruitment. Circulation is found to be a driver of larval recruitment if larvae have behavior that

reduces their dispersal distance, allowing them to come from source sites near to the

recruitment sites. Overall, we hypothesize that larval behavior which reduces dispersal improves

recruitment and that the timing of recruitment and reproduction can predict the impact of

temperature on recruitment.

1.0 Introduction

Extreme changes in ocean conditions, such as temperature, circulation, and primary productivity, pose significant threats to marine ecosystems and are predicted to alter the distribution and productivity of marine species, in part by affecting larval recruitment (Doney et al. 2012; Sydeman et al. 2014). Improving our understanding of larval recruitment can explain and predict shifts in the dynamics of populations of fished and ecologically important species (Botsford et al. 1998; Myers 1998; Pineda et al. 2010; Stige et al. 2013). Thus, quantifying how ocean conditions shape spatial and temporal variation in larval dispersal and recruitment is critical to our understanding how climate drivers influence marine populations. In this study, we explore the impacts of changing ocean conditions on larval recruitment by investigating the relationship between larval recruitment of purple sea urchins and ocean conditions of circulation, temperature, and food supply in the Southern California Bight (SC Bight, Fig. 1) using 3D biophysical modeling.

The SC Bight, the southern component of the California Current System, is a transition zone lying at the interface between two very different oceanographic regimes, the cold nutrient-rich waters of the south-flowing California Current and the warm nutrient-poor equatorial waters of the north-flowing Southern California Countercurrent (Lynn and Simpson 1987; Hickey 1993; Winant et al. 2003; Dong et al. 2009). The meeting of these two regimes in the SC Bight creates a year-round temperature gradient with colder water in the north and warmer water in the south and a seasonal chlorophyll gradient with high levels in the north and low levels in the south during the spring and early summer (Otero and Siegel 2004; Henderikx Freitas et al. 2017; Simons and Catlett 2023). As a result, the SC Bight is particularly sensitive to shifts in the

California Current, which creates dramatic changes in ocean circulation, temperature, and

primary productivity in the SC Bight and are often correlated with El Niño-Southern Oscillation

(ENSO) events (Lynn and Bograd 2002; Dong et al. 2009). These features, in combination with an

abundance of long-term ecological data, make the SC Bight an ideal system for studying the

relationship between changing ocean conditions and larval recruitment.

In the California Current, changes in abundance of the purple sea urchins,

Strongylocentrotus purpuratus, indirectly regulates the species assembly of nearshore rocky

coastlines through voracious herbivory on kelp (Tegner and Dayton 1991; Filbee-Dexter and

Scheibling 2014). A substantial body of research describes how predation and disease impact

adult Strongylocentrotid urchin populations (Mann and Breen 1972; Estes and Duggins 1995;

Lafferty 2004; Filbee-Dexter and Scheibling 2014; Burt et al. 2018) and documents climate-

related booms and busts of purple urchins along the California Coast in recent years (Rogers-

Bennett and Catton 2019; Okamoto et al. 2020; Rogers-Bennett and Okamoto 2020; Shanks et

al. 2020). However, despite the widespread recognition of the importance of larval transport

and recruitment in controlling population fluctuations in many marine species (Morgan 2001;

Underwood and Keough 2001; Pineda et al. 2007; White et al. 2019), no studies have evaluated

how changes in the myriad of ocean conditions influence larval recruitment of purple urchins in

the California Current. The purple sea urchin (*S. purpuratus*) is used for this study because long-

term, spatially replicated recruitment data is available at six locations in the SC Bight and prior

work suggests that high temperatures and ENSO events in the SC Bight are linked to poor

recruitment years (Okamoto et al. 2020) in part through changes in reproduction (Okamoto et

al. Accepted).

The main goal of this study is to assess the impacts of changes in circulation,

temperature and food supply on larval recruitment of purple sea urchins in the SC Bight. A

biophysical model of the SC Bight, an ocean circulation model with particle tracking, is used to

estimate larval recruitment at the recruitment sites where long-term urchin recruitment data

has been collected (Fig. 1, Okamoto et al. 2020). The impact of changing circulation,

temperature, and food supply on larval recruitment is quantified by deriving five indices that

represent ocean conditions for each recruitment site. Two indices represent circulation, larval

dispersal distance (LDD) and source zone strength (SZS). LDD is the distance a particle travels

from a source site to a recruitment site. SZS is the relative probability that larvae would

originate from nine coastal zones in the SC Bight (Fig. 1). Two indices represent temperature,

larval temperature (T_L) and source site temperature (T_S). T_L is the temperature that larvae are

exposed to as they are transported from the source sites to the recruitment sites. T_S is the

temperature that adult urchins are exposed to at the spawning or larval source sites during

reproduction. One index represents food availability (Chl), which is estimated using surface

chlorophyll concentrations from satellite data along the particle tracks. The five ocean condition

indices (OCI) are calculated for an 18-year modeling period from 1996-2013.

To accurately model larval transport and recruitment, the larval life history of the species

must be realistically represented in the biophysical model (Paris et al. 2007; Cowen and

Sponaugle 2009; Metaxas and Saunders 2009). However, empirical data on larval life history

traits, such as vertical migration behavior, of many species is not available, and thus many

studies do not include species-specific larval life history when modeling larval transport and

connectivity (Swearer et al. 2019). To address this problem, we evaluate the sensitivity of the

five OCI to variations in three larval life history parameters, plankton larval duration (PLD), larval

behavior, and nearshore retention. Two types of PLD are simulated, a PLD of 42 days, which is

an intermediate value from literature for purple sea urchins (Cameron and Schroeter 1980;

Miller and Emlet 1999; Tegner 2001), and a temperature dependent PLD based on experimental

studies . Two types of larval behavior are simulated passive drifting and diel vertical migration,

which occurs when the larvae migrate to the surface at night and to depth during the day. For

passive drifting, particles are moved by circulation only. Although evidence of vertical migration

for echinoderms is very limited (Metaxas 2020; Doll et al. 2022), diel vertical migration has been

widely observed for invertebrate larvae (Cohen and Forward 2009; Bandara et al. 2021) and is

thus included in this study. With a 1 km^2 horizontal scale, the ocean circulation model does not

reproduce circulation patterns that are smaller than 1 km, including nearshore circulation which

has been hypothesized to retain larvae (Morgan et al. 2009, 2018; Hagerty et al. 2018; Yamhure

et al. 2021). Thus, a relationship is derived from the number of times a particle passes through a

source site to represent nearshore retention.

Using all possible combinations of the three larval life history parameters, 12 simulations

are conducted over the 18-year modeling period for each of five OCI, LDD, SZS, T_L, T_S and Chl,

determining how sensitive the OCI are to variations in the larval life history parameters. All OCI

are found to be relatively insensitive to static versus dynamic PLDs, but showed varying levels of

sensitivity to both nearshore retention and larval behavior. LDD and SZS are found to be more

sensitive to nearshore retention and larval behavior than Chl, T_S, and T_L. To determine which OCI

are the most likely drivers of larval recruitment in the SC Bight, all simulations of the OCI are

statistically compared to a larval recruitment index derived from larval recruitment data

(Okamoto et al. 2020). T_S is found to be the mostly likely driver to larval recruitment. LDD and

SZS are found to potential drivers of larval recruitment if larvae are conducting DVM and/or

influenced by nearshore retention, while T_L and Chl are unlikely to be drivers of larval

recruitment.

2.0 Methods

2.1 Biophysical Model

To simulate larval transport, a 3D biophysical model consisting of two parts, an ocean

circulation model (OCM) and a particle tracking model (PTM), was used. The OCM was a high-

resolution Regional Ocean Modeling System (ROMS) applied to the SC Bight region (Dong and

McWilliams 2007; Dong et al. 2009). The model domain, shown in Fig. 1, contained the SC Bight

and had 1 km horizontal resolution with 40 vertical levels. The model was thoroughly validated

for mesoscale circulation (Dong et al. 2009) and has successfully reproduced upwelling events in

the SC Bight (Dong et al. 2011; Simons and Catlett 2023) and eddy circulation in the Santa

Barbara Channel (SBC, Simons et al. 2015). The temperature from the OCM was validated using

temperature data from 26 stations in the SC Bight (see Supplementary Information (SI)). Six-

hour averaged 3D flow fields and temperature solutions produced by the OCM from 1996-2013

were used for this study.

Larval transport was simulated using the 3D PTM driven by 3D flow fields produced by

the OCM (Mitarai et al. 2009; Simons et al. 2013). For this project, the particle tracking was run

in reverse time, which was possible since the ROMS flow fields and temperature solutions were

stored offline. Reverse particle tracking was used to simulate the arrival of larvae at five

recruitment sites in the SC Bight, Gaviota, Ellwood, Stearns, Anacapa, and Scripps, where larval

recruitment has been recorded since 1993 (Fig. 1, Schroeter et al. 2017; Okamoto et al. 2020).

The PTM has been used extensively for larval transport in the SC Bight (Mitarai et al. 2009;

Simons et al. 2016; Nishimoto et al. 2019; Page 2019), and validated against observational data

from drifter experiments (Ohlmann and Mitarai 2010).

Particles were released from 45 horizontal locations evenly spaced every square

kilometer around each recruitment site. This distribution of horizontal release points was

selected to capture the mesoscale circulation in the area surrounding the recruitment sites. At

each of the 45 horizontal release points, particles were released vertically every meter from 1 to

30 meters below the surface, the maximum depth where purple sea urchins are typically found

in high densities. By evenly distributing particles over the water column, the particle distribution

reflected average vertical flow conditions as recommended for robustness in biophysical models

(Simons et al. 2013). Particles were released every 6 hours from the 1,350 vertical and

horizontal locations at each recruitment site for 1996-2013 and tracked for 8 weeks. The

tracking time of 8 weeks was selected because it is the maximum possible plankton larval

duration under idealized conditions (Munstermann et al. In Review). The location and

temperature of every particle trajectory was recorded every 6 hours. The particle release

frequency was selected to meet the particle number criteria for model robustness (Simons et al.

2013).

The particles arriving at the recruitment sites must originate from a source site in the

study area (Fig. 1). Potential larval source sites in the SC Bight were identified by location of

suitable spawning habitat for sea urchins. Suitable spawning habitat for sea urchins consists of

rocky reef substrate located within areas that can hosts kelp or are algal dominated. Thus, the

spatial distribution of larval source sites in the SC Bight was derived from satellite observations

of giant kelp canopy (Bell et al. 2015). Using the mean distribution of giant kelp canopy from

1984-2022, suitable spawning habitat was found in 926 of the 1-km^2 model grid cells, which

were designated as potential larval source sites (Fig. 1). To represent the peak recruitment

season for purple sea urchins, which runs from March to June in the SC Bight (Okamoto et al.

2020), particles were released every 6 hours from March 1st to June 30th for every year of the

modeling period (1996-2013) at the release locations for each recruitment site. Based on the

particle release locations and frequency, a total of 12 million particles were released from each

recruitment site over the 18-year modeling period.

To quantify the horizontal dispersal of particles, the 3D particle tracks were transformed

into 2D particle density maps (PDMs). Using the three-dimensional particle tracks for each

recruitment month and site, 2D monthly PDMs were produced by compiling the horizontal

locations of all particle tracks from the particles released over the month into a single dataset

and then summing the number of particle locations within each model grid cell of 1 km^2,

creating a 2D particle density matrix the same size as the model grid. Example PDMs for

Gaviota and Scripps sites are shown in Fig. 2(a-f).

2.2 Larval life history parameters

To accurately simulate the transport of purple sea urchin larvae in the SC Bight, larval life

history parameters, which include larval behavior, planktonic larval duration (PLD), and

nearshore retention, must be incorporated into the biophysical model. This section describes

how each of these parameters was modeled.

Three types of larval behavior were simulated with the PTM, passive drifting, diel

vertical migration to 20 m depth (DVM20), and diel vertical migration to 40 m depth (DVM40).

To simulate passive drifting, particles were moved only by 3D currents. Diel vertical migration

(DVM) occurs when larvae migrate to the surface at night and to depth during the day to feed

and/or avoid predation and is hypothesized to be triggered by light availability (Cohen and

Forward 2009; Bandara et al. 2021). As the vertical movement or distribution of sea urchin

larvae has not been studied in the SC Bight, depths of 20 m and 40 m were selected to test DVM

behavior. As the average depth of the euphotic zone in the SC Bight is ~30 m (Brzezinski and

Washburn 2011; Krause et al. 2013), depths of 20 m and 40 m were selected such that particles

would migration within the euphotic zone and below euphotic zone respectively. DVM20 and

DVM40 were modeled by particles remaining at near-surface depths (~3 m) at night (18:00 hr to

06:00 hr) and at deeper depths of 20 m or 40 m during the day (06:00 hr to 18:00 hr). A basic

random walk was used to vertically distribute the particles in a gaussian pattern around the set

depths, near-surface, 20 m, and 40 m.

PLD is represented in the PTM by the length of time particles are tracked. Two types of

PLD were used for this study. The first PLD was 6 weeks, which was an intermediate value based

on both 4literature and experimental data (Cameron and Schroeter 1980; Miller and Emlet

1999; Tegner 2001) and was termed "static PLD". The second PLD was a temperature dependent

PLD based on experimental studies , which was termed "dynamic PLD". To model the static PLD,

the particle tracking time was set to six weeks or 42 days. To calculate the dynamic PLD, the

following two-step process was used. First, using all particle tracks for one recruitment site and

month of the modeling period, the mean particle temperature was calculated for daily tracking

times of 1 to 56 days. Second, the dynamic PLD was determined by the intersection of two lines

of temperature versus PLD, one line from model results and the other line from the

experimental data (example shown Fig. S1). Experimental data indicates PLD as a function of

temperature does not vary with food availability (Munstermann et al. In Review).This process

was repeated for the three larval behaviors, five recruitment sites, and four months of the

recruitment period for the 1996-2013 modeling period. The dynamic PLDs ranged from 26 to 56

days with a mean of 36 days and are displayed Fig. S2.

To quantify the effect of nearshore circulation on larval retention, two contrasting

simulations, one without nearshore retention and one with nearshore retention, were

produced. The simulation without nearshore retention assumed that particles travelled

continuously from a source site to a recruitment site during their PLD and were not retained in a

source site even if they passed through a source site during their PLD. As no information exists

on the probability of larvae being retained along the shoreline, the simulations with nearshore

retention assumed that the number of particles released during a recruitment month that

passed through a source site at any time during their PLD was directly proportional to the

probability of particles being retained within the source site. Thus, the simulations with

nearshore retention contained all particle tracks that passed through a source site at any time

during their PLD. If a particle track did not originate from a source site at the beginning of their

PLD, the particle track was truncated at the first source site it entered, and the particle was then

assumed to be retained within this source site for the portion of the PLD that it was not being

transported to the recruitment site. Consequently, the PDMs without nearshore retention were

created using only the particle tracks that start at a source site at the beginning of the their PLD,

while the PDMs with nearshore retention were created using particle tracks that start a source

site anytime during their PLD. This means that PDMs with nearshore retention include many

particle tracks that are shorter than the PLD and assume that these particles are retained within

the source site at the beginning of their PLD until they are transported to the recruitment site.

2.3 Ocean condition indices

This section describes how the five ocean condition indices (OCI), larval dispersal

distance, source zone strength, larval food supply, larval temperature, and temperature at the

source sites, were derived from particle tracking results. All OCI are calculated on a monthly

timescale for the modeling period of 1996-2013. Larval dispersal distance, source zone strength,

larval food supply, and larval temperature are calculated for the recruitment period of March to

June. Temperature at the source sites is calculated for the reproductive period of September to

November. For the larval life history parameters described in the previous section, only

nearshore retention changed the particle tracking times to less than the PLD. This change is

particle tracking times affected how source zone strength and larval temperature were

calculated with and without nearshore retention and is explained in this section.

Larval dispersal distance (LDD)

To compare larval dispersal between simulations, a monthly LDD for each recruitment

month and site was produced. The LDD was derived by calculating the distance between the

source site and recruitment site for each particle and then averaging this distance for all

particles tracked over the month. Thus, a monthly time-series of LDD for the recruitment period

was produced for each recruitment site in units of kilometers. To derive a mean LDD for the

entire 18-year study period, the LDD was averaged over all months and years of the study

period and is termed "mean LDD".

Source zone strength (SZS)

When using a reverse particle tracking method, source site strength is defined as the

probability that a larva arriving at a recruitment site originated from a specified source site. For

each recruitment sites, source site strength for the recruitment period was derived by summing

the number of particles that originated from or passed through a source site for one month of

the modeling period. Thus, the number of particles at each of the 926 source sites represents

the relative probability or strength of the source sites. Without nearshore retention, source site

strength was calculated using only the particles that originated at a source site at the beginning

of the PLD and by counting each particle one time at its source site. With nearshore retention,

source site strength was calculated using particles that originated at a source site anytime

during the PLD and by counting the particle every time its track passed through a source site.

Thus, the method for implementing nearshore retention resulted in particles being counted

multiple times for different source sites. The source sites were then geographically grouped into

nine source zones (Fig. 1). The nine source zones consisted of (1) the mainland of the SBC, (2)

the north side of the North Channel Islands, (3) the south side of the North Channel Islands, (4-

7) groups of contiguous source sites along the central and southern mainland of the SC Bight,

(8) Catalina Island, and (9) three South Channel Islands. For each recruitment site and month,

the percentage of particles in each zone was calculated, producing a monthly time-series of

source zone strength (SZS) for each zone and recruitment site over the recruitment period.

Larval food availability (Chl)

To estimate food availability during larval transport, surface chlorophyll was used as a

proxy for phytoplankton, which is what urchin larvae use for food (Strathmann 1975). Surface

chlorophyll for the SC Bight was calculated from satellite-derived 5-day composites of surface

chlorophyll-a concentrations (Kahru et al. 2012, 2015, https://spg-satdata.ucsd.edu) and was

available from 1998 to 2013, 16 years of 18-year modeling period. As the satellite data is two-

dimensional and does not represent the chlorophyll in the sub-surface, we recognize that this is

a limitation of our analysis. However, surface chlorophyll has been shown to reflect the overall

seasonal and interannual changes in chlorophyll in the SC Bight (Otero and Siegel 2004;

Henderikx Freitas et al. 2017; Simons and Catlett 2023) and can provide some insight into the

correlation between food availability and larval recruitment. In addition, satellite data was the

only source of chlorophyll data that provided the spatial and temporal coverage needed to use

with the particle tracking model results.

The 5-day composites of surface chlorophyll were averaged over each recruitment

month of March-June to produce monthly surface chlorophyll of the SC Bight from 1998-2013.

For each recruitment month and site, particle exposure to surface chlorophyll was calculated by

first interpolating the model grid onto the monthly surface chlorophyll and then multiplying the

surface chlorophyll in each model grid cell by the corresponding fraction of particles in the grid

cell derived from the PDMs. Using this method, monthly time-series of larval exposure to

chlorophyll (Chl) for each recruitment site was derived in units of mg m^{-3}.

<u>Larval temperature (T_L)</u>

Larval temperature (T_L) represents the temperature that larvae are exposed to during

their PLD. T_L was derived from two sources of temperature produced by the OCM for the

recruitment period. The first source was the particle temperature, which was recorded every 6

hours along the particle track. The second source was the temperature at the source sites,

which was calculated every six hours at 5 m depth in the model grid cell where the source site

was located. A depth of 5 m was selected because it is a common depth of sea urchins in the SC

Bight (Okamoto, D., personal observation). For simulations without nearshore retention, only

particle temperature was used to calculate T_L as particles were tracked for the entire duration of

the PLD. In this case, T_L was calculated by first averaging the particle temperature for each

particle track, producing a mean particle temperature, and then averaging the mean particle

temperatures for all the particles from one recruitment site and one month of the modeling

period, producing a monthly time-series of T_L for each recruitment site in units of $°C$ for the

recruitment period. For simulations with nearshore retention, both particle temperature and

temperature at the source sites were used to calculate T_L as many particles were retained at a

source site for part of the PLD before being transported to a recruitment site. For the particles

that were retained at a source site, the mean particle temperature was calculated by averaging

the temperature at the source site for the time that the particle was retained there and the

particle temperature for the time that the particle was tracked from the source site to the

recruitment site. For the particles that were not retained at a source site, the mean particle

temperature for particles was calculated the same way as the simulations without nearshore

retention. Using both particles that were and were not retained at a source site, T_L for

simulations with nearshore retention was derived by averaging the mean particle temperatures

from one recruitment site and one month of the modeling period, producing a monthly time-

series of T_L for each recruitment site in units of $°C$ for the recruitment period.

<u>Source site temperature (T_S)</u>

The reproductive period for purple urchins in the SC Bight occurs from September to

November. To assess the impact of temperature on adult urchins during the reproductive

period, temperature at the source sites (T_S), the location of the adult urchins, was needed. T_S

was calculated using the following four-step method. First, the monthly mean temperature at 5

m below the surface was calculated from the OCM at each of 926 source sites for the

reproductive period from 1996-2013. Second, the monthly source site strength for the each of

the 926 source sites was averaged over the recruitment period (March-June) for each year,

producing yearly source site strength for each of the 18 years of the modeling period and each

source site. Third, the yearly source site strength for each site was then divided by the sum of

the yearly source site strength for all source sites, producing a fraction for each source site and

year. As reproduction occurs prior to recruitment, the T_S during the reproductive period was

calculated from the source site strength for the following year and thus, limited to a 17-year

modeling period of 1996-2012. Fourth, using the fractions derived from source site strength and

temperature at the source sites, a weighted mean temperature or T_S was calculated for each

recruitment site and month of the reproductive period, producing a 17-year time-series of

monthly T_S for each recruitment site in units of °C.

2.4 Sensitivity study of larval life history parameters

Sensitivity of the five OCI, LDD, SZS, Chl, T_L, and T_S, to variations in three larval life history

parameters of PLD, larval behavior, and nearshore retention was tested. PLD had two options,

static and dynamic. Behavior had three options, passive drifting, DVM20, and DVM40.

Nearshore retention had two options, with and without nearshore retention. For each OCI, 12

simulations were created using all possible combinations of the three parameters and are listed

in Table 1. Simulation 1 with passive drifting behavior, no nearshore retention, and a static PLD

was designated as the baseline simulation as it represents how larval transport is simulated for

many other biophysical modeling studies (Swearer et al 2019).

Simulation Number	Larval Life History Parameters		
	Larval Behavior	Nearshore Retention	PLD
1	Passive Drifting	No	Static
2	Passive Drifting	No	Dynamic
3	Passive Drifting	Yes	Static
4	Passive Drifting	Yes	Dynamic
5	DVM20	No	Static
6	DVM20	No	Dynamic
7	DVM20	Yes	Static
8	DVM20	Yes	Dynamic
9	DVM40	No	Static
10	DVM40	No	Dynamic
11	DVM40	Yes	Static
12	DVM40	Yes	Dynamic

Table 1: Simulations created for each of the five ocean condition indices (OCI), LDD, SZS, Chl, T_L,

and T_S, with three larval life history parameters of larval behavior, nearshore retention, and PLD.

For each ocean condition index, simulations were compared using three statical

measurements, root-mean-square deviation (RMSD), Pearson's correlation coefficient (*r*), and

mean error (ME). The RMSD was used to assess the absolute magnitude of deviation between

the simulations. The *r* was used to assess the linear correlation between the simulations. ME

was used to quantify the increase or decrease in OCI between the simulations. To calculate

RMSD, *r*, and ME for two simulations, the OCI must be in the form of a 1D time-series, which is

the case for LDD, T_L, Chl, and T_S. However, the SZS has 9 1D time-series, with each time-series

corresponding to one zone. In order to calculate RMSD and *r* for two SZS simulations, the 9 SZS

time-series for each simulation were arranged into a 2D matrix with zone versus time and then

linearized into a 1D vector.

To test the sensitivity of each parameter, RMSD, *r*, and ME were calculated for two

simulations where one parameter varied and the other two parameters were held constant. The

two simulations are termed a 'simulation pair'. For PLD, 6 simulations pairs were used where

one simulation had a static PLD and the other had a dynamic PLD while larval behavior and

nearshore retention were the same for both simulations. The 6 simulation pairs for PLD were:

(1) simulations 1 and 2, (2) simulations 3 and 4, (3) simulations 5 and 6, (4) simulations 7 and 8,

(5) simulations 9 and 10, and (6) simulations 11 and 12. For nearshore retention, 6 simulation

pairs were used where one simulation included nearshore retention and the other did not while

larval behavior and PLD were the same for both simulations. The 6 simulations pairs for

nearshore retention were: (1) simulations 1 and 3, (2) simulations 2 and 4, (3) simulations 5 and

7, (4) simulations 6 and 8, (5) simulations 9 and 11, and (6) simulations 10 and 12. For larval

behavior, 12 simulations pairs were used where one simulation had passive drifting and the

other had DVM20 or DVM40 or where one simulation had DVM20 and the other had DVM40

while PLD and nearshore retention was the same for both pairs. The 12 simulation pairs for

larval behavior were: (1) simulations 1 and 5, (2) simulations 2 and 6, (3) simulations 3 and 7, (4)

simulations 4 and 8, (5) simulations 1 and 9, (6) simulations 2 and 10, (7) simulations 3 and 11,

(8) simulations 4 and 12, (9) simulations 5 and 9, (10) simulations 6 and 10, (11) simulations 7

and 11, and (12) simulations 8 and 12.

2.5 Correlation of ocean condition indices to larval recruitment index

To determine which ocean condition index was most likely driving larval recruitment in

the SC Bight, the OCI were compared to an annual larval recruitment index that was derived

from larval recruitment data. Recruitment indexes in the form of standardized larval settlement

anomaly were available for the 5 recruitment sites including an additional site, Ocean Beach,

which was located close to the Scripps recruitment site (Okamoto et al. 2020). A larval

recruitment index was calculated for the northern SC Bight by averaging the indices from the

four SBC recruitment sites, Gaviota, Ellwood, Stearns, and Anacapa, and for southern SC Bight

by averaging the indexes from the recruitment sites for Scripps and Ocean Beach sites (Table

S1). Northern and southern SC Bight recruitment indexes were developed due to the

synchronous patterns of recruitment at the northern versus the southern recruitment sites

observed by Okamoto et al. (2020), indicating that forces driving recruitment are on a regional

versus a local scale. As the recruitment indices were on an annual time scale, the monthly time-

series' of the OCI were averaged over the recruitment period for LDD, SZS, Chl, and T_L, and over

reproductive period for T_S for each year to produce annual time-series'. The annual time-series

of the OCI were then compared to the northern recruitment index for the Gaviota, Ellwood,

Stearns, and Anacapa recruitment sites and the southern recruitment index for the Scripps

recruitment site by calculating *r* for the 12 simulations. LDD, T_L, Chl, and T_S were in the form 1D

time-series and could be used directly to calculate *r*, but SZS was a 2D time-series and had to be

converted into a 1D time-series to compare to the 1D larval recruitment index. To do this, 1D

SZS for each simulation and site was calculated by summing SZS over all possible combination of

one to three zones. For the 9 zones, 129 combinations of 1 to 3 zones were used, producing 129

1D SZS time-series for each simulation and site. Then r was calculated between the 129 1D SZS

time-series and the larval recruitment index. To indicate that any of the five OCI may be driving

recruitment, r would have to be of a similar high magnitude and sign across all recruitment sites

for a single simulation.

3.0 Results

Our results are presented in two parts: first, the sensitivity of OCI to the larval life history

parameters and second, the correlation of OCI to the larval recruitment index. The five OCI,

which are LDD, SZS, Chl, T_L, and T_S, fall into three categories: ocean circulation represented by

LDD and SZS, food supply represented by Chl, and temperature represented by T_L and T_S. The

larval life history parameters include PLD, nearshore retention, and behavior. PLD has two

options, a static PLD of 42 days and a dynamic PLD, which is as a function of temperature.

Nearshore retention has two options, included or not included in the simulation. Behavior has

three options, passive drifting, DVM with daytime depth of 20m (DVM20), and DVM with a

daytime depth of 40m (DVM40). Using all possible combinations of these larval life history

parameters, 12 particle tracking simulations, shown in Table 1, are produced for each of the OCI

and the recruitment sites over the 18-year modeling period of 1996-2013. Due to their

location, the Gaviota, Ellwood, Stearns, and Anacapa sites are referred to collectively as the SBC

sites, and the Gaviota, Ellwood, and Stearns sites are referred to collectively as the SBC

mainland sites. Figures of the five OCI for the five recruitment sites and the 12 simulations are

included in the SI (Fig. S3-S11).

3.1 Sensitivity of ocean condition indices to larval life history parameters

<u>Larval dispersal distance</u>

Larval dispersal distance (LDD) represents the mean distance that the larvae have

traveled from the source sites to the recruitment sites and reflects changes in circulation. For

the baseline simulation of passive drifting behavior, a static PLD, and no nearshore retention,

LDD is shown for the recruitment period of March-June in Fig. 3(a,b) for Gaviota and Scripps.

Gaviota and Scripps are shown because they are the recruitment sites that are farthest apart

and represent the northern versus southern conditions in the SC Bight. For the baseline

simulation, LDD is extremely variable with each recruitment site ranging an order of magnitude

over the modeling period. Baseline LDD for the modeling period has a mean and standard

deviation of 79.5 km and 51.6 km for Gaviota, 84.8 km and 39.6 km for Ellwood, 99.7 km and

30.8 km for Stearns, 71.1 km and 25.3 km for Anacapa, and 88.4 km and 47.8 km for Scripps.

Planktonic larval duration

To examine the sensitivity of LDD to PLD, the six simulations pairs with static versus

dynamic PLD are compared and show very strong agreement for all recruitment sites. For all

recruitment sites, RMSD ranges 1.8-16.9 km, *r* ranges from 0.95-0.99, and ME ranges from 0.01-

10 km.

Nearshore retention

For nearshore retention, the sensitivity testing of LDD for all recruitment sites shows a

strong linear correlation between all simulation pairs with and without nearshore retention for

all sites, resulting in a *r* ranging from 0.73-0.96 with a mean of 0.88, but displays a large range of

agreement in magnitude among the simulation pairs as a function of behavior and site location.

For the SBC sites, the weakest agreement for LDD is observed for the two passive drifting

simulation pairs. The two simulation pairs with passive drifting have RMSD ranging from 33.5-

44.1 km for Gaviota, Ellwood, and Stearns and from 19.2-19.8 km for Anacapa. ME shows a

decrease in LDD of 28.2-34.6 km for Gaviota, Ellwood, and Stearns and 14.0-14.5 km for

Anacapa when nearshore retention is included in the simulation with passive drifting. The

decrease in LDD for Gaviota when nearshore retention is included with passive drifting is shown

in Fig. 3(a). For SBC sites, stronger agreement is observed for the four DVM20 and DVM40

simulation pairs with RMSD ranging from 11.4-25.6 km. ME for the four DVM20 and DVM40

simulations pairs shows a decrease in LDD of 4.6-10.9 km when nearshore retention is included,

which is a smaller decrease in LDD than for the two passive drifting pairs. For Scripps, LDD

shows similar magnitudes of agreement for all six simulation pairs with RMSD ranging from

55.7-67.0 km. ME for the four simulation pairs with passive drifting and DVM20 show a

decrease in LDD of 44.9-54.0 km when nearshore retention is included, but the two simulation

pairs with DVM40 show an increase in LDD of 19.2-21.8 km when nearshore retention is

included. The decrease in LDD for Scripps when nearshore retention is included with passive

drifting is shown in Fig. 3(b).

Larval behavior

For the SBC mainland sites, LDD shows different patterns of sensitivity in magnitude and

linear correlation as functions of behavior and nearshore retention. For magnitude, LDD shows

the weakest agreement for the four simulation pairs comparing passive drifting behavior to

DVM20 or DVM40 without nearshore retention, medium agreement for the four simulation

pairs comparing passive drifting behavior to DVM20 or DVM40 with nearshore retention, and

the strongest agreement in LDD for the four simulation pairs comparing DVM20 to DVM40 with

and without nearshore retention. The four simulation pairs comparing passive drifting behavior

to DVM20 or DVM40 without nearshore retention have the largest RMSD ranging from 60.2-

74.3 km. Fig. 3(a) shows LDD for Gaviota for simulations with passive drifting and DVM20

without nearshore retention. The four simulation pairs comparing passive drifting behavior to

DVM20 or DVM40 with nearshore retention have smaller RMSD ranging from 27.9-42.9 km. The

four simulation pairs comparing DVM20 to DVM40 with and without nearshore retention have

the smallest RMSD ranging from 14.8-32.4 km. For the eight simulation pairs comparing passive

drifting with DVM with and without nearshore retention, r ranges widely from 0.30 and 0.69,

and the strength of correlation is not associated with larval behavior or nearshore retention.

The four simulation pairs comparing DVM20 to DVM40 with and without nearshore retention

have the highest r ranging from 0.71-0.91. For the eight simulations pairs with passive drifting

and DVM, LDD has a ME ranging from 17.8-54.6 km, indicating that LDD is smaller for

simulations with DVM than for simulations with passive drifting. The absolute ME between the

four simulations pairs with DVM20 and DVM40 ranges from 4-12 km and does not show a

consistent increase or decrease in LDD, indicating that LDD has similar magnitudes for DVM20

and DVM40 simulations.

For Anacapa, LDD shows similarly strong agreement for all simulations pairs with RMSD

ranging from 20.1-28.7 km and r ranging from 0.56-0.75. The absolute ME for all simulations

pairs is small, ranging from 0 to 15.4 km with a mean of 8.1 km. This small ME implies that

there is no consistent increase or decrease in LDD associated with a specific behavior or

nearshore retention.

For Scripps, LDD shows the largest range of agreement in magnitude and linear

correlation for the simulation pairs of all the five recruitment sites. The four simulation pairs

comparing passive drifting to DVM20 or DVM40 with nearshore retention have the weakest

agreement with RMSD ranging from 115-150 km and r ranging from 0.29-0.42. The four

simulation pairs comparing passive drifting to DVM20 or DVM40 without nearshore retention

have stronger agreement with RMSD ranging from 55-79 km and r ranging from 0.49-0.63. Fig.

3(b) shows LDD for Scripps for simulations with passive drifting and DVM20 without nearshore

retention. The four simulation pairs comparing DVM20 to DVM40 with and without nearshore

retention have the strongest agreement with RMSD ranging from 27-40 km and r ranging from

0.79-0.88. The ME for Scripps shows an increase in LDD of 20-120 km for all simulation pairs,

which means that LDD is larger for simulations with DVM20 or DVM40 than passive drifting and

for simulations with DVM40 than DVM20 regardless if nearshore retention is present. The

magnitude of ME for Scripps does not correlate with a specific behavior or nearshore retention.

<u>Larval source zone strength</u>

Larval source zone strength (SZS) is defined as the probability that larvae have originated

from a specified zone and represents how ocean circulation drives spatial and temporal

variation in SZS. The nine source zones are identified in the SC Bight (Fig. 1). The units for SZS

are the percentage of particles per zone, which is directly proportional to SZS. SZS for the

baseline simulation of passive drifting, no nearshore retention, and a static PLD is shown for the

recruitment period of March-June in Fig. 4(a,d) for Gaviota and Scripps. For all recruitment sites,

baseline SZS is extremely variable in time and space over the 9 zones. Baseline SZS for Gaviota is

similar to Ellwood, Stearns, and Anacapa with RMSD ranging from 5-9% and r ranging from 0.72-

0.90, but very different to Scripps with a RMSD of 18% and a *r* of 0.02. For the baseline

simulation, the maximum SZS per month and zone has a mean and standard deviation of 36%

and 18% for Gaviota, 34% and 14% for Ellwood, 34% and 13% for Stearns, 32% and 13% for 40%

for Anacapa and 20% for Scripps. Maximum SZS occurs predominantly in Zones 1, 3, 8, and 9 for

Gaviota, Ellwood, Stearns, and Anacapa and Zones 6, 7, and 9 for Scripps.

Planktonic larval duration

The sensitivity testing of SZS to PLD shows strong agreement for all simulation pairs with

a static versus a dynamic PLD for all recruitment sites. For all simulation pairs and recruitment

sites, the RMSD ranges 1-9%, the *r* ranges from 0.88-1.0, and the ME ranges from 0-8%.

Nearshore retention

For the SBC mainland sites, SZS shows similar patterns of agreement in magnitude and

linear correlation as function of behavior. SZS shows the weakest agreement for the two

simulation pairs comparing passive drifting with and without nearshore retention, producing

RMSD ranging from 19-21% and *r* ranging from 0.21-0.37. As shown in Fig. 2(g,h) and Fig. 4(a,b)

for Gaviota, the addition of nearshore retention to passive drifting produces in the highest SZS

occurring in the Zone 1, where the recruitment sites are located. For the two passive drifting

simulation pairs, the ME in Zone 1 shows SZS increasing by 46-53% when nearshore retention is

present. SZS shows the strongest agreement for the four simulation pairs comparing DVM20 or

DVM40 with and without nearshore retention, producing RMSD ranging from 12-15% and *r*

ranging from 0.85-0.91. For the four DVM simulation pairs, the ME in Zone 1 shows SZS

increasing by only 18-26% when nearshore retention is present.

For Anacapa, SZS shows similar agreement in magnitude for all simulation pairs with a

RMSD ranging from 11-14%. The SZS for Anacapa shows a weaker linear correlation for the two

passive drifting simulation pairs of r from 0.43-0.44 than the four DVM20 and DVM40

simulation pairs of r from 0.73-0.81. ME for Zone 2, where Anacapa is located, displays an

increase in SZS by 28-29% when nearshore retention is added to the passive drifting

simulations. ME for Zone 2 shows a smaller increase in SZS of 11-17% when nearshore

retention is added the DVM simulations.

For Scripps, the magnitude of SZS is very similar for all simulations pairs with RMSD

ranging from 18-20%. The linear correlation of SZS is slightly higher for the passive drifting

simulation pairs, ranging from 0.59-0.69, than the DVM simulation pairs, ranging from 0.43-

0.49. For passive drifting and DVM, the SZS for Zone 7 increases when nearshore retention is

present as shown in Fig. 2(j,k) and Fig. 4(d,e). ME for Zone 7 shows a similar increase in SZS of

39-48% when nearshore retention is added the passive drifting and DVM simulations.

Larval behavior

For the SBC mainland sites, the sensitivity testing of SZS to behavior shows the weakest

agreement for the four simulation pairs comparing passive drifting to DVM20 or DVM40

without nearshore retention, producing RMSD ranging from 22-26% and r ranging from 0.10-

0.36. The passive drifting simulations without nearshore retention produce similarly low levels

of SZS across all zones (Fig. 2(g) and Fig. 4(a) for Gaviota), while the DVM20 and DVM40

simulations without nearshore retention produce the highest SZS in Zone 1, the zone where the

recruitment site is located (Fig. 2(i) and Fig. 4(c) for Gaviota). The ME in Zone 1 shows that SZS

for DVM without nearshore retention is higher by 44-55% than SZS for passive drifting without

nearshore retention. The sensitivity testing of SZS displays the strongest agreement between

the eight simulation pairs comparing DVM20 to DVM40 with and without nearshore retention

and passive drifting to DVM20 or DVM40 with nearshore retention, producing a RMSD ranging

from 4-13% and a r ranging from 0.90-99. The ME in Zone 1 shows that SZS from DVM

simulations with nearshore retention is higher by 17-19% than SZS from passive drifting

simulation with nearshore retention and that SZS from DVM40 simulations is higher by 1-8%

than SZS from DVM20 simulations with and without nearshore retention.

For Anacapa, testing the sensitivity of SZS to behavior shows the strongest agreement

for the four simulations pairs comparing DVM20 to DVM40 with and without nearshore

retention, producing RMSD ranging from 8-12% and r ranging from 0.80-0.88. The ME in Zone 2

shows that SZS from DVM40 simulations is higher by only 2-7% than SZS from DVM20

simulations. For the eight simulations pairs comparing passive drifting to DVM20 or DVM40 with

and without nearshore retention, SZS for Anacapa shows strong agreement in magnitude with

RMSD ranging from 14-18%, but weak linear correlation with r ranging from 0.35-0.60. The ME

in Zone 2 shows that SZS from DVM simulations is smaller by 8-27% than SZS from passive

drifting simulations. This pattern is the opposite of what is observed for the SBC mainland sites

where SZS in Zone 1, the location of the recruitment sites, is higher for DVM than passive

drifting. For Anacapa, the ME for Zones 1 and 3 shows that SZS from DVM simulations is higher

by 17-25% and 6-13% respectively than SZS from passive drifting simulations.

For Scripps, the sensitivity testing of SZS shows the weakest agreement for the four

simulations pairs comparing passive drifting to DVM20 or DVM40 without nearshore retention,

producing RMSD ranging from 16-18% and r ranging from 0.43-0.50. For these four simulation

pairs, the ME is largest for Zone 7 and shows SZS decreasing by a 16-22% when DVM is present

versus passive drifting without nearshore retention. SZS shows strong agreement of similar

magnitude for the eight simulations pairs comparing passive drifting to DVM20 or DVM40 with

nearshore retention and DVM20 to DVM40 with or without nearshore retention, producing

RMSD ranging from 4-13% and r ranging from 0.78-0.98. The ME in Zone 7 shows that SZS is

higher for DVM simulations with nearshore retention by 22-27% for passive drifting simulations

with nearshore retention and that SZS for DVM40 simulations is only 1-2% than DVM20

simulations.

<u>Larval food availability</u>

Larval food is represented by surface chlorophyll (Chl), which was available from 1998-

2013, 16 years of the modeling period. For the five recruitment sites, Chl for the baseline

simulation of passive drifting behavior, a static PLD, and no nearshore retention, is shown in Fig.

3(c,d) for the recruitment period of March-June for Gaviota and Scripps. Baseline Chl is most

similar between the Gaviota site and the Ellwood and Stearns sites with a RMSD of 0.70 mg m^{-3}

and 0.75 mg m^{-3} and a r of 0.94 and 0.90 respectively and least similar between the Gaviota site

and the Anacapa and Scripps sites with a RMSD of 1.24 mg m^{-3} and 1.57 mg m^{-3} and a r of 0.88

and 0.74 respectively. Averaged over the modeling period, mean baseline Chl is 2.95 mg m^{-3} for

Gaviota, 2.65 mg m^{-3} for Ellwood, 2.74 mg m^{-3} for Stearns, 2.06 mg m^{-3} for Anacapa, and 1.84

mg m^{-3} for Scripps. The similar mean baseline Chl between the SBC mainland sites of Gaviota,

Ellwood, and Stearns and the decreasing mean baseline Chl from Gaviota to Anacapa to Scripps

reflects the mean north-south gradient of Chl in the SC Bight (Fig. 5(a)).

Planktonic larval duration

The sensitivity testing of Chl to PLD shows strong agreement between all simulation

pairs with a static PLD versus a dynamic PLD. For the six simulation pairs tested, RMSD ranges

0.10-0.37 mg m^{-3}, r ranges from 0.98-1.0, and ME ranges from 0.01-0.20 mg m^{-3}.

Nearshore retention

For the six simulation pairs with and without nearshore retention, the sensitivity testing

of Chl for all recruitment sites shows strong linear correlations with r ranging from 0.91-1.0, but

displays a range of agreement in magnitude among the simulation pairs as a function of

behavior. For all recruitment sites, Chl shows a slightly weaker agreement in magnitude for the

two passive drifting simulations pairs with RMSD ranging from 0.49-0.64 mg m^{-3} for the SBC

sites and 0.76-0.81 mg m^{-3} for Scripps (Fig. 3(c,d)) than the four DVM20 and DVM40 simulation

pairs with RMSD ranging from 0.29-0.46 mg m^{-3} for the SBC sites and 0.44-0.66 mg m^{-3} for

Scripps. For all sites, the ME for Chl shows that the passive drifting simulations with nearshore

retention are higher by 0.22-0.68 mg m^{-3} than the passive drifting simulations without

nearshore retention. For the SBC sites, Chl is similar in magnitude for DVM simulations with and

without nearshore retention with ME ranging from 0.02-0.09 mg m^{-3}. For Scripps, the ME

shows that Chl from the DVM simulations with nearshore retention are higher by 0.19-0.43 mg

m^{-3} than Chl from the DVM simulations without nearshore retention.

Larval behavior

Chl shows strong linear correlations for the 12 larval behavior simulation pairs for all

recruitment sites with r ranging from 0.80-0.99, but differences in magnitude as a function of

behavior, nearshore retention, and site location. For the SBC sites, Chl shows the weakest

agreement for the four simulation pairs comparing passive drifting to DVM20 or DVM40

without nearshore retention, producing RMSD ranging from 1.23-1.62 mg m^{-3}. For these four

simulation pairs, the ME shows that Chl for the DVM simulations is greater by 0.32-0.85 mg m^{-3}

than the passive drifting simulations. For the SBC sites, Chl shows the slightly stronger

agreement for the four simulation pairs comparing passive drifting to DVM20 or DVM40 with

nearshore retention, producing RMSD ranging from 0.67-1.45 mg m^{-3}. For these four simulation

pairs, the ME shows that Chl for the DVM simulations is greater by 0.03-0.52 mg m^{-3} than the

passive drifting simulations. For the SBC sites, Chl shows the strongest agreement for the four

simulation pairs comparing DVM20 and DVM40 with RMSD ranging from 0.04-0.38 mg m^{-3} and

ME from 0.01-0.12 mg m^{-3}. For Scripps, Chl shows a range of weak to strong agreement for the

12 simulations pairs, but the magnitude of agreement is not correlated with larval behaviors or

nearshore retention. For Scripps, Chl shows a RMSD ranging from 0.27-0.74 mg m^{-3} and ME

ranging from 0.10-0.41 mg m^{-3}.

<u>Larval temperature</u>

Larval temperature (T_L) is defined as the temperature that larvae are exposed to during

their PLD. T_L for the baseline simulation of passive drifting behavior, a static PLD, and no

nearshore retention is shown in Fig. 3(e,f) for the recruitment period of March-June for Gaviota

and Scripps. For the baseline simulation, T_L for all months and recruitment sites is below 16°C

and ranges from 9.7-15.6°C. For all recruitment sites, the yearly change in baseline T_L over the

recruitment period averages 1.5°C with a maximum of 2.8°C and does not show a consistent

pattern of warming or cooling. Baseline T_L is very similar between mainland SBC sites of Gaviota,

Ellwood, and Stearns. Comparing baseline T_L for the Gaviota site to the Ellwood and Stearns

sites, RMSD and *r* range from 0.28-0.45°C and 0.92-0.96 respectively. Greater differences in

baseline T_L are observed between Gaviota and Anacapa with a RMSD of 0.75°C and a r of 0.86

and between Gaviota and Scripps with a RMSD of 1.4°C and a r of 0.59. Averaging over the

modeling period, mean baseline T_L is 12.4°C, 12.9°C, and 13.5°C for Gaviota, Anacapa, and

Scripps respectively, reflecting the mean north-south temperature gradient in the SC Bight (Fig.

5(b)).

Planktonic larval duration

The sensitivity testing of T_L to PLD shows strong agreement between the six simulation

pairs comparing static versus dynamic PLD. For the six simulation pairs and all recruitment sites,

the RMSD ranges 0.13-0.36°C, the r ranges from 0.96-0.99, and ME ranges from 0.00-0.18°C.

Nearshore retention

The sensitivity testing of T_L to nearshore retention shows strong linear correlations for all

simulation pairs and recruitment sites with r ranging from 0.83-0.98. However, differences in

magnitude of T_L are observed with the weakest agreement for the simulation pairs with passive

drifting (Fig. 3(e,f)) and the strongest agreement for the simulation pairs with DVM20 or DVM40

for all recruitment sites. T_L for the two simulation pairs with passive drifting have a RMSD

ranging from 0.54-0.80°C for the SBC sites and a RMSD ranging from 0.87-1.0°C for Scripps. In

contrast, the four simulation pairs with DVM20 or DVM40 have smaller RMSD ranging from

0.21-0.44°C for the SBC sites and a RMSD ranging from 0.55-0.82°C for Scripps. ME for T_L did

not show consistent patterns as a function of nearshore retention or recruitment site.

Larval behavior

The sensitivity testing of T_L to behavior shows strong linear correlations for all simulation

pairs and recruitment sites with r ranging from 0.85-0.99, but varies in magnitude of agreement

as a function of behavior and site location. For the SBC mainland sites, T_L shows the weakest

agreement for the four simulation pairs comparing passive drifting to DVM20 or DVM40

without nearshore retention, which have RMSD ranging from 0.44-0.91°C, and the strongest

agreement for the four simulation pairs comparing DVM20 to DVM40 with or without

nearshore retention, which had a RMSD ranging from 0.23-0.35°C. The strength of agreement

of the four simulation pairs comparing passive drifting to DVM20 or DVM40 with nearshore

retention falls between the simulations with the weakest and strongest agreement and has a

RMSD ranging from 0.37-0.48°C. For Anacapa and Scripps, T_L has the weakest agreement for

the two simulation pairs comparing passive drifting to DVM20 without nearshore retention,

which had a RMSD ranging from 0.49-0.54°C and 0.72-0.80°C respectively. T_L shows slightly

stronger agreement for the four simulation pairs comparing passive drifting to DVM40 and

DVM20 to DVM40 without nearshore retention with a RMSD ranging from 0.35-0.48°C for

Anacapa and from 0.43-0.53°C for Scripps. T_L shows the strongest agreement for the six

simulations pairs that included nearshore retention with a RMSD ranging from 0.24-0.34°C for

Anacapa and from 0.34-0.43°C for Scripps. ME for T_L did not show consistent patterns as a

function of behavior, nearshore retention, or recruitment site.

<u>Source site temperature</u>

Source site temperature (T_S) is defined as the temperature that adult urchins at the

source sites are exposed to during the reproductive period of September to November. Since

the reproductive period occurs in the fall while spawning and recruitment occurs in the

following spring, the modeling period for T_S is limited to 17 years from 1996-2012. T_S for the

baseline simulation of passive drifting, a static PLD, and no nearshore retention is shown in Fig.

3(g,h) for Gaviota and Scripps. For all months and recruitment sites, T_S for the baseline

simulation ranges from 14.2-20.5°C. Mean baseline T_S over the modeling period is 17.2°C for

Gaviota, 17.3°C for Ellwood, 17.4°C for Stearns, 17.2°C for Anacapa, and 17.5°C for Scripps. For

all years except 1998, the baseline T_S decreases over the reproductive period with the change in

T_S ranging from 0.5°C to 4°C and averaging 2.4°C. When baseline T_S at Gaviota is compared to

the other recruitment sites, T_S is found to have a strong linear correlation for all sites with r

ranging from 0.97 to 1.0. Baseline T_S at Gaviota is similar in magnitude to Ellwood, Stearns, and

Anacapa with RMSD ranging from 0.14-0.25°C. The largest difference in magnitude of baseline

T_S is between Gaviota and Scripps with a RMSD of 0.49°C, reflecting the mean north-south

temperature gradient in the SC Bight (Fig. 5(c)).

Planktonic larval duration

The sensitivity testing of T_S to PLD shows strong agreement for simulation pairs with

static and dynamic PLD for all recruitment sites. For the six simulation pairs tested for PLD, the

RMSD ranges from 0.02-0.22°C, the r ranges from 0.99-1.00, and the ME ranges from 0.01-

0.10°C.

Nearshore retention

Testing the sensitivity of T_S to nearshore retention shows strong linear correlations for all

simulation pairs and recruitment sites with r ranging from 0.91-1.00, but displays a range of

differences in magnitude as a function of behavior and site location. For the SBC mainland sites,

the sensitivity testing of T_S to nearshore retention shows the weakest agreement for the two

passive drifting simulation pairs, which produce RMSD ranging from 0.68-1.1°C, and the

strongest agreement for the four DVM20 and DVM40 simulation pairs, which produce RMSD

from 0.19-0.45°C. The ME for the SBC mainland sites shows that that T_S for the passive drifting

simulations with nearshore retention is smaller by 0.53-0.91 °C than the passive drifting

simulations without nearshore retention and that T_S for the DVM simulations with nearshore

retention is smaller by 0.02-0.33 °C than the DVM without nearshore retention. For Anacapa,

the sensitivity testing of T_S to nearshore retention shows a similar but less pronounced pattern

as the SBC mainland sites with the two passive drifting simulation pairs producing RMSD from

0.43-0.45°C and the four DVM20 and DVM40 simulation pairs producing RMSD from 0.17-0.27°.

The ME for Anacapa shows that that T_S for the passive drifting simulations with nearshore

retention is smaller by 0.24-0.30 °C than the passive drifting simulations without nearshore

retention and that T_S for the DVM simulations with nearshore retention is smaller by 0.05-0.0.8

°C than the DVM without nearshore retention. For Scripps, sensitivity testing of T_S to nearshore

retention does not show changes in magnitude that correlate with behavior and produces

RMSD from 0.38-0.52°C for all simulation pairs. The ME for Scripps shows that that T_S for all

simulations with nearshore retention are greater by 0.31-0.40 °C than all simulations without

nearshore retention, the opposite of the SBC sites.

Larval behavior

Testing the sensitivity of T_S to behavior shows strong linear correlations for all simulation

pairs and recruitment sites with *r* ranging from 0.86-0.99 and differences in magnitude as a

function of behavior, nearshore retention, and site location. For the SBC mainland sites,

sensitivity testing of T_S to behavior shows the weakest agreement for the four simulation pairs

comparing passive drifting to DVM20 or DVM40 without nearshore retention, producing RMSD

from 0.97-1.2°C. T_S for the SBC mainland sites shows the strongest agreement for the four

simulation pairs comparing DVM20 to DVM40 with and without nearshore retention, producing

RMSD from 0.07-0.31°C. T_S agreement for the four simulations pairs comparing passive drifting

to DVM20 or DVM40 with nearshore retention fall in between the simulations pairs with the

strongest and weakest agreement, producing RMSD from 0.32-0.48°C. The ME for the SBC

mainland sites shows that that T_S for the DVM simulations without nearshore retention is

smaller by 0.73-0.95°C than the passive drifting simulations without nearshore retention and

that T_S for the DVM simulations with nearshore retention is smaller by 0.24-0.39 °C than the

passive drifting simulations with nearshore retention. The ME of 0.01-0.12°C suggests that

there is little difference in T_S between DVM20 and the DVM40 simulations with and without

nearshore retention.

For Anacapa and Scripps, agreement of T_S between simulations pairs is function of larval

behavior and nearshore retention similar to the SBC mainland sites, but with overall stronger

agreement between the simulation pairs than the SBC mainland sites. T_S for the four simulation

pairs comparing passive drifting to DVM20 or DVM40 without nearshore retention produces the

weakest agreement with RMSD ranging from 0.60-0.67°C for Anacapa and from 0.28-0.33°C for

Scripps. T_S for the four simulation pairs comparing DVM20 to DVM40 with and without

nearshore retention produces the strongest agreement with RMSD ranging from 0.13-0.32°C for

Anacapa and from 0.05-0.18°C for Scripps. T_S for the four simulation pairs comparing DVM20 to

DVM40 with nearshore retention produces medium agreement with RMSD ranging from 0.32-

0.25°C for Anacapa and from 0.16-0.22°C for Scripps. For Anacapa, the ME shows that that T_S

for the DVM simulations is smaller than T_S for the passive drifting simulations by 0.43-0.47°C

without nearshore retention and by 0.22-0.26 C with nearshore retention. For Scripps, the ME

shows that the T_S for the DVM simulations with and without nearshore retention is smaller than

T_S for the passive drifting simulations with and without nearshore retention by 0.11-0.18°C. For

both Anacapa and Scrips, the ME of 0.01-0.03°C suggests that there is little difference in T_S

between the DVM20 and the DVM40 simulations with and without nearshore retention.

3.2 Comparison of ocean condition indices to larval recruitment index

To determine the influence of ocean conditions on larval recruitment, the 12 simulations

for each ocean condition index are linearly correlated to a larval recruitment index for all

recruitment sites. A meaningful linear correlation between an ocean condition index and the

larval recruitment index should have strong agreement or high values of r. A northern SC Bight

recruitment index is used for the Gaviota, Ellwood, Stearns, and Anacapa sites, which has five

poor recruitment years of 1998 and 2004-2007 and 13 average or above-average recruitment

years. A southern SC Bight recruitment index is used for the Scripps site, which has 13 poor

recruitment years and five average and above-average recruitment years of 1996 and 1999-

2002.

<u>Larval dispersal distance</u>

The strength of correlation between LDD simulations and the larval recruitment index

varies as a function of larval behavior and recruitment site. For the SBC sites, LDD for all passive

drifting simulations have weak and inconsistent correlations with r ranging from -0.23 to 0.24.

For the SBC mainland sites, LDD for the DVM20 simulations show a weak but consistent inverse

correlation with r ranging from -0.15 to -0.28. For Anacapa, LDD for the DVM20 simulations has

very weak correlations with r ranging from -0.07 to 0.0. For the SBC sites, LDD for the DVM40

simulations have the strongest correlations with r ranging from -0.18 to -0.54. For Scripps, LDD

has similarly strong and consistently inverse correlations for all simulations with r ranging from -

0.48 to -0.73. Although ranging in correlation strength, the consistent presence of negative r for

the DVM20 and DVM40 simulations for the SBC mainland and Scripps sites suggests that LDD is

inversely proportional to the larval recruitment index and that shorter LDD is correlated with

higher recruitment.

<u>Source zone strength</u>

For SZS, zone combinations are identified that display a positive correlation with the

larval recruitment index, implying that more larvae originate from these zones in average and

above-average recruitment years than in poor recruitment years. For all sites, simulations, and

zone combinations, SZS displays a very wide range of positive r from 0.01-0.76. The two

simulations with DVM20 behavior, no nearshore retention, and a static or dynamic PLD display

the strongest correlations across all sites with r ranging from 0.48-0.55 for the SBC sites and

from 0.67-0.70 for the Scripps site. The zones associated with these correlations are Zones 1

and 3 for the SBC sites and Zones 6, 7 and 8 for the Scripps site.   For the SBC sites, SZS is higher

for Zone 1 than Zone 3, but the difference between Zones 1 and 3 is greater for the SBC

mainland sites than Anacapa. For the SBC mainland sites, the mean SZS for the modeling period

ranges from 59-70% for Zone 1 and 11-12% for Zone 3. For Anacapa, the mean SZS ranges from

35-38% for Zone 1 and 20-21% for Zone 3.   For Scripps, SZS is higher for Zone 6 than Zones 7

and 8 and is lower for Zones 6-8 than Zone 1 for the SBC sites. The mean SZS for the modeling

period ranges from 22-30% for Zone 6, 15-17% for Zone 7, and 12-17% for Zone 8.

To display the linear correlation between SZS and the larval recruitment index, Fig. 6(a,b)

shows plots of SZS versus larval recruitment index for the simulation with DVM20, no nearshore

retention, and a static PLD for Gaviota and Scripps. Using Zones 1 and 3 for Gaviota, this

simulation has a *r* value of 0.55. Zone 1 is located on the SBC mainland and contains Gaviota,

and Zone 3 is located on the southern coast of the North Channel Islands (Fig. 1). As shown in

Fig. 6(a) for Gaviota, the SZS has a mean of 80% and a standard deviation of 23% for average

and above-average recruitment years and a mean of 54% and a standard deviation of 23% in the

poor recruitment years. SZS for 1998 is an outlier with poor recruitment and high SZS. Using

Zones 6, 7, and 8 for Scripps, the simulation with DVM20, no nearshore retention, and a static

or dynamic PLD has a *r* value of 0.70. Zone 7 contains the Scripps site and is located on the

southern most part of the SC Bight mainland. Zone 6 is located directly north of Zone 7 along

the SC Bight mainland. Zone 8 contains the Catalina Island coastline. As shown in Fig. 6(b) for

Scripps, the SZS has a mean of 81% and a standard deviation of 20% for average and above-

average recruitment years and a mean of 54% and a standard deviation of 23% for the poor

recruitment years.

<u>Food supply</u>

Comparing Chl to the larval recruitment index produces only a weak and inconsistent

linear correlations for all recruitment sites and simulations. For Chl, *r* ranges from -0.52 to 0.05

with a mean of -0.13 for all simulations and recruitment sites. These results imply that that

larval food supply is unlikely to be driver of larval recruitment of purple sea urchins in the SC

Bight.

<u>Larval temperature</u>

When T_L is compared to the larval recruitment index, only weak and inconsistent linear

correlations are found across all recruitment sites and simulations. For T_L, *r* ranges from -0.35 to

0.29 with a mean of -0.13 for all simulations and recruitment sites. These results suggest that

that larval temperature is not a likely driver of larval recruitment of purple sea urchins in the SC

Bight.

<u>Source site temperature</u>

Out of all the OCI, T_S shows the highest correlation with the larval recruitment index,

producing *r* with a mean of -0.69 and a standard deviation of 0.08 for all sites and simulations.

The negative *r* means that high T_S produces to low recruitment and that low T_S produces

average to above-average recruitment. For the SBC mainland sites, T_S simulations that have

DVM and/or nearshore retention produce stronger correlations with *r* ranging from -0.66 to -

0.78 than the T_S simulations that have passive drifting without nearshore retention with *r*

ranging from -0.44 to -0.55. For Anacapa and Scripps, T_S produces correlations with *r* ranging

from -0.58 to -0.80 that do not correlate in strength with larval behavior or nearshore retention.

To display the linear correlation between T_S and larval recruitment index, Fig. 6(c,d)

shows plots of T_S versus the larval recruitment index for the simulation with DVM20, no

nearshore retention, and a static PLD for Gaviota and Scripps. This simulations produces a *r* of -

0.75 for Gaviota and -0.58 for Scripps. For Gaviota, the 13 years with average to above-average

recruitment correlate with lower T_S ranging from 14.9-16.6°C and averaging 16.0°C. Three of

the poor recruitment years (1998, 2005, 2007) correlate with high T_S of 17.8-18.0°C, but the

poor recruitment years of 2004 and 2006 deviate from the linear correlation with lower T_S of

16.4°C and 16.5°C respectively. For Scripps, the 13 poor recruitment years have higher T_S,

ranging from 17.6-19.9°C and averaging 18.2°C. The four years with average to above-average

recruitment have lower T_S ranging from 15.9-17.6°C and averaging 17.1°C.

4.0 Discussion

This study assesses the impacts of changes in circulation, temperature, and food supply

on larval recruitment of purple sea urchins. Changes in circulation are quantified using LDD and

SZS, which are calculated directly from particle tracks. Temperature is quantified through T_L and

T_S, which are calculated from the OCM's temperature and particle tracks. Food supply is

quantified using Chl, which is calculated from surface chlorophyll from satellite data and particle

tracks. To accurately model larval recruitment, the larval life history of purple sea urchins must

be accurately represented in the biophysical model. Although some aspects of the larval life

history of purple sea urchins are known, such as the spawning and reproductive periods and

potential source sites, uncertainty remains for three important aspects of larval life history, PLD,

nearshore retention, and larval behavior. To address this uncertainty, sensitivity of the five OCI

to these three larval life history parameters is discussed in Section 4.1. Derived from the 12

simulations with all possible combinations of larval life history parameters, each ocean

condition index is then compared to a larval recruitment index to identify the potential drivers

of larval recruitment, which is discussed in Section 4.2.

4.1 Sensitivity of ocean condition indices to larval life history parameters

<u>Planktonic larval duration</u>

All OCI shows very strong agreement between the simulation pairs with static and

dynamic PLDs. This result initially appears surprising as the static PLD of 42 days is significantly

different from the dynamic PLD, which ranges from 26-56 days with a mean of 37 days and

standard deviation of 8 days (Fig. S2). However, Simons et al. (2013) found that particle

dispersal in the SC Bight is much more sensitive to changes in particle tracking times or PLDs

when PLDs are greater than ~20 days and concluded that the changes in particle dispersal

patterns decrease with increasing PLD because the particles become more evenly mixed over

increasingly larger areas. To determine if this could be true for our study, r is calculated between

the PDM with a PLD of 42 days and daily PDMs with PLDs of 1-56 days for each month,

simulation, and recruitment site, producing 4,320 values of r for each day from 1-56 days. As

PDMs are 2D particle density matrices, PDMs are vectorized for a grid cell to grid cell calculation

of r following the methods in Simons et al. (2013). Fig. 7 displays mean $r \pm 1$ standard deviation

versus PLD for all r values and shows that there is minimal difference between the PDMs with a

42-day PLD and PDMs with PLDs greater than ~25 days. As all OCI are derived from the particle

tracks, which are used to create the PDMs, this similarity between the PDM with a PLD of 42-

days and PDMs with PLDs greater than ~25 days explains why the OCI are relatively insensitive

to the static PLD of 42 days versus the dynamic PLD with a range of 26-56 days.

<u>Nearshore retention and larval behavior</u>

Two main trends are observed in the sensitivity testing of OCI to nearshore retention

and larval behavior. The first trend is that the OCI representing circulation, LDD and SZS, show

much higher sensitivity to nearshore retention and larval behavior than the OCI representing

food supply, Chl, and temperature, T_L, and T_S, for all simulations and recruitment sites. The

second trend is that simulation pairs for nearshore retention and larval behavior show similar

levels of sensitivity between the SBC mainland sites for all OCI and varying levels of sensitivity

between the SBC mainland sites, the Anacapa site, and the Scripps sites.

Both trends appear to derive from differences in horizontal patterns of circulation,

temperature, and chlorophyll within SC Bight. Fig. 5(a) shows the mean surface chlorophyll from

satellite data averaged temporally over the recruitment period for the modeling period of 1996-

2013. Fig. 5(b,c) displays the mean temperature from the OCM averaged vertically over the top

50 m of the water column and temporally over the recruitment and reproductive periods

respectively for the modeling period. Fig. 5(d) shows the mean current from the OCM averaged

vertically over the top 50 m of the water column and temporally over the recruitment period for

the modeling period. The top 50 m of the water column is selected because ~70% of the

particles with passive drifting remain the top 50 m of the water column and 100% of the

particles with DVM remain in the top 40 m of the water column. As shown in Fig. 5, the

circulation, temperature, and chlorophyll for the areas near the recruitment sites are described

below.

In the northern SC Bight, the mean circulation in the SBC, where the SBC mainland sites

are located, is dominated by a cyclonic eddy, which has the strongest mean velocities in the SC

Bight up to 21 cm s^{-1} and penetrates throughout the top 100 m of the water column (Oey et al.

2004; Dong et al. 2009). The strong eddy circulation creates a region of high residence times

and particle accumulation in the SBC (Simons et al. 2015; Simons and Catlett 2023). The SBC is

the coldest region in the SC Bight with mean temperatures 4-5°C lower than the southern SC

Bight. The mean temperature immediately surrounding the SBC is ~1.5°C higher than the mean

temperature within the SBC. Driven by spring upwelling and high residence times in the SBC

(Simons and Catlett 2023; Brokaw et al. 2024), the mean surface chlorophyll in the SBC is the

highest in SC Bight with a maximum of 2.7 mg m^{-3} and drops to low levels of 0.1 - 1 mg m^{-3}

outside of the SBC.

Located on Anacapa Island at the edge of the SBC, the Anacapa site is surrounded by the

weakest mean velocities, ~3 cm s^{-1}, of the five recruitment sites and sits at the junction between

flows entering the SBC and the large cyclonic eddy in the central SC Bight. The Anacapa site is

located in the transition zone between the SBC and the central and southern SC Bight, where

strong gradients in mean temperature and surface chlorophyll are present.

In the southern SC Bight, the circulation near the Scripps site shows southward flow

along the coastline with mean velocities up to 10 cm s^{-1} and which decreases an order of

magnitude to the north and east. Unlike the SBC which has consistent mean circulation to at

least 100 m depth, mean circulation along the southern coastline varies with depth, displaying

stronger southward flows in the top 50 m of the water column and weaker northward flows

from 50-100 m depth (Dong et al. 2009). Defined by an open coastline with a narrow shelf,

circulation in the region near the Scripps site produces shorter residence times and greater

offshore dispersal than the SBC (Dong et al. 2012). Except for ~5 km adjacent to the coastline,

the southern SC Bight displays smaller north-south mean temperature gradients of 0.1-0.2°C

per 10 km than the area surrounding the SBC of 0.3-0.4°C per 10 km. The mean surface

chlorophyll throughout the southern SC Bight is very low from 0.1-1 mg m^{-3}.

In order to compare the level of sensitivity between OCI, which have different units,

RMSD for each ocean condition index is normalized by dividing the RMSD by the mean ocean

condition index for both simulations in the pair, producing a RMSD percentage for each

simulation pair. RMSD percentage is calculated for the nearshore retention and larval behavior

simulation pairs with the highest sensitivity for the five OCI. For nearshore retention, the

simulations pair with the highest sensitivity consists of the passive drifting simulation without

nearshore retention and the passive drifting simulation with nearshore retention. For larval

behavior, the simulation pair with the highest sensitivity consists of the passive drifting

simulation without nearshore retention and the DVM20 simulation without nearshore

retention. SZS has the highest RMSD percent for all sites and both simulation pairs, ranging

from 110-210%. LDD has the next highest RMSD percent for the nearshore retention simulation

pair for all sites, ranging from 30-110%, and for the behavior simulations pair for for the SBC

mainland site and the Scripps site, ranging from 41-109%. RMSD percent for Chl is lower than

LDD and SZS for the nearshore retention simulation pair for all sites, ranging from 21-37% and

for the behavior simulations pair for the SBC mainland site and the Scripps site, ranging from

24-51%. Anacapa is the only site that has similar levels sensitivity to LDD and Chl.   For all sites

and both simulation pairs, T_L and T_S have the lowest RMSD percent by 1-2 orders of magnitude

compared SZS, LDD, and Chl, ranging from 2-7% with a mean of 5%. For all OCI, the SBC

mainland sites have similar levels of RMSD percentage, but distinctly different levels of RMSD

percentage from the Anacapa and Scripps sites.

To understand why OCI and recruitment sites show different levels of sensitivity, we

examine the relationship of particle dispersal to circulation, temperature, and surface

chlorophyll. To quantify particle dispersal, mean PDMs for the modeling period are displayed in

Fig. 2 for the Gaviota and Scripps sites for the three simulations used in the nearshore retention

and larval behavior simulation pairs with the highest sensitivity. The three simulations are (1)

passive drifting without nearshore retention (Fig. 2(a,d)), (2) passive drifting with nearshore

retention (Fig. 2(b,e)), and (3) DVM20 without nearshore retention (Fig. 2(c,f)). All simulations

shown have a static PLD. The Gaviota and Scripps sites are selected because their results reflect

the difference in circulation between the SBC and the southern SC Bight.

Circulation (LDD and SZS) and particle dispersal:

When nearshore retention is included in a simulation, the number particles coming from

source sites near the recruitment site automatically increases as simulations with nearshore

retention includes all particle tracks from a source site to a recruitment site that are at or below

the PLD, e.g. PLDs less than or equal to 42 days. In comparison, simulations without nearshore

retention only include particle tracks from a source site to a recruitment site at the specified

PLD, e.g. PLDs equal to 42 days. The increase in particle concentrations near the Goleta and

Scripps sites when nearshore retention is applied to the passive drifting simulation is shown in

Fig. 2(a-b,d-e). As particles with shorter tracking times have a shorter dispersal distance, the

LDD decreases by an average of 47% for all sites when nearshore retention is applied to the

passive drifting simulation. For the same reason, the number particles in the source zones

where recruitment sites are located increases when nearshore retention is applied to the

passive drifting simulation. As a results, the pattern of SZS changes for all sites from being more

evenly spread throughout the 9 source zones (Fig. 2(g,j)) to being dominated by the zone where

the recruitment site is located (Fig. 2(h,k)). In the zones where the recruitment sites are located,

the SZS increases by an average of 44% for all sites.

When DVM is included in a simulation, particle dispersal changes as new particle tracks

are produced. When compared to the passive drifting simulation without nearshore retention,

the DVM20 simulation without nearshore retention decreases particle dispersal for the SBC

mainland sites, which results in a decrease of LDD by an average of 82%. The decrease in

particle dispersal for the Gaviota site can be seen in Fig. 2(a,c). For the SBC mainland sites, long

residence times in the SBC appear to limit particle dispersal for particle tracks with DVM, but

not for particle tracks with passive drifting. This decrease in particle dispersal for DVM20

behavior increases particle concentrations near the SBC mainland sites and SZS by an average of

46% in Zone 1, where the recruitment sites are located. The Anacapa site shows a smaller

increase in particle dispersal and LDD than the SBC mainland sites for the behavior simulation

pair. Compared to the regions near the other recruitment sites, the circulation in the region of

the Anacapa Island, where the Anacapa site is located, has the lowest velocities and shallowest

bathymetry, which may be restricting particle dispersal regardless of larval behavior. In contrast,

DVM20 behavior increases particle dispersal for the Scripps site compared to passive drifting,

which results in an increase in LDD by an average of 20%. The increase in particle dispersal for

DVM20 behavior appears to be driven by the open coastline and short retention times of the

southern SC Bight. For the Scripps site, the increase in particle dispersal for DVM20 versus

passive drifting is shown in Fig. 2(d,f). For the Scripps site, the increase in particle dispersal

from DVM20 decreases the SZS for the Zone 7, where the Scripps site is located, by an average

of 16% and increases the SZS for Zones 6 and 8, which are near Zone 7, by an average of 5% and

6% respectively.

Temperature (T_L and T_S) and particle dispersal:

The OCI representing temperature, T_L and T_S, show the lowest sensitivity to nearshore

retention and larval behavior. T_L is calculated from particle tracks and OCM temperature, and T_S

is calculated from SZS and OCM temperature. Although both nearshore retention and larval

behavior produce significant changes in particle dispersal, temperature is relatively consistent

over the horizontal scale of the changes in particle dispersal. This can be seen by comparing the

areas of high particle concentrations in the mean PDMs (Fig. 2(a-f)) with the mean temperature

for the recruitment and reproductive periods in Fig. 5(b,c)). As a result, RMSD for T_L and T_S for

all recruitment sites is relatively small, ranging from 0.49-0.99°C and 0.31-1.1°C respectively for

the simulation pairs with the highest sensitivity. However, it is important to note that although

the RMSD for T_L and T_S are small relative to their absolute values, a difference of ~1°C could

have be important if the temperature is near a threshold that is detrimental for larval

development or survival.

Food Supply (Chl) and particle dispersal:

The ocean condition index representing food supply, Chl, shows higher sensitivity than T_L

and T_S and lower sensitivity than SZS and LDD for all recruitment sites. Chl is calculated directly

from particle tracks and surface chlorophyll from satellite data. The highest sensitivity for Chl

comes from the SBC mainland sites for the two simulation pairs comparing passive drifting with

and without nearshore retention and passive drifting to DVM20 without nearshore retention. As

the highest Chl in the SC Bight is confined to the SBC (Fig. 5(a)), Chl for SBC mainland sites will

increase when nearshore retention increases particle concentrations in the SBC and when DVM

limits particle dispersal to the SBC region (Fig. 2(c)). In contrast, Chl is much less sensitive to

changes in particle dispersal outside of the SBC, including the southern SC Bight, where Chl is

consistently low.

Summary

The higher sensitivity of LDD and SZS to nearshore retention and larval behavior than T_L,

T_S, and Chl (trend 1) appears to be driven by the fact that circulation and subsequently particle

dispersal in the SC Bight are much more horizontally variable than temperature and surface

chlorophyll. As LDD and SZS are derived directly from particle tracks driven by circulation, they

are more sensitive to the spatial changes in circulation than T_L, T_S, and Chl, which are derived

from particle tracks with temperature or surface chlorophyll.

All OCI show similar levels of sensitivity between the SBC mainland sites and varying

levels of sensitivity between the SBC mainland sites, the Anacapa site, and the Scripps sites

(trend 2). This result appears to be driven by the strong consistent circulation in the SBC. When

nearshore retention or DVM is included in a simulation for the SBC mainland sites, the

circulation creates a region of high particle concentrations in the SBC, which produces similar

changes in OCI for all SBC mainland sites. Due to very different circulation and retention near

the Anacapa and Scripps sites, the addition of nearshore retention or DVM to the simulation for

these sites does not produce changes in OCI consistent with the other recruitment sites.

The high sensitivity of LDD and SZS to nearshore retention and variable sensitivity of LDD

and SZS to larval behavior as a function location within the SC Bight have important implications

for the use of larval connectivity calculated from biophysical models for scientific research as

well as policy decisions for environmental protection (Cowen and Sponaugle 2009; Metaxas and

Saunders 2009; Swearer et al. 2019). These results show that selection of larval life history

parameters can dramatically change the location of predicted larval source sites by 100s of

kilometers. Thus, larval connectivity predictions that are based on a single set of larval life

history parameters could provide misleading information for policy decisions, such as those for

marine protected areas. For this reason, our study shows the importance of conducting

sensitivity studies for larval life history parameters that can't be derived from existing empirical

data.

4.2 Identifying ocean conditions indices that drive of larval recruitment

Circulation, food supply, and temperature are discussed as potential drivers of larval

recruitment of purple sea urchins in the SC Bight. OCI are considered potential drivers of

interannual changes in larval recruitment if a strong linear correlation is found between the

ocean condition index and the larval recruitment index for one or more simulations and all

recruitment sites.

Circulation (LDD and SZS)

The OCI representing circulation, LDD and SZS, displayed a range of correlation strength

as function of larval behavior. The simulations with the strongest correlations for both LDD and

SZS included DVM. For simulations with DVM, LDD is found to be weakly to moderately

inversely correlated to the larval recruitment index for all recruitment sites, implying that larval

dispersal occurs over smaller areas in the average and above-average recruitment years than in

the poor recruitment years. For the zones containing and close to the recruitment sites, SZS for

the DVM20 simulations without nearshore retention shows the strongest linear correlation with

the larval recruitment index, suggesting that a higher percentage of larvae come from these

zones in the average and above-average recruitment years than in the poor recruitment years.

From these results, we hypothesize that circulation in the SC Bight is similar among the average

and above-average recruitment years and among the poor recruitment years, but different

between the average and above-average recruitment years and poor recruitment years.

However, our results suggest that larval recruitment is only connected to circulation when

larvae have behavior, such as DVM, that limits their dispersal to the region near the recruitment

sites. Although this study only compares passive drifting to two simple DVM patterns, urchin

larvae in the SC Bight may performing a different behavior that also reduces larval dispersal in

average and above-average recruitment years.

<u>Food Supply (Chl)</u>

Chl is used as a proxy for larval food supply in the SC Bight because it represents amount

of phytoplankton present, which is the food source for sea urchin larvae. The correlation

between Chl and the recruitment index is very low and inconsistent in both magnitude and sign

across all simulations, implying that Chl is not driving the interannual changes in larval

recruitment. For all sites, the lowest values of Chl occur during 2000, a very high recruitment

year, and 1998, a very low recruitment years (Fig. 3, Table S1), implying that larvae are not food

limited in the SC Bight even in years with lower primary production. It is important to note that

of the five OCI in this study, Chl has the most uncertainty as it is based on surface chlorophyll

and does not account for vertical changes chlorophyll.

<u>Temperature (T_L and T_S)</u>

In prior work, Okamoto et al. (2020) analyzed 27-year larval recruitment datasets of

purple sea urchins in the northern and southern SC Bight and found that recruitment was

negatively correlated with sea surface temperature at the recruitment sites and positively

correlated with the multivariate ENSO index (MEI), which tracks El Niño and La Niña events in

the Pacific Ocean. Throughout the SC Bight, high temperatures and El Niño events were

observed to correspond with low larval recruitment. Okamoto et al. (2020) found that

recruitment was highly synchronous within the northern and southern SC Bight regions,

implying that recruitment was being driven by regional trends in oceanography and not by local

conditions at the recruitment sites.

1077   The OCI that represent temperature are T_L and T_S. Although Okamoto et al. (2020) found

that sea surface temperature was a linearly correlation with larval recruitment, T_L and T_S show

extremely different correlation strengths. T_L is weak and inconsistently correlated with the larval

recruitment index for all sites and simulations, while T_S is strongly correlated with the larval

recruitment index for all sites and simulations. These results imply that T_S is likely driving the

interannual changes in larval recruitment, but T_L is not. To understand why T_L and T_S produce

divergent results, T_L and T_S are compared to documented thresholds for larval survival and

reproduction for the purple sea urchin in the SC Bight.

1085   In the SC Bight, the threshold temperature for lethal conditions of purple sea urchin

larvae increases as temperature approaches 18-20°C (Munstermann et al. In Review). T_L for all

simulations and recruitment sites is below this threshold of ~18°C. For the modeling period of

1996-2013, the highest recorded temperatures in the SC Bight occurred during a El Niño event

that lasted from July 1997 to June 1998 (Lynn and Bograd 2002; McClatchie et al. 2016).

However, during the 1998 recruitment period of March-June, T_L is far below the lethal level with

maximum T_L of 15.9°C for all simulations and recruitment sites. Fig. 8 shows the mean seasonal

temperature cycle at the Gaviota and Scripps recruitment sites using temperatures from the

OCM averaged over the 18-year modeling period. These recruitment sites represent

temperature extremes in SC Bight with Gaviota in the cooler northern region and Scripps in the

warmer southern region (Fig. 5(b,c)). In Fig. 8, the recruitment period of March-June shows

mean temperatures at Gaviota ranging from 12.0-14.3°C and Scripps ranging from 14.4-16.6°C.

The dashed lines from March-June in Fig. 8 represent the temperatures at Gaviota and Scripps

during the 1998 El Niño event over the recruitment period, which range from 12.9-16.6°C and

show that even during a strong El Niño event, the temperatures during the recruitment period

do not vary significantly from the mean annual cycle. Thus, strong El Niño events, although

known for producing high temperatures, appear to have minimal effect on temperatures in the

SC Bight during the recruitment period, remaining well below the lethal limit of 18°C. As

warming produced by marine heatwaves, such as the 1997-1998 El Niño event, does not raise T_L

above the lethal limit for urchin larvae during the recruitment period, we hypothesize that T_L is

most likely not driver of larval recruitment for purple sea urchins in the SC Bight.

Okamoto et al. (Accepted) evaluated the impact of temperature on the reproduction of

purple sea urchins and found that simulating historical heatwave conditions of temperatures

greater than ~18°C suppressed female reproduction. For the simulations with DVM and/or

nearshore retention, T_S for SBC mainland sites is below the threshold of 18°C for average and

above-average recruitment years with a mean and standard deviation per simulation ranging

from 15.5-16.4°C and 1.1-1.3°C. For the same simulations, T_S for SBC mainland sites is warmer

and close to the 18°C threshold for the poor recruitment years with a mean and standard

deviation per simulation ranging from 17.2-17.8°C and 1.2-1.4°C. For Anacapa and Scripps, the

difference in T_S between the average and above-average recruitment years and poor

recruitment years is smaller than the SBC sites, but still shows that the good recruitment years

are colder below threshold of 18°C and the poor recruitment years are warmer near or above

the threshold. For Anacapa, T_S has a mean and standard deviation per simulation ranging from

16.3-16.6°C and 1.3-1.4°C for average and above-average recruitment years and from 17.5-

17.8°C and 1.1-1.3°C for poor recruitment years. For Scripps, T_S has a mean and standard

deviation per simulation ranging from 16.6-17.2°C and 1.0-1.3°C for average and above-average

recruitment years and from 17.7-18.3°C and 1.5-1.6°C for poor recruitment years and. These

results imply that T_S is a likely driver of larval recruitment and that recruitment along the SBC

mainland is more strongly tied to T_S than recruitment near Anacapa Island and the southern SC

Bight mainland.

The SBC mainland sites display the strongest correlation between T_S and the larval

recruitment index for the simulations with DVM and/or nearshore retention. DVM and

nearshore retention both increase the concentrations of particles in Zone 1, where the SBC

mainland sites are located, which consequently increases the SZS for Zone 1 and more heavily

weights T_S by the temperature close to the recruitment sites. This result implies that for the SBC

mainland sites, larval sources in near the recruitment sites are more likely driving recruitment at

these sites than larval sources in zones farther away from the recruitment sites. For the

Anacapa and Scripps sites, the inclusion of DVM and nearshore retention in the simulations did

not produce a higher correlation between T_S and the larval recruitment index, which suggest

that the larval sources in the zone nearest these sites may not be as important in driving larval

recruitment than the SBC mainland sites.

Okamoto et al. (2020) also found that the larval recruitment index correlated strongly

with MEI. To explore this correlation, *r* is calculated for mean T_S and mean MEI over the

reproductive period. Yearly mean T_S and MEI are found have a strong positive linear correlation

with *r* ranging from 0.63-0.82 with a mean of 0.81 for all simulations and sites. These results

suggest that El Niño events, represented by a positive MEI, are correlated with high T_S and poor

recruitment. The strongest El Niño event of the modeling period occurred from July 1997 to

June 1998 and had a mean MEI of 2.1 over this time period. In Fig. 8, the dashed lines from

September to November represent the temperatures at the Gaviota and Scripps sites during the

1997 reproductive period and show that the temperatures in the SC Bight were significantly

higher than the annual average. Fig. 9(a) shows the larval recruitment index as a function of T_S

versus MEI for the simulation with DVM20, no nearshore retention, and a static PLD for all

recruitment sites. The data in Fig. 9(a) suggests that T_S greater than ~17.5°C and MEI greater

than ~-0.5 are likely to produce poor larval recruitment. Fig. 9(b) shows the larval recruitment

index as a function of T_S versus SZS for the Zones 1 and 3 for the SBC sites and Zones 6, 7, and 8

for the Scripps site for the simulation with DVM20, no nearshore retention, and a static PLD.

Fig. 9(b) suggests that in average recruitment year, T_S is less than ~17°C and the SZS for the

zones near the recruitment site is greater than ~60%. In Fig. 9(b), 1998 for the SBC mainland

sites is an outlier as it is a poor recruitment year, but shows high SZS for the Zones 1 and 3. SZS

for the 1998 recruitment period correlates with T_S for 1997 reproductive period. T_S in 1997 is

the highest of the 18-yr modeling period, ranging from 17.9-18.2°C, for the SBC mainland sites

and the only year where monthly T_S does not decrease over the reproductive period (Fig. 3(g)).

Thus for 1998, we hypothesize that although the circulation in the SBC correlates with average

to above-average recruitment, the high temperatures during the 1997 reproductive period

reduced larval supply in 1998 to the extent that circulation is no longer a factor in recruitment.

<u>Summary</u>

T_S is hypothesized to be a primary driver of the interannual variation in larval

recruitment of purple sea urchins in the SC Bight. Our results suggest that mean T_S over the

three-month reproductive period greater than 17-18°C produces poor recruitment by

suppressing reproduction in adult sea urchins at the source sites, which is supported by

experimental data (Okamoto et al. Accepted). Circulation, through LDD and SZS, is found to be a

potential driver of larval recruitment if the larvae have behavior that allows them to remain 35-

45 km from a recruitment site inside the SBC and 80-100 km from a recruitment site in the

central and southern SC Bight. As T_S for the SBC mainland sites had a stronger correlation with

larval recruitment for the simulations that included larval life history parameters that limited

dispersal distance, we hypothesize that urchin larvae in the SBC are likely behaving in a way

which reduces their dispersal distance and consequently that urchin larvae are more likely to

come from sources near the recruitment sites. Many prior studies attribute the settlement

success to larvae coming from areas near their spawning sites due to migration and swimming

behavior, even in strong upwelling systems (Tapia and Pineda 2007; Morgan 2014; Morgan et al.

2018; James et al. 2023; Meyer et al. 2024).

We found that T_S during the reproductive period of September to November is likely an

important driver of larval recruitment, but T_L during the recruitment period of March to June is

not, even in years with marine heat waves and strong El Niño events. These results suggest that

the timing of species recruitment and reproduction is a critical factor in predicting the impact of

temperature on larval recruitment. We hypothesize that marine heatwaves which occur during

time of year when temperatures are at their annual maximum are more likely to produce

temperatures high enough to negatively impact recruitment and reproduction than those which

occur during the time of year when temperatures are at their annual minimum. Species that

recruit or reproduce during the time of year when the system temperatures are at their annual

maximum are more susceptible to marine heat waves than those that recruit or reproduce at

other times of the year. As the frequency of marine heatwaves increases (Sydeman et al. 2014;

Di Lorenzo and Mantua 2016; Bograd et al. 2023), understanding the relationship between the

timing of marine heatwaves and species recruitment and reproduction becomes even more

important.

1190

1191   **Acknowledgements**

1192   This research was supported by the National Science Foundation Biological Oceanography

1193   Program (OCE-2023693 and OCE-2023649).

1194

References

Bandara, K., Ø. Varpe, L. Wijewardene, V. Tverberg, and K. Eiane. 2021. Two hundred years of

zooplankton vertical migration research. Biol. Rev. **96**: 1547–1589.

doi:10.1111/brv.12715

Bell, T. W., K. C. Cavanaugh, D. C. Reed, and D. A. Siegel. 2015. Geographical variability in the

controls of giant kelp biomass dynamics. J. Biogeogr. **42**: 2010–2021.

doi:10.1111/jbi.12550

Bograd, S. J. and others. 2023. Climate change impacts on Eastern Boundary Upwelling Systems.

Ann Rev Mar Sci **15**: 303–328. doi:10.1146/annurev-marine-032122-021945

Botsford, L. W., S. R. Wing, and J. L. Largier. 1998. Population dynamics and management

implications of larval dispersal. South Afr. J. Mar. Sci. **19**: 131–142.

doi:10.2989/025776198784126845

Brokaw, R. J., D. A. Siegel, and L. Washburn. 2024. Physical Drivers of Surface Water Retention in

the Santa Barbara Channel. J. Geophys. Res. Oceans **129**: e2024JC021246.

doi:10.1029/2024JC021246

Brzezinski, M. A., and L. Washburn. 2011. Phytoplankton primary productivity in the Santa

Barbara Channel: Effects of wind-driven upwelling and mesoscale eddies. J. Geophys.

Res. **116**: C12013. doi:10.1029/2011JC007397

Burt, J. M., M. T. Tinker, D. K. Okamoto, K. W. Demes, K. Holmes, and A. K. Salomon. 2018.

Sudden collapse of a mesopredator reveals its complementary role in mediating rocky

reef regime shifts. Proc. R. Soc. B Biol. Sci. **285**: 20180553. doi:10.1098/rspb.2018.0553

Cameron, R., and S. Schroeter. 1980. Sea Urchin Recruitment: Effect of Substrate Selection on

Juvenile Distribution. Mar. Ecol. Prog. Ser. **2**: 243–247. doi:10.3354/meps002243

Cohen, J. H., and R. B. Forward. 2009. Zooplankton diel vertical migration — a review of

proximate control, p. 77–110. *In* J.D.M. Gordon, R.J.A. Atkinson, and R.N. Gibson [eds.],

Oceanography and marine biology: an annual review: Volume 47. Taylor & Francis.

Cowen, R. K., and S. Sponaugle. 2009. Larval dispersal and marine population connectivity. Annu

Rev Mar Sci **1**: 443–466. doi:10.1146/annurev.marine.010908.163757

Di Lorenzo, E., and N. Mantua. 2016. Multi-year persistence of the 2014/15 North Pacific marine

heatwave. Nat. Clim. Change **6**: 1042–1047. doi:10.1038/nclimate3082

Doll, P. C., C. F. Caballes, A. S. Hoey, S. Uthicke, S. D. Ling, and M. S. Pratchett. 2022. Larval

Settlement in Echinoderms: A Review of Processes and Patterns, p. 433–494. *In*

Oceanography and Marine Biology: An Annual Review, Volume 60. CRC Press.

Doney, S. C. and others. 2012. Climate Change Impacts on Marine Ecosystems. Annu. Rev. Mar.

Sci. **4**: 11–37. doi:10.1146/annurev-marine-041911-111611

Dong, C. M. and others. 2012. Three-dimensional oceanic eddy analysis in the Southern

California Bight from a numerical product. J. Geophys. Res.-Oceans **117**.

doi:10.1029/2011jc007354

Dong, C. M., E. Y. Idica, and J. C. McWilliams. 2009. Circulation and multiple-scale variability in

the Southern California Bight. Prog. Oceanogr. **82**: 168–190.

doi:10.1016/j.pocean.2009.07.005

Dong, C. M., and J. C. McWilliams. 2007. A numerical study of island wakes in the Southern

California Bight. Cont. Shelf Res. **27**: 1233–1248. doi:10.1016/j.csr.2007.01.016

Dong, C. M., J. C. McWilliams, A. Hall, and M. Hughes. 2011. Numerical simulation of a synoptic

event in the Southern California Bight. J. Geophys. Res.-Oceans **116**: C05018.

doi:10.1029/2010jc006578

Estes, J. A., and D. O. Duggins. 1995. Sea Otters and Kelp Forests in Alaska: Generality and

Variation in a Community Ecological Paradigm. Ecol. Monogr. **65**: 75–100.

doi:10.2307/2937159

Filbee-Dexter, K., and R. Scheibling. 2014. Sea urchin barrens as alternative stable states of

collapsed kelp ecosystems. Mar. Ecol. Prog. Ser. **495**: 1–25. doi:10.3354/meps10573

Hagerty, M., N. Reyns, and J. Pineda. 2018. Constrained nearshore larval distributions and

thermal stratification. Mar. Ecol. Prog. Ser. **595**: 105–122. doi:10.3354/meps12561

Henderikx Freitas, F., D. A. Siegel, S. Maritorena, and E. Fields. 2017. Satellite assessment of

particulate matter and phytoplankton variations in the Santa Barbara Channel and its

surrounding waters: Role of surface waves. J. Geophys. Res. Oceans **122**: 355–371.

doi:10.1002/2016JC012152

Hickey, B. M. 1993. Chapter 2. Physical Oceanography, p. 19–70. *In* M.D. Dailey, D.J. Reish, and

J.W. Anderson [eds.], Ecology of the Southern California Bight. University of California

Press.

James, M. K., J. A. Polton, C. G. Mayorga-Adame, K. L. Howell, and A. M. Knights. 2023. Assessing

the influence of behavioural parameterisation on the dispersal of larvae in marine

systems. Ecol. Model. **476**: 110252. doi:10.1016/j.ecolmodel.2022.110252

Kahru, M., R. M. Kudela, C. R. Anderson, and B. G. Mitchell. 2015. Optimized merger of ocean

chlorophyll algorithms of MODIS-Aqua and VIIRS. IEEE Geosci. Remote Sens. Lett. **12**:

2282–2285. doi:10.1109/LGRS.2015.2470250

Kahru, M., R. M. Kudela, M. Manzano-Sarabia, and B. Greg Mitchell. 2012. Trends in the surface

chlorophyll of the California Current: Merging data from multiple ocean color satellites.

Deep Sea Res Part II Top Stud Ocean. **77–80**: 89–98. doi:10.1016/j.dsr2.2012.04.007

Krause, J. W., M. A. Brzezinski, D. A. Siegel, and R. C. Thunell. 2013. Biogenic silica standing stock

and export in the Santa Barbara Channel ecosystem. J. Geophys. Res. Oceans **118**: 736–

749. doi:10.1029/2012JC008070

Lafferty, K. D. 2004. Fishing for lobsters indirectly increases epidemics in sea urchins. Ecol. Appl.

14: 1566–1573. doi:10.1890/03-5088

Lynn, R. J., and S. J. Bograd. 2002. Dynamic evolution of the 1997–1999 El Niño–La Niña cycle in

the southern California Current System. Prog. Oceanogr. **54**: 59–75. doi:10.1016/S0079-

6611(02)00043-5

Lynn, R. J., and J. J. Simpson. 1987. The California Current system: The seasonal variability of its

physical characteristics. J. Geophys. Res. **92**: 12947. doi:10.1029/JC092iC12p12947

Mann, K. H., and P. A. Breen. 1972. The Relation Between Lobster Abundance, Sea Urchins, and

Kelp Beds. J. Fish. Res. Board Can. **29**: 603–605. doi:10.1139/f72-104

McClatchie, S., M. G. Jacox, M. D. Ohman, L. M. Sala, and R. Goericke. 2016. State of the

California Current 2015–16: Comparisons with the 1997–98 El Nino. Calif. Coop. Fish.

Invest. Rep. **57**: 5–61.

Metaxas, A. 2020. Larval ecology of echinoids, p. 77–93. *In* Developments in Aquaculture and

Fisheries Science. Elsevier.

Metaxas, A., and M. Saunders. 2009. Quantifying the "Bio-" components in biophysical models

of larval transport in marine benthic invertebrates: advances and pitfalls. Biol. Bull. **216**:

257–272. doi:10.2307/25548159

Meyer, A. D., A. Hastings, and J. L. Largier. 2024. Making Your Own Luck: Weak Vertical

Swimming Improves Dispersal Success for Coastal Marine Larvae. Bull. Math. Biol. **86**:

23. doi:10.1007/s11538-023-01252-2

Miller, B. A., and R. B. Emlet. 1999. Development of newly metamorphosed juvenile sea urchins

(Strongylocentrotus franciscanus and S. purpuratus): morphology, the effects of

temperature and larval food ration, and a method for determining age. J. Exp. Mar. Biol.

Ecol. **235**: 67–90. doi:10.1016/S0022-0981(98)00164-6

Mitarai, S., D. A. Siegel, J. R. Watson, C. Dong, and J. C. McWilliams. 2009. Quantifying

connectivity in the coastal ocean with application to the Southern California Bight. J.

Geophys. Res.-Oceans **114**: C10026. doi:10.1029/2008JC005166

Morgan, S. G. 2001. The larval ecology of marine communities, p. 159–181. *In* M.D. Bertness,

S.D. Gaines, and M.E. Hay [eds.], Marine community ecology. Sinauer Associates.

Morgan, S. G. 2014. Behaviorally Mediated Larval Transport in Upwelling Systems. Adv.

Oceanogr. **2014**: 1–17. doi:10.1155/2014/364214

Morgan, S. G., J. L. Fisher, S. H. Miller, S. T. McAfee, and J. L. Largier. 2009. Nearshore larval

retention in a region of strong upwelling and recruitment limitation. Ecology **90**: 3489–

3502. doi:10.1890/08-1550.1

Morgan, S. G., S. H. Miller, M. J. Robart, and J. L. Largier. 2018. Nearshore Larval Retention and

Cross-Shelf Migration of Benthic Crustaceans at an Upwelling Center. Front. Mar. Sci. **5**:

161. doi:10.3389/fmars.2018.00161

Munstermann, M., S. Karelitz, R. Ferraro, L. Rogers-Bennett, R. Simons, and D. Okamoto. In

Review. Food limitation alters the thermal response of survival but not rates of

development in larvae of the purple sea urchin Strongylocentrotus purpuratus. Ecology.

doi:10.22541/au.173469634.44386797/v1

Myers, R. A. 1998. When do environment–recruitment correlations work? Rev. Fish Biol. Fish. **8**:

285–305. doi:10.1023/A:1008828730759

Nishimoto, M. M., R. D. Simons, and M. S. Love. 2019. Offshore oil production platforms as

potential sources of larvae to coastal shelf regions off southern California. Bull. Mar. Sci.

95: 535–558. doi:10.5343/bms.2019.0033

Oey, L.-Y., C. Winant, E. Dever, W. R. Johnson, and D.-P. Wang. 2004. A Model of the Near-Surface

Circulation of the Santa Barbara Channel: Comparison with Observations and Dynamical

Interpretations. J. Phys. Oceanogr. **34**: 23–43. doi:10.1175/1520-

0485(2004)034%3C0023:AMOTNC%3E2.0.CO;2

Ohlmann, J. C., and S. Mitarai. 2010. Lagrangian assessment of simulated surface current

dispersion in the coastal ocean. Geophys. Res. Lett. **37**: L17602.

doi:10.1029/2010GL044436

Okamoto, D. K., S. C. Schroeter, and D. C. Reed. 2020. Effects of ocean climate on spatiotemporal

variation in sea urchin settlement and recruitment. Limnol. Oceanogr. **65**: 2076–2091.

doi:10.1002/lno.11440

Okamoto, D. K., N. B. Spindel, B. Collicutt, M. Munstermann, S. Karelitz, and Gimenez. Accepted.

Thermal suppression of gametogenesis explains historical collapses in larval recruitment.

Commun. Biol. doi:https://doi.org/10.1101/2023.09.28.559919

Otero, M., and D. Siegel. 2004. Spatial and temporal characteristics of sediment plumes and

phytoplankton blooms in the Santa Barbara Channel. Deep Sea Res. Part II Top. Stud.

Oceanogr. **51**: 1129–1149. doi:10.1016/S0967-0645(04)00104-3

Page, H. 2019. Distribution and potential larval connectivity of the non-native Watersipora

(Bryozoa) among harbors, offshore oil platforms, and natural reefs. Aquat. Invasions **14**:

615–637. doi:10.3391/ai.2019.14.4.04

Paris, C., L. Chérubin, and R. Cowen. 2007. Surfing, spinning, or diving from reef to reef: effects

on population connectivity. Mar. Ecol. Prog. Ser. **347**: 285–300. doi:10.3354/meps06985

Pineda, J., J. Hare, and S. Sponaugle. 2007. Larval transport and dispersal in the coastal ocean

and consequences for population connectivity. Oceanogr **20**: 22–39.

doi:10.5670/oceanog.2007.27

Pineda, J., F. Porri, V. Starczak, and J. Blythe. 2010. Causes of decoupling between larval supply

and settlement and consequences for understanding recruitment and population

connectivity. J. Exp. Mar. Biol. Ecol. **392**: 9–21. doi:10.1016/j.jembe.2010.04.008

Rogers-Bennett, L., and C. A. Catton. 2019. Marine heat wave and multiple stressors tip bull kelp

forest to sea urchin barrens. Sci. Rep. **9**: 15050. doi:10.1038/s41598-019-51114-y

Rogers-Bennett, L., and D. Okamoto. 2020. Mesocentrotus franciscanus and Strongylocentrotus

purpuratus, p. 593–608. *In* Developments in Aquaculture and Fisheries Science. Elsevier.

Schroeter, S. C., J. D. Dixon, T. Ebert, and J. Richards. 2017. SBC LTER: Time series of settlement

of urchins and other

invertebrates.doi:10.6073/PASTA/A7BA12263D75BE25CE526A2E20B0E6F3

Shanks, A. L. and others. 2020. Marine heat waves, climate change, and failed spawning by

coastal invertebrates. Limnol. Oceanogr. **65**: 627–636. doi:10.1002/lno.11331

Simons, R. D., and D. Catlett. 2023. Regulation of a surface chlorophyll hotspot by wind-driven

upwelling and eddy circulation in the Santa Barbara Channel, Southern California. Prog.

Oceanogr. **217**: 103096. doi:10.1016/j.pocean.2023.103096

Simons, R. D., M. M. Nishimoto, L. Washburn, K. S. Brown, and D. A. Siegel. 2015. Linking

kinematic characteristics and high concentrations of small pelagic fish in a coastal

mesoscale eddy. Deep Sea Res Part Ocean. Res Pap. **100**: 34–47.

doi:10.1016/j.dsr.2015.02.002

Simons, R. D., H. M. Page, S. Zaleski, R. Miller, J. E. Dugan, D. M. Schroeder, and B. Doheny. 2016.

The effects of anthropogenic structures on habitat connectivity and the potential spread

of non-native invertebrate species in the offshore environment M. (Gee) G. Chapman

[ed.]. PLOS ONE **11**: e0152261. doi:10.1371/journal.pone.0152261

Simons, R. D., D. A. Siegel, and K. S. Brown. 2013. Model sensitivity and robustness in the

estimation of larval transport: A study of particle tracking parameters. J. Mar. Syst. **119**:

19–29. doi:10.1016/J.Jmarsys.2013.03.004

Stige, L., M. Hunsicker, K. Bailey, N. Yaragina, and J. Hunt GL. 2013. Predicting fish recruitment

from juvenile abundance and environmental indices. Mar. Ecol. Prog. Ser. **480**: 245–261.

doi:10.3354/meps10246

Strathmann, R. R. 1975. Larval Feeding in Echinoderms. Am. Zool. **15**: 717–730.

doi:10.1093/icb/15.3.717

Swearer, S. E., E. A. Treml, and J. S. Shima. 2019. A Review of Biophysical Models of Marine

Larval Dispersal, p. 325–356. *In* S.J. Hawkins, A.L. Allcock, A.E. Bates, L.B. Firth, I.P. Smith,

S.E. Swearer, and P.A. Todd [eds.], Oceanography and Marine Biology. CRC Press.

Sydeman, W. J., M. García-Reyes, D. S. Schoeman, R. R. Rykaczewski, S. A. Thompson, B. A. Black,

and S. J. Bograd. 2014. Climate change and wind intensification in coastal upwelling

ecosystems. Science **345**: 77–80. doi:10.1126/science.1251635

Tapia, F., and J. Pineda. 2007. Stage-specific distribution of barnacle larvae in nearshore waters:

potential for limited dispersal and high mortality rates. Mar. Ecol. Prog. Ser. **342**: 177–

190. doi:10.3354/meps342177

Tegner, M., and P. Dayton. 1991. Sea urchins, El Ninos, and the long term stability of Southern

California kelp forest communities. Mar. Ecol. Prog. Ser. **77**: 49–63.

doi:10.3354/meps077049

Tegner, M. J. 2001. The ecology of Strongylocentrotus franciscanus and Strongylocentrotus

purpuratus, p. 307–331. *In* Developments in Aquaculture and Fisheries Science. Elsevier.

Underwood, A. J., and M. J. Keough. 2001. Supply-side ecology: the nature and consequences of

variations in recruitment of intertidal organisms, p. 183–200. *In* M.D. Bertness, S.D.

Gaines, and M.E. Hay [eds.], Marine community ecology. Sinauer Associates.

White, J. W. and others. 2019. Connectivity, Dispersal, and Recruitment: Connecting Benthic

Communities and the Coastal Ocean. Oceanography **32**: 50–59.

doi:10.5670/oceanog.2019.310

Winant, C. D., E. P. Dever, and M. C. Hendershott. 2003. Characteristic patterns of shelf

circulation at the boundary between central and southern California. J. Geophys. Res.

Oceans **108**: 3021. doi:10.1029/2001JC001302

Yamhure, G. M., N. Reyns, and J. Pineda. 2021. High Larval Concentrations and Onshore

Transport of Barnacle Cyprids Associated With Thermal Stratification. Front. Mar. Sci. **8**:

748389. doi:10.3389/fmars.2021.748389

1394

1395

Figures

1397

1398
Figure 1: Southern California Bight: study area and model domain. Black diamonds are the 5
larval recruitment sites: Gaviota, Ellwood, Stearns, Anacapa, and Scripps. Red and yellow
coastline locations are the nine source zones, which are identified by the black numbers: (1)
mainland of the Santa Barbara Channel, (2) north side of the North Channel Islands, (3) south
side of the North Channel Islands, (4-7) groups of contiguous source sites along the central and
southern mainland of the Southern California Bight, (8) Catalina Island, and (9) three South
Channel Islands

1406
1407
1408

1409
Figure 2: (A-F) Mean particle density maps (PDMs) and (G-L) mean source zone strength (SZS)
for three following simulations with static PLDs: (1) passive drifting without nearshore retention
for Gaviota (A,G) and for Scripps (D,J), (2) passive drifting with nearshore retention for Gaviota
(b,h) and Scripps (E,K), and (3) DVM20 without nearshore retention for Gaviota (C,F) and for
Scripps (F,L). PDMs and SZS have been averaged over all months of the modeling period (1996-
2013). PDMs are plotted on a log scale.
1416

1417

Figure 3: Ocean conditions for three simulations: (1) passive drifting behavior, no nearshore
retention, and static PLD (yellow line), (2) passive drifting behavior, nearshore retention, and
static PLD (blue line), and (3) DVM20 behavior, nearshore retention, and static PLD (red line).
(A) LDD (km) for Gaviota, (B) LDD (km) for Scripps, (C) Chl (mg m^{-3}) for Gaviota, (D) Chl (mg m^{-3})
for Scripps, (E) T_L (°C) for Gaviota, (F) T_L (°C) for Scripps, (G) T_S (°C) for Gaviota, and (H) T_S (°C) for
Scripps. LDD, Chl, and T_L are shown for the larval recruitment period of March-June. T_S is shown
for the reproductive period of September-November.

1425

1426

1427

1428

1429

1430

1431
Figure 4: SZS for three simulations, (S1) passive drifting behavior, no nearshore retention, and
static PLD, (S2) passive drifting behavior, nearshore retention, and static PLD, and (S3) DVM20
behavior, nearshore retention, and static PLD: (A) S1 for Gaviota, (B) S2 for Gaviota, (C) S3 for
Gaviota, (D) S1 for Scripps, (E), S2 for Scripps, and (F) S3 for Scripps. SZS is shown for the
recruitment period of March-June. Units for SZS is in percentage of particles. Zones 1-9 are
shown in Figure 1.

Figure 5: (A) Mean surface chlorophyll-a (mg m^{-3}) from satellite data for the recruitment period,
(B) mean temperature (°C) averaged over top 50 m of water column for the recruitment period,
(C) mean temperature (°C) averaged over top 50 m of water column for the reproductive
period, and (D) mean velocity (cm s^{-1}) averaged over the top 50 m of the water column for the
recruitment period. The solid black circles represent the recruitment sites: (1) Gaviota, (2)
Ellwood, (3) Stearns, (4) Anacapa, and (5) Scripps. All data is averaged over the modeling period
of 1996-2013. The recruitment period is from March to June. The reproductive period is from
September to November.
1453

1454
1455 Figure 6: SZS (%) versus the larval recruitment index for (A) Gaviota (Zones 1 and 3) and (B)
1456 Scripps (Zones 6, 7, and 8). T_S (°C) versus the larval recruitment index for (C) Gaviota and (D)
1457 Scripps. The simulation shown has DVM20 behavior, no nearshore retention, and a static PLD.
1458 Each solid black circle represents one year. The solid black line shows the linear regression for
1459 all years.
1460
1461

1462
1463 Figure 7: Pearson's r versus PLD (days). r is calculated between the particle density map (PDM)
1464 with a PLD of 42 days and the PDMs with PLDs of 1-56 days for all simulations, recruitment sites,
1465 and months of the modeling period. Solid black line is mean r. Gray area is ±1 standard
1466 deviation of r. Dashed black line identifies a static PLD of 42 days.

1467

1468
1469 Figure 8: Mean annual temperature cycle $\pm$ 1 standard deviation (°C) at 5m depth for Gaviota
1470 (red) and Scripps (blue). Dashed lines represent the reproductive period (September-
1471 November) for 1997 and the recruitment period (March-June) for 1998. Temperature is derived
1472 from the ocean circulation model and is averaged over the modeling period of 1996-2013.
1473
1474
1475

1476
1477 Figure 9: Larval recruitment index as a function of (a) T_S (°C) and MEI for all recruitment sites
1478 and (b) T_S (°C) and SZS (%) for Gaviota, Ellwood, Stearns, and Scripps. All data show is for the
1479 simulation with DVM20 behavior, no nearshore retention, and a static PLD. A circle represents
1480 one year. SZS is calculated from Zones 1 and 3 for Gaviota, Ellwood, and Stearns and Zones 6, 7,
1481 and 8 for Scripps.
1482
1483