1 **Main Manuscript for**

3 **Long lasting non-cellular reactions in sterile soils recapitulate most**
4 **of the intermediates of the Krebs cycle**

6 Clémentin Bouquet[1], Mounir Traikia[2,3], Fanny Perrière[1], Benoit Kéraval[1], Gaël Alvarez[4], Loïc Louis[5], Martin
7 Leremboure[2], Guillaume Voyard[2], Sébastien Fontaine[4] and Anne-Catherine Lehours[1*]
8
9
[1]Université Clermont Auvergne, CNRS, LMGE, F-63000 Clermont-Ferrand, France
[2]Université Clermont Auvergne, CNRS, ICCF, F-63000 Clermont-Ferrand, France
[3]Université Clermont Auvergne, PFEM, MetaboHUB Clermont, Clermont-Ferrand, France
[4]Université Clermont Auvergne, INRAE, VetAgro Sup, UMR Ecosystème Prairial, 63000 Clermont-
Ferrand, France
[5]Université de Lorraine, AgroParisTech, INRAE, SILVA, F-54000 Nancy, France
16
17

* Correspondence to: Anne-Catherine Lehours

Email: a-catherine.lehours@uca.fr

Author Contributions: CB, BK and ACL have designed the work. CB, MT, FP, BK, GV, LL contributed
to the experimental data acquisition. All authors contributed to data set analysis and interpretation of
data. CB and ACL drafted the work, and all authors have revised it. ACL, MT and SF supervised the
project.

Competing Interest Statement: The authors declare that they have no conflicts of interest.
25

Classification: Biological Sciences/ Environmental Sciences

Keywords: extracellular metabolisms (EXOMETs), stable carbon isotope, chromatography, mass
spectrometry

This file includes:

Main Text
Figure 1
33

Abstract

Over the past decade, chemical evidence has emerged that non-enzymatic metabolic pathways, such

as the Krebs cycle, may have existed before cellular life on the primitive Earth. However, the question

of whether non-cellular reactions analogous to cell respiration metabolism are still "active" in today's

biosphere and whether they contribute to CO_2 emissions in contemporary ecosystems remains open. In

the present study, we investigated the long-term fate (> 6 months) of organic substrates supplied in

sterilised soils in which cell life was not detectable. Through a series of analytical studies performed on

the water-extractable fraction of soil exometabolites using chromatography, mass spectrometry and

isotope labelling, we demonstrate that sterile soil matrix incubated with $[^{13}C_6]$-glucose and $[^{13}C_6]$-citrate

can spontaneously generate intermediates of the Krebs cycle, alongside by-products such as acetate

and formate. These findings support the hypothesis that extracellular metabolisms (EXOMETs) form a

network of non-cellular reactions resembling metabolic pathways involved in aerobic respiration and

anaerobic fermentation of cells. This research not only provides insights into the chemical continuity

between chemistry and biochemistry but also raises questions about the implications of non-cellular

pathways for soil ecosystem functioning and carbon fluxes in the contemporary biosphere.

Main Text

Introduction

Metabolism, along with replication and information transfer, is one of the most critical features of living

systems. Most of the key reactions in the metabolic network are catalysed by protein enzymes, whose

complex molecular structures are the product of Darwinian evolution. This has contributed to the view,

which is backed by 'genetics-first' theories, that the topologies of metabolic pathways might have

developed through the evolution of enzymes. According to this line of thought, there should be no

environmental chemistry that mimics metabolism **(1)**.

However, chemical evidence has recently emerged for the existence of non-enzymatic metabolic

pathways. One of the first was the serendipitous discovery of glycolysis-like reactions catalysed by the

metal cation Fe^{2+} **(2)**. Then, a series of non-enzymatic reactions resembling those of the tricarboxylic

acid (TCA) cycle, also known as the Krebs cycle, have been repeatedly documented **(3, 4, 5, 6)**,

including the identification of TCA intermediates in a carbonaceous meteorite **(7)**. These compelling

arguments support that the topologies of the metabolic pathways have their roots in non-enzymatic

chemistry **(8)**. From an ecological and biogeochemical perspective, these findings also raise the

questions of whether non-cellular reactions analogous to metabolism are still 'active' in today's

biosphere, and whether they contribute to respiration and, hence, CO_2 emissions in contemporary

ecosystems.

These questions arise from the recurrent observations of substantial CO_2 emissions that persist in soils

where living biomass has been reduced to undetectable levels by exposure to toxics, gamma irradiation,

or autoclaving **(9, 10, 11)**. Various non-cellular processes have been proposed as potential contributors

to CO_2 emissions in sterile soils including single catalytic reactions driven by reactive oxygen species

and/or metals such as iron **(12)** or supported by decarboxylases released during cell lysis **(10, 13)**.

However, these proposals do not explain how $[^{13}C_6]$-glucose is completely oxidised to ^{13}C-CO_2 in sterile

soils **(10, 11, 14)**. A decade ago, we proposed the hypothesis of extracellular metabolisms (EXOMETs),

consisting of multiple coupled redox reactions that reconstitute an equivalent of a cellular respiration

process **(10, 11)**. Recently, we have shown that in sterile soils, electron fluxes operate in conjunction

with a series of time-ordered reactions controlled by the nature of the substrate, which produce and

consume exometabolites and lead to the complete oxidation of organic substrates ($[^{13}C_6]$- glucose and

-citrate) with oxygen as the final electron acceptor. We have also shown that the soil matrix can sustain

the complete oxidation of organic compounds for years, a period much longer than the half-life of
enzymes, revealing the involvement of stable non-enzymatic catalysts in EXOMET (14). However, to
what extent EXOMET activities follow pathways close to central cell metabolism, such as the Krebs
cycle, is not yet known.
In the present study, we show that in soils devoid of cell life, citrate and glucose spontaneously generate
reactions that recapitulate most of the intermediates of the core biological pathways of the Krebs cycle.
In addition, we observe that glucose enhances the synthesis of formate and acetate in sterile soils,
supporting the hypothesis that many additional cellular metabolism-like reactions remain to be
discovered in contemporary soils.

The analyses were carried out on soil sample extracts obtained from the experimental setup of Bouquet
et al. (14). Briefly, soils (pH = 6.2) sterilised by gamma irradiation at 45 kGy and non-sterilised soils were
incubated in the dark at 20 °C and a water potential of -100 kPa for 163 days. Five different substrate
treatments were applied to sterilised soils: no substrate addition (IS for irradiated soil), the addition of
$[^{13}C_6]$-labelled citrate (^{13}C-IS), unlabelled citrate (^{12}C-IS), $[^{13}C_6]$-labelled glucose (^{13}G-IS) or unlabelled
glucose (^{12}G-IS). All treatments were replicated four times (n = 4). The water-extractable fraction of
exometabolites with a molecular weight < 3 kDa (hereafter referred to as WEM) was extracted from
each replicate sample at each sampling time during the incubation period, which involved sampling of
the soils at various intervals (t= 0.2, 3, 6, 17, 23, 66, 100, 163 days). (*SI Appendix*).
First,we analysed WEM using ion chromatography-mass spectrometry (IC-MS) with comparisons
against authentic standards to achieve absolute quantification of seven Krebs cycle intermediates (*SI*
Appendix, **Figure 1A**). In non-sterilised soil samples, none of the target Krebs cycle intermediates were
detected. By contrast, all of the Krebs cycle target intermediates (citrate, cis-aconitate, α-ketoglutarate,
succinate, fumarate, malate and oxaloacetate, **Figure 1A**) were detected in the sterilised soils, with the
exception of the citrate which was below the detection threshold at all sampling times for the IS, ^{12}G-IS
and ^{13}G-IS samples. Although the α-ketoglutarate was detected in some samples, its abundance was
systematically below the limit of quantification (*SI Appendix*). The temporal dynamics of the
abundances of the remaining five Krebs cycle intermediates were monitored (**Figure 1A**). It should be
noted that ^{13}C-labelling did not impact these dynamics since the abundance of the targeted metabolites
at each sampling time did not differ significantly between the ^{13}C-IS and ^{12}C-IS treatments, or the ^{13}G-
IS and ^{12}G-IS treatments. Therefore, to improve statistical robustness, we computed concentrations for
each sampling date using the pooled data for each substrate treatment (**Figure 1**).
During incubation, the production of Krebs metabolites was significantly stimulated ($p < 0.01$) in sterilised
soils that had been amended with citrate, compared to soils that had not been amended and soils that
had been amended with glucose. (**Figure 1B**). Although these Krebs compounds exhibited distinct
temporal dynamics, they accumulated much more in citrate-amended soils than in the other treatments
(**Figure 1C**). For instance, succinate and malate accumulated significantly more ($p < 0.01$) in citrate-
amended soils (1.6 ± 0.1 µg.g^{-1} and 1.3 ± 0.1 µg.g^{-1}, respectively) than in unamended soils (0.8 ± 0.2
µg.g^{-1} and 0.2 ± 0.05 µg.g^{-1}, respectively) and in glucose-amended soils (0.7 ± 0.1 µg.g^{-1} and 0.2 ± 0.02
µg.g^{-1}, respectively) (**Figure 1C**). In contrast, fumarate accumulation was much less pronounced with
small differences between treatments (**Figure 1C**). Together with the fact that α-ketoglutarate is below

the quantification threshold (*SI Appendix*), these observations suggest a possible shunting of metabolic
pathways in sterilised soils with respect to the classical cellular Krebs cycle. Future studies should
explore whether non-cellular reactions in sterilised soils also replicate the glyoxylate shunt, a two-step
pathway generally regarded as an auxiliary pathway of the Krebs cycle (**Figure 1A**). This is an intriguing
prospect, given that Keller et al. [4] have demonstrated the ability of non-enzymatic reactions to
reproduce the glyoxylate shunt in experiments simulating the archaean oceans under conditions (e.g.
heating to 70°C) that differ greatly from those of the present study [4, 8].
In addition to being the most abundant, oxaloacetate is the metabolite that accumulates the most. By
the end of the incubation period, its concentration had reached 49.6 ± 3.1 $\mu g.g^{-1}$ in citrate-amended
sterilised soils (**Figure 1C**). This is more than 2.5 times higher than in sterilised soils or in glucose-
amended sterilised soils and between 25 and 130 times higher than the other Krebs metabolites (**Figure
1C**). In the enzyme-catalysed Krebs cycle occurring in cells, oxaloacetate - the end product of a previous
round of the cycle - forms citrate by condensing with acetyl-CoA (**Figure 1A**). To date, this step has not
been reproduced in non-enzymatic experiments [4, 8]. Therefore, the accumulation of oxaloacetate,
combined with the lack of citrate detection in unamended and glucose-amended sterilised soils,
suggests that oxaloacetate production may represent a dead end for EXOMETs.
As the addition of glucose to sterilised soils did not stimulate the production of Krebs cycle intermediates
(**Figure 1B**), we examined the dynamics of acetate and formate, which are considered classic markers
of fermentative reactions. The abundances of these two compounds were significantly ($p < 0.05$)
stimulated by the addition of glucose, particularly during the last incubation period (> 66 days, **Figure
1D**). After 163 days, the glucose-amended soils had accumulated 198 % and 173 % more acetate and
256 % and 192 % more formate than the unamended and citrate-amended sterilised soils (**Figure 1D**).
These results suggest that EXOMETs may be engaged in additional non-cellular metabolic processes
besides aerobic respiration, such as fermentation.

Secondly, to verify that the accumulation of Krebs cycle intermediates was at least partly due to the
conversion of labelled or unlabelled glucose and citrate in sterilised soils, the $\delta^{13}C$ isotope ratios of
metabolites in the sterilised soils were measured using gas chromatography-isotope ratio mass
spectrometry (GC-C-IRMS) (*SI Appendix*). Each sample was also concurrently analysed using gas
chromatography-mass spectrometry (GC-MS) to identify and determine the retention times of the target
compounds (*SI Appendix*). Prior to analysis, the samples were derivatised, resulting in the loss of
malate during the procedure. In citrate-amended soils, the $\delta^{13}C$ values of cis-aconitate, succinate,
fumarate and oxaloacetate differed significantly ($p < 0.01$) between ^{13}C-IS and ^{12}C-IS. For cis-aconitate,
the average $\delta^{13}C$ values were 1752.2 ± 283.2 and $- 17.5 \pm 4.6$ in ^{13}C-IS and ^{12}C-IS, respectively (**Figure
1E**). Due to the low amounts of Krebs intermediates observed in glucose-amended soils, it was not
always possible to detect these intermediates by GC-C-IRMS, depending on the metabolite and day of
incubation (**Figure 1E**). However, the $\delta^{13}C$ values of the Krebs intermediates in each measured sample
were significantly higher ($p < 0.001$) in ^{13}G-IS than in ^{12}G-IS showing that glucose also contributes to the
formation of succinate and oxaloacetate (**Figure 1E**). This suggests that non-cellular reactions in soils
can trigger cascades of reactions capable of replicating all or part of glycolysis.

Conclusions

Until far, research on non-cellular processes in soils has focused on CO_2 emissions from sterilised soils (e.g. **9, 10, 11, 14, 15**). We open a "black box" by providing the first evidence that citrate and glucose in sterile soils trigger non-cellular convergent reactions that lead to the formation of several Krebs cycle intermediates, as well as additional fermentation products such as acetate and formate. These observations extend those of Bouquet et al. **(14)** who demonstrated a series of time-ordered reactions controlled by the nature of the substrate in sterile soils. Together, these results challenge the dogma of the cell as the minimal unit for carrying out metabolic reactions by showing that, under mild conditions (20°C, pH 6.2) in soils, a network of non-cellular reactions recapitulates most of the intermediates of the Krebs cycle.

Strengthening the idea of a real chemical continuity between prebiotic chemistry and biochemistry **(8)**, our observations bring up a number of issues not only regarding the diversity of these non-cellular metabolic networks (e.g. respiration, glycolysis, fermentation) but also their fundamental architecture (e.g. connection between metabolic intermediates, possible shunting of pathways). This will be pursued using specific positional labelling to track isotopomers and isotopologues as well as fluxomic approaches to study the kinetics and potential cyclisation of these processes.

From a biogeochemical perspective, while previous estimates suggest that EXOMETs could contribute significantly to CO_2 emissions in contemporary soils **(10, 15)**, our study indicates that they could also contribute to CO_2 emissions in anaerobic environments such as marine or freshwater sediments, through non-cellular fermentation.

Materials and Methods

Supplementary methods appear in *SI Appendix*.

Data Availability

GC-MS and GC-C-IRMS data are available in Figshare: https://figshare.com/s/5470d388bbc43f84ff69

Acknowledgments

Clémentin Bouquet was supported by PhD fellowship from the French Ministry of Education and Research. The authors thank the financial support of (i) the CNRS through the Mission pour les Initiatives transverses et interdisciplinaires (MITI) and through Projet Exploratoire Premier Soutien (PEPS) and the funding of the ISO-EXOMET and EXCEED projects, (ii) CAP 20-25 Isite of the University Clermont Auvergne and the funding of the EXOMET project. The author(s) would like to thank SILVATECH (Silvatech, INRAE, 2018. Structural and functional analysis of tree and wood Facility, doi: 10.15454/1.5572400113627854E12) from UMR 1434 SILVA, 1136 IAM, 1138 BEF and 4370 EA LERMAB from the research center INRAE Grand-Est Nancy for it contribution to isotopic analysis and methodological development. SILVATECH facility is supported by the French National Research Agency through the Laboratory of Excellence ARBRE (ANR-11-LABX-0002-01)

References

1. L. Orgel, Origin of life. A simpler nucleic acid. *Science.* 290,1306-1307 (2000).

2. M. A. Keller, A. V. Turchyn, M. Ralser. Non-enzymatic glycolysis and pentose phosphate pathway-like reactions in a plausible Archean ocean. *Mol. Syst. Biol.* 10, 725 (2014).

3. D. Zubarev, D. Rappoport, A. Aspuru-Guzik. Uncertainty of prebiotic scenarios: the case of the non-enzymatic reverse tricarboxylic acid cycle. *Sci. Rep.* 5, 8009 (2015).

4. M. A. Keller, D. Kampjut, S. A. Harrison, M. Ralser M. Sulfate radicals enable a non-enzymatic Krebs cycle precursor. *Nat. Ecol. Evol.* 1, 83 (2017).

5. K. B. Muchowska, S. J. Varma, J. Moran. Synthesis and breakdown of universal metabolic precursors promoted by iron. *Nature.* 569,104-107 (2019).

6. R. T. Stubbs, M. Yadav, R. Krishnamurthy, *et al.* A plausible metal-free ancestral analogue of the Krebs cycle composed entirely of α-ketoacids. *Nat. Chem.* 12, 1016-1022 (2020).

7. G. Cooper, N. Kimmich, W. Belisle, *et al.* Carbonaceous meteorites as a source of sugar-related organic compounds for the early Earth. *Nature* 414, 879-883 (2001).

8. M. Ralser. An appeal to magic? The discovery of a non-enzymatic metabolism and its role in the origins of life. *Biochem. J.* 475, 2577-2592 (2018).

9. S. J. Kemmitt, C. V. Lanyon, I. S. Waite, et al. Mineralization of native soil organic matter is not regulated by the size, activity or composition of the soil microbial biomass-a new perspective. *Soil Biol. Biochem.* 40, 61-73 (2008).

10. V. Maire, G. Alvarez, J. Colombet, et al. An unknown oxidative metabolism substantially contributes to soil CO_2 emissions. *Biogeosciences* 10, 1155-1167 (2013).

11. B. Kéraval, A.-C. Lehours, J. Colombet, et al. Soil carbon dioxide emissions controlled by an extracellular oxidative metabolism identifiable by its isotope signature. *Biogeosciences* 13, 6353-6362 (2016).

12. G.-H Yu, Y. Kuzyakov. Fenton chemistry and reactive oxygen species in soil: abiotic mechanisms of biotic processes, controls and consequences for carbon and nutrient cycling. *Earth-Sci. Rev.* 214, 103525 (2021).

13. J. C. Blankinship, C. A. Becerra, S. M. Schaeffer, J. P. Schimel. Separating cellular metabolism from exoenzyme activity in soil organic matter decomposition. *Soil Biol. Biochem.* 71, 68–75 (2014).

14. C. Bouquet, B. Kéraval, M. Traikia, et al. Non-living respiration: another breath in the soil? *bioRxiv* 2025.07.03.662961(2025).

15. B. Kéraval, S. Fontaine, A. Lallemant, S. Revaillot, H. Billard, G. Alvarez, F. Maestre, C. Amblard, A.-C. Lehours. Cellular and non-cellular mineralization of organic carbon in soils with contrasted physicochemical properties. *Soil Biol. Biochem.* 125, 286–289 (2018).

Legends of Figures

Figure 1. Dynamics of the concentration and carbon isotopic composition of the Krebs cycle

intermediates as well as formate and acetate concentrations in sterile soils.

(A)   Schematic representation of the cellular enzyme-catalysed Krebs cycle, as it occurs in yeast,

and the glyoxylate shunt (modified from **(8)**), as well as two products (acetate and formate) that can be

produced by the fermentation of glucose. The inner (black) and outer (red) circles of the diagram circle

indicate the compounds analysed by ionic chromatography coupled to mass spectrometry (IC-MS) and

gas chromatography-isotope ratio mass spectrometry (GC-C-IRMS), respectively. Glucose and citrate,

coloured pink, were used as substrates in different sterilised soil treatments. **(B)** Quantitative stimulation

of Krebs cycle intermediates by citrate or glucose amendment compared to unamended soils. The colour

scale represents the results of the two-tailed paired p-value (n=8) for all metabolites, except for cis-

aconitic acid for which a one-sample t-test was used (n=8). Asterisks denote p-values as follows: ns: p

≥ 0.05; *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$. **(C)** Temporal dynamics of the concentrations, analysed by

IC-MS, over 163 days of incubation of (from the top to the bottom of the diagram): Cis-aconitate,

succinate, fumarate, malate, oxaloacetate in unamended sterilised soils (in red), in glucose-amended

sterilised soils (in purple) and in citrate-amended soils (in orange). **(D)** Temporal dynamics of carbon

stable isotope ratios (δ^{13}C) analysed by GC-C-IRMS of cis-aconitate, succinate, fumarate and

oxaloacetate in sterilised soils amended with labelled (solid circles) or unlabelled (open circles) glucose

or citrate. For **(C)**, **(D)** and **(E)**, analyses were performed after 0.2, 3, 6, 17, 23, 66, 100, and 163 days

of incubation.