Defence recognition of a stripe rust fungal effector is uncoupled

from disease outcomes in wheat

Eric C. Pereira[1‡]**, Bayantes Dagvadorj**[1‡]**, Rita Tam**[1]**, Haoran Li**[1]**, Danish Ilyas**

Baig[1]**, Mareike Möller**[1]**, Miraclemario Raphael**[1]**, Simon Williams**[1]**, Sambasivam**

Periyannan[1,2,3]**, Florence Danila**[1]**, John P. Rathjen**[1]**, and Benjamin Schwessinger**[1]

[1]Research School of Biology, The Australian National University, Canberra, ACT, 2601

Australia

[2]Commonwealth Scientific and Industrial Research Organisation Agriculture and

Food, Canberra ACT 2601, Australia

[3]School of Agriculture and Environmental Science, Centre for Crop Health, University

of Southern Queensland, Toowoomba, QLD 4350, Australia

‡ **shared co-first authors**

*Corresponding author: *benjamin.schwessinger@anu.edu.au*

Abstract

Plant resistance (*R*) and pathogen avirulence (*Avr*) gene interactions are central to

pathogen recognition and disease resistance in crops. Functional characterisation of

recognised *Avr* effectors of *Puccinia striiformis* f. sp. *tritici* (*Pst*) lags other key fungal

pathogens of wheat. Here, we used a wheat protoplast-based screen to identify *Avr/R*

interactions via the proxy of effector-induced defence responses in a set of diverse

wheat cultivars. We identified an *Avr* candidate, termed *AvrPstB48,* that triggers

Eric C. Pereira
MPMI

defence responses in 16 out of 24 cultivars tested. *AvrPstB48* is hemizygous, and the

Pst genome carries four divergent paralogs within a gene cluster. Analysis of these

paralogs revealed partial redundancy in their ability to activate wheat defences and

enabled us to identify a single amino acid in *AvrPstB4*8 that is necessary but not

sufficient for defence activation. Notably, the activation of defence signalling by

AvrPstB48 in protoplasts did not directly correlate with disease outcomes. Whole-plant

infection assays revealed that some cultivars which exhibited strong defence

activation in the protoplast assay are susceptible to the *Pst* isolate Pst104E137A- from

which *AvrPstB48* is derived. Comparison of infection dynamics of two wheat cultivars

that differ in their *AvrPstB48* recognition capacity revealed a delay in disease

progression in the recognising cultivar Avocet *S* compared to the non-recognising

cultivar Morocco. While correlative only, our observations, combined with other recent

reports, support a 'recognize-then-suppress' model of plant-pathogen interaction

where disease outcomes are driven not only by simple *Avr/R* interactions but also by

pathogen effectors that suppress defence signalling downstream of effector

recognition.

Keywords: *Puccinia striiformis* f. sp. *tritici*, effectors, avirulence genes, protoplast

screening, wheat resistance, plant defence.

INTRODUCTION

Plant pathogenic fungi are major threats to crop yields globally. This is exemplified by

wheat stripe (yellow) rust caused by *Puccinia striiformis* f. sp. *tritici* (*Pst*), which causes

significant crop losses driven by the ongoing intercontinental spread and rapid host

adaptation of this fungus (Hovmøller et al., 2010; 2025). On the molecular level,

disease is caused by a large set of effector proteins that are secreted during wheat

(Wang et al., 2023). These effectors modulate host cellular processes to facilitate host

colonisation, including nutrient acquisition and suppression of defence responses

(Zheng et al., 2023). The suppression of defence responses is especially important for

biotrophic plant pathogens such as *Pst* which require living host tissue for prolonged

periods to complete their lifecycles (Schwessinger, 2017). A subset of effectors

designated avirulence (*Avr*) effectors are detected by host resistance (*R*) genes in

certain genotypes and cultivars (Xia et al., 2020). *Avr* recognition triggers a myriad of

defence responses that restrict pathogen colonisation, including transcriptional

reprogramming and, at times, programmed cell death (Dodds & Rathjen, 2010).

Several *Avr* genes from other wheat rust fungi have been cloned including *AvrSr13,*

AvrSr22, AvrSr27, AvrSr35, AvrSr50, and *AvrSr62* from *P. graminis* f. sp. *tritici* (Arndell

et al., 2024; Chen et al., 2017; Chen et al., 2024; Outram et al., 2024; Upadhyaya et

al., 2021; Salcedo et al., 2017) and *AvrLr15* and *AvrLr21* from *P. triticina* (Cui et al.,

2024; Shen et al., 2024). Similar advances have been reported for *Blumeria graminis*,

with the identification of *AvrPm2* (Praz et al., 2017) and *AvrPm3* (Bourras et al., 2019),

and *Zymoseptoria tritici* with the identification of *AvrStb6*, *Avr3D1* and *AvrStb9*

(Amezrou et al., 2023; Meile et al., 2018; Zhong et al., 2017).

Disease outcomes are not always defined by *Avr/R* recognition events alone. Some

effectors suppress defence signalling or modulate host metabolic pathways to subvert

defence activation (Oliveira-Garcia et al., 2023; Ramírez-Zavaleta et al., 2022; Rhodes

et al., 2022). Defence suppressive effectors can be categorised into causing effector-

triggered susceptibility (ETS) or effector-triggered susceptibility II (ETS II), depending

on the molecular targets and their sequential effect during plant infection (Jones &

Eric C. Pereira
MPMI

Dangl, 2006; Sohn et al., 2007; Win et al., 2012). ETS is defined as effector-

suppression of the defence response downstream of the recognition of conserved

pathogen molecules via pattern recognition receptors (Miya et al., 2007). ETS II is

defined as effector-suppression of defence signalling downstream of *Avr/R* recognition

events and can also be referred to as a 'recognize-then-suppress' model of plant-

pathogen interactions.

Despite significant progress in cloning wheat stripe rust *R* genes, the identification of

corresponding *Avr* genes in *Pst* has lagged far behind. To overcome these challenges,

we adapted and optimised a rapid protoplast-based screening system for functional

testing of *Pst* effector candidates (Wilson et al., 2024). We focused on hemizygous

effectors identified in the fully phased genome assembly of the Australian *Pst* isolate

Pst104E137A- (Tam et al., 2025). These effectors are of particular interest because

hemizygosity implies that only a single allele of an effector needs to undergo a

mutational event to avoid recognition in case of an avirulence effector. By screening

these previously identified hemizygous effectors, we isolated a single avirulence

effector candidate that is differentially recognised by a diverse set of wheat cultivars.

This study characterises this effector termed *AvrPstB48* and its four paralogs using a

protoplast screening system and diverse wheat cultivars. In addition to confirming their

recognition, we analysed *AvrPstB48* sequence diversity, expression patterns during

infection, recognition specificity and their potential impacts on disease outcome during

infection.

MATERIAL AND METHODS

Candidate *Pst Avr* selection and gene synthesis

Hemizygous effectors were previously identified by analysing a time course of long-

read transcriptomic datasets of Pst104E137A- as detailed in Tam et al. (2025). Briefly,

transcripts were sampled from six conditions, each with four replicates, including

ungerminated urediniospores and infected wheat leaves at 4, 6, 8, 10, and 12 days

post-infection (dpi). All samples were subjected to direct Oxford Nanopore

Technologies (ONT) cDNA sequencing. DESeq2 (Love et al., 2014) differential

expression analysis was conducted on transcript read abundance to identify

secretome genes upregulated ($\log_2$ fold change ≥ 2) early during infection (4 and 6

dpi) relative to dormancy. Inter-haplotype sequence analysis was then conducted to

shortlist the candidates based on hemizygosity. The selected candidates were

synthesised as gene blocks to facilitate precise assembly through Golden Gate cloning

(Engler et al., 2014). Here, we describe the functional characterisation of one of these

hemizygous effector candidates, ID Pst104E137_010232, referred to as *AvrPstB48*.

Identification of *AvrPstB48* paralogs

The tblastn v2.16.0+ was used to search for *AvrPstB48* paralogs using its protein

sequence against the *Pst*104E genome assembly (-evalue 1e-5) (Camacho et al.,

2009). We applied a low identity cut-off of 0.4, considering effector proteins are

typically sequence-divergent yet structurally conserved (Yu et al., 2023). The search

results were used to locate the annotated gene hits. Multiple protein sequence

alignments of *AvrPstB48* and the paralogs Pst104E137_025766 (*AvrPstB48_P1*),

Pst104E137_025767 (*AvrPstB48_P2*) and Pst104E137_010231 (*AvrPstB48_P3*),

were performed using MAFFT v7.490 (Katoh & Standley, 2013) and visually inspected

in Geneious to confirm overall similarities in physicochemical properties of individual

amino acids in the alignment.

Effector prediction

We used full-length protein sequences of *AvrPstB48, AvrPstB48_P1,*

AvrPstB48_P2/P4, and *AvrPstB48_P3* with SignalP6 to predict the presence and

location of signal peptides (Teufel et al., 2022). We used respective protein sequences

without signal peptide with EffectorP3 to predict the probability of the proteins to

encode for effectors, including preference for cytoplasmic and/or apoplastic effector

location (Sperschneider & Dodds, 2022).

In silico **structural modelling**

The predicted tertiary structures of *AvrPstB48* and its paralogs were generated using

the AlphaFold Server v3.0 (DeepMind, UK) (Abramson et al., 2024). Models were

generated based on the protein sequences with the signal peptides removed. The

predicted Template Modelling (pTM) score can be used to assess global model quality

and domain orientation, where scores ≥0.5 are generally considered acceptable for

reliable domain-level predictions (Jumper et al., 2021). The initial scores for all

structural models were <0.5 (Table S1). However, we observed from those models that

~50 residues at the N-terminus of each protein were unstructured. We also noted that

the sequences of the proteins were rich in conserved cysteine residues. In other rust

effectors, these have been shown to co-ordinate Zn^{2+} (Zhang et al., 2018; Outram et

al., 2024). We therefore repeated the predictions with the N-terminal truncated and

Zn^{2+} included. We observed an overall improvement in the pTM scores for all models

>0.5 when the N-terminal region was removed, and the scores generally improved

further when Zn was added to the prediction. The interface prediction score for Zn

binding was, in most cases>0.7, indicating acceptable interaction/binding reliability

(Abramson et al., 2024) (Table S1). The resulting models were visualised and

analysed using PyMOL and ChimeraX v1.8 (UCSF) (Meng et al., 2023). Structural

comparison of *AvrPstB48* and its paralogs *(P1, P2/P4, and P3)* was performed via

superimposition in PyMOL, using *AvrPstB48* as the reference structure to identify

conformational differences that may underlie functional divergence (Table S1).

Golden Gate assembly of expression constructs

Synthetic double-stranded DNA fragments (gBlocks Gene Fragments; Integrated DNA

Technologies (IDT), Coralville, IA, USA) encoding the hemizygous effector candidate

AvrPstB48, its paralogs *AvrPstB48_P1*, *AvrPstB48_P2*, *AvrPstB48_P3*, and the

mutants *AvrPstB48_*D133G and *AvrPstB48_P3_*G133D were synthesised with

flanking *BsaI* recognition sites to generate an AATG overhang downstream of the long

intergenic region and a GCTT overhang upstream of the short intergenic region to

facilitate level 1 Golden Gate assemblies. The digested fragments were ligated into

the pWDV1 expression vector, which had also been digested with *BsaI*. The reaction

was cycled between 37°C for 3 minutes (to allow digestion) and 16°C for 4 minutes

(for ligation) for 25 cycles, followed by a final incubation at 50°C for 5 minutes and

80°C for 5 minutes to inactivate the *BsaI* enzyme. Following the Golden Gate reaction,

5 µL of the reaction mixture was transformed into competent *Escherichia coli* cells.

The cells were incubated on ice for 30 minutes, heat-shocked at 42°C for 30 seconds,

then returned to ice for 5 minutes. SOC medium (200 µL; containing 2% tryptone, 0.5%

yeast extract, 10 mM NaCl, 2.5 mM KCl, 10 mM $MgCl_2$, 10 mM $MgSO_4$, and 20 mM

glucose) was added, and the mixture was incubated at 37°C for 1 hour with shaking

at 250 rpm.

Eric C. Pereira
MPMI

Transformed *E. coli* cells were plated on LB agar supplemented with 35 µg/mL

chloramphenicol and incubated overnight at 37°C. Individual colonies were screened

by colony PCR using gene-specific primers to confirm the presence of *Avr* candidate

inserts (Table S2). PCR-positive clones were cultured in LB broth for plasmid

extraction using a standard miniprep protocol (Qiagen). Extracted plasmids were

subjected to restriction enzyme digestion with insert-specific restriction sites to validate

correct plasmid architecture. Clones showing the expected digestion patterns were

then selected for full-length plasmid verification via ONT sequencing. Whole-plasmid

sequencing was performed using ONT MinION flow cells, and raw reads were

analysed in Geneious Prime v2025.1.3 using Minimap alignment against the plasmid

reference map to confirm correct insert orientation and base pair sequence integrity

(Li, 2018) (Table S3).

Plant materials and growth conditions for protoplast isolation

For each wheat cultivar (cv), Chinese 166, Avocet/*Yr1*, Heines VII, Nord Desprez,

Triticum spelta, Avocet/*Yr5*, Heines Peko, Heines Kolben, Avocet/*Yr6,* Reichersberg

42, Avocet/*Yr7*, Compare, Avocet/*Yr8,* Clement, Avocet/*Yr9*, *Yr15*, Avocet/*Yr15*,

Spalding prolific, Avocet/*YrSp*, Avocet S, Breakwell, Gabo, Coorong and Morocco, 15

seeds were sown in each pot (14 cm in diameter, 10 cm in depth, and approximately

1.5 L in volume). The plants were grown in controlled-environment growth cabinets at

18°C under a 16 h light/8 h dark photoperiod and a light intensity of 300 µmol/m²/s.

Relative humidity was maintained at 60% during the light period and 70% during the

dark period. Ten grams of soluble fertiliser (HORTICO) was dissolved in 4.5 L of water

and applied to each tray of plants. Plants were watered every two days. At 8 days after

sowing, leaves were harvested for protoplast isolation.

Protoplast isolation

Wheat protoplasts were isolated using the protocol described by Wilson et al. (2023,

2024), with minor modifications. Briefly, protoplasts were isolated by making shallow

incisions in the leaf epidermis of the first and second true leaf, followed by peeling the

epidermis to reveal the underlying mesophyll cells. The excised leaf segments were

incubated with the peeled surface in contact with a 0.6 M mannitol solution for 10

minutes. Subsequently, the segments were transferred to a Petri plate (4.5 cm)

containing an enzyme solution composed of MES buffer (pH 5.7, 20 mM), mannitol

(0.8 M), KCl (10 mM), cellulase (Yakult R-10) 1.5% w/v, macerozyme (Yakult R-10)

0.75% w/v, $CaCl_2$ (10 mM), and BSA 0.1% w/v. The plate was wrapped in aluminium

foil to prevent light exposure and placed on an orbital shaker set to 60 rpm for 3 h at

room temperature. Following enzyme digestion, released protoplasts were filtered

through a 40 µm nylon cell strainer. Protoplasts were then collected from the enzyme

solution by gentle centrifugation at 100 g for 3 minutes using a 30 mL round-bottom

tube, resuspended and washed in W5 solution (MES buffer, pH 5.7, 2 mM; KCl, 5 mM;

$CaCl_2$, 125 mM; NaCl, 154 mM) and incubated at 4°C for 40 minutes to allow settling.,

A 10 µl aliquot of the concentrated protoplasts was taken for cell quantification and

quality assessment using a hemocytometer and light microscope. The protoplasts

were left at 4°C for 40 minutes to allow settling. The W5 solution was then removed,

and the protoplasts were resuspended to a final concentration of 300,000–350,000

cells/mL using MMG solution (MES buffer, pH 5.7, 4 mM; mannitol, 0.8 mM; $MgCl_2$, 15

mM). The protoplasts were used immediately for subsequent transfection.

Protoplast transfection

Eric C. Pereira
MPMI

Plasmid DNA (1 µg/µL) was prepared using standard mini and maxi-prep protocols

(SV Wizard kits, Promega, Madison, WI, USA). Protoplast isolation and PEG-mediated

transfection followed the method described by Wilson et al. (2024), adapted to a 96-

well plate format to increase scalability and throughput (Khan et al., 2022; Lloyd et al.,

2022) with additional optimizations (Pereira et al., 2025).

All expression constructs were based on the self-replicating pWDV1 vector backbone

and were driven by the maize ubiquitin (UBI) promoter (Table S3). This included (i)

individual *Avr* effector candidates, (ii) wheat resistance (*R*) genes of positive control

Sr50, and (iii) luciferase reporter constructs. Reporter activity was quantified using a

dual-luciferase system adapted from Wilson et al. (2024), in which a defence-inducible

reporter plasmid (UBI-pDefense) and a constitutive normalisation plasmid (UBI-pRedf)

were co-transfected. These reporters use the same substrate but encode luciferases

with distinct emission spectra, which were read using dedicated Lumi Green band-

pass (~505 – 560 nm) and Red NB long-pass (cut-on ~600 nm) filters on a

luminescence plate reader (Tecan, Mannedorf, Switzerland; Infinite 200Pro) to allow

ratiometric normalisation of defence induction.

For each transfection, three classes of reactions were prepared in parallel: (i) *Avr*

candidates under test, (ii) a positive control (*AvrSr50* + Sr50), and (iii) a negative

control (*AvrSr50* alone, no matching *R* gene). In positive-control wells, 0.5 µg of the

AvrSr50 effector plasmid, 2 µg of the Sr50 resistance gene plasmid, and 5 µg of each

reporter plasmid (UBI-pDefense and UBI-pRedf) were co-transfected per well.

Negative-control wells received 0.5 µg of the *AvrSr50* plasmid plus 5 µg of each

reporter plasmid but no *Sr50* plasmid, to confirm that defence reporter activation

required a cognate *R* gene. *Avr*-only reactions for candidate effectors were assembled

analogously, using 1 µg of the *Avr* candidate plasmid and 5 µg of each reporter

plasmid, without an *R* gene. Plasmid DNA mixes were dispensed into sterile 96-well

V-bottom plates (Costar). Freshly isolated wheat protoplasts (50 μL per well) were

added, followed by 50 μL of PEG solution (40% w/v PEG 4000, 0.2 M mannitol, 100

mM CaCl$_2$). The suspension was gently mixed by pipetting up and down five times,

then incubated for 10 min at room temperature to allow DNA uptake. Transfection was

stopped by adding 200 μL of W5 solution (154 mM NaCl, 125 mM CaCl$_2$, 5 mM KCl,

2 mM MES, pH 5.7). The plates were then incubated under low-light conditions at room

temperature for 16 h to permit expression prior to luminescence quantification.

Monitoring luciferase expression

After the 16h incubation, protoplasts were prepared for luminescence measurement.

The W5 supernatant was removed gently using a multichannel pipette without

disturbing the settled protoplast pellet. Fifty μL of 1× cell lysis buffer (#E3971;

Promega) was added, and the samples were gently mixed by pipetting 5 times, then

incubated for 15 minutes to ensure complete lysis. Fifty μL of lysate was collected and

transferred to an opaque, flat-bottom, white, solid polystyrene 96-well plate (Corning

96-well NBS Microplate, #CLS3600). The total luminescence was measured using a

plate reader (Tecan, Mannedorf, Switzerland; Infinite 200Pro, 1000 ms integration,

0 ms settle) without adding substrate to measure background signal. Immediately after

this baseline read, 50 μL of Steady-Glo luciferase substrate (Promega, #E2520) was

added to each well, and the luminescence was measured again. The plate reader was

programmed first to measure total luminescence without filters, using an integration

time of 1000 ms and a settle time of 0 ms. Subsequently, luminescence was recorded

sequentially through the installed Lumi Green band-pass (~505–560 nm) and Red NB

long-pass (cut-on ~600 nm) filters under the same acquisition settings. These filtered

measurements allowed separation of the defence-inducible reporter (UBI-pDefense)

and the constitutive normalisation reporter (UBI-pRedf), which use the same substrate

but emit at different wavelengths.

Protoplast RT-PCR

Wheat protoplasts were prepared as described above and dispensed at 900 µL per

replicate (3.5×10^5 cells mL^{-1} in MMG buffer) for transfection. For each transfection,

9 µg of the candidate expression plasmid was co-transfected with 5 µg of each

reporter. Transfections were performed in four independent biological replicates. After

16 h at room temperature under low light, 50 µL of each suspension was removed for

luciferase quantification, and the remaining protoplasts were pelleted by centrifugation

at 300 x g for 3 min. The supernatant was removed, snap-frozen in liquid nitrogen, and

stored at −80 °C for RNA extraction.

Total RNA was isolated with the Direct-zol RNA MiniPrep Plus kit (Zymo Research)

following the manufacturer's instructions, including on-column DNase I treatment.

RNA integrity was verified by agarose gel electrophoresis, and concentration and

purity were measured with a Qubit fluorometer and a NanoDrop spectrophotometer.

For cDNA synthesis, 1 µg of total RNA was reverse-transcribed using the High-

Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific) according to the

manufacturer's protocol.

Gene-specific primers targeting *AvrPstB48* and the paralog *AvrPstB48_P3* were

designed in Geneious. Wheat 18S rRNA (Ta-18S) primers served as an endogenous

reference control (Table S2).

Conventional endpoint RT-PCR was performed with GoTaq DNA polymerase

(Promega) in 10 µL reactions with 3 µL cDNA template. Thermal cycling used a

standard program (1 cycle of 95 °C for 1 min and 30 s; 30 cycles of 95 °C for 30 s,

primer-specific annealing temperature of 50 °C for 30 s, and 72 °C for 2 min; final

extension 72 °C for 5 min). Empty-vector (EV) transfections were included as negative

biological controls; no-template controls (NTC) were run for each primer pair.

Amplicons (5 µL per reaction) were resolved on 1% agarose in 1× TAE, stained, and

imaged alongside a 1 kb Plus DNA Ladder (NEB #N3200).

Inoculation of wheat plants with *Pst*

Fresh spores of *Pst* isolate Pst104E137A- (Schwessinger et al., 2018) were collected

from infected leaf tissue. Spores were suspended in 3M Novec solution and measured

using a CellDrop Automated Cell Counter (DeNovix Inc.) to a concentration of

approximately $5×10^5$ spores/mL. Using a paintbrush, spores were inoculated on the

upper surface of the first-emerged leaf of wheat cv Avocet *S* and Morocco 14 days

after sowing at the same conditions described above. Infected plants were lightly

sprayed with clean deionised water before being placed in an opaque, 25-L plastic

container (33 cm x 22.5 cm x 47 cm) with a lid. Containers with infected plants were

wrapped with black plastic bags and incubated at 10°C in the dark to initiate *Pst*

germination. After 48 hours, the infected plants were removed from the container and

transferred back to the growth cabinet. Each infected leaf was assessed according to

Dracatos et al. (2016) over 10 or more days post-infection (dpi).

Hyphae staining of infected wheat leaves

Staining was performed as described by Redkar *et al.* (2018). Freshly harvested leaf

tissues were placed in 2 mL tubes containing 100% ethanol for three days or until the

majority of the chlorophyll had been removed. When the tissues were nearly devoid of

colour, ethanol was replaced with 10% aqueous potassium hydroxide and incubated

on a dry heating block at 85°C for two hours. Cleared tissues were carefully washed

10 times in 1x phosphate-buffered saline (PBS Thermo Fisher Scientific, 28237), pH

7.4. Propidium iodide (PI; Sigma-Aldrich, P4170) was mixed with wheat germ

agglutinin conjugated with AlexaFluor488 (WGA-AF488; Thermo Fisher Scientific,

W11261) to create a PI-WGA AF488 staining solution (20 μg/mL PI, 10 μg/mL WGA-

AF488, and 0.1% Tween 20 in 1× PBS, pH 7.4). After discarding the last PBS wash,

PI-WGA-AF488 staining solution was added to the 2 mL tubes to fully submerge the

tissues. Vacuum infiltration was applied at 7 millibars, repeated 5 times for 2 minutes

each, with 2-minute vacuum release in between. Stained tissues were washed twice

in 1x PBS to remove excess staining solution, transferred to fresh 1x PBS and stored

at 4°C in the dark.

Callose staining of infected wheat leaves

Freshly harvested leaf tissues were placed in 2 mL tubes containing 1x PBS, pH 7.4

and vacuum infiltrated at 7 millibars until they sank to the bottom of the tube. Prior to

staining, leaf tissues were hand-sectioned paradermally using a razor blade. Eight

pieces of sectioned leaf tissue were placed in each well of the 96-well plate and

washed with 200 μL 1x PBS. Tissues were stained with 0.1% aniline blue (AB; Sigma-

Aldrich, 415049) in 1x PBS for 60 mins with vacuum infiltration and washed twice in

1x PBS. WGA-AF488 staining solution (10 μg/L WGA-AF488 and 0.1% Tween 20 in

1× PBS, pH 7.4) was added to the samples and vacuum infiltrated at 7 millibars for 5

times, 2 minutes each, with 2-minute vacuum release in between. Samples were

washed twice in 1x PBS, transferred to fresh 1x PBS and stored at 4°C in the dark.

Confocal fluorescence microscopy

Stained leaf tissues were mounted onto a glass slide with 1x PBS and imaged under

a TCS SP8 confocal microscope (Leica Microsystems) using 20x (PI-WGA-AF488)

and 63x (AB-WGA-AF488) water-immersion objectives. WGA-AF488-stained fungal

structures were visualised at 500-540 nm following excitation with an Argon laser at

488 nm. PI-stained cell wall was visualised at 580-630 nm following excitation with a

DPSS laser at 561 nm. AB-stained callose was visualised at 430-500 nm following UV

excitation at 405 nm. To obtain a composite image with greater depth of field, Z-

stacking was performed with each optical section captured at a 0.1 μm interval. Images

were processed (e.g., maximum projection, brightness and contrast) using Leica

Application Suite X software (LASX, Leica Microsystems).

Data collection and statistical analysis

Differences in normalised luminescence values were assessed using one-way

analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference

(HSD) post hoc test (Wilson et al., 2024). The infectious hyphae cover area per leaf

surface area was quantified using images obtained from PI-WGA-AF488-stained leaf

tissues. Meanwhile, the callose signal area was quantified using images obtained from

AB-WGA-AF488-stained leaf tissues. These quantifications were performed using

ImageJ software (National Institutes of Health) on a Wacom Cintiq graphics tablet

(Wacom Technology). Statistical analysis of data obtained from confocal images was

performed using Welch's t-test.

RESULTS

A hemizygous stripe rust candidate effector triggers defence signalling in wheat

protoplasts

The hemizygous effector candidates were identified previously from the nuclear

haplotype-phased genome of the *Pst* reference isolate Pst104E137A- (Tam et al.,

2025). Here we describe the functional characterisation of one of these hemizygous

effector candidates referred to here as *AvrPstB48*. The name *AvrPstB48* reflects its

predicted avirulence function "Avr" in *Pst*, with "B" representing its broad recognition

pattern across diverse wheat cultivars, and "48" corresponding to its internal candidate

ID in the screening pipeline. Additional hemizygous effector candidates, including

Pst104E137_008104 (#47) and *Pst104E137_024713* (#49), were selected using the

same criteria and served as comparisons for validation assays (Tam et al., 2025).

Because *AvrPstB48* expression pattern aligned with a function during plant infection

(Tam et al., 2025), we cross-validated its role as effector using the machine learning

tool EffectorP3 (Sperschneider & Dodds, 2022). *AvrPstB48* was predicted as an

effector with a most likely intracellular localisation. We therefore decided to perform

transient expression of *AvrPstB48* in wheat protoplasts with its signal peptide removed

to allow for intracellular localisation. Using a dual luciferase-based defence reporter

assay with the D14 defence promoter (Wilson et al., 2024), we observed a robust and

specific increase in ratiometric luminescence following co-expression of *AvrPstB48*

and the defence reporter, indicating a recognition-dependent defence response in

Avocet *Yr5* and Avocet *S* (Figure 1). The increase in ratiometric luminescence

observed when expressing *AvrPstB48* was similar to the *AvrSr50/Sr50* gene pair,

which served as a positive control to validate the assay performance (Wilson et al.,

2024). By contrast, two other hemizygous effector candidates, #47 and #49, did not

induce measurable reporter activity being similar to the negative control, *AvrSr50*

(Figure 1). Together, these results are consistent with *AvrPstB48* encoding a

recognised effector activating host defence responses in specific wheat cultivars

(Table S5).

Given the strong defence promoter activation observed for *AvrPstB48*, further analysis

was conducted to identify and characterise its paralogs. Sequence homology analysis

was performed by tblastn and revealed two divergent paralogs located in close

proximity on each haplotype of chromosome 10, *AvrPstB48*_P1, *AvrPstB48*_P2,

AvrPstB48_P3 and *AvrPstB48*_P4. Among these allelic pairs, *AvrPstB48*_P2 and

*AvrPstB48*_P4 share 100% nucleotide identity, whereas *AvrPstB48*_P1 and

AvrPstB48_P3 exhibit 98.2% nucleotide identity. Relative to AvrPstB48, the *P2/P4* pair

is more closely related (90.8%), whereas the *P1/P3* pair is more divergent (57.6%

identity to *AvrPstB48_P1*) (Figure 2a). Consistent with this, protein sequence

alignments show that AvrPstB48_P2/P4 are highly similar to AvrPstB48, whereas

AvrPstB48_P1 and AvrPstB48_P3 carry multiple non-synonymous substitutions and

are more divergent at the amino-acid level (Figure 2b). AlphaFold3 modelling

supported a shared fold among AvrPstB48, P1, P2/4, and P3. Backbone superposition

yielded pairwise RMSDs of ~0.5–1.5 Å, with divergence concentrated in the C-terminal

helical region and adjacent loops (Fig. 2c). Global pTM values (≈0.50–0.57) indicate

these are hypothesis-generating rather than definitive models; nevertheless, they are

consistent with a conserved core and local surface variation between the paralogs.

The locations of these variations correspond to clusters of substitutions in the multiple

sequence alignment (Fig. 2b) and may modulate protein stability or interactions with

host targets. Temporal expression profiling across infection stages revealed distinct

expression patterns among the paralogs (Figure 2d). *AvrPstB48* exhibited peak

expression at 8 days post-inoculation (dpi), reaching its highest levels relative to the

other paralogs. All paralogs also peaked at 8 dpi, although expression intensity varied,

with *P1*, *P2/P4*, and *P3* remaining moderately expressed throughout the infection time

course. None of the paralogs were expressed in ungerminated spores, supporting their

likely function in host-pathogen interactions.

AvrPstB48 recognition is conserved across multiple wheat cultivars

To further assess defence promoter activation triggered by *AvrPstB48*, we tested their

activities in additional wheat cultivars including Avocet S, Morocco, *Triticum spelta Yr5*,

Chinese 166 and Gabo, using three previously validated defence reporter promoters

(D2, D14, and D15) (Wilson et al., 2024). Significant activation of each promoter was

observed in response to *AvrPstB48* expression in Avocet *S* (Figure 3a), *T. spelta Yr5*

and Gabo (Figures S1a, S1b), indicating effector recognition. In contrast, no increase

in ratiometric luminescence signal was detected in Morocco (Figure 3b) or Chinese

166 (Figure S1c), suggesting a lack of *AvrPstB48* recognition in these cultivars.

Additional screening of 16 wheat cultivars confirmed consistent activation in all near-

isogenic Avocet lines, while eight cultivars showed no measurable response (Table

S5). Importantly, the positive control (*AvrSr50/Sr50*) consistently induced strong

luminescence across all tested backgrounds, validating assay performance.

AvrPstB48 paralogs AvrPstB48_P1 and AvrPstB48_P2/P4, but not

AvrPstB48_P3, are recognized in multiple wheat cultivars

To assess the functional redundancy of *AvrPstB48*, we tested recognition of its

paralogs in the same set of wheat cultivars, Avocet S, Morocco, *T. spelta Yr5*, Gabo,

and Chinese 166, using activation of the D14 promoter as a readout. Defence

activation was detected for *AvrPstB48_*P1 and *AvrPstB48_*P2/P4 in Avocet *S* (Figure

4a), *T. spelta Yr5,* and Gabo (Figures S2), suggesting a similar recognition spectrum

to *AvrPstB48*. *AvrPstB48_P3* failed to activate the D14 defence promoter in any of the

tested cultivars. No induction was observed for any of the paralogs tested in Morocco

(Figure 4b) and Chinese 166 (Figure S2).

Given that *AvrPstB48_P3* shares overall structural and sequence similarity with

AvrPstB48 (Figure 2), we investigated whether it could interfere with *AvrPstB48-*

triggered defence signalling. In co-expression assays in Avocet *S* protoplasts,

AvrPstB48 was delivered either alone or together with *AvrPstB48_*P3 at a 1:1 plasmid

ratio. Co-expression did not reduce relative ratiometric luminescence signal induction

when compared to *AvrPstB48* alone, indicating that *AvrPstB48_P3* does not act as a

competitive suppressive paralog when co-expressed in wheat protoplasts (Figure 4c).

To verify effector transcript accumulation for both *AvrPstB48* and *AvrPstB48_P3*, we

performed endpoint RT-PCR on cDNA from protoplasts transfected with the

corresponding expression constructs. Gene-specific primer pairs yielded the expected

amplicons for *AvrPstB48* and *AvrPstB48_P3* only in the matched transfection. No

bands were detected in empty vector (EV) controls amplified with either primer set.

The wheat 18S rRNA control amplified in all samples, supporting cDNA quality (Figure

S3).

Aspartic acid at position 134 is necessary but not sufficient for defence

activation

To investigate why *AvrPstB48_P3* does not trigger defence promoter activation in

Avocet S, we explored potential structural and biochemical differences among

AvrPstB48 and its paralogs. Structural predictions using AlphaFold3 revealed that all

Eric C. Pereira
MPMI

paralogs adopt a broadly conserved fold (Figure 2d). Closer inspection revealed

several amino acid substitutions specific to *AvrPstB48_P3*, particularly in surface-

exposed loops that may be involved in effector recognition and receptor interactions.

One notable difference occurs at position 134, where *AvrPstB48, AvrPstB48_P1*, and

AvrPstB48_P2 encode an aspartic acid (D), while *AvrPstB48_P3* encodes a glycine

(G) (Figure 5a). This substitution replaces a negatively charged residue (D) with a

small, nonpolar amino acid (G) at a position that is predicted to be surface-exposed

on a protruding loop. This makes the D134G substitution a prime candidate for

functional testing.

To further investigate the functional significance of the amino acid variation at position

134, synthetic gene fragments were designed to encode *AvrPstB48* carrying an

aspartic acid to glycine substitution (D134G), and *AvrPstB48_P3* carrying the reverse

substitution (G134D). These constructs were tested for their ability to induce defence

promoter activation in Avocet *S* using the D14 promoter-luciferase reporter system in

wheat protoplasts. AvrPstB48_D134G failed to activate the defence reporter when

compared to the increase of ratiometric luminescence when expressing wild-type

AvrPstB48 (Figure 5b). This indicates that the D134G substitution is sufficient to

disrupt recognition. The reciprocal mutant AvrPstB48_P3_G134D also failed to induce

defence promoter activity, suggesting that the aspartic acid at position 134 is not the

only residue contributing to the avoidance of recognition in *AvrPstB48_P3*.

AvrPstB48*-triggered reporter activation in protoplasts does not correlate fully

with *Pst* resistance *in planta*

We assessed infection outcomes using the Australian pathotype Pst104E137A-

(Schwessinger et al., 2018) to evaluate whether the *AvrPstB48*-triggered reporter

activation observed in wheat protoplasts corresponded to resistance *in planta*. Wheat

lines previously tested in the protoplast assay were inoculated and disease severity

scored at 10 days post-inoculation (dpi) using a 0–9 infection type (IT) scale

(Hovmøller et al., 2017). Reporter activation in protoplasts did not correlate with

disease outcomes in all cultivars tested. While *T. spelta Yr5* and Chinese 166 exhibited

full resistance (IT = 0), Gabo, Morocco, and Avocet *S* were highly susceptible (IT = 9),

despite *AvrPstB48* triggering defence promoter activation in Gabo and Avocet *S*

(Figure S4).

AvrPstB48 recognition correlates with delayed infection progression in the
susceptible wheat cultivar Avocet S

To explore whether *AvrPstB48* recognition might influence infection dynamics even in

a susceptible genotype, we compared Avocet *S* and Morocco. Both of these are

susceptible to the *Pst* isolate Pst104E137A-, but Avocet *S* responds to *AvrPstB48*

expression in the protoplast assay, whereas Morocco does not. At 8 dpi, neither line

exhibited macroscopic ureodiniospore formation. By 10 dpi, both displayed visible rust

symptoms, however, Morocco developed more extensive ureodiniospore coverage,

indicating a higher sporulation capacity and greater disease severity (Figure 6A).

Confocal microscopy supported these observations, revealing more abundant and

earlier hyphal growth in Morocco compared to Avocet *S* at 8 dpi (Figure 6B). Although

both lines were heavily colonised by 10 dpi, Avocet *S* retained localised areas with

reduced fungal presence, consistent with greater disease severity. Quantification of

infectious hyphal cover over time showed broadly similar colonisation dynamics

between the two genotypes, with Morocco tending to exhibit slightly higher mean

hyphal coverage at later time points (Figure 6D).

To assess host immune responses underlying these differences, we examined callose

deposition, a hallmark of plant defence. At 8 dpi, Avocet *S* exhibited greater callose

accumulation than Morocco, particularly around infection sites (Figure 6C), and this

was supported by quantitative analysis (Figure 6E).

Discussion

Recent advances in high-quality genome assemblies of *Pst* (Wang et al., 2024; Tam

et al., 2025), coupled with advances in effector identification and functional genomics,

have significantly enhanced our capacity to investigate host–pathogen interactions of

Pst and other rust wheat rust fungi (Arndell et al., 2024; Chen et al., 2017; Chen et al.,

2024; Cui et al., 2024; Outram et al., 2024; Salcedo et al., 2017; Shen et al., 2024;

Upadhyaya et al., 2021; Wilson et al., 2024). Additionally, we have gained deeper

insights into wheat defence responses by integrating comprehensive effector libraries

with protoplast transient expression methods (Arndell et al., 2024; Wilson et al., 2024).

Within this framework, we identified *AvrPstB48* as a hemizygous avirulence candidate

effector that triggers defence signalling in a wheat cultivar-specific manner. Using

wheat protoplasts, we deliberately focused on intracellular expression of *AvrPstB48,*

removing the signal peptide encoding sequence from the expression clone based on

localisation prediction made with EffectorP3. In cultivars such as Avocet *S, T. spelta*

Yr5 and Gabo, intracellular expression of *AvrPstB48* elicited strong activation of

defence reporters, comparable in magnitude to the canonical *Avr/R* control

AvrSr50/Sr50, whereas Morocco and Chinese 166 showed no detectable response to

AvrPstB48. This clear cultivar-specific response and its intracellular recognition align

with intracellular *R* gene-mediated recognition in wheat protoplasts and mirror prior

observations *for AvrSr35, AvrSr27, AvrSr13, AvrSr22* and *AvrPm2a*, all of which

activate the same defence reporters in protoplasts when expressed in the presence of

their cognate *R* genes (Wilson et al., 2024). While we have not identified a cognate *R*

gene for *AvrPstB48*, we hypothesise that it is recognised by a broadly conserved *R*

gene that is common to the wheat cultivars that respond to *AvrPstB48*. Importantly,

the cultivar-specific recognition of *AvrPstB48* did not correlate with full disease

resistance when using the *AvrPstB48*-encoding isolate Pst104E137A– for infection

assays. Several cultivars that responded to *AvrPstB48* with defence signalling in

protoplasts, such as Avocet S and Gabo, were fully susceptible in whole plant infection

assays at 14 dpi.

We chose Avocet *S* and Morocco for more detailed phenotypic comparisons of two

fully susceptible wheat cultivars that varied in their response to *AvrPstB48*. We

selected them because they are both widely used as 'universal' susceptible *Pst*

cultivars*,* with the key difference that there are recent reports of *Pst* isolates that are

avirulent on Avocet *S* (Nazari & El Amil, 2014) but not on Morocco. Based on this

observation, there was justification to postulate that Avocet *S* carries a *R* gene

designated *YrAvS* that is observable only during whole plant infection assays and with

certain *Pst* isolates. We did observe a lower sporulation and hyphal coverage in Avocet

S when compared to Morocco in our detailed macroscopic and microscopic time

course infection comparison using Pst104E137A-. This infection delay in Avocet *S* is

consistent with its ability to recognise *AvrPstB48,* upon which we propose the

presence of the uncharacterised *YrAvS*. We do not know if *AvrPstB48* recognition

contributes to infection delay and if it is recognised by *YrAvS*. An infection delay could

also be caused by the recognition of another effector by *YrAvS* or any other unrelated

genetic differences between Avocet *S* and Morocco. However, it is tempting to

speculate that the recognition of *AvrPstB48* contributes to the infection delay, because

recent segregation analysis of a *PstS8* selfing population suggests the presence of an

effector that can suppress the recognition of *PstS8* on Avocet *S*, rendering *PstS8*

virulent despite the presence of an avirulence effector (Livbjerg et al., 2025). Similar

observations of "inhibitor" loci that suppress *Avr* recognition have been made for the

flax rust fungus *Melampsora lini* (Lawrence et al., 1981) and the barley powdery

mildew fungus *Blumeria graminis* f. sp. *hordei* (Caffier et al., 1996). These are

consistent with a recent observation in which some *Blumeria graminis* f. sp. *tritici*

isolates employ the effector SvrPm4 to suppress the recognition of AvrPm4 by Pm4

(Bernasconi et al., 2025). We therefore hypothesise that the *AvrPstB48*-Avocet *S*

phenotype is explained by a 'recognise–then–suppress' scenario, fitting within the

concept of ETSII. In ETSII, effectors that are poised to trigger *R* gene dependent

disease resistance are attenuated by other effectors that suppress defence signalling,

allowing disease progression despite recognition (Sohn et al., 2007; Wu & Derevnina,

2023). Therefore, the infection outcomes of the *Pst*-wheat pathosystem are likely more

complex than Flor's gene-for-gene hypothesis suggests (Flor, 1971), reflecting the

complex coevolution between the obligate biotroph pathogen and its host.

Paralog analysis revealed that *AvrPstB48* belongs to a small effector family with four

out of five members triggering host defence. Two paralogs (*P1* and *P2/P4*) trigger

defence activation in *AvrPstB48* responsive cultivars, whereas *P3* consistently did not.

AvrPstB48_P3 provides an example of how local sequence changes within a shared

scaffold mediate escape from recognition. Structural modelling and mutational

analyses identified the single amino acid at position 134 as a key determinant of

recognition. In AvrPstB48, P1 and P2/P4, this position encodes an aspartic acid

(D134), whereas P3 encodes a glycine (G134). AlphaFold-based models predict that

residue 134 lies within a solvent-exposed loop projecting from a compact, cysteine-

rich core. Replacing D134 with G134 in AvrPstB48 abolished defence activation,

whereas the reciprocal G134 to D134 change in P3 failed to restore recognition. This

asymmetry indicates that D134 is necessary but not sufficient for recognition. The

predicted *AvrPstB48* fold, a compact, cysteine-rich core likely stabilised by disulphide

bonds and Zn^{2+} coordination decorated with variable surface loops, is consistent with

emerging paradigms for rust effectors such as *AvrSr27*, in which conserved cysteines

and metal-binding centres maintain the core, while modular loops control host

specificity and escape (Outram et al., 2024).

Acknowledgments

We acknowledge the contribution of the Plant Pathogen 'Omics Initiative consortium

in the generation of data used in this publication. The Initiative is supported by funding

from Bioplatforms Australia, enabled by the Commonwealth Government National

Collaborative Research Infrastructure Strategy (NCRIS).

Author contributions: B.S., B.D., R.T., and J.P.R. conceptualized the project. B.S.,

R.T. and J.P.R. acquired funding. B.S. supervised the work. E.C.P., B.D., R.T., H.L.,

D.I.B., M.M., M.R., S.W., S.P., and F.D. acquired experimental data and conducted

data analysis. E.C.P. and B.S. drafted the manuscript, and all authors contributed to

writing, reviewing, and editing.

Data availability: All supporting data is available on Zenodo (doi:

10.5281/zenodo.15845412, link: 10.5281/zenodo.15845412).

Funding: This work was supported by a Discovery Project grant from the Australian

Research Council (DP230100941) to B. S. and J. P. R. This work was supported by

computational resources provided by the Australian Government through the National

Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme. R.T. was

supported by a Grains Research and Development Corporation Graduate Research

Scholarship.

References

Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,

Willmore, L., Ballard, A.J., Bambrick, J., Bodenstein, S.W., Evans, D.A., Hung,

C.C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A.,

Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M.,

Cowen-Rivers, A.I., Cowie, A., Figurnov, M., Fuchs, F.B., Gladman, H., Jain, R.,

Khan, Y.A., Low, C.M.R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A.,

Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E.D., Zielinski, M., Žídek, A.,

Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., and Jumper, J.M. 2024.

Addendum: Accurate structure prediction of biomolecular interactions with

AlphaFold 3. Nature. 636: E4.

Amezrou, R., Audeon, C., Compain, J., Gelisse, S., Ducasse, A., Saintenac, C.,

Lapalu, N., Louet, C., Orford, S., Croll, D., Amselem, J, Fillinger, S, and Marcel,

T.C. 2023. A secreted protease-like protein in *Zymoseptoria tritici* is responsible

for avirulence on Stb9 resistance gene in wheat. PLoS Pathog. 19: e1011376.

Arndell, T., Chen, J., Sperschneider, J., Upadhyaya, N.M., Blundell, C., Niesner, N.,

Outram, M.A., Wang, A., Swain, S., Luo, M., Ayliffe, M.A., Figueroa, M.,

Vanhercke, T., and Dodds, P.N. 2024. Pooled effector library screening in

protoplasts rapidly identifies novel *Avr* genes. Nat. Plants 10: 572–580.

Bernasconi, Z., Herger, A.G., Caro, M.D.P., Kunz, L., Müller, M.C., Stirnemann, U.,

Outram, M.A., Widrig, V., Neidhart, M., Isaksson, J., Schudel, S., Rösli, S., Wicker,

T., Bender, K.W., Zipfel, C., Dodds, P.N., Figueroa, M., Sánchez-Martín, J., and

Keller, B. 2025. Virulence on Pm4 kinase-based resistance is determined by two

divergent wheat powdery mildew effectors. bioRxiv 2025.07.04.663163.

doi:10.1101/2025.07.04.663163

Bourras, S., Kunz, L., Xue, M., Praz, C.R., Muller, M.C., Kälin, C., Schläfli, M.,

Ackermann, P., Flückiger, S., Parlange, F., Menardo, F., Schaefer, L.K., Ben-

David, R., Roffler, S., Oberhaensli, S., Widrig, V., Lindner, S., Isaksson, J., Wicker,

T., Yu, D., and Keller, B. 2019. The *Avr*Pm3-Pm3 effector-NLR interactions control

both race-specific resistance and host specificity of cereal mildews on wheat. Nat.

Commun. 10: 2292.

Caffier, V., de Vallavieille-Pope, C., and Brown, J.K.M. 1996. Segregation of

avirulences and genetic basis of infection types in *Erysiphe graminis f. sp. hordei*.

Phytopathol. 86:1112–1121.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and

Madden, T.L. 2009. BLAST+: architecture and applications. BMC Bioinform. 10:

421.

Chen, J., Upadhyaya, N.M., Ortiz, D., Sperschneider, J., Li, F., Bouton, C., Breen, S.,

Dong, C., Xu, B., Zhang, X., Mago, R., Newell K, Xia, X., Bernoux, M., Taylor, J.M.,

Steffenson, B., Jin, Y., Zhang, P., Kanyuka, K., Figueroa, M., Ellis, J.G., Park, R.F.,

and Dodds, P.N. 2017. Loss of *Avr*Sr50 by somatic exchange in stem rust leads

to virulence for Sr50 resistance in wheat. Science 358: 1607–1610.

Chen, R., Chen, J., Outram, M.A., Powell, O.R., Arndell, T., Gajendiran, K., Wang, Y.L.,

Ayliffe, M.A., Blundell, C., Figueroa, M., Sperschneider, J., Vanhercke, T., Tang,

D., Xu, Y., Zhong, G., Gardener, C., Yu, G., Gourdoupis, S., Jaremko, Ł., Matny,

O., Steffenson, B.J., Boshoff, W.H.P., Meyer, W.B., Arold, S.T., Dodds, P.N., and

Wulff, B.B.H. 2024. A wheat tandem kinase sensor activates an NLR helper to

trigger immunity. Science 387: 1402-1408.

Cui, Z., Shen, S., Meng, L., Sun, X., Jin, Y., Liu, Y., Liu, D., Ma, L., and Wang, H. 2024.

Evasion of wheat resistance gene *Lr15* recognition by the leaf rust fungus is

attributed to the coincidence of natural mutations and deletion in *AvrLr15* gene.

Mol. Plant Pathol. 25: e13490.

Derbyshire, M.C., and Raffaele, S. 2023. Surface frustration re-patterning underlies

the structural landscape and evolvability of fungal orphan candidate effectors. Nat

Commun. 14:5244.

Dodds, P.N., and Rathjen, J.P. 2010. Plant immunity: Towards an integrated view of

plant-pathogen interactions. Nat. Rev. Genet. 11: 539–548.

Dracatos, P.M., Zhang, P., Park, R.F., McIntosh, RA, Wellings, CR. 2016.

Complementary resistance genes in wheat selection 'Avocet R' confer resistance

to stripe rust. Theor. Appl. Genet. 129: 65–76.

Engler, C., Youles, M., Gruetzner, R., Ehnert, T.M., Werner, S., Jones, J.D., Patron,

N.J., and Marillonnet, S. 2014. A golden gate modular cloning toolbox for plants.

ACS Synth Biol. 3: 839-43.

Flor, H.H. (1971). Current status of the gene-for-gene concept. Annual Review of

Phytopathology, 9, 275–296.

Eric C. Pereira
MPMI

Hovmøller, M.S., Walter, S., and Justesen, A.F. 2010. Escalating threat of wheat rusts.

Science 329: 369.

Hovmøller, M.S., Rodriguez-Algaba, J., Thach, T., and Sørensen, C.K. (2017). "Race

typing of *Puccinia striiformis* on wheat," in Wheat rust diseases, methods and

protocols. Editor S. Periyannan (N.Y., U.S.A: Springer Protocols Methods in

Molecular Biology), 29–40.

Hovmøller, M.S., Thach, T., Rodriguez-Algaba, J., Hansen, J.G., Meyer, M., Hodson,

D., Nazari, K., Park, R.F., Tam, R., Möller, M., Schwessinger, B., Rathjen, J.P.,

Silva, P., Riella, V., and Justesen, A.F. 2025. Long-term surveillance suggests

multiple hybridization events by nuclear reassortment and accelerated

intercontinental spread of wheat yellow rust. bioRxiv 2025.05.22.655633. doi:

10.1101/2025.05.22.655633

Jones, J., and Dangl, J. 2006. The plant immune system. Nature 444: 323–329.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. 2021. Highly

accurate protein structure prediction with AlphaFold. Nature, 596: 583–589.

Katoh, K, Standley, D.M. 2013. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. Mol. Biol. Evol. 30: 772-80.

Khan, A., Lloyd, J.P.B., Kidd, B., Lister, R. 2022. Protoplast isolation and transfection

in a 96-well plate. protocols.io. doi: 10.17504/protocols.io.yxmvmn1d6g3p/v1

Lawrence, G.J., Mayo, G.M.E., and Shepherd, K.W. 1981. Interactions between genes

controlling pathogenicity in the flax rust fungus. Phytopathology 71: 12–19.

Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.

34: 3094–3100.

Livbjerg, M., Rodriguez-Algaba, J., Brown, J.K.M., Sørensen, C.K., Hovmøller, M.,

Justesen, A.F. 2025. Genetics of avirulence and interacting inhibitors in *Puccinia*

striiformis f. sp. tritici. Fungal Genet. Biol. 181: 104037.

Lloyd, J.P.B., Ly, F., Gong, P., Pflueger, J., Swain, T., Pflueger, C., Fourie, E., Khan,

M.A., Kidd, B.N., and Lister, R. 2022. Synthetic memory circuits for stable cell

reprogramming in plants. Nat. Biotechnol. 40: 1862–1872.

Love, M.I., Huber, W., and Anders, S. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome Biol. 15: 550.

Meile, L., Croll, D., Brunner, P.C., Plissonneau, C., Hartmann, F.E., McDonald, B.A.,

and Sánchez-Vallet, A. 2018. A fungal avirulence factor encoded in a highly plastic

genomic region triggers partial resistance to septoria tritici blotch. New Phytol.

219: 1048-1061.

Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H.,

and Ferrin, T.E. 2023. UCSF ChimeraX: Tools for structure building and analysis.

Protein Sci. 32: e4792.

Miya, A., Albert, P., Shinya, T., Desaki, Y., Ichimura, K, Shirasu, K, Narusaka, Y.,

Kawakami, N, Kaku, H, and Shibuya, N. 2007. CERK1, a LysM receptor kinase,

is essential for chitin elicitor signaling in Arabidopsis. PNAS.104:19613-8.

Nazari, K., and El Amil, R. 2013. First report of resistance of wheat line Avocet '*S*' to

stripe rust caused by *Puccinia striiformis* f. sp. *tritici* (*Pst*) in Syria. Plant Dis. 97:

996.

Oliveira-Garcia, E., Yan, X., Oses-Ruiz, M., de Paula, S., and Talbot, N.J. 2023.

Effector-triggered susceptibility by the rice blast fungus *Magnaporthe oryzae*. New

Phytol. 24: 1007-1020.

Outram, M.A., Chen, J., Broderick, S., Li, Z., Aditya, S., Tasneem, N., Arndell, T.,

Blundell, C., Ericsson, D.J., Figueroa, M., Sperschneider, J., Dodds, P.N., and

Williams, S.J. 2024. *Avr*Sr27 is a zinc-bound effector with a modular structure

important for immune recognition. New Phytol. 243: 314-329.

Pereira, E., Baig, D.I., and Schwessinger, B. 2025. Wheat protoplast preparation and

transformation in a 96-well plate. protocols.io. doi:

10.17504/protocols.io.5qpvod1ndg4o/v1

Praz, C.R., Bourras, S., Zeng, F., Sanchez-Martın, J., Menardo, F., Xue, M., Yang, L.,

Roffler, S., Boni, R., Herren, G., McNally, K.E., Ben-David, R., Parlange, F.,

Oberhaensli, S., Flückiger, S., Schäfer, L.K., Wicker, T., Yu, D., and Keller, B. 2017.

*Avr*Pm2 encodes an RNaselike avirulence effector which is conserved in the two

different specialized forms of wheat and rye powdery mildew fungus. New Phytol.

213: 1301–1314.

Ramírez-Zavaleta, C.Y., García-Barrera, L.J., Rodríguez-Verástegui, L.L., Arrieta-

Flores, D., and Gregorio-Jorge, J. 2022. An overview of PRR- and NLR-mediated

immunities: conserved signaling components across the plant kingdom that

communicate both pathways. Int. J. Mol. Sci. 23: 12974.

Redkar, A., Jaeger, E., and Doehlemann, G. 2018. Visualization of growth and

morphology of fungal hyphae in planta using WGA-AF488 and propidium iodide

co-staining. Bio-Protocol. 8: e2942.

Rhodes, J., Zipfel, C., Jones, J.D.G., and Ngou, B.P.M. 2022. Concerted actions of

PRR- and NLR-mediated immunity. Essays Biochem. 66: 501-511.

Salcedo, A., Rutter, W., Wang, S., Akhunova, A., Bolus, S., Chao, S., Anderson, N.,

De Soto, M.F., Rouse, M., Szabo, L., Bowden, R.L., Dubcovsky, J., Akhunov, E.

2017. Variation in the AvrSr35 gene determines Sr35 resistance against wheat

stem rust race Ug99. Science. 358: 1604-1606.

Schwessinger, B. (2017). Fundamental wheat stripe rust research in the 21st century.

New Phytol. 213: 1625-1631.

Schwessinger, B., Sperschneider, J., Cuddy, W.S., Garnica, D.P., Miller, M.E., Taylor,

J.M., Dodds, P.N., Figueroa, M., Park, R.F., and Rathjen, J.P. 2018. A near-

complete haplotype-phased genome of the dikaryotic wheat stripe rust fungus

Puccinia striiformis f. sp. *tritici* reveals high interhaplotype diversity. mBio 9:

e02275-17

Shen, S., Wang, F., Cui, Z. Yuan, S., Meng, L., Liu, D., Ma, L., and Wang, H., 2024.

Puccinia triticina avirulence protein *AvrLr21* directly interacts with wheat

resistance protein *Lr21* to activate wheat immune response. Commun. Biol. 7:

1170.

Sohn, K.H., Lei, R., Nemri, A., and Jones, J.D. 2007. The downy mildew effector

proteins *ATR1* and *ATR13* promote disease susceptibility in *Arabidopsis thaliana*.

Plant Cell. 19: 4077-90.

Sperschneider, J., and Dodds, P.N. 2022. EffectorP 3.0: Prediction of apoplastic and

cytoplasmic effectors in fungi and oomycetes. Mol. Plant Microbe. 35: 146-156.

Tam, R., Möller, M., Luo, R., Luo, Z., Jones, A., Periyannan, S., Rathjen, J.P.,

Schwessinger, B. 2025. Long-read genomics reveal extensive nuclear-specific

evolution and allele-specific expression in a dikaryotic fungus. Genome Res. 11:

gr.280359.124.

Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gíslason, M.H., Pihl, S.I.,

Tsirigos, K.D., Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. 2022.

SignalP 6.0 predicts all five types of signal peptides using protein language
models. Nat. Biotechnol. 40: 1023-1025.

Upadhyaya, N.M., Mago, R., Panwar, V., Hewitt, T., Luo, M., Chen, J., Sperschneider,
J., Nguyen-Phuc, H., Wang, A., Ortiz, D., Hac, L., Bhatt, D., Li, F., Zhang, J., Ayliffe,
M., Figueroa, M., Kanyuka, K., Ellis, J.G., and Dodds, P.N. 2021. Genomics
accelerated isolation of a new stem rust avirulence gene-wheat resistance gene
pair. Nat Plants. 7: 1220-1228.

Wang, J., Chen, T., Tang, Y., Zhang, S., Xu, M., Liu, M., Zhang, J., Loake, G.J., and
Jiang, J. 2023. The Biological Roles of *Puccinia striiformis* f. sp. *tritici* Effectors
during Infection of Wheat. Biomolecules. 13: 889.

Wang, J., Xu, Y., Peng, Y., Wang, Y., Kang, Z., and Zhao, J. 2024. A fully haplotype-
resolved and nearly gap-free genome assembly of wheat stripe rust fungus. Sci.
Data. 11: 508.

Wilson, S., Dagvadorj, B., Tam, R., Schwessinger, B. (n.d.). Wheat protoplast
preparation and transformation (Version 1). protocols.io. doi:
10.17504/protocols.io.q26g7r3zkvwz/v1

Wilson, S., Dagvadorj, B., Tam, R., Murphy, L., Schulz-Kroenert, S., Heng, N., Crean,
E., Greenwood, J., Rathjen, J.P., and Schwessinger, B. 2024. Multiplexed effector
screening for recognition by endogenous resistance genes using positive defense
reporters in wheat protoplasts. New Phytol. 241: 2621-2636.

Win, J., Chaparro-Garcia, A., Belhaj, K., Saunders, D.G., Yoshida, K., Dong, S.,
Schornack, S., Zipfel, C., Robatzek, S., Hogenhout, S.A., and Kamoun, S. 2012.
Effector biology of plant-associated organisms: concepts and perspectives. Cold
Spring Har. Symp. Quant. Biol. 77: 235–247.

Eric C. Pereira
MPMI

Wu, C.H., and Derevnina, L. 2023. The battle within: How pathogen effectors suppress

NLR-mediated immunity. Curr. Opin. Plant Biol. 74: 102396.

Xia, C., Lei, Y., Wang, M., Chen, W., and Chen, X. 2020. An avirulence gene cluster in

the wheat stripe rust pathogen (*Puccinia striiformis* f. sp*. tritici*) identified through

genetic mapping and whole-genome sequencing of a sexual population. mSphere

5: 10.1128/msphere.00128-20.

Yu, D.S., Outram, M.A., Smith, A., McCombe, C.L., Khambalkar, P.B., Rima, S.A., Sun,

X., Ma, L., Ericsson, D.J., Jones, D.A., and Williams, S.J. 2024. The structural

repertoire of *Fusarium oxysporum* f. sp. *lycopersici* effectors revealed by

experimental and computational studies. Elife.12: RP89280.

Zhang, X., Farah, N., Rolston, L., Ericsson, D.J., Catanzariti, A.M., Bernoux, M., Ve,

T., Bendak, K., Chen, C., Mackay, J.P., Lawrence, G.J., Hardham, A., Ellis, J.G.,

Williams, S.J., Dodds, P.N., Jones, D.A., and Kobe, B. 2018. Crystal structure of

the *Melampsora lini* effector AvrP reveals insights into a possible nuclear function

and recognition by the flax disease resistance protein. P. Mol. Plant Pathol. 19:

1196-1209.

Zheng, P., Liu, M., Pang, L., Sun, R., Yao, M., Wang, X., Kang, Z., and Liu, J. 2023.

Stripe rust effector *Pst*21674 compromises wheat resistance by targeting

transcription factor TaASR3. Plant Physiol. 193: 2806-2824.

Zhong, Z., Marcel, T.C., Hartmann, F.E., Ma, X., Plissonneau, C., Zala, M., Ducasse,

A., Confais, J., Compain, J., Lapalu, N., Amselem, J., McDonald, B.A., Croll, D.,

Palma-Guerrero, J. 2017. A small secreted protein in *Zymoseptoria tritici* is

responsible for avirulence on wheat cultivars carrying the *Stb6* resistance gene.

New Phytol. 214, 619-631.

Figures

Figure 1 – *pDefense* **D14 promoter activation triggered by** *AvrPstB48* **in Avocet** *Yr5* **and Avocet**

S **protoplasts**. Dual luciferase-based reporter assays were performed using wheat protoplasts from

wheat cultivars Avocet *Yr5* **a)** and Avocet *S* **b)** to assess defence signalling. The x-axis shows the

plasmid combinations tested, including candidate effectors *Pst104E137_008104* (#47),

Pst104E137_010232 (*AvrPstB48*), and *Pst104E137_024713* (#49), along with the positive control pair

AvrSr50/Sr50. The y-axis represents normalised luminescence values, measured as the ratio of

luciferase activity driven by the D14 promoter to the internal control UBI-pRed. *AvrPstB48* triggered a

significant increase in reporter activity, indicating activation of defence responses. Statistical differences

were analysed using one-way ANOVA followed by Tukey's Honestly Significant Difference (HSD) test;

Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure 2 – Genomic, evolutionary, and structural characterisation of the candidate effector gene

AvrPstB48 **and its paralogs. a)** Genomic organisation and synteny of *AvrPstB48*

(*Pst104E137_010232*) and its paralogs (P1: *Pst104E137_025766*; P2: *Pst104E137_025767*; P3:

Pst104E137_010231; P4: *Pst104E137_010234*) across homologous regions of Chromosomes 10A and

10B, with genes shown as red or blue arrows, respectively. Grey-shaded blocks and contained numbers

represent pairwise nucleotide sequence identity (%) between all paralogs. **b)** Multiple sequence

alignment of *AvrPstB48* and its paralogs, showing conservation at the amino acid level. Identical or

similar residues are highlighted using the Clustal X colour scheme, which assigns colours based on

amino acid conservation and properties. Conserved motifs are evident across the sequences. Dashes

indicate gaps in the alignment. Pink shading marks the predicted N-terminal signal peptide; orange bars

denote predicted α-helical segments; red arrows denote predicted β-strands mapped from the

AlphaFold3 structural model onto the multiple-sequence alignment of *AvrPstB48* and its paralogs (*P1,*

P2/P4, P3). **c)** Superposition of AlphaFold3 models for *AvrPstB48* and its paralogs. Global backbone

similarity is high (pairwise RMSD ~0.5–1.5 Å), with most divergence at the C-terminal helical region.

Predicted model confidences (pTM ~0.50–0.57) indicate hypothesis-generating structures rather than

definitive models. Colour key: *AvrPstB48* (green), *P1* (orange), *P2/4* (pink), *P3* (light blue). The N-

terminus are represented by green arrows and the C-terminus by red arrows, and includes *AvrPstB48*

Eric C. Pereira
MPMI

(C48) paralog (PX) labelling. **d)** Expression profile of *AvrPstB48* and its paralogs in ungerminated

urediniospores (UG) and infected wheat leaves at 4, 6, 8, 10 and 12 days post-infection (dpi) *in planta*.

RNA-seq abundance was normalised using DESeq's median of ratio method and plotted as box plots

with four biological replicates. Analysis of variance (ANOVA) was performed using a two-way ANOVA

with sampling condition (UG, 4, 6, 8, 10 and 12 dpi) and effector identity (*AvrPstB48, P1, P2, P3* and

P4) as fixed factors, followed by post hoc Tukey's Honestly Significant Difference (HSD) test. Sampling

condition –effector combinations marked with distinct letters indicate that they are significantly different

($p < 0.05$) according to Tukey's post hoc multiple-comparisons test. Boxes represent the interquartile

range with the horizontal line inside the box indicating the median.

Figure 3 – Differential activation of defence reporters by *AvrPstB48* reveals genotype-dependent

recognition. Normalised luminescence signals from dual luciferase-based reporter assays using three

promoter constructs (pD14, pD2, pD15) and UBI-pRedf in **a)** Avocet S and **b)** Morocco. *AvrSr50* and

AvrSr50:Sr50 were used as negative and positive controls, respectively. One-way analysis of variance

(ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used to compare

treatments. Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure 4 – Differential activation of defence reporters by *AvrPstB48* paralogs reveals genotype-

dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays

using the pD14 promoter construct and UBI-pRedf in **a)** Avocet S, **b)** Morocco and **c)** Co-expression of

AvrPstB48 and *AvrPstB48_P3* (#48:P3) and *AvrPstB48* and Empty Vector (#48:ev) at a 1:1 plasmid

ratio in Avocet S. *AvrSr50* and *AvrSr50:Sr50* were included as negative and positive controls. One-way

analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used

to compare treatments. Conditions with different letters are significantly different from each other ($p <$

0.05).

Figure 5 – A single amino acid (D134) is necessary but not sufficient for recognition of *AvrPstB48*

and its paralog *AvrPstB48_P3*. a) Predicted structures of *AvrPstB48* and *AvrPstB48*_P3 were

generated using AlphaFold3. The surface-exposed amino acid position 134 is highlighted where

AvrPstB48, *P1*, and *P2/P4* encode an aspartic acid (D), while *P3* encodes a glycine (G). **b)** Normalised

luminescence signals from dual luciferase-based reporter assays using the pD14 promoter and UBI-

pRedf. One-way analysis of variance (ANOVA) followed by the Tukey Honestly Significant Difference

(HSD) test was used to compare the treatments. Conditions with different letters are significantly

different from each other ($p < 0.05$).

Figure 6. The susceptible wheat cultivar Avocet S displays a delayed infection time course to

***Pst*104E137A- when compared to Morocco. a)** Leaf phenotypes at 8 and 10 days post-infection (dpi).

b) Representative confocal micrographs showing the extent of infection inside the leaves at 8 and 10

dpi. Magenta corresponds to propidium iodide-stained cell walls, and green to fungal structures stained

with wheat germ agglutinin. Bars = 200 μm. **c)** Representative confocal micrographs showing callose

deposition (cyan) associated with infectious hyphae (magenta). Bars = 25 μm. **d)** Infection progression

and **e)** callose deposition through time in infected leaves. Asterisks denote significant differences

according to Welch's t-test at p-value < 0.01 (**) and p-value < 0.001 (***).

Supplementary Figures

Figure S1 - Differential activation of defence reporters by *AvrPstB48* reveals genotype-

dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays

using three promoter constructs (pD14, pD2, pD15) and UBI-pRedf in **a)** Triticum spelta *Yr5*, **b)** Gabo

and **c)** Chinese 166. AvrSr50 and AvrSr50:Sr50 were used as negative and positive controls,

respectively. One-way analysis of variance (ANOVA) followed by Tukey's Honestly Significant

Difference (HSD) test was used to compare treatments. Conditions with different letters are significantly

different from each other ($p < 0.05$).

Figure S2 - Differential activation of defence reporters by *AvrPstB48* paralogs reveals genotype-

dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays

using the pD14 promoter construct and UBI-pRedf in **a)** *Triticum spelta Yr5*, **b)** Gabo and **c)** Chinese

166. AvrSr50 and AvrSr50:Sr50 were included as negative and positive controls, respectively. One-way

analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used

to compare treatments. Conditions with different letters are significantly different from each other ($p <$

0.05).

Figure S3 – Endpoint RT-PCR confirmation of transgene expression in wheat protoplasts. Total

RNA from transfected protoplasts was DNase-treated and reverse-transcribed. Gene-specific primers

amplified *AvrPstB48*, *AvrPstB48_P3*, and wheat 18S rRNA (Ta 18s; loading control). Five microlitres of

each amplicon were resolved on a 1% agarose gel and visualised under UV. M, 1 kb Plus DNA Ladder

(NEB #N3200).

Figure S4 – Infection types of selected wheat cultivars inoculated with Pst104E137A–. Infection

types (IT) on a 0-9 scale on the wheat cultivars *Triticum spelta Yr5*, Avocet *S*, Chinese 166, Morocco

and Gabo at 12 days post-infection (DPI). 0: no visible disease symptoms (immune), 1: minor chlorotic

and necrotic flecks, 2: chlorotic and necrotic flecks without sporulation, 3–4: chlorotic and necrotic areas

with limited sporulation, 5–6: chlorotic and necrotic areas with moderate sporulation, 7: abundant

sporulation with moderate chlorosis, 8–9: abundant and dense sporulation without notable chlorosis

and necrosis.

Supplementary Tables

Table S1 - Comparative sequence alignment and AlphaFold modelling metrics for AvrPstB48 and

paralogs *(P1, P2/P4, P3*). The AlphaFold metrics include ipTM (interface predicted TM-score) and

pTM (predicted TM-score), which estimate confidence in the predicted structures and inter-domain

orientations, with higher values indicating greater reliability. The "Zn" column indicates whether zinc

ions were included during modelling, as zinc can stabilise cysteine-rich proteins and influence predicted

folding.

Table S2 – Primers used for Avr insert verification and RT–PCR. Primer names, functions and

sequences used to confirm the presence of Avr inserts (DB34, DB35) after cloning and for cDNA

amplification of AvrPstB48, AvrPstB48_P3 and the wheat reference gene Ta18S in RT–PCR.

Table S3 - List of plasmids.

Table S4 - List of Avirulence (Avr) screening in wheat protoplasts of various differential wheat

lines to *Puccinia striiformis* f. sp. *tritici* (*Pst*) Pst104E137A-infection in a protoplast-based

defence reporter assay.

Table S5 – Resume of promoter activation triggered by *AvrPstB48* and Infection types (IT) on a

0-9 scale for the isolate 104E137A- inoculated in different cultivars. Response of various

differential wheat lines to *Puccinia striiformis* f. sp. *tritici* (*Pst*) *Pst*104E137A-infection and the induction

of *AvrPstB48* in a protoplast-based defence reporter assay. "*Pst*-resistant gene" refers to the known Yr

gene(s) present in each line. "*AvrPstB48* induction" indicates whether candidate *AvrPstB48* triggered a

defence response (+) or not (-) in the protoplast assay. "*Pst* infection *outcomes*" represents the infection

types (IT) on a 0-9 scale on the wheat cultivars at 12 days post-infection (DPI). 0: no visible disease

symptoms (immune), 1: minor chlorotic and necrotic flecks, 2: chlorotic and necrotic flecks without

sporulation, 3–4: chlorotic and necrotic areas with limited sporulation, 5–6: chlorotic and necrotic areas

with moderate sporulation, 7: abundant sporulation with moderate chlorosis, 8–9: abundant and dense

sporulation without notable chlorosis and necrosis.

Figures

Figure 1 – *pDefense* D14 promoter activation triggered by *AvrPstB48* in Avocet *Yr5* and Avocet *S* protoplasts. Dual luciferase-based reporter assays were performed using wheat protoplasts from wheat cultivars Avocet *Yr5* **a)** and Avocet *S* **b)** to assess defence signalling. The x-axis shows the plasmid combinations tested, including candidate effectors *Pst104E137_008104* (#47), *Pst104E137_010232* (*AvrPstB48*), and *Pst104E137_024713* (#49), along with the positive control pair *AvrSr50/Sr50*. The y-axis represents normalised luminescence values, measured as the ratio of luciferase activity driven by the D14 promoter to the internal control UBI-pRed. *AvrPstB48* triggered a significant increase in reporter activity, indicating activation of defence responses. Statistical differences were analysed using one-way ANOVA followed by Tukey's Honestly Significant Difference (HSD) test; Conditions with different letters are significantly different from each other ($p < 0.05$).

a)

Pst104E137_010231
(AvrPstB48_P3)

Pst104E137_010232
(AvrPstB48)

Pst104E137_010234
(AvrPstB48_P4)

Chr10A

3,960,000 - 3,985,000 bp

98.2%   57.6%   90.8%   100%

Chr10B

3,990,000 - 4,015,000 bp

Pst104E137_025766
(AvrPstB48_P1)

Pst104E137_025767
(AvrPstB48_P2)

b)

c)

d)

Figure 2 – Genomic, evolutionary, and structural characterisation of the candidate effector gene *AvrPstB48* and its paralogs. a) Genomic organisation and synteny of *AvrPstB48* (*Pst104E137_010232*) and its paralogs (P1: *Pst104E137_025766*; P2: *Pst104E137_025767*; P3: *Pst104E137_010231*; P4: *Pst104E137_010234*) across homologous regions of Chromosomes 10A and 10B, with genes shown as red or blue arrows, respectively. Grey-shaded blocks and contained numbers represent pairwise nucleotide sequence identity (%) between all paralogs. **b)** Multiple sequence alignment of *AvrPstB48* and its paralogs, showing conservation at the amino acid level. Identical or similar residues are highlighted using the Clustal X colour scheme, which assigns colours based on amino acid conservation and properties. Conserved motifs are evident across the sequences. Dashes indicate gaps in the alignment. Pink shading marks the predicted N-terminal signal peptide; orange bars denote predicted α-helical segments; red arrows denote predicted β-strands mapped from the AlphaFold3 structural model onto the multiple-sequence alignment of *AvrPstB48* and its paralogs (*P1, P2/P4, P3*). **c)** Superposition of AlphaFold3 models for *AvrPstB48* and its paralogs. Global backbone similarity is high (pairwise RMSD ~0.5–1.5 Å), with most divergence at the C-terminal helical region. Predicted model confidences (pTM ~0.50–0.57) indicate hypothesis-generating structures rather than definitive models. Colour key: *AvrPstB48* (green), *P1* (orange), *P2/4* (pink), *P3* (light blue). The N-terminus are represented by green arrows and the C-terminus by red arrows, and includes *AvrPstB48* (C48) paralog (PX) labelling. **d)** Expression profile of *AvrPstB48* and its paralogs in ungerminated urediniospores (UG) and infected wheat leaves at 4, 6, 8, 10 and 12 days post-infection (dpi) *in planta*. RNA-seq abundance was normalised using DESeq's median of ratio method and plotted as box plots with four biological replicates. Analysis of variance (ANOVA) was performed using a two-way ANOVA with sampling condition (UG, 4, 6, 8, 10 and 12 dpi) and effector identity (*AvrPstB48, P1, P2, P3* and *P4*) as fixed factors, followed by post hoc Tukey's Honestly Significant Difference (HSD) test. Sampling condition –effector combinations marked with distinct letters indicate that they are significantly different ($p < 0.05$) according to Tukey's post hoc multiple-comparisons test. Boxes represent the interquartile range with the horizontal line inside the box indicating the median.

Figure 3 – Differential activation of defence reporters by *AvrPstB48* reveals genotype-dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays using three promoter constructs (pD14, pD2, pD15) and UBI-pRedf in **a)** Avocet S and **b)** Morocco. *AvrSr50* and *AvrSr50:Sr50* were used as negative and positive controls, respectively. One-way analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used to compare treatments. Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure 4 – Differential activation of defence reporters by *AvrPstB48* paralogs reveals genotype-dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays using the pD14 promoter construct and UBI-pRedf in **a)** Avocet S, **b)** Morocco and **c)** Co-expression of *AvrPstB48* and *AvrPstB48_P3* (#48:P3) and *AvrPstB48* and Empty Vector (#48:ev) at a 1:1 plasmid ratio in Avocet S. *AvrSr50* and *AvrSr50:Sr50* were included as negative and positive controls. One-way

analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used to compare treatments. Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure 5 – A single amino acid (D134) is necessary but not sufficient for recognition of *AvrPstB48*

and its paralog *AvrPstB48_P3*. a) Predicted structures of *AvrPstB48* and *AvrPstB48*_P3 were

generated using AlphaFold3. The surface-exposed amino acid position 134 is highlighted where

AvrPstB48, *P1*, and *P2/P4* encode an aspartic acid (D), while P3 encodes a glycine (G). **b)** Normalised

luminescence signals from dual luciferase-based reporter assays using the pD14 promoter and UBI-

pRedf. One-way analysis of variance (ANOVA) followed by the Tukey Honestly Significant Difference (HSD) test was used to compare the treatments. Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure 6. The susceptible wheat cultivar Avocet S displays a delayed infection time course to
Pst**104E137A- when compared to Morocco. a)** Leaf phenotypes at 8 and 10 days post-infection (dpi).
b) Representative confocal micrographs showing the extent of infection inside the leaves at 8 and 10
dpi. Magenta corresponds to propidium iodide-stained cell walls, and green to fungal structures stained
with wheat germ agglutinin. Bars = 200 µm. **c)** Representative confocal micrographs showing callose
deposition (cyan) associated with infectious hyphae (magenta). Bars = 25 µm. **d)** Infection progression
and **e)** callose deposition through time in infected leaves. Asterisks denote significant differences
according to Welch's t-test at p-value < 0.01 (**) and p-value < 0.001 (***).

Supplementary Figures

Figure S1 - Differential activation of defence reporters by *AvrPstB48* reveals genotype-dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays using three promoter constructs (pD14, pD2, pD15) and UBI-pRedf in **a)** Triticum spelta *Yr5*, **b)** Gabo and **c)** Chinese 166. AvrSr50 and AvrSr50:Sr50 were used as negative and positive controls,

respectively. One-way analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used to compare treatments. Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure S2 - Differential activation of defence reporters by *AvrPstB48* paralogs reveals genotype-dependent recognition. Normalised luminescence signals from dual luciferase-based reporter assays using the pD14 promoter construct and UBI-pRedf in **a)** *Triticum spelta Yr5*, **b)** Gabo and **c)** Chinese 166. AvrSr50 and AvrSr50:Sr50 were included as negative and positive controls, respectively. One-way analysis of variance (ANOVA) followed by Tukey's Honestly Significant Difference (HSD) test was used to compare treatments. Conditions with different letters are significantly different from each other ($p < 0.05$).

Figure S3 – Endpoint RT-PCR confirmation of transgene expression in wheat protoplasts. Total RNA from transfected protoplasts was DNase-treated and reverse-transcribed. Gene-specific primers amplified *AvrPstB48*, *AvrPstB48_P3*, and wheat 18S rRNA (Ta 18s; loading control). Five microlitres of each amplicon were resolved on a 1% agarose gel and visualised under UV. M, 1 kb Plus DNA Ladder (NEB #N3200).

Figure S4 – Infection types of selected wheat cultivars inoculated with Pst104E137A–. Infection types (IT) on a 0-9 scale on the wheat cultivars *Triticum spelta Yr5*, Avocet *S*, Chinese 166, Morocco and Gabo at 12 days post-infection (DPI). 0: no visible disease symptoms (immune), 1: minor chlorotic and necrotic flecks, 2: chlorotic and necrotic flecks without sporulation, 3–4: chlorotic and necrotic areas with limited sporulation, 5–6: chlorotic and necrotic areas with moderate sporulation, 7: abundant sporulation with moderate chlorosis, 8–9: abundant and dense sporulation without notable chlorosis and necrosis.