1

2   Automatic pairing of real-world stimuli in younger and older adults: An event-

3   related potential study

5   Petia Kojouharova[1]*, István Czigler[1], Boglárka Nagy[1], Zsófia Anna Gaál[1]

7   [1]Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for
8   Natural Sciences, Budapest, Hungary
9

10   * Corresponding author

11   Email: kojouharova.petia@ttk.hu (PK)

Abstract

Contextual information can influence object perception, even when it is irrelevant to the task. When two unrelated stimuli frequently occur in close temporal succession, they may become associated. This process might be stronger in older adults due to age-related decline in inhibitory control, leading to deeper processing of irrelevant context. Automatic associations and their violations can be studied using visual mismatch negativity (vMMN), an event-related potential component reflecting the detection of violations of regularity in unattended stimuli.

In our study, younger (n=18, M=21.2 ± 2.1 yrs) and older (n=17, M=69.8 ± 2.8 yrs) adults viewed pairs of images presented in succession: a scene (forest or street) followed by an emotional face (happy or angry). One scene–emotional face combination occurred frequently, the other rarely, forming a contextual oddball sequence. All individual stimuli appeared equally often. Participants performed an unrelated colour-change detection task, ensuring that the scene–emotional face pairs were unattended. We expected a vMMN to the emotional face in rare pairings, especially in older adults if they formed stronger associations. A control oddball condition with only emotional faces verified automatic emotion discrimination.

In the control condition, vMMN emerged for both deviant emotions (in various ranges within the 94–276 ms post-stimulus time window). However, no vMMN was observed in the scene-emotional face pair condition. Instead, younger adults showed an early posterior positivity (90–161 ms) to rare pairings, while older adults exhibited a later negativity (356–384 ms). These results suggest that task-irrelevant, thematically unrelated visual events can become associated through temporal proximity. However, violations of these associations evoke neural responses distinct from vMMN and vary across age groups, with older adults relying on later, potentially semantic-related mechanisms rather than early visual processes to detect the regularity.

Introduction

Task-irrelevant information such as context may implicitly link to a target stimulus, facilitating or impeding our performance on a task. When searching for a familiar pub in a town we have visited before, the spatial arrangement of landmarks can help us, while a recently demolished building might mislead us. Similarly, anticipating sequential events – such as noticing the gradual darkening of clouds and unconsciously expecting rain – can support our behaviour. While the neural mechanisms of spatial context in guiding behaviour are well-documented, much less is known about whether consecutive unattended stimuli become automatically associated. This may be particularly relevant for older adults, for whom certain cognitive functions decline with age, but contextual information and automatic processes could serve as compensatory mechanisms. In this study, we investigated whether an association can form between two thematically unrelated, consecutively presented visual events that are not attended, and examined the electrophysiological correlates of violating such an association and tested whether these processes differ between younger and older adults.

In real-world settings, it is rare for objects to appear without any spatial or temporal context. A large body of research has demonstrated that regular spatial context has a considerable effect on object identification (e.g., [1,2]). Similarly, temporal regularity is helpful in the processing of objects and their features. Applying neurophysiological methods in studies investigating the effects of expected (regular) vs. surprising (irregular) successive attended events, surprising events elicited increased BOLD activity (e.g., [3]). In ERP studies where face stimuli were used, unpredictable alternating stimuli elicited larger responses than predictable ones (e.g., [4]), underscoring the presence of an expectation effect. Temporal regularity facilitates object processing even when such regularities are task-irrelevant (i.e. outside the focus of attention). Electrophysiological results from both auditory and visual modalities show that stimuli violating temporal regularities (i.e., deviant events) elicit distinct

event-related responses – namely, the auditory mismatch negativity (MMN, for a review see

[5]) and the visual mismatch response (vMMR, or visual mismatch negativity, vMMN, when

the response is negative, for reviews see [6,7]).

MMN and vMMN are usually investigated using the passive oddball paradigm. In the

visual version of this paradigm, standard events are constructed from stimuli with identical

features or from stimuli belonging to the same category, and deviant events violate this

regularity by differing in features or category. According to the predictive coding framework,

(e.g., [7,8]), standard stimuli establish a memory-based prediction of expected stimuli, and the

MMN/vMMN reflects a cascade of processing triggered by the detection of a surprising deviant

event.

In the present study, our aim was to investigate a more complex form of temporal

regularity: a particular scene (either a forest or a street) consistently preceded and was paired

with faces expressing a specific emotion (happiness or anger). However, the pairings were

unbalanced: one combination (e.g., forest scenes with happy faces and street scenes with angry

faces) occurred frequently and thus served as the standards, while the reverse combination (e.g.,

forest scenes with angry faces and street scenes with happy faces) occurred rarely and served

as the deviants. Together, these frequent and rare pairings formed a contextual oddball

sequence (scene-emotional face pair oddball). Importantly, both the scenes and the emotional

faces were task-irrelevant, meaning that any association formed between them would have

occurred automatically. A vMMN-like response to these infrequent, reversed pairings would

indicate the automatic detection of an unexpected event. Previous research has shown that rare

conjunctions of two features (i.e. deviant objects) can elicit a vMMN response [9]. The present

study goes a step further: here the deviant conjunction is distributed across two sequential,

thematically distinct visual events. So, we compared event-related potentials (ERPs) elicited

by face stimuli in the frequent pairings (e.g., expected happy face after the forest scene) versus

the rare pairings (e.g., unexpected happy face after the street scene).

To examine potential age-related differences in automatic temporal association

learning, we compared older and younger participants. Both the formation of automatic

associations between events and the detection of violation of such associations are important

for adaptive behaviour. While many cognitive processes show age-related decline (e.g., [10–

14]), various types of implicit learning are thought to remain relatively preserved in older adults

(but see [15,16]). For example, results from studies using the contextual cueing paradigm

suggest that this type of automatic learning is preserved with age [17–20]. When impairments

have been reported [21], they were often attributed to a slower learning rate in older adults

[22,23], or to the fact that older adults tend to acquire only a subset of repeated contexts [23].

Moreover, Yao and colleagues [23] found that, when tested at least four weeks after initial

exposure, older adults retained contextual memory more robustly than younger adults, who

required re-learning. At the same time, the two age groups may rely on different mechanisms

to perform the task. While the spatial contextual cueing effect was behaviourally similar across

age groups, underlying neural mechanisms differed in Kojouharova et al.'s [18] experiment: in

younger adults, the effect was driven by enhanced early attention and categorization processes,

whereas in older adults, it was associated with enhanced late, response-related activity.

As a control, we also included a traditional passive oddball condition using the same

emotional faces from the contextual oddball task, to examine whether participants

automatically discriminated between the two facial emotions (emotional-face-only oddball.

Facial emotions are frequently and successfully applied in vMMN research (e.g., [24–26]).

However, to the best of our knowledge, no study has yet investigated vMMN responses to

emotional faces in older adults. As with other types of stimuli, vMMN results show that the

component is intact, though often diminished in older adults compared to younger adults; for

example, for checkerboard stimuli [27], and disappearing events [28], or see Kremláček et al.

[29] for a review. These results were interpreted as indicating that younger adults process the

deviant stimuli more extensively. However, when the integration of successively presented

fragments of pseudo-letters was required to detect deviance, with a longer delay between the

members of the pairs, older adults performed better than younger adults [30]. This result

suggests that older adults' memory representation of the fragments persists for a longer time.

In the present study, we anticipated the following outcomes. First, if scenes and

emotional faces are automatically associated, ERPs should differ for faces in the frequently

paired scene-emotional face combinations compared to the same faces appearing in rare

pairings. Second, if older adults process context and unattended faces more effectively due to

impaired inhibitory control, the association (i.e. context effect) should be more likely to

develop, and ERPs related to unexpected faces should emerge with greater amplitude. Finally,

we expected vMMN to be elicited in both age groups for both emotional expressions in the

emotional-face-only oddball paradigm. However, further comparisons between age groups or

emotional expressions were beyond the scope of the present study.

Methods

130   The data that support the findings of this study are openly available in OSF at

131 https://doi.org/10.17605/OSF.IO/UNDJA.

Participants

Twenty-six younger and twenty-three older adults participated in the experiment. The

younger adults were recruited through a school cooperative, while the older adults were

recruited from our own database. All participants received payment for their participation.

Three older participants did not complete the experiment. The data of five younger and three

older participants were excluded from analysis because of noisy EEG data. Thus, the final

sample consisted of 18 participants in the younger group (13 women, 1 transgender man,

M=21.17 years, SD=2.06, 2 left-handed) and 17 participants in the older group (10 women,

M=69.82 years, SD=2.81, all right-handed). All had normal or corrected-to-normal vision and

no reported neurological or psychiatric disorders. The size of the sample was determined by

using G*Power [31] and was based on a previous study of vMMN to emotional faces. In

Stefanics et al. [26], vMMN was elicited posteriorly to deviant emotional (happy and fearful)

faces in three post-stimulus time windows. For each time window, η^2 was reported as the effect

size for the main effect of stimulus (standard or deviant): 70-120 ms: $\eta^2 = 0.17$ (for fearful

faces only), 170-220 ms: $\eta^2 = 0.32$, and 250-360 ms: $\eta^2 = 0.39$. Thus, taking into account the

lowest effect size of $\eta^2 = 0.17$, a sample of 12 participants is required for power = 80%, alpha

= 0.05.

To exclude dementia-related differences between the age groups, we measured

intelligence with four subtests of the Hungarian version of Wechsler Adult Intelligence Scale

(WAIS IV [34]) representing the four major components: Similarities – verbal comprehension;

Digit Span – working memory; Matrix Reasoning – perceptual reasoning; Coding – processing

speed. The scaled scores (where the age-group average is 10) achieved by the younger group

were as follows: Similarities: $M = 13.5$ ($SD = 2.03$); Digit Span: $M = 10.06$ ($SD = 3.14$); Matrix

Reasoning: $M = 11.06$ ($SD = 1.78$); Coding: $M = 11.22$ ($SD = 1.36$). The older group achieved

the following scores on the four subtests: Similarities: $M = 13.88$ ($SD = 2.19$); Digit Span: M

$= 12.35$ ($SD = 2.83$); Matrix Reasoning: $M = 11.65$ ($SD = 2.99$); Coding: $M = 14.59$ ($SD =$

2.59).

Ethics statement

The study was approved by the United Ethical Review Committee for Research in Psychology (Hungary). Data collection was conducted from 8th December, 2022 until 29th June, 2023. Written informed consent was obtained from all individual participants included in the study.

Stimuli and procedure

A summary of the stimuli and the procedure can be found on Fig 1.

Fig 1. Stimuli and procedure. (A) Stimuli presented in the scene-emotional face pair oddball and in the emotional-face-only oddball. (B) Presentation of the stimuli in the scene-emotional face pair and in the emotional-face-only oddballs. The emotional face images presented on the figure were obtained from the FACES database [33].

The participants performed a task while task-irrelevant stimulus pairs were presented in sequences on a grey background (RGB 0.5, 0.5, 0.5, 45.8 cd/m^2). The stimuli were presented on a 24" monitor (ASUS VG245, resolution 1920×1080 px, refresh rate 60 Hz) located at a viewing distance of 140 cm. The task-relevant stimulus was a continuously present coloured rectangle frame (measuring 496×334 px and 5.6° × 3.7° of visual angle, line width 10 px) centred on the screen. The frame switched colours between red (RGB 1, 0, 0, 42.4 cd/m^2) and green (RGB 0, 1, 0, 154.2 cd/m^2) randomly every 5 to 15 seconds, and the participant had to indicate the colour switch by pressing Space on a computer keyboard. Responses were recorded as correct only if they were within 800 ms after the colour change. Key presses outside this time frame were not recorded. The purpose of the task was to distract attention from the task-irrelevant stimuli. After each sequence the participant was given feedback about their percentage of correct responses and their reaction time.

In the first half of the experimental session the task-irrelevant stimuli were image pairs. The first stimulus of the pair was a photo of a forest or a street. The second stimulus was an emotional face displaying either happiness or anger. We used one forest scene photo and one street scene photo. Both photos (see Fig 1A) were chosen from Unsplash (https://unsplash.com/), and their size was 486×324 px (5.5° × 3.6° of visual angle). The emotional face images were obtained from photos from the FACES database [33]. There were four emotional faces for both emotions: a younger woman, a younger man, an older woman, and an older man (see Fig 1A). The size of the images was 259×324 px (2.9° × 3.6° of visual angle). The images were centred on the screen and presented within the coloured frame. The stimuli pairs were presented in oddball sequences of 80 pairs. In each sequence each image was presented an equal number of times, i.e., on their own, the context forest and street images and the emotional happy and angry faces had a 0.5 probability. However, there were frequent

(standard) pairs with a 0.8 probability and rare (deviant) pairs with a 0.2 probability. Half of

the participants saw the forest-happy face and street-angry face as the standard pairs and the

forest-angry face and street-happy face as the deviant pairs. For the other half of the participants

the pairs' roles were reversed. A sequence always started with 6 standard pairs (3 of each).

There were 10 sequences (presented in five blocks) with 64 standard pairs and 16 deviant pairs

per sequence for a total of 640 standard pairs and 160 deviant pairs. Between two deviant pairs

there were 2-6 standard pairs. Stimulus duration was 140 ms for both the scene and the

emotional face, while the intra-pair interval was 360±17 ms. The inter-pair interval was 720±51

ms (in up to three steps of 17 ms). Each sequence was approximately 2 minutes long.

In the second part of the session only emotional faces were presented in oddball

sequences. In half of the sequences the happy faces were the standard, and the angry faces were

the deviant stimuli, and in the rest of the sequences the roles were reversed. In other words, we

applied a reverse control where each emotional face was both a standard and a deviant. The

order was counterbalanced between participants. The probability of the standard emotional face

was 0.8, and the probability of the deviant emotional face was 0.2. A sequence always started

with 6 standards, and there were between 2 and 6 standards between deviants. There were 10

sequences (presented in five blocks) with 64 standard emotional faces and 16 deviant emotional

faces in each sequence for a total of 320 standard happy faces, 80 deviant happy faces, 320

standard angry faces, and 80 deviant angry faces. The stimulus duration was 140 ms, and the

interstimulus interval was 720±51 ms (in up to three steps of 17 ms). Each sequence was

approximately 1.2 minutes long.

The stimuli presentation and response recording were realized using the Cogent 2000

[34] and Cogent Graphics [35] toolboxes in MATLAB [36].

Recording and measuring of the electrophysiological activity

Brain electric activity was recorded (bandwidth: DC-70 Hz; sampling rate 1000 Hz; BrainVision actiCHamp amplifier, BrainVision Recorder, BrainVision Products GMBH) with Ag/AgCl active electrodes placed at 59 locations according to the extended 10–20 system using an elastic electrode cap (EasyCap, Brain Products GmbH). The reference electrode was placed on the nose. Eye movements were recorded with four electrodes placed around the eyes. Horizontal EOG was recorded with a bipolar configuration between electrodes positioned lateral to the outer canthi of the eyes (one electrode on each side). Vertical eye movement was monitored with a bipolar montage between two electrodes, one placed above and one below the left eye. The impedance of all electrodes was kept below 30 kΩ.

Data analysis

Behavioural analysis

Mean accuracy (percentage of hits) and mean reaction time (RT, reaction time to the hits) were calculated for each sequence for a participant, and were then averaged for the two oddball conditions separately. A mixed analysis of variance (ANOVA) was conducted for both measures with ODDBALL (scene–emotional face pair or emotional-face-only) as the within-subject factor and AGE GROUP (younger or older) as the between-subject factor to investigate whether performance on the task differed depending on age and on the task-irrelevant stimuli. Slower RT was expected in the older compared to the younger group because of age-related slowing.

ERP analysis

The EEG signal was band-pass-filtered offline with a non-causal Kaiser-windowed FIR filter (lowpass filter parameters: cutoff frequency of 30 Hz, beta of 12.2653, a transition band of 10 Hz; highpass filter parameters: cutoff frequency of 0.1 Hz, beta of 5.6533, a transition band of 0.2 Hz). Stimulus-locked epochs were extracted for each emotional face in both conditions with a duration of 900 ms, including a 100 ms pre-stimulus interval.

In the case of the scene-emotional face pair oddball, standard epochs were the ones for the emotional faces that were expected after the scene image (i.e. from the standard pairs), and deviant epochs were the emotional faces that were unexpected after the scene image (i.e. from the deviant pairs). To equate the number of epochs for standards and deviants, only emotional faces from the standard pairs preceding deviant pairs were used for averaging. Because brain activity from the preceding (scene) stimulus did not return to baseline within the allotted 500 ms to allow for the use of the 100 ms pre-stimulus interval as a baseline, the 700-800 ms post-stimulus interval was used as a baseline instead.

In the case of the emotional-face-only oddball, epochs were extracted for the standard happy, standard angry, deviant happy, and deviant angry emotional faces. To equate the number of epochs for standards and deviants, only standard emotional faces preceding a deviant emotional face were used for averaging. Here, the 100 ms pre-stimulus interval served as the baseline.

The monitor lag (the time difference between the trigger and the appearance of a stimulus on the screen) was measured by a photodiode before the experiment. The lag was the same across events (67 ms), and thus the latencies of the events were corrected in the data after data collection. Epochs with an amplitude change exceeding 100 µV on any posterior (parietal, parietooccipital, or occipital) channel were excluded from further analysis. The average number of epochs for each analysis can be found in Table 1.

Table 1. Average number of epochs with standard deviations in parentheses and ranges in the second row for deviant and standard stimuli preceding the deviants in each group and condition.

		scene-emotional face pair oddball	emotional-face-only oddball	
		emotional face	happy	angry
Younger group	**deviant**	132.4 (9.4) 109-145	66.4 (5.8) 56-76	65.8 (10.5) 44-98
	standard	128.5 (11.5) 109-148	68.3 (9.3) 55-99	67.5 (4.8) 56-76
Older group	**deviant**	131.3 (12.3) 95-143	62.7 (8.9) 34-71	65.1 (8.2) 42-73
	standard	132 (14.9) 86-145	65 (8.7) 41-73	65.1 (9.6) 35-74

In the case of the scene-emotional face pair oddball, because our focus was on the second stimulus in the pair being differing from what should be expected, the epochs were pooled together independent of emotion (happiness and anger) or preceding scene (forest or street), separately for standards and deviants, and were then averaged. A difference potential was then computed by subtracting the ERP for the standards from the ERP for the deviants.

In the emotional-face-only oddball, for each participant the epochs were averaged separately for the happy standard faces, deviant happy faces, angry standard faces and deviant angry faces. Then difference potentials were computed as a deviant-minus-standard ERP separately for happy (happy deviant minus happy standard) and angry (angry deviant minus angry standard) emotional faces. The purpose of this analysis was to show that vMMN is elicited for the emotional faces used in this study, in line with previous vMMN studies.

For all difference potentials we conducted point-by-point one-sample t-tests to 0 over the 50-400 ms post-stimulus time window (which was slightly longer that the expected 100-

350 ms post-stimulus time window of vMMN) at the posterior regions of interest (ROI). This

analysis is a necessary step before a direct comparison between the groups can be performed

for the following reasons: 1) we need to establish the presence of vMMN within each group,

and 2) if vMMN is observed, we need to establish whether it is observed in different ranges in

each group – in older adults components may be in a different range (usually delayed)

compared to younger adults. The following ROIs were formed: occipital (O1, Oz, O2),

parietooccipital (PO3, POz, PO4), left parietooccipital (PO7, P7, P5), and right parietooccipital

(PO8, P8, P6). The occipital and parietooccipital electrode sites are the expected sites of

vMMN, while the left and right parietooccipital electrode sites are usually implicated in the

processing of face stimuli, including in vMMN studies. At least twenty consecutive statistically

significant negative differences (20 ms duration) at any of the four ROIs within the 50-400 ms

post-stimulus time window were considered to show the elicitation of vMMN (cf. [37]).

For any such interval observed in either of the age groups in the scene-emotional face

pair condition, we calculated mean amplitude, and then performed a repeated measures

ANOVA with ROI (occipital, parietooccipital, left parietooccipital, right parietooccipital) as a

within-subject factor and GROUP (younger, older) as a between-subject factor.

EEG data were preprocessed with MATLAB R2019b [38] and EEGLAB 2020.0 and

2024.0 [39]. JASP [40] was used for the analysis of the behavioural results.

Results

Behavioural results

Mean accuracy and RT are displayed in Table 2. There was a main effect of AGE

GROUP on performance, $F(1,33) = 6.289$, $p = 0.017$, $\eta^2_p = 0.16$, with lower performance in

the older group. The main effect of ODDBALL, $F(1,33) = 1.806$, $p = 0.188$, $\eta^2_p = 0.052$, and

the interaction, $F(1,33) = 1.046$, $p = 0.314$, $\eta^2_p = 0.031$, were not significant. While the older

adults had lower accuracy in the task than the younger adults, their performance was still above

90% which suggests adequate task engagement. The older adults were overall slower than the

younger adults, $F(1,33) = 6.09$, $p = 0.019$, $\eta^2_p = 0.156$. Furthermore, both groups responded

faster in the emotional-face-only oddball condition than in the scene-emotional face pair

oddball condition, $F(1,33) = 14.394$, $p < 0.001$, $\eta^2_p = 0.304$. The interaction was not significant,

$F(1,33) = 0.003$, $p = 0.96$, $\eta^2_p < 0.001$. Overall slower RTs are the norm for older adults

compared to younger adults when performing the same task. An additional analysis with

BLOCK (10 Blocks) as a within-subject factor and GROUP as a between-subject factor

revealed a main effect of BLOCK, $F(3.945,130.186) = 6.05$, $p < 0.001$, $\eta^2_p = 0.155$, but no

BLOCK×GROUP interaction, $F(3.945,130.186) = 1.457$, $p = 0.22$, $\eta^2_p = 0.042$, suggesting that

the difference between the oddball conditions was due to an improvement in performance

during the session as the oddball conditions were always presented in the same order.

Table 2. Mean accuracy and mean reaction time (RTs) for each group and condition.

	Mean accuracy (SD) in percentage		Mean RT (SD) in ms	
	scene-emotional face pair oddball	emotional-face-only oddball	scene-emotional face pair oddball	emotional-face-only oddball
Younger group	97.75% (0.04)	97.46% (0.02)	440.2 (35.3)	420.5 (48.3)
Older group	94.3% (0.07)	92.15% (0.08)	479.7 (48.5)	459.5 (58.4)

ERP results

Figs 2-4 show the ERPs and the difference potentials in both in the scene-emotional

face pair oddball and in the emotional-face-only oddball conditions. Latency ranges with

significant difference (20 consecutive difference at significant t-tests at a posterior ROI) are

marked in the difference potentials. Scalp distributions in these ranges are also shown on Figs

2-4. As the deviant-minus-standard difference potentials indicates, in the scene-emotional face

pair oddball condition in the younger group the only difference emerged in the 90-161 ms

range, as a positivity. In the older group a negative difference appeared in the 356-384 ms

range. Table 3 shows the ranges in the various ROIs.

The difference potentials differed significantly between the groups in the 90-161 ms

time interval, $F(1,33) = 4.583$, $p = 0.04$, $\eta^2_p = 0.122$. There was no main effect of ROI, $F(3,99)$

$= 1.598$, $p = 0.195$, $\eta^2_p = 0.046$, and no interaction, $F(3,99) = 2.078$, $p = 0.108$, $\eta^2_p = 0.059$.

There were neither ROI, $F(2.217,73.172) = 0.556$, $p = 0.645$, $\eta^2_p = 0.017$, nor GROUP main

effects, $F(1,33) = 0.666$, $p = 0.42$, $\eta^2_p = 0.02$, for the 356-384 ms time interval. The

ROI×GROUP interaction was significant, $F(2.217,73.172) = 6.646$, $p = 0.002$, $\eta^2_p = 0.168$. The

post hoc tests revealed only a significant difference in mean amplitude between the

parietooccipital and the right parietooccipital ROI in the younger group ($p = 0.016$), with the

amplitude being more negative in the right parietooccipital ROI.

Fig 2. Average ERPs and deviant-minus-standard difference potentials for each group and region of interest (ROI) in the scene-emotional face pair oddball condition. Red lines depict ERPs to the deviant stimuli, light blue lines depict ERPs to the standard stimuli preceding the deviants, blue lines depict the deviant-minus-standard difference potentials, with the 95% CI interval shown by the blue shaded area. Scalp distributions for the ranges in which significant differences were found are shown on the right.

Fig 3. Average ERPs and deviant-minus-standard difference potentials for each group and region of interest (ROI) in the happy emotional-face-only oddball condition. Red lines depict ERPs to the deviant stimuli, light blue lines depict ERPs to the standard stimuli preceding the deviants, blue lines depict the deviant-minus-standard difference potentials, with the 95% CI interval shown by the blue shaded area. Scalp distributions for the ranges in which significant differences were found are shown on the right.

Fig 4. Average ERPs and deviant-minus-standard difference potentials for each group and region of interest (ROI) in the angry emotional-face-only oddball condition. Red lines depict ERPs to the deviant stimuli, light blue lines depict ERPs to the standard stimuli preceding the deviants, blue lines depict the deviant-minus-standard difference potentials, with the 95% CI interval shown by the blue shaded area. Scalp distributions for the ranges in which significant differences were found are shown on the right.

Table 3. Ranges of significant differences in the deviant-minus-standard difference

potentials for each group, condition, and region of interest (ROI).

		Occipital ROI	Parietooccipital ROI	Left parietooccipital ROI	Right parietooccipital ROI
Younger group	**scene-emotional face pair oddball**	94 - 116 ms	91 - 151 ms	90 - 119 ms, 129 - 141 ms	94 - 115 ms, 155 - 161 ms
	happy emotional-face-only oddball	260 - 276 ms	262 - 274 ms	262 - 275 ms	264 - 273 ms
	angry emotional-face-only oddball	182 - 205 ms	187 - 204 ms	179 - 215 ms	144 - 166 ms, 177 - 206 ms, 239 - 260 ms
Older group	**scene-emotional face pair oddball**	362 - 378 ms	358 - 384 ms,	356 - 369 ms,	
	happy emotional-face-only oddball	95 - 162 ms, 194 - 269 ms	94 - 164 ms, 201 - 268 ms	98 - 117 ms, 125 - 167 ms, 197 - 291 ms	127 - 156 ms, 174 - 220 ms, 227 - 269 ms
	angry emotional-face-only oddball	152 - 206 ms	153 - 207 ms	144 - 203 ms	170 - 209 ms

In the emotional-face-only condition negative deviant-minus-standard difference appeared in various ranges within 144-276 ms in the younger group, and within 94-269 ms in the older group. Details of the ranges in the various ROIs and for happy and angry deviants are shown on Table 3. The negative difference can be considered as vMMN.

Discussion

In this study, we investigated whether participants automatically form associations between two temporally linked, thematically unrelated visual events – specifically, a scene and an emotional facial expression – when these events are not task-relevant. We also examined

whether violations of these associations elicit distinct electrophysiological responses, and whether these responses differ between younger and older adults.

In the traditional oddball design using happy and angry faces, we observed ERP differences between deviant and standard stimuli: at posterior sites, deviant-related ERPs were more negative than those elicited by standard stimuli. These differences were comparable across the two age groups. In the younger group, deviant happy faces elicited only short negativity (9-16 ms), however, it was elicited at all ROIs. This result is putatively indicating the sensitivity to violated regularity of repetitive emotional category, i.e., the emergence of vMMN (e.g., [24-26]).

Contrary to the results of the traditional oddball paradigm, no unambiguously vMMN-related activity emerged in the scene-emotional face pair oddball condition. Thus, we found no evidence for the engagement of similar mechanisms when the regular pairing of scene and facial emotion was violated by a rare, deviant pairing – that is, when the face was preceded by the alternative scene. However, we cannot conclude that the regular contingency remained undetected. In the younger group, the deviant facial emotion elicited an early positivity, while in the older group, a later negativity. The emergence of early posterior positivity in response to stimuli that differ from the first element of a pair is a well-documented phenomenon, even when the varying visual features are task-irrelevant (e.g., [41–43]). A novel finding of the present study is that a similar early positivity can also be elicited by physically different, but regularly associated, visually complex events. As an age-related difference, we recorded no such early effect in the older group. However, in a later ERP time window, older adults also showed sensitivity to the violation of the association between the scene and the facial expression. One study [44] found a difference to deviant faces of old models among standard faces of young models in the 330-370 ms range in older adults, which the authors interpreted as a delayed vMMN. However, this interpretation remains debated, as responses in this latency

range may reflect higher-level evaluative processes rather than classical mismatch detection.

In the present study we did not observe a comparable delayed response to emotional faces in

the emotional-face-only condition. One possibility is that violations involving more complex

or abstract associations (e.g., age in [44], or cross-category associations in the present study)

may not elicit a canonical vMMN in older adults, but instead engage later processing stages.

At this stage of stimulus processing, it is likely that mechanisms other than predictive coding

are involved. Indeed, the detection of incongruence in this latency range is well documented in

other domains, most prominently in the N400 component, which is typically observed in

response to unexpected verbal stimuli (for a review see [45]). Age-related differences in change

detection under cognitively more complex conditions have also been observed using the

reference-back working memory paradigm [46]. The study reported a late anterior positivity in

younger adults, and an even later posterior positivity in older adults on certain types of deviant

trials.

What are the fundamental differences between the traditional passive oddball paradigm

– where vMMN typically emerges even when the deviant is a rare conjunction of features such

as colour and grating orientation (e.g., [9]) – and the current paradigm, in which no vMMN

was observed? The simplest explanation could be the lack of an association between the scene

and the emotional face in the absence of task relevance or attention. However, this explanation

is challenged by the early positivity observed in the younger group, and the later negativity

observed in the older group, both indicating that the deviant pairing was processed differently.

From the perspective of predictive coding accounts, one might expect vMMN-like activity to

emerge, given that the deviant facial emotion was unexpected. However, according to more

nuanced interpretations of the predictive theory (e.g., [7]), vMMN reflects a process involved

in the identification – or attempted identification – of surprising events (whether successful or

not, see [46]). In the scene-emotional face pairing condition, while the deviant emotional face

was indeed unpredicted, it belonged to a stimulus category (emotional faces) presented with

equal overall probability. Thus, its identification did not require the full cascade of processes

hypothesised by predictive coding theory as necessary for eliciting vMMN. The early positivity

has been interpreted as reflecting feature-specific, memory-based change detection, possibly

related to the vMMN [41–43], i.e. in the younger group an early mismatch between expected

features was sufficient to signal a violation of the association. Another important consideration

is the nature of the manipulated regularity. VMMN in the context of complex stimuli remains

largely unexplored (e.g., [47]). Unlike classical vMMN paradigms, the scene-emotional face

pair condition involved category-level alternations between complex stimuli, which may not

give rise to stable low-level predictions required for eliciting a classical vMMN. These

alternations may engage higher-level categorical or semantic processing, not captured by early

vMMN components. This may be reflected in the later negativity in older adults, where the

association appears to have been encoded as a more complex regularity. This may explain the

absence of a vMMN-like response in our data.

To summarize the present results, our findings suggest that frequent pairings of two

thematically unrelated visual events can become associated, even when these events are not

task-relevant. However, the ERP response to the violation of such associations differs from the

neural mechanisms underlying the detection of these violations appear to differ between older

and younger adults.

Acknowledgements

463 We thank Krisztina Czinkóczi for her technical assistance.

References

1. Biederman I, Mezzanotte RJ, Rabinowitz JC. Scene perception: Detecting and judging objects undergoing relational violations. Cognit Psychol. 1982;14: 143–177. doi:10.1016/0010-0285(82)90007-x

2. Zhu W, Drewes J. Effects of object-scene congruency with and without awareness. J Vis. 2025;25: 3. doi:10.1167/jov.25.7.3

3. Amado C, Hermann P, Kovács P, Grotheer M, Vidnyánszky Z, Kovács G. The contribution of surprise to the prediction based modulation of fMRI responses. Neuropsychologia. 2016;84: 105–112. doi:10.1016/j.neuropsychologia.2016.02.003

4. Feuerriegel D, Churches O, Coussens S, Keage HAD. Evidence for spatiotemporally distinct effects of image repetition and perceptual expectations as measured by event-related potentials. NeuroImage. 2018;169: 94–105. doi:10.1016/j.neuroimage.2017.12.029

5. Schröger E. Mismatch Negativity: A Microphone into Auditory Memory. J Psychophysiol. 2007;21: 138–146. doi:10.1027/0269-8803.21.34.138

6. Czigler I, Kojouharova P. Visual Mismatch Negativity: A Mini-Review of Non-pathological Studies With Special Populations and Stimuli. Front Hum Neurosci. 2022;15: 78123. doi:10.3389/fnhum.2021.781234

7. Stefanics G, Kremláček J, Czigler I. Visual mismatch negativity: a predictive coding view. Front Hum Neurosci. 2014;8. doi:10.3389/fnhum.2014.00666

8. Garrido MI, Kilner JM, Stephan KE, Friston KJ. The mismatch negativity: A review of underlying mechanisms. Clin Neurophysiol. 2009;120: 453–463. doi:10.1016/j.clinph.2008.11.029

488 9. Winkler I, Czigler I, Sussman E, Horváth J, Balázs L. Preattentive Binding of Auditory
489   and Visual Stimulus Features. J Cogn Neurosci. 2005;17: 320–339.
490   doi:10.1162/0898929053124866

491 10. Cabeza R. Hemispheric asymmetry reduction in older adults: The HAROLD model.
492   Psychol Aging. 2002;17: 85–100. doi:10.1037/0882-7974.17.1.85

493 11. Craik FIM, Byrd M. Aging and Cognitive Deficits. In: Craik FIM, Trehub S, editors.
494   Aging and Cognitive Processes. Boston, MA: Springer US; 1982. pp. 191–211.
495   doi:10.1007/978-1-4684-4178-9_11

496 12. Davis SW, Dennis NA, Daselaar SM, Fleck MS, Cabeza R. Que PASA? The Posterior-
497   Anterior Shift in Aging. Cereb Cortex. 2008;18: 1201–1209. doi:10.1093/cercor/bhm155

498 13. Reuter-Lorenz PA, Park DC. How Does it STAC Up? Revisiting the Scaffolding Theory
499   of Aging and Cognition. Neuropsychol Rev. 2014;24: 355–370. doi:10.1007/s11065-014-
500   9270-9

501 14. Salthouse TA. The processing-speed theory of adult age differences in cognition. Psychol
502   Rev. 1996;103: 403–428. doi:10.1037/0033-295X.103.3.403

503 15. Rieckmann A, Bäckman L. Implicit Learning in Aging: Extant Patterns and New
504   Directions. Neuropsychol Rev. 2009;19: 490–503. doi:10.1007/s11065-009-9117-y

505 16. Ward EV, Shanks DR. Implicit Memory and Cognitive Aging. Oxford Research
506   Encyclopedia of Psychology. Oxford University Press; 2018.
507   doi:10.1093/acrefore/9780190236557.013.378

508 17. Howard JH, Howard DV, Dennis NA, Yankovich H, Vaidya CJ. Implicit Spatial
509   Contextual Learning in Healthy Aging. Neuropsychology. 2004;18: 124–134.
510   doi:10.1037/0894-4105.18.1.124

511 18. Kojouharova P, Nagy B, Czigler I, Gaál ZA. Mechanisms of spatial contextual cueing in
512   younger and older adults. Psychophysiology. 2023;60: e14361. doi:10.1111/psyp.14361

19.   Merrill EC, Conners FA, Roskos B, Klinger MR, Klinger LG. Contextual Cueing Effects

Across   the   Lifespan.   J   Genet   Psychol.   2013;174:   387–402.

doi:10.1080/00221325.2012.694919

20.   Nagy B, Kojouharova P, Protzner AB, Gaál ZA. Investigating the Effect of Contextual

Cueing with Face Stimuli on Electrophysiological Measures in Younger and Older

Adults. J Cogn Neurosci. 2024;36: 776–799. doi:10.1162/jocn_a_02135

21.   Smyth AC, Shanks DR. Aging and implicit learning: Explorations in contextual cuing.

Psychol Aging. 2011;26: 127–132. doi:10.1037/a0022014

22.   Preuschhof C, Sharifian F, Rosenblum L, Pohl TM, Pollmann S. Contextual cueing in

older adults: Slow initial learning but flexible use of distractor configurations. Vis Cogn.

2019;27: 563–575. doi:10.1080/13506285.2019.1668516

23.   Yao Y, Luo R, Fan C, Qian Y, Zang X. Age-related contextual cueing features are more

evident in reaction variability than in reaction time. Q J Exp Psychol. 2024;

17470218241241954. doi:10.1177/17470218241241954

24.   Astikainen P, Hietanen JK. Event-related potentials to task-irrelevant changes in facial

expressions. Behav Brain Funct. 2009;5: 30. doi:10.1186/1744-9081-5-30

25.   Kreegipuu K, Kuldkepp N, Sibolt O, Toom M, Allik J, Näätänen R. vMMN for schematic

faces: automatic detection of change in emotional expression. Front Hum Neurosci.

2013;7. doi:10.3389/fnhum.2013.00714

26.   Stefanics G, Csukly G, Komlósi S, Czobor P, Czigler I. Processing of unattended facial

emotions: A visual mismatch negativity study. NeuroImage. 2012;59: 3042–3049.

doi:10.1016/j.neuroimage.2011.10.041

27.   Sulykos I, Gaál ZA, Czigler I. Automatic Change Detection in Older and Younger

Women: A Visual Mismatch Negativity Study. Gerontology. 2018;64: 318–325.

doi:10.1159/000488588

28. Sulykos I, Gaál ZA, Czigler I. Visual mismatch negativity to vanishing parts of objects in
younger and older adults. Malmierca MS, editor. PLOS ONE. 2017;12: e0188929.
doi:10.1371/journal.pone.0188929

29. Kremláček J, Kreegipuu K, Tales A, Astikainen P, Põldver N, Näätänen R, et al. Visual
mismatch negativity (vMMN): A review and meta-analysis of studies in psychiatric and
neurological disorders. Cortex. 2016;80: 76–112. doi:10.1016/j.cortex.2016.03.017

30. Gaál ZA, Bodnár F, Czigler I. When Elderly Outperform Young Adults—Integration in
Vision Revealed by the Visual Mismatch Negativity. Front Aging Neurosci. 2017;9.
doi:10.3389/fnagi.2017.00015

31. Faul F, Erdfelder E, Lang A-G, Buchner A. G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods.
2007;39: 175–191. doi:10.3758/BF03193146

32. Rózsa S, Kő N, Kuncz E, Mészáros A, Mlinkó R. WAIS-IV. Wechsler Adult Intelligence
Scale—Fourth Edition. Tesztfelvételi és pontozási kézikönyv. 2010.

33. Ebner NC, Riediger M, Lindenberger U. FACES—A database of facial expressions in
young, middle-aged, and older women and men: Development and validation. Behav Res
Methods. 2010;42: 351–362. doi:10.3758/BRM.42.1.351

34. Cogent 2000 team. Cogent 2000. Wellcome Department of Imaging Neuroscience, LON;
2003. Available: http://www.vislab.ucl.ac.uk/cogent.php

35. Romaya J. Cogent Graphics. Wellcome Department of Imaging Neuroscience, LON;
2008. Available: http://www.vislab.ucl.ac.uk/cogent.php

36. MathWorks, Inc. MATLAB. Natick, Massachusetts, United States; 2014.

37. Guthrie D, Buchwald JS. Significance testing of difference potentials. Psychophysiology.
1991;28: 240–244. doi:10.1111/j.1469-8986.1991.tb00417.x

38. MathWorks, Inc. MATLAB. Natick, Massachusetts, United States; 2019.

39.   Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG

dynamics including independent component analysis. J Neurosci Methods. 2004;134: 9–

21. doi:10.1016/j.jneumeth.2003.10.009

40.   JASP Team. JASP. 2023. Available: https://jasp-stats.org/

41.   Kimura M, Katayama J, Murohashi H. Positive difference in ERPs reflects independent

processing of visual changes. Psychophysiology. 2005;42: 369–379. doi:10.1111/j.1469-

8986.2005.00297.x

42.   Kimura M, Katayama J, Murohashi H. An ERP study of visual change detection: Effects

of magnitude of spatial frequency changes on the change-related posterior positivity. Int

J Psychophysiol. 2006;62: 14–23. doi:10.1016/j.ijpsycho.2005.11.005

43.   Wang Y, Tian S, Wang H, Cui L, Zhang Y, Zhang X. Event-related potentials evoked by

multi-feature conflict under different attentive conditions. Exp Brain Res. 2003;148: 451–

457. doi:10.1007/s00221-002-1319-y

44.   Csizmadia P, Petro B, Kojouharova P, Gaál ZA, Scheiling K, Nagy B, et al. Older Adults

Automatically Detect Age of Older Adults' Photographs: A Visual Mismatch Negativity

Study. Front Hum Neurosci. 2021;15: 707702. doi:10.3389/fnhum.2021.707702

45.   Kutas M, Federmeier KD. Thirty Years and Counting: Finding Meaning in the N400

Component of the Event-Related Brain Potential (ERP). Annu Rev Psychol. 2011;62:

621–647. doi:10.1146/annurev.psych.093008.131123

46.   Gaál ZA, Nagy B, Czigler I, Csizmadia P, Petró B, Kojouharova P. Decomposing working

memory subprocesses with the reference-back paradigm: Event-related potentials and

age-related differences. Wang J, editor. PLOS ONE. 2024;19: e0307351.

doi:10.1371/journal.pone.0307351

47.   Kojouharova P, File D, Sulykos I, Czigler I. Visual mismatch negativity and stimulus-

587   specific adaptation: the role of stimulus complexity. Exp Brain Res. 2019 [cited 24 Mar

588   2019]. doi:10.1007/s00221-019-05494-2