From leaves to defenders: how the amount and dispersion of leaf

damage affect extrafloral nectar production and ant-mediated

protection in wild cotton

Abstract

1. Extrafloral nectar (EFN) drives mutualistic interactions between plants and ants.

However, EFN production is costly, and its induction is predicted to vary with herbivore-

related factors. We examine how the amount and within-plant damage spatial uniformity

(i.e., dispersed and concentrated damage) affect EFN production and ant-mediated

defence in wild cotton (*Gossypium hirsutum*). Previous work suggests that severe and

concentrated damage is costliest to plants, leading us to hypothesise that such damage

would induce greater EFN production and, consequently, stronger ant attraction and

protection.

2. We conducted a greenhouse experiment in which plants were subjected to one of the

following mechanical damage treatments: control (no damage), low damage (two

damaged leaves, 30% of area removed each), high concentrated damage (two damaged

leaves, 60% each), and high dispersed damage (four damaged leaves, 30% each; i.e.,

same total area than concentrated damage). The day after treatments, we measured EFN

volume and sugar content from one nectary per plant (a per-nectary EFN response), and

the number of secreting nectaries per plant (a proxy for whole-plant level EFN response).

Subsequently, we performed a field experiment using the same treatments to evaluate

EFN-mediated ant recruitment and ant-provided protection by placing caterpillars on

plants.

3. Both high-damage treatments increased EFN volume per nectary, whereas low damage

did not. However, against our prediction, dispersed damage was the only treatment that

increased the number of secreting nectaries. Consistently, once placed in the field, plants

with dispersed damage recruited more ants and caterpillars on them were more likely to

be attacked by ants.

4. Contrary to expectations, plants subjected to dispersed damage exhibited a greater total

investment in EFN, presumably via local induction of more nectaries per plant, and

thereby enhancing ant-mediated defence. This study highlights the importance of

herbivory intensity and within-plant spatial uniformity in shaping ant-plant interactions.

Keywords: ant-plant interactions, predation, mutualism, secondary defence, induced defence,

herbivory variability

Introduction

Extrafloral nectaries are specialised structures present in more than 100 families of plants

(Weber & Keeler, 2012) that secrete a sugary, energy-rich solution. This secretion can occur

constitutively (always present, i.e., basal levels) on leaves and stems but is also highly inducible

(i.e., increased production over and above basal levels) in response to herbivory (Rico-Gray &

Oliveira, 2008; Heil, 2015). Extrafloral nectar (EFN) serves as an important food source for ants

and other insects, which in turn, often protect the plant by deterring or attacking herbivores (Heil

& McKey, 2003; Rico-Gray & Oliveira, 2008; Rosumek et al., 2009). However, EFN induction

is highly variable and context-dependent, shaped by both biotic and abiotic factors, and

modulated by the balance of its benefits (via indirect defence) and costs to the plant across

different ecological contexts (Bronstein, 1998; Bronstein et al., 2006; Kessler & Heil, 2011;

Marazzi et al., 2013; Interian-Aguiñaga et al., 2025). Addressing this context dependency is key

for understanding the ecological impacts and evolutionary stability of EFN-mediated

mutualisms.

Multiple herbivory-related factors regulate EFN induced responses. Commonly, EFN

induction increases in a dose-dependent manner as a function of the amount of herbivory (Rios et

al., 2008; Xu et al., 2014; Huang et al., 2015; Calixto et al., 2021), suggesting that plants can

fine-tune their defensive investment based on perceived herbivore pressure. Here studies have

found that higher levels of leaf damage often trigger increased EFN production, which in turn

attracts more ant defenders to help mitigate further herbivory (Heil et al., 2001; Ness, 2003;

Nogueira, 2025). This relationship is not always linear, with some studies reporting threshold

effects where EFN secretion significantly increases only after damage surpasses a certain level

(Pimenta et al., 2025). Similar patterns have been shown for other plant defensive traits, such as

chemical defences, where induction thresholds as well as saturation (no further increase in

induction beyond a given damage level) can constrain plant induced responses (e.g. Underwood,

2010). These findings suggest that induction of plant defences, including EFN, operates within

physiological or ecological limits. Within this context, research on EFN responses to varying

levels of herbivory intensities has mounted in recent years (e.g., Xu et al., 2014; Abdala-Roberts

et al., 2019b; Calixto et al., 2021; Interian-Aguiñaga et al., 2025; Pimenta et al., 2025), but

further studies are needed to clarify how these patterns influence ant-plant interactions.

The spatial uniformity of herbivore damage within a plant can also play a crucial role in

shaping plant fitness and the induction of defensive traits (Wetzel et al., 2023). Early work

suggests that plants tolerate dispersed damage better than concentrated damage, implying higher

fitness costs associated with the latter (Marquis, 1992; 1996; Mauricio et al., 1993). The reduced

tolerance to concentrated damage has been proposed to stem from the modular nature of plants

and physiological limitations for resource movement between modules. When damage is

concentrated on specific branches (or leaves), resource limitations in those areas cannot be easily

compensated for by other parts of the plant, leading to localised stress and reduced whole-plant

level re-growth (Marquis, 1996). In accordance, Optimal Defence Theory predicts that plants

should induce stronger defensive responses under concentrated damage compared to dispersed

damage, in order to offset the greater cost associated with more localised herbivory (Holland et

al., 2009; Moreira et al., 2012; Calixto et al., 2021). However, most research to date has focused

on tolerance mechanisms associated with sectoriality (i.e. resource reallocation constraints

between plant modules limiting regrowth or compensatory growth) whereas work examining

how damage spatial uniformity affects the induction of resistance-related traits have been largely

ignored. In one of the few exceptions. Ohnmeiss et al. (1997) found that concentrated damage

increased nicotine levels (a direct defence) to a greater extent than dispersed damage in

Nicotiana sylvestris. It has been hypothesised (Marquis, 1996) and shown empirically by at least

one study that stronger induced resistance responses to concentrated damage (e.g., local

induction) contribute to the spread of subsequent damage throughout the canopy (e.g., Karban &

Yang, 2020), reducing subsequent herbivory costs. Alternatively, defensive responses could be

modulated by damage thresholds within plant organs. Once damage exceeds critical thresholds, a

plant may no longer invest in defending a compromised organ (Wetzel et al., 2023), and instead

allocate more defences to less damaged or intact organs. These ideas remain largely untested,

and, to our knowledge, no study to date has evaluated how within-plant damage spatial

uniformity influences EFN induction and ant-plant interactions, or disentangled these effects

from other herbivory-related factors (e.g., total damage amount, see above).

Here, we used wild cotton (*Gossypium hirsutum* L.; Malvaceae), an extrafloral nectary-

bearing, myrmecophytic shrub distributed along the northern coastline of Yucatan (Mexico), to

evaluate the effect of leaf damage amount and within-plant dispersion on the production of EFN.

We also examined how these EFN induction patterns influence ant recruitment, as well as how

effectively ants deter or prey upon herbivores. Previous work with wild cotton has shown that

EFN induction enhances ant recruitment and defensive responses against herbivores (Vázquez-

Barrios et al., 2021), suggesting that EFN can provide fitness benefits via indirect defence,

though not universally (Reyes-Hernández et al., 2022). Wild cotton plants also show cotton

plants are less able to compensate (in terms of regrowth) after high levels of damage as well as in

response to concentrated (vs. dispersed) damage (Quijano-Medina et al., 2021, Ramírez-Delgado

et al., 2025), implying higher fitness costs under these herbivory scenarios. Based on this, we

hypothesised that high and concentrated damage would elicit stronger EFN induction (in

accordance with expected higher costs due to these types of damage), leading to greater ant

recruitment and attack rates on herbivores. Overall, by disentangling the effects of herbivory

intensity and within-plant damage spatial uniformity on EFN induction and ant-plant

interactions, this study provides novel insights into how herbivory-related factors may shape

indirect plant defence.

Methods

Study species

Wild cotton (*Gossypium hirsutum* L.) is a coastal myrmecophytic shrub native to Mexico,

Central America, and the Caribbean, that is particularly abundant in the Yucatan Peninsula

(Coppens d'Eeckenbrugge & Lacape, 2014). Leaf chewers, such as grasshoppers and

caterpillars, cause most of the damage to this species, with the specialist *Alabama argillacea*

Hübner, 1823 (Erebidae) being particularly abundant and damaging at some sites (Nascimento et

al., 2011). Leaf damage exhibits considerable variability—up to 3.5-fold difference among plants

within populations, and up to three-fold variation in damage distribution across leaves within

individual plants (Abdala-Roberts et al., 2019a). In addition, there is some evidence suggesting

that high and concentrated damage are costlier for wild cotton plants (Quijano-Medina et al.,

2019; Ramírez-Delgado et al., 2025). Wild cotton secretes EFN from specialised nectaries

located along the mid-vein and sometimes on secondary veins. Several studies have shown that

EFN production in wild cotton increases primarily in response to mechanical leaf damage

(Abdala-Roberts et al., 2019b; Briones-May et al., 2023), with similar findings reported for

cultivated cotton (Wäckers & Bonifay, 2004; Llandres et al., 2019). In turn, a diverse ant

community is associated with wild cotton in situ, including species of *Camponotus*,

Pseudomyrmex, Monomorium, Pheidole Brachymyrmex, *Dorymyrmex*, and *Crematogaster*

(Vázquez-Barrios et al., 2021, Reyes-Hernandez et al., 2022). Many of these ant species

consume EFN and can play a defensive role by deterring or consuming insect herbivores

(Vázquez-Barrios et al., 2021). Also, in a previous work, Abdala-Roberts et al., (2019b) found

that higher levels of damage more strongly induced EFN, but to date, the effects of within-plant

damage spatial uniformity on EFN induction have not been tested or disentangled from the

effects of damage amount, and the consequences of these damage patterns for ant attraction and

ant-plant-herbivore interactions have not been investigated.

Plant material and propagation

We collected seeds from seven adult plants (hereafter referred to as genotypes) located at a

coastal scrubland site near the port of Sisal, Yucatán, Mexico (21°11' 45.3" N, 89°57' 18" W).

We germinated seeds and transplanted them into 2-litre polyethene bags filled with a soil mixture

of sand, native deciduous forest soil, and perlite (1:2:1). We maintained plants in a greenhouse at

the Campus de Ciencias Biológicas y Agropecuarias of the Universidad Autónoma de Yucatán

(20° 51' 57" N, 89° 37' 20" W), where they were watered three times per week with 300 ml of

water. At the time the experiments were conducted, plants were four months old.

Greenhouse experiment: treatment effects on extrafloral nectar

We used a total of 79 plants to evaluate the effects of leaf damage treatments on EFN induction.

We placed plants 90 cm apart and subjected them to one of the following mechanical damage

treatments using serrated scissors, with genotypes evenly represented across groups: a) control:

no leaves were damaged—18 plants; b) low damage: one side was removed from each of two

leaves (leaves 2 and 4, counting downward from the apical meristem)—20 plants; c) high

concentrated damage: two sides were removed from each of two leaves (2 and 4)—22 plants;

and d) high dispersed damage: one side was removed from each of four leaves (2 to 5)—19

plants. Damage treatments were applied over two consecutive days to mimic the gradual

progression of damage from feeding patterns by insects, such as lepidopterans and orthopterans,

which are the main herbivores for *G. hirsutum* (Abdala-Roberts et al., 2019a). We did not

damage leaf 1 since it is often still under expansion and this may influence local induced

responses compared to mature leaves. To standardize damage application, we first calibrated the

technique using leaves from non-experimental plants of a similar age and verifying the

percentage of leaf area removed by analysing photographs of these leaves analysed in ImageJ

software (Schneider et al., 2012). On the first day, we removed 15% of the area per leaf by

cutting one side of each of two or four leaves per plant for low and high dispersed damage

treatments, respectively, or 15% of the area for both sides of each of two leaves per plant for the

high concentrated damage treatment (30% of the total area of each leave). The following day, we

removed an additional 15% from each previously damaged side of the same leaves. As a result,

plants in the low and high dispersed damage treatments had two or four leaves damaged,

respectively, with 30% area removed per leaf in both cases. In contrast, plants in the high

concentrated damage treatment had two leaves damaged, each with 60% of the area removed.

Thus, highly damaged plants experienced twice the total amount of damage compared to low-

damaged plants. At the same time, the two high-damage treatments differed only in the spatial

uniformity of damage, not in total damage amount (Fig. S1, supplementary material).

Comparisons between the low- and high-damage treatments allowed us to test for an effect of

damage amount, while the comparison between the two high-damage treatments (concentrated

vs. dispersed) allowed us to test for an effect of within-plant damage spatial uniformity (Martín-

Cacheda et al., 2025). It is important to note that in cotton, mechanical damage alone is sufficient

to elicit EFN induction of a magnitude comparable to that caused by natural herbivory (Wäckers

& Wunderlin, 1999). This includes induction levels similar to those observed under varying

degrees of damage amount and within-plant damage dispersion caused by natural herbivory

(Ramírez-Delgado et al., 2025). Thus, mechanical damage provides a realistic proxy for testing

EFN defence induction, while offering greater precision and control over the amount and

distribution of leaf damage (Abdala-Roberts et al., 2019b). In other instances, however, as for

chemical defences, elicitors from insect oral secretions have been shown to influence cotton

chemical defences (Arce et al., 2021).

On the morning following the completion of damage treatments, between 06:00 and

07:00, we collected EFN from leaf 2 of each plant using 5-µL capillary tubes (micropipettes

Blaubrand ® intraMARK, colour code white, Germany). Twenty-four hours prior to nectar

sampling, we removed nectar from all nectaries with wetted cotton, ensuring a baseline for EFN

measurements during the sampling day. We selected leaf 2 as it was the most apical leaf with

damage, and EFN induction is typically strongest in younger leaves (1-3) and at the site of

damage (i.e., local induction; Abdala-Roberts et al., 2019b). After measuring EFN volume in the

capillary tubes, we determined sugar concentration (° Brix) using a refractometer (Atago Master,

Japan). We then calculated total sugar produced per sample as: total sugar = nectar volume ×

nectar sugar concentration (° Brix). One-degree Brix (1 ° Brix) is equivalent to 1 gram of sugar

dissolved in 100 grams of solution (Kimball, 1991). Finally, we recorded the number of active

(secreting) nectaries per leaf by visual inspection. To ensure that nectaries classified as inactive

were truly not secreting, we gently probed each one with a capillary tube to confirm the absence

of EFN. In addition to per-nectary responses reflecting local induction (volume, sugars), the

number of secreting nectaries provided a measure of whole-plant response, involving not only

local but likely also systemic induction (EFN produced by non-damaged leaves).

Field experiment: treatment effects on ant recruitment and their anti-herbivore responses

A week after the greenhouse experiment, we transported a second group of 101 plants (same age

as those used in the greenhouse EFN induction experiment) to coastal scrubland site near Sisal,

Yucatan. We placed plants at nearby wild cotton plants from which seeds for the experiments

were sourced. Plants were spaced one meter apart in an open area surrounded by natural

vegetation. We assigned plants randomly to the same leaf damage treatments described above

(control: 28 plants; low damage: 24 plants; high concentrated damage: 27 plants; high dispersed

damage: 22 plants), using the exact same methodology, with genotypes evenly represented

across treatment groups. Plant positions were also randomised. Throughout this field experiment,

we watered the plants twice a week. Two days after placement, we applied the mechanical

damage treatments. The following morning, we recorded ant abundance for 2 min per plant

between 9:00 and 11:00 AM over two consecutive days. We collected ant specimens for

taxonomic identification in the laboratory. The day after the second ant survey, we conducted ant

attack bioassays by placing caterpillars over two consecutive days. For this, each morning at 9:00

AM, we placed a frozen third- or fourth-instar caterpillar of *A. argillacea* on each plant, gently

securing it to an apical leaf with a metal clip. The use of dead caterpillars allowed us to

standardise caterpillar position and eliminate behavioural and chemical variability associated

with live specimens (Lövei & Ferrante, 2017; Nagy et al., 2020). We observed no saprophagous

insects on the plants during the bioassays. Per day, and based on prior bioassays of this type, we inspected each plant 100 minutes after caterpillar placement to determine whether the caterpillar had been attacked (scored as 1) or not attacked (scored as 0). An attack was defined as any instance in which one or more ants made direct contact with the caterpillar and exhibited aggressive behaviour or attempted to remove it. Cases in which the caterpillar was missing for unknown reasons (< 6% of observations) were excluded from statistical analyses.

Statistical analyses

We used generalised linear mixed models (GLMMs) to analyse nectar volume, total sugar production, the proportion of active nectaries (number of leaves with active nectaries relative to the total number of leaves per plant), the number of patrolling ants, and attack by ants. For both ant-related response variables, we pooled the counts across all observed ant species for increased power and to test for community-level ant responses. In all models, we included treatment as the main fixed effect and plant height as a covariate. We included plant genotype as a random effect to account for genetic or maternal effects. For the models analysing ant abundance and attack by ants, we also included survey day and plant ID as random factors to account for temporal variation and repeated measures taken on the same plants across sampling days, respectively. For nectar volume and total sugar production, we used a Tweedie distribution with a log link function, which is suitable for continuous, non-negative, and right-skewed data that include zeros (Hasan & Dunn, 2015). The proportion of active nectaries was analysed using a beta distribution with a logit link function, ant abundance (pooled across species was modelled using a negative binomial distribution with a log link function to account for overdispersion, while the probability of attack by ants was modelled using a binomial distribution with a logit link function which estimates the natural logarithm of the odds (log-odds) of an attack occurring. These log-odds were then back-transformed to provide an estimate of the probability of ant attack.

All statistical analyses were performed in R 4.4.2 (R Core Team, 2024). We used the

"glmmTMB" package (Brooks et al., 2017) to fit the GLMMs, and the "DHARMa" package

(Hartig, 2024) to evaluate models fit. To test for main effects, we employed the *Anova*() function

from the "car" package (Fox & Weisberg, 2019), using a Wald χ^2 tests. When treatment effects

were significant, we used the "emmeans" package (Lenth, 2024) to perform Tukey pairwise

comparisons between groups and to extract back-transformed estimated marginal means (EMMs)

with their 95% confidence intervals (95% CIs). We visualised these results using the "ggplot2"

package (Wickham, 2016). Model estimates are reported throughout the Results section as

EMMs with 95% CI, in the format: EMM [2.5% lower CI, 97.5% upper CI].

Results

Extrafloral nectar

Results from the greenhouse experiment showed that leaf damage significantly influenced EFN

volume (Table 1). Both high damage treatments—concentrated (0.589 [0.409, 0.850] µL) and

dispersed (0.694 [0.479, 1.01] µL)—resulted in significantly more EFN volume produced

compared to the control group (0.197 [0.107, 0.36] µL) (Fig. 1a). The low damage treatment

(0.457 [0.3, 0.697] µL) did not differ significantly from either the control or the high damage

treatments (Fig. 1a). Leaf damage also had a significant effect on total sugar production (Table

1). In this case, all damage treatments—high concentrated (0.140 [0.097, 0.2] mg), high

dispersed (0.162 [0.113, 0.233] mg), and low (0.115 [0.077, 0.172] mg)—produced significantly

more sugar than the control group (0.036 [0.019, 0.069] mg), with no significant differences

among the damage treatments (Fig. 1b). Finally, the proportion of secreting nectaries was also

significantly affected by leaf damage (Table 1), with only the high dispersed damage treatment

(0.527 [0.454, 0.599]) differing significantly from the control (0.288 [0.226, 0.359]), as well as

from the low (0.378 [0.312, 0.449]) and high concentrated (0.355 [0.294, 0.421]) damage

treatments (Fig. 1c). Statistics for Tukey pairwise comparisons for all nectar responses are

presented in Table S1 A-C, of the supplementary material.

Ant recruitment and anti-herbivore defensive response

We identified 10 ant species on the experimental cotton plants, with *Monomorium ebeninum*,

Brachymyrmex australis, and *Forelius pruinosus* being the most abundant (Table S2,

Supplementary Material). The leaf damage treatment significantly affected ant abundance (Table

1). Specifically, plants subjected to the high dispersed damage treatment attracted significantly

more ants (5.45 [3.86, 7.7] ants) than controls (2.33 [1.61, 3.35] ants) and plants exposed to high

concentrated damage (2.45 [1.69, 3.55] ants). The low damage treatment (3.01 [2.07, 4.36] ants)

did not differ significantly from either the control or the high damage treatments (Fig. 2a; Table

S1). Consistently, caterpillars on plants with high dispersed damage in turn had the highest

probability of being attacked by ants (0.886 [0.713, 0.961]), which was significantly greater than

the probability of attack on controls (0.442 [0.27, 0.629]) and plants with low damage (0.499

[0.293, 0.705]) treatments. The attack probability for the high concentrated damage treatment

(0.629 [0.429, 0.794]) exhibited an intermediate value and did not differ significantly from

controls or the other damage treatments (Fig. 2b; Table S1).

Discussion

Effects of damage amount on wild cotton EFN induction varied as a function of the trait measured, with low damage significantly inducing (relative to controls) EFN sugar content and behaving similarly to the high damage treatments, but did not induce either EFN volume or the proportion of active nectaries. Within-plant dispersion of damage, on the other hand, had no detectable effects on nectar volume or sugar content (i.e., no difference between concentrated vs. dispersed treatments), but did have a clear effect on the proportion of secreting nectaries whereby only plants in the high dispersed damage group differed significantly from controls as well as from the other damage groups, whereas the latter did not differ from controls. Consistently, dispersed damage was the only treatment to significantly increase ant abundance and the probability of attack relative to controls. Overall, these findings suggest that variation in the spatial uniformity of leaf damage within the plant (i.e., concentrated vs. dispersed damage), rather than damage amount, is the main driver of ant recruitment and indirect defence via EFN induction. These results highlight a previously underappreciated role of variability in the spatial uniformity of herbivore damage in shaping ant-plant mutualisms and call for greater attention to how herbivory patterns influence the induction of plant defences.

Extrafloral nectar

Leaf damage amount has been shown to correlate positively with EFN induction in other plant species (Rios et al., 2008; Xu et al., 2014; Calixto et al., 2021), suggesting that plants are able to allocate to induced responses proportionally to perceived herbivore threats (e.g., as a function of herbivory severity). Our findings provide partial support for this. Both high damage treatments, but not the low damage treatment, resulted in significantly more EFN volume secreted compared to controls, suggesting that damage amount matters for this trait. In contrast, low damage

increased (significantly) total sugar production to a similar extent relative to the high damage

treatments, i.e., the induction of this trait was not contingent on damage amount. This latter result

is similar to that observed, for example, in the shrub *Banisteriopsis malifolia*, for which leaf

damage increases EFN sugar secretion to similar degree regardless of damage severity (Pimenta

et al., 2025). In particular, previous work with wild cotton found that EFN volume induction

behaved similarly across levels of leaf damage (Abdala-Roberts et al., 2019b), albeit involving

overall lower damage levels (relative to the present study) and the combination of mechanical

damage with oral secretions from the generalist *Spodoptera frugiperda*. However, in a recent

study using natural herbivory by *A. argillacea* we found that low herbivory significantly induced

EFN volume (Ramírez-Delgado et al., 2025). This suggests that differences in the induction of

EFN volume between natural (but contingent on the caterpillar species) and mechanical damage

can have an important influence on EFN induction. In that same study, however, low herbivory

did not induce EFN sugar content, which together with our present results, suggests that the

induction of sugar content behaves similarly irrespective of damage amount, i.e., EFN traits are

not induced equally by different degrees of herbivory severity and based on whether natural vs.

mechanical damage is applied. Further experimentation is needed to assess better the role of

herbivory amount on wild cotton, such as the inclusion of a broader range and number of damage

levels combined with mechanical vs. natural damage treatments, to better understand herbivory

amount's effect on the induction of multiple EFN-associated traits.

Our results also revealed that within-plant variability in the spatial uniformity of leaf

damage significantly affected EFN induction, specifically by altering the proportion of secreting

nectaries. Contrary to predictions, plants subjected to high dispersed damage were the only group

that differed significantly from controls and other damage treatments for this EFN response. The

fact that recent work shows that cotton plants are less able to compensate for concentrated than

dispersed damage (Ramírez-Delgado et al., 2025), suggests that EFN induced responses are not

aligned with differential resource reallocation costs from contrasting herbivory patterns but

rather respond to other factors such as ant foraging patterns. We come back to this later on and

next speculate on two proximal, non-exclusive induction-related processes. First, high dispersed

damage presumably locally induced a greater number of nectaries (i.e., by spreading damage

across more leaves per plant, and despite lower damage amount per leaf), while maintaining

similar EFN production per nectary compared to the other damage treatments (see also Wäckers

et al., 2001). Second, dispersed damage may have also triggered a stronger systemic response,

leading to increased nectar production in undamaged leaves. Supporting the latter, we observed

that the dispersed damage treatment was the only one for which all plants exhibited nectar

secretion in at least two non-damaged leaves, whereas for low damage and concentrated damage

nectar induction was restricted to damaged leaves for most plants. Teasing apart the relative

contributions of EFN local vs. systemic induced responses would be a worthwhile goal in future

work, including also a more detailed investigation of physiological mechanisms driving systemic

EFN induced responses to dispersed damage in a defensive trait previously shown to be

preeminently shaped by local induction patterns (Abdala-Roberts et al., 2019b). These may

include, for example, within-plant volatile signalling, i.e., among stems or leaves (Hernández-

Zepeda et al., 2018) or signal transport through the xylem, which may reach undamaged leaves

and upregulate nectar production (Chatt et al., 2021). However, these explanations are merely

speculative at this stage. Finally, considering the role of herbivore chemical elicitors using

natural damage by different insect herbivores would also be useful, as even though EFN

responses are often strongly determined by mechanical damage per se, herbivore-specific factors

merit attention in studying the mechanisms of local vs. systemic induction of EFN under

contrasting herbivory patterns.

Ant abundance and anti-herbivore response

The ant species recorded during the field experiment have been previously reported on wild

cotton, though involving different species composition and relative abundances in some cases.

For example, Reyes-Hernández et al., (2022) reported high abundances of *Crematogaster* on

wild cotton in a tropical dry forest gap, while Vázquez-Barrios et al., (2021) found that

Camponotus planatus and *Paratrechina longicornis* were dominant species in a coastal

scrubland. In contrast, our study recorded high abundances of *Monomorium ebeninum*,

Brachymyrmex australis, and *Forelius pruinosus*. Although these small-bodied species are not

strictly predatory (Díaz-Castelazo et al., 2004), their persistent presence on plants may

discourage herbivores through disturbance or harassment, potentially leading to reduced overall

damage (Bentley, 1977; Rudgers et al., 2003). Larger, more aggressive species, including

predatory ants in *Pseudomyrmex* and *Camponotus*, were much less abundant but have been

previously documented on wild cotton at coastal sites (Vázquez-Barrios et al., 2021). Ant

competitive dynamics and the availability of alternative plant resources or prey may influence

ant community composition (Byk & Del-Claro, 2011; Nogueira et al., 2015) and warrant further

examination across wild cotton populations.

Mirroring effects on the proportion of secreting nectaries, ant abundance (pooled across

species) was highest on plants subjected to high dispersed damage. This was likely due to an

increased availability of secreting nectaries per plant under this treatment, enhancing ant

recruitment and patrolling activity. Consistent with this finding, attack rates on caterpillars were

highest on plants subjected to dispersed damage, suggesting increased ant abundance led to

higher attack rates, i.e., a density-mediated effect (Abdala-Roberts et al., 2012). It is also

possible that dispersed damage increased ant aggressiveness or foraging efficiency,

independently of ant numbers, as previous work in other systems has reported that plants

influence ant behaviour through differences in nectar composition (Ness, 2003; Ness et al., 2006;

Shenoy et al., 2012), i.e., a trait-mediated effect (Abdala-Roberts & Mooney, 2015). Regardless

of the mechanism, and while this finding suggests dispersed damage could lead to heightened

indirect defence, it is important to note that recorded attacks on caterpillars may not be

representative of actual ant effectiveness in reducing herbivory. In addition, herbivory levels are

highly fluctuating and are overall often low at cotton populations (Abdala-Roberts et al., 2019a),

suggesting that the benefits to plant fitness of short-term ant responses may only accrue under a

specific set of conditions (e.g., periods of high and consistent damage over several weeks and ant

communities dominated by more aggressive species). It is important to also note that although

using dead caterpillars has proven to be a useful as well as realistic method for testing ant

predatory or deterrence behaviour against insect herbivores (Grof-Tisza et al., 2024), this

approach may underestimate attack rates (as live caterpillars may move and attract more

attention) or overestimate successful deterrence and linkages to indirect defence on plants as live

caterpillars can actively defend themselves (Lövei & Ferrante, 2017; Nagy et al., 2020). Within

this context, an important challenge will be to disentangle ant-plant-herbivore interaction

mechanisms via experimental procedures such as ant exclusions across sites with contrasting

herbivore and ant abundances and composition, and testing for such effects over longer periods

of time during the growing season.

An ecological-based mechanism linking cotton direct and indirect defences in response to

herbivory variability

The implications of our results can be interpreted within the broader context of cotton's

extensive defensive arsenal, including direct chemical defences. Concentrated herbivory

typically imposes greater fitness costs on plants (Marquis, 1992; Wetzel et al., 2023), a notion

that is consistent with recent findings with wild cotton (Ramírez-Delgado et al., 2025). We

predicted that direct chemical defences (e.g., phenolics, terpene aldehydes) should be more

strongly induced under such attacks. If this is the case, in theory, it would not only reduce

herbivory but also promote its spread from leaf-to-leaf, further contributing to dampen the costs

of localised damage (Marquis, 1996; Karban & Yang, 2020). Once damage becomes dispersed,

the plant's strategy can shift to enhancing indirect defence, aligning with our results of increased

EFN production and ant recruitment. Based on this, a selective feedback loop could be in place

whereby selection for direct and indirect defences becomes entrained: the induction of chemical

defences against concentrated attacks is reinforced because it leads to dispersed damage, which

in turn promotes stronger or more effective EFN-mediated ant defence. Current research in our

group is aimed at testing the effects of variability in the spatial uniformity of herbivore damage

on both chemical defences and EFN, as well as assessing the fitness costs of different herbivory

spatial uniformity patterns. From this, we can then expand and evaluate these responses across

populations that experience contrasting ecological conditions (e.g., varying ant composition,

herbivore pressure, and abiotic stress). In closing, our results underscore the complexity of plant-

insect multitrophic interactions in natural ecosystems and emphasise the need to understand how

plants optimise their defensive investments across varying herbivory contexts in order to

maximise the benefits of indirect defence derived from mutualistic interactions with predators.

References

Abdala-Roberts, L. & Mooney, K. A. (2015) Plant and herbivore evolution within the trophic
sandwich. In: Hanley, T. & La Pierre, K. (Eds.) *Trophic interactions: bottom-up and top-*
down interactions in aquatic and terrestrial ecosystems. Cambridge University Press, pp.
340–364.

Abdala-Roberts, L., Quijano-Medina, T., Moreira, X., Vázquez-González, C., Parra-Tabla, V.,
Berny Mier Y Terán, J. C., Grandi, L., Glauser, G., Turlings, T. C. J. & Benrey, B.
(2019a) Bottom-up control of geographic variation in insect herbivory on wild cotton
(*Gossypium hirsutum*) by plant defenses and climate. *American Journal of Botany*, 106:
1059–1067.

Abdala-Roberts, L., Reyes-Hernández, M., Quijano-Medina, T., Moreira, X., Francisco, M.,
Angulo, D. F., Parra-Tabla, V., Virgen, A. & Rojas, J. C. (2019b) Effects of amount and
recurrence of leaf herbivory on the induction of direct and indirect defences in wild
cotton. *Plant Biology*, 21: 1063–1071.

Arce, C. M., Besomi, G., Glauser, G. & Turlings, T. C. J. (2021) Caterpillar-induced volatile
emissions in cotton: the relative importance of damage and insect-derived factors.
Frontiers in Plant Science, 12: 709858.

Bentley, B. L. (1977) Extrafloral nectaries and protection by pugnacious bodyguards. *Annual*
Review of Ecology, Evolution and Systematics, 8: 407–427.

Briones-May, Y., Quijano-Medina, T., Perez-Nino, B., Benrey, B., Turlings, T. C. J., Bustos-
Segura, C. & Abdala-Roberts, L. (2023) Soil salinization disrupts plant-plant signaling
effects on extra-floral nectar induction in wild cotton. *Oecologia,* 202: 313–323.

Bronstein, J. (1998) The contribution of ant plant protection studies to our understanding of
mutualism. *Biotropica*, 30: 150–161.

Bronstein, J. L., Alarcon, R. & Geber, M. (2006) The evolution of plant-insect mutualisms. *New*
Phytologist, 172: 412–428.

Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A.,
Skaug, H. J., Maechler, M. & Bolker, B. M. (2017) glmmTMB balances speed and

flexibility among packages for zero-inflated generalized linear mixed modeling. *The R*
Journal, 9: 378–400.

Byk, J. & Del-Claro, K. (2011) Ant–plant interaction in the Neotropical savanna: direct
beneficial effects of extrafloral nectar on ant colony fitness. *Population Ecology*, 53:
327–332.

Calixto, E. S., Lange, D., Bronstein, J., Torezan-Silingardi, H. M. & Del-Claro, K. (2021)
Optimal defense theory in an ant–plant mutualism: extrafloral nectar as an induced
defence is maximized in the most valuable plant structures. *Journal of Ecology*, 109:
167–178.

Calixto, E. S., Del-Claro, K., Lange, D. & Bronstein, J. (2023) Time course of inducibility of
indirect responses in an ant-defended plant. *Ecology*. 104: e4029.

Chatt, E. C., Mahalim, S.-N., Mohd-Fadzil, N.-A., Roy, R., Klinkenberg, P. M., Horner, H. T.,
Hampton, M., Carter, C. J. & Nikolau, B. J. (2021) Nectar biosynthesis is conserved
among floral and extrafloral nectaries. *Plant Physiology*, 185: 1595–1616.

Coppens d'Eeckenbrugge, G. & Lacape, J.-M. (2014) Distribution and differentiation of wild,
feral, and cultivated populations of perennial upland cotton (*Gossypium hirsutum* L.) in
Mesoamerica and the Caribbean. *PLoS ONE,* 9: 1–19.

Díaz-Castelazo, C., Rico-Gray, V., Oliveira, P. & Cuautle, M. (2004) Extrafloral nectary-
mediated ant-plant interactions in the coastal vegetation of Veracruz, Mexico: Richness,
occurrence, seasonality, and ant foraging patterns. *Écoscience*, 11: 472–481.

Fox, J. & Weisberg, S. (2019) An R Companion to Applied Regression. Sage, Thousand Oaks.

Grof-Tisza, P., Turlings, T. C. J., Bustos-Segura, C. & Benrey, B. (2024) Field evidence for the
role of plant volatiles induced by caterpillar oral secretion in prey localization by
predatory social wasps. *Biology Letters*, 20: 20240384.

Hartig, F. 2024. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models.
https://cran.r-project.org/web/packages/DHARMa/vignettes/DHARMa.html

Hasan, M. M. & Dunn, P. K. (2015) Seasonal rainfall totals of Australian stations can be
modelled with distributions from the Tweedie family. *International Journal of*
Climatology, 35: 3093–3101.

Heil, M. (2015) Extrafloral nectar at the plant-insect interface: a spotlight on chemical ecology,
phenotypic plasticity, and food webs. *Annual Review of Entomology*, 60: 213–232.

Heil, M. & McKey, D. (2003) Protective ant-plant interactions as model systems in ecological
and evolutionary research. *Annual Review of Ecology, Evolution and Systematics*, 34:
425–553.

Heil, M., Koch, T., Hilpert, A., Fiala, B., Boland, W. & Linsenmair, K. E. (2001) Extrafloral
nectar production of the ant-associated plant, *Macaranga tanarius*, is an induced,
indirect, defensive response elicited by jasmonic acid. *Proceedings of the National*
Academy of Sciences, 98: 1083–1088.

Hernández-Zepeda, O. F., Razo-Belman, R. & Heil, M. (2018) Reduced responsiveness to
volatile signals creates a modular reward provisioning in an obligate food-for-protection
mutualism. *Frontiers in Plant Science*, 9: 1076.

Holland, J. N., Chamberlain, S. A. & Horn, K. C. (2009) Optimal defence theory predicts
investment in extrafloral nectar resources in an ant–plant mutualism. *Journal of Ecology*,
97: 89–96.

Huang, W., Siemann, E., Carrillo, J. & Ding, J. (2015) Below-ground herbivory limits induction
of extrafloral nectar by above-ground herbivores. *Annals of Botany*, 115: 841–846.

Interian-Aguiñaga, J., Solís-Rodriguez, U., Briones-May, Y., Guadarrama-Chávez, P., Mamin,
M., Clancy, M. V., Turlings, T. C. J., Moreira, X. & Abdala-Roberts, L. (2025) Soil
salinity effects on inter-plant signalling in wild cotton are contingent on herbivore load
and differentially shape direct and indirect defences. *Planta*, 262: 64.

Jones, I. M. & Koptur, S. (2015) Dynamic extrafloral nectar production: The timing of leaf
damage affects the defensive response in *Senna mexicana* var. *chapmanii* (Fabaceae).
American Journal of Botany, 102: 58–66.

Karban, R. & Yang, L. H. (2020) Feeding and damage-induced volatile cues make beetles
disperse and produce a more even distribution of damage for sagebrush. *Journal of*
Animal Ecology, 89: 2056–2062.

Kessler, A. & Heil, M. (2011) The multiple faces of indirect defences and their agents of natural
selection: Multiple faces of indirect defences. *Functional Ecology*, 25: 348–357.

Kimball, D. (1991) Brix and soluble solids. In: *Citrus Processing*. Springer, Dordrecht, pp. 7–33.

Lenth, R. V. (2024) emmeans: Estimated Marginal Means, aka Least-Squares Means.
https://rvlenth.github.io/emmeans/

Lövei, G. L. & Ferrante, M. (2017) A review of the sentinel prey method as a way of quantifying
invertebrate predation under field conditions. *Insect Science*, 24: 528–542.

Llandres, A. L., Verdeny-Vilalta, O., Jean, J., Goebel, F.-R., Seydi, O. & Brévault, T. (2019)
Cotton extrafloral nectaries as indirect defence against insect pests. *Basic and Applied*
Ecology, 37: 24–34.

Marazzi, B., Bronstein, J. L. & Koptur, S. (2013) The diversity, ecology and evolution of
extrafloral nectaries: current perspectives and future challenges. *Annals of Botany*, 111:
1243–1250.

Marquis, R. J. (1992) A bite is a bite is a bite? Constraints on response to folivory in *Piper*
arieianum (Piperaceae). *Ecology*, 73: 143–152.

Marquis, R. J. (1996) Plant architecture, sectoriality and plant tolerance to herbivores. *Vegetatio*,
127: 85–97.

Martín-Cacheda, L., Briones-May, Y., Bustos-Segura C., Turlings, T. C. J., Ramírez-Delgado, V.
H. & Abdala-Roberts, L. (2025) Effects of herbivory amount and within-plant variability
by a specialist caterpillar on volatile emissions mediating inter-plant signalling in wild
cotton. *Ecological Entomology*, 50: 909–920.

Mauricio, R., Bowers, M. D. & Bazzaz, F. A. (1993) Pattern of leaf damage affects fitness of the
annual plant *Raphanus sativus* (Brassicaceae). *Ecology*, 74: 2066–2071.

Moreira, X., Zas, R. & Sampedro, L. (2012) Differential allocation of constitutive and induced chemical defenses in pine tree juveniles: a test of the optimal defense theory. *PLoS ONE*, 7: e34006.

Nagy, R. K., Schellhorn, N. A. & Zalucki, M. P. (2020) Fresh, frozen or fake: A comparison of predation rates measured by various types of sentinel prey. *Journal of Applied Entomology*, 144: 407–416.

Nascimento, A., Ramalho, F. S., Azeredo, T. L., Fernandes, F. S., Júnior, J. L. N., Silva, C. & Malaquias, J. B. (2011) Feeding and life history of *Alabama argillacea* (Lepidoptera: Noctuidae) on cotton cultivars producing colored fibers. *Annals of the Entomological Society of America*, 104: 613–619.

Ness, J. (2003) *Catalpa bignonioides* alters extrafloral nectar production after herbivory and attracts ant bodyguards. *Oecologia*, 134: 210–218.

Ness, J. H., Morris, W. F. & Bronstein, J. L. (2006) Integrating quality and quantity of mutualistic service to contrast ant species protecting *Ferocactus wislizeni*. *Ecology*, 87: 912–921.

Nogueira, A., Rey, P. J., Alcántara, J. M., Feitosa, R. M. & Lohmann, L. G. (2015) Geographic mosaic of plant evolution: extrafloral nectary variation mediated by ant and herbivore assemblages. *PLoS ONE*, 10: e0123806.

Nogueira, R. R. (2025) Mutualistic interactions between plants with extrafloral nectaries and ants: ecological impacts, ant sharing among neighboring plants, and the effects of environmental conditions. *Community Ecology*, 26: 437–447.

Ohnmeiss, T. E., McCloud, E. S., Lynds, G. Y. & Baldwin, I. T. (1997) Within-plant relationships among wounding, jasmonic acid, and nicotine implications for defence in *Nicotiana sylvestris*. *New Phytologist*, 137: 441–452.

Pimenta, I. C. D. O., Calixto, E. S. & Del-Claro, K. (2025) Impacts of leaf damage intensity on ant–plant protection mutualism and plant fitness. *Plants*, 14: 837.

Quijano-Medina, T., Covelo, F., Moreira, X. & Abdala-Roberts, L. (2019) Compensation to simulated insect leaf herbivory in wild cotton (*Gossypium hirsutum*): responses to multiple levels of damage and associated traits. *Plant Biology*, 21: 805–812.

Quijano-Medina, T., Turlings, T. C. J., Sosenski, P., Grandi, L., Cervera, J. C., Moreira, X. &
Abdala-Roberts, L. (2021) Effects of soil salinity on the expression of direct and indirect
defences in wild cotton *Gossypium hirsutum*. *Journal of Ecology*, 109: 354–368.

R Core Team 2024. R: A language and environment for statistical computing. https://www.r-
project.org/

Ramírez-Delgado, V. H., Briones-May, Y., Sánchez-Durán, J., Martín-Cacheda, L., Moreira, X.,
Turlings, T. C. J. & Abdala-Roberts, L. (2025) Effects of varying amounts and
distribution of herbivory on resistance- and tolerance-related responses in wild cotton.
bioRxiv, 2025.07.14.664844.

Reyes-Hernández, M., Angulo-Pérez, D., Quijano-Medina, T., Moreira, X., Parra-Tabla, V.,
Vásquez-Bolaños, M. & Abdala-Roberts, L. (2022) An experimental test of ant effects on
herbivory and pathogen infection on wild cotton (*Gossypium hirsutum* L.). *Arthropod-*
Plant Interactions, 16: 77–86.

Rico-Gray, V. & Oliveira, P. S. (2008) The ecology and evolution of ant-plant interactions.
University of Chicago Press, Chicago.

Rios, R. S., Marquis, R. J. & Flunker, J. C. (2008) Population variation in plant traits associated
with ant attraction and herbivory in *Chamaecrista fasciculata* (Fabaceae). *Oecologia*,
156: 577–588.

Rosumek, F. B., Silveira, F. A. O., De S. Neves, F., De U. Barbosa, N. P., Diniz, L., Oki, Y.,
Pezzini, F., Fernandes, G. W. & Cornelissen, T. (2009) Ants on plants: a meta-analysis of
the role of ants as plant biotic defenses. *Oecologia*, 160: 537–549.

Rudgers, J. A., Hodgen, J. G. & White, J. W. (2003) Behavioral mechanisms underlie an ant-
plant mutualism. *Oecologia*, 135: 51–59.

Rudgers, J. A. & Strauss, S. Y. (2004) A selection mosaic in the facultative mutualism between
ants and wild cotton. *Proceedings of the Royal Society of London. Series B: Biological*
Sciences, 271: 2481–2488.

Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. (2012) NIH Image to ImageJ: 25 years of
image analysis. *Nature Methods*, 9: 671–675.

Shenoy, M., Radhika, V., Satish, S. & Borges, R. M. (2012) Composition of extrafloral nectar
influences interactions between the myrmecophyte *humboldtia brunonis* and its ant
associates. *Journal of Chemical Ecology*, 38: 88–99.

Underwood, N. (2010) Density dependence in insect performance within individual plants:
induced resistance to *Spodoptera exigua* in tomato. *Oikos*, 119: 1993–1999.

Vázquez-Barrios, V., Boege, K., Sosa-Fuentes, T. G., Rojas, P. & Wegier, A. (2021) Ongoing
ecological and evolutionary consequences by the presence of transgenes in a wild cotton
population. *Scientific Reports*, 11: 1959.

Wäckers, F. L. & Wunderlin, R. (1999) Induction of cotton extrafloral nectar production in
response to herbivory does not require a herbivore-specific elicitor. *Entomologia*
Experimentalis et Applicata, 91:149–154.

Wäckers, F. L. & Bonifay, C. (2004) How to be sweet? Extrafloral nectar allocation by
Gossypium hirsutum fits optimal defense theory predictions. *Ecology*, 85: 1512–1518.

Wäckers, F. L., Zuber, D., Wunderlin, R. & Keller, F. (2001) The effect of herbivory on
temporal and spatial dynamics of foliar nectar production in cotton and castor. *Annals of*
Botany, 87: 365–370.

Weber, M. G. & Keeler, K. H. (2012) The phylogenetic distribution of extrafloral nectaries in
plants. *Annals of Botany*, 111: 1251–1261.

Wetzel, W. C., Inouye, B. D., Hahn, P. G., Whitehead, S. R. & Underwood, N. (2023)
Variability in plant–herbivore interactions. *Annual Review of Ecology, Evolution and*
Systematics, 54: 451–474.

Wickham, H. (2016) ggplot2: Elegant graphics for data analysis. Springer-Verlag New York,
New York.

Xu, F. F., Chen, J. & Husson, J. (2014) Leaf area lost, rather than herbivore type, determines the
induction of extrafloral nectar secretion in a tropical plant (*Clerodendrum philippinum*).
Arthropod-Plant Interactions, 8: 513–518.

Yamawo, A. & Suzuki, N. (2018) Induction and relaxation of extrafloral nectaries in response to

simulated herbivory in young *Mallotus japonicus* plants. *Journal of Plant Research*, 131:

255–260.

Tables

Table 1. Wald χ² tests for the GLMMs assessing the effects of leaf damage treatments and plant

height on EFN (volume, total sugar, and proportion of active nectaries) and ant activity

(patrolling, and attack on caterpillars).

Response variable	Effect	df	χ^2	P
EFN volume	Damage treatment	3	13.69	**0.0003**
	Plant height	1	1.41	0.235
Sugar production	Damage treatment	3	16.95	**0.0007**
	Plant height	1	0.21	0.6394
Proportion of nectaries active	Damage treatment	3	23.39	**< 0.0001**
	Plant height	1	1.246	0.2643
Patrolling ants	Damage treatment	3	14.99	**0.0018**
	Plant height	1	3.12	0.0773
Attack on caterpillars	Damage treatment	3	10.92	**0.0121**
	Plant height	1	1.53	0.2454

Figures legends

Figure 1. EFN volume production and sugar content for leaf 2 (damaged leaf, except for control plants) of wild cotton (*Gossypium hirsutum*) plants under different leaf damage treatments. Panels show: (a) per-nectary EFN volume, (b) per-nectary EFN sugar content, and (c) the proportion of active nectaries per plant. Black dots and error bars indicate EMMs and 95% CIs based on GLMMs. Single values are plotted (open colour circles). Different letters indicate significant differences between treatments according to Tukey's post hoc test ($p < 0.05$).

Figure 2. Ant activity on wild cotton (*Gossypium hirsutum*) plants under different leaf damage treatments. Panels: (a) mean number of patrolling ants per plant, and (b) probability of ant attack on caterpillars (i.e., indirect defence bioassays). Black dots and error bars indicate EMMs and 95% CIs based on GLMMs. Single values are plotted (open colour circles). Different letters indicate significant differences between treatments according to Tukey's post hoc test ($p < 0.05$).

(a)

Patrolling ants by plant

(b)

Probability of ant attack

Control (no damage)   Low   High concentrated   High dispersed

Damage treatments