Molecular evolution of terpene synthase underlying the diversification of isoprene emission in Fagaceae

Yuka Ikezaki[1], Shuichi N. Kudo[1], Taichi Nakata[1], Sora Koita[2], Ryosuke Munakata[2], Kazufumi Yazaki[2], Takeshi Torimaru[3], Nobuhiro Tomaru[4], Sachiko Isobe[5,6], Hideki Hirakawa[5,7], Junko Kusumi[8]*, Akiko Satake[1]*

1. Department of Biology, Faculty of Science, Kyushu University, Fukuoka 819-0395, Japan.
2. Research Institute for Sustainable Humanosphere, Kyoto University, Gokasho, Uji, 611-0011 Japan.
3. Graduate School of Bioresources, Mie University, Tsu, 514-8507 Japan.
4. Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya 464-8603 Japan.
5. Kazusa DNA Research Institute, 2-6-7 Kazusa-kamatari, Kisarazu, Chiba 292-0818, Japan.
6. Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, 113-8657, Japan.
7. Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University, Fukuoka 819-0395, Japan.
8. Department of Environmental Changes, Faculty of Social and Cultural Studies, Kyushu University, Fukuoka 819-0395, Japan.

*Corresponding author:
Akiko Satake
Email: satake.akiko.269@m.kyushu-u.ac.jp
Kusumi Junko
Email: jkusumi@scs.kyushu-u.ac.jp

Author Contributions: Y.I. and A.S. designed research; Y.I., S.N.K., N.T., S.K., R.M., K.Y., S.I., H.H., J.K. and A.S. performed research; Y.I., S.N.K., N.T., S.K., R.M., K.Y., S.I., H.H., J.K. and A.S. analyzed data; T.T. and N.T. provided genome data; Y.I. and A.S. wrote the paper with inputs from all the authors.

Competing Interest Statement: The authors declare no conflict of interest.

Classification: Biological Sciences, Evolution

Keywords: Biogenic volatile compounds, isoprene synthase, ancestral sequence reconstruction, positive selection, convergent evolution

Abstract

Plants emit a wide range of volatile organic compounds, among which isoprene is the most abundant and atmospherically influential. Although oak species are major contributors to isoprene emission, there is considerable variation in isoprene

emission capacity within the Fagaceae family. To unravel the evolutionary origins of isoprene emission, we investigated the molecular evolution of terpene synthase (TPS) genes across eight species from four genera within the Fagaceae. We identified a Fagaceae-specific TPS-b subclade in which potential isoprene synthase (IspS) activity evolved independently in two gene lineages within subgenus *Quercus*. Seasonal gene expression patterns showed that one lineage exhibited high expression particularly in summer, whereas the other displayed consistently low expression across all seasons. Ancestral sequence reconstruction revealed that the acquisition of a diagnostic amino acid residue for IspS function (a substitution from Thr338 to Phe338) arose convergently in these lineages and was subject to positive selection, suggesting adaptive evolution. Ancestral-enzyme assays targeting the gene lineage with high gene expression revealed that the early protein primarily produced monoterpenes—particularly *trans*-β-ocimene— from geranyl diphosphate (GPP), whereas their descendants shifted substrate preference to dimethylallyl diphosphate (DMAPP), evolving into dedicated isoprene synthases. Our results indicate that IspS activity was not ancestral in Fagaceae, but evolved approximately 56 million years ago within the subgenus *Quercus*, and has been retained ever since. These findings emphasize the roles of enzyme structural innovation and regulatory shifts in the diversification of volatile terpenoid biosynthesis.

Significance Statement

Isoprene, the most abundantly emitted biogenic volatile organic compound, influences atmospheric chemistry and protects plants from thermal stress. Yet, its evolutionary origins remain unclear. By analyzing TPS genes across eight Fagaceae species, we demonstrated that high isoprene synthase activity was not ancestral in Fagaceae, but evolved approximately 56 million years ago within the subgenus *Quercus*. Ancestral sequence reconstruction revealed that IspS activity arose through a convergent amino acid substitution under positive selection. Enzyme assays further showed that this functional shift was accompanied by substrate preference changes. Our study provides the genomic evidence of convergent evolution of isoprene biosynthesis in trees and highlights the molecular basis underpinning ecological adaptation through specialized metabolism.

Main Text

Introduction
Plants release a wide spectrum of biogenic volatile organic compounds (BVOCs), which include molecules with diverse structural and physicochemical characteristics (1-3). These compounds, emitted from various plant tissues such as leaves, flowers, fruits, and roots, enter the atmosphere in significant quantities (4). BVOCs serve as a dynamic interface between terrestrial ecosystems and the atmosphere, playing an important role in climate and atmospheric chemistry (5). The annual global emission of BVOCs is estimated to be around 1000 Tg C, with terpenoids—including isoprene, monoterpenes, and sesquiterpenes (6, 7). Among

these, isoprene alone accounts for roughly half of the total BVOC, underscoring its
ecological and atmospheric significance (6).
In the plant kingdom, isoprene emission is distributed heterogeneously
across plant taxa (8-10). It is particularly common in fast-growing plants, especially
hardwood trees such as oaks (*Quercus* spp.), *Populus* spp. (11), and *Eucalyptus*
spp. (10). Isoprene emission has also been reported in mosses (12), ferns (13),
and legumes (14), indicating its occurrence across diverse plant groups. Isoprene
is synthesized by isoprene synthase (IspS), a class I terpenoid synthase (TPS)
enzyme that converts dimethylallyl diphosphate (DMAPP) into isoprene at a key
terminal branching point of the methylerythritol phosphate (MEP) pathway (15, 16).
The TPS family likely originated in ancestral land plants following their divergence
from green algae (17). In angiosperms, IspS belongs to the TPS-b subfamily (15,
18). In contrast, recent research has shown that IspS in mosses belongs to the
TPS-c subfamily (19), suggesting that IspS has multiple evolutionary origins within
the plant kingdom. In angiosperms, IspS appears to have evolved multiple times,
likely from closely related terpene synthases. Among these, *trans*-β-ocimene
synthases are considered a key precursor, reflecting the functional diversification
of terpene synthases in plants (15, 18).
An intriguing and well-studied example of the uneven distribution of
isoprene emission within a plant clade can be found in the Fagaceae family (8, 9).
In this family, isoprene emission is particularly prominent in the genus *Quercus*
(oaks), whereas other genera, such as *Fagus*, *Castanopsis*, and *Lithocarpus*
exhibit little to no isoprene emission (9, 20, 21). Intriguingly, within the genus
Quercus, isoprene emission capacity exhibits significant variation, with a mix of
strong isoprene emitters, such as *Q. rubra* and *Q. robur*, and non-emitters, such
as *Q. ilex* and *Q. suber* (22). Based on analyses associating isoprene emission
capacity with molecular phylogeny within the Fagaceae, Loreto (20) and Monson
et al. (9) concluded that isoprene emission was likely an ancestral trait in the genus
Quercus and was subsequently lost and not restored in certain species within the
Quercus sections *Cerris* and *Ilex*. These findings lay a critical foundation for
reevaluating the evolutionary history of isoprene emission in the Fagaceae. The
growing availability of high-quality genomic and transcriptomic data, even for non-
model Fagaceae species (23-25), presents a valuable opportunity to reconstruct
the evolutionary trajectory of isoprene emission by examining genome evolution
and the expression patterns of *IspS*.
In the present study, we investigated the evolutionary trajectory of isoprene
emission in eight species representing four genera within the Fagaceae family.
Using comparative genomics, phylogenetics, and ancestral sequence
reconstruction, we uncovered a convergent evolutionary origin of *IspS*, shedding
light on a novel evolutionary pathway to isoprene emission in angiosperms.
130
131 **Results**
132 **Identification of TPS genes in Fagaceae.** To identify TPS genes in Fagaceae,
133 we first obtained genome sequences of eight Fagaceae trees from four genera
134 using public databases (Table S1). These included five species from the genus
135 *Quercus* (*Q. robur*, *Q. lobata*, *Q. rubra*, *Q. acutissima*, and *Q. glauca*), and one

species each from the genera *Lithocarpus* (*L. edulis*), *Castanopsis* (*C. hystrix*), and *Fagus* (*F. crenata*) (Table S1). Isoprene emission factors vary significantly among these eight species (Table S2), highlighting their suitability for studying the evolution of isoprene emission. A comparative genomic analysis of these species revealed a highly conserved syntenic structure across genera, including *Quercus*, *Lithocarpus*, and *Castanopsis* (Fig. 1*A*). However, the genomic structure of the genus *Fagus* differed significantly from that of the other genera (Fig. 1*A*), suggesting extensive genome rearrangements following its divergence from the rest of the Fagaceae.

To identify TPS genes in our target Fagaceae species, we conducted HMMER searches using the annotated protein sequences from the genomes of eight Fagaceae species against the metal-binding domain (PF03936) and the N-terminal domain (PF01397). Each TPS candidate gene was further examined and filtered based on exon-intron structure, including the number and locations of exons (Fig. S1), by comparing with previously reported TPSs (26), as well as total sequence length (Fig. S2). The number of identified TPS genes varied across species, ranging from a minimum of 24 in *C. hystrix* to a maximum of 67 in *Q. glauca* (Fig. S3).

The rooted TPS phylogenetic tree constructed from our TPS dataset along with sequences from *Solanum lycopersicum* (tomato) and *Arabidopsis thaliana*, revealed a clear distinction among the TPS-a, TPS-b, TPS-c, TPS-e/f, and TPS-g subfamilies (Fig. 1*B*). TPS-a members in Fagaceae formed a distinct clade, separate from those of tomato and *A. thaliana*, but showing a closer evolutionary relationship to tomato TPS-a genes (Fig. 1*B*). The TPS-c, TPS-g, and TPS-e/f subfamilies were present in all Fagaceae species, except for the TPS-e/f subfamily, which was absent in *Q. acutissima* (Fig. 1*B*).

The TPS-b subfamily represents the largest gene group within Fagaceae species (Fig. 1*B*). The three TPS-b genes in tomato form a distinct clade from other tomato and Arabidopsis TPS-b genes (red asterisk in Fig. 1*B*). These three tomato genes are known to synthesize acyclic monoterpenes, with one of them, TPS47, encoding a cytosolic isoprene synthase (27).

Many genes in the TPS-a and TPS-b subfamilies are arranged in tandem gene arrays on chromosomes 8 and 9, respectively, suggesting their origin from tandem gene duplication events (Fig. S4*A*). Overall, the genomic locations of these genes were largely conserved across *Quercus*, *Lithocarpus*, and *Castanopsis*. However, in *Fagus*, significant genome rearrangements were observed, indicating a distinct evolutionary trajectory (Fig. S4*B*).

Isoprene score of TPS-b genes in Fagaceae. To investigate the evolutionary history of *IspS* in Fagaceae, we focused on the TPS-b subfamily that includes known *IspS* from other plant species. We designated the clade containing the tomato *IspS* as TPS-b-target1 (TPS-b-t1) and the clade containing the *Q. serrata* *IspS* as TPS-b-target1 (TPS-b-t3) (Fig. 2*A*). Between the TPS-b-t1 and TPS-b-t3, we identified a distinct clade, which we designate as TPS-b-target2 (TPS-b-t2) (Fig. 2*A*).Notably, both TPS-b-t2 and TPS-b-t3 consist exclusively of genes from

Fagaceae, suggesting that these clades are specific to this family and potentially
involved in the evolution of *IspS* in Fagaceae.
Sequences of genes included in these three clades were aligned and used
to calculate an isoprene score (Fig. 2*B*). The isoprene score serves as a useful
screening tool to identify enzymes with exclusive IspS activity based on sequence
information and the structural analysis of IspS (15). This score is calculated using
the four conserved amino acid residues (known as the diagnostic tetrad) that are
characteristic of IspS. Two of the residues (F338 and F485 in Fig. 2*B*) were
suggested to restrict the size of the substrate-binding pocket and play a key role
in determining enzyme specificity (15). While F338 and F485 of the diagnostic
tetrad residues are considered highly specific to IspS, the other positions are
considered redundant (18). In IspS in some angiosperm families, valine (V446) is
found instead of serine (18). Therefore, we treated both serine and valine as
conserved residues at this position and denoted it as S/V446. The isoprene score
ranges from 0 to 4, with a score of 4 representing full conservation of the diagnostic
tetrad and thus a high likelihood of exclusive IspS activity.
The multiple sequence alignment revealed that the isoprene score for
sequences in the TPS-b-t1 clade was consistently one, whereas the IspS
sequence in tomato had a score of 3 (Fig. 2*B*). In the TPS-b-t2 and TPS-b-t3
clades, sequences from the species in the subgenus *Quercus* (*Q. robur*, *Q. lobata*,
Q. rubra, and *Q. serrata*) were found to have an isoprene score of 4. Specifically,
Q. robur contained three, *Q. lobata* had two, and *Q. rubra* had one sequence with
an isoprene score of 4. In contrast, the maximum isoprene score for sequences
from *Lithocarpus*, *Castanopsi*s, and *Fagus* was only three. Many sequences
lacked F338, an amino acid considered crucial for isoprene synthase activity (15).
These patterns in isoprene score align with isoprene emission capacities, where
Q. robur, *Q. lobata*, *Q. rubra*, and *Q. serrata* are known to exhibit high isoprene
emission, while other species show lower emission levels (Table S2)
Within the twin clades TPS-b-t2 and TPS-b-t3, *Fagus* sequences were
found only in TPS-b-t2, whereas sequences from other species were present in
both clades. Most genes within TPS-b-t2 and TPS-b-t3 were located on
chromosome 8, with the exception of *F. crenata* (Fig. S4). Interestingly, two genes
in *Q. robur* and one in *Q. lobata*, all with an isoprene score of 4, were located on
chromosome 3 (Fig. S4), suggesting chromosomal translocation events.
215
Seasonal expression patterns of TPS genes. To investigate the seasonal
expression patterns of TPS genes, we conducted RNA-seq on leaf and bud tissues
collected monthly over a two-year period from *Q. acutissima* growing under natural
conditions in Fukuoka, Japan. Additionally, we incorporated previously published
field transcriptome data from *Q. glauca* and *L. edulis* (25), which were sampled at
the same study site as *Q. acutissima*. On average, 36% and 41% of TPS genes
were expressed in leaves (Fig. 3*A*) and buds (Fig. S6*A*), respectively. Within the
TPS-b clade, 42% of genes were expressed in leaves, including 6 out of 14 in *Q.*
acutissima, 10 out of 28 in *Q. glauca*, and 2 out of 24 in *L. edulis*. Seasonal
expression profiles in leaves revealed that most expressed genes exhibited peak
expression in summer, although a small fraction of genes showed peaked

expression during autumn–winter (e.g. Qa12 and Qg1 in the TPS-a clade in Fig.
3*A*). Notably, expression patterns differed markedly among the three target TPS
subclades i.e., TPS-b-t1, t2, and t3. In leaves, only genes in the TPS-b-t3 clade
showed clear and consistent summer expression peaks (Fig. 3*D*), whereas
expression levels in the TPS-b-t1 and TPS-b-t2 clades were low in all species (Fig.
3*B, C*). These results suggest that TPS-b-t3 genes are primarily responsible for
terpenoid biosynthesis. Furthermore, the expression pattern of *Q, serrata IspS* (28)
within the TPS-b-t3 clade also showed strong expression (Fig. 3*B-D*), providing
additional support for the functional role of this clade in isoprene biosynthesis. In
contrast, in bud tissues, expression levels of genes within the TPS-b-t3 clade were
consistently low, and genes in the other two clades, TPS-b-t1 and t2 were relatively
high (Fig. S6*B-D*), implying a tissue-specific regulation of TPS gene expression.
239
Convergent evolution and positive selection in *IspS* candidates of Fagaceae.
To determine whether the genes with an isoprene score of 4 were gained or lost
during evolution, we employed the ancestral sequence reconstruction (ASR)
method. This approach relies on statistical principles to infer the sequences of
putative ancestral genes based on the amino acid or nucleotide sequences of
extant species. For this analysis, we used nucleotide sequences from 17 Fagales
species belonging to the TPS-b clade (Table S3) and reconstructed putative
ancestral sequences using codeml implemented in PAML
(https://github.com/abacus-gene/paml).
The estimated ancestral sequences of the last common node of the TPS-b-
t1, TPS-b-t2, and TPS-b-t3 clades (node N2; Fig. 4*A*) had an isoprene score of 2,
with the conserved residuals S/V446 and F485 (Fig. 4*B*). Subsequently, N505 was
gained in the common ancestor of the TPS-b-t2 and TPS-b-t3 clades (node N4;
Fig. 4*A, B*), prior to their duplication. Following this duplication, F338 was acquired
independently in each of the TPS-b-t2 and TPS-b-t3 clades (nodes N6 and 8; Fig.
4*A B*), specifically in the lineage leading to the three species within subgenus
Quercus (*Q. robur*, *Q lobata*, and *Q. rubra*).
Notably, the gain of F338 occurred independently in both the TPS-b-t2 and
the TPS-b-t3 clades, despite requiring two nucleotide substitutions for the
transition from ACT (T) to TTT (F). We also employed GRaSP
(https://github.com/charles-abreu/GRaSP) to infer the ancestral sequences and
reached the same conclusion that the gain of F338 occurred independently in both
the TPS-b-t2 and TPS-b-t3 clades (Fig. S7). Although the inferred ancestral
sequences for the distant ancestor at nodes N1–N3 differed slightly between
PAML and GRaSP, the evolutionary pattern across the remaining nodes was
consistent (Fig. S7). These results suggest that the acquisition of isoprene-
producing function occurred through the stepwise accumulation of mutations,
potentially driven by convergent molecular evolution in the TPS-b-t2 and TPS-b-t3
clades.
We next tested for evidence of positive selection in the clades that acquired
F338 (predefined as foreground branches) by applying the branch-site model in
codeml (29, 30). This approach identifies amino acid sites where the
nonsynonymous to synonymous substitution rate ratio (ω) exceeds one, indicative

of positive selection. The analysis revealed significant evidence of positive
selection in the foreground branches (likelihood ratio test, P = 1.027×10⁻⁶) with
four residuals identified as candidates under positive selection (Bayes Empirical
Bayes posterior probability > 0.95) (Table S4; Fig. S8). Notably, the residue
corresponding to F338 was included among these candidates, suggesting that the
convergent gain of this residue in the TPS-b-t2 and TPS-b-t3 clades was likely
driven by positive selection.
280
***in vitro* enzyme assays demonstrated that the *IspS* evolved from**
monoterpene synthase. To evaluate the evolutionary trajectory of enzymatic
function, ancestral TPS sequences were artificially synthesized (Dataset S1) and
subjected to enzymatic characterization using an *Escherichia coli* expression
system. Because genes within the TPS-b-t3 clade exhibited high expression
levels, we focused our analyses on this clade. Enzyme assays using purified
recombinant proteins were conducted with either DMAPP (for IspS activity) or GPP
(for mono-TPS activity) as a substrate. The ancestral enzyme at node N7 exhibited
mono-TPS activity, specifically producing *trans*- and *cis*-β-ocimene, while the
enzyme at node N8 exhibited IspS activity only (Fig. 4*C*, *D*). Ancestral proteins
reconstructed using both PAML and GRaSP showed identical enzymatic activities
at nodes N7 and N8 (Fig. 4*C*, *D*; Fig. S7*C*, *D*). These results indicate that *IspS* with
an isoprene score of 4 evolved from β-ocimene synthases, supporting the
functional shift during evolution.
295
Discussion
Using comparative genomics, ancestral-sequence reconstruction, and *in vitro*
assays of ancestral enzymes, we found that *IspS* candidates with an isoprene
score of 4 arose independently twice in the subgenus *Quercus*—once in the TPS-
b-t3 clade and once in the TPS-b-t2 clade—within the TPS-b subfamily in
Fagaceae. Because the subgenus *Quercus* diverged from its common ancestor
roughly 56 million years ago (31), we infer that the onset of substantial isoprene
emission in Fagaceae to the Eocene. Moreover, in all species we examined (*Q.*
robur, *Q lobata*, and *Q. rubra*), once the IspS activity evolved, it has been retained.
This contrasts with rosids, where a single early gain of isoprene synthase was
followed by multiple independent losses (32). Fagaceae therefore provide a
complementary pattern—multiple, convergent gains coupled with long-term
retention—highlighting diverse evolutionary routes to isoprene emission in
angiosperms.
The early TPS-b-t3 proteins preferred GPP and mainly produced β-ocimene,
whereas their descendants evolved to favour DMAPP and became dedicated
isoprene synthases in the subgenus *Quercus*. The shift would correlate with
pocket-narrowing substitutions—most notably the substitution from Thr to Phe at
the 338 residue—that exclude the larger GPP substrate. These results confirmed
the previous view in which the ocimene synthases is ancestral TPS for which IspS
evolved by active site size reduction following acquisition of F338 (33). The
combination of branch-level and site-level likelihood tests detected positive
selection at this residue, indicating adaptive convergence toward isoprene

emission. We also identified three additional residues under positive selection, suggesting that other sites may contribute to isoprene emission, although the effects of these sites on substrate specificity are currently unknown. Isoprene provides a threefold heat shield—membrane stabilization, ROS scavenging, and Ca^{2+}-mediated heat-shock signaling (34)—mechanisms that likely protect photosynthesis and sustain growth in *Quercus* species with thin, deciduous leaves under rising temperatures.

Terpenoids represent the largest and most diverse class of natural products in terrestrial plants (35). Comparative and functional genomics can chart the continuous evolutionary innovation of terpenoid-biosynthetic pathways with unprecedented resolution. Such insight will deepen our understanding of how chemical diversity has shaped plant adaptation and diversification on the earth.

Materials and Methods
Genome synteny analysis
To compare the genome structures of Fagaceae species across genera, published genome sequences of eight Fagaceae species from four genera were obtained from the public databases and used for subsequent analyses (Table S1). We first performed an all-vs-all BLASTp search using protein sequences with a cut-off e-value of 1E-5 to identify homologous relationships across the genomes. To investigate genomic synteny, we detected syntenic blocks within or between genomes using MCScanX (https://github.com/wyp1125/MCScanX), based on the results of the above BLASTp searches. The synteny blocks were visualized using SynVisio (https://synvisio.github.io). To clarify syntenic relationships across chromosomes in each species, we renumbered chromosomes according to the chromosomal order in *Q. robur*. Additionally, chromosomes oriented opposite to those of *Q. robur* were reversed. These adjusted chromosomes were used for subsequent analyses.

Phylogenetic analysis of TPS genes
Alignment and phylogenetic tree construction were conducted using the build function of ETE3 v3.1.3 (https://etetoolkit.org). Sequence alignment was performed with MAFFT v6.861b (https://mafft.cbrc.jp), alignment trimming with trimAl v1.4.rev (https://vicfero.github.io/trimal), and phylogenetic tree construction with RAxML v8.2.11 (https://cme.h-its.org/exelixis/web/software/raxml) using the GAMMA JTT model and 500 bootstrap replications. The resulting phylogenetic tree was visualized using the online tool iTOL (https://itol.embl.de). The analyzed TPS genes were classified into five subclade groups: TPS-a, TPS-b, TPS-c, TPS-e/f, and TPS-g, based on reference genes from Arabidopsis and tomato. The chromosomal locations of these classified genes were visualized using TBtools-II v2.2.10 (https://github.com/CJ-Chen/TBtools-II).

Identification of TPS family members
We retrieved candidate genes of the TPS family based on two conserved motifs: the metal-binding domain (PF03936) and the N-terminal domain (PF01397). Protein sequences from eight Fagaceae species were used for a domain search

against the Pfam-A database ver 35.0 (36) using HMMER ver 3.3.2
(http://hmmer.org) with a cut-off e-value of 1E-5. For genes with more than two
isoforms, the longest isoform was selected.
To exclude candidate genes with short sequence lengths, we utilized the
conserved exon-intron arrangement patterns found in TPS genes (26). Across a
broad range of plant species, these patterns are well-documented and categorized
into three classes: Class I (12–14 introns), Class II (9 introns), and Class III (6
introns) (26). Notably, Class II TPS genes are specific to gymnosperms, meaning
that only Class I and Class III TPS genes are present in the angiosperm species
analyzed in this study. To identify putative TPS genes, we generated histograms
depicting the intron count for each gene within each species. The majority of genes
exhibited exon-intron patterns consistent with either Class I or Class III (Fig. S1).
Based on this distribution, we selected genes containing more than five exons.
Furthermore, some genes remained too short for detailed analysis. To
address this, genes encoding proteins shorter than 400 amino acids were excluded
(Fig. S2). To classify the subgroups of TPS genes, we incorporated protein
sequences of well-characterized TPS genes from *S. lycopersicum* and *A. thaliana*
as references for phylogenetic analysis. These reference sequences were
obtained from Table S7 in Zhou and Pichersky (27).
Additionally, *IspS* and *IspS-like* from *Q. serrata* were obtained from previous
study (28) and included in the analysis. To root the phylogenetic tree, we included
the *ent-copalyl diphosphate synthase* sequence from *Picea sitchensis* as an
outgroup.
388

Phylogenetic analysis of TPS genes
Alignment and phylogenetic tree construction were conducted using the build
function of ETE3 v3.1.3 (https://etetoolkit.org). Sequence alignment was
performed with MAFFT v6.861b (https://mafft.cbrc.jp), alignment trimming with
trimAl v1.4.rev (https://vicfero.github.io/trimal), and phylogenetic tree construction
with RAxML v8.2.11 (https://cme.h-its.org/exelixis/web/software/raxml) using the
GAMMA JTT model and 500 bootstrap replications. The resulting phylogenetic tree
was visualized using the online tool iTOL (https://itol.embl.de). The analyzed TPS
genes were classified into five subclade groups: TPS-a, TPS-b, TPS-c, TPS-e/f,
and TPS-g, based on reference genes from Arabidopsis and tomato. The
chromosomal locations of these classified genes were visualized using TBtools-II
v2.2.10 (https://github.com/CJ-Chen/TBtools-II).
401

Isoprene synthase gene analysis
To investigate molecular evolution of *IspS* in Fagaceae, we utilized published *IspS*
sequences from *S. lycopersicum* (27) and *Q. serrata* (28), both belonging to the
TPS-b subfamily. Since an adjacent subclade was located near the subclades
containing these sequences, we focused on the three subclades that include 44
Fagaceae genes.
These genes of three subclades were aligned by MAFFT v7.520
(https://mafft.cbrc.jp) with the reference *IspS* sequence from *Populus alba*

(AB198180) (37), *S. lycopersicum* (27) and *Q. serrata* (28) to calculate an isoprene
score (15).
412
Sample collection for RNA sequencing
To evaluate expression of TPS genes in natural conditions, we analyzed
transcriptome data obtained from leaf and bud tissues in three Fagaceae species,
Q. acutissima, *Q. glauca*, and *L. edulis*. We generated data for *Q. acutissima*,
whereas those for *Q. glauca* and *L. edulis* were obtained from the previous
study (25). For all data, samples were collected every four weeks spanning from
2020–2021 in *Q. acutissima*, 2022–2023 in *Q. glauca* and 2019–2021 in *L. edulis*
in Fukuoka prefectures in Japan. For each species, we targeted three individuals
and each sample was collected from three branches per individual, except for *L.*
edulis. Given the consistent expression patterns observed among individuals in a
previous study (24), we focused on only one individual for *L. edulis*. Study sites for
Q. acutissima and *Q. glauca* is the biodiversity reserve on the Ito campus of
Kyushu University (33º35′ N, 130º12′ E, 20 to 57 m a.s.l). *L. edulis* were studied at
the Imajyuku Field Activity Center (33º33′ N, 130º16′ E, 84 to 111 m a.s.l). Leaf
samples were collected from spring to summer in the deciduous species *Q.*
acutissima, whereas in the evergreen species *Q. glauca* and *L. edulis*, sampling
was conducted throughout the year.
RNA-seq samples were collected and preserved in 2 ml microtubes
containing 1.5 ml of RNA-stabilizing reagent (RNAlater; Thermo Fisher Scientific,
Waltham, MA, USA), then stored at −80°C until RNA extraction, following the
protocol described by Kudo et al. (25).
434
RNA extraction and RNA-seq analysis
Total RNA was extracted from leaf samples following the protocol described
previously (38). RNA integrity was assessed using the method outlined previously
(25). For each sample, 5–6 µg of total RNA was sent to Hangzhou Veritas Genetics
Medical Institute Co., Ltd., where cDNA libraries were prepared using the NEBNext
Ultra II RNA Library Prep Kit for Illumina. Transcriptome sequencing (150 bp
paired-end reads) was performed on an Illumina NovaSeq 6000 platform (Illumina,
San Diego, CA, USA). Gene expression quantification followed a three-step
pipeline: (1) quality filtering was performed using fastp v0.20.0 with default settings
(https://github.com/OpenGene/fastp); (2) RNA-seq reads were aligned to the
reference genome using TopHat v2.1.1
(https://ccb.jhu.edu/software/tophat/index.shtml); and (3) transcript abundance
was quantified using StringTie v1.3.5 (https://ccb.jhu.edu/software/stringtie).
448
Seasonal expression patterns of TPS genes in Fagaceae
To investigate the seasonal expression patterns of TPS genes, we extracted TPS
gene transcripts from RNA-seq data of *Q. acutissima*, *Q. glauca*, and *L. edulis*. For
each gene, we calculated the mean expression level in transcripts per million
(TPM) across three individuals, except for *L. edulis*, for which data were available
from only one individual (25). TPM values were transformed using $\log_2$(TPM +
0.01). Genes with a mean TPM > 1 across all samples were considered to be

expressed. A heatmap was generated in R ver. 4.3.1 using the 'pheatmap'
package, with gene order arranged according to TPS clades (Fig. 3*A*). In addition
to these three species, we also obtained expression data for *IspS* and *IspS*-like in
Q. serrata (28), which were used to plot seasonal changes in expression levels.
To generate the heatmap, we calculated the average gene expression levels for
each month by aggregating data from different individuals over a two-year period
for each species.
463
Ancestral amino acid sequence inference
To infer the ancestral amino acid sequences of putative *IspS* in *Quercus* species,
we analyzed 95 representative nucleotide sequences, which have no ambiguity,
of TPS-b subfamily members from Fagales. These sequences include genes
identified through the genomic survey of eight Fagaceae species described above,
genes from *Q. serrata*, and additional genes retrieved via BLAST searches against
the database. The BLAST search identified genes from *Q. suber*, *L. litseifolius*,
Castanea mollissima, *Juglans regia*, *J. maicrocarpa x J. regida*, *Carya illinoinensis*,
Corylus avellana, and *Alnus glutinosa*, resulting in a dataset comprising genes
from 17 Fagales species. (Table S3). Additionally, seven TPS-g subfamily genes
were used as an outgroup.
Sequence alignment was performed using MACSE v2 (https://www.agap-
ge2pop.org/macse), which can account for the underlying codon structure of
protein-coding sequences and generate both nucleotide (NT) and amino acid (AA)
alignments. Phylogenetic reconstruction and ancestral amino acid sequence
inference were conducted separately for NT and AA alignments. ML phylogenetic
trees were constructed with RAxML v8.2.11 (https://cme.h-
its.org/exelixis/web/software/raxml) with 1000 bootstrap replications. The GAMMA
GTR model (GTR+I+G) and the GAMMA JTT model (JTT+I+G) were used for the
NT and AA alignment, respectively. The resulting ML trees exhibited compatible
topology, particularly the splits of the major clusters. Two strongly supported
clusters of *IspS* with the maximum score were identified in both trees. Ancestral
amino acid sequences were inferred at each internal node using codeml in PAML
(https://github.com/abacus-gene/paml) with the F3x4 codon model for the NT
alignment. We also employed GRaSP (https://github.com/charles-abreu/GRaSP)
to infer the ancestral sequences from the AA alignment and estimate the location
of gaps in the ancestral sequences.
491
Detection of positive selection
To investigate whether amino acid sites in putative *IspS* (foreground) have
undergone positive selection compared with the other TPS-b subfamily members
(background), we applied the branch-site test for positive selection (29). The NT
alignment and MLtree used for the ancestral sequence inference were employed
in this analysis. Alignment gaps were removed prior to the analyses. Previous
studies have shown that gene conversion and recombination events could lead to
false positives (39, 40). To account for this, we applied the GARD method
implemented in HYPHY (https://hyphy.org) to detect potential gene conversion or
recombination events. No significant evidence of gene conversion or

recombination was found in the genes, so subsequent analyses were conducted
using the full-length alignments.
All branches within two clades of the genes with the isoprene score of 4
were set as foreground branches and the branch-site tests were conducted using
codeml (30). The null model was set to model = 2, NSsites = 2, fix_omega = 1 and
omega = 1, and the alternative model (Model A) was with model = 2, NSsites = 2
and fix_omega = 0 (to estimate ω_2). The Bayes empirical Bayes (BEB) method
was used to identify sites potentially under positive selection on the foreground
branches.
511
Materials and reagents for *in vivo* enzymatic characterization
DMAPP and GPP were used as substrates, with DMAPP purchased from
TargetMol (Boston, USA) and GPP from Sigma-Aldrich (St. Louis, USA). As
standard compounds, isoprene was obtained from Tokyo Chemical Industry Co.,
Ltd. (Tokyo, Japan), and β-ocimene was purchased from Cayman Chemical (Ann
Arbor, USA).
518
Construction of *E. coli* expression vectors
To express the ancestral sequences estimated at nodes N7 and N8 by PAML
(*N7_paml* and *N7_paml*) and GRaSP (*N7_grasp* and *N8_grasp*) in *E. coli*, their
putative transit peptides at N-terminus were removed and the truncated CDSs
were further codon-optimized for *E. coli*. The optimized sequences were
synthesized by GeneArt (Thermo Fisher Scientific). These sequences were
amplified by PCR using KOD One® (Toyobo, Osaka, Japan) with the primer pairs
shown in Table S5. The amplification program is composed of an initial
denaturation at 94°C for 1 min followed by 40 cycles of denaturation at 98°C for
10 sec, annealing at 51°C for 5 sec, and extension at 68°C for 10 sec, with a final
extension at 68°C for 5 min. The amplicons were individually inserted into
pET22b(+) vector (Novagen, Madison, US) by In-Fusion reaction (Takara Bio,
Kusatsu, Japan) using NdeI and BamHI sites, which enables the production of
proteins with the histidine tag (6×His). The sequence of *N8_paml* was constructed
by introducing a single mutation to *N8_grasp* by PCR using KOD One® (Toyobo)
and the primer pair *N8_paml_*Fw and *N8_paml_*Rv. This amplification was carried
out with an initial denaturation at 94°C for 1 min, followed by 40 cycles of
denaturation at 98°C for 10 sec, annealing at 51°C for 5 sec, and extension at
68°C for 40 sec, with a final extension at 68°C for 5 min. The amplicon was
introduced into the pET22b(+) vector in the same process as described above.
539
Enzymatic characterization
Constructs *ΔTP-N7_paml-6×His*, *ΔTP-N7_grasp-6×His*, *ΔTP-N8_paml-6×His*,
and *ΔTP-N8_grasp-6×His* were individually introduced into *E. coli* BL21 (DE3) for
recombinant protein production. Protein preparation was performed as previously
described (Koita et al.) (28). Protein concentrations were measured using a Qubit®
2.0 Fluorometer and the Qubit Protein Assay Kit (Thermo Fisher Scientific).
A standard reaction mixture (200 µl) contained 0.5 mM DMAPP or GPP, 20
mM $MgCl_2$, and 160 µl of purified enzyme. For assays using DMAPP, 27–32 µg of

enzyme were used, whereas 1.4–1.7 µg were used for GPP-based assays.
Reactions were incubated at 37 °C for 30 minutes in 2 ml glass vials (Thermo
Fisher Scientific). GC-MS analysis of the enzymatic products was carried out as
described in Koita et al. (28).
552

553 **Data availability**
554 The RNAseq data are available from the PRJDB35868.
555

Acknowledgments
We thank Kayoko Ohta for her technical support with RNA extraction. We also
thank Koharu Yamaguchi, Ryota Aoyagi, and Ryosuke Imai for their help with
sample collection for RNAseq, Sangwan Kim for his support with registration of
RNA sequence data to DDBJ, and Kenji Fukushima for his insightful comments on
the bioinformatic analyses. This study was supported by grant-in-Aid for
Transformative Research Areas "Plant-Climate Feedback" (no. JP23H04965 to
A.S., K.Y., R.M., S.I., H.H; JP23H04966 to A.S., JP23H04967 to K.Y., R.M).
564

References
1. J. Kesselmeier, M. Staudt, Biogenic volatile organic compounds (VOC): An
overview on emission, physiology and ecology. *J. Atmos. Chem.* **33**, 23-88
(1999).

2. J. Laothawornkitkul, J. E. Taylor, N. D. Paul, C. N. Hewitt, Biogenic volatile
organic compounds in the Earth system. *New Phytologist.* **183**, 27-51 (2009).

3. A. Satake, *et al.*, Plant Molecular Phenology and Climate Feedbacks
Mediated by BVOCs. *Annu Rev Plant Biol.* **75**, 605-627 (2024).

4. M. Antonelli, *et al.*, Forest volatile organic compounds and their effects on
human health: A state-of-the-art review. *Int. J. Environ. Res. Public Health.*
17, 6506 (2020).

5. N. Lee Ng, *et al.*, Nitrate radicals and biogenic volatile organic compounds:
Oxidation, mechanisms, and organic aerosol. *Atmos Chem Phys.* **17**, 2103-
2162 (2017).

6. A. Guenther, A global model of natural volatile organic compound emissions.
J. Geophys. Res.:Atmos. **100**, 8873-8892 (1995).

7. A. B. Guenther, *et al.*, The Model of Emissions of Gases and Aerosols from
Nature version 2.1 (MEGAN2.1): an extended and updated framework for
modeling biogenic emissions. *Geosci. Model Dev.* **5**, 1471–1492 (2012).

8. P. C. Harley, R. K. Monson, M. T. Lerdau, Ecological and evolutionary
aspects of isoprene emission from plants. *Oecologia.* **118**, 109-123 (1999).

9. R. K. Monson, R. T. Jones, T. N. Rosenstiel, J. P. Schnitzler, Why only some
plants emit isoprene. *Plant Cell Environ.* **36**, 503–516 (2013).

10. K. G. S. Dani, I. M. Jamie, I. C. Prentice, B. J. Atwell, Evolution of isoprene
emission capacity in plants. *Trends Plant Sci.* **19**, 439-446 (2014).

11. B. Miller, C. Oschinski, W. Zimmer, First isolation of an isoprene synthase
gene from poplar and successful expression of the gene in Escherichia coli.
Planta. **213**, 483-487 (2001).

12. D. T. Hanson, S. Swanson, L. E. Graham, T. D. Sharkey, Evolutionary
significance of isoprene emission from mosses. *Am. J. Bot.* **86**, 634–639
(1999).

13. D. T. Tingey, R. C. Evans, E. H. Bates, M. L. Gumpertz, Isoprene emissions
and photosynthesis in three ferns – The influence of light and temperature.
Physiol. Plant. **69**, 609-616 (1987).

14. P. Harley, *et al.*, Variation in potential for isoprene emissions among
Neotropical forest sites. *Glob Chang Biol.* **10**, 630-650 (2004).

15. T. D. Sharkey, D. W. Gray, H. K. Pell, S. R. Breneman, L. Topper, Isoprene
synthase genes form a monophyletic clade of acyclic terpene synthases in
the Tps-b terpene synthase family. *Evolution.* **67**, 1026–1040 (2013).

16. G. M. Silver, R. Fall, Characterization of aspen isoprene synthase, an
enzyme responsible for leaf isoprene emission to the atmosphere. *J. Biol.*
Chem. **270**, 13010-13016 (1995).

17. Q. Jia, *et al.*, Origin and early evolution of the plant terpene synthase family.
Proc. Natl. Acad. Sci. U.S.A. **119**, e2100361119 (2022).

18. M. Li, *et al.*, In Planta Recapitulation of Isoprene Synthase Evolution from
Ocimene Synthases. *Mol. Biol. Evol.* **34**, 2583–2599 (2017).

19. T. Kawakami, S. Miyazaki, H. Kawaide, Molecular characterization of a moss
isoprene synthase provides insight into its evolution. *FEBS Lett.* **597**, 2133–
2142 (2023).

20. F. Loreto, Distribution of isoprenoid emitters in the Quercus genus around
the world: Chemo-taxonomical implications and evolutionary considerations
based on the ecological function of the trait. *Perspect. Plant Ecol. Evol. Syst.*
5, 185-192 (2002).

21. A. Tani, N. Masui, T. W. Chang, M. Okumura, Y. Kokubu, Basal emission
rates of isoprene and monoterpenes from major tree species in Japan:
interspecies and intraspecies variabilities. *Prog. Earth Planet. Sci.* **11**, 42
(2024).

22. F. Loreto, S. Fineschi, Reconciling functions and evolution of isoprene
emission in higher plants. *New Phytol.* **206**, 578-582 (2015).

23. A. Kremer, *et al.*, Genomics of Fagaceae. *Tree Genet. Genomes.* **8**, 583-610
(2012).

24. A. Satake, K. Ohta, N. Takeda-Kamiya, K. Toyooka, J. Kusumi, Seasonal
gene expression signatures of delayed fertilization in Fagaceae. *Mol. Ecol.*
32, 4801-4813 (2023).

25. S. N. Kudo, *et al.*, The evolution of gene expression in seasonal
environments. bioRxiv [Preprint] (2025).
http://biorxiv.org/lookup/doi/10.1101/2025.03.25.645377 (accessed 20 April
2025).

26. S. C. Trapp, R. B. Croteau, Genomic Organization of Plant Terpene
Synthases and Molecular Evolutionary Implications. *Genetics.* **158**, 811–832.
(2001).

27. F. Zhou, E. Pichersky, The complete functional characterisation of the
terpene synthase family in tomato. *New Phytol.* **226**, 1341–1360 (2020).

28. S. Koita, *et al.*, Molecular basis behind the isoprene emission diversity in
Fagaceae. bioRxiv [Preprint] (2025).
http://biorxiv.org/lookup/doi/10.1101/2025.05.17.654689 (accessed 20 May
2025).

29. J. Zhang, R. Nielsen, Z. Yang, Evaluation of an improved branch-site
likelihood method for detecting positive selection at the molecular level. *Mol.*
Biol. Evol. **22**, 2472–2479 (2005).

30. Z. Yang, PAML 4: Phylogenetic analysis by maximum likelihood. *Mol. Biol.*
Evol. **24**, 1586–1591 (2007).

31. B. F. Zhou, *et al.*, Phylogenomic analyses highlight innovation and
introgression in the continental radiations of Fagaceae across the Northern
Hemisphere. *Nat. Commun.* **13**, 1320 (2022).

32. J. Shang, *et al.*, Evolution of the biosynthetic pathways of terpene scent
compounds in roses. *Curr. Biol.* **34**, 3550-3563.e8 (2024).

33. D. W. Gray, S. R. Breneman, L. A. Topper, T. D. Sharkey, Biochemical
characterization and homology modeling of methylbutenol synthase and
implications for understanding hemiterpene synthase evolution in plants. *J.*
Biol. Chem. **286**, 20582-20590 (2011).

34. Z. Zuo, S. M. Weraduwage, T. Huang, T. D. Sharkey, How volatile
isoprenoids improve plant thermotolerance. *Trends Plant Sci.*, in press. doi:
10.1016/J.TPLANTS.2025.05.004.

35. F. Chen, D. Tholl, J. Bohlmann, E. Pichersky, The family of terpene
synthases in plants: A mid-size family of genes for specialized metabolism

that is highly diversified throughout the kingdom. *Plant J.* **66**, 212–229
(2011).

36. J. Mistry, *et al.*, Pfam: The protein families database in 2021. *Nucleic Acids*
Res. **49**, D412–D419 (2021).

37. K. Sasaki, K. Ohara, K. Yazaki, Gene expression and characterization of
isoprene synthase from Populus alba. *FEBS Lett.* **579**, 2514-2518 (2005).

38. Y. Miyazaki, *et al.*, Nitrogen as a key regulator of flowering in Fagus crenata:
Understanding the physiological mechanism of masting by gene expression
analysis. *Ecol. Lett.* **17**, 1299-1309 (2014).

39. M. Anisimova, R. Nielsen, Z. Yang, Effect of recombination on the accuracy
of the likelihood method for detecting positive selection at amino acid sites.
Genetics. **164**, 1229-1236 (2003).

40. C. Casola, M. W. Hahn, Gene conversion among paralogs results in
moderate false detection of positive selection using likelihood methods. *J.*
Mol. Evol. **68**, 679–687 (2009).

676
677

Figures and Tables

Figure 1. Syntenic relationships of Fagaceae genomes and the phylogenetic analysis of TPS genes. (*A*) Syntenic relationships among eight Fagaceae genomes. The phylogenetic tree was constructed based on whole-genome sequences from eight Fagaceae species, with bootstrap values shown at each branch. Chromosomes of each species are arranged along the corresponding branches of the phylogenetic tree. Syntenic relationships are illustrated with black and gray lines, where black indicates reverse alignments and gray represents standard (collinear) alignments. Chromosome numbers are labeled accordingly.

(*B*) Molecular phylogenetic tree of TPS genes in the eight Fagaceae species. TPS
genes from *S. lycopersicum* and *A. thaliana* were used as references to define five
TPS subclades (TPS-a, -b, -c, -e/f, and -g). The three TPS-b genes in tomato
including isoprene synthase were shown by red asterisk. The *ent*-copalyl
diphosphate (*ent*-CPP) synthase gene from *Picea sitchensis* was used as the
outgroup. The branches whose bootstrap values were 0.9 or more were shown by
purple circles.
696
697

Figure 2. Phylogenetic tree and multiple sequence alignment of TPS genes in the TPS-b Subclade. (*A*) Molecular phylogenetic tree of TPS genes classified within the TPS-b subclade. The phylogeny was inferred with RAxML v8.2.11 under the GAMMA JTT substitution model, using 500 bootstrap replicates to assess node support. A TPS-g gene from *S. lycopersicum* (TPSgSlTPS37) was used as an outgroup. Two genes previously characterized with high isoprene synthase activity—score 3 in *S. lycopersicum* and score 4 in *Q. serrata*—are marked with red stars. (*B*) Multiple sequence alignment of TPS genes from the three TPS-b subclades: TPS-b-t1, TPS-b-t2, and TPS-b-t3. Four conserved amino acid positions used to compute isoprene scores are highlighted by red and yellow

709   shading, red means conserved amino acid at this position and yellow means
710   another. Genes with the highest isoprene score (score 4) are indicated by asterisks.
711
712

713
Figure 3. Seasonal expression patterns of TPS genes in leaves. (*A*) Heatmap
showing the seasonal expression patterns of TPS genes in three Fagaceae
species: *Q. acutissima* (blue), *Q. glauca* (light blue), and *L. edulis* (green). TPS-c
genes are not included in the heatmap, as no TPS-c genes were consistently
expressed in all three species. Gray squares indicate missing data due to leaf
shedding. Genes were considered expressed if the mean of TPM values across
all observation periods was ≥1. (*B-D*) Seasonal expression patterns of TPS genes
within the subclades TPS-b-t1, TPS-b-t2, and TPS-b-t3. In addition to the three
species mentioned above, data from *Q. serrata* were incorporated from a previous
study (Koita et al. 2025).
724
725

Figure 4. Results of ancestral sequence reconstruction of Fagaceae TPS genes
using PAML. (*A*) Maximum-likelihood phylogeny of the TPS gene set used for
ancestral reconstructions. The phylogeny was inferred with RAxML v8.2.11 under
the GAMMA GTR substitution model, using 1,000 bootstrap replicates to assess
node support. Internal nodes N1–N8 mark the bifurcation points analysed. Branch
colours denote the inferred isoprene score—blue (2), purple (3) and red (4). The
three focal clades (TPS-b-t1, TPS-b-t2 and TPS-b-t3) are highlighted in distinct
colours for ease of reference. (*B*) Inferred amino-acid states at the diagnostic
tetrad and the corresponding isoprene score for each ancestral node (N1–N8). The
posterior probability (PP) of the most likely codon/amino-acid state was as follows:
N1 – S/V446: GCA (Ala, PP = 0.263) → GTA (Val, PP = 0.600); F485: CTT (Leu,
PP = 0.523) → TTT (Phe, PP = 0.405). N4 – T505: AAA (Lys, PP = 0.774) → AAT
(Asn, PP = 0.956). N6 – F338: ACT (Thr, PP = 0.999) → TTT (Phe, PP = 0.970).
N8 – F338: ACT (Thr, PP = 0.996) → TTT (Phe, PP = 0.990). (*C*) and (*D*) Enzyme
assays of the ancestral sequence products using DMAPP and GPP as substrates.
The reaction products were detected at *m/z* 67 for isoprene (*C*) and *m/z* 93 for
monoterpenes (*D*) as the substrate. The *cis* or *trans* configurations of β-ocimene
were identified based on spectral annotation. The chromatograms, except for the

745 β-ocimene standard, are plotted on the same scale. Asterisks represent peaks of
746 impurities.
747
748
749 **Additional files**
750
751 **Supplementary file 1**
752 Supplementary figures (Fig. S1 to Fig. S8) and supplementary tables (Table S1 to
753 Table S5) were included in this file.
754
755 **Supplementary Dataset**
756 **Dataset S1**
757 Multiple sequence alignment of ancestral sequences of Fagaceae TPS genes
758 using PAML and GRaSP. The four conserved amino acid residues used for
759 calculating isoprene score were shown by black frame.
760