Title: Larger hybrid clutch size could drive regional displacement of native *Iguana* populations
across the Lesser Antilles

Short running title: Hybrids displace native *Iguana* populations

Matthijs P. van den Burg[1,2,*], Sarah A. van der Horn[3], P-A Åhlen[4], Lara Jansen[5], Kai Wulf[6], Misha
Paauw[7], and Adolphe O. Debrot[5,*]

[1] IUCN SSC Iguana Specialist Group, Gland, Switzerland

[2] Burg Biologica, The Hague, The Netherlands

[3] Nature and Environmental Policy team, Public Entity Saba, Saba, Caribbean Netherlands

[4] Swedish Association for Hunting and Wildlife Management, Nyköping, Sweden

[5] Marine Animal Ecology group, Wageningen University and Research, P.O. Box 338, 6700AH
Wageningen, The Netherlands

[6] Saba Conservation Foundation, Saba, Caribbean Netherlands

[7] Green Life Sciences, Swammerdam Institute for Life Sciences, University of Amsterdam,
Amsterdam, The Netherlands

* Corresponding authors: thijs.burg@gmail.com, dolfi.debrot@wur.nl

Abstract

Native *Iguana* populations across the Caribbean Lesser Antilles are being replaced by non-native
green iguanas (NNGI) once these establish themselves, leading to eventual extinction. Despite this
knowledge, and much effort towards NNGI removal and prevention of incursions, the underlying
mechanisms of the observed genetic swamping are poorly understood. Previous studies show that
larger clutch size is a key reproductive trait of mainland as opposed to insular lizard populations.
Higher clutch size in NNGI and their progeny could be a driver in the genetic swamping of insular
Iguana populations. We collected data on snout-vent length, tail length, clutch size and genetics
opportunistically during an eradication program of hybrid and NNGIs on the island of Saba, home to
a native *Iguana iguana* population. Data of native and hybrid iguanas from Saba were combined with
similar published data from native insular and mainland *Iguana* populations. Generalized linear
modeling revealed that climatic variables could not explain differences in clutch size, in contrast to
genetic origins. Our preliminary results show that hybrid iguanas lay significantly more eggs (~40%)
than similar-sized native Saba iguanas. Therefore, we suggest that higher hybrid clutch size may be a
contributing factor hastening the genetic swamping and declines in native *Iguana* populations across
the Lesser Antilles. Further study is needed to clarify and better understand the drivers of the rapid

genetic swamping and loss of Antillean iguana diversity, which should also focus on *Iguana*

delicatissima and utilize genomic approaches to pinpoint genes under selection.

Keywords: Competition, Displacement, Hybridization, Iguana iguana, Lesser Antilles.

Introduction

Global biodiversity is threatened significantly by invasive alien species (IAS) that negatively impact

native species through a range of processes like resource competition, introduction of diseases,

predation, and hybridization (e.g., Allendorf et al. 2001; Strubbe and Matthysen 2009). Hybridization

and introgression with IAS (i.e. anthropogenic hybridization; Allendorf et al. 2001) can have different

diminishing impacts on native species (e.g., loss of genetic diversity, wasted reproductive effort, see

Allendorf et al. 2001), and has even led to local extirpations (Rhymer & Simberloff 1996; Todesco et

al. 2016). Conservationists are especially challenged when hybridization occurs between a rare

species and closely-related IAS. Not only because this process can easily go undetected (e.g.,

Vanhaecke et al. 2012; Hata et al. 2019), but also since protecting the rare endangered species

especially requires efforts to prevent hybridization. However, the mechanisms of decline due to

hybridization remain poorly understood.

Native iguanas within the genus *Iguana* were once present across islands in the Caribbean

Lesser Antilles, but have been either extirpated from numerous islands or have dramatically

decreased in population size due to a range of direct and indirect anthropogenic causes (Knapp et al.

2014, 2021; Vuillaume et al. 2015; van den Burg et al. 2018a; Breuil et al. 2019). Within this insular

region the current most severe threat to remaining native *Iguana* populations is the region-wide

presence of non-native green iguanas (NNGI) (Breuil et al. 2019; Knapp et al. 2021; van den Burg et

al. 2023). Their initial arrival to the region on the islands of Les Saintes dates to the 1860's (Vuillaume

et al. 2015), while more recently, natural (hurricanes, Censky et al. 1998) and unnatural (e.g., as

stowaways in cargo containers) intra-regional translocations have occurred resulting in the

establishment of non-native populations (van den Burg et al. 2018b, 2020).

Impacted native iguanas include *Iguana delicatissima* Laurenti, 1768 (from Anguilla to

Martinique, except Saba and Montserrat) and members of the *Iguana iguana* species complex

(Linnaeus, 1758) (sensu Iguana Taxonomy Working Group, 2016). Considering the entire *I. iguana*

complex, these populations belong to one of the four major mtDNA clades (named clade IIb in

Stephen et al. 2013; *I. iguana* in van den Burg et al. 2021; clade IV van den Burg et al. 2026).

Following a number of recent manuscripts on the taxonomy of these insular Lesser Antillean *I.*

iguana populations (Breuil et al. 2019, 2020, 2022), currently the subspecies of *I. iguana insularis* (St.

Vincent and the Grenadines and Grenada) and *I. iguana sanctaluciae* (St. Lucia) are recognized, but

not the species *I. melanoderma* (also not recognized as subspecies; Saba and Montserrat) (Iguana

Taxonomy Working Group, 2022; van den Burg et al. 2026). Additionally, the Iguana Taxonomy

Working Group (2022) currently recognizes *I. iguana iguana* and *I. iguana rhinolopha*; see geographic

limitation for *I. i. rhinolopha* in van den Burg et al. (2021a).

Non-native green iguanas within the Lesser Antilles have a diverse origin, which have been

traced through genetic assessments of NNGIs from several islands using geographic referenced

datasets from the native range (see van den Burg et al. 2026 for clade codes). These studies indicate

origins from all four mtDNA clades, especially from Central America (Clade III) and northwestern

South America (Clade II), with only few non-native populations including iguanas with (partial) origins

from eastern South America (Clade IV) and the Curaçao clade (Clade I) (Vuillaume et al. 2015; van

den Burg et al. 2018b, 2023, 2024; Pounder et al. 2020). However, a complete overview is still

lacking, as several islands lack assessments or include only limited analyzed samples.

The arrival of NNGIs to Lesser Antillean islands has repeatedly led to anthropogenic

hybridization and introgression with native *Iguana* populations (Day and Thorpe 1996; Vuillaume et

al. 2015; van den Burg et al. 2018a; Breuil et al. 2019). Since this process leads to genetic swamping,

including complete admixture without selection against hybrids with eventual extinction of the

native population (Allendorf et al. 2001), much effort is being invested, through morphological

(Breuil 2013; van den Burg et al. 2023, 2024) and genetic methods (Day and Thorpe 1996; Valette et

al. 2013; Pounder et al. 2020; van den Burg et al. 2021a; van Kuijk et al. 2025), to rapidly identify and

remove NNGIs. However, the mechanisms behind genetic swamping have received less research

attention and are mostly limited to anecdotal data from *I. delicatissima* populations invaded by

NNGIs. Vuillaume et al. (2015) indicate that hybrid and introgressed iguanas are "bigger and longer"

than *I. delicatissima* and non-native males are more "powerful," but without providing data. Using

morphometric data, van den Burg et al. (2024) showed that NNGI on St. Maarten/Martin attain

larger overall size compared to *I. delicatissima* on St. Eustatius, which also holds for native green

iguanas from Saba (*I. i. iguana*; van den Burg et al. 2026), but a broader geographic study is needed

to confirm this as there is variation in both parental species (Day and Thorpe 1996; Bock et al. 2018).

Whether native X non-native hybrids also attain larger overall size compared to native iguanas has

remained unassessed. Additionally, Vuillaume et al. (2015) suggested that the difference in

reproductive timing between native and non-native iguanas could also play a role.

Clutch size has long been recognized as a key life-history trait affecting fitness and

competitive ability (Godfray et al. 1991). While clutch size is often also correlated to other variables

like egg weight and hatchling mass, these relationships are largely overrated, even in bird guilds

where the relationship between clutch and egg parameters has been most intensively studied

(Rohwer 1988). Thus while clutch size has several other correlates of significance like egg size and

brood frequency, the key difference in reproductive traits between mainland and insular lizard

populations is the difference in size-specific clutch size (Novosolov and Meiri 2013; Meiri et al. 2020).

Hence, considering the observed genetic swamping of native island *Iguana* populations and thus a

decline in future population size, clutch size is an important life-history trait to understand, as it

directly reflects reproductive output and can influence future population growth. Especially, insights

on potential differences between native and non-native iguanas and their hybrids are of great

interest to understand future population trends and potential implications of biological invasions.

Whilst it is known that *I. delicatissima* lays fewer eggs than (most of the) populations in the *I. iguana*

species complex (Knapp et al. 2016; Bock et al. 2018). Very little data is available from native

populations of the *I. iguana* species complex from islands in the Lesser Antilles (but see Blankenship

1990). Regarding hybridization, only a single published record of clutch size for an F1 *I. delicatissima* x

I. iguana that suggests a larger clutch size in hybrids compared to *I. delicatissima* (see van

Wagensveld and van den Burg 2018 and references therein). To contribute to a better understand the

underlying mechanism(s) of the regional competitive displacement of native insular *Iguana*

populations, we studied the difference in clutch size between native, invasive and hybrid *I. i. iguana*

on the Lesser Antillean island of Saba.

Materials and Methods

On Saba, the presence of NNGIs was first identified in 2021 (van den Burg et al. 2025a), which

catalyzed a program to map their spread and remove as many non-native individuals as possible.

Efforts within this program were sporadic during 2022 and 2023, but became more structural with a

major NNGI removal effort during the reproductive season between December 2024 and March

2025. During these efforts we surveyed for and culled pure non-native and hybrid iguanas following

morphological assignment based on the presence (native assignment)/absence (non-native or hybrid

assignment) of the melanistic lateral spot between the eye and the tympanum (van den Burg et al.

2025a), the relative length of the head and dorsal spines and presence of enlarged nasal scales (van

den Burg et al. 2023; van den Burg et al. 2025b). Continuous genetic assessment of removed iguanas

indicates that only a minor percentage >5% was incorrectly identified (van den Burg et al. 2023,

2025b, unpublished data), and that these can correctly be identified after they become roughly two

years old. We also opportunistically captured native iguanas to collect morphological data, which

were thereafter released at their capture location. Additionally, we culled a few native females given

they were part of a harem that included a large non-native dominant male and the high probability

that mating had already taken place. Prior to December 2024, iguanas were either culled using an air

rifle or through euthanasia after noose/hand captures (lethal injection of pentobarbital by

veterinarian), whilst most iguanas were shot using either a .22 or using different types of high-caliber

ammunition during the 2024-2025 culling effort; .223, .243 and 6.5X55 with soft point ammunition

for longer distances). This change of methodology was due to efficiency since approaching iguanas

for close-proximity removal can often result in them hiding. At longer distances, and given windy

conditions on Saba, accurate bullet placement was aided using the Vortex Fury HD 5000 AB distance

measuring feature and the Fury HD Vortex application, and all shots preluded a thorough assessment

of backstop presence and orientation.

This culling effort allowed us to collect data on snout-vent length (SVL; in mm), tail length

(TL; in mm), and clutch size, except for those bodies that could not be retrieved or when these

variables were affected by bullet impact. From each captured or retrieved iguana we also sampled

blood (using non-heparinized syringe) or tissue (using single-use scalpel), which were stored in

Longmire buffer (Longmire et al. 1992) or >95% ethanol, respectively. Thereafter, we performed

necropsies on all females to determine reproductive state and count the number of developing eggs

(>1 cm diameter) to score clutch size. When SVL or the number of eggs was affected by bullet impact

we excluded the iguana from our dataset (n = 3). Lastly, prior to 2024, we collected SVL and clutch

size data opportunistically from a single native female that was hit by a car in 2021 (van den Burg et

al. 2025). Fieldwork permits were obtained through the Public Entity Saba (permits BC663.21,

BC515.2023, and BC428.2024), and we worked following recommendations from the IUCN SSC

Iguana Specialist Group (IUCN SSC Iguana Specialist Group 2017) and euthanasia guidelines from the

American Veterinary Medical Association.

We constructed two datasets to assess differences in total size (SVL + TL from animals with

complete and unbroken tails) and clutch size among different groups. We assessed differences in

total size among native iguanas from Saba (n = 25), NNGIs from St. Maarten (n = 22, van den Burg et

al. 2024, 2025b), and genetically-confirmed hybrid iguanas from Saba (n = 17; for genetics see below

and Appendix 1). Hereafter we retrieved residuals from a SVL-tail length regression, and compared

the means of these SVL-corrected tail length differences using an analysis of variance (ANOVA)

including subsequent posthoc Tukey test to assess different results between the groups. Considering

clutch size, we assessed SVL-clutch size relationships among native and non-native iguanas from

Saba, as well as previously published data from an *I. i. iguana* population in Panama (Rand 1984;

Miller 1987), an *I. i. rhinolopha* population Nicaragua (Fitch & Henderson 1977), one population of *I.*

delicatissima (Knapp et al. 2016), and the only reported *I. delicatissima* x *I. iguana* F1 hybrid

datapoint (van Wagensveld & van den Burg 2018). First, we assessed data normality and

homogeneity of variances using a Shapiro-Wilk and Levene's test, respectively. We then assessed

differences in regression slopes for the SVL – clutch size relation among these groups (excluding the

F1 hybrid record) through a linear model and pairwise comparisons of regression coefficients and

least-square means. Then, for groups with equal slopes, we used generalized linear modeling (GLM)

to assess whether differences in SVL and clutch size among these groups are explained by

genetic/population status or climatic variables. We ran GLM analyses on two datasets, both with

clutch size as the response variable. For the first analysis, we included the predictor variables SVL and

genetic status, whilst for the second analysis we also included the WorldClim variables. We used

three bioclimatic variables from the WorldClim 2 dataset (Fick & Hijmans 2017) of importance

considering variation in reptilian clutch sizes (BIO1, annual mean temperature; BIO4, temperature

seasonality; BIO15, precipitation seasonality; Meiri et al. 2020). Since clutch size is count data we

used Poisson models. For this, we used analysis of deviance (AoD) (with Chisquare-based estimate)

to assess relations between the response and predictor variables, and individual coefficient results

for the relations of between the response and categories within the predictor variables. Both models

(with and without the bioclimatic variables) were compared using the Akaike information criterion

(AIC). We used visual assessments of residuals, quantile-quantile plot and Cook's statistics to assess

model normality, assumptions and disproportionality. In case bioclimatic variables had no effect on

clutch size difference, we tested for differences in SVL-corrected clutch size among the different

genetic groups using an overall ANOVA and individual *t*-tests between all combinations; *p*-values

were corrected using the BY-FDR method (White et al. 2019).

Blood/tissue samples were used to assess species/hybrid status for each female iguana and

all iguanas with complete SVL+tail data, following van den Burg et al. (2023, 2024). After DNA

extraction using a DNeasy Kit (QIAGEN, Germany), we performed Sanger sequencing (forward and

reverse) at the University of Amsterdam where laboratory methods were implemented following

Malone et al. (2017) and van den Burg et al. (2018b) to amplify the mitochondrial NADH

dehydrogenase subunit 4 (ND4) and the nuclear MutL homolog 3 (MLH3). Using Geneious Prime

2024.0.4 (Kearse et al. 2012), we quality-checked chromatograms and compared consensus

sequences per individual to published GenBank records and unpublished sequences. Additionally, for

the female iguanas only, we generated microsatellite data for 17 loci commonly used for *Iguana*

(Valette et al. 2013; Vuillaume et al. 2015; van den Burg et al. 2021a) at the Qualyse laboratory in

France. Generated allelic data were compared to the IguanaBase database (van den Burg et al.

2021b) using discriminant analyses of principal components (DAPC) using the *adegenet* package

(Jombart 2008) in the R environment (R Core Team 2025, version 4.3.2; RStudio Team 2025, version

2023.12.1+402). Since very few genetic data are available from St. Maarten/Martin, we were not

able to employ a program to analytically predict the status of hybrid individuals (F1/F2), which we

instead tentatively did using the mtDNA and nDNA haplotypes.

Results

For clutch size comparisons, we obtained useful data from culled gravid female iguanas during Dec

2024–March 2025, indicating that both native, hybrid and non-native iguanas are reproductively

active in the same period (Table 1). The variables SVL and clutch size had normal distributions (SVL, *p*

= 0.059; eggs, *p* = 0.691) and equal variances (SVL, *p* = 0.245; eggs, *p* = 0.114).

We obtained ND4 and MLH3 sequences data from all 13 gravid females (Table 1); five

individuals only had haplotypes native to Saba (ND4 and MLH3 haplotypes correspond to HM352505

and OQ556119, respectively), seven tentative hybrid individuals had native haplotype(s) and one or

two haplotypes of Clade I or III, and one individual only had haplotypes of Clade II (mtDNA,

PV974931) and III (nDNA, OQ556117 and PV974930). Out of the seven tentative hybrids, all except

SAB91 show mtDNA/nDNA signs of F1 hybrids, with SAB91 being either a F2 hybrid or backcross.

Considering non-native haplotype origin, the two unique mtDNA haplotypes originated from Clade II

(PV974931) and III (HM352508), and the four unique nDNA haplotypes from Clade I (KX610637) and

III (OQ556117, OQ556118, PV974928, and PV974930) (Table 1).

We also generated microsatellite allelic data from all but two samples (SAB103+104; Figure

1). DAPC clustering for these microsatellite alleles showed that samples either clustered within

"*Iguana iguana* melanistic" (which were considered native given previous assessments; van den Burg

et al. 2021a, 2023), or were placed outside of the "*Iguana iguana* melanistic" cluster. These later

samples showed alleles not observed in native Saba iguanas (Breuil et al. 2020; van den Burg et al.

2021a, 2023), and were thus considered non-native (Table 1). Combined, for the 13 gravid female

iguanas, these genetic data indicated that five were native Saba iguanas (SAB45, 103, 104, 108, 114),

six were native X non-native F1 hybrids (SAB80, 97, 100, 101, 102, 117), and one iguana was a F2

hybrid (SAB91). SAB88 was a non-native iguana with mixed geographic origins and was therefore not

included in SVL-clutch size comparisons.

SVL-clutch size regression slopes differed significantly only between *I. delicatissima – I. i.*

iguana Panama (p = 0.0003) and *I. delicatissima – I. i. rhinolopha* Nicaragua (p = 0.0302) (Figure 2A).

Therefore, *I. delicatissima* (as well as the single *I. delicatissima* x *I. i. iguana* record) was not included

in the generalized linear modeling datasets. Results of both GLM analyses indicated a significant

effect of SVL on clutch size of native Saba iguanas (Table 2 and 3). For the model excluding

bioclimatic variables (AIC score of 517.66), the analysis of deviance indicated a significant effect of

SVL ($p < 2.2^{e-16}$) and Status ($p = 0.00047$) on clutch size. Specifically, we found a significant effect of

SVL on clutch size of native Saba iguanas, and a significant relation between clutch size of native Saba

iguanas and hybrid iguanas on Saba (Table 2). Post-hoc analyses of pairwise comparisons between

clutch sizes of the different groups (Saba native, Saba hybrid, Panama, and Nicaragua) showed that

clutch size of hybrid iguanas on Saba was significantly larger than native Saba iguanas (with Tukey

correction = 0.752, SE = 0.074, $p = 0.018$) and iguanas from Nicaragua (with Tukey correction = 1.282,

SE = 0.084, $p = 0.001$), and clutch size of iguanas from Panama was significantly larger than iguanas

from Nicaragua (with Tukey correction = 0.893, SE = 0.038, $p = 0.043$). Thus an average native female

of 33.75 SVL females had an average clutch size of 30 eggs whereas F1 hybrids had an average clutch

size of 42 eggs (40% higher) (Figure 2A).

For the model including bioclimatic variables, the analysis of deviance indicated a significant

effect of SVL ($p < 2.2^{e-16}$) and Status ($p = 0.00047$) on clutch size, and no effect of BIO1 ($p = 0.607$),

BIO4 ($p = 0.526$), and BIO15 ($p = 0.869$). Results of the GLM coefficients were also not significant for

the bioclimatic variables (Table 2). The model including the bioclimatic variables had an AIC score of

522.99, higher than the model excluding the bioclimatic variables. Visual assessments of residuals,

quantile-quantile plot and Cook's statistics showed no clear deviations of model normality,

assumptions and disproportionality of observations (Supplementary figures 1 and 2). Since the

model excluding the bioclimatic variables had a lower AIC, we used an ANOVA to assess the pairwise

comparisons of SVL-corrected clutch size. The ANOVA indicated significant differences among SVL-

corrected clutch size among means of the four groups (Figure 2B), $p = 0.0011$; Saba (native), Saba

(hybrid), *I. i. rhinolopha* Nicaragua, and *I. i. iguana* Panama. HY-FDR adjusted *p*-values for pairwise

comparisons were significant for all but two comparisons (Table 4): Saba (native) – *I. i. rhinolopha*

Nicaragua and Saba (hybrid) – *I. i. iguana* Panama.

Considering comparisons of total size, ANOVA results indicated significant differences of

mean SVL-corrected tail lengths among native Saba iguanas (*n* = 25), hybrid iguanas from Saba (*n* =

17), and non-native iguanas from St. Maarten (n = 22); p = 0.0312 (Figure 3). Tukey posthoc test

showed only a significant difference between native Saba and non-native St. Maarten iguanas (p =

0.0265), and non-significant differences between Saba hybrids and the other two groups (p > 0.29).

Discussion

In light of the ongoing threat of hybridization and genetic swamping by NNGI to native *Iguana*

populations in the Lesser Antilles, clutch size has remained an unstudied life-history trait. Here we

provide a first preliminary report on clutch size difference between native Saba iguanas, hybrid

iguanas from Saba, and two mainland *Iguana* populations (from different mtDNA clades, II and III;

see van den Burg et al. 2026). First, our results indicate that some mainland populations lay larger

clutches compared to native Saba iguanas. This result aligns with the findings that insular lizards

generally have smaller clutch size than to continental lizard populations (Novosolov and Meiri 2013;

Meiri et al. 2020). Importantly, although using a small dataset, we find that hybrid iguanas on Saba

appear to produce larger clutches than native Saba iguanas with a similar snout-vent length, which

may facilitate their genetic swamping of native *Iguana* populations as witnessed throughout the

Lesser Antilles.

Our NNGI removal campaign targets all sighted hybrid and nonnative iguanas on Saba, which

appear to be unequivocally identifiable after about two years of age (van den Burg et al. 2023,

2025b, unpublished data), when their detectability stabilizes to that of other adult animals. The

clutch size data were, necessarily of an opportunistic nature, since the aim of the campaign is

complete removal of all hybrid and non-native iguanas, preferably before they successfully nest. Data

opportunities were additionally constrained by the limited number of hybrid iguanas due the

invasion process having started less than 10 years ago (van den Burg et al. 2023, 2025b, 2025c,

2025d, unpublished data). This resulted in a relative small clutch size dataset, for which data could

furthermore only be collected from adult females that successfully copulated and were removed

during January-early March.

Clutch size in *Iguana* shows a high variation among sampled populations (e.g., Blankenship

1990; Nosolov et al. 2012; Knapp et al. 2016; Bock et al. 2022 and references therein). Presumably

clutch size is at least partially affected by local climate and environmental conditions similar to other

reptiles (e.g., Diaz et al. 2012; Roitberg et al. 2013), as well as island-mainland differences (Silliceo &

Díaz 2010). There are similar signs from within *Iguana* (e.g., van Marken Lichtenbelt & Albers 1993;

Knapp et al. 2016), but no study has of yet placed a narrow focus on this topic. Considering our case

study, non-native iguanas on Saba originate from St. Maarten/Martin (van den Burg et al. 2018b,

2023), where a wide-spread non-native iguana population is present that has a genetic origin

comprised of at least three of the four major mitochondrial clades within the *Iguana iguana* species

complex (Clade I, II, and III; see van den Burg et al. 2026). Equally, our generated genetic data from

hybrid and non-native individuals on Saba indicates similar genetic origins (Table 1 and Appendix 1).

Currently there are no data on clutch sizes from non-native iguanas on St. Maarten/Martin, which

would be interesting to better interpret the clutch size differences in our dataset. Considering clutch

sizes of the three major mtDNA clades, although there likely is ample variation within each clade, our

dataset includes data from Clades II and III (Fitch & Henderson 1977; Rand 1984; Miller 1987), but

not from Clade I (van Marken Lichtenbelt & Albers 1993 did not provide the raw data). However, van

Marken Lichtenbelt & Albers (1993) compared their SVL-clutch size data to those of Rand (1984),

finding that clutch size for equal SVLs was significantly lower for Clade I compared to Clade II (see

table 2 in van Marken Lichtenbelt & Albers 1993). Their assessment combined with our comparison

of SVL-clutch size residuals (Table 4) suggests hybrid iguanas on Saba lay more eggs compared to

native Saba iguanas as well as iguanas from Nicaragua and Curaçao, but not Panama despite a

potential trend (p = 0.073). Clutch size data combined with genomic approaches could shed light on

genes involved in fecundity and selective pressure in (hybrid) iguanas, as well as potential novel gene

complexes and local adaptation in (the mostly mixed-origin) populations of NNGIs across the Lesser

Antilles (Martin et al. 2015; Vuillaume et al. 2015; van den Burg et al. 2018b, 2023, 2024; Pounder et

al. 2020).

For hybridization to occur, mating periods of native and non-native iguana should at least

partially overlap. Equally variable compared to clutch size is the timing of reproduction within both *I.*

delicatissima and *I. iguana* that follows local annual climate patterns (van den Burg et al. 2018a;

Breuil et al. 2019). On Saba, we found that reproductive timing was similar between non-

native/hybrid and native Saba iguanas, which is in contrast to some Lesser Antillean islands native to

I. delicatissima (Breuil et al. 2019). However, since non-native iguanas from different geographic

origins are present within the region and also hybridize (see Table 1; e.g., Vuillaume et al. 2015; van

den Burg et al. 2023), future assessments of reproductive timing and geographic origin of non-native

populations should be considered (e.g., on St. Martin), as well as potential changes in reproductive

timing when NNGI arrive (plasticity) and when hybridization results in hybrid swarms. Understanding

these population characteristics could inform tailored conservation strategies to bolster native

populations against the threats posed by non-native iguanas, especially when targeting gravid

females (with potential hybrid clutches) at nesting sites.

In conclusion, our results suggest that the observed decline in native *Iguana* populations is at

least partially driven by genetic swamping, not only by a larger clutch size of pure NNGIs, but also

through larger clutch size of hybrid iguanas compared to native iguanas (i.e., hybrid vigor). Because

native iguanas can live to 24 years in the wild (Warret Rodrigues et al. 2021), and thus can remain

present long enough after incursions and establishment of NNGIs for (genetic) identification to make

it possible to safeguard their genetic lineage for conservation purposes. We emphasize that despite

the low variation in mtDNA across the range of *I. delicatissima* (Martin et al. 2015), nuclear markers

do indicate genetic structure among the different island populations (van den Burg et al. 2021). Since

the maintenance of intraspecific genetic diversity is critical to long-term conservation of a species

(e.g., Lacy 1997), identifying and capturing remaining native iguanas in hybridized and introgressed

populations is critical for native *Iguana* conservation. We therefore urge regional partners to identify

remaining native *Iguana* survivors within areas that have been heavily impacted by hybridization and

introgression with non-native green iguanas (e.g., Grenada, Guadeloupe, and Martinique; Angin

2017; Breuil et al. 2019) so that their protection can be improved.

Acknowledgments

We thank Peter Kuperus for performing all laboratory procedures and analyses, and the Public Entity

Saba and Saba Conservation Foundation for on-island logistic support. We thank Charles Knapp for

providing feedback and recommendations that greatly improved an earlier version of this

manuscript. We also remain indebted to four anonymous reviewers for their critical feedback that

allowed essential improvements to our manuscript. We are grateful for the financial support that

allowed us to perform this study from the Dutch Iguana Foundation, the Iguanaland Conservation

Fund, the Public Entity Saba within their invasive species project, and the Ministry of Agriculture,

Fisheries, Food Security and Nature (LVVN) through the Wageningen University BO research program

(BO-43-117-006) under Wageningen University and Research project number 4318100346-1.

Fieldwork on Saba was approved under permits BC663.21, BC515.2023 and BC428.2024.

References

Allendorf, F.W., Leary, R.F., Spruell, P. & Wenburg, J.K. (2001). The problems with hybrids: setting
conservation guidelines. *Trends in Ecology & Evolution* **16**, 613–622.

Angin, B. (2017). Plan National d'Actions Pour le Rétablissement de L'iguane des Petites Antilles,
Iguana delicatissima, 2018–2022; Ministère de la Transition Écologique et Solidaire: Schoelcher,
Martinique, 2017; p. 69 + annexes.

Blankenship, J.R. (1990). The Wildlife of Montserrat including an annotated bird list for the island.
Montserrat National Trust, Montserrat, West Indies.

Bock, B., Malone, C.L., Knapp, C., Aparicio, J., Avila-Pires, T.C.S., Cacciali, P., Caicedo, J.R., Chaves, G.,
Cisneros-Heredia, D.F., Gutiérrez-Cárdenas, P., Lamar, W., Moravec, J., Perez, P., Porras, L.W., Rivas, G.,
Scott, N., Solórzano, A. & Sunyer, J. (2022). *Iguana iguana* (amended version of 2020
assessment). *The IUCN Red List of Threatened Species* **2022**: e.T174481A218317281. Accessed on 09
January 2026.

Breuil, M., Vuillaume, B., Schikorski, D., Krauss, U., Morton, M.N., Haynes, P., Daltry, J.C., Corry, E.,
Gaymes, G., Gaymes, J., Bech, N., Jelic, M. & Grandjean, F. (2019). A story of nasal horns: two new

subspecies of *Iguana* Laurenti, 1768 (Squamata, Iguanidae) in Saint Lucia, St Vincent & the
Grenadines, and Grenada (southern Lesser Antilles). *Zootaxa* **4608**, 201–232.

Breuil, M., Schikorski, D., Vuillaume, B., Krauss, U., Morton, M.N., Corry, E., Bech, N., Jelić, M. &
Grandjean, F. (2020) Painted black: *Iguana melanoderma* (Reptilia, Squamata, Iguanidae) a new
melanistic endemic species from Saba and Montserrat islands (Lesser Antilles). *ZooKeys* **926**, 95–131.

Breuil, M., Schikorski, D., Vuillaume, B., Krauss, U., Daltry, J.C., Gaymes, G., Gaymes, J., Lepais, O.,
Bech, N., Jelic, M. & Becking, T. (2022). *Iguana insularis* (Iguanidae) from the southern Lesser Antilles:
an endemic lineage endangered by hybridization. *ZooKeys* **1086**, 137–161.

Censky, E.J., Hodge, K. & Dudley, J. (1998). Over-water dispersal of lizards due to hurricanes. *Nature*
395, 556.

Day, M.L. & Thorpe, R.S. (1996). Population Differentiation of *Iguana delicatissima* and *Iguana iguana*
in the Lesser Antilles. In Contributions to West Indian Herpetology: A Tribute to Albert Schwartz;
Powell, R., Henderson, R.W., Eds.; Society for the Study of Amphibians and Reptiles: Plymouth, MN,
USA; pp. 136–137.

Díaz, J.A., Iraeta, P., Verdú-Ricoy, J., Siliceo, I. & Salvador, A. (2012). Intraspecific variation of
reproductive traits in a Mediterranean lizard: clutch, population, and lineage effects. *Evolutionary*
Biology **39**, 106–115.

Fick, S.E. & Hijmans, R.J. (2017). WorldClim 2: new 1km spatial resolution climate surfaces for global
land areas. *International Journal of Climatology* **37**(12), 4302–4315.

Fitch, H.S. & Henderson, R.W. (1977). Age and sex differences, reproduction, and conservation of
Iguana iguana (No. 13). *Contributions in Biology and Geology, Milwaukee Public Museum* **13**, 1–21.

Godfray, H.C.J., Partridge, L. & Harvey, P.H. (1991). Clutch size. *Annual Review of Ecology and*
Systematics **22**, 409–429.

Hata, H., Uemura, Y., Ouchi, K. and Matsuba, H. (2019). Hybridization between an endangered
freshwater fish and an introduced congeneric species and consequent genetic introgression. *PLoS*
ONE **14**(2), e0212452.

Iguana Taxonomy Working Group; Buckley, L.J., de Queiroz, K., Grant, T.D., Hollingsworth, B.D.,
Iverson, J.B., Pasachnik, S.A. & Stephen, C.L. (2016). A checklist of the iguanas of the world
(Iguanidae; Iguaninae). In: Iverson, J.B., Grant, T.D., Knapp, C.R. & Pasachnik, S.A. (eds), Iguanas:
Biology, Systematics, and Conservation. *Herpetological Conservation and Biology* **11**, 4–46. University
of California Press, US.

Iguana Taxonomy Working Group; Buckley, L.J., de Queiroz, K., Grant, T.D., Hollingsworth, B.D.,
Malone, C.L., Pasachnik, S.A., Reynolds, R.G. & Zarza, E. (2022). A checklist of the iguanas of the
world (Iguanidae; Iguaninae) 2022 supplement to: 2016. *Herpetol. Conserv. Biol.* **11**, 4– 46 and 2019
Supplement. Available from: https://www.iucnisg.org/wp-
content/uploads/2022/05/ITWG_Checklist_2022_Supplement.pdf.

IUCN SSC Iguana Specialist Group (2017) Position Statement of the IUCN SSC Iguana Specialist Group
on Non-Native Invasive Iguanas. Version 2.1. Available from: http://www.iucn-isg.org/wp-
content/uploads/2017/12/IUCN_SSC_ISG_Invasive_Iguana_Position_Statement_Dec2017.pdf

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics **24**(11), 1403–1405.

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A.,
Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P. & Drummond, A. (2012). Geneious basic:
An integrated and extendable desktop software platform for the organization and analysis of
sequence data. *Bioinformatics* **28**, 1647–1649.

Knapp, C.R., Breuil, M., Rodrigues, C. & Iverson, J.B. (2014). Lesser Antillean Iguana, *Iguana*
delicatissima: conservation Action Plan, 2014—2016. Gland (Switzerland): IUCN SSC Iguana Specialist
Group.

Knapp, C.R., Prince, L. & James, A. (2016). Movements and nesting of the Lesser Antillean Iguana
(*Iguana delicatissima*) from Dominica, West Indies: implications for conservation. In Iguanas: biology,
systematics, and conservation. Iverson, J.B., Grant, T.D., Knapp, C.R. & Pasachnik, S.A. (eds.). Open
Access Publishing, *Herpetological Conservation and Biology* **11**(Monograph 6): 154–167.

Knapp, C.R., Grant, T.D., Pasachnik, S.A., Angin, B., Boman, E., Brisbane, J., Buckner, S.D., Haakonsson,
J.E., Harlow, P.S., Mukhida, F., Thomas-Moko, N., van den Burg, M.P. & Wasilewski, J.A. (2021). The
global need to address threats from invasive alien iguanas. *Animal Conservation* **24**, 717–719.

Lacy, R.C. (1997). Importance of genetic variation to the viability of mammalian populations. *Journal*
of Mammalogy **78**, 329–335.

Longmire, J.L., Gee, G.F., Hardekopf, C.L. & Mark, G.A. (1992). Establishing paternity in whooping
cranes (*Grus americana*) by DNA analysis. *Auk* **109**, 522–529.

Malone, C.L., Reynoso, V.H. & Buckley, L. (2017). Never judge an iguana by its spines: systematics of
the Yucatan spiny tailed iguana, *Ctenosaura defensor* (Cope, 1866). *Molecular Phylogenetics and*
Evolution **115**, 27–39.

Martin, J.L., Knapp, C.R., Gerber, G.P., Thorpe, R.S. & Welch, M.E. (2015). Phylogeography of the
endangered Lesser Antillean iguana, *Iguana delicatissima*: a recent diaspora in an archipelago known
for ancient herpetological endemism. *Journal of Heredity* **106**(3), 315–321.

Meiri, S., Avila, L., Bauer, A.M., Chapple, D.G., Das, I., Doan, T.M., Doughty, P., Ellis, R., Grismer, L.,
Kraus, F., Morando, M., Oliver, P., Pincheira-Donoso, D., Ribeiro-Junior, M.A., Shea, G., Torres-
Carvajal, O, Slavenko, A. & Roll, U. (2020). The global diversity and distribution of lizard clutch
sizes. *Global Ecology and Biogeography* **29**(9), 1515–1530.

Miller, T.J. (1987). Artificial incubation of eggs of the green iguana (*Iguana iguana*). *Zoo biology* **6**(3),
225–236.

Novosolov, M., Raia, P. & Meiri, S. (2013). The island syndrome in lizards. *Global Ecology and*
Biogeography 22(2), 184–191.

Pounder, K.C., Mukhida, F., Brown, R.P., Carter, D., Daltry, J.C., Fleming, T., Goetz, M., Halsey, L.G.,
Hughes, G., Questel, K., Saccheri, I.J., Williams, R. & Soanes, L.M. (2020). Testing for hybridisation of
the critically endangered *Iguana delicatissima* on Anguilla to inform conservation efforts.
Conservation Genetics **21**, 405–420.

R Core Team (2025). R: a language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing. https://www.R-project.org/.

Rand, A.S. (1984). Clutch size in *Iguana iguana* in central Panama. In: Vertebrate Ecology and
Systematics - A Tribute to Henry S. Fitch, p. 115–122. Seigel, R.A., Hunt, L.E., Knight, J.L., Malaret, L.,
Zuschlag, N.L., Eds., University Kansas Mus. Nat. Hist. Spec. Publ. No. 10.

Rhymer, J.M. & Simberloff, D. (1996). Extinction by hybridization and introgression. *Annual Review of*
Ecology, Evolution, and Systematics **27**, 83–109.

Rohwer, F.C. (1988). Inter and intraspecific relationships between egg size and clutch size in
waterfowl. *The Auk* **105**(1), 161–176.

Roitberg, E.S., Kuranova, V.N., Bulakhova, N.A., Orlova, V.F., Eplanova, G.V., Zinenko, O.I.,
Shamgunova, R.R., Hofmann, S. & Yakovlev, V.A. (2013). Variation of reproductive traits and female
body size in the most widely-ranging terrestrial reptile: testing the effects of reproductive mode,
lineage, and climate. *Evolutionary Biology* **40**, 420–438.

RStudio Team (2015). RStudio: integrated development for R. Boston, MA: RStudio, Inc.
http://www.rstudio.com/.

Siliceo, I. & Díaz, J.A. (2010). A comparative study of clutch size, range size, and the conservation
status of island vs. mainland lacertid lizards. *Biological Conservation* **143**(11), 2601–2608.

Strubbe, D. & Matthysen, E. (2009). Experimental evidence for nest-site competition between
invasive ring-necked parakeets (*Psittacula krameri*) and native nuthatches (*Sitta europaea*). *Biological*
Conservation **142**(8), 1588–1594.

Todesco, M., Pascual, M.A., Owens, G.L., Ostevik, K.L., Moyers, B.T., Hübner, S., Heredia, S.M., Hahn,
M.A., Caseys, C., Bock, D.G. & Rieseberg, L.H. (2016). Hybridization and extinction. *Evolutionary*
Applications **9**, 892–908.

Vanhaecke, D., Garcia de Leaniz, C., Gajardo, G., Young, K., Sanzana, J., Orellana, G., Fowler, D.,
Howes, P., Monzon-Arguello, C. & Consuegra, S. (2012). DNA barcoding and microsatellites help
species delimitation and hybrid identification in endangered galaxiid fishes. *PLoS ONE*, **7**(3), e32939.

Valette, V., Filipova, L., Vuillaume, B., Cherbonnel, C., Risterucci, A.M., Delaunay, C., Breuil, M. &
Grandjean, F. (2013). Isolation and characterization of microsatellite loci from *Iguana delicatissima*
(Reptilia: Iguanidae), new perspectives for investigation of hybridization events with *Iguana iguana*.
Conservation Genetics Resources **5**, 173–175.

van den Burg, M., Breuil, M. & Knapp, C. (2018a). Lesser Antillean iguana: *Iguana delicatissima*. *The*
IUCN Red List of Threatened Species **2018**: eT10800A122936983. Available from:
https://doi.org/10.2305/IUCN.UK.2018-1.RLTS. T10800A122936983.en. Accessed on 17 April 2025.

van den Burg, M.P., Meirmans, P.G., van Wagensveld, T. P., Kluskens, B., Madden, H., Welch, M.E. &
Breeuwer, J.A.J. (2018b). The Lesser Antillean Iguana (*Iguana delicatissima*) on St. Eustatius:
genetically depauperate and threatened by ongoing hybridization. *Journal of Heredity* **109**, 426–437.

van den Burg, M.P., Madden, H., van Wagensveld, T.P. & Buma, C. (2018c). Anthropogenic mortality in
the critically endangered lesser antillean iguana (*Iguana delicatissima*) on St. Eustatius. *Reptiles &*
Amphibians **25**(2), 120–124.

van den Burg, M.P., Brisbane, J.L.K. & Knapp, C.R. (2020). Post-hurricane relief facilitates invasion and
establishment of two invasive alien vertebrate species in the Commonwealth of Dominica, West
Indies. *Biological Invasions* **22**, 195–203.

van den Burg, M.P., Grandjean, F., Schikorski, D., Breuil, M. & Malone, C.L. (2021a). A genus-wide
analysis of genetic variation to guide population management, hybrid identification, and monitoring
of invasions and illegal trade in *Iguana* (Reptilia: Iguanidae). *Conservation Genetic Resources* **13**, 435–
445.

van den Burg, M.P., Grandjean, F., Schikorski, D., Breuil, M. & Malone, C.L. (2021b). IguanaBase: a live
microsatellite database to guide research and conservation in the genus *Iguana* (Reptilia: Iguanidae).
figshare. Available at: https://doi.org/10.6084/m9.figshare.13584923.v4.

van den Burg, M.P., Goetz, M., Brannon, L., Weekes, T.S., Ryan, K.V. & Debrot, A.O. (2023). An
integrative approach to assess non-native iguana presence on Saba and Montserrat: are we losing all
native *Iguana* populations in the Lesser Antilles? *Animal Conservation* **26**, 813–825.

van den Burg, M. P., Kappelhof, J., Mitchell, A. & Debrot, A.O. (2024). Exploring the potential of
morphometrics to inform the conservation of *Iguana delicatissima*. *Conservation* **4**, 617–626.

van den Burg, M.P., Madden, H. & Debrot, A.O. (2025a). Population estimate and conservation of the
melanistic *Iguana iguana* population on Saba, Caribbean Netherlands. *Herpetological Journal* **35**,
176–186.

van den Burg, M.P., Kappelhof, J. & Debrot, A.O. (2025b). Morphometrics aid identification of non-
native iguanas to improve conservation of native Saba Green Iguanas*, Iguana iguana* (Linnaeus,
1758). *Herpetology Notes* **18**, 1031–1037.

van den Burg, M.P., Madden, H. & Debrot, A.O. (2025c). Conservation State of the Saba Green Iguana
in Saba. In: State of nature report for the Caribbean Netherlands 2024. Debrot, A.O., Henkens,
R.J.H.G., Verweij, P.J.F.M., van den Burg, M.P. & Meesters, E.H., pp. 205–212. Wageningen Marine
Research report C001/25.

van den Burg, M.P., Paredes-León, R., Angin, B., Brannon, L., López-Delgado, J., Cornwell, T.O., Jones,
L., Kuperus, P. & Debrot, A.O. (2025d). Exploring the co-evolutionary host-parasite complex as a novel
tool for monitoring iguanid invasions. *Biological Invasions* **27**(8), 173.

van den Burg, M.P., de Queiroz, K., Ventayol, M., Debrot, A.O. & Malone, C.L. (2026). Consideration of
range-wide variation is critical when splitting widely distributed species: the case of the proposed
Iguana melanoderma. *Zootaxa* **5748**(3), 387–399.

van Kuijk, J.L., van den Burg, M.P., Didaskalou, E.A., de Boer, M., Debrot, A.O., Wielstra, B. & Stewart,
K.A. (2025). Safeguarding *Iguana* diversity: enabling rapid and low-effort tracking of non-native
iguanas through terrestrial eDNA innovations. *Biological Invasions* **27**(1), 63.

van Marken Lichtenbelt, W.D. & Albers, K.B. (1993). Reproductive adaptations of the green iguana on
a semiarid island. *Copeia* **1993**, 790–798.

Vuillaume, B., Valette, V., Lepais, O., Grandjean, F. & Breuil, M. (2015). Genetic evidence of
hybridization between the endangered native species *Iguana delicatissima* and the invasive *Iguana*
iguana (Reptilia: Iguanidae) in the Lesser Antilles: Management implications. *PLoS ONE* **10**, 0127575.

Warret Rodrigues, C., Angin, B. & Besnard, A. (2021). Favoring recruitment as a conservation strategy
to improve the resilience of long-lived reptile populations: Insights from a population viability
analysis. *Ecology and Evolution* **11**(19), 13068–13080.

White, T., van der Ende, J. & Nichols, T.E. (2019). Beyond Bonferroni revisited: Concerns over inflated
false positive research findings in the fields of conservation genetics, biology, and medicine.
Conservation Genetics **20**, 927–937.

Tables

Table 1. Collected data on snout-vent length (SVL) and egg count from female iguanas on Saba, including status assignment that was based on the combined morphological and genetic assignment for mitochondrial (mtDNA), nuclear (nDNA), and microsatellite (msat) data. Unless native, clade numbers in the assignment columns refer to mtDNA clades within the *Iguana iguana* species complex following van den Burg et al. (2026). Asterisk: SAB88 was not included in further analyses since it lacked native haplotypes. Superscripted B: tails that were not complete.

Individual	SVL (mm)	Tail (mm)	Eggs	Assignment	Morphology	mtDNA	nDNA	msat
SAB45	325	NA	29	native	native	native	native / native	native
SAB103	305	450 [B]	21	native	native	native	native / native	NA
SAB104	345	500 [B]	29	native	native	native	native / native	NA
SAB108	354	701 [B]	34	native	native	native	native / native	native
SAB114	381	955	44	native	native	native	native / native	native
SAB97	357	811	42	F1 hybrid	hybrid	Clade III: HM352508	native / Clade III: OQ556118	nonnative
SAB100	369	921	50	F1 hybrid	hybrid	native	native / Clade I: KX610637	nonnative
SAB102	340	880	45	F1 hybrid	hybrid	native	native / Clade III: PV974928	nonnative
SAB117	345	843	42	F1 hybrid	hybrid	native	native / Clade III: OQ556117	nonnative
SAB80	361	875	40	F1 hybrid	nonnative	native	native / Clade III: PV974928	nonnative
SAB101	379	855 [B]	58	F1 hybrid	nonnative	Clade III: HM352508	native / Clade III: OQ556117	nonnative
SAB91	311	802	37	F2 hybrid	hybrid	Clade III: HM352508	native / native	nonnative
SAB88*	371	901	48	nonnative	nonnative	Clade II: PV974931	Clade III: OQ556117 / Clade III: PV974930	nonnative

Table 2. Results of the generalized linear mixed model (GLM) for factors affecting clutch size of native Saba iguanas, dataset excluding bioclimatic variables

	Estimate	SE	z	P
Intercept	0.622	0.196	3.167	0.002
SVL	0.008	0.001	15.856	< 2 e-16
Saba hybrids	0.286	0.098	2.920	0.004
Nicaragua	0.037	0.086	0.432	0.666
Panama	0.150	0.084	1.779	0.075

Table 3. Results of the generalized linear mixed model (GLM) for factors affecting clutch size of native Saba iguanas, dataset including bioclimatic variables; BIO1, annual mean temperature; BIO4, temperature seasonality; BIO15, precipitation seasonality (Fick & Hijmans 2017).

	Estimate	SE	z	P
Intercept	-0.762	8.042	-0.095	0.925
SVL	0.008	0.001	15.698	< 2 e-16
Saba hybrids	0.316	0.108	2.914	0.004
Nicaragua	-3.365	21.510	-0.156	0.876
Panama	-1.305	11.982	-0.109	0.913
BIO1	-0.057	0.648	-0.087	0.930
BIO4	0.008	0.030	0.264	0.792
BIO15	0.055	0.333	0.165	0.869

567 **Table 4.** Results of pairwise *t*-test for differences in SVL-corrected residuals in clutch size for female

568 iguanas across four groups; Saba (native), Saba (nonnative), *I. i. rhinolopha* Nicaragua, and *I. i. iguana*

569 Panama. *P*-values are HY-FDR adjusted. Non-significant results are indicated by n.s. These results

570 exclude *I. delicatissima* and the only reported *I. delicatissima* x *I. iguana* hybrid (van Wagensveld &

571 van den Burg 2018) as also in Figure 2A. SAB88 was also not included.

	Saba (native)	Saba (hybrid)	*I. i. rhinolopha* Nicaragua
Saba (native)			
Saba (hybrid)	0.0017		
I. i. rhinolopha Nicaragua	0.073 (n.s.)	0.0065	
I. i. iguana Panama	0.0017	0.073 (n.s.)	0.016

Figure legends

596 **Figure 1.** Discriminant analysis of a principal components scatterplot representing 17 microsatellite

597 loci from reference IguanaBase individuals (colored circles) and 11 gravid females from Saba; three

598 native iguanas (circles), and eight hybrid (triangles) and nonnative (star) iguanas. Clade codes follow

599 van den Burg et al. (2026).

Figure 2. Relationships between snout-vent length (SVL) and clutch size among native and hybrid iguanas on Saba and different populations of *Iguana* spp., including data from Fitch & Henderson (1977), Rand (1984), Miller (1987), Knapp et al. (2016), and van Wagensveld & van den Burg (2018). **A)** SVL-clutch size regression including all groups. **B)** Boxplots of SVL-clutch size residuals among groups with regression slopes that were not significantly different. SAB88 is not included.

Figure 3. Boxplots showing differences between SVL-corrected tail length (overall size) between native Saba iguanas, hybrid iguanas from Saba, and non-native iguanas on St. Maarten.

Appendix

Appendix 1. Collected data on snout-vent length (SVL) and tail length from iguanas on Saba, including genetic assignment for mitochondrial (mtDNA; one haplotype), nuclear (nDNA; two haplotypes), and microsatellite (msat) data. Unless native, clade numbers in the assignment columns refer to mtDNA clades within the *Iguana iguana* species complex following van den Burg et al. (2026). Asterisk indicate data published in van den Burg et al. (2023). Native ND4 and MLH3 haplotypes correspond to HM352505 and OQ556119, respectively. All individuals had complete tails.

Individual	Fate	SVL (mm)	Tail (mm)	mtDNA	nDNA
SAB01 (M)*	Alive	230	595	native	native/native
SAB03 (M)*	Alive	405	1022	native	native/native
SAB11 (F)*	Alive	340	821	native	native/native
SAB15 (M)*	Alive	389	959	native	native/native
SAB17 (F)*	Alive	350	828	native	native/native
SAB18 (F)*	Alive	326	795	native	native/native
SAB22 (M)*	Alive	310	725	native	native/native
SAB23 (F)*	Alive	267	697	native	native/native
SAB24 (M)*	Alive	265	708	native	native/native
SAB28 (M)*	Alive	401	965	native	native/native
SAB30 (F)*	Alive	316	776	native	native/native
SAB31 (F)*	Alive	300	774	native	native/native
SAB41 (F)*	Alive	362	846	native	native/native
SAB44 (M)*	Alive	305	757	native	native/native
SAB51 (F)*	Alive	363	810	native	native/native
SAB60 (F)	Alive	361	838	native	native/native
SAB61 (M)	Alive	391	988	native	native/native
SAB70 (F)	Alive	357	902	native	native/native
SAB75 (M)	Alive	311	830	native	native/native
SAB76 (M)	Alive	371	957	native	native/native
SAB82 (M)	Dead	340	896	native	native/native
SAB83 (M)	Alive	383	907	native	native/native
SAB84 (M)	Alive	327	844	native	native/native
SAB112 (M)	Dead	271	675	native	native/native
SAB114 (F)	Dead	381	955	native	native/native
SAB08 (M)*	Dead	345	888	Clade III: HM352508	native/Clade III: OQ556118
SAB57 (F)*	Dead	359	838	Clade III: HM352508	native/Clade III: OQ556117
SAB63 (M)	Dead	410	1104	Clade I: OQ556120	native/ Clade I: KX610637_a
SAB69 (M)	Dead	267	697	Clade I: OQ556120	native/ Clade I: KX610637_a
SAB80 (F)	Dead	361	875	native	native/Clade III: PV974928
SAB85 (M)	Dead	440	1073	Clade I: HM352499	native/Clade II: PV974929
SAB87 (M)	Dead	400	991	Clade III: HM352508	native/Clade III: OQ556117
SAB90 (M)	Dead	312	835	Clade III: HM352508	native/Clade III: OQ556118
SAB91 (F)	Dead	311	802	Clade III: HM352508	native/native
SAB94 (M)	Dead	315	855	native	native/Clade III: OQ556117
SAB97 (F)	Dead	357	811	Clade III: HM352508	native/Clade III: OQ556118
SAB100 (F)	Dead	369	921	native	native/Clade I: OQ556119
SAB102 (F)	Dead	340	880	native	native/Clade III: PV974928
SAB105 (F)	Dead	255	659	Clade III: HM352508	native/native
SAB115 (M)	Dead	189	503	Clade III: HM352508	native/native
SAB116 (F)	Dead	259	633	Clade III: HM352508	native/Clade III: OQ556117
SAB117 (F)	Dead	345	843	native	native/Clade III: OQ556117

629