Protein trafficking and synaptic demand configure complex and dynamic synaptome architectures of individual neurons

Oksana Sorokina[1], Edita Bulovaite[2,3], Anatoly Sorokin[4], Seth G.N. Grant[2,3,5,6,7]& J. Douglas Armstrong[1,8].

[1]Institute for Adaptive and Neural Computation, School of Informatics, University of Edinburgh

[2]Genes to Cognition Programme, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh EH16 4SB, UK

[3]Institute of Neuroscience and Cardiovascular Research, University of Edinburgh, Edinburgh EH16 4SB, UK

[4]Okinawa Institute of Science and Technology, Japan

[5]Universitat Autònoma de Barcelona, Bellaterra (Cerdanyola del Vallès), Spain.

[6]Euan MacDonald Centre, University of Edinburgh, Edinburgh EH16 4SB, UK

[7]Simons Initiative for the Developing Brain (SIDB), Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh EH8 9XD, UK

[8]Computational Biomedicine Institute (IAS-5 / INM-9), Forschungszentrum Jülich, Jülich, Germany.

Abstract

Excitatory synapses are the most abundant synapse type in the brain and are essential for behaviour and implicated in hundreds of brain disorders. These synapses exhibit striking structural and functional diversity, arising from differences in their proteomic composition and turnover. Synaptome mapping at single-synapse resolution reveals that this diversity is spatially organised along the dendritic tree of individual neurons. However, the cell biological mechanisms underlying the generation of these spatial synaptic patterns and their variation across neuron types and throughout the lifespan remain poorly understood. To investigate the contributions of somatic and dendritic protein synthesis, protein trafficking, and local regulatory mechanisms such as activity-dependent degradation, we developed computational models simulating these processes and compared their predictions with empirical synaptome data. We found that an extended sushi-belt model for spatial trafficking and local synaptic demand for proteins was sufficient to explain the complex profiles of synaptic protein distributions observed in young, mature and old mice and in different cell types. Our findings suggest the highly complex and dynamic synaptome architecture of the brain is an emergent property of a minimal set of cell biological processes. Our model sets the stage for simulations of brain tissue incorporating molecularly diverse neuronal and synaptic types in a synaptome and connectome architecture.

Key words

Synapse, synaptome, dendrite, neuron, simulation.

Introduction

The hallmark anatomical feature of neurons is their highly elaborate tree-like branches that maximise the number of synaptic contacts[1]. A single neuron may receive direct input on its dendritic tree from thousands of other neurons. Every synaptic contact contains up to hundreds of thousands of individual proteins encoded by several thousand genes[2]. These proteins assemble into multiprotein complexes[3] that are molecular machines mediating release of neurotransmitters from the presynaptic terminal and signal integration in the postsynaptic terminal. The

crucial role of these proteins is exemplified by the hundreds of gene mutations that result in brain and behavioural disorders[4,5].

The majority of synapses in the mammalian brain employ glutamate as a neurotransmitter and are excitatory in nature. However, underneath the common transmitter phenotype there is a hidden degree of complexity at the individual synapse level with vastly diverse protein compositions. The proteomes of individual synapses differ depending on their location on the dendritic tree, neuron type, age, neuronal activity, genetics and other factors [6-8]. Additionally, synapse proteins have to be replaced continuously and the time course of their turnover differs between synapses, further contributing to the molecular diversity of synapses [9,10].

The spatial distribution of molecularly diverse synapses on the dendritic tree is non-random. For example, synaptome mapping reveals the synaptic scaffolding proteins PSD95 and SAP102[6] are differentially distributed on the dendrites of CA1 pyramidal neurons (PyNs) and dentate gyrus granule neurons (DGs) in the intact mouse brain. Single-synapse resolution mapping of PSD95 turnover also indicates the presence of complex spatial gradients across the dendritic tree [9]. However, these gradients are not simple continuums from soma to distal dendrite but appear to be a combination of such continuums together with local discontinuities that correlate with the presence of pathway-specific axonal inputs [9]. This suggests the interplay of two mechanisms: a cell-autonomous mechanism that establishes a distance-dependent gradient and a non-cell autonomous mechanism that operates in local regions of the dendrite [6,7,9].

To explore the potential mechanisms underlying complex gradients of diverse synapses in neurons, we adapted existing computational models of dendritic molecule distributions to the synaptome mapping data that reveals these gradients in mice of different ages. An extensive literature[11] invokes many mechanisms including somatic and dendritic protein synthesis, active and passive trafficking of proteins to their destination and local protein degradation. While simulations incorporating diffusion, active transport and local translation have been explored, none have been applied to the complex molecular and synaptic gradients observed in the intact brain.

We used a model that incorporates Doyle and Kiebler's 'sushi-belt' theory of molecular transport[12,13], which assumes that molecules synthesised in the soma are continuously trafficked throughout the entire dendritic tree, imitating a conveyor belt. Local demand for proteins, which mimics the activity/plasticity state of the local synapse population, generates hot spots of "hungry" synapses that sequester and anchor these molecules. This model can be conveniently simulated with the neuron geometry defined in a format compatible with NEURON, a biophysical multi-compartment model[13,14], accommodating the intricate branching of the neuron type in question. We show that the complex gradients of synapse composition and dynamics can be simulated in two major classes of excitatory neurons – pyramidal and granule cell neurons – using a model assuming only somatic protein synthesis and a sushi belt that distributes proteins to synapses with different demand and degradation rates. Our simulation recapitulates the synaptome architecture observed in mice of different ages across the lifespan, implicating the regulation of specific cellular processes that change with age. The arising picture is that the spatial organisation of molecular diverse synapses could be an emergent property of a few basic cell biological processes.

Results

Establishing the spatiotemporal distribution of PSD95 in hippocampal pyramidal neurons.

Bulovaite et al, 2022[9] measured the quantity of fluorescently pulse-labelled synaptic PSD95 puncta in 20 individual subregions of CA1 pyramidal neurons in the mouse brain and tracked the labelled protein over seven days. These data revealed complex distributions of synapse densities in neurons. Further, they showed a complex dynamic process where intensities at different regions of the neuron changed over several days post-labelling (a term labelled Synapse Protein Lifetime (SPL); Figure 1, A shows distribution at day 0). In apical dendrites, SPL appeared to increase with distance from the soma. However, the curve was not linear and included a dip at the intersection between stratum radiatum and stratum lacunosum of the CA1. A gradient of increasing SPL with distance from soma was also observed in the more proximal parts of stratum oriens before levelling and reversing in the

most distal regions. These complex distributions of SPL paralleled the distribution of synaptic puncta density across the CA1 dendritic trees.

Assuming the total concentration of (labelled and unlabelled) PSD95 to be more or less constant in any given puncta, the SPL of labelled synaptic PSD95 across the subcellular regions may be a compound of at least two processes: First, there is the actual protein turn-over, or molecular decay rate, where labelled protein is degraded and subsequently replaced by new unlabelled protein, which acts to reduce SPL. Second, there will be a rate of replenishment of degraded protein with pre-existing labelled PSD95 from the finite pool existing elsewhere, *e.g.* from molecular transport mechanisms such as a sushi-belt conveyor, which is expected to increase SPL. One or both of these factors needs to vary along the neuron to account for the spatial differences in observed decay rates in labelled protein. Here we re-used the experimental data of PSD95 to interrogate computational models that describe possible mechanisms for PSD95 distribution in CA1 neurons.

Extending the sushi-belt model for protein distribution simulations

Williams et al's (2016)[13] model (Figure 1, B) was designed to predict molecular distribution given a set of parameters, based on an assumption that the molecule pool moves from soma to the points of high local demand. Here, we face the reverse challenge; can we calculate the model parameters that best fit the observed molecular distributions? Therefore, we implemented a new model, using the same underlying mathematical principles, but enabling simulations that can be used to explore the parameter space and allow us to compare predicted profiles against observed data (further details in Methods).

In common with Williams et al (2016), our PSD95 cargo is bound to motor proteins, which can move the cargo in both directions along the dendritic branches. The model splits these branches into virtual compartments and considers cargo movement between adjoining compartments. The reconstructed 3D pyramidal neuron model (from Migliore and Migliore, 2012)[15] is divided into 742 compartments of varying length from 5 to 34 microns. Therefore, as a first step, we had to align them to the 20 subregions from the original experimental data, where each compartment in the

model inherits a corresponding values for the PSD95 puncta density (Table 1) based on their distance from the soma (see also Figure 1 and Methods).

The parameter *u* represents the concentration of PSD95 'cargo' attached to the microtubules and is indexed by compartment. PSD95 can move to neighbouring compartments with rate constants *a* or *b,* for forwards or backwards movement, respectively. Cargo can also irreversibly detach from the microtubule with detachment constant *c,* thus providing a concentration of detached cargo *u*,* which – given the nature of the data – we assume to be mostly 'synaptic' (Figure 1, B).

As was suggested in Williams at al[13], the interplay between the different transport processes available in the model result in behaviours that lie on a spectrum between two extreme modes: One end is dominated by demand-dependent traffic (DDT), where the rate of movement depends on the local demand and the rate of detachment of cargo from the transport system is equal across all compartments. At the other end, demand-dependent detachment (DDD), where the rate of movement is uniform across the cell, while the rate of detachment varies with local demand. The ratio of these modes is represented by the parameter *F* (see Appendix).

In contrast to the Williams model, where all cargo starts at the soma, we already have fully distributed cargo at the outset. Therefore, we extended the model by adding an additional parameter *mProp*, which defines the ratio of cargo:synaptic PSD95 at simulation time 0. Moreover, the original model does not consider degradation rates at synapses, which is clearly a fundamental process affecting the time course of the experimental dataset. Therefore, we further extended the model to add another parameter *d* describing the molecular degradation rate for synaptic PSD95 protein. This extended model can simulate the impact of varying rates of trafficking, detachment (*a,b* and *c*) and degradation *d,* to assess how well these can account for the distribution of protein observed in the biologically obtained datasets.

Inferring the local demand profile from protein distribution data

Our simulations are based on a series of assumptions as follows: First, we assume that puncta density, as measured by Bulovaite et al (2022)[9], correlates with the underlying local demand for PSD95 at the location of the signal. Thus, we can

estimate local demand values from fitting the sushi-belt model to the PSD95 puncta density profile. Second, we assume that protein bound to the polymerized microtubule bundles (*i.e.* cargo) is not measured as it forms the background of the original imaging data. Therefore the measured puncta represent visible pools of PSD95 which we assume to be mostly synaptic. Moreover, in the experimental data, only proteins expressed during day 0 are available to the HaloTag ligand and can subsequently be measured by imaging. Therefore, we can safely ignore the influence of local and somatic translation mechanisms in explaining observed changes in distribution of labelled PSD95 protein between day 0 and day 7. Next, we assume that the neuron is, on average, in a steady state across the 7 days; the total PSD95 at each synapse remains relatively constant but the ratios of labelled vs unlabelled protein change over the time course resulting in the values for SPL. For simplicity, we assume that molecular degradation occurs only at the detached protein (*e.g.* synaptic) and does not affect cargo protein in the active transport system. While our model can support reattachment of synaptic protein back onto the cargo pool, again for simplicity we have excluded this from simulations.

Spatiotemporal distribution of PSD95 is tuned by local transport.

We tested how well the model captures the observed changes in SPL of PSD95 protein across the 20 dendritic regions. We used the Day 0 data to indicate the initial abundance of synaptic PSD95. We then simulated the distribution of synaptically localised protein at Day 7 by adjusting the parameters for traffic, detachment and degradation. Finally, we compared the obtained protein distribution with experimental values (after normalisation for non-uniform number of compartments and Day 0 distribution, for details of the optimisation see Methods).

Given the original complexity of the Migliore and Migliore (2012)[15] model (742 compartments), we tried to minimise model overfitting by keeping the parameter number as low as possible. We started from a model with just three parameters for demand, assuming that demand values might be uniform within each of the three major CA1 regions: CA1so, CA1sr and CA1slm (Figure 1, B). For this simulation, the degradation rate was set to a constant for the whole cell. In addition to the three local demand values and a global degradation rate *d,* we also fit detachment rate *Ctau*, *F*

(ratio between DDD and DDT modes) and *mProp* (ratio between bound and unbound protein at time 0), resulting in seven parameters in total.

We assess each iteration of model for how well the best fit configuration matches the relevant biological dataset (effectively mean squared difference, see Methods). The best fit using this first configuration (Figure 2, A; cost = 0.00024 orange simulation line)   does not precisely follow the experimental day 7 data (blue): Deviations are more obvious in the more distant subregions of apical dendrites, where the trends of experimental and simulation curves critically diverge. Therefore, we concluded that the three regions demand model with uniform degradation is not fully sufficient to describe the experimental data. The optimal value for *F* obtained in this simulation is 0, corresponding to a DDD mode, which means that the model achieves the best fit to observed protein distribution by adjusting the detachment rate while keeping traffic rates uniform.

Next, in the second configuration, we added three distinct degradation rates, one for each of the regions, with its own demand value thus increasing the number of parameters to nine. As shown in Figure 2, B the fit of this model is more accurate (*cost* = 0.00015). The trends for measured and modelled protein distributions at day 7 are now more similar across the proximal regions of the neuron. However, the model still fails to describe the most distant apical dendrite subregions accurately. The obtained value for *F* in this second set is close to 1 (0.99), which means that DDT (demand-dependent traffic) becomes now the dominant process. We noted that in this set of simulations we found that degradation rates correlate positively with demand.

In the final (third) set of simulations, we increased the number of parameters for demand, making it equal to the number of subregions (20) in the experimental data where each of the subregions has its own characteristic value for PSD95 puncta density and SPL. To limit the number of parameters needed we did not implement 20 individual degradation rates, rather we used a linear model with two parameters based on our second simulation results, that degradation is positively correlated with demand:

$$dV_i = dv^A + dv^B * demand_i$$

Here, *demand*$_i$ is the demand value for a specific compartment *i*. dv^A and dv^B are the rates for degradation with dv^A a constant degradation component across the whole neuron and dv^B_i represents a subregion specific degradation component that correlates with local demand.

With this increase in complexity (Figure 2, C), the optimised fit perfectly matches the experimental curve (the orange and blue curves are effectively overlaid). The resulting parameter values are presented in Table 2, with *F* now taking a value of 0.8, which indicates that the system operates primarily in DDT mode, so that redistribution of proteins is achieved mainly by tuning local transportation rate. Meanwhile, the value for *mProp* is 0.3, corresponding to a distribution of 30% of cargo PSD95 and 70% synaptic (Figure 2, D).

Preferential transport of PSD95 to the distal ends of the dendritic tree.

When we estimate the global degradation rate (dv^A) from the optimised model, we get a value corresponding to a half-life of 4.365 days; which aligns well with other experimentally measured values for PSD95 protein (*e.g.* Cohen et al., 2013, etc)[16]. The value for the distance/subregion dependant component dv^B (1.0e-18) is very small, which suggests that the differences in molecular degradation rate across the compartments are likely to be minimal.

Bulovaite et al., (2022)[9] reported an "apparent" protein half-life directly from the experimental imaging data (assuming that the change in concentrations between Day7 and Day 0 fits an exponential decay function within each subregion and that there is no influx of labelled protein). The apparent half-life follows a gradient of increasing PSD95 protein half-life observed in stratum radiatum, followed by significant reduction in stratum lacunosum and then, again an increase towards the most distant subregions of stratum moleculare, (see Figures 1A, 2E green line & 2F blue line).

Given that our model indicates a flat distribution for protein half-life (Figure 2, F, orange line) and that the values for local demand do not closely follow the apparent protein half-life reported, we asked what else could account for the redistribution of dendritically localised fluorescently labelled PSD95 protein between Day 0 and Day 7. From a correlation dendrogram of parameter values (Figure 3, A), we can see that the profile of apparent half-life ($t_{1/2}$) correlates with the parameters *a* (forward traffic rate constant) and *c* (detachment rate), which suggests that the labelled PSD95 distribution can be explained by tuning the movement of bound protein towards the distal dendritic compartment, its sequestration by synapses, followed by degradation.

Figure 3, B and C show how the predicted concentration of labelled PSD95 protein changes in each of the considered neuron subregions from Day 0 to Day 7. The only two compartments where the amount of PSD95 decays exponentially is CA1so_5 and CA1slm_1, and there, the apparent and model half-lives are similar to each other (Figure 2, F). In other compartments, the amount of protein increases from Day 0 to Day 2 due to transport and then decreases by Day 7. The highest increase at Day 2 can be seen in subregions with the highest apparent half-life, such as CA1sr_7-9, and CA1slm_3-5.

PSD95 protein half-life varies with age.

We tested if the same model parameters could also predict the distribution of PSD95 protein in mice of different ages. According to the data from Bulovaite et al., 2022[9], the apparent half-life of PSD95 increases from that seen in juveniles (3-week old mice) to older adults (18-month old mice). We took the best fit model (for 3-month data) and its associated parameters and applied it directly to the data from 3-week and 18-month old animals (Figure 4, A and B), again using their respective puncta density values as initial conditions.

The model predicts a distribution for 3-week old animals that has a similar shape/profile. However, it overestimates the abundance of PSD95 by a considerable margin (Figure 4, A orange model vs blue data). At 18-months, the model prediction is actually very close to the observed data (Figure 4, B). The similar profiles indicate that the core processes are relatively stable with age. The most likely explanation for

the difference at 3 weeks would be faster a molecular degradation rate. Therefore, we adjusted the model to see if allowing degradation rates dv^A and dv^B to vary would allow a better fit. In the 3-week data, a change in dv^A from 1.837e-06 to 5.41e-06 allows an optimal fit and suggests a change in protein half-life between 1.54 days at 3 weeks old compared to 4.3 days observed in mature animals. For a best fit at 18 months, dv^A remains largely unchanged while dv^B has a modest increase from 1.002e-18 to 1.0e-08, indicating that that while global cell rate of degradation rate is constant, there is a trend tends towards degradation being slightly more sensitive to local demand.

The sushi-belt model can explain PSD95 distribution in other types of neurons

We tested if the model was sufficiently generic to be relevant to other neuron types/architectures. Specifically, we focused on the distribution of PSD95 protein in dentate gyrus (DG) granule cells. In contrast to CA1 PyNs, there are few obvious boundaries in DG cells. Here, the PSD95 imaging data suggests a small linear increase in SPL scaling with distance from soma (with the exception of the most distal regions). The linear increase was barely visible in 3-week old animals.

The experimental data was divided into 10 sections of even width from the soma (Figure 5, A). The model we selected (Neuromorphe ID: NMO_80550), reconstructed from the DG of 2 month-old mice[17] was divided into 32 compartments (which equalled the number of branches). To make the DG and CA1 PyN models more comparable, we increased the number of compartments, in proportion to the branch length, to 287. As with the CA1 PyN model, these compartments inherited the experimental values based on the data from equivalent distances from the soma.

Given the relatively simpler cell morphology compared to CA1 PyNs together with a basic gradient of synaptic puncta density distribution, we decided to start from a simpler model of 10 subregions with 10 local demand values and one degradation rate. Figure 5, B shows this model already fits the experimental data (*cost* = 1.51e-

07) and did not require any increase in complexity. Resulting parameters are presented in *F*.

From Table 3, it can be seen that the best fit model operates in the DDD regime (F=0.01), with demand-dependent detachment playing the major role in protein distribution. mProp is 0.49, which means that PSD95 is equally distributed between mobile and detached pools at the start of the study. The predicted degradation rate of PSD95 is higher than that for CA1. Therefore, the model predicted half-life is shorter (1.7 days against 4.3 days in CA1). This matches well with the observation of Bulovaite and colleagues (2022) for faster degradation in granule cells.

We then tested if we were to predict the distribution of puncta density in 3-week and 18-month old mice simply by adjusting the degradation rates. As in the CA1 case, we fixed all model parameters except the degradation rates and refitted the model to the respective experimental data. Figure 5, C and D show the fit of the 3-month model to 3-week data without (*cost* = 0.014) and with (*cost* = 3.94e-16) adjusted degradation rates, respectively. Figure 5, E and F show the fit of the model to the 18-month data, again without (*cost* = 2.98e-05) and with (*cost* = 1.9e-09) adjusted degradation rates, respectively. As with the CA1 PyN model, we found that rate of degradation is much faster (*i.e.* shorter protein half-life) in 3-week old mice (varying in a range of hours instead of days), and only slightly longer in old mice compared to adult 3-month old mice (Table 3).

These results demonstrate consistency of the model effectiveness to explain the protein distribution in different types of neuron cell.

Discussion

How developmental biology and genetics combine to build an organ as complex as the mammalian brain has been a major focus of neuroscience for many decades. The sheer scale and complexity of neuronal networks in brains is multiplied by the cellular diversity of neurons and molecular diversity of synapses that underpin each of these connections. It is highly unlikely that the spatial organisation of diverse synapses is encoded by a genetic program that specifies each synapse and its location. Rather, a few developmental programs could build the synaptome

architecture within the dendritic tree of a neuron and this architecture is then modified by neuronal activity and molecular interactions from the presynaptic terminal of input neurons. There have been remarkable steps in understanding how this happens at the network level. The more recent discovery of the hidden complexity of the synaptome architecture of the brain, where synapses of diverse shapes and molecular compositions are not distributed uniformly along the neuron and different synaptic subtypes can be observed at different localisations of the brain[7], added a new dimension to the conceptual challenge of building and maintaining a complex brain.

Several computational models have previously attempted to describe the dynamics of molecular distribution in neurons (reviewed in Wagle et al., 2022[11]). Models that assume diffusion as the prominent mode of molecular motion are not sufficient to describe experimentally observed protein distributions in dendritic spines, especially when considering the branched neuron structure[18-20]. This shortcoming can be addressed by including active transport of molecules along microtubules, which serve as railroads to provide uni- or bidirectional transport of both molecules and organelles along the long distances in axon and dendrites at a relatively fast velocity[21]. A conceptual model for active transport based on a "tug-of-war" between opposing forces of bi-directional motion was suggested by Hirokawa and Takemura, 2005[22] and Muller et al., 2008[16]. Of particular interest for us was the sushi-belt model of demand-driven movement of molecules [12,13,19,23], which resolved the protein distribution with the minimal number of parameters for the whole realistic heavily-branched neuron morphology. The original sushi-belt model suggested by Doyle and Kiebler in 2011 [12] was designed to explain the conditional distribution of RNA and its supportive protein complex (RNPs)[24]. The model can be applied to other cargo including, synaptic vesicles, mitochondria, pigment granules, liquid droplets, viruses and intermediate filaments[21], and proteins, notably PDZ domain-containing proteins in general[25] and the PSD95 protein itself in particular[26].

Here, we tested a tractable model that aims at describing a molecular basis for creating and maintaining synaptic diversity based on observed data. We

demonstrated that our implementation of the sushi-belt model is sufficient for explaining the spatiotemporal distribution of PSD95: It can accurately describe the changes observed in pulse-labelled PSD95 distribution over 7 days, and, importantly, it can accurately predict the distribution of the protein in young and old animals by simply adjusting the protein degradation rate. We also showed that the model can be successfully applied to different neuron types, showing examples for CA1 pyramidal neurons and DG granule cells. Although we have specified a model for PSD95 data, the code built around this is generally applicable to any protein/RNA data and can be used with minor modifications.

As was suggested in Williams at al[13], the sushi-belt model can operate in different modes, balancing the contribution of traffic and detachment to achieve more rapid or accurate distribution. We observed, for example, that our model prefers a demand-dependent trafficking (DDT) regime in CA1 neurons and demand-dependent detachment (DDD) in DG granule cell neurons. This suggests that resulting balance of the demand-dependent processes is flexible and is regulated by geometry or the molecular genetics of the cell.

Local demand in individual postsynaptic terminals is a dominant factor in our simulations but how this is regulated across the dendritic tree to produce the complex gradients is unclear. One possibility is that the postsynaptic synapse-specific differences in local demand could be induced by inputs from presynaptic terminals. These input projections would likely have different presynaptic protein compositions[27], which in turn influence the clustering and localisation of postsynaptic proteins via trans-synaptic adhesion proteins. Thus, a postsynaptic cell-autonomous mechanism involving the sushi-belt distributes proteins and local demand is regulated by a presynaptic non-cell-autonomous process. Together these two processes could produce the complex synaptome architecture of individual neurons. By simply varying the postsynaptic processes of demand and degradation could result in the observed age-related changes.

Another factor regulating local demand could be the geometry of the dendritic tree. Williams at al., 2016 [13] described so-called "bottlenecks", sites with very low demand, where the trafficking rates tend towards zero. Indeed, in our data we

observed two sites with very low demand, namely CA1so_5 and CA1slm_1. Demand could be modulated by the activity state of the local synaptic environment. For example, locations with particularly low synapse density or very low branch number passing through these areas could dictate the urgency of molecular requirements. In this case, the demand structure should be less stable and will change according to the active states of specific synapses. Exploration of this hypothesis would require activity induced datasets. The likely scenario is that both components (synaptic structure and activity) may influence the final protein distribution, with a balance specific for the respective developmental stage and neuronal cell type.

Our results offer an enticingly simple framework that accounts for synapse specific delivery of molecules without resorting to more complex models where specific proteins are selectively trafficked to specific synapses. Individual synapses can effectively place a local request for more of a specific molecule from a cell-wide delivery system: The global transport mechanisms would provide a pool of molecules to meet local demand. If the global pool is depleted, then the cell just needs to act through generic homeostatic mechanisms to top the global pool back up to 'normal'. This does not preclude synaptic events leading to messages being sent back to the nucleus (or locally) to refine gene expression. Although we have not included local protein synthesis in our model, our simulation suggests it is not required for the observed spatial organisation of the synaptome architecture. We cannot exclude that local protein synthesis could be a fine tuning mechanism by contributing to general protein supply or regulating some other feature of synaptome architecture that was not incorporated into our model.

The framework presented here may help explain how many different diseases result in changes in the synaptome architecture and ultimately in abnormal behaviour. Defects in active transport have been linked to almost every neurodegenerative disorder (e.g. Liu et al 2013)[28]. Moreover, mutations in genes regulating proteostasis are a common cause of autism, intellectual disability and synaptic dysfunction[29]. Mutations in synaptic proteins themselves can influence the turnover of synaptic proteins[30]. These results would imply that any genetic or environmental insult to the

active transport system and the mechanisms of demand and degradation would rapidly impact the neuron's ability to maintain its normal synaptome architecture. Understanding these issues could be facilitated by building from our current simulations models to ones containing diverse interconnected neurons in a realistic a tissue architecture, such as the cortical column, with include simulation of synaptic electrophysiological properties, thereby integrating models of synaptome[31] and connectome architectures at the structural and functional levels.

Methods

Mapping molecular data onto computational model.

In the Bulovoite et al (2022) experimental data, PSD95 puncta density was quantified across a series of windows starting from the soma of the pyramidal neuron in CA1 through to the distal basal dendrites in stratum oriens (so) and to the distal apical dendrites in stratum radiatum (sr), stratum lacunosum (sl), and stratuam moleculare (sm) (Figure 1, A). Moreover, each of these regions was further subdivided: 5 in so, 10 in sr and 5 in sl and sm together (slm in Figure 1, B); giving a total of 20 subregions each containing a value for quantified PSD95 puncta density (schematic representation of the CA1 neuron structure is presented in Figure 1, A, upper panel and B, left panel).

In the model of Migliore and Migliore (2012), which was also used in Williams et al, (2016) there are 742 compartments of five types: "soma", "dend", "axon", "apic" and "dend5". Given that we do not have PSD95 values for the soma or axon, we only considered the "dend", "dend5" and "apic" compartment types. Those three types were mapped onto the experimental data using their relative distance from the soma (Table 1) . Given the highly branched structure of the neuron (Figure 1, B, right panel), compartments the same distance from the soma all inherit the same value for puncta density. Code for mapping is available as a part of Jupyter notebook on GitHub.

Model development

Initiating the data structures to support the neuron morphology model, preparation of the matrix of parameters and part of simulation code was adopted using Python code extracted directly from the Jupyter notebooks accompanying the Williams et al (2016) paper (https://github.com/ahwillia/Williams-etal-Synaptic-Transport). This code was then developed into Python package and extended to include the capacity for experimental data mapping and parameter extraction. All code is available on a GitHub repository (https://github.com/oksankas/Sushi_belt_PSD95).

Model initiation

Initial conditions of the system were assigned as follows: For each compartment, the normalized value $x_i = \frac{V_{seg}}{L_{seg} \Sigma \quad V_{seg}}$ was calculated, where V_{seg} is the experimental value for the segment or subregion at Day 0 and L_{seg} is number of compartments in the segment or subregion. Then it was normalized to the total sum: $\widetilde{x_i} = \frac{x_i}{\Sigma \quad x_i}$. Cargo protein concentrations were assigned value $p_i^{m0} = mProp \times \widetilde{x_i}$, while synaptic protein concentrations were assigned value $p_i^{f0} = (1 - mProp) \times \widetilde{x_i}$.

Simulation and parameter optimisation.

Parameter identification was performed using hybrid optimization techniques originally developed for the Systems Biology Software Infrastructure (SBSI)[32]. In short, 50 rounds of Particle Swarm Optimisation (PSO) were performed for a fast initial exploration of the parameter space, followed by 2000 rounds of Genetic Algorithm (GA) optimisation initiated using the best solutions found by the PSO step. Every 10 rounds of GA the best genome was optimized with 100 rounds of the Nelder-Mead algorithm before next generation construction to accelerate global optimum identification.

Population size for PSO was calculated as $N_{PSO} = max(15, 5 \times N_{par})$, where N_{par} is the number of parameters to identify. Population for GA was selected as $N_{GA} = 2 \times N_{par}$.

A cost function for the optimisation algorithm was calculated as the mean square difference between predicted concentration of the free protein pool at Day7 (p_i^{f7}) and normalized experimental value at that day ($\widetilde{y_i}$): $cost = \sum \left(\widetilde{y_i} - p_i^{f7}\right)^2$. Normalized experimental values were calculated similarly to initial condition protein concentrations: $y_i = \frac{T_{seg}}{L_{seg}\sum V_{seg}}, \widetilde{y_i} = \frac{y_i}{(1-mProp)\sum y_i}$, where T_{seg} is the experimental value for the segment or subregion at the Day 7.

Note that normalisation of the Day 7 experimental value against the sum of Day 0 values requires us to take into account the decay of the total PSD95. Therefore, we used an additional normalisation factor $(1 - mProp)$ because we are fitting only the synaptic component of the protein.

For the DG cell model each topological segment was divided into compartments of length 3.7 μm to be comparable with the CA1 model.

Optimization was performed using PSO and GA implementations from the package scikit-opt version 0.6.6. Nelder-Mead optimizations were performed using the package scipy.optimize version 1.11.3.

Neuron topology import and visualisation were done with NEURON version 8.2.3 and PyNeuron-Toolbox version 0.2 (Alex H Williams. (2014). PyNeuron-Toolbox. *GitHub Repository*. https://github.com/ahwillia/PyNeuron-Toolbox DOI:10.5281/zenodo.12576).

All calculations were performed on Edinburgh Compute and Data Facility (ECDF) the HPC cluster using python 3.10.12.

Aknowledgements

OS, JDA and SGNG were supported by BBSRC Grant number BB/X009343/1, 'Development of a computational model of synaptome architecture'. EB and SGNG were supported by European Research Council (ERC) under the European Union's

Data availability

All code is available on a GitHub repository (https://github.com/oksankas/Sushi_belt_PSD95).

References

CAJAL, D. S. R. Y. Nueva concepta de la histologia de los centros nervosos. *Annals of Surgery* **18**, 122 (1893).
Bayes, A. *et al.* Characterization of the proteome, diseases and evolution of the human postsynaptic density. *Nat Neurosci* **14**, 19-21 (2011). https://doi.org/10.1038/nn.2719
Frank, R. A. W. *et al.* NMDA receptors are selectively partitioned into complexes and supercomplexes during synapse maturation. *Nature Communications* **7**, 11264 (2016). https://doi.org/10.1038/ncomms11264
Sorokina, O., Dominic, D., Bayés, À., Armstrong, J. D. & Grant, S. G. N. SynaptopathyDB: a resource for studying the genetic and synaptic basis of nervous system disorders. *bioRxiv*, 2025.2006.2027.661932 (2025). https://doi.org/10.1101/2025.06.27.661932
Bayes, A. *et al.* Characterization of the proteome, diseases and evolution of the human postsynaptic density. *Nat Neurosci* **14**, 19-21 (2011). https://doi.org/10.1038/nn.2719
Zhu, F. *et al.* Architecture of the Mouse Brain Synaptome. *Neuron* **99**, 781-799.e710 (2018). https://doi.org/10.1016/j.neuron.2018.07.007
Cizeron, M. *et al.* A brainwide atlas of synapses across the mouse life span. *Science* **369**, 270-275 (2020). https://doi.org/10.1126/science.aba3163
Distler, U. *et al.* Proteomic Analysis of Brain Region and Sex-Specific Synaptic Protein Expression in the Adult Mouse Brain. *Cells* **9** (2020). https://doi.org/10.3390/cells9020313
Bulovaite, E. *et al.* A brain atlas of synapse protein lifetime across the mouse lifespan. *Neuron* **110**, 4057-4073.e4058 (2022). https://doi.org/10.1016/j.neuron.2022.09.009

Morris, K. *et al.* Sequential replacement of PSD95 subunits in postsynaptic supercomplexes is slowest in the cortex. *Elife* **13** (2024). https://doi.org/10.7554/eLife.99303

Wagle, S., Kraynyukova, N., Hafner, A. S. & Tchumatchenko, T. Computational insights into mRNA and protein dynamics underlying synaptic plasticity rules. *Mol Cell Neurosci* **125**, 103846 (2023). https://doi.org/10.1016/j.mcn.2023.103846

Doyle, M. & Kiebler, M. A. Mechanisms of dendritic mRNA transport and its role in synaptic tagging. *Embo j* **30**, 3540-3552 (2011). https://doi.org/10.1038/emboj.2011.278

Williams, A. H., O'Donnell, C., Sejnowski, T. J. & O'Leary, T. Dendritic trafficking faces physiologically critical speed-precision tradeoffs. *Elife* **5** (2016). https://doi.org/10.7554/eLife.20556

McDougal, R. A., Hines, M. L. & Lytton, W. W. Reaction-diffusion in the NEURON simulator. *Front Neuroinform* **7**, 28 (2013). https://doi.org/10.3389/fninf.2013.00028

Migliore, M. & Migliore, R. Know your current I(h): interaction with a shunting current explains the puzzling effects of its pharmacological or pathological modulations. *PLoS One* **7**, e36867 (2012). https://doi.org/10.1371/journal.pone.0036867

Cohen, L. D. *et al.* Metabolic turnover of synaptic proteins: kinetics, interdependencies and implications for synaptic maintenance. *PLoS One* **8**, e63191 (2013). https://doi.org/10.1371/journal.pone.0063191

Revest, J. M. *et al.* Adult hippocampal neurogenesis is involved in anxiety-related behaviors. *Mol Psychiatry* **14**, 959-967 (2009). https://doi.org/10.1038/mp.2009.15

Li, T. P., Song, Y., MacGillavry, H. D., Blanpied, T. A. & Raghavachari, S. Protein Crowding within the Postsynaptic Density Can Impede the Escape of Membrane Proteins. *J Neurosci* **36**, 4276-4295 (2016). https://doi.org/10.1523/jneurosci.3154-15.2016

Bressloff, P. C. & Earnshaw, B. A. Diffusion-trapping model of receptor trafficking in dendrites. *Phys Rev E Stat Nonlin Soft Matter Phys* **75**, 041915 (2007). https://doi.org/10.1103/PhysRevE.75.041915

Sartori, F. *et al.* Statistical Laws of Protein Motion in Neuronal Dendritic Trees. *Cell Rep* **33**, 108391 (2020). https://doi.org/10.1016/j.celrep.2020.108391

Welte, M. A. Bidirectional transport along microtubules. *Curr Biol* **14**, R525-537 (2004). https://doi.org/10.1016/j.cub.2004.06.045

Hirokawa, N. & Takemura, R. Molecular motors and mechanisms of directional transport in neurons. *Nat Rev Neurosci* **6**, 201-214 (2005). https://doi.org/10.1038/nrn1624

Newby, J. M. & Bressloff, P. C. Quasi-steady state reduction of molecular motor-based models of directed intermittent search. *Bull Math Biol* **72**, 1840-1866 (2010). https://doi.org/10.1007/s11538-010-9513-8

Kiebler, M. A. & Bauer, K. E. RNA granules in flux: dynamics to balance physiology and pathology. *Nat Rev Neurosci* **25**, 711-725 (2024). https://doi.org/10.1038/s41583-024-00859-1

Setou, M., Nakagawa, T., Seog, D. H. & Hirokawa, N. Kinesin superfamily motor protein KIF17 and mLin-10 in NMDA receptor-containing vesicle transport. *Science* **288**, 1796-1802 (2000). https://doi.org/10.1126/science.288.5472.1796

Yoo, K.-S. *et al.* Postsynaptic density protein 95 (PSD-95) is transported by KIF5 to dendritic regions. *Molecular Brain* **12**, 97 (2019). https://doi.org/10.1186/s13041-019-0520-x

Reig-Viader, R. *et al.* Synaptic proteome diversity is shaped by the levels of glutamate receptors and their regulatory proteins. *bioRxiv*, 2024.2004.2004.588090 (2025). https://doi.org/10.1101/2024.04.04.588090

Liu, X. A., Rizzo, V. & Puthanveettil, S. V. PATHOLOGIES OF AXONAL TRANSPORT IN NEURODEGENERATIVE DISEASES. *Transl Neurosci* **3**, 355-372 (2012). https://doi.org/10.2478/s13380-012-0044-7

Louros, S. R. & Osterweil, E. K. Perturbed proteostasis in autism spectrum disorders. *Journal of neurochemistry* **139**, 1081-1092 (2016).

Bulovaite, E. *et al.* A brain atlas of synapse protein lifetime across the mouse lifespan. *Neuron* (2022). https://doi.org/https://doi.org/10.1016/j.neuron.2022.09.009

Zhu, F. *et al.* Architecture of the Mouse Brain Synaptome. *Neuron* **99**, 781-799 e710 (2018). https://doi.org/10.1016/j.neuron.2018.07.007

Adams, R. *et al.* SBSI: an extensible distributed software infrastructure for parameter estimation in systems biology. *Bioinformatics* **29**, 664-665 (2013). https://doi.org/10.1093/bioinformatics/btt023

Figure 1. A. Model of a pyramidal CA1 neuron colour coded by values for normalised distribution of PSD95 protein taken on day 0 of the Halo-ligand injection (from Bulovaite et al., 2022). The apical dendrite traverses three subregions of the hippocampus (stratum radiatum, lacunosum and moleculare) and the basal dendrites occupy the stratum oriens. Each subregion of the CA1 receives distinct afferent inputs: CA1 stratum radiatum receives projections from CA3, while stratum

lacunosum is targeted by entorhinal cortex layer 2 and stratum moleculare by entorhinal cortex layer 3. Stratum oriens receives inputs from CA2 and CA3 pyramidal neurons. B. Upper: schematic representation of four neighbouring sushi-belt model compartments, with cargo (u1, u2, u3, u4) and synaptic (u1*, u2*, u3*, u4*) PSD95 in each compartment. Lower; mass-action model with respective kinetic constants taken from Williams et al (2016).

Figure 2. Fitting of the sushi-belt model to the experimental data from three month old mice; D7M3 is a normalized protein distribution obtained from Day 7 puncta density data. A. Best fit for a simple model with three compartments and one degradation rate (blue line with dots – experimental data, orange line – simulation) B. Fitting of model with three compartments and three degradation rates. C Fitting results of final model of 20 compartments and two parameters for degradation rates (blue and orange lines are congruent). D. for the best fit as presented in C shown the both the synaptic protein pool obtained from model (green, not normalised, see Methods for details) and cargo / belt-bound protein pool (orange, 30 % of total as corresponds to mProp = 0.3). E. Apparent half-life estimated from experimental data based on an assumption of exponential decay (green line) is plotted against estimated local demand for each compartment (blue line). F. Apparent half-life estimated from experimental data (blue line) plotted against model half-life (orange line) estimated from simulations.

Figure 3. Results of model optimisation. A. Heat map of z-normalized parameters (shown as row names), which are dependent of local demand, including the apparent half-life and the local demand itself. B. Time series of changes in synaptic PSD95 abundance estimated for each compartment based on model prediction C. Similarly to B, changes in synaptic PSD95 are plotted individually.

Figure 4. Modelling PSD95 in mice younger (3 week) and older (18 months). A. D7W3 is the normalized protein distribution obtained from 3-week old mice 7 days post labelling (blue) against the simulation (orange) using the best parameters from the 3-month old simulations. B. As Panel A but using data from 18-month old mice. C. As panel A (3-week old) except with a re-optimised degradation rate. D. As B (18-month old) except with a re-optimised degradation rate.

Figure 5. Sushi-belt model application to the PSD95 distribution in Dentate Gyrus cell. A. Schematic image of granule cell obtained from Neuromorpho model (ID: NMO_80550). Panel A shows the 10 subregions considered in the cell. Colour coding indicates normalised puncta density measured in 3-month old mice seven days after labelling. B. Result of fitting the sushi-belt model (orange line) to the experimental data (blue line) of 3-month old mice. C. As B, except comparing a 3-month simulation to data from 3-week old animals. D. As C, except allowing optimisation of the degradation rate. E. As B, except comparing 3-month simulation to data from 18-month old mice. F. As E, except allowing optimisation of the degradation rate.

Neuron subregion number	Neuron subregion name	Subregion abbreviation	Distance from the soma	Model compartment name	Number of model compartments
1	CA1so	CA1so_1	-165	dend	9
2	CA1so	CA1so_2	-127.5	dend	85
3	CA1so	CA1so_3	-90	dend	134
4	CA1so	CA1so_4	-52.5	dend	100
5	CA1so	CA1so_5	0	dend	16
6	CA1sr	CA1sr_1	10	apic	3
7	CA1sr	CA1sr_2	52.5	apic	6
8	CA1sr	CA1sr_3	94.286	apic	29
9	CA1sr	CA1sr_4	136.43	apic	37
10	CA1sr	CA1sr_5	178.57	apic	48
11	CA1sr	CA1sr_6	220.71	apic	44
12	CA1sr	CA1sr_7	262.86	apic	45
13	CA1sr	CA1sr_8	305	apic	38
14	CA1sr	CA1sr_9	347.14	apic	21
15	CA1sr	CA1sr_9	389.29	apic	9
16	CA1slm	CA1slm_1	431.43	dend5	13
17	CA1slm	CA1slm_2	473.57	dend5	24
18	CA1slm	CA1slm_3	515.71	dend5	35
19	CA1slm	CA1slm_4	557.86	dend5	29
20	CA1slm	CA1slm_5	600	dend5	9

Table 1. Mapping of CA1 neuron subregions to the model of reconstructed pyramidal neuron. Abbreviations for CA1 neuron subregions mapped on the model compartments from Migliore and Migliore (2012). Each of 20 subregions measured in experiment correspond to 6-134 model compartments located at the equal distance from the soma.

Parameter name	Definition	Fitted value 3m	Fitted value 3w	Fitted value

				18m
F	Ratio between DDT (1) and DDD (0) modes	0.8861	0.8861	0.8861
Ctau	Coefficient for demand-dependent detachment	4.38e-06	4.38e-06	4.38e-06
mProp	Ratio between mobile and detached polls protein pools	0.312	0.312	0.312
dvA	Intercept for degradation linear approximation	1.837e-06	**5.21e-06**	1.63e-06
dvB	Slope for degradation linear approximation	1.002e-18	1e-18	**9.57e-08**
Demand_CA1so_1	Local demand value for st.oriens subregion 1	0.214	0.214	0.214
Demand_CA1so_2	Local demand value for st.oriens subregion 2	0.061	0.061	0.061
Demand_CA1so_3	Local demand value for st.oriens subregion 3	0.047	0.047	0.047
Demand_CA1so_4	Local demand value for st.oriens subregion 4	0.045	0.045	0.045
Demand_CA1so_5	Local demand value st.oriens subregion 5	1.0e-07	1.0e-07	1.0e-07
Demand_CA1sr_1	Local demand value for st.radiatum subregion 1	0.773	0.773	0.773
Demand_CA1sr_2	Local demand value for st.radiatum subregion 2	0.554	0.554	0.554
Demand_CA1sr_3	Local demand value for st.radiatum subregion 3	0.147	0.147	0.147
Demand_CA1sr_4	Local demand value for st.radiatum subregion 4	0.147	0.147	0.147
Demand_CA1sr_5	Local demand value for st.radiatum subregion 5	0.131	0.131	0.131
Demand_CA1sr_6	Local demand value for st.radiatum subregion 6	0.154	0.154	0.154
Demand_CA1sr_7	Local demand value for st.radiatum subregion 7	0.172	0.172	0.172
Demand_CA1sr_8	Local demand value for st.radiatum subregion 8	0.202	0.202	0.202
Demand_CA1sr_9	Local demand value for st.radiatum subregion 9	0.373	0.373	0.373
Demand_CA1sr_10	Local demand value for st.radiatum subregion 10	0.524	0.524	0.524
Demand_CA1slm_1	Local demand value for st.lacunose and moleculare subregion 1	0.043	0.043	0.043
Demand_CA1slm_2	Local demand value for st.lacunose and moleculare subregion 2	0.196	0.196	0.196
Demand_CA1slm_3	Local demand value for st.lacunose and moleculare subregion 3	0.196	0.196	0.196
Demand_CA1slm_4	Local demand value for st.lacunose and moleculare subregion 4	0.275	0.275	0.275
Demand_CA1slm_5	Local demand value for st.lacunose and moleculare subregion 5	0.999	0.999	0.999

Table 2. List of parameters obtained in the optimisation for 3-month, 3-week and 18-month old mice in CA1 neuron.

Parameter name	Definition	Value for	Value for 3W	Value for 18M

		3M		
F	Ratio between DDT (1) and DDD (0) modes	0.01	0.01	0.01
Ctau	Coefficient for demand-dependent detachment	2.3e-06	2.3e-06	2.3e-06
mProp	Ratio between mobile and detached polls protein pools	0.49	0.49	0.49
demand_rd1	Local demand value for rd1	1.03e-07	1.03e-07	1.03e-07
demand_rd2	Local demand value for rd2	0.029	0.029	0.029
demand_rd3	Local demand value for rd3	0.012	0.012	0.012
demand_rd4	Local demand value for rd4	0.041	0.041	0.041
demand_rd5	Local demand value for rd5	0.067	0.067	0.067
demand_rd6	Local demand value for rd6	0.089	0.089	0.089
demand_rd7	Local demand value for rd7	0.151	0.151	0.151
demand_rd8	Local demand value for rd8	0.610	0.610	0.610
demand_rd9	Local demand value for rd9	0.543	0.543	0.543
demand_rd10	Local demand value for rd10	0.281	0.281	0.281
dv	Degradation rate	4.6e-06	**5.52e-04**	**4.53e-06**
$T_{1/2}$	PSD95 protein half-life, days	1.74	**0.014**	**1.77**

Table 3. List of parameters obtained in the optimisation for 3-month, 3-week and 18-month old mice in DG neuron.