Batch culture and species effects modulate the decoupling between diatom frustule-bound and biomass nitrogen isotope signatures

Jochem Baan[1]*, Moritz F. Lehmann[1], Anja S. Studer[1]†

[1]University of Basel, Department of Environmental Sciences, Bernoullistrasse 30, 4056 Basel, Switzerland

†Now at Laboratory of Ion Beam Physics, ETH Zurich, Otto-Stern-Weg 5, 8093 Zurich, Switzerland

*Corresponding author e-mail: jochem.baan@unibas.ch

Abstract

The stable nitrogen (N) isotope composition of organic matter encapsulated in diatom silica frustules ($\delta^{15}N_{DB}$) from sedimentary records has been used as a proxy for reconstructing N consumption dynamics in the ocean over geologic timescales. This proxy relies on the assumption that $\delta^{15}N_{DB}$ tracks biomass $\delta^{15}N$ without being affected by internal N-isotope fractionation. However, recent ground-truthing efforts have shown that $\delta^{15}N_{DB}$ can diverge from biomass $\delta^{15}N$ values, though the extent and mechanisms behind this decoupling remain unclear.

In this study, we cultured two freshwater and two marine diatom species in batch cultures to test whether $\delta^{15}N_{DB}$ (1) is subject to species-dependent internal ^{15}N fractionation, and (2) reflects the $\delta^{15}N$ of source nitrate to the same extent as biomass $\delta^{15}N$ values, assessing potential asynchronous integration of the N isotope signal. We monitored the N-isotope systematics during diatom growth by measuring $\delta^{15}N$ values of nitrate, bulk biomass and frustule-bound organic N throughout batch culture progression. We found that $\delta^{15}N_{DB}$ did not follow typical Rayleigh fractionation dynamics and remained relatively stable, while biomass $\delta^{15}N$ increased predictably with progressive fractional nitrate consumption. The observed divergence could only be partially explained by asynchronous integration of source-nitrate $\delta^{15}N$ values into biomass versus frustule-bound organic N (i.e., delayed incorporation into frustule-bound material), as newly formed frustules predominantly recorded the $\delta^{15}N$ of ^{15}N-labeled nitrate added during growth. This demonstrates that $\delta^{15}N_{DB}$ values capture the isotopic signature of newly assimilated nitrate rather than N derived from internal, or legacy, pools. We hypothesize that shifts in growth conditions during batch culture progression alter the coupling between carbon and nitrogen metabolism, leading to physiologically driven variation in internal ^{15}N fractionation and corresponding offsets between $\delta^{15}N_{DB}$ and biomass $\delta^{15}N$. Such sensitivity to internal isotope fractionation during

34 biosynthesis implies that the interpretation of sedimentary $\delta^{15}N_{DB}$ records is more intricate than

35 previously assumed.

37 **Keywords:** biomass, diatom, frustule-bound organic compounds, isotope fractionation,

38 nitrate, nitrogen stable isotopes

1. Introduction

The nitrogen (N) isotopic composition ($\delta^{15}N$) of diatom frustule-bound organic matter ($\delta^{15}N_{DB}$) from sedimentary archives has been used as a promising proxy indicator for reconstructing nitrate utilization across various timescales in the ocean (Sigman et al., 1999; Robinson et al., 2004; Ren et al., 2015; Studer et al., 2015; Ai et al., 2020), and more recently, in lacustrine environments for paleoenvironmental reconstructions (Studer et al., 2024). The main advantage of this proxy is that $\delta^{15}N_{DB}$ is thought to be physically protected from diagenetic alteration and bacterial degradation, unlike the $\delta^{15}N$ of organic matter in bulk sediment which is known to be susceptible to post-depositional alterations (Lehmann et al., 2002; Robinson et al., 2012). Despite the fact that the $\delta^{15}N_{DB}$ proxy has been applied in palaeoceanographic research for more than two decades, the mechanisms by which the $\delta^{15}N$ signature of assimilated inorganic N is transferred into diatom biomass, and ultimately preserved in $\delta^{15}N_{DB}$, remain poorly understood.

When diatoms grow, their cell walls become silicified, forming opaline frustule structures. This silicification occurs after mitosis, through the synthesis of new valves, and during cell expansion, when girdle bands are formed (Kröger and Poulsen, 2008). Frustule formation is facilitated by several N-containing organic compounds (e.g., silaffins, silacidins, and polyamines), which act as organic templates for silicate precipitation (Poulsen et al., 2003; Bridoux and Ingalls, 2010; Richthammer et al., 2011). These compounds become encapsulated within the frustule and can persist over long time scales. The N in these frustule-bound organic compounds originates from the intracellular organic N pool in the diatom, which in turn derives from the assimilation of inorganic N in the surrounding water. Under this framework, the $\delta^{15}N$ values of assimilated inorganic N should be transferred into the cellular organic N pool and therefore reflected by $\delta^{15}N_{DB}$. Accordingly, $\delta^{15}N_{DB}$ values have been interpreted as proxies for nitrate utilization

(Robinson et al., 2005; Brunelle et al., 2007), nitrogen cycling in general (Kalansky et al., 2011),

and N eutrophication (Studer et al., 2024).

Efforts to ground-truth the diatom frustule-bound $\delta^{15}N$ paleo-proxy through controlled

batch culture experiments have shown that $\delta^{15}N_{DB}$ values can differ substantially from biomass

$\delta^{15}N$ values (Horn et al., 2011; Jones et al., 2022; Robinson et al., 2025). Horn et al. (2011) found

$\delta^{15}N_{DB}$ values lower than those of biomass, with species-specific offsets, whereas Jones et al.

(2022) reported the opposite pattern. More recently, Robinson et al. (2025) observed both positive

and negative offsets, depending on species, and suggested that the nitrate-to-silicate uptake ratio

may play a modulating role. Overall, these findings suggest that $\delta^{15}N_{DB}$ values can vary

substantially and do not consistently track biomass $\delta^{15}N$ values, and that the underlying drivers of

the offset remain poorly understood.

Culturing diatoms under controlled environmental conditions enables detailed observation

of N isotope transfer from source nitrate into diatom biomass and frustule-bound material. In batch

cultures, the $\delta^{15}N$ values of nitrate and accumulated biomass typically follow Rayleigh-type

substrate-product dynamics (Mariotti et al., 1981; Waser et al., 1998). Once assimilated, N is

processed into various proteins and other N-containing biomolecules, including those found in

diatom frustules (Sumper and Brunner, 2008; Kröger and Poulsen, 2008; Bridoux and Ingalls,

2010; Richthammer et al., 2011; Kirkham et al., 2017). As diatoms synthesize frustules over the

course of a batch culture, the frustule-bound organic N pool accumulates. It might therefore be

expected that $\delta^{15}N_{DB}$ values reflect the integrated isotopic composition of the accumulating

frustule-bound organic matter, similar to the integrated $\delta^{15}N$ signal in bulk biomass (Mariotti et

al., 1981). However, this expectation rests on two key assumptions: (1) that ^{15}N fractionation

associated with the synthesis of frustule-bound organic compounds is constant, and (2) that the

integration time window of the isotope-signal incorporation from the inorganic N source is the

same for biomass and frustule-bound organic N. With regards to the first assumption, internal N-

isotope fractionation during biosynthesis may vary due to metabolism-driven biochemical

reactions (Werner and Schmidt, 2002; Tcherkez and Hodges, 2008). Additionally, the $\delta^{15}N$ values

of amino acids (the building blocks of proteins and polyamines) can vary between compounds and

across species (Macko et al., 1987; Werner and Schmidt, 2002), suggesting that $\delta^{15}N_{DB}$ values

may also depend on (species-specific) differences in the frustule-bound organic N

matrix/composition (Sumper and Brunner, 2008; Bridoux and Ingalls, 2010). As for the second

assumption, frustule-bound organic N may derive from an internal N pool with a distinct isotopic

signature, potentially leading to asynchronous $\delta^{15}N$-signal integration relative to bulk biomass. To

date, neither of these assumptions have been experimentally verified, yet the observation that

$\delta^{15}N_{DB}$ values can be both higher and lower than biomass $\delta^{15}N$ values (Horn et al., 2011; Jones et

al., 2022; Robinson et al., 2025) implies that at least one of them may not hold. As such, it remains

uncertain whether $\delta^{15}N_{DB}$ values consistently reflect biomass and/or nitrate $\delta^{15}N$ values, and thus

exhibit predictable Rayleigh-type fractionation behavior.

Diatom culture studies aimed at ground-truthing the $\delta^{15}N_{DB}$ proxy have so far exclusively

focused on marine species. However, since the proxy has recently also been applied in lacustrine

systems (Studer et al., 2024), it is important to investigate the relationship between nitrate, biomass

and frustule-bound $\delta^{15}N$ values in freshwater diatoms. Moreover, freshwater species generally

exhibit higher silicate uptake rates compared to most marine species (Conley et al., 1989).

Therefore, further insight into whether and how silicate-to-nitrate uptake ratios influence $\delta^{15}N_{DB}$

relative to biomass $\delta^{15}N$ (Robinson et al., 2025) may be gained by including freshwater taxa in the

experimental design.

In this study, we cultured two marine and two freshwater diatom species under controlled

environmental conditions, with nitrate as the limiting nutrient. These batch cultures were closely

monitored, and $\delta^{15}N$ values of nitrate, bulk biomass and frustule-bound organic N were monitored

throughout the progression of each culture. This experimental setup allowed us to specifically test

whether variation in frustule-bound $\delta^{15}N$ could be explained by that of nitrate and/or biomass, and

whether the observed patterns were consistent (1) among species, (2) between ecosystems

(freshwater vs. marine), and (3) over time during batch culture progression. In a separate ^{15}N-

labeling experiment we manipulated the $\delta^{15}N$ of the nitrate source during growth, to assess if bulk-

biomass and frustule-bound $\delta^{15}N$ values capture the nitrate isotope signal concurrently, or whether

differences in isotopic integration timing exist between these N pools. Overall, our results highlight

that $\delta^{15}N_{DB}$ values do not strictly follow Rayleigh-type dynamics in batch culture. However, this

deviation from expected patterns is not due to (species-specific) differences in the timing of nitrate-

$\delta^{15}N$ incorporation into diatom biomass versus frustule-bound N. Instead, our findings suggest

that the observed offset is more likely driven by variable, growth-condition-dependent biochemical

^{15}N fractionation.

2. Methods

2.1. Diatom cultures

Two colony-forming freshwater species were selected: the centric *Aulacoseira granulata* and the pennate *Fragilaria capucina*, both isolated from Swiss lakes and kindly provided by Marta Reyes (Swiss Federal Institute for Aquatic Research, Dubendorf, Switzerland). Two marine diatom species forming individual cells were obtained from the Culture Collection of Algae & Protozoa (CCAP, Scottish Association for Marine Science, Dunbeg, Scotland, United Kingdom): *Chaetoceros calcitrans* (CCAP 1085/3), a small centric diatom, and *Coscinodiscus radiatus* (CCAP 1013/11), a large centric diatom.

The growth medium for freshwater diatoms was based on the CCAP 'Diatom Medium' recipe, with some modifications, resulting in the following starting concentrations: 102 μM $Ca(NO_3)_2$, 100 μM K_2HPO_4, 101 μM $MgSO_4$, 200 μM $KHCO_3$, 200-350 μM Na_2SiO_3, 36 μM Fe-EDTA, 5 μM $MnCl_2$, 0.2 μM $CuSO_4$, 0.9 μM $ZnSO_4$, 0.5 μM Na_2MoO_4, 40 μM H_3BO_3, and vitamins (0.04 mg/L cyanocobalamin, thiamine HCl, and biotin each).

The medium for marine diatoms was based on the ESAW artificial seawater recipe (Harrison et al., 1980; Berges et al., 2001), with some modifications. The base seawater composition followed ESAW with the exception that NaBr was used instead of KBr. Micronutrient composition also followed ESAW, except $CoSO_4$ was replaced with CoCl. Since copper is not typically included in ESAW, but was added to the freshwater medium, 0.1 μM of $CuSO_4$ was also added here. Iron was provided as 36 μM Fe-EDTA. The same vitamin concentrations were used as in the freshwater medium. Lastly, nitrate, phosphate and silicate were added at starting concentrations of 254 μM $Ca(NO_3)_2$, 103 μM K_2HPO_4 and 150-350 μM Na_2SiO_3. In order to

obtain sufficient frustule material from *C. calcitrans*, which is weakly silicifying, a higher initial

nitrate concentration (~508 μM) was used. The pH of the artificial seawater medium was adjusted

to 8.1 ± 0.1 using a 0.5 M NaOH prior to filtration.

All media were filter-sterilized using sterile 0.2 μm polyether sulfone (PES) membranes

into autoclaved containers. Since the Na_2SiO_3 stock solution had a high pH (12-13), its addition

influenced the pH of the medium and caused precipitation and filtration problems. Therefore,

silicate was added aseptically *after* filtration. In some freshwater batch cultures, 400 mg/L N-

Tris(hydroxymethyl)methyl-2-aminoethanesulphonic acid (TES) buffer was added to stabilize pH

during growth, and to counteract the pH rise from Na_2SiO_3 addition (Table 1). For marine cultures

and freshwater batch cultures 5 & 6, silicate was added as a neutralized colloid (Table 1): a 50

mM Na_2SiO_3 stock solution was titrated with 1M HCl to pH 7.5-8.5 while stirring, forming a

Na_2SiO_3 colloid that was then aseptically added post-filtration.

Diatom culture stocks were maintained in 100-200 mL Erlenmeyer flasks, filled with 50-

100 mL of medium and covered with aluminum foil. Cultures were grown at 20 °C under a 12h/12h

light/dark cycle with ~140 μmol m^{-2} s^{-1} photons from LED lighting in the growth chambers

(MLR-350, SANYO, Japan). Stocks were manually stirred daily, and refreshed every 2-4 weeks,

either by addition of medium, or by transferring 5 mL of culture into fresh 50 mL medium in a

new flask.

The batch culture experiments were conducted in 20-L clear polycarbonate carboys filled

with sterile-filtered medium. Each carboy was fitted with three ports: one for sampling, one for air

inflow to aerate the medium, and one for air outflow. Air was supplied via a compressed air system

through 0.2 μm hydrophobic PTFE filters. Carboys were placed in a large custom-built walk-in

phytotron growth chamber at 20 °C, with ceiling-mounted LEDs providing ~140 μmol m^{-2} s^{-1}

photons. A 12h/12h light/dark cycle was used, with gradual light-intensity ramp-up and ramp-

down over the first and last hour of the light period, respectively. Carboys were equilibrated with

ambient air for at least 24 h prior to inoculation. Cultures were inoculated with 20-40 mL of

recently transferred stock culture. This relatively small inoculum delayed biomass sample

collection in early incubation stages but minimized disruption of expected closed-system isotope

dynamics. Each species was cultured in at least two replicate batch cultures.

Media were regularly sampled to monitor growth, pH, and nutrient (i.e., nitrate and silicate)

consumption. Growth was tracked via *in vivo* fluorescence using an AquaFluor handheld

fluorometer (Turner Designs, CA, USA). Samples were filtered through 0.22 μm PES syringe

filters. In freshwater cultures, nitrate concentrations were measured by spectrophotometric

analysis using LCK339 nitrate test kits (Hach, Germany). In artificial seawater, nitrate was

measured using a chemiluminescence method (described below) due to interference from high ion

concentrations with the nitrate test. For dissolved silica analysis, filtered samples were acidified to

~pH 2 using 1 μL 6N HCl per mL of sample. Silicate concentrations were determined

photometrically (Strickland and Parsons, 1972), using molybdate-3, citric acid, and amino acid F

reagents, and calibrated against a 1 mg/l SiO_2 standard (Hach, Germany). When silicate became

limiting before a substantial nitrate fraction was consumed, additional silicate was added to the

batch culture. Samples for nitrate concentration and $\delta^{15}N$ analysis were stored frozen at -20 °C.

Silicate samples were stored at room temperature to avoid polymerization at high silicate

concentrations.

Once biomass reached sufficient density – based on nitrate and silicate consumption – 1-4

L samples were harvested at regular time intervals. Biomass was concentrated by centrifugation

and split into two aliquots: one for bulk biomass $\delta^{15}N$ and $\delta^{13}C$ measurements and elemental (N

and C) analysis, and one for the analysis of frustule-bound $\delta^{15}N$ values. Bulk-biomass aliquots

were frozen at -20 °C, freeze-dried, and stored in the dark at room temperature prior to analysis.

Frustule-bound $\delta^{15}N$ samples were stored frozen at -20 °C until further processing.

Two replicate batch-cultures of each freshwater diatom species (culture numbers 5 & 6;

Table 1) were grown as described above, except that after the first 2-4 biomass samples had been

harvested, and once nitrate had decreased to ~50 % of its initial concentration, a 4 mM solution of

^{15}N-labelled nitrate ($\delta^{15}N \approx 4000$ ‰) was added. The volume added was calculated based on

mass-balance targeting an 80-100 ‰ increase in nitrate $\delta^{15}N$, with minimal change in nitrate

concentration (1-2 µM). After the ^{15}N-nitrate spike, cultures were monitored and harvested

regularly until all nitrate was consumed. In this experiment, the relative abundance of empty

frustules in harvested biomass was assessed microscopically to determine whether a potential

decoupling between biomass and frustule-bound $\delta^{15}N$ value could result from an increased

contribution of older, unlabeled frustules.

2.2. Sample processing and isotope analysis

2.2.1. Nitrate concentration measurements

Nitrate concentrations were determined using a chemiluminescence method. Briefly,

samples were injected into a heated 0.1 M vanadium(III) in 2N HCl solution to convert sample

nitrate to NO (Braman and Hendrix, 1989), followed by chemiluminescent detection of the NO

gas after reaction with ozone in a TELEDYNE T200 NO/NO$_2$/NOx analyzer (TELEDYNE API,

CA, USA). Sample nitrate concentrations were determined by comparison to multiple injections

of KNO$_3$ reference standards of known nitrate concentrations.

2.2.2.   Biomass $\delta^{15}N$ and $\delta^{13}C$ values, and N and C content

Aliquots of 1-3 mg of freeze-dried diatom biomass were weighed into tin cups. Biomass

$\delta^{15}N$ and $\delta^{13}C$ values, as well as total N and C contents were determined using a Delta V Plus

isotope ratio mass spectrometer (IRMS; Thermo Fisher Scientific) coupled to a Flash Elemental

Analyzer (EA). Scale normalization for $\delta^{15}N$ values was performed using USGS61 ($\delta^{15}N$ = -2.87

‰; $\delta^{13}C$ = -35.05 ‰), and USGS62 ($\delta^{15}N$ = 20.17 ‰; $\delta^{13}C$ = -14.79 ‰) international reference

materials. For samples from the ^{15}N-labeled nitrate addition experiments, we additionally included

USGS41 ($\delta^{15}N$ = 47.57 ‰; $\delta^{13}C$ = 37.63 ‰). Scale normalization for $\delta^{13}C$ values was based on

USGS40 ($\delta^{15}N$ = -4.52 ‰; $\delta^{13}C$ = -26.39 ‰), USGS62, and USGS64 ($\delta^{15}N$ = 1.76 ‰; $\delta^{13}C$ = -

40.81 ‰) standards. Elemental N and C contents were determined from IRMS peak areas using a

calibration curve that was established based on the analysis of a wheat flour standard (1.36 % N;

39.38 % C; Säntis analytical, Switzerland) and EDTA (9.58 % N; 41.06 % C). Analytical

reproducibility was assessed by repeated analysis (n = 5) of biomass from a commercial diatom

culture (*Thalassiosira weissflogii*, TW1800, Reed Mariculture, CA, USA), yielding mean ± SD

$\delta^{15}N$ and $\delta^{13}C$ values of 6.9 ± 0.3 ‰ and -10.50 ± 0.1 ‰, respectively.

2.2.3.   Frustule-bound and ambient nitrate $\delta^{15}N$ analysis

All glassware used in frustule-bound $\delta^{15}N$ analysis was cleaned by combustion in a muffle

oven at 450 °C for 6 h. Diatom frustules were cleaned as described by (Morales et al., 2013).

Briefly, diatom samples underwent at least two cycles of sonication in 2 % (w/v) sodium dodecyl

sulfate (SDS), followed by centrifugation, decanting, sonication in purified water, centrifugation

and decanting. Non-frustule-bound biomass was then further removed through subsequent steps

of oxidation by sulfuric acid and potassium permanganate ($KMnO_4$), 14 % perchloric acid for 1 h

at 100 °C, and 70 % perchloric acid for 1 h at 100 °C. Cleaned diatom frustules were collected on

2 µm polycarbonate filter membranes and rinsed thoroughly with purified water. Diatom samples

were then dried at 60 °C in a dedicated, low-N oven. Dissolution of the opal frustules and oxidation

of the released frustule-bound organic N was performed within one week of drying.

Approximately 2-7 mg of dried frustules were weighed into 4 mL glass vials, but samples

with lower weights were transferred to glass vials using purified water. Persulfate oxidizing

reagent (POR) solution used to dissolve diatom frustules and oxidize organic N to nitrate was

added (1.5ml per vial). POR consisted of potassium persulfate (5% w/v) and sodium hydroxide

(5% w/v). The potassium persulfate salt was pre-purified to remove residual N via multiple

dissolution and crystallization cycles.

Vials were autoclaved for 90 minutes at 121 °C. Following oxidation, sample pH was

adjusted to 4 - 7 using ultrapure HCl. Nitrate concentrations of oxidized frustule samples were

determined using the chemiluminescence method described above and the $\delta^{15}N$ of the frustule-

derived nitrate was measured using the denitrifier method (see below; Sigman *et al.*, 2001). Each

sample batch included vials with known concentrations of amino acid standards (USGS64,

USGS65, USGS40 and USGS41) and procedural blanks to assess recovery, oxidation

completeness, and blank contribution. Procedural blank size and $\delta^{15}N$ values were used for post-

correction of frustule-bound $\delta^{15}N$ values using isotope mass-balance considerations.

Consistency of the frustule-cleaning and oxidation method was assessed by routinely

including material from a commercial diatom culture of *T. weissflogii* (TW1800, Reed

Mariculture). This regularly co-processed material yielded $\delta^{15}N_{DB}$ values of 0.4 ± 0.3 ‰ (n = 10).

Additionally, to further determine the consistency of the oxidation method, diatom frustule

material from two reference sources - a commercial *Thalassiosira pseudonana* culture (TP1800,

Reed Mariculture) and a phytoplankton sample collected in Lake Murten, Switzerland (2023) -

was cleaned once in bulk, stored in sealed vials isolated from the atmosphere, and regularly co-

analyzed across runs. These samples yielded frustule-bound $\delta^{15}N$ values of -6.1 ± 0.5 ‰ (n = 9),

and 8.4 ± 0.4 ‰ (n = 8), respectively. Across all diatom culture samples, the mean procedural

blank contribution was 1.5 %, with a maximum of 5.6 %. Corresponding corrections to frustule-

bound $\delta^{15}N$ values were, on average, 0.13 ‰ with a maximum of 0.63 ‰.

Ten nanomoles of dissolved nitrate (from culture medium or from oxidized diatom

frustules) was converted to N_2O using the denitrifier method (Sigman et al., 2001). The $\delta^{15}N$ of

the resulting N_2O was measured by gas-chromatography isotope ratio mass spectrometry (GC-

IRMS; Delta V Plus, Thermo Fisher Scientific) coupled to a Gasbench used for N_2O purification.

Scale normalization was performed using nitrate reference materials with known $\delta^{15}N$: USGS34 (-

1.8 ‰), IAEA N3 (4.7 ‰) and a lab-internal standard UBN-1 (14.15 ‰). For the analysis of nitrate

samples from the ^{15}N-labelling experiment, the USGS32 standard ($\delta^{15}N$ = 180 ‰) was also

included. Analytical precision was assessed through repeated analyses of nitrate from a Deep-

Pacific-Ocean water sample (5.2 ± 0.1 ‰, n = 28).

2.3. Data analysis

All $\delta^{15}N$ values are reported relative to AIR-N_2, and $\delta^{13}C$ values relative to the Vienna Pee

Dee Belemnite (V-PDB) standard. All calculations and statistical analyses were performed in R

version 4.3.0 (R Core Team, 2023). For each batch culture, the nitrate-to-silicate uptake ratio was

determined from the slope of a linear regression between nitrate and silicate concentrations, using

only data points collected before the addition of supplemental silicate to the culture.

Assuming that ^{15}N fractionation during nitrate assimilation in the closed-system batch

cultures follows Rayleigh-type behavior/systematics, the N-isotope fractionation (i.e., isotope

effect), $\varepsilon_{biomass/NO3}$, was estimated as the slope of the linear relationship between the natural

logarithm of the N isotopic ratio of nitrate ($^{15}N/^{14}N$) and the fraction of nitrate remaining in the

medium (f) (Mariotti et al., 1981). This relationship was modelled with ordinary least squares

regression (OLS) using the *lm* function in R. The intercept of this regression approximates the

initial nitrate $\delta^{15}N$ value, i.e., when $f = 1$ ($\delta^{15}N_{f1}$). Using $\varepsilon_{biomass/NO3}$ and $\delta^{15}N_{f1}$ we calculated the

expected $\delta^{15}N$ value of biomass (i.e., the accumulated product) as a function of f, using the

Rayleigh-based equation (Mariotti et al., 1981):

$$\delta^{15} N_{pred.\,biomass} = \delta^{15} N_{f1} - \varepsilon_{biomass\,/\,NO3} \times \frac{f\,ln f}{1 - f}$$

To evaluate whether the integration of nitrate $\delta^{15}N$ values into biomass followed Rayleigh

dynamics predictably, we tested if the slope of the relationship between predicted and measured

biomass $\delta^{15}N$ values differed significantly from 0 and 1. Additionally, to assess how faithfully

biomass $\delta^{15}N$ values are recorded in frustule-bound N, we tested whether the slope of the

regression between $\delta^{15}N$ values of frustule-bound N and those of bulk biomass differed

significantly from 0 and 1. These (linear) regressions were also conducted using the *lm* function in

R. The N-isotopic offset between $\delta^{15}N$ values of frustule-bound N and biomass was calculated as

$\varepsilon_{DB/biomass} = \alpha - 1$, and $\alpha = (\delta^{15}N_{DB} + 1)/(\delta^{15}N_{biomass} + 1)$, expressed in ‰ (Coplen, 2011). We then

related $\varepsilon_{DB/biomass}$ values to the nitrate drawdown (f) to determine whether $\varepsilon_{DB/biomass}$ changed with

batch culture progression. Lastly, the slopes of these relationships were compared to $\varepsilon_{biomass/NO3}$,

but this analysis was limited to batch cultures with at least four $\varepsilon_{DB/biomass}$ values (Table 1).

All the above analyses excluded samples from the ^{15}N-labeled nitrate addition experiments.

For these experiments, the biomass $\delta^{15}N$ values after label addition were estimated by isotope mass

balance, considering f values and the $\delta^{15}N$ values of both the newly synthesized biomass (after

addition of the ^{15}N-labeled nitrate), and the pre-existing accumulated biomass (before ^{15}N-label

addition).

3. Results

3.1. Nutrient concentrations during batch cultures

Batch cultures lasted 12-15 days for *A. granulata*, 15-22 days for *F. capucina*, 7-8 days for *C. calcitrans*, and 47 days for *C. radiatus* (Table 1; Fig. S1 and S2). In 14 out of 17 batch-culture experiments, nitrate was consumed to concentrations corresponding to *f* values <0.1. Silicate-to-nitrate uptake ratios were highly variable among species: for the freshwater species they ranged from 1.5 and 2.9, for *C. calcitrans* from 0.4 and 0.7. For *C. radiatus* the highest Si:NO_3^- uptake ratios were observed, ranging between 4.1 and 4.4 (Table 1).

Table 1: Overview of initial and final nitrate concentrations, Si:NO_3^- uptake ratios, frustule N contents, C and N contents, and batch culture durations for experiments with different diatom cultures. Values are reported as mean ± SD where applicable. bSi, biogenic silica

Species	No.	Starting [NO_3^-] (uM)	Final [NO_3^-] (uM)	Si:NO_3^- uptake ratio	Frustule N content (nmol N mg^{-1} bSi)	Biomass C fraction (g g^{-1})	Biomass N fraction (g g^{-1})	Number of frustule samples	Culture duration (d)	Notes
Aulacoseira granulata	1	195	56	2.6	14.2 ± 1.6	0.19 ± 0.02	0.031 ± 0.001	3	13	
	2	196	14	2.1	12.4 ± 0.5	0.22 ± 0.04	0.035 ± 0.002	3	13	
	3	191	0	2.6	12.8 ± 0.7	0.20 ± 0.02	0.037 ± 0.006	4	14	With TES buffer
	4	196	0	1.8	15.7 ± 5.9	0.18 ± 0.02	0.035 ± 0.004	5	15	With TES buffer
	5	203	0	2.3	9.0 ± 0.5	0.16 ± 0.01	0.031 ± 0.003	6	15	[15]N-labeled nitrate addition
	6	208	1	1.9	9.9 ± 0.7	0.18 ± 0.02	0.033 ± 0.002	6	12	[15]N-labeled nitrate addition
Fragilaria capucina	1	194	45	2.2	13.2 ± 1.0	0.23 ± 0.01	0.030 ± 0.002	3	22	
	2	194	21	2.3	14.3 ± 2.6	0.26 ± 0.03	0.032 ± 0.001	3	21	
	3	185	1	2.9	9.3 ± 1.3	0.20 ± 0.03	0.028 ± 0.002	5	15	With TES buffer
	4	194	6	2.6	8.8 ± 1.2	0.19 ± 0.03	0.028 ± 0.002	6	17	With TES buffer
	5	205	2	1.5	7.1 ± 2.3	0.21 ± 0.03	0.030 ± 0.005	6	21	[15]N-labeled nitrate addition
	6	200	0	1.9	6.0 ± 1.3	0.22 ± 0.02	0.028 ± 0.003	6	21	[15]N-labeled nitrate addition
Chaetoceros calcitrans	1	512	4	0.5	17.6 ± 1.5	0.15 ± 0.03	0.028 ± 0.005	4	8	
	2	502	4	0.4	15.6 ± N/A	0.14 ± 0.06	0.025 ± 0.011	4	8	
	3	500	2	0.7	4.1 ± 0.7	0.13 ± 0.04	0.027 ± 0.007	4	7	
Coscinodiscus radiatus	1	509	27	4.1	2.2 ± 0.4	0.11 ± 0.01	0.016 ± 0.002	5	47	
	2	513	2	4.4	1.7 ± 0.3	0.12 ± 0.02	0.016 ± 0.002	5	47	

314 ### 3.2. Fractionation of nitrate N isotopes during nitrate assimilation

315   The isotope effect associated with nitrate assimilation ($\varepsilon_{biomass/NO3}$), varied substantially

316 among species (Fig. 1). The largest $\varepsilon_{biomass/NO3}$ values were found for *C. radiatus* (-13.8 ±0.2 ‰),

317 followed by *A. granulata* (-10.8 ±1.0 ‰), and *C. calcitrans* (-3.1 ±0.5 ‰), with the smallest values

318 observed for *F. capucina* (-1.9 ±0.2 ‰).

Figure 1: Nitrate ^{15}N fractionation ($\varepsilon_{biomass/NO3}$) during assimilation by different diatom species. ε values given in the legend represent species means ± SD, based on slopes of the linear regression, where the SD is derived from ε values obtained from culture replicates. Note that ε values were calculated from nitrogen isotope ratios rather than $\delta^{15}N$ values; however, for ease of interpretation, the regressions are shown on the $\delta^{15}N$ scale.

3.3. Biomass δ15N and δ13C values

Predicted biomass δ15N values – calculated using the cumulative-product equation by

(Mariotti et al., 1981) and based on the observed nitrate isotope fractionation during nitrate

assimilation – were very similar to measured biomass δ15N values (Fig. 2) with a regression slope

of 0.95 (p < 0.001 for slope ≠ 0), close to but significantly different from 1 (p = 0.026 for slope ≠

1). The explanatory power of this relationship was very high (R^2 = 0.97), indicating that predicted

biomass δ15N values explain nearly all of the observed variation in measured biomass δ15N values.

Figure 2: Comparison of predicted versus measured biomass δ15N values. Predicted biomass δ15N values were calculated using the cumulative-product equation (Mariotti et al. 1981), based on the observed 15N fractionation during nitrate assimilation and the fraction of nitrate remaining at the time of sampling.

Consistent with the species-specific $\varepsilon_{biomass/NO3}$, biomass δ15N values at high *f* values were

lowest in *A. granulata* and *C. radiatus*, around -11 to -12 ‰, while those of *F. capucina* and *C.*

calcitrans were never lower than -5 ‰ (Fig. 3). Upon (near) complete consumption of nitrate in

the medium, for all species, biomass $\delta^{15}N$ approached values close to the starting nitrate $\delta^{15}N$ value

of -1.4 ‰ (Fig. 3).

For *A. granulata*, *F. capucina* and *C. calcitrans*, biomass $\delta^{13}C$ values increased with

decreasing *f* (Fig. S3), with the magnitude of the increase being larger for species that more rapidly

consumed the nitrate in the medium (Table 1; Fig. S1). For *C. radiatus*, there was no strong

increase in biomass $\delta^{13}C$ values during the batch culture, although the values were elevated

compared to the lowest biomass $\delta^{13}C$ values in the other species.

Figure 3: $\delta^{15}N$ values of nitrate (blue circles), biomass (green diamonds) and frustule-bound N (white triangles) plotted against the fraction of nitrate remaining in the medium (f). Panels represent results from single batch cultures, organized by species (columns) and culture replicates (rows). Solid, dashed, and dotted lines represent theoretical $\delta^{15}N$ values for nitrate, accumulated biomass, and starting nitrate, respectively. For A. granulata *and* F. capucina, *culture replicates 3 and 4 were grown in medium containing TES buffer.*

3.4. Diatom frustule-bound δ^{15}N values

Linear regression of δ^{15}N$_{DB}$ values against biomass δ^{15}N values across all species shows a significant relationship (p = 0.003 for slope $\neq$ 0; Fig. 4). However, the slope of this relationship was significantly different from 1 (p < 0.001 for slope $\neq$ 1), and the explanatory power was relatively low (R^2 = 0.32). Moreover, both the significance and explanatory power of this relationship decreased when the data were analyzed at the species- or culture-specific level. This is further illustrated by increasing biomass δ^{15}N values with decreasing f, while δ^{15}N$_{DB}$ values did not show substantial change (Fig. 3).

Figure 4: Comparison between frustule-bound and biomass δ^{15}N values. The linear regression (dashed line) was determined across all species and batch cultures.

The isotopic offset between δ^{15}N$_{DB}$ values and biomass δ^{15}N values, expressed as $\varepsilon_{DB/biomass}$, was highly variable, from +10.3 to -8.4 ‰ depending on the species (Fig. 5). In nearly all cultures, $\varepsilon_{DB/biomass}$ decreased with decreasing f. The magnitude of this decrease differed among

349   species, with the strongest shifts observed for *A. granulata* and *C. radiatus*, and weaker decreases

350   in *F. capucina* and *C. calcitrans*. In *A. granulata*, $\varepsilon_{DB/biomass}$ decreased from +10 to +6 ‰ at $f \approx 0.8$

351   to around 0 ‰ when f approached zero, whereas in *C. radiatus* it decreased from +5 to +1 ‰ at f

352   = 0.7 to -8 ‰ when f approached zero. In *F. capucina* and *C. calcitrans*, $\varepsilon_{DB/biomass}$ decreased by

353   roughly 4 ‰ with f decreasing from 0.8 to 0.

Figure 5: Isotopic offset between biomass and frustule-bound N (expressed as $\varepsilon_{DB/biomass}$) progressing along the batch culture, shown as a function of the fraction of nitrate remaining (f). Points represent individual measurements, and lines connect sequential measurements from a single batch culture.

We further quantified the change in $\varepsilon_{DB/biomass}$ along the f gradient, by calculating the slope

of a linear regression for each batch culture, where the slope represents the rate of change of

$\varepsilon_{DB/biomass}$. When these slope values were plotted against the corresponding $\varepsilon_{biomass/NO3}$ values (Fig.

S4), a strong inverse relationship emerged. More precisely, cultures with less negative (i.e., higher)

$\varepsilon_{biomass/NO3}$ values exhibited a less pronounced slope, whereas those with more negative (i.e., lower)

$\varepsilon_{biomass/NO3}$ values displayed a stronger decrease in $\varepsilon_{DB/biomass}$. This indicates that the extent to which

frustule-bound N diverges from biomass $\delta^{15}N$ is closely linked to the species-specific N isotope

effect during nitrate assimilation.

C and N content (Table 1), and their ratio, could not explain variation in $\epsilon_{DB/biomass}$ within

most species, except for *A. granulata*, where lower C contents and C/N ratios were associated

with lower $\epsilon_{DB/biomass}$ values (p = 0.045 and p = 0.049, respectively).

3.5. Changes in biomass and frustule-bound $\delta^{15}N$ values in response to high-$\delta^{15}N$ nitrate

additions

In an experiment to assess the incorporation of N isotope signatures from nitrate into

biomass and frustule-bound N, we manipulated medium-nitrate $\delta^{15}N$ values when roughly half of

the initial nitrate pool had been consumed. Thereby the nitrate concentrations increased by only 1-

Figure 6: $\delta^{15}N$ values of nitrate, biomass and frustule-bound N in response to the addition of high-$\delta^{15}N$ nitrate (targeted nitrate $\delta^{15}N$ 90‰) when roughly half of total nitrate had been consumed. Dashed lines represent estimated biomass $\delta^{15}N$ values (accumulated product), calculated using an isotope mass balance approach following the addition of the nitrate $\delta^{15}N$ spike.

2 μM, and spiked nitrate $\delta^{15}N$ values upon ^{15}N-labeled nitrate addition were between 85.5 ‰ and

94.4 ‰ (Fig. 6). Both biomass and diatom frustule-bound $\delta^{15}N$ values clearly responded to the

addition of ^{15}N-labeled nitrate during batch cultures. Biomass $\delta^{15}N$ values increased in close

agreement with predictions for an accumulated product. Frustule-bound $\delta^{15}N$ values also

increased, but they remained consistently lower than those of biomass. The resulting $\varepsilon_{DB/biomass}$

values were species-specific, with *A. granulata* showing smaller offsets compared to *F. capucina*.

Upon complete nitrate consumption $\varepsilon_{DB/biomass}$ values ranged from -6.4 to -8.0‰ for *A. granulata*,

and from -23.6 to -23.8‰ for *F. capucina*. Using i) the $\delta^{15}N$ values of samples collected before

and shortly after addition of ^{15}N-labeled nitrate, and ii) $\delta^{15}N$ values after the spike addition, the

relationship between biomass and frustule-bound $\delta^{15}N$ values were assessed. Initially, the biomass-

vs.-frustule-bound $\delta^{15}N$ slopes were relatively shallow for both species, indicating that biomass

$\delta^{15}N$ values increased more strongly than frustule-bound $\delta^{15}N$ values (Fig. 7). After these initial

samples, the slope of this relationship approached 1 for *A. granulata* (0.93, p = 0.55 for slope ≠1),

whereas for *F. capucina* the slope remained significantly lower than 1 (0.66, p = 0.04 for slope

≠1). For *A. granulata*, the first after-spike samples were taken at roughly 3.6 and 5.5h after the

^{15}N-nitrate addition, corresponding to a fractional nitrate consumption of 0.07*f* (Fig. S2). By

contrast, for *F. capucina*, the first two samples were taken after 21.6 and 22.8h after spiking,

corresponding to a fractional consumption of 0.19*f*. Throughout the experiment, empty frustule

counts relative to frustules containing biomass were consistently low (< 3 %; Table S1).

Figure 7: Frustule-bound δ15N values plotted against biomass δ15N values of samples taken in the experiment with 15N-labeled nitrate. Slopes (dashed lines) were determined on a species-specific basis and are distinguished by color. Initial slopes were calculated using the samples collected before the addition of 15N-labeled nitrate and the first two samples collected afterward. The 'final slopes' were calculated using only data from samples collected after the addition of 15N-labeled nitrate. The solid line represents a slope of 1 with an intercept 0. Dotted lines represent a slope of 1 with intercepts sequentially offset by 10 ‰.

4. Discussion

Meaningful interpretation of $\delta^{15}N$ values of diatom frustule-bound organic N ($\delta^{15}N_{DB}$) stored in sedimentary records relies on proper understanding of how N stable isotopes from the source nitrate pool are integrated into diatom biomass and subsequently into frustule-bound organic N. Here we present new data from mono-specific diatom cultures that illustrates the relationship among $\delta^{15}N$ values of nitrate, biomass, and frustule-bound organic N in two freshwater and two marine diatom species. The results from these cultures show that $\delta^{15}N_{DB}$ values can differ substantially from diatom biomass $\delta^{15}N$ values, and that these differences depend on species as well as on the progression of the batch culture. Moreover, by experimentally manipulating nitrate $\delta^{15}N$ values during batch culture incubations, we found that integration of the nitrate $\delta^{15}N$ into biomass occurred more rapidly than into frustule-bound organic N, and the extent of this difference was species-dependent. However, these manipulations could only explain a small part of the discrepancies between biomass and frustule-bound $\delta^{15}N$ values observed in the natural-abundance experiments, suggesting that the observed offsets are primarily driven by internal biochemical ^{15}N fractionation.

4.1. Incorporation of nitrate $\delta^{15}N$ signatures into biomass and frustule-bound N

Overall, we find that the ^{15}N fractionation upon assimilation of nitrate ($\varepsilon_{biomass/NO3}$) can vary substantially among species, but it is relatively stable within species under the conditions tested (Fig. 1), consistent with previous studies (Needoba et al., 2003). Moreover, measured biomass $\delta^{15}N$ values agreed well with predicted biomass $\delta^{15}N$ values calculated as accumulated product based on $\varepsilon_{biomass/NO3}$ (Fig. 2 & 3). Thus, ^{15}N fractionation of nitrate and subsequent incorporation into biomass followed predictable Rayleigh-type fractionation behavior in our batch cultures, as would be expected based on past research (Waser et al., 1998).

Previous studies using field-collected samples from the natural environment showed that,

although biomass and frustule-bound $\delta^{15}N$ values may be correlated with one another, neither was

correlated with ambient nitrate $\delta^{15}N$ values - likely due to substantial uptake of ammonium with

distinct $\delta^{15}N$ signatures (Morales et al., 2014; Robinson et al., 2020). In controlled batch cultures

where nitrate is the only inorganic N source, both biomass $\delta^{15}N$ and $\delta^{15}N_{DB}$ would therefore be

expected to follow the isotopic evolution of the nitrate pool, with biomass $\delta^{15}N$ increasing

according to $\varepsilon_{biomass/NO3}$ and $\delta^{15}N_{DB}$ anticipated to also track these changes. Indeed, across species

and culture replicates, we find a significant relationship between $\delta^{15}N$ values of frustule-bound N

and those of biomass (Fig. 4). However, it is important to note that the slope of this relationship is

significantly different from 1, and when the same comparison is made within species or within

culture replicates, the relationship disappears completely. Moreover, the unexpectedly low

explanatory power of this relationship ($R^2 = 0.32$) indicates that only about one third of the

observed variation in $\delta^{15}N_{DB}$ values is explained by biomass $\delta^{15}N$ values. Thus, the increase in

biomass $\delta^{15}N$ values induced by progressive nitrate consumption was not strictly mirrored by a

corresponding increase in $\delta^{15}N_{DB}$. This implies that $\delta^{15}N_{DB}$ does not directly represent the $\delta^{15}N$ of

an accumulated product in batch cultures, and therefore it cannot be assumed to straightforwardly

reflect biomass $\delta^{15}N$ values.

4.2. The offset between frustule-bound and biomass $\delta^{15}N$ changes with batch culture

progression

We observed that the isotopic offset between frustule-bound and biomass $\delta^{15}N$ values

($\varepsilon_{DB/biomass}$) was not constant but decreased with batch culture progression (Fig. 5). This pattern

primarily reflects the species-specific increase in biomass $\delta^{15}N$ with fractional nitrate

consumption, while $\delta^{15}N_{DB}$ values remained comparatively stable (Fig. 3). Such relatively stable

$\delta^{15}N_{DB}$ values could imply that the organic N preserved in frustules originates from an N pool that

remains unaltered throughout the batch culture. It has been proposed that the synthesis of

frustule-bound organic N molecules can occur well before frustule formation, which may explain

the low $\delta^{15}N_{DB}$ values observed in *Cheatoceros* resting spores relative to frustules from vegetative

cells (Dove et al., 2025). In batch culture experiments with natural-abundance nitrate $\delta^{15}N$ levels,

it is difficult to assess whether organic N in newly formed frustules undergoes variable

biochemical ^{15}N fractionation or isotopically reflects early culture stages, which is further

discussed below. Irrespective of the underlying mechanism, the observed change in $\varepsilon_{DB/biomass}$

provides useful context for interpreting batch culture results reported in previous studies.

Results from Horn et al. (2011) suggest a positive relationship between $\varepsilon_{DB/biomass}$ and

$\varepsilon_{biomass/NO3}$. Similarly, our results also show that $\delta^{15}N_{DB}$ values can be strongly offset from biomass

$\delta^{15}N$ values in species with large $\varepsilon_{biomass/NO3}$ values, whereas the offsets are generally small in

species with small $\varepsilon_{biomass/NO3}$ values (Fig. 1 & 5). However, it is important to consider that

$\varepsilon_{DB/biomass}$ was not constant in our cultures (Fig. 3 & 5), and the magnitude of change in $\varepsilon_{DB/biomass}$

during batch culture progression appears to be proportional to $\varepsilon_{biomass/NO3}$ (Fig. S4). This suggests

that processes driving variable $\varepsilon_{DB/biomass}$ values could be related to physiological mechanisms

involved in nitrogen assimilation and/or intracellular metabolism.

As silicification likely controls the frustule-bound organic compound matrix (Bridoux et

al., 2012), and their synthesis timing (Kotzsch et al., 2017), silicification rates may affect

incorporation of $\delta^{15}N$ values into frustule-bound-N. The species used in our study represent a large

range of silicification levels, with *C. radiatus* exhibiting $Si:NO_3^-$ uptake ratios greater than 4,

compared to 0.4 to 0.7 for *C. calcitrans*, and 1.5 to 2.9 for the freshwater species (Table 1). In line

with previous suggestions (Robinson et al., 2025), the heavily silicified *C. radiatus* was the only

species whose $\delta^{15}N_{DB}$ values were consistently lower than biomass $\delta^{15}N$ values (Fig. 4 & 5).

However, even though *A. granulata* and *F. capucina* had similar $Si:NO_3^-$ uptake ratios (Table 1),

their observed $\varepsilon_{DB/biomass}$ values differ substantially (Fig. 5). Moreover, $\varepsilon_{DB/biomass}$ values were not

consistent within single batch cultures, shifting from as high as +10 ‰ to around 0 ‰ in *A.*

granulata, and from nearly +5 ‰ to -8 ‰ in *C. radiatus*. Depending on species, the silicification

rates may change during batch culture progression as Si concentrations decrease (Kudo, 2003).

However, we observed substantial shifts in $\varepsilon_{DB/biomass}$ even under conditions where Si

concentrations changed only modestly (Fig. 5; Fig. S1 & S2), and there was no clear relation with

the replenishments of Si during the batch culture. Thus, our batch culture results do not support

the proposed relationship between $Si:NO_3^-$ uptake ratios and $\varepsilon_{DB/biomass}$ suggested by Robinson et

al. (2025).

In previous studies, it was difficult to gauge whether $\varepsilon_{DB/biomass}$ values were stable or

variable throughout batch cultures, since cultures were often terminated before complete nitrate

consumption (Horn et al., 2011), or only a single sample was collected for $\delta^{15}N_{DB}$ analyses (Jones

et al., 2022; Robinson et al., 2025). Consequently, interpretations from previous culture studies

may be confounded if $\varepsilon_{DB/biomass}$ values cannot be assumed constant over the course of the batch

culture. Importantly, our results imply that $\delta^{15}N_{DB}$ values do not clearly follow the dynamics of an

accumulating product in a Rayleigh-type system. The observed changes in $\varepsilon_{DB/biomass}$ with batch

culture progression are likely driven by (1) changes in ^{15}N fractionation associated with shifts in

biochemical reactions during N metabolism, and/or (2) differences in the timing of N isotope

integration between $\delta^{15}N_{DB}$ and biomass $\delta^{15}N$ values through the use of intracellular N-reservoirs

established early in the batch culture. To further explore the importance of the latter process, we

conducted separate batch culture experiments in which ^{15}N-labeled nitrate was added when

roughly half of the initial nitrate remained, and traced its incorporation into biomass and frustule-

bound N.

4.3. Slower nitrate isotope incorporation into the frustule-bound N pool does not explain $\varepsilon_{DB/biomass}$ variation

The relatively constant $\delta^{15}N_{DB}$ values observed during batch culture progression (Fig. 3)

could suggest that $\varepsilon_{DB/biomass}$ is mostly driven by the utilization of an isotopically stable N reservoir

for the synthesis of frustule-bound organic N. To test this hypothesis, we conducted an experiment

where nitrate $\delta^{15}N$ values were manipulated during the batch culture. In this experiment, biomass

$\delta^{15}N$ values increased along the predicted Rayleigh path, whereas $\delta^{15}N_{DB}$ values also increased

but remained consistently lower than biomass $\delta^{15}N$ values (Fig. 6; Fig. 7). One possible

explanation for this offset could be a high or variable number of empty frustules relative to biomass

-filled frustules; however, the abundance of empty frustules was consistently low (Table S1).

After addition of ^{15}N-labeled nitrate, the distinctly lower $\delta^{15}N$ values of frustule-bound N

compared to biomass indicate that a portion of the organic N incorporated into frustules originates

from before the $\delta^{15}N$ manipulation. Moreover, the magnitude of this effect appears to be species-

specific: frustule-bound $\delta^{15}N$ values were substantially lower relative to biomass in *F. capucina*

(-23.6 ‰) compared to *A. granulata* (-7.2 ‰) upon complete nitrate consumption. Additionally,

the slope between biomass $\delta^{15}N$ and $\delta^{15}N_{DB}$ never reached 1 during the experiment time for *F.*

capucina, whereas for *A. granulata* this slope approached 1 after only a few hours. Therefore,

assimilation of ^{15}N-labeled nitrate affects biomass $\delta^{15}N$ values essentially instantaneously, while

the pool of organic N used for frustule-bound organic N synthesis requires time to be replaced

with newly assimilated, ^{15}N-enriched N. An alternative explanation is that no new frustules were

synthesized during the period when $\delta^{15}N_{DB}$ values remained stable despite large change in biomass

$\delta^{15}N$ values; however, at least in *F. capucina* some new frustules must have formed during this

time, as indicated by the observed Si consumption (Fig. S2). Thus, our results imply that

frustule-bound organic N is derived from an intracellular N reservoir that integrates newly

assimilated nitrate more slowly, leading to asynchronous incorporation of $\delta^{15}N$ signals into

frustule-bound N compared to bulk biomass.

Diatom-internal N reservoirs relevant to $\delta^{15}N_{DB}$ values could take several forms:

intracellular nitrate (Stief et al., 2022); monomers such as amino acids (Admiraal et al., 1986);

N/C-storage compounds including urea and creatine-P (Allen et al., 2006; Parker et al., 2008); or

the organic molecules (proteins and polyamines) involved in silicate precipitating prior to their

incorporation into the frustule (Kotzsch et al., 2017; Tesson et al., 2017). Stored nitrate originating

from early in the culture progression is unlikely to influence $\delta^{15}N_{DB}$ values independently of

biomass, since nitrate assimilation into organic compounds is not expected to occur via a

temporally or spatially distinct pathway dedicated specifically to frustule-bound organic N

synthesis.

Once assimilated into amino acids – the building blocks of proteins and polyamines,

including those integrated into diatom frustules – N can be stored in various organic forms. In

higher plants, amino acid turnover times can be less than one hour (Racusen and Foote, 1960), and

in diatoms, substantial ^{15}N enrichment in amino acids has been observed within one hour after ^{15}

N-label addition (Smith et al., 2019). These rapid N turnover rates imply that amino acids are

unlikely to retain isotopically distinct signatures over extended periods. Therefore, it is improbable

that metabolically active amino acids constitute the source of delayed or asynchronous

incorporation of ^{15}N-labeled nitrate into frustule-bound organic N compared to biomass.

Many products of the diatom urea cycle can act as N-reservoirs, including urea and

creatine-P (Allen et al., 2006; Parker et al., 2008). The consequences for frustule-bound $\delta^{15}N$

values of these N-reservoirs depend on the extent to which such pools accumulate and are

subsequently mobilized, processes that presumably vary with species identity and growth

conditions. Assuming that utilization of these organic N-reservoirs occurs mainly under N-limiting

conditions, it is unlikely that they were a major source of N for frustule-bound organic N synthesis,

because N-limitation only developed toward the very end of the batch culture.

Pools of certain frustule-bound organic compounds may be established prior to frustule

synthesis (Kotzsch et al., 2017; Tesson et al., 2017). If substantial amounts of these organic

compounds are synthesized before frustule formation, a pool-size- and synthesis-rate-dependent

amount of time would be required for them to be replaced with newly assimilated ^{15}N-labeled N,

thereby affecting the integration of nitrate $\delta^{15}N$ values into the frustule-bound N pool.

Overall, the results from the ^{15}N-labeled nitrate addition experiments (Fig. 6 & 7) imply

that the organic N incorporated into frustules is derived from an organic N pool that is relatively

quickly replaced by newly assimilated N. However, this time-window is clearly species-

dependent, as slower-growing species required longer periods to replace these reservoirs with

^{15}N-enriched N (Table 1, Fig. 7). Therefore, the incorporation of nitrate $\delta^{15}N$ values into frustule-

bound $\delta^{15}N$ will lag behind their incorporation into biomass $\delta^{15}N$ by a species-specific amount of

time. This consideration is important for future batch culture experiments, as $\delta^{15}N_{DB}$ values cannot

be assumed to integrate over the exact same time window as biomass $\delta^{15}N$ values.

With the slower integration of nitrate $\delta^{15}N$ values into frustule-bound N compared to

biomass, $\delta^{15}N_{DB}$ values would still be expected to reflect the increase in biomass $\delta^{15}N$ values in

batch cultures – albeit with a temporal or f-related offset. Yet, in our batch cultures with natural-

abundance nitrate $\delta^{15}N$ values, $\delta^{15}N_{DB}$ values showed hardly any increase at all (Fig. 3). Thus, to

explain the observed changes in $\varepsilon_{DB/biomass}$ values (Fig. 5), internal ^{15}N fractionation must have

varied with batch culture progression, which will be discussed next.

4.4. Potential biochemical ^{15}N fractionation effects on frustule-bound $\delta^{15}N$ values

Changes in growth conditions are an inherent feature of batch cultures, as nutrient

concentrations decrease while cellular demand for nutrients (including inorganic C) increases with

rising cell density. Increased cell density also causes self-shading, reducing light intensity in the

culture. We found that biomass $\delta^{13}C$ values progressively increased (or remained consistently

elevated in the case of *C. radiatus*) (Fig. S3), which implies (1) a higher degree of inorganic C

consumption (Degens et al., 1968; Deuser, 1970; Fry, 1996; Burkhardt et al., 1999), (2) induction

of a carbon-concentration mechanism (Sharkey and Berry, 1985), and/or (3) enhanced C fixation

via phosphoenolpyruvate carboxylase in response to increased ammonium assimilation (Guy et

al., 1989). Thus, the observed biomass $\delta^{13}C$ patterns illustrate that the progressive changes in

growth conditions that inherently occur during batch culture progression affect diatom physiology.

Overall, the initially higher, and subsequently decreasing, frustule-bound $\delta^{15}N$ values relative to

biomass (Fig. 5) could be a result of growth-condition-driven changes in (1) the frustule-bound

organic compound matrix, (2) ^{15}N-fractionating reactions during frustule-bound organic

compound synthesis, and/or (3) ^{15}N-fractionating reactions within amino acid synthesis pathways.

In the following, we address potential mechanisms by which $\varepsilon_{DB/biomass}$ may – or may not – have

changed.

Regarding (1): If the frustule-bound organic compound matrix changed during the batch

cultures, associated with a shift from N-rich precursors with high $\delta^{15}N$ values to precursors with

lower $\delta^{15}N$ values, this would be reflected in $\delta^{15}N_{DB}$. However, it is currently unknown whether

573 the composition of the frustule-bound organic matrix is plastically regulated within species (i.e.,

574 whether it varies within species in response to growth conditions). Yet, because the frustule-bound

575 polyamine composition is highly species-specific (Bridoux and Ingalls, 2010; Bridoux et al.,

576 2012), it is more likely that the frustule-bound organic compound matrix is governed primarily by

577 diatom phylogeny rather than by growth conditions.

578   Regarding (2): For proteins, ribosomal biosynthesis from amino acids is generally assumed

579 not to introduce isotopic discrimination (Werner and Schmidt, 2002). However, in higher plants,

580 polyamine $\delta^{15}N$ values have been found to be substantially higher than those of their amino acid

581 precursors, suggesting considerable potential for ^{15}N fractionation during polyamine biosynthesis

582 (Yoneyama et al., 1998). Moreover, in diatoms, methylation of amine groups in frustule-bound

583 silaffins and polyamines has also been documented (Sumper and Brunner, 2008; Kröger and

584 Poulsen, 2008; Bridoux and Ingalls, 2010), and since such reactions involve the breaking of N-H

585 and forming of N-C bonds, they may introduce ^{15}N fractionation. This isotopic fractionation could

586 be expressed if methylation occurs at branching points within the synthesis pathway, and/or if the

587 substrate-product reaction is incomplete. Increased methylation has been shown to enhance the

588 ability of polyamines to precipitate silicic acid (Robinson et al., 2007), and the degree of

589 methylation varies among species (Bridoux et al., 2012). However, whether changes in the degree

590 of methylation of frustule-bound organic-N can occur *within* a species, thereby producing variable

591 ^{15}N fractionation and contributing to the observed changes in $\varepsilon_{DB/biomass}$, remains unclear.

592   Regarding (3): To identify potential ^{15}N fractionation processes prior to the synthesis of

593 frustule-bound organic compounds, it is important to understand how these molecules are

594 synthesized, and from which precursors. Polyamines are synthesized from a putrescine or ornithine

595 backbone and from polypropylene-imine (PPI) chains of species-dependent lengths, which are

ultimately derived from methionine synthesized from aspartate (Bridoux and Ingalls, 2010; Lin et

al., 2024). Thus, for polyamines composed of more than two PPI units, methionine represents the

dominant source of N. By comparison, frustule-bound proteins such as silaffins and silacidins are

rich in serine (Poulsen et al., 2003; Richthammer et al., 2011), which is produced primarily during

photorespiration, whereas other frustule-associated proteins have been shown to be enriched in

glutamine and asparagine (Kotzsch et al., 2017), amino acids more closely related to N

assimilation. Assuming that metabolic fluxes influence the extent of isotopic fractionation

(Tcherkez, 2010), diatom metabolism likely plays an important role in shaping amino acid $\delta^{15}N$

values (Fig. 8) and, consequently, the $\delta^{15}N$ of frustule-bound organic N compounds.

Understanding the variation among amino acid $\delta^{15}N$ values and the metabolic processes

influencing them may therefore provide a comprehensive framework for identifying potential

biochemical controls on frustule-bound $\delta^{15}N$.

Transamination and deamination reactions in the photorespiration cycle lead to relatively

low $\delta^{15}N$ values of serine in photosynthetic organisms (Fig. 8) (Tcherkez and Hodges, 2008;

Gauthier et al., 2013; McCarthy et al., 2013; McMahon and McCarthy, 2016; Takizawa et al.,

2017). Although photorespiration strongly impacts the $\delta^{15}N$ values of amino acids relevant for

frustule-bound protein synthesis (Fig. 8) (Tcherkez and Hodges, 2008; Tcherkez, 2010), variation

in photorespiratory flux in diatoms is likely minimized by carbon-concentrating mechanisms

(Clement et al., 2016) and the increased specificity of the RubisCO enzyme for CO_2 compared to

O_2 (i.e., which reduces oxygenation reactions and thereby suppresses photorespiration; Tortell,

2000; Kroth et al., 2008). However, photorespiration in diatoms can be induced under high-light

conditions (Parker et al., 2004), indicating that variable photorespiratory flux is possible, and could

therefore affect the $\delta^{15}N$ values of amino acids used in the synthesis of frustule-bound organic

compounds.

Figure 8: Schematic, simplified overview of the interaction between diatom C and N metabolism. The main areas of interaction are N-assimilation (pink), photorespiration (blue), and the urea cycle (purple). Reactions involving N-containing organic compounds are annotated with ε and a sign prefix, where -ε indicates a reaction resulting in lower product δ15N values, and +ε indicates the opposite. In cases where the 15N fractionation is unknown, 'ε?' is shown. In the figure, distinctions are made between GS-GOGAT located in the plastid (GSII-GOGAT) involved in N-assimilation and GS-GOGAT located in the mitochondrion (GSIII-GOGAT) involved in ammonium recycling. Abbreviations: 2-OG, 2-oxoglutarate; 3PGA, 3-phosphoglycerate; Arg, arginine; Asp, aspartate; CBB cycle, Calvin-Benson-Bassham cycle; Gln, glutamine; Glu, glutamate; GOGAT, glutamate synthase; GS, glutamine synthetase; Gly, glycine; Met, methionine; Orn, ornithine; OAA, oxaloacetate; RuBP, ribulose bisphosphate; TCA cycle, tricarboxylic acid cycle; TP, triose-phosphate. Information compiled from multiple publications (Werner & Schmidt, 2002; Kroth et al., 2008; Tcherkez & Hodges, 2008; Smith et al., 2019).

Methionine δ^{15}N values in photosynthetic organisms are typically substantially lower than

those of glutamate (Ishikawa et al., 2018). Considering that aspartate δ^{15}N values are generally

higher than those of glutamate (Tcherkez and Hodges, 2008), the relatively low methionine δ^{15}N

values imply strong ^{15}N fractionation during its synthesis or utilization. Diatoms possess a

cobalamin (vitamin B_{12})-dependent methionine synthase enzyme, MetH, and certain species also

contain a cobalamin independent form, MetE (Bertrand et al., 2012). The latter is considerably

less efficient than the cobalamin-dependent enzyme. Thus, depending on whether MetH or MetE

is used, substantial ^{15}N fractionation may occur during methionine synthesis and processing,

reflecting vitamin B_{12} availability in the medium. Our batch cultures contained cobalamin

concentrations substantially higher than those in the F/2 medium used in previous studies (Horn et

al., 2011; Robinson et al., 2025), making it unlikely that vitamin B_{12} levels were low enough to

influence methionine $\delta^{15}N$ values. Nevertheless, the oceanographic relevance of vitamin B12 and

its potential effect on methionine synthesis should be carefully considered in $\delta^{15}N_{DB}$ related

research.

An important source of N in the urea cycle is carbamoyl phosphate, formed from

bicarbonate and ammonium (Fig. 8). Transcripts of the carbamoyl phosphate synthase gene

(unCPS) are more abundant under high-light compared to low-light conditions (Bender et al.,

2012), indicating sensitivity of N metabolism to light. Moreover, unCPS transcript abundance

shown in Bender et al. (2012) correlates with transcripts for glutamine synthesis in both the

chloroplast and mitochondrion, linking N-assimilation to the urea cycle (Fig. 8). Unlike higher

plants, in which ammonium released during photorespiration and amino acid catabolism is

predominantly re-assimilated in the chloroplast, in diatoms this ammonium can also be re-

assimilated via carbamoyl phosphate synthesis and subsequently processed through the urea cycle

(Allen et al., 2006, 2011; Parker et al., 2008). Assuming that deamination reactions releasing

ammonium impart isotopic fractionation (Tcherkez and Hodges, 2008; Takizawa et al., 2017), this

pathway could shuttle N with metabolism-imparted distinct $\delta^{15}N$ signatures into the urea-cycle

products, which can then be redistributed to other amino acids.

Another notable result is the observed relationship between $\varepsilon_{biomass/NO3}$ values and the

temporal change in $\varepsilon_{DB/biomass}$ over the course of batch culture progression (Fig. S4). During nitrate

assimilation, nitrate is converted to nitrite and subsequently to ammonium. The conversion of

nitrate to nitrite by nitrate reductase (NR) is typically the limiting step in this sequence. The

magnitude of the associated ^{15}N fractionation, expressed as $\varepsilon_{biomass/NO3}$, depends on the balance

between nitrate influx and efflux across the cell membrane (Granger et al., 2010; Chen et al., 2024).

Because $\varepsilon_{biomass/NO3}$ is largely governed by transport/diffusion and reduction kinetics, it is unlikely

to directly cause the variation observed in $\varepsilon_{DB/biomass}$. Instead, the variable relationship between the

two likely reflects species-specific differences in N metabolism and the degree of internal N-

isotope fractionation during biosynthetic processing of N. This interpretation is supported by the

relation between amino acid δ^{15}N values and N-assimilation rates in higher plants (Tcherkez,

2010), and the connection between N-assimilation pathways and the urea cycle in diatoms (Bender

et al., 2012).

Ultimately, based on our experiments alone, the specific mechanism underlying changes

in $\varepsilon_{DB/biomass}$ could not be determined. Nevertheless, our results strongly suggest that frustule-bound

δ^{15}N values are sensitive to changes in growth conditions (such as light intensity and inorganic C

availability) that influence metabolically driven ^{15}N fractionation during intracellular processing

of organic N. We thus hypothesize that alterations in metabolic fluxes within the N-assimilation,

photorespiration, and urea-cycle pathways may play particularly important roles in determining

^{15}N fractionation of amino acids used for frustule-bound organic compound synthesis.

4.5. Implications for the diatom frustule-bound nitrogen isotope paleo-proxy

The observed slower integration of nitrate δ^{15}N values into frustule-bound organic N

compared to bulk biomass is likely not important for paleorecords, because the timescale of this

lag - on the order of a few hours to at most a few days - is negligible relative to the multiple-year

to millennial integration represented in sedimentary $\delta^{15}N_{DB}$. More importantly, for *A. granulata*,

we showed that after this brief lag period, $\delta^{15}N_{DB}$ values follow those of biomass relatively closely

(Fig. 7), confirming that $\delta^{15}N_{DB}$ values can, in principle, faithfully track changes in nitrate $\delta^{15}N$

across a wide range of values.

Although this short-term time lag is negligible for paleo-applications, our results also show

that $\delta^{15}N_{DB}$ may diverge from biomass $\delta^{15}N$ for reasons other than timing alone. Historically,

$\delta^{15}N_{DB}$ values have been interpreted as direct reflections of biomass $\delta^{15}N$ values. However, the

present work reveals substantial ambiguity in how closely $\delta^{15}N_{DB}$ values track biomass $\delta^{15}N$,

consistent with observations from other diatom culture studies (Horn et al., 2011; Jones et al.,

2022; Robinson et al., 2025). We hypothesize that species-specific and growth-condition-

dependent interactions between diatom C and N metabolism and the associated ^{15}N fractionation

can lead to variability in $\delta^{15}N$ values of frustule-bound organic N that is not directly predictable

from biomass $\delta^{15}N$ values alone. Whether such interactions also impact frustule-bound $\delta^{15}N$ values

under natural environmental conditions remains to be determined.

In sedimentary records from the Southern Ocean, diatom frustule-bound $\delta^{15}N$ has been

shown to vary on glacial/interglacial timescales, with $\delta^{15}N_{DB}$ values up to 4 ‰ higher during

glacial periods compared to interglacials (Robinson et al., 2004; Studer et al., 2015).

Glacial/interglacial cycles are characterized by large changes in atmospheric CO_2 concentrations

and marine biological productivity, with both being lower in the Antarctic Ocean during glacial

periods (Mortlock et al., 1991; Petit et al., 1999; Studer et al., 2015). The elevated glacial $\delta^{15}N_{DB}$

values have generally been attributed to reduced nitrate supply, leading to increased fractional

nitrate consumption and thus higher $\delta^{15}N$ values incorporated into frustule-bound organic N

(Robinson et al., 2004; Studer et al., 2015). In contrast to the natural environment, the batch culture

experiments in our study were conducted under near-unlimited nutrient conditions until nitrate

concentrations approached complete consumption. Despite this, $\varepsilon_{DB/biomass}$ was highly variable,

suggesting that $\varepsilon_{DB/biomass}$ was not directly driven by external nutrient status, but rather by other

factors, such as light conditions, inorganic C supply, and/or cell density. A previous study

highlighted that the isotope effect of nitrate assimilation in the Antarctic Ocean is insensitive to

mixed-layer depth and thus, light availability (Fripiat et al., 2019). Moreover, improved light

conditions are unlikely to contribute to the observed glacial increase in frustule-bound $\delta^{15}N$ in

paleorecords, because light-driven increases in nitrate consumption would have been associated

with higher productivity, opposite to what is observed in the Antarctic Ocean during glacial periods

(Studer et al., 2015; Ai et al., 2020).

With respect to atmospheric CO_2, it is important to consider whether diatom growth could

have been C-limited during glacial periods, potentially inducing changes in diatom metabolic

pathways relevant to internal biochemical ^{15}N fractionation. Although phytoplankton growth is

often assumed not to be limited by inorganic C in the natural environment, diatoms can experience

CO_2 limitation during dense blooms, especially in air-equilibrated surface ocean waters during

glacial periods when atmospheric CO_2 concentrations were ~200 ppm (Riebesell et al., 1993).

Since our results suggest that the generation of frustule-bound $\delta^{15}N$ signals can be sensitive to

growth conditions, this aspect should be carefully considered when interpreting sedimentary

$\delta^{15}N_{DB}$ records.

The continued development of the $\delta^{15}N_{DB}$ proxy should include systematic characterization

of how diatom metabolic processes affect frustule-bound $\delta^{15}N$ values. As discussed above,

compound-specific ^{15}N-isotope analysis of biomass-derived amino acids will provide vital

information linking diatom metabolism to the isotopic fractionation mechanisms that shape

$\delta^{15}N_{DB}$. Additionally, $\delta^{15}N$ measurements of amino acids derived from frustule-bound organic N

could be highly informative for distinguishing between shifts in source-N $\delta^{15}N$ values (where all

amino acid $\delta^{15}N$ values change similarly) and metabolically driven biochemical ^{15}N fractionation

effects, where amino acid $\delta^{15}N$ values change relative to one another. Such approaches have

already been applied to mollusk shells from sediment archives (Vokhshoori et al., 2023), and

extending them to diatom frustules would provide significant new insights for refining the $\delta^{15}N_{DB}$

paleo-proxy.

5. Conclusion

The results presented here demonstrate that diatom biomass and frustule-bound $\delta^{15}N$ values

can become decoupled in batch cultures, and that this decoupling cannot be explained by

differences in isotopic integration timing. We show that $\delta^{15}N_{DB}$ is indeed driven by $\delta^{15}N$ values of

assimilated nitrate in ^{15}N-labeled nitrate experiments; however, under natural-abundance

conditions, ^{15}N fractionation during intracellular processing of organic N substantially influences

$\delta^{15}N_{DB}$. We hypothesize that this offset arises from growth-condition-dependent shifts in metabolic

fluxes, which can influence ^{15}N fractionation during amino acid synthesis and the formation of

frustule-bound organic compounds. Future ground-truthing efforts should therefore aim to

quantify the extent and drivers of decoupling between bulk biomass and frustule-bound $\delta^{15}N$ under

controlled varying environmental conditions. Such experiments would improve our understanding

of how $\delta^{15}N_{DB}$ values respond to physiological variability during diatom growth, ultimately

enhancing the reliability of the $\delta^{15}N_{DB}$ paleo-proxy for reconstructing N cycling and environmental

change in aquatic systems over long time scales.

Author contributions

Jochem Baan: Conceptualization, Data curation, Investigation, Formal analysis, Methodology, Visualization, Writing – original draft, Writing – review and editing; **Moritz F. Lehmann**: Conceptualization, Resources, Supervision, Writing – review and editing; **Anja S. Studer**: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Writing – review and editing.

Data availability

The resulting dataset from this study is available on the *Zenodo* repository (https://doi.org/10.5281/zenodo.16994169)

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

Funding for this study was provided by the Swiss National Science Foundation (Grant 200021_200766 to A.S.S.) and by resources from the University of Basel. We would like to thank Marta Reyes (Eawag) for providing freshwater diatom cultures and help with diatom culturing techniques; Thomas Kuhn (University of Basel) for help with Flash-EA IRMS analyses of biomass samples and general laboratory support; Klaus Schläppi (University of Basel) for providing access to SANYO climate chambers; Ansgar Kahmen (University of Basel), Günter Hoch (University of Basel), and Georges Grün (University of Basel) for access to, and setting up phytotron growth chambers; Gregory de Souza (ETH Zürich) for help with silicate concentration analysis; and

760 Guillaume Tcherkez (University of Angers), Roland Werner (ETH Zürich) and Peter Kroth

761 (University of Konstanz) for helpful discussions of biochemical [15]N fractionation and diatom

762 metabolism.

Appendix A. Supplementary material

Supplementary Table S1: Percentage of empty frustules for each sample collected during

batch cultures spiked with [15]N-labeled nitrate.

Supplementary Figure S1: Silicate concentration, fraction of nitrate remaining in the

medium (*f*) and pH over time during batch culture progression.

Supplementary Figure S2: Silicate concentration, fraction of nitrate remaining in the

medium (*f*) and pH over time during batch culture progression, where [15]N-labeled nitrate was

added after approximately half of the initial nitrate pool had been consumed.

Supplementary Figure S3: Biomass $\delta^{13}C$ values plotted against *f*.

Supplementary Figure S4: Slopes of the linear regression between $\varepsilon_{DB/biomass}$ and fraction

of nitrate remaining in the medium (*f*) plotted against $\varepsilon_{biomass/NO3}$ values.

References

Admiraal W., Peletier H., Laane R. W. P. M., 1986. Nitrogen metabolism of marine planktonic diatoms; excretion, assimilation and cellular pools of free amino acids in seven species with different cell size. J. Exp. Mar. Biol. Ecol. 98, 241–263.

Ai X. E., Studer A. S., Sigman D. M., Martínez-García A., Fripiat F., Thöle L. M., Michel E., Gottschalk J., Arnold L., Moretti S., Schmitt M., Oleynik S., Jaccard S. L., Haug G. H., 2020. Southern Ocean upwelling, Earth's obliquity, and glacial-interglacial atmospheric CO_2 change. Science 370, 1348–1352.

Allen A. E., Dupont C. L., Oborník M., Horák A., Nunes-Nesi A., McCrow J. P., Zheng H., Johnson D. A., Hu H., Fernie A. R., Bowler C., 2011. Evolution and metabolic significance of the urea cycle in photosynthetic diatoms. Nature 473, 203–207.

Allen A. E., Vardi A., Bowler C., 2006. An ecological and evolutionary context for integrated nitrogen metabolism and related signaling pathways in marine diatoms. Curr. Opin. Plant Biol. 9, 264–273.

Bender S. J., Parker M. S., Armbrust E. V., 2012. Coupled Effects of Light and Nitrogen Source on the Urea Cycle and Nitrogen Metabolism over a Diel Cycle in the Marine Diatom *Thalassiosira pseudonana*. Protist 163, 232–251.

Berges J. A., Franklin D. J., Harrison P. J., 2001. Evolution of an Artificial Seawater Medium: Improvements in Enriched Seawater, Artificial Water Over the Last Two Decades. J. Phycol. 37, 1138–1145.

Bertrand E. M., Allen A. E., Dupont C. L., Norden-Krichmar T. M., Bai J., Valas R. E., Saito M. A., 2012. Influence of cobalamin scarcity on diatom molecular physiology and identification of a cobalamin acquisition protein. Proc. Natl. Acad. Sci. 109, E1762–E1771.

Braman R. S., Hendrix S. A., 1989. Nanogram nitrite and nitrate determination in environmental and biological materials by vanadium (III) reduction with chemiluminescence detection. Anal. Chem. 61, 2715–2718.

Bridoux M. C., Ingalls A. E., 2010. Structural identification of long-chain polyamines associated with diatom biosilica in a Southern Ocean sediment core. Geochim. Cosmochim. Acta 74, 4044–4057.

Bridoux M. C., Keil R. G., Ingalls A. E., 2012. Analysis of natural diatom communities reveals novel insights into the diversity of long chain polyamine (LCPA) structures involved in silica precipitation. Org. Geochem. 47, 9–21.

Brunelle B. G., Sigman D. M., Cook M. S., Keigwin L. D., Haug G. H., Plessen B., Schettler G., Jaccard S. L., 2007. Evidence from diatom-bound nitrogen isotopes for subarctic Pacific stratification during the last ice age and a link to North Pacific denitrification changes. Paleoceanography 22.

Burkhardt S., Riebesell U., Zondervan I., 1999. Stable carbon isotope fractionation by marine phytoplankton in response to daylength, growth rate, and CO_2 availability. Mar. Ecol. Prog. Ser. 184, 31–41.

Chen Y., Yang J. T., Tang J., Hong H., Kao S., Dai M., Shi D., 2024. Changes in isotope fractionation during nitrate assimilation by marine eukaryotic and prokaryotic algae under different pH and CO_2 conditions. Limnol. Oceanogr. 69, 1045–1055.

Clement R., Dimnet L., Maberly S. C., Gontero B., 2016. The nature of the CO_2 -concentrating mechanisms in a marine diatom, *Thalassiosira pseudonana*. New Phytol. 209, 1417–1427.

Conley D. J., Kilham S. S., Theriot E., 1989. Differences in silica content between marine and freshwater diatoms. Limnol. Oceanogr. 34, 205–212.

Coplen T. B., 2011. Guidelines and recommended terms for expression of stable-isotope-ratio and gas-ratio measurement results. Rapid Commun. Mass Spectrom. 25, 2538–2560.

Degens E. T., Guillard R. R. L., Sackett W. M., Hellebust J. A., 1968. Metabolic fractionation of carbon isotopes in marine plankton—I. Temperature and respiration experiments. Deep Sea Res. Oceanogr. Abstr. 15, 1–9.

Deuser W. G., 1970. Isotopic Evidence for Diminishing Supply of Available Carbon during Diatom Bloom in the Black Sea. Nature 225, 1069–1071.

Dove I. A., Bishop I. W., Crosta X., Riedinger N., Kelly R. P., Robinson R. S., 2025. *Chaetoceros* Resting Spores Do Not Significantly Bias Sedimentary Diatom-Bound Nitrogen Isotope Records Despite Distinctly Low Values. Paleoceanogr. Paleoclimatology 40, e2024PA005041.

Fripiat F., Martínez-García A., Fawcett S. E., Kemeny P. C., Studer A. S., Smart S. M., Rubach F., Oleynik S., Sigman D. M., Haug G. H., 2019. The isotope effect of nitrate assimilation in the Antarctic Zone: Improved estimates and paleoceanographic implications. Geochim. Cosmochim. Acta 247, 261–279.

Fry B., 1996. $^{13}C/^{12}C$ fractionation by marine diatoms. Mar. Ecol. Prog. Ser. 134, 283–294.

Gauthier P. P. G., Lamothe M., Mahé A., Molero G., Nogués S., Hodges M., Tcherkez G., 2013. Metabolic origin of $\delta^{15}N$ values in nitrogenous compounds from Brassica napus L. leaves. Plant Cell Environ. 36, 128–137.

Granger J., Sigman D. M., Rohde M. M., Maldonado M. T., Tortell P. D., 2010. N and O isotope effects during nitrate assimilation by unicellular prokaryotic and eukaryotic plankton cultures. Geochim. Cosmochim. Acta 74, 1030–1040.

Guy R. D., Vanlerberghe G. C., Turpin D. H., 1989. Significance of Phosphoenolpyruvate Carboxylase during Ammonium Assimilation: Carbon Isotope Discrimination in Photosynthesis and Respiration by the N-Limited Green Alga *Selenastrum minutum*. Plant Physiol. 89, 1150–1157.

847 Harrison P. J., Waters R. E., Taylor F. J. R., 1980. A Broad Spectrum Artificial Sea Water Medium
848    for Coastal and Open Ocean Phytoplankton. J. Phycol. 16, 28–35.

849 Horn M. G., Robinson R. S., Rynearson T. A., Sigman D. M., 2011. Nitrogen isotopic relationship
850    between diatom-bound and bulk organic matter of cultured polar diatoms.
851    Paleoceanography 26, 2010PA002080.

852 Ishikawa N. F., Chikaraishi Y., Takano Y., Sasaki Y., Takizawa Y., Tsuchiya M., Tayasu I.,
853    Nagata T., Ohkouchi N., 2018. A new analytical method for determination of the nitrogen
854    isotopic composition of methionine: Its application to aquatic ecosystems with mixed
855    resources. Limnol. Oceanogr. Methods 16, 607–620.

856 Jones C. A., Closset I., Riesselman C. R., Kelly R. P., Brzezinski M. A., Robinson R. S., 2022.
857    New Constraints on Assemblage-Driven Variation in the Relationship Amongst
858    Diatom-Bound, Biomass, and Nitrate Nitrogen Isotope Values. Paleoceanogr.
859    Paleoclimatology 37, e2022PA004428.

860 Kalansky J. F., Robinson R. S., Popp B. N., 2011. Insights into nitrogen cycling in the western
861    Gulf of California from the nitrogen isotopic composition of diatom-bound organic matter.
862    Geochem. Geophys. Geosystems 12.

863 Kirkham A. R., Richthammer P., Schmidt K., Wustmann M., Maeda Y., Hedrich R., Brunner E.,
864    Tanaka T., van Pée K.-H., Falciatore A., Mock T., 2017. A role for the cell-wall protein
865    silacidin in cell size of the diatom *Thalassiosira pseudonana*. ISME J. 11, 2452–2464.

866 Kotzsch A., Gröger P., Pawolski D., Bomans P. H. H., Sommerdijk N. A. J. M., Schlierf M.,
867    Kröger N., 2017. Silicanin-1 is a conserved diatom membrane protein involved in silica
868    biomineralization. BMC Biol. 15, 65.

869 Kröger N., Poulsen N., 2008. Diatoms—From Cell Wall Biogenesis to Nanotechnology. Annu.
870    Rev. Genet. 42, 83–107.

871 Kroth P. G., Chiovitti A., Gruber A., Martin-Jezequel V., Mock T., Parker M. S., Stanley M. S.,
872    Kaplan A., Caron L., Weber T., Maheswari U., Armbrust E. V., Bowler C., 2008. A Model
873    for Carbohydrate Metabolism in the Diatom *Phaeodactylum tricornutum* Deduced from
874    Comparative Whole Genome Analysis. PLOS ONE 3, e1426.

875 Kudo I., 2003. Change in the uptake and cellular Si:N ratio in diatoms responding to the ambient
876    Si:N ratio and growth phase. Mar. Biol. 143, 39–46.

877 Lehmann M. F., Bernasconi S. M., Barbieri A., McKenzie J. A., 2002. Preservation of organic
878    matter and alteration of its carbon and nitrogen isotope composition during simulated and
879    in situ early sedimentary diagenesis. Geochim. Cosmochim. Acta 66, 3573–3584.

880 Lin H.-Y., Liu C.-H., Kang Y.-T., Lin S.-W., Liu H.-Y., Lee C.-T., Liu Y.-C., Hsu M.-C., Chien
881    Y.-Y., Hong S.-M., Cheng Y.-H., Hsieh B.-Y., Lin H.-J., 2024. Enhancing the Spermidine
882    Synthase-Based Polyamine Biosynthetic Pathway to Boost Rapid Growth in Marine
883    Diatom Phaeodactylum tricornutum. Biomolecules 14, 372.

Macko S. A., Fogel M. L., Hare P. E., Hoering T. C., 1987. Isotopic fractionation of nitrogen and
carbon in the synthesis of amino acids by microorganisms. Chem. Geol. Isot. Geosci. Sect.
65, 79–92.

Mariotti A., Germon J. C., Hubert P., Kaiser P., Letolle R., Tardieux A., Tardieux P., 1981.
Experimental determination of nitrogen kinetic isotope fractionation: Some principles;
illustration for the denitrification and nitrification processes. Plant Soil 62, 413–430.

McCarthy M. D., Lehman J., Kudela R., 2013. Compound-specific amino acid δ^{15}N patterns in
marine algae: Tracer potential for cyanobacterial vs. eukaryotic organic nitrogen sources
in the ocean. Geochim. Cosmochim. Acta 103, 104–120.

McMahon K. W., McCarthy M. D., 2016. Embracing variability in amino acid δ<sup>15</supN
fractionation: mechanisms, implications, and applications for trophic ecology. Ecosphere
7, e01511.

Morales L. V., Granger J., Chang B. X., Prokopenko M. G., Plessen B., Gradinger R., Sigman D.
M., 2014. Elevated ^{15}N/^{14}N in particulate organic matter, zooplankton, and diatom
frustule-bound nitrogen in the ice-covered water column of the Bering Sea eastern shelf.
Deep Sea Res. Part II Top. Stud. Oceanogr. 109, 100–111.

Morales L. V., Sigman D. M., Horn M. G., Robinson R. S., 2013. Cleaning methods for the isotopic
determination of diatom-bound nitrogen in non-fossil diatom frustules. Limnol. Oceanogr.
Methods 11, 101–112.

Mortlock R. A., Charles C. D., Froelich P. N., Zibello M. A., Saltzman J., Hays J. D., Burckle L.
H., 1991. Evidence for lower productivity in the Antarctic Ocean during the last glaciation.
Nature 351, 220–223.

Needoba J. A., Waser N. A., Harrison P. J., Calvert S. E., 2003. Nitrogen isotope fractionation in
12 species of marine phytoplankton during growth on nitrate. Mar. Ecol. Prog. Ser. 255,
81–91.

Parker M. S., Armbrust E. V., Piovia-Scott J., Keil R. G., 2004. Induction of Photorespiration by
Light in the Centric Diatom *Thalassiosira weissflogii* (bacillariophyceae): Molecular
Characterization and Physiological Consequences. J. Phycol. 40, 557–567.

Parker M. S., Mock T., Armbrust E. V., 2008. Genomic Insights into Marine Microalgae. Annu.
Rev. Genet. 42, 619–645.

Petit J. R., Jouzel J., Raynaud D., Barkov N. I., Barnola J.-M., Basile I., Bender M., Chappellaz J.,
Davis M., Delaygue G., Delmotte M., Kotlyakov V. M., Legrand M., Lipenkov V. Y.,
Lorius C., PÉpin L., Ritz C., Saltzman E., Stievenard M., 1999. Climate and atmospheric
history of the past 420,000 years from the Vostok ice core, Antarctica. Nature 399,
429–436.

Poulsen N., Sumper M., Kröger N., 2003. Biosilica formation in diatoms: Characterization of
native silaffin-2 and its role in silica morphogenesis. Proc. Natl. Acad. Sci. 100,
12075–12080.

R Core Team, 2023. R: A language and environment for statistical computing.

Racusen D., Foote M., 1960. Amino acid turnover and protein synthesis in leaves. Arch. Biochem.
Biophys. 90, 90–95.

Ren H., Studer A. S., Serno S., Sigman D. M., Winckler G., Anderson R. F., Oleynik S., Gersonde
R., Haug G. H., 2015. Glacial-to-interglacial changes in nitrate supply and consumption in
the subarctic North Pacific from microfossil-bound N isotopes at two trophic levels.
Paleoceanography 30, 1217–1232.

Richthammer P., Börmel M., Brunner E., van Pée K.-H., 2011. Biomineralization in Diatoms: The
Role of Silacidins. ChemBioChem 12, 1362–1366.

Riebesell U., Wolf-Gladrow D. A., Smetacek V., 1993. Carbon dioxide limitation of marine
phytoplankton growth rates. Nature 361, 249–251.

Robinson D. B., Rognlien J. L., Bauer C. A., Simmons B. A., 2007. Dependence of amine-
accelerated silicate condensation on amine structure. J. Mater. Chem. 17, 2113.

Robinson R. S., Brunelle B. G., Sigman D. M., 2004. Revisiting nutrient utilization in the glacial
Antarctic: Evidence from a new method for diatom-bound N isotopic analysis.
Paleoceanography 19.

Robinson R. S., Jones C. A., Dove I. A., Kelly R. P., Brzezinski M. A., 2025. Monospecific Diatom
Cultures Suggest Potential Interspecies Variation of Diatom-Bound Nitrogen Isotope
Signatures Associated With Silica Acquisition. Geochem. Geophys. Geosystems 26,
e2024GC011923.

Robinson R. S., Jones C. A., Kelly R. P., Love A., Closset I., Rafter P. A., Brzezinski M., 2020. A
Test of the Diatom-Bound Paleoproxy: Tracing the Isotopic Composition of Nutrient-
Nitrogen Into Southern Ocean Particles and Sediments. Glob. Biogeochem. Cycles 34,
e2019GB006508.

Robinson R. S., Kienast M., Luiza Albuquerque A., Altabet M., Contreras S., De Pol Holz R.,
Dubois N., Francois R., Galbraith E., Hsu T.-C., Ivanochko T., Jaccard S., Kao S.-J., Kiefer
T., Kienast S., Lehmann M., Martinez P., McCarthy M., Möbius J., Pedersen T., Quan T.
M., Ryabenko E., Schmittner A., Schneider R., Schneider-Mor A., Shigemitsu M., Sinclair
D., Somes C., Studer A., Thunell R., Yang J.-Y., 2012. A review of nitrogen isotopic
alteration in marine sediments. Paleoceanography 27.

Robinson R. S., Sigman D. M., DiFiore P. J., Rohde M. M., Mashiotta T. A., Lea D. W., 2005.
Diatom-bound ^{15}N/^{14}N: New support for enhanced nutrient consumption in the ice age
subantarctic. Paleoceanography 20.

Sharkey T. D., Berry J. A., 1985. Carbon isotope fractionation of algae as influenced by an
inducible CO_2 concentrating mechanism. Inorg. Carbon Uptake Aquat. Photosynth. Org.,
389–401.

Sigman D. M., Altabet M. A., Francois R., McCorkle D. C., Gaillard J.-F., 1999. The isotopic
composition of diatom-bound nitrogen in Southern Ocean sediments. Paleoceanography
14, 118–134.

Sigman D. M., Casciotti K. L., Andreani M., Barford C., Galanter M., Böhlke J. K., 2001. A
Bacterial Method for the Nitrogen Isotopic Analysis of Nitrate in Seawater and Freshwater.
Anal. Chem. 73, 4145–4153.

Smith S. R., Dupont C. L., McCarthy J. K., Broddrick J. T., Oborník M., Horák A., Füssy Z., Cihlář
J., Kleessen S., Zheng H., McCrow J. P., Hixson K. K., Araújo W. L., Nunes-Nesi A.,
Fernie A., Nikoloski Z., Palsson B. O., Allen A. E., 2019. Evolution and regulation of
nitrogen flux through compartmentalized metabolic networks in a marine diatom. Nat.
Commun. 10, 4552.

Stief P., Schauberger C., Lund M. B., Greve A., Abed R. M. M., Al-Najjar M. A. A., Attard K.,
Bonaglia S., Deutzmann J. S., Franco-Cisterna B., García-Robledo E., Holtappels M., John
U., Maciute A., Magee M. J., Pors R., Santl-Temkiv T., Scherwass A., Sevilgen D. S., De
Beer D., Glud R. N., Schramm A., Kamp A., 2022. Intracellular nitrate storage by diatoms
can be an important nitrogen pool in freshwater and marine ecosystems. Commun. Earth
Environ. 3, 154.

Strickland J. D. H., Parsons T. R., 1972. A Practical Handbook of Seawater Analysis, 2nd edition.

Studer A. S., Sigman D. M., Martínez-García A., Benz V., Winckler G., Kuhn G., Esper O., Lamy
F., Jaccard S. L., Wacker L., Oleynik S., Gersonde R., Haug G. H., 2015. Antarctic Zone
nutrient conditions during the last two glacial cycles. Paleoceanography 30, 845–862.

Studer A. S., Wörmer L., Vogel H., Dubois N., Bartosiewicz M., Hinrichs K.-U., Lepori F.,
Lehmann M. F., 2024. First lacustrine application of the diatom-bound nitrogen isotope
paleo-proxy reveals coupling of denitrification and N fixation in a hyper-eutrophic lake.
Limnol. Oceanogr. 69, 1797–1809.

Sumper M., Brunner E., 2008. Silica Biomineralisation in Diatoms: The Model Organism
Thalassiosira pseudonana. ChemBioChem 9, 1187–1194.

Takizawa Y., Dharampal P. S., Steffan S. A., Takano Y., Ohkouchi N., Chikaraishi Y., 2017.
Intra-trophic isotopic discrimination of $^{15}N/^{14}N$ for amino acids in autotrophs: Implications
for nitrogen dynamics in ecological studies. Ecol. Evol. 7, 2916–2924.

Tcherkez G., 2010. Natural $^{15}N/^{14}N$ isotope composition in C_3 leaves: are enzymatic isotope effects
informative for predicting the 15N-abundance in key metabolites? Funct. Plant Biol. 38,
1–12.

Tcherkez G., Hodges M., 2008. How stable isotopes may help to elucidate primary nitrogen
metabolism and its interaction with (photo)respiration in C_3 leaves. J. Exp. Bot. 59,
1685–1693.

Tesson B., Lerch S. J. L., Hildebrand M., 2017. Characterization of a New Protein Family
Associated With the Silica Deposition Vesicle Membrane Enables Genetic Manipulation
of Diatom Silica. Sci. Rep. 7, 13457.

Tortell P. D., 2000. Evolutionary and ecological perspectives on carbon acquisition in
phytoplankton. Limnol. Oceanogr. 45, 744–750.

Vokhshoori N. L., Rick T. C., Braje T. J., McCarthy M. D., 2023. Preservation of stable isotope
signatures of amino acids in diagenetically altered Middle to Late Holocene archaeological
mollusc shells. Geochim. Cosmochim. Acta 352, 36–50.

Waser N. A. D., Harrison P. J., Nielsen B., Calvert S. E., Turpin D. H., 1998. Nitrogen isotope
fractionation during the uptake and assimilation of nitrate, nitrite, ammonium, and urea by
a marine diatom. Limnol. Oceanogr. 43, 215–224.

Werner R. A., Schmidt H.-L., 2002. The in vivo nitrogen isotope discrimination among organic
plant compounds. Phytochemistry 61, 465–484.

Yoneyama T., Fujihara S., Yagi K., 1998. Natural abundance of [15]N in amino acids and polyamines
from leguminous nodules: unique [15]N enrichment in homospermidine. J. Exp. Bot. 49,
521–526.

1010

1011

1012