Essential role of astrocytes in rapid activity-dependent restructuration of the axon initial segment

Rafael Sanz-Gálvez[1,2]†, Yanis Inglebert[1,2]†, Arlette Kolta[1,2,3]

[1]Department of Neurosciences, Université de Montréal, Montréal, Canada

[2]Centre Interdisciplinaire de recherche sur le cerveau et l'apprentissage (CIRCA), Montréal, Québec, Canada

[3]Faculté de Médecine Dentaire, Université de Montréal, Montréal, Québec, Canada.

†These authors contributed equally to this work and share first authorship

Corresponding authors: arlette.kolta@umontreal.ca

Keywords: Astrocytes, Intrinsic Excitability, LTP-IE, Visual Cortex, S100β, Axon Initial Segment

34 pages, 82 references, 5 figures, 4 Supplementary figures

ABSTRACT

One key regulator of neuronal excitability is the axon initial segment (AIS), a highly specialized axonal region, enriched in ion channels, where action potentials are initiated. The AIS can undergo significant morphological changes to fine-tune neuronal excitability in response to external perturbations. Long considered solely a homeostatic mechanism operating over long timescales (hours to days) to adjust excitability, we show here that this phenomenon can also occur rapidly, within minutes, following a brief period of high activity in layer 5 pyramidal neurons of the visual cortex. Because astrocytes have been known to regulate neuronal excitability, we explored the effects of gliotransmitters on this process and identified the calcium-binding protein S100β from astrocytes to be required for the rapid reorganization of the AIS.

INTRODUCTION

The plasticity of the input-output function is regulated by diverse plasticity rules to adapt neuron' excitability in response to external or internal perturbations. One powerful regulator is the axon initial segment (AIS), a specialized subcellular region of the proximal axon where action potentials are shaped and initiated (Debanne et al., 2011). Structural plasticity of the AIS can influence neuronal output as demonstrated by changes in AIS length in response to external perturbation to maintain appropriate neuronal excitability (Yamada and Kuba, 2016; Fréal et al., 2023). The length of the AIS is a reliable indicator of neuronal excitability, where increased length enhances excitability and decreased length lowers it. Plasticity of the AIS is highly used in different sensory processing brain regions to fine-tune neuronal excitability or during development to maintain stable network excitability (Kuba et al., 2006; Schlüter et al., 2017). For instance, in the visual cortex, the AIS is developmentally regulated by visual experience during critical period (Gutzmann et al., 2014; Höfflin et al., 2017). Historically, AIS plasticity has been considered a homeostatic mechanism operating over long timescales (days to weeks), allowing neurons to maintain their firing set point in response to network perturbations. This view largely stems from studies conducted in neuronal cultures following chronic network manipulations (Grubb and Burrone, 2010). This perspective has evolved

over time, with plasticity being described at progressively shorter timescales, from hours to

minutes (Grubb et al., 2011; Evans et al., 2015). The latest findings, using newly developed

genetic tools (Thome et al., 2025) that label the scaffolding protein Ankyrin G (AnkG), a

specific marker of the AIS (Leterrier, 2018) reveal that AIS reorganization can occur within

minutes following long-term depression in the hippocampus (Fréal et al., 2023). This is the

first evidence showing that the synergistic excitability changes observed following synaptic

plasticity could be accompanied by AIS plasticity (Campanac and Debanne, 2007, 2008).

However, synaptic plasticity is not the sole mechanism of memory storage; intrinsic

excitability plasticity can also occur in response to non-homeostatic stimulation and could

be associated as well with rapid AIS plasticity, a possibility that, until now, has never been

tested. Here, we provide initial evidence of a close relationship between intrinsic excitability

changes and AIS plasticity. In Layer 5 cortical pyramidal neurons (L5PNs), repetitive bursts

high-frequency firing induce long-term potentiation of intrinsic excitability (LTP-IE) a

fundamental mechanism to refine visual circuit (Cudmore and Turrigiano, 2004; Nataraj et

al., 2010; Duménieu et al., 2025). In cortical circuit, LTP-IE is dependent on extracellular

calcium influx and modulation of ion channels (Cudmore and Turrigiano, 2004). Among the

various external factors that could influence LTP-IE, astrocytes have long been overlooked

despite increasing evidence of their role as powerful modulators of input-output functions

and sensory processing (Sanz-Gálvez et al., 2024). Recent findings show that astrocytes

directly interact with the axon or the AIS to modulate excitability through the release of

gliotransmitters (Debanne and Rama, 2011; Lezmy et al., 2021). Moreover, in murine models

of Alzheimer's disease, AIS integrity and length deteriorate due to transcriptomic and

proteomic alterations in astrocytes (Benitez et al., 2024). One relatively understudied

gliotransmitter, S100β, is particularly intriguing as it functions both as a calcium (Ca^{2+})-

binding protein to regulate the concentration of extracellular calcium and a modulator of

various voltage-gated ion channels (Hermann et al., 2012; Bancroft et al., 2022).

Extracellularly, it primarily binds to membrane receptors such as the multi-ligand Receptor

for Advanced Glycation End Products (RAGE), triggering intracellular signaling cascades that,

depending on its concentration, may lead to physiological or pathological outcomes (Donato

et al., 2009). We have previously shown that S100β and stimulation of astrocytes

(presumably through release of S100β) elicit spiking in L5PNs from the visual cortex and

trigeminal primary afferent axons or promote rhythmic activity in trigeminal sensori-motor

circuit through the enhanced activation of voltage-gated sodium (NaV) 1.6 channels

(Morquette et al., 2015; Ryczko et al., 2021; Gaudel et al., 2025). Interestingly, NaV 1.6 is one

of the main isoforms of NaV channels expressed at the AIS in excitatory neurons which

distribution is reorganized during plasticity (Fréal et al., 2023). Here, we first test whether LTP-

IE is associated with rapid AIS plasticity in L5PNs of the visual cortex and then examine if

astrocytes regulate this process through the release of the gliotransmitter S100β.

RESULTS

Long-lasting potentiation of intrinsic excitability (LTP-IE) is associated with rapid AIS plasticity

Layer 5 pyramidal neurons (L5PNs) were recorded in acute slices from the primary visual

cortex (V1) obtained from Vglut2-AnkG-GFP young mice (P14–P30) (see Methods and **Fig.**

1A), and their intrinsic excitability was monitored by a 500-ms constant direct current pulse

every 10 seconds for 5 min after which repetitive current injection (500-ms depolarizing

pulses, 60 times at 4-second interval, **Fig. 1B bottom)** was used to induce long-lasting

potentiation of intrinsic excitability (LTP-IE). All experiments were conducted in the presence

of glutamatergic and GABAergic receptors blockers to prevent interferences from

synaptically released fast excitatory and inhibitory neurotransmitters. As previously

described (Cudmore and Turrigiano, 2004), this protocol induced a robust and long-lasting

increase in intrinsic excitability (151% ± 12, n=6, **Fig. 1B**), which was monitored every 10

seconds for 30 minutes after induction. After confirming our immunohistochemical results

showing that the GFP signal colocalized with the master AIS protein, AnkG (**Fig. 1A top**), we

proceeded to monitor AIS remodeling in real-time before and after the induction protocol in

the recorded neuron filled with Alexa Fluor 647 (**Fig. 1A bottom**). Fluorescence images of

AnkG-GFP taken 30 minutes after induction revealed a rapid and significant elongation in the

distal part of the AIS, as confirmed by fluorescence intensity measurements (mean AIS

length: 29.68 ± 1.25 μm before induction and 33.29 ± 1.35 μm after induction, n=6, t-test,

P<0.001; **Fig. 1C**). To avoid rapid deterioration due to phototoxicity, imaging of the AIS was

minimized, but in a separate set of experiments (n=5), images were captured every 5 minutes,

revealing that in some cells, the increase began as early as 10–15 minutes after induction

(Fig. S1). In the absence of the induction protocol, no LTP-IE (102% ± 2, n=5, **Fig. 1B**) or AIS

remodeling was observed (mean AIS length: 34.31 ± 0.95 μm before and 34.21 ± 0.92 μm

after, n=5, t-test, P>0.05; **Fig. 1F**). To determine changes in neuronal excitability, the number

of APs for each increase in injected current was counted, establishing input-output curves

before and after the induction protocol. Additionally, rheobase, an important parameter of

neuronal excitability, was measured. In neurons subjected to the brief repetitive firing period,

a significant increase in intrinsic excitability was observed across other supra-threshold

current amplitudes (from 30 pA to 100 pA, t-test, P<0.01; P<0.05; **Fig. 1D**), whereas no

significant changes were observed in control neurons (all steps; t-test, P>0.05; **Fig. 1G**).

Moreover, in neurons subjected to the induction protocol, but not in control neurons (**Fig.**

1H), a significant decrease in rheobase was observed (30 ± 12 pA before vs. 21 ± 9 pA after

induction, n=6, t-test, P<0.01; **Fig. 1E**). Together, these data demonstrate that a period of

high-frequency activation as brief as 4 min suffice to induce remodeling of the AIS and LTP-

IE in L5PNs.

LTP-IE and AIS plasticity involves peri-AIS astrocyte signaling

Astrocytes are involved in various forms of synaptic plasticity (Yang et al., 2003; Andersson

et al., 2007; Falcón-Moya et al., 2020; Sanz-Gálvez et al., 2024); however, their potential

contribution to intrinsic plasticity has not been assessed. Therefore, we examined whether

peri-AIS astrocytes (astrocytes near the AIS) are necessary for LTP-IE and the consequent

structural remodeling of the AIS in L5PN neurons. To address this, we preincubated acute

brain slices for 5 minutes RT with Sulforhodamine (SR101, 1 μM), a widely used astrocyte

marker in different brain regions (**Fig. 2A**) (Watanabe et al., 2023). Next, a single astrocyte

was loaded with 20 mM of the Ca^{2+} chelator BAPTA via a patch pipette to inhibit Ca^{2+}-

dependent vesicular release and gliotransmitter signaling (Araque et al., 1998) and a nearby

L5PN (distance between the AIS of the recorded neuron and the patched astrocyte; 37.02 ±

8.74 μm, **Fig. 2B**) was targeted for whole-cell recording after allowing time (15 min) for BAPTA

to diffuse in the astrocytic syncitium. Astrocytic inactivation with BAPTA prevented the

induction of LTP-IE (103% ± 4, n=5; **Fig. 2B**) and the associated AIS lengthening (mean AIS

length: 33.4 ± 0.95 µm before induction vs. 32.61 ± 0.94 µm after induction, n=5, t-test,

P>0.05; **Fig. 2C**). No significant changes were observed in input-output curves or rheobase

values before and after induction (**Fig. 2D,E**). Taken together, these results clearly indicate

that peri-AIS astrocytes are required for the induction of this form of intrinsic excitability

plasticity and AIS remodeling.

LTP-IE and AIS plasticity depend on the astrocytic protein S100β

In the visual cortex, LTP-IE, as other types of synaptic plasticity, depends on extracellular Ca^{2+}

(Inglebert et al., 2020), since it is abolished in 0mM Ca^{2+} medium (Cudmore and Turrigiano,

2004). Our previous studies demonstrated that astrocytes could tightly regulate extracellular

Ca^{2+} levels through the release of S100β (Morquette et al., 2015) and that optogenetic

activation of layer 5 astrocytes in the visual cortex decreases extracellular Ca^{2+} via S100β

release (Ryczko et al., 2021). Based on these findings, we investigated whether this Ca^{2+}-

binding astrocytic protein could locally modulate LTP-IE and AIS structural remodeling. To

this end, we locally administered an anti-S100β antibody in the vicinity of the AIS for 5

minutes prior to induction to prevent the endogenous protein from exerting its effects and

from binding Ca^{2+} (Morquette et al., 2015). Applications of the same antibody, but

denatured by preheating to 100 degrees Celsius for 20 minutes, served as controls. The

S100β antibody effectively blocked LTP-IE, whereas the denatured antibody did not preclude

its induction (S100β antibody: 106% ± 17, n=5 vs. S100β antibody denatured: 180% ± 9, n=6,

(MW) U test, P<0.05; **Fig. 3A**). Confocal imaging further revealed that the S100β antibody

prevented AIS elongation (mean AIS length: 34.76 ± 0.41 µm before induction and 34.48 ±

0.48 µm after induction, n=5, t-test, P>0.05; **Fig. 3B**). Conversely, with the denatured

antibody, significant AIS elongation was observed (mean AIS length: 32.45 ± 1.67 µm before

induction and 34.63 ± 1.62 µm after induction, n=6, t-test, P<0.001; **Fig. 3E**) similar to what

is observed in control condition (**Fig 1C**). The S100β antibody application was found to cause

a small decrease of excitability as measured in the input-output curves, but this change was

not statistically significant (all steps; t-test, P>0.05; **Fig. 3C**). This effect is likely attributable

to an increase in extracellular Ca^{2+} levels induced by S100β blockade, which is consistent

with prior findings indicating that elevated extracellular Ca^{2+} levels suppress neuronal

excitability (Forsberg et al., 2019, 2025). In contrast, the denatured antibody application is

associated with an upward shift of the input-output curve (pA40, pA60-pA100, t-test, P<0.05;

Fig. 3F). No significant changes were observed in rheobase under any condition (**Fig. 3D,G**).

Collectively, these data underscore the necessity of the Ca^{2+}-binding astrocytic protein

S100β for the induction of LTP-IE and AIS morphological modifications in LPN5s.

Optogenetic activation of peri-AIS astrocytes induces LTP-IE and AIS plasticity through S100β release

Having established that astrocytes and S100β are critically involved in mediating LTP-IE and

AIS remodeling, we next aimed to determine whether selective activation of peri-AIS

astrocytes alone could suffice to induce LTP-IE. To address this, neonatal VGlut2-AnkG-GFP

mice (P0) were bilaterally injected intracerebrally with an adeno-associated virus (AAV)

expressing Channelrhodopsin-2 (ChR2) fused to the mCherry reporter and targeted to

astrocytes using the GFAP promoter (**Fig. 4A left**). To verify that viral transfection was

specifically restricted to astrocytes, immunohistochemistry was performed on V1 sections

using an antibody against S100β. Immunostaining revealed that viral expression was targeted

to astrocytes, as evidenced by the colocalization of mCherry/ChR2-GFAP with S100β (**Fig.**

4A right). Next, in acute brain slices, patch-clamp recordings demonstrated that transfected

astrocytes responded to light pulses (using a combination of 440 nm and 488 nm lasers from

a confocal microscope) by evoking membrane depolarizations (**Fig. S2**). This response was

verified in representative experiments (**see Methods**). Additionally, we confirmed that viral

transfection did not affect the real-time visualization of AIS morphology (**Fig. 4B bottom left**).

Subsequently, excitability of L5PNs was assessed as above, but the induction protocol was

substituted by photostimulation of astrocytes (short light pulses of 2 seconds on, followed

by 2 seconds off, repeated 60 times; **Fig. S2**) which was as efficient for induction of LTP-IE

(**Fig. 4B**). Importantly, local administration of the S100β antibody for 5 minutes prior to

optogenetic stimulation of peri-AIS astrocytes abolished LTP-IE (opto Astro: 160% ± 10, n=6

vs. S100β antibody + opto Astro: 101% ± 3, n=6; (MW) U test, P<0.01; **Fig. 4B**). Confocal

imaging revealed a significant elongation of the AIS length 30 minutes after optogenetic

stimulation of peri-AIS astrocytes (mean AIS length: 31.96 ± 2.05 µm pre-stimulation vs.

35.25 ± 2.20 µm post-stimulation, n=6, t-test, P<0.01; **Fig. 4C**), which was prevented when

the S100β antibody was applied locally (mean AIS length: 33.14 ± 1.98 µm pre-stimulation

vs. 32.85 ± 2.15 µm post-stimulation, n=6, t-test, P>0.05; **Fig. 4D**). Input-output curves

showed that responses to suprathreshold current amplitudes significantly increased after

optogenetic stimulation of peri-AIS astrocytes (pA30-pA80, t-test, P<0.01; P<0.05; **Fig. 4E**),

whereas no changes were observed when the S100β antibody was applied prior to optical

stimulation (all steps; t-test, P>0.05; **Fig. 4F**). Changes in rheobase values were observed in

recorded neurons after optogenetic activation of peri-AIS astrocytes (33 ± 15 pA before vs.

25 ± 10 pA after stimulation, n=6, t-test, P<0.05; **Fig. 4E**), whereas no significant changes

were detected following prior application of the S100β antibody before activation (**Fig. 4F**). It

might be expected that S100β is not the only gliotransmitter released by astrocytes to

regulate this form of LTP-IE and rapid AIS plasticity. Given that ATP/adenosine released by

astrocytes can alter AIS excitability (Lezmy et al., 2021), we decided to test the local

administration of Suramin (100 µM) (a purinergic antagonist) for 5 minutes before

optogenetic stimulation of astrocytes. However, our results showed that the administration

of Suramin did not abolish either LTP-IE or the morphological changes in the AIS (**Fig. S3**).

These findings demonstrate that selective activation of peri-AIS astrocytes is sufficient to

induce LTP-IE and AIS remodeling, highlighting the crucial involvement of astrocyte-derived

S100β in regulating neuronal excitability and axonal plasticity.

The astrocytic protein S100β rescues LTP-IE and rapid AIS plasticity

To further confirm the key role of S100β in LTP-IE and rapid AIS remodeling observed in V1

L5PNs, we repeated the previous experiment of astrocytic network inhibition but with local

application of S100β (129 µM) for 5 minutes prior to the induction protocol. This approach

allowed us to fully saturate the local environment without inducing spiking during the

protocol itself, which could otherwise alter the results. Consistent with our previous findings,

application of S100β to the proximal axon, while peri-AIS astrocytes were inactivated, was

sufficient to rescue LTP-IE (157% ± 12, n=6 vs 102% ± 6, n=5, (MW) U test, P<0.01, **Fig. 5A**).

Confocal imaging also revealed rescue of AIS remodeling with S100β application (mean AIS

length: 33.45 ± 1.78 µm before induction and 36.57 ± 2.10 µm after induction, n=6, t-test,

P<0.01; **Fig. 5B**). Input-output curves showed that responses to suprathreshold current

amplitudes significantly increased after application of S100β (pA40-pA100, t-test, P<0.05;

Fig. 5C). Additionally, significant differences were observed in rheobase values (45 ± 16 pA

before vs. 31 ± 14 pA after induction, n=6, t-test, P<0.05; **Fig. 5D**). To insure that the effects

of the locally applied S100β resulted from its Ca^{2+} chelating action, we confirmed these

findings with the local extracellular application of BAPTA (5 mM) which exerts similar effects

on Ca^{2+} (**Fig. S4**). Together, these results further support the role of S100β as a key regulator

of LTP-IE and rapid AIS remodeling.

DISCUSSION

Previous studies have shown that the AIS is a highly dynamic structure that contributes to

the fine regulation of neuronal output (Grubb and Burrone, 2010; Evans et al., 2013).

However, until now, plastic changes in the AIS have been limited to static post-hoc snapshots

in large, differentially treated populations, preventing the detection of small-scale changes

that may be obscured due to intercellular heterogeneity. Here, we have leveraged a new

AnkG-GFP mouse model (Thome et al., 2025) to monitor the AIS in real time at the single-cell

level. We found that in the visual cortex, the long-lasting increase of intrinsic excitability (LTP-

IE) produced by a brief period of high frequency activity as previously described by Cudmore

and Turrigiano (2004) is associated with rapid structural plasticity of the AIS, predominantly

characterized by a distal elongation of its length (**Fig. 1**). This observation aligns with previous

studies (Kuba et al., 2015; Jamann et al., 2021) showing that AIS elongation is linked to

increased intrinsic excitability and a lower action potential (AP) threshold, highlighting a

coordinated interplay between functional and structural plasticity of the AIS. Astrocytes have

long been recognized as regulators of synaptic transmission, plasticity, and excitability

through gliotransmitter release and local ion homeostasis. However, to our knowledge, their

potential involvement in the plasticity of intrinsic excitability, including AIS morphological

changes, has not yet been explored. We have demonstrated that peri-AIS astrocytes

(proximal or adjacent to the AIS) can regulate the structural and functional plasticity of the

AIS following brief periods of intrinsic spiking activity. This regulation depends on the release

of the astrocytic calcium-binding protein S100β.

LTP-IE is associated with increases AIS

Our results demonstrate that LTP-IE in L5PNs of the visual cortex is associated with a rapid

increase in AIS length. This finding is significant, as it highlights the AIS's ability to very rapidly

(within minutes) adjust its length to regulate neuronal excitability, while it was believed to

occur only on longer timescale (hours to days) acting mostly as a homeostatic mechanism.

These results demonstrate that AIS morphology plasticity is also a mechanism allowing a

neuron to adapt rapidly its excitability in addition to others such as regulation of ion channels

to modulate spike thresholds or EPSP amplification (Debanne and Russier, 2019). LTP-IE in

non-homeostatic form has been described in other brain regions including the hippocampus

and the brainstem (Nelson et al., 2003; Sourdet et al., 2003) and it would be interesting to

know if it is accompanied by changes in AIS morphology as well. The mechanisms described

for LTP-IE so far seem to vary from a region to another involving Protein Kinase A in the visual

cortex (Cudmore and Turrigiano, 2004), metabotropic glutamate receptor subtype 5

(mGluR5) in the somatosensory cortex, NMDA receptors in the hippocampus (Sourdet et al.,

2003; Xu et al., 2005), down-regulation of Kv1 channels in the dorsal lateral geniculate

nucleus (Duménieu et al., 2025) or reduction in BK-type calcium-activated potassium

currents in the brainstem (Nelson et al., 2003). However, all these changes have ultimately

been associated with the modulation (increase or decrease) of intracellular or extracellular

calcium, as they are inhibited by calcium chelation. This could also implicate the astrocytic

calcium-binding protein S100β, which is strongly expressed in all these plastic regions, and

that we demonstrated is required for LTP-IE in the visual cortex. But S100β is much more than

a simple Ca^{2+} chelator (Sorci et al., 2010).

S100β: a multifaceted protein

Given that extracellular calcium is a key regulator of all forms of plasticity, S100β, which

reduces $[Ca^{2+}]_e$, is expected to have a significant impact on these processes. Surprisingly, our

results demonstrated that S100β does not inhibit LTP-IE or AIS plasticity in the visual cortex;

rather, it is essential for these processes despite that both have been shown to depend on calcium elevation and Ca^{2+}-activated molecules such as calcineurin and cyclin-dependent kinase 5 (cdk5) (Chand et al., 2015). This underscores the complex and multifaceted role of S100β, which extends beyond calcium regulation. S100β has been shown to interact directly or indirectly with various voltage gated channels, including Ca^{2+}, K^+, and Na^+ channels (Hermann et al., 2012). For example, it facilitates Ca^{2+} flux across artificial membranes, activates L-type Ca^{2+} channels, and inhibits K^+ channels such as the ether-à-go-go (hEAG1) and Kv4.2 channels (Bancroft et al., 2022). Our previous results demonstrated that S100β enhances the spiking duration and frequency of L5PNs through NaV1.6 potentiation (Ryczko et al., 2021; Gaudel et al., 2025). This is explained by the fact that Na^+ channels are transiently clogged by Ca^{2+} ions occupying the channel lumen (Armstrong, 1999; Armstrong and Cota, 1999). Notably, LTP-IE requires neurons to fire at approximately 20–30 Hz during the induction protocol. The release of S100β could further promote high-frequency firing and facilitate LTP-IE induction. This idea is further supported by the fact that BAPTA rescues LTP-IE **(Fig S4)**. In addition, activity-dependent relocation of the AIS has been shown to be mediated by T- and L- type calcium channels (Grubb and Burrone, 2010). Similar to what has been described for S100A1, another member of the S100 protein family, S100β could directly bind to L-type calcium channels and increase their opening probability through protein kinase A (PKA) activation (Treves et al., 1997) which is required for LTP-IE activity in the visual cortex (Cudmore and Turrigiano, 2004). S100β secretion is tightly linked to neuronal activity (Nardin et al., 2009), and could serve as a gatekeeper of intrinsic plasticity to allow rapid AIS plasticity in response to high periods of activity, a processus which could be really important during development, especially in the visual cortex (Gutzmann et al., 2014). In addition, S100β could provide a direct mechanism for astrocytes to modulate synaptic plasticity through Ca^{2+} regulation. S100β has long been considered as a direct competitor of calmodulin, another Ca^{2+}-binding protein that plays a key role in synaptic plasticity (Xia and Storm, 2005), which can tilt the balance toward LTD or LTP depending on the level of Ca^{2+} elevation. In line with this idea, S100β perfusion enhanced LTP in the hippocampus (Nishiyama et al., 2002). Taken together, these findings suggest that S100β may also

influence other forms of plasticity beyond LTP-IE. Given the well-established interplay

between synaptic and intrinsic plasticity (Debanne et al., 2019), it would be interesting to

explore whether the interplay between synaptic and intrinsic plasticity involves structural

modifications at the AIS, and whether S100β, or other gliotransmitters, act as key mediators

of these coordinated forms of plasticity.

S100β, NaV 1.6 and AIS reorganization

Na^+ channels are transiently clogged by Ca^{2+} ions occupying the channel lumen. Reduction

of extracellular Ca^{2+} relieves this partial block of the Na^+ channel and allows for a larger influx

of Na^+ currents (Armstrong, 1999; Armstrong and Cota, 1999), and an enhanced sodium

persistent current (I_{NaP}), which is associated with the control of neuronal excitability and

rhythmic activity (Agrawal et al., 2001; Wu et al., 2005). I_{NaP} is most commonly associated to

Na^+-channels including the Nav1.6 α-subunit (Drouillas et al., 2023), which is abundantly

located at the distal end of the AIS (Van Wart et al., 2007; Hu et al., 2009; Drouillas et al.,

2023), where the initiation of APs in L5PNs has been experimentally demonstrated (Palmer

and Stuart, 2006), coinciding with the region where we observed structural changes in the

AIS in our experiments (**Fig. 1C**). Furthermore, according to computational and experimental

analyses, Nav1.6 α-subunits determine the lowest threshold for AP initiation in the distal AIS

of cortical pyramidal neurons (Hu et al., 2009). We observed a reduction in the threshold

potential for evoking APs (**Fig. 1E**), which could indicate an increase in the density and/or

reinsertion of these channels after inducing LTP-IE. The absence of changes in passive cell

properties (see Methods) further supports that LTP-IE associated with AIS elongation is

mediated by changes in ionic conductances. Molecular trafficking of protein complexes in

the AIS can follow two routes during development: (1) either insertion of proteins such as

neurofascin 186 (Nfasc186) or the Kv7.3 channel into the somatodendritic and/or axonal

membrane followed by bidirectional diffusion until immobilization by interaction with

AnkyrinG (AnkG) (Ghosh et al., 2020); or (2) preferential vesicular insertion directly into the

AIS, allowing binding to AnkG, as in the case of Nav1.6 (Akin et al., 2015). The maintenance

of Nav1.6 channels has not yet been thoroughly studied, but so far, it appears that once

anchored, they do not migrate to adjacent membranes. Therefore, although one study

showed that prolonged activity caused lateral diffusion of other receptors (GABA$_A$) in the AIS

of hippocampal neurons (Muir and Kittler, 2014), we hypothesize that the origin of these

structural changes was rather a vesicular insertion of new Nav1.6 channels into the distal AIS

minutes after the period of elevated activity. This is consistent with the results of Kuba et al.,

(2010), who showed that AIS elongation was accompanied by increased expression of Na$^+$

channels and enhanced axonal I$_{NaP}$ and membrane excitability in the auditory pathway.

Moreover, a computational model predicted that the distal AIS exhibits greater variation in

terms of Na$^+$ conductance and that small changes in the density of functional Na$^+$ channels

in this region would have a strong impact on neuronal excitability in L5PNs (Baranauskas et

al., 2013). We do not rule out the simultaneous contribution of other VGCCs, such as the

combination of T-type and/or L-type VGCCs, which were necessary for activity-dependent

AIS relocation (Grubb and Burrone, 2010; Kuba et al., 2015), or K$^+$ channels, such as Kv7.2,

whose activity-dependent expression can increase in the AIS, reducing the shunt current

around the AP threshold (Jamann et al., 2021) and enhancing the electrotonic isolation of the

distal AIS, a factor that had also been computationally predicted as a key modulator of

neuronal excitability (Baranauskas et al., 2013), or Kv2.1, whose phosphorylation,

localization in the AIS (non-canonical pathway), and function are regulated by neuronal

activity and can drastically affect excitability and intrinsic plasticity (Misonou et al., 2005;

Jensen et al., 2017). Even other interactions between key protein partners could have

contributed to this activity-driven intrinsic plasticity, such as the phosphorylation of casein

kinase 2 (CK2), which regulates the interaction between Na$^+$ channels and AnkG (Bréchet et

al., 2008), or the interaction of Nav1.6 with the microtubule-associated protein Map1b

(Mtap1b), which facilitates the trafficking of these channels to the cell surface (O'Brien et al.,

2012; Solé et al., 2019). Leveraging the opportunities offered by novel genome-editing

techniques and super-resolution microscopy (Willems et al., 2020; Liu et al., 2022), future

studies should investigate the exocytic machinery in the AIS in contexts of activity-

dependent intrinsic plasticity, perhaps starting with endogenous tagging of Nav1.6 in L5PNs.

AIS plasticity in critical periods

Brain plasticity driven by activity or experience is enhanced during critical periods. For over 40 years, the primary model for studying critical period plasticity has been the primary visual cortex (Wiesel and Hubel, 1963; Hensch, 2005). Specifically, the critical visual period in mice has been defined from eye opening (P13) to approximately P35. The P13-P19 phase is considered a precritical period, independent of visual experience; P19-P28 represents the critical period with the highest sensitivity to imbalances; and P28-P35 corresponds to the consolidation and closure of the critical period (Gordon and Stryker, 1996; Corlew et al., 2007). Interestingly, the most significant elongation and reduction in AIS development in the visual cortex occur approximately one week before the opening and closing of the critical period, respectively (Gutzmann et al., 2014). This structural refinement of the AIS may be necessary to prepare the cortical network for the subsequent initiation and termination of the critical period. However, what determines the duration of these plasticity windows? GABAergic inhibition has gathered substantial evidence as a key mechanism regulating the onset and duration of the critical period in the visual cortex (Hensch, 2004, 2005). However, it is not the only factor. A recent study demonstrated that astrocytic connexin 30 controls the timing of critical period closure in the visual cortex (Ribot et al., 2021), and remarkably, the reintroduction of immature cultured astrocytes reinstates a period of heightened plasticity in adult animals, resembling the critical visual period (Müller and Best, 1989). Moreover, in other brain regions such as the hippocampus and somatosensory cortex, astrocytes have also been implicated in the closure of the critical plasticity period (Falcón-Moya et al., 2020; Martínez-Gallego et al., 2022). While we have not explored the role of astrocytes in AIS plasticity beyond the visual critical period, it would be interesting to investigate whether astrocytic manipulation could reinstate a period of AIS plasticity in adulthood, when its length is presumed to be stable.

CONCLUSION

The million-dollar question is how astrocytes detect neuronal activity and, more importantly, how they determine when to release the appropriate gliotransmitters, specifically, S100β in this case, at the right time. As previously mentioned, astrocytes can detect neuronal activity through changes in the extracellular ion composition, including K^+, Ca^{2+}, and Na^+. In

particular, the sodium/calcium exchanger (NCX) is predominantly expressed in the distal processes and terminal feet of astrocytes and can be easily activated by an increase in astrocytic $[Na^+]_i$ driven by neuronal activity. Given that certain astrocytic processes can contact the AIS, it is tempting to speculate that intense neuronal activity increases $[Na^+]_i$ signals in astrocytes, allowing Ca^{2+} influx from the extracellular space through the NCX exchanger. In future experiments, it would be interesting to determine whether $[Na^+]_i$ and $[Ca^{2+}]_i$ transients in astrocytes are closely related to specific protocols of intrinsic plasticity.

MATERIAL AND METHODS

Experimental model

All experiments were conducted in accordance with the rules of the Canadian Institutes of Health Research and were approved by the Animal Care and Use Committee of the University of Montreal (Protocol #23-207). We crossed mice from the newly generated AnkG-GFP-lox knock-in line (B6;129SV-ank3tm1DCI/HD) (Thome et al., 2025) with Vglut2-cre mice (B6J.129S6(FVB)-Slc17a6tm2(cre)Lowl/Mwar). This allowed Cre-dependent expression of AnkG-GFP driven by the VGlut2 promoter, a genetic marker of glutamatergic signaling. VGlut2-AnkG-GFP were fed ad libitum and housed with a 12 h light/dark cycle. Male and female mice were used for experiments.

V1 slice preparation

Coronal slices (350 µm) containing layer 5 of the V1 were prepared from 14- to 30-day-old Vglut2-AnkG-GFP mice. Mice were anesthetized with isoflurane (Pharmaceutical Partners of Canada Inc., Richmond Hill, ON, Canada) and subsequently decapitated. Following decapitation, the skull was quickly removed, and the brain was transferred to modified artificial cerebrospinal fluid (aCSF, in mM: 3 KCl, 1.25 KH2PO4, 4 MgSO4, 26 NaHCO3, 10 dextrose, 0.2 CaCl2, 219 sucrose; pH 7.3-7.4, 300-320 mOsmol/kg) that was ice-cold (0-4°C) and oxygenated with a mixture of 95% O_2 and 5% CO_2. Brain slices were then sectioned using a VT1000S vibratome (Leica). After a 15-minute incubation at 30°C in a temperature-controlled water bath within a slice holder containing oxygenated artificial cerebrospinal fluid (aCSF, in mM: 124 NaCl, 3 KCl, 1.25 KH2PO4, 1.3 MgSO4, 26 NaHCO3, 10 dextrose, and

1.2 CaCl2; pH 7.3-7.4, 294-300 mOsmol/kg) bubbled with 95% O2 and 5% CO2, slices were

transferred to the recording chamber at room temperature. A recovery period of at least one

hour was allowed for the slices before recording.

Electrophysiology

Recordings were conducted at room temperature in a submerged chamber continuously

perfused with artificial CSF (in mM: 124 NaCl, 3 KCl, 1.25 KH2PO4, 1.3 MgSO4, 26 NaHCO3,

10 dextrose, and 1.2 CaCl2; pH 7.3-7.4, 294-300 mOsmol/kg) bubbled with 95% O2 and 5%

CO2. Patch microelectrodes were fabricated from borosilicate glass capillaries (1.5 mm

outside diameter, 1.12 mm inside diameter, World Precision Instruments) using a P-97 puller

(Sutter Instruments). For neuronal recordings, pipettes with resistance ranging from 6 to 12

MΩ were filled with an internal solution containing the following (in mM): 140 K-gluconate, 5

NaCl, 2 MgCl2, 10 HEPES, 0.5 EGTA, 2 Tris ATP salt, 0.4 Tris GTP salt, pH 7.2-7.3, 280-

300mOsmol/kg. To visualize neuronal and axonal morphologies during the experiment, Alexa

Fluor 647 was added to the internal solution. Layer 5 pyramidal neurons of the

extratelencephalic subtype were selected based on their morphological characteristics

(large soma and thick apical dendrite) and electrophysiological properties (burst spiking). All

recordings were performed using a Multiclamp 700A amplifier, Digidata 1550B interface

coupled to a computer equipped with pClamp 11 software (Molecular Devices, San Jose,

CA). After the establishment of a gigaseal, the membrane potential was held at -65 mV, and

the membrane patch was ruptured by suction. During the experiments, the changes in Vm

(1–6 mV) were corrected by imposing continuous current to maintain the membrane

potential constant at -65mV. Recordings were discarded if the resting membrane potential

shifted by >6 mV or if resting input resistance (measured with a 50pA hyperpolarizing pulse

lasting 50 ms, delivered every 10 seconds) varied >20%. Astrocytes were initially identified

morphologically after incubating the slices in the marker Sulforhodamine 101 (SR101) for 5

minutes, based on their size and shape. This identification was subsequently confirmed by

their electrophysiological characteristics, including a low resting membrane potential and

passive responses to negative and positive current injections. For experiments involving

astrocyte inactivation, BAPTA tetrapotassium salt (20 mM) was added to the internal solution

filling the recording pipette. All recordings were performed in the presence of antagonists

of N-methyl-d-aspartate (NMDA), AMPA/kainate, and $GABA_A$ receptors [2-amino-5-

phosphonovaleric acid (d-APV), 50 μM; 6-cyano-7-nitroquinoxalene-2,3-dione (CNQX), 20

μM; and Gabazine, 20 μM, respectively, which were all purchased from Tocris Biosciences

(Ellisville, Missouri, USA)].

Plasticity Protocol and Analysis

After achieving whole-cell access by rupturing the membrane with negative pressure, a

waiting period of 5 to 10 minutes was allowed for the cell to dialyze with the pipette solution

and for the Alexa 647 fluorescent dye to diffuse. Throughout the recording, intrinsic

excitability was assessed every 10 seconds using a small, constant-amplitude depolarizing

pulse (500 ms, 10–60 pA), which was selected to evoke 2–3 APs during the baseline period (5

minutes) and then maintained constant throughout the recording. To generate input-output

curves and determine the current threshold for AP generation, a series of depolarizing

current injections (0–100 pA in 10 pA increments) was applied before the baseline period and

at the end of the recording. Each series was delivered at least twice per cell, and responses

were averaged. The induction stimulus consisted of 500 ms depolarizing pulses (150–180 pA)

delivered 60 times at 4-second intervals, with an amplitude selected to evoke a sustained

firing rate of 18–30 Hz. Only neurons with a stable baseline, consistent recording parameters,

and a post-induction recording duration of at least 20 minutes were included in the analysis.

The degree of LTP-IE was calculated as the ratio of firing frequency during the last 10 minutes

of the post-induction recording to the baseline. Intra- and intercellular comparisons were

performed by extracting the average response per minute (6 pulses every 15 seconds), and

the mean firing rate during the 5-minute baseline period was used to normalize data before

and after induction. Control cells remained at rest during the induction stimulus period, and

analyses were conducted similarly in this condition.

Optogenetic Stimulation

Astrocytes peri-AIS in VGlut2-AnkG-GFP mice, which received bilateral intracerebral viral

administration at P0 of an AAV2/5-GFAP-hChR2(H134R)-mCherry (3.40E+12) expressing

GFAP-promoter-driven Channelrhodopsin-2 (ChR2) fused to the mCherry reporter (provided

by the CNP Viral Vector Core at the CERVO Research Center, RRID: SCR_016477), were

optogenetically stimulated (beginning two weeks after viral injection) using two lasers (440

nm and 488 nm) simultaneously in the SIM light path of an FV1000 microscope (Olympus).

The SIM scanner was used in square scanning mode to photoactivate small manually

delineated areas within astrocytes near the AIS of the recorded neurons. Optogenetic

stimulations were applied after the baseline period using 60 pulses of 2 s ON and 2 s OFF

(10–20% laser power / 8.6–14.9 µW for the 440 nm laser / 8.7–15.9 µW for the 488 nm laser).

To assess the efficacy of the stimulation, laser pulses were applied while recording the

electrophysiological activity of astrocytes. These recordings confirmed that the

photostimulation reliably evoked small membrane depolarizations, averaging 7.4 mV and

ranging between 6 and 9 mV.

Drugs

Chemicals used in this study were purchased from Sigma-Aldrich (Oakville, Ontario,

Canada), Tocris Biosciences (Ellisville, Missouri, USA) and Abcam (Cambridge, UK). The

following drugs were bath-applied using a syringe pump: D,L-2-amino-5-phosphonovaleric

acid (APV, 75 µM), CNQX (10 µM), SR 95531 hydrobromide (Gabazine, 20 µM) and Suramin

(100 µM). In some experiments, the Ca^{2+}-binding protein S100β (129 µM) (Produced in the

Department of Biochemistry of the Université de Montreal as described in Morquette et al.,

2015) or 1,2-bis(o-aminophenoxy) ethane-N,N,N',N'-tetraacetic acid tetrasodium salt

(BAPTA, 5 mM) were locally applied using glass micropipettes (tip diameter ~1 µm) with 2–20

psi pressure pulses for 5 minutes (Picospritzer III, Parker Instrumentation, Fairfield, NJ, USA).

Monoclonal anti-S100β antibodies (mouse anti-S100β, Sigma Aldrich #S2532 or rabbit anti-

S100β, Abcam ab56642), as well as their denatured form, were applied locally with large-tip

(10-20 µm) glass micropipettes carefully lowered near the AIS recorded neuron with 0.1-2 psi

pressure pulses lasting 5 minutes. All bath-applied chemicals were diluted in water at 100X

their final concentrations and further diluted through the perfusing aCSF at their final

concentrations.

Immunohistochemistry and confocal microscopy

For the immunohistochemistry against AnkG and NeuN, coronal brainstem slices (350 μm)

were prepared from 14 to 30-day-old VGlut2-AnkG-GFP mice using a vibratome VT 1000S

(Leica), immediately immersed in a solution of 4% (wt/vol) paraformaldehyde in PBS, and

kept for one hour at 4°C. The sections were rinsed 3 times for 10 minutes in PBS and

incubated in a blocking solution containing 0.3% Triton X-100 and 5% Normal Donkey Serum

(Jackson ImmunoResearch #017-000-121) in PBS for one hour at room temperature. The

sections were then rinsed 3 times for 10 minutes in PBS and incubated overnight at 4°C in a

mix of the primary antibodies (chicken anti-AnkG, EnCor Biotech #PCAC-ANK3, dilution

1:1000; mouse anti-NeuN, Millipore #MAB377, dilution 1:500). The following day, the slices

were rinsed 5 times for 10 minutes in PBS and incubated in the relevant secondary antibodies

mix (donkey anti-chicken Alexa Fluor 647, Jackson Immunoresearch #703-605-155, dilution

1:250; donkey anti-mouse Alexa Fluor 594, Jackson Immunoresearch #715-585-151, dilution

1:250) diluted in the blocking solution for 120 minutes in a dark chamber at room

temperature. The sections were then rinsed 4 times for 15 minutes in PBS and mounted on

ColorFrost Plus slides (Fisher Scientific, Ottawa, Ontario, Canada) using Fluoromount-G

(Southern Biotech, Birmingham, Alabama, USA). A similar procedure was used for the

immunohistochemistry against S100β, except that the slices were immersed in 4% (wt/vol)

paraformaldehyde in PBS overnight at 4°C. The staining was performed using mouse anti-

S100β (Sigma #S2532, 1:400) as the primary antibody and donkey anti-mouse Alexa Fluor

647 (Abcam #ab150107, 1:250) as the secondary antibody. The washing steps, blocking

conditions, and mounting procedure were identical to those described above. Slides imaging

was performed using either a Zeiss LSM-900 Confocal Airyscan microscope or an FV1000

confocal microscope (Olympus). AnkG-GFP was visualized in real-time during

electrophysiology experiments using the FV1000, operated with FV10-ASW imaging software

(Version 03.00). Z-stacks were acquired at a speed of 10 μs/pixel and a resolution of 800 ×

800 pixels, with 1 μm step sizes and a zoom range of 2 to 4×. To minimize phototoxicity and

photobleaching, AIS z-stacks were captured at two time points: 5 minutes before induction

or drug exposure, as applicable, and 30 minutes afterward. AIS length was measured

manually and by fluorescence intensity along a 1-pixel-wide line drawn from the soma down

the axon (1 pixel ≈ 0.206 μm), using Fiji software (ImageJ 1.50e, NIH, USA). Fluorescence

profiles were smoothed using a 10-point (~2 μm) sliding mean and normalized between 1

(maximum) and 0 (minimum). The AIS end was defined at a 33% fluorescence threshold

(method adapted from Grubb and Burrone, 2010). No difference was observed between the

two methods. In a subset of experiments (**Fig. S1**), images were acquired using a two-photon

microscope equipped with a polygon scanner (BliQ Photonics), enabling high-speed imaging

while minimizing phototoxicity from repeated AIS imaging during recording.

Quantification and statical analysis

Data are presented as mean ± standard error to the mean (SEM) and as proportions (%).

Statistical analyses were performed depending on the experimental design. All paired

comparisons before and after treatment were confirmed to follow a normal distribution

(Shapiro–Wilk test, $P>0.05$) and were analyzed using paired t-tests. Comparisons between

independent groups, for which normality could not be assumed, were performed using the

non-parametric Mann–Whitney U test. Statistical significance was set at *$P<0.05$, **$P<0.01$,

and ***$P<0.001$; NS indicates not significant. Data analysis was performed using SPSS (IBM

SPSS Statistics).

DECLARATIONS

<u>**Author contributions**</u>

R.S.G., Y.I and A.K wrote the paper. R.S.G and Y.I collected and analyzed the data and built

the figures. All authors contributed to the article and approved the submitted version.

<u>**Acknowledgements**</u>

We thank Dr. Dorly Verdier for critically reading a preliminary version of the manuscript and
members of A.K laboratories for comments on the manuscript and technical support. We are
especially grateful to our esteemed colleagues, Dr. James G. Omichinski and Dr. Haytham
Wahba, from the Department of Biochemistry at the Université de Montréal, for their
invaluable contribution to the production of the S100β protein, and to Dr. Paul Jenkins, from
Michigan Neuroscience Institute Affiliate, for generously providing the AnkG-GFP-lox mouse
line. The graphical abstract incorporates elements from SciDraw (Federico Claudi:
doi.org/10.5281/zenodo.3925905; Andrew Hardaway: 10.5281/zenodo.5348394; Roberta
Schellino: 10.5281/zenodo.10390020), an open-access platform for scientific illustrations.

Funding

This work was supported by the Canadian Institutes of Health Research (Grant # 197855),
Natural Sciences and Engineering Research Council of Canada (RGPIN/05255-2020), CIRCA
Post-doctoral fellowship to Y.I.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or
financial relationships that could be construed as a potential conflict of interest.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Preprint

This manuscript was posted on a preprint.

REFERENCES

Agrawal, N., Hamam, B. N., Magistretti, J., Alonso, A., and Ragsdale, D. S. (2001). Persistent sodium
channel activity mediates subthreshold membrane potential oscillations and low-threshold spikes
in rat entorhinal cortex layer V neurons. *Neuroscience* 102, 53–64. doi: 10.1016/S0306-
4522(00)00455-3

Akin, E. J., Solé, L., Dib-Hajj, S. D., Waxman, S. G., and Tamkun, M. M. (2015). Preferential Targeting
of Nav1.6 Voltage-Gated Na+ Channels to the Axon Initial Segment during Development. *PLOS ONE*
10, e0124397. doi: 10.1371/journal.pone.0124397

Andersson, M., Blomstrand, F., and Hanse, E. (2007). Astrocytes play a critical role in transient
heterosynaptic depression in the rat hippocampal CA1 region. *The Journal of Physiology* 585, 843–
852. doi: 10.1113/jphysiol.2007.142737

Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1998). Glutamate-dependent astrocyte
modulation of synaptic transmission between cultured hippocampal neurons. *European Journal of*
Neuroscience 10, 2129–2142. doi: 10.1046/j.1460-9568.1998.00221.x

Armstrong, C. M. (1999). Distinguishing surface effects of calcium ion from pore-occupancy effects
in Na+ channels. *Proceedings of the National Academy of Sciences* 96, 4158–4163. doi:
10.1073/pnas.96.7.4158

Armstrong, C. M., and Cota, G. (1999). Calcium block of Na+ channels and its effect on closing rate.
Proceedings of the National Academy of Sciences 96, 4154–4157. doi: 10.1073/pnas.96.7.4154

Bancroft, E. A., De La Mora, M., Pandey, G., Zarate, S. M., and Srinivasan, R. (2022). Extracellular
S100B inhibits A-type voltage-gated potassium currents and increases L-type voltage-gated calcium
channel activity in dopaminergic neurons. *Glia* 70, 2330–2347. doi: 10.1002/glia.24254

Baranauskas, G., David, Y., and Fleidervish, I. A. (2013). Spatial mismatch between the Na$^+$ flux and
spike initiation in axon initial segment. *Proceedings of the National Academy of Sciences* 110, 4051–
4056. doi: 10.1073/pnas.1215125110

Benitez, M. J., Retana, D., Ordoñez-Gutiérrez, L., Colmena, I., Goméz, M. J., Álvarez, R., et al. (2024).
Transcriptomic alterations in APP/PS1 mice astrocytes lead to early postnatal axon initial segment
structural changes. *Cell. Mol. Life Sci.* 81, 444. doi: 10.1007/s00018-024-05485-9

Bréchet, A., Fache, M.-P., Brachet, A., Ferracci, G., Baude, A., Irondelle, M., et al. (2008). Protein
kinase CK2 contributes to the organization of sodium channels in axonal membranes by regulating
their interactions with ankyrin G. *Journal of Cell Biology* 183, 1101–1114. doi:
10.1083/jcb.200805169

Campanac, E., and Debanne, D. (2007). Plasticity of neuronal excitability: Hebbian rules beyond the
synapse. *Arch Ital Biol* 145, 277–287.

Campanac, E., and Debanne, D. (2008). Spike timing-dependent plasticity: a learning rule for
dendritic integration in rat CA1 pyramidal neurons. *The Journal of Physiology* 586, 779–793. doi:
10.1113/jphysiol.2007.147017

Chand, A. N., Galliano, E., Chesters, R. A., and Grubb, M. S. (2015). A Distinct Subtype of
Dopaminergic Interneuron Displays Inverted Structural Plasticity at the Axon Initial Segment. *J.*
Neurosci. 35, 1573–1590. doi: 10.1523/JNEUROSCI.3515-14.2015

Corlew, R., Wang, Y., Ghermazien, H., Erisir, A., and Philpot, B. D. (2007). Developmental Switch in
the Contribution of Presynaptic and Postsynaptic NMDA Receptors to Long-Term Depression. *J.*
Neurosci. 27, 9835–9845. doi: 10.1523/JNEUROSCI.5494-06.2007

Cudmore, R. H., and Turrigiano, G. G. (2004). Long-Term Potentiation of Intrinsic Excitability in LV
Visual Cortical Neurons. *Journal of Neurophysiology* 92, 341–348. doi: 10.1152/jn.01059.2003

Debanne, D., Campanac, E., Bialowas, A., Carlier, E., and Alcaraz, G. (2011). Axon Physiology.
Physiological Reviews. doi: 10.1152/physrev.00048.2009

Debanne, D., Inglebert, Y., and Russier, M. (2019). Plasticity of intrinsic neuronal excitability. *Current
Opinion in Neurobiology* 54, 73–82. doi: 10.1016/j.conb.2018.09.001

Debanne, D., and Rama, S. (2011). Astrocytes Shape Axonal Signaling. *Science Signaling* 4, pe11–
pe11. doi: 10.1126/scisignal.2001884

Debanne, D., and Russier, M. (2019). The contribution of ion channels in input-output plasticity.
Neurobiology of Learning and Memory 166, 107095. doi: 10.1016/j.nlm.2019.107095

Donato, R., Sorci, G., Riuzzi, F., Arcuri, C., Bianchi, R., Brozzi, F., et al. (2009). S100B's double life:
Intracellular regulator and extracellular signal. *Biochimica et Biophysica Acta (BBA) - Molecular Cell
Research* 1793, 1008–1022. doi: 10.1016/j.bbamcr.2008.11.009

Drouillas, B., Brocard, C., Zanella, S., Bos, R., and Brocard, F. (2023). Persistent Nav1.1 and Nav1.6
currents drive spinal locomotor functions through nonlinear dynamics. *Cell Reports* 42, 113085.
doi: 10.1016/j.celrep.2023.113085

Duménieu, M., Fronzaroli-Molinieres, L., Naudin, L., Iborra-Bonnaure, C., Wakade, A., Zanin, E., et
al. (2025). Visual activity enhances neuronal excitability in thalamic relay neurons. *Science
Advances* 11, eadp4627. doi: 10.1126/sciadv.adp4627

Evans, M. D., Dumitrescu, A. S., Kruijssen, D. L. H., Taylor, S. E., and Grubb, M. S. (2015). Rapid
Modulation of Axon Initial Segment Length Influences Repetitive Spike Firing. *Cell Reports* 13, 1233–
1245. doi: 10.1016/j.celrep.2015.09.066

Evans, M. D., Sammons, R. P., Lebron, S., Dumitrescu, A. S., Watkins, T. B. K., Uebele, V. N., et al.
(2013). Calcineurin Signaling Mediates Activity-Dependent Relocation of the Axon Initial Segment. *J.
Neurosci.* 33, 6950–6963. doi: 10.1523/JNEUROSCI.0277-13.2013

Falcón-Moya, R., Pérez-Rodríguez, M., Prius-Mengual, J., Andrade-Talavera, Y., Arroyo-García, L. E.,
Pérez-Artés, R., et al. (2020). Astrocyte-mediated switch in spike timing-dependent plasticity during
hippocampal development. *Nat Commun* 11, 4388. doi: 10.1038/s41467-020-18024-4

Forsberg, M., Seth, H., Björefeldt, A., Lyckenvik, T., Andersson, M., Wasling, P., et al. (2019). Ionized
calcium in human cerebrospinal fluid and its influence on intrinsic and synaptic excitability of
hippocampal pyramidal neurons in the rat. *Journal of Neurochemistry* 149, 452–470. doi:
10.1111/jnc.14693

Forsberg, M., Zhou, D., Jalali, S., Faravelli, G., Seth, H., Björefeldt, A., et al. (2025). Evaluation of
mechanisms involved in regulation of intrinsic excitability by extracellular calcium in CA1 pyramidal
neurons of rat. *Journal of Neurochemistry* 169, e16209. doi: 10.1111/jnc.16209

Fréal, A., Jamann, N., Ten Bos, J., Jansen, J., Petersen, N., Ligthart, T., et al. (2023). Sodium channel
endocytosis drives axon initial segment plasticity. *Science Advances* 9, eadf3885. doi:
10.1126/sciadv.adf3885

Gaudel, F., Giraud, J., Morquette, P., Couillard-Larocque, M., Verdier, D., and Kolta, A. (2025).
Astrocyte-induced firing in primary afferent axons. *iScience* 0. doi: 10.1016/j.isci.2025.112006

Ghosh, A., Malavasi, E. L., Sherman, D. L., and Brophy, P. J. (2020). Neurofascin and Kv7.3 are
delivered to somatic and axon terminal surface membranes en route to the axon initial segment.
eLife 9, e60619. doi: 10.7554/eLife.60619

Gordon, J. A., and Stryker, M. P. (1996). Experience-Dependent Plasticity of Binocular Responses in
the Primary Visual Cortex of the Mouse. *J. Neurosci.* 16, 3274–3286. doi: 10.1523/JNEUROSCI.16-
10-03274.1996

Grubb, M. S., and Burrone, J. (2010). Activity-dependent relocation of the axon initial segment fine-
tunes neuronal excitability. *Nature* 465, 1070–1074. doi: 10.1038/nature09160

Grubb, M. S., Shu, Y., Kuba, H., Rasband, M. N., Wimmer, V. C., and Bender, K. J. (2011). Short- and
long-term plasticity at the axon initial segment. *J Neurosci* 31, 16049–16055. doi:
10.1523/JNEUROSCI.4064-11.2011

Gutzmann, A., Erguel, N., Grossmann, R., Schultz, C., Wahle, P., and Engelhardt, M. (2014). A period
of structural plasticity at the axon initial segment in developing visual cortex. *Front. Neuroanat.* 8.
doi: 10.3389/fnana.2014.00011

Hensch, T. K. (2004). Critical period regulation. *Annu Rev Neurosci* 27, 549–579. doi:
10.1146/annurev.neuro.27.070203.144327

Hensch, T. K. (2005). Critical period plasticity in local cortical circuits. *Nat Rev Neurosci* 6, 877–888.
doi: 10.1038/nrn1787

Hermann, A., Donato, R., Weiger, T. M., and Chazin, W. J. (2012). S100 Calcium Binding Proteins and
Ion Channels. *Front Pharmacol* 3, 67. doi: 10.3389/fphar.2012.00067

Höfflin, F., Jack, A., Riedel, C., Mack-Bucher, J., Roos, J., Corcelli, C., et al. (2017). Heterogeneity of
the Axon Initial Segment in Interneurons and Pyramidal Cells of Rodent Visual Cortex. *Front. Cell.*
Neurosci. 11. doi: 10.3389/fncel.2017.00332

Hu, W., Tian, C., Li, T., Yang, M., Hou, H., and Shu, Y. (2009). Distinct contributions of Nav1.6 and
Nav1.2 in action potential initiation and backpropagation. *Nat Neurosci* 12, 996–1002. doi:
10.1038/nn.2359

Inglebert, Y., Aljadeff, J., Brunel, N., and Debanne, D. (2020). Synaptic plasticity rules with
physiological calcium levels. *Proceedings of the National Academy of Sciences* 117, 33639–33648.
doi: 10.1073/pnas.2013663117

Jamann, N., Dannehl, D., Lehmann, N., Wagener, R., Thielemann, C., Schultz, C., et al. (2021).
Sensory input drives rapid homeostatic scaling of the axon initial segment in mouse barrel cortex.
Nat Commun 12, 23. doi: 10.1038/s41467-020-20232-x

Jensen, C. S., Watanabe, S., Stas, J. I., Klaphaak, J., Yamane, A., Schmitt, N., et al. (2017). Trafficking
of Kv2.1 Channels to the Axon Initial Segment by a Novel Nonconventional Secretory Pathway. *J.*
Neurosci. 37, 11523–11536. doi: 10.1523/JNEUROSCI.3510-16.2017

Kuba, H., Ishii, T. M., and Ohmori, H. (2006). Axonal site of spike initiation enhances auditory
coincidence detection. *Nature* 444, 1069–1072. doi: 10.1038/nature05347

Kuba, H., Oichi, Y., and Ohmori, H. (2010). Presynaptic activity regulates Na+ channel distribution at
the axon initial segment. *Nature* 465, 1075–1078. doi: 10.1038/nature09087

Kuba, H., Yamada, R., Ishiguro, G., and Adachi, R. (2015). Redistribution of Kv1 and Kv7 enhances
neuronal excitability during structural axon initial segment plasticity. *Nat Commun* 6, 8815. doi:
10.1038/ncomms9815

Leterrier, C. (2018). The Axon Initial Segment: An Updated Viewpoint. *J Neurosci* 38, 2135–2145. doi:
10.1523/JNEUROSCI.1922-17.2018

Lezmy, J., Arancibia-Cárcamo, I. L., Quintela-López, T., Sherman, D. L., Brophy, P. J., and Attwell, D.
(2021). Astrocyte Ca2+-evoked ATP release regulates myelinated axon excitability and conduction
speed. *Science* 374, eabh2858. doi: 10.1126/science.abh2858

Liu, H., Wang, H.-G., Pitt, G., and Liu, Z. (2022). Direct Observation of Compartment-Specific
Localization and Dynamics of Voltage-Gated Sodium Channels. *J Neurosci* 42, 5482–5498. doi:
10.1523/JNEUROSCI.0086-22.2022

Martínez-Gallego, I., Pérez-Rodríguez, M., Coatl-Cuaya, H., Flores, G., and Rodríguez-Moreno, A.
(2022). Adenosine and Astrocytes Determine the Developmental Dynamics of Spike Timing-
Dependent Plasticity in the Somatosensory Cortex. *J Neurosci* 42, 6038–6052. doi:
10.1523/JNEUROSCI.0115-22.2022

Misonou, H., Mohapatra, D. P., and Trimmer, J. S. (2005). Kv2.1: A Voltage-Gated K+ Channel Critical
to Dynamic Control of Neuronal Excitability. *NeuroToxicology* 26, 743–752. doi:
10.1016/j.neuro.2005.02.003

Morquette, P., Verdier, D., Kadala, A., Féthière, J., Philippe, A. G., Robitaille, R., et al. (2015). An
astrocyte-dependent mechanism for neuronal rhythmogenesis. *Nat Neurosci* 18, 844–854. doi:
10.1038/nn.4013

Muir, J., and Kittler, J. T. (2014). Plasticity of GABAA receptor diffusion dynamics at the axon initial
segment. *Front. Cell. Neurosci.* 8. doi: 10.3389/fncel.2014.00151

Müller, C. M., and Best, J. (1989). Ocular dominance plasticity in adult cat visual cortex after
transplantation of cultured astrocytes. *Nature* 342, 427–430. doi: 10.1038/342427a0

Nardin, P., Tortorelli, L., Quincozes-Santos, A., de Almeida, L. M. V., Leite, M. C., Thomazi, A. P., et al.
(2009). S100B Secretion in Acute Brain Slices: Modulation by Extracellular Levels of Ca2+ and K+.
Neurochem Res 34, 1603–1611. doi: 10.1007/s11064-009-9949-0

Nataraj, K., Le Roux, N., Nahmani, M., Lefort, S., and Turrigiano, G. (2010). Visual Deprivation
Suppresses L5 Pyramidal Neuron Excitability by Preventing the Induction of Intrinsic Plasticity.
Neuron 68, 750–762. doi: 10.1016/j.neuron.2010.09.033

Nelson, A. B., Krispel, C. M., Sekirnjak, C., and du Lac, S. (2003). Long-lasting increases in intrinsic
excitability triggered by inhibition. *Neuron* 40, 609–620. doi: 10.1016/s0896-6273(03)00641-x

Nishiyama, H., Knopfel, T., Endo, S., and Itohara, S. (2002). Glial protein S100B modulates long-term
neuronal synaptic plasticity. *Proc Natl Acad Sci U S A* 99, 4037–4042. doi: 10.1073/pnas.052020999

O'Brien, J. E., Sharkey, L. M., Vallianatos, C. N., Han, C., Blossom, J. C., Yu, T., et al. (2012).
Interaction of Voltage-gated Sodium Channel Nav1.6 (SCN8A) with Microtubule-associated Protein
Map1b *. *Journal of Biological Chemistry* 287, 18459–18466. doi: 10.1074/jbc.M111.336024

Palmer, L. M., and Stuart, G. J. (2006). Site of Action Potential Initiation in Layer 5 Pyramidal
Neurons. *J. Neurosci.* 26, 1854–1863. doi: 10.1523/JNEUROSCI.4812-05.2006

Ribot, J., Breton, R., Calvo, C.-F., Moulard, J., Ezan, P., Zapata, J., et al. (2021). Astrocytes close the
mouse critical period for visual plasticity. *Science* 373, 77–81. doi: 10.1126/science.abf5273

Ryczko, D., Hanini-Daoud, M., Condamine, S., Bréant, B. J. B., Fougère, M., Araya, R., et al. (2021).
S100β-mediated astroglial control of firing and input processing in layer 5 pyramidal neurons of the
mouse visual cortex. *The Journal of Physiology* 599, 677–707. doi: 10.1113/JP280501

Sanz-Gálvez, R., Falardeau, D., Kolta, A., and Inglebert, Y. (2024). The role of astrocytes from
synaptic to non-synaptic plasticity. *Front. Cell. Neurosci.* 18. doi: 10.3389/fncel.2024.1477985

Schlüter, A., Del Turco, D., Deller, T., Gutzmann, A., Schultz, C., and Engelhardt, M. (2017).
Structural Plasticity of Synaptopodin in the Axon Initial Segment during Visual Cortex Development.
Cerebral Cortex 27, 4662–4675. doi: 10.1093/cercor/bhx208

Solé, L., Wagnon, J. L., Akin, E. J., Meisler, M. H., and Tamkun, M. M. (2019). The MAP1B Binding
Domain of Nav1.6 Is Required for Stable Expression at the Axon Initial Segment. *J. Neurosci.* 39,
4238–4251. doi: 10.1523/JNEUROSCI.2771-18.2019

Sorci, G., Riuzzi, F., Arcuri, C., Bianchi, R., Brozzi, F., Tubaro, C., et al. (2010). The many faces of
S100B protein: when an extracellular factor inactivates its own receptor and activates another one.
Ital J Anat Embryol 115, 147–151.

Sourdet, V., Russier, M., Daoudal, G., Ankri, N., and Debanne, D. (2003). Long-term enhancement of
neuronal excitability and temporal fidelity mediated by metabotropic glutamate receptor subtype 5.
J Neurosci 23, 10238–10248. doi: 10.1523/JNEUROSCI.23-32-10238.2003

Thome, C., Janssen, J. M., Karabulut, S., Acuna, C., D'Este, E., Soyka, S. J., et al. (2025). Live imaging
of excitable axonal microdomains in ankyrin-G-GFP mice. *eLife* 12, RP87078. doi:
10.7554/eLife.87078

Treves, S., Scutari, E., Robert, M., Groh, S., Ottolia, M., Prestipino, G., et al. (1997). Interaction of
S100A1 with the Ca2+ Release Channel (Ryanodine Receptor) of Skeletal Muscle. *Biochemistry* 36,
11496–11503. doi: 10.1021/bi970160w

Van Wart, A., Trimmer, J. S., and Matthews, G. (2007). Polarized distribution of ion channels within
microdomains of the axon initial segment. *Journal of Comparative Neurology* 500, 339–352. doi:
10.1002/cne.21173

Watanabe, A., Guo, C., and Sjöström, P. J. (2023). The developmental profile of visual cortex
astrocytes. *iScience* 26, 106828. doi: 10.1016/j.isci.2023.106828

Wiesel, T. N., and Hubel, D. H. (1963). Single-cell responses in striate cortex of kittens deprived of
vision in one eye. *Journal of Neurophysiology* 26, 1003–1017. doi: 10.1152/jn.1963.26.6.1003

Willems, J., Jong, A. P. H. de, Scheefhals, N., Mertens, E., Catsburg, L. A. E., Poorthuis, R. B., et al.
(2020). ORANGE: A CRISPR/Cas9-based genome editing toolbox for epitope tagging of endogenous
proteins in neurons. *PLOS Biology* 18, e3000665. doi: 10.1371/journal.pbio.3000665

Wu, N., Enomoto, A., Tanaka, S., Hsiao, C.-F., Nykamp, D. Q., Izhikevich, E., et al. (2005). Persistent
Sodium Currents in Mesencephalic V Neurons Participate in Burst Generation and Control of
Membrane Excitability. *Journal of Neurophysiology* 93, 2710–2722. doi: 10.1152/jn.00636.2004

Xia, Z., and Storm, D. R. (2005). The role of calmodulin as a signal integrator for synaptic plasticity.
Nat Rev Neurosci 6, 267–276. doi: 10.1038/nrn1647

Xu, J., Kang, N., Jiang, L., Nedergaard, M., and Kang, J. (2005). Activity-Dependent Long-Term
Potentiation of Intrinsic Excitability in Hippocampal CA1 Pyramidal Neurons. *J. Neurosci.* 25, 1750–
1760. doi: 10.1523/JNEUROSCI.4217-04.2005

Yamada, R., and Kuba, H. (2016). Structural and Functional Plasticity at the Axon Initial Segment.
Front. Cell. Neurosci. 10. doi: 10.3389/fncel.2016.00250

Yang, Y., Ge, W., Chen, Y., Zhang, Z., Shen, W., Wu, C., et al. (2003). Contribution of astrocytes to
hippocampal long-term potentiation through release of d-serine. *Proceedings of the National
Academy of Sciences* 100, 15194–15199. doi: 10.1073/pnas.2431073100

FIGURE LEGENDS

Figure 1. LTP-IE is associated with rapid AIS plasticity

A. *Top*. High-resolution images showing strong co-localization of AnkG (red, *left*) and GFP

(green, *middle*) in pyramidal neurons (cyan, *right*) from layer 5 (L5PNs) of the visual cortex.

Bottom left. Photomicrography of the axon initial segment (AIS) (AnkG-GFP) in L5PNs filled

with Alexa 647. *Bottom right*. Membrane response of the whole-cell recorded neuron at the

soma during the induction protocol. **B.** *Left*. Time course of normalized spike number before

and after (representative traces shown above) induction (red) and in control conditions

(black). *Right*. Quantification of spike number changes in the last 10 minutes, with individual

values for each cell represented as red (induction) and black (control) open circles. **C.** *Left*.

Live image of an AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after

induction. Black arrowheads indicate the start and end of the AIS. *Middle*. Induction of

intrinsic excitability potentiation elongates AIS length (left, manual measurement of

individual cells; right, mean normalized fluorescence intensity profiles before (black) and 30

min after induction (red). Arrowheads indicate AIS end positions where the fluorescence

profile crosses the 33% threshold, showing AIS elongation). **D.** *Left*. Current-number of spike

relationships for each condition (before and after induction). Induction caused a leftward

shift in the mean curve. *Right*. Representative traces of individual current injections at

different amplitudes. Note the difference in the spike number with the same amplitude

before and after induction. **E.** Decrease in Rheobase after induction. **F.** *Left*. Live image of an

AnkG-GFP-positive neuron fixed with Alexa Fluor 647 before and after the control condition.

Black arrowheads indicate the start and end of the AIS. *Middle*. No AIS length change

conditions (left: manual measurements; right: fluorescence profiles before and after control,

with no shift at the 33% threshold). **G.** *Left*. Current-number of spike relationships for each

condition (before and after the control period). No shift in the mean curve was observed.

Right. Representative traces of individual current injections at different amplitudes in the

control condition, with no changes observed. **H.** No change in Rheobase in the control

condition.

Figure 2. Astrocytic inactivation prevents LTP-IE and AIS plasticity

A. Confocal imaging of Sulforhodamine (SR101) labeling in astrocytes of layer 5 of the visual

cortex. **B.** *Left*. Experimental paradigm. Stimulation was performed by current injection into

the whole-cell recorded neuron, after targeting a neighboring peri-AIS astrocyte for recording

with a BAPTA filled pipette for its inactivation. *Middle*. Time course of normalized spike

number before and after induction (red) and during peri-AIS astrocyte network inactivation

(green), with representative traces shown above. *Right*. Quantification of spike number

changes in the last 10 minutes, with individual values for each cell represented as red

(induction) and green (astrocytes inactivated) open circles and distance of the inactivated

astrocytes from the soma of the recorded neuron. **C.** Live image of an AnkG-GFP-positive

neuron filled with Alexa Fluor 647 before and after induction during peri-AIS astrocyte

network inactivation, filled with Alexa Fluor 594. Black arrowheads indicate the start and end

of the AIS. *Right.* Peri-AIS astrocyte inactivation prevented the change in AIS length. **D.** *Left.*

Current-number of spike relationships for each condition (before and after induction during

peri-AIS astrocyte network inactivation). No shift in the mean curve was observed. *Right.*

Representative traces of individual current injections at different amplitudes, with no

changes observed. **E.** No change in Rheobase in the recorded neuron during peri-AIS

astrocyte network inactivation.

Figure 3. Blockade of the astrocytic protein S100β impairs LTP-IE and AIS plasticity

A. *Left.* Experimental paradigm. The induction protocol was performed by current injection

into the whole-cell recorded neuron 5 minutes after a local application of either an S100β

antibody or its denatured form. *Middle.* Time course of normalized spike number before and

after induction, with prior application of S100β antibody (purple) and its denatured form

(pink), with representative traces shown above. *Right.* Quantification of spike number

changes in the last 10 minutes, with individual values for each cell represented as purple

(S100β antibody) and pink (S100β antibody denatured) open circles. **B.** *Left*. Live image of an

AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after the application of

S100β antibody, followed by the induction protocol. Black arrowheads indicate the start and

end of the AIS. *Right.* The application of the S100β antibody, prior to induction, prevents the

change in AIS length. **C.** *Left.* Current-number of spike relationships for each condition

(before and after S100β antibody application, followed by the induction protocol). Induction

caused a rightward shift in the mean curve. *Right.* Representative traces of individual current

injections at different amplitudes. Note the difference in the spike number with the same

amplitude before and after induction with prior application of S100β antibody. **D.** No change

in Rheobase in the recorded neuron with the application of S100β antibody prior to induction.

E. *Left*. Live image of an AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after the application of S100β antibody denatured, followed by the induction protocol. *Right*. The application of denatured S100β antibody prior to induction does not prevent the change in AIS length. **F.** *Left*. Current-number of spike relationships for each condition (before and after S100β antibody denatured application, followed by the induction protocol). Induction caused a leftward shift in the mean curve. *Right*. Representative traces of individual current injections at different amplitudes. Note the difference in the spike number with the same amplitude before and after induction with prior application of S100β antibody denatured. **G.** No change in Rheobase in the recorded neuron with the application of S100β antibody denatured prior to induction.

Figure 4. Optogenetic activation of peri-AIS astrocyte reproduces LTP-IE and AIS plasticity

A. *Left*. Schematic representation of the strategy used for the optogenetic activation of astrocytes in layer 5 of the cortex visual. *Right*. Confocal imaging of immunohistochemistry-confirmed AAV GFAP-ChR2 (cyan) expression in astrocytes (S100β, magenta) in AnkG-GFP-positive mice (green). **B.** *Left*. Photomicrography of the AIS (AnkG-GFP) in an L5PN filled with Alexa 647 alongside an astrocyte expressing AAV GFAP-ChR2. **B.** *Middle*. Time course of normalized spike number before and after optogenetic activation of astrocytes, with (purple) or without (cyan) S100β antibody application, with representative traces above. *Right*. Quantification of spike number changes in the last 10 minutes, with individual values for each cell represented as cyan (optogenetic activation of astrocytes), purple (S100β antibody followed by optogenetic activation of astrocytes). **C.** *Left*. Live image of an AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after optogenetic activation of an astrocyte expressing ChR2 (cyan). Black arrowheads indicate the start and end of the AIS. *Right*. The optogenetic activation of astrocytes significantly elongates the length of the AIS. **D.** *Left*. Live image of an AnkG-GFP-positive neuron fixed with Alexa Fluor 647 before and after optogenetic activation of a ChR2-expressing astrocyte (cyan), with prior S100β antibody

application. Black arrowheads indicate the start and end of the AIS. *Right*. The application of the S100β antibody, prior to optogenetic peri-AIS astrocyte activation, prevents changes in AIS length. **E.** *Left*. Current-number of spike relationships for each condition (before and after optogenetic activation of astrocytes). Optogenetic activation caused a leftward shift in the mean curve. *Top*. Representative traces of individual current injections at different amplitudes. Note the difference in the spike number with the same amplitude before and after optogenetic activation of astrocytes. *Right*. Change in Rheobase in the recorded neuron during optogenetic peri-AIS astrocyte activation. **F.** *Left*. Current-number of spike relationships for each condition (before and after optogenetic astrocyte activation with prior S100β antibody application). *Top*. Representative traces of individual current injections at different amplitudes, with no changes observed. *Right*. No change in Rheobase in the recorded neuron during optogenetic peri-AIS astrocyte activation with prior S100β antibody application.

Figure 5. S100β rescues LTP-IE and AIS plasticity

A. *Left*. Experimental paradigm. Stimulation was performed by current injection into the whole-cell recorded neuron, with BAPTA-filled peri-AIS astrocyte inactivation and S100β protein applied 5 minutes before induction. *Middle*. Time course of normalized spike number before and after induction during peri-AIS astrocyte network inactivation (green) and following S100β application (orange), with representative traces shown above. *Right*. Quantification of spike number changes in the last 10 minutes, with individual values for each cell represented as green (astrocytes inactivated) and orange (astrocytes inactivated with prior S100β application) open circles. **B.** *Left*. Live image of an AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after induction during peri-AIS astrocyte (filled with Alexa Fluor 594) network inactivation, with prior S100β application. Black arrowheads indicate the start and end of the AIS. *Right*. S100β rescues the elongation of AIS length. **C.** *Left*. Current-number of spike relationships for each condition (before and after induction, during peri-AIS astrocyte network inactivation, with prior S100β application). Induction

caused a leftward shift in the mean curve. *Right*. Representative traces of individual current

injections at different amplitudes. Note the difference in the spike number with the same

amplitude before and after induction, during peri-AIS astrocyte network inactivation, with

prior S100β application. **D.** Change in Rheobase in the recorded neuron during peri-AIS

astrocyte network inactivation with prior S100β application.

Supplementary Figure 1. AIS plasticity linked to LTP-IE emerges within minutes post-

induction

Left. Time course of normalized spike count before and after induction (red), with

representative traces shown above. Time course of normalized AIS length (green) measured

every 5 minutes post-induction. Note the immediate change in excitability following

induction and the increase in AIS length occurring 10–15 minutes later. *Right*. Individual AIS

length measurements over time, with light gray representing individual cells and red

indicating the average. Example images are shown at baseline (0 min, before) and 30 minutes

(after) post-induction.

Supplementary Figure 2. Representation of the astrocyte optogenetic stimulation

protocol

A. Photomicrograph of a patched astrocyte in whole-cell configuration, filled with Alexa Fluor

594 and expressing ChR2 (cyan) for the data presented in **B**. The ROI drawing represents the

optogenetic stimulation area. **B.** *Top*. Membrane responses of the astrocyte in **A** to

optogenetic stimulation (ROI), representing the complete stimulation protocol used in our

experiments with VGlut2-AnkG-GFP mice injected with AAV GFAP-ChR2. *Bottom*. Expansion

of a part of the stimulation protocol clearly showing how each scanning episode (2s) triggers

a membrane depolarization.

Supplementary Figure 3. ATP/adenosine inhibition does not affect LTP-IE or AIS plasticity

A. *Left*. Experimental paradigm. Stimulation was performed by optogenetic activation of an astrocyte near the AIS of the recorded whole-cell neuron, after a 5-minute application of the ATP/adenosine antagonist Suramin. *Middle*. Time course of normalized spike number before and after induction during peri-AIS astrocyte optogenetic activation in the presence of Suramin (blue), with representative traces shown above. *Right*. Quantification of spike number changes in the last 10 minutes, with individual values for each cell represented as open blue circles (optogenetic activation of astrocytes in the presence of Suramin). **B.** *Left*. Live image of an AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after optogenetic activation of an astrocyte expressing ChR2 (cyan), with Suramin present. Black arrowheads indicate the start and end of the AIS. *Right:* Suramin does not affect the elongation of AIS length. **C.** *Left*. Current-number of spike relationships for each condition (before and after optogenetic activation of astrocytes, under Suramin exposure). Optogenetic activation caused a leftward shift in the mean curve. *Right*. Representative traces of individual current injections at different amplitudes. Note the difference in the spike number with the same amplitude before and after optogenetic activation of astrocytes, while Suramin was applied. **D.** No change in Rheobase in the recorded neuron during optogenetic activation of peri-AIS astrocyte network in the presence of Suramin.

Supplementary Figure 4. BAPTA extracellular rescues LTP-IE and AIS plasticity

A. *Left*. Experimental paradigm. Stimulation was performed by current injection into the whole-cell recorded neuron, with BAPTA-filled peri-AIS astrocyte for inactivation of the astrocytic syncytium and local extracellular application of BAPTA to mimic the Ca2+-chelating action of S100β, 5 minutes before induction. *Middle*. Time course of normalized spike number before and after induction during peri-AIS astrocyte network inactivation (green) and following extracellular BAPTA application (brown), with representative traces shown above. *Right*. Quantification of spike number changes in the last 10 minutes, with

individual values for each cell represented as green (astrocytes inactivated) and brown

(astrocytes inactivated with prior BAPTA extracellular application) open circles. **B.** Live image

of an AnkG-GFP-positive neuron filled with Alexa Fluor 647 before and after induction during

peri-AIS astrocyte (filled with Alexa Fluor 594) network inactivation, with prior extracellular

BAPTA application. Black arrowheads indicate the start and end of the AIS. *Right*. BAPTA

extracellular rescues the elongation of AIS length. **C.** *Left*. Current-number of spike

relationships for each condition (before and after induction, during peri-AIS astrocyte

network inactivation, with prior extracellular BAPTA application). Induction caused a leftward

shift in the mean curve. *Right*. Representative traces of individual current injections at

different amplitudes. Note the difference in the spike number with the same amplitude

before and after induction, with prior BAPTA extracellular application. **D.** No change in

Rheobase in the recorded neuron during peri-AIS astrocyte network inactivation with prior

BAPTA extracellular application.

FIGURE 1

FIGURE 2

A SR101 50µm

B Before / After
20mV | 0.1s
20pA
BAPTA
I stim
ASTROCYTE   L5PN

Number of Spikes (Normalized)
Induction
Control (n=6)
BAPTA Astro (n=5)
Time (min)

**
Number of Spikes (%Ctrl)
Ctrl   BAPTA Astro

Distance from AIS (µm)

C Confocal AnkG-GFP
Before After AF647 AF594
10µm
ns
Length AIS (µm)
Before After

D Before / After
all steps; ns
Number of Spike
Current (pA)
-67 mV +30pA
I m Before
0.2s | 20mV
40pA
-68 mV +30pA
I m After

E ns
Rheobase (pA)
Before After

FIGURE 3

FIGURE 4

FIGURE 5

A

Before   After

20mV | 0.1s
20pA

ASTROCYTE   BAPTA   I$_{stim}$
L5PN

S100β

Number of Spikes (Normalized)

S100β   Induction

○ BAPTA Astro (n=5)
○ BAPTA Astro + S100β (n=6)

Time (min)

**

Number of Spikes (%Ctrl)

BAPTA
Astro

BAPTA
Astro
+
S100β

B

Confocal AnkG-GFP

Before   After

AF647 AF594   10μm

**

Length AIS (μm)

Before   After

C

**

Number of Spike

■ Before
■ After

Current (pA)

-66mV   +40pA
I m   Before

-65mV   +40pA
I m   After

0.2s | 20mV
40pA

D

*

Rheobase (pA)

Before   After

Figure S1

FIGURE S2

A

B

FIGURE S3

FIGURE S4

A

ASTROCYTE BAPTA I_stim L5PN BAPTA

Before   After

20mV | 0.1s
20pA

Number of Spikes (Normalized)

BAPTA Extracell Induction

○ BAPTA Astro (n=5)
○ BAPTA Astro + BAPTA Extracell (n=5)

Time (min)

Number of Spikes (%Ctrl)

**

BAPTA Astro   BAPTA Astro + BAPTA Extracell

B

Confocal AnkG-GFP

Before After

AF647 AF594 10µm

Length AIS (µm)

**

Before After

C

Number of Spike

Before
After

Current (pA)

-66mV +40pA
I m Before

0.2s | 20mV
40pA

-67mV +40pA
I m After

D

Rheobase (pA)

ns

Before After

Graphical abstract

V1
Pyramidal neuron

L5PN
Astrocyte Peri-AIS
S100β

Rheobase

Number of Spike
0 min
30 min
Current (pA)

AIS

Brief period of
high-frequency
activation

Rheobase

Number of Spike
0 min
30 min
Current (pA)

Longer AIS

Normal neuronal activity

Increased intrinsic excitability