Cellulose Synthase Complex and Remorin Nanodomains Mediate Stress Resilience Through Cell Wall-Plasma Membrane Attachments

Authors: Yue Rui[1,*], Andres V. Reyes[2,3], Tarabryn S. Grismer[2,3], Nikolaj B. Abel[4], William P. Dwyer[1], Thomas Ott[5,6], Joseph J. Kieber[7], Shou-Ling Xu[2,3], and José R. Dinneny[1,8,*]

Affiliations:
[1]Department of Biology, Stanford University, Stanford, CA 94305, USA
[2]Division of Biosphere Science and Engineering, Carnegie Institution for Science, Stanford, CA 94305, USA
[3]Carnegie Mass Spectrometry Facility, Carnegie Institution for Science, Stanford, CA 94305, USA
[4]Department of Molecular Biology and Genetics, Aarhus University, Aarhus, 8000, Denmark
[5]Faculty of Biology, University of Freiburg, 79104 Freiburg, Germany
[6]CIBSS - Centre for Integrative Biological Signalling Studies, University of Freiburg, 79104 Freiburg, Germany
[7]Department of Biology, University of North Carolina, Chapel Hill, NC 27599, USA
[8]Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305, USA
*Correspondence: yuerui@stanford.edu (Y.R.), dinneny@stanford.edu (J.R.D.)

Highlights

- The plasma membrane forms attachments to the plant cell wall that are revealed by hyperosmotic shock and correlate to the tolerance of root growth to stress.
- Cellulose Synthase Complex (CSC) and REMORIN (REM) nanodomains localize to cell wall-plasma membrane attachment sites.
- CSC density at the plasma membrane determines the extent of cell wall-plasma membrane attachment under hyperosmotic stress.
- REM nanodomains form rapidly under hyperosmotic stress and harbor the CSC exocytosis inhibitors SHOU4/4L, which limit CSC density at the plasma membrane.

Summary

The outer cell surface of an organism is the frontline for detecting and responding to environmental stimuli. In plants, this interface consists of the plasma membrane that lies beneath the cell wall and remains associated with it through attachment sites. These wall-membrane attachments become evident upon hyperosmotic shock, when severe water loss causes the membrane to retract from the wall. Despite their long-standing observation, the molecular identity and function of these attachments remain poorly understood. Here, we identified two nanodomain-mediated mechanisms governing wall-membrane attachments: one dependent on the Cellulose Synthase Complex (CSC), whose density at the plasma membrane positively correlates with resistance to hyperosmotic stress, and the other on REMORIN (REM), which acts antagonistically to the CSC mechanism. Using proximity-labeling proteomics, we identified SHOU4/4L as REM-associated proteins that mediate this antagonism. Together, our

45   findings reveal how membrane nanodomains pattern wall-membrane attachments to mediate
46   plant cell resilience under water stress.
47
48   **Introduction**
49
Living organisms must sense and respond to changing environments for their survival. The
outer cell surface represents the frontline in this process and comprises an extracellular matrix
that surrounds the cell and interfaces with the plasma membrane, where signaling is integrated
and communicated to the cell through receptors and other membrane-associated proteins. A
fundamental understanding of the organization, dynamics, and composition of this outer cell
surface will shed light on the molecular underpinnings of stress perception and acclimation. In
animal cells, the plasma membrane is anchored to the extracellular matrix by integrins,[1]
transmembrane receptors that serve as structural and signaling hubs for sensing and
transmitting mechanochemical cues at the cell surface.[2] Integrins have not been found in
plants,[3,4] however, and it remains unclear what molecules in the cell wall and the plasma
membrane mediate their attachment, how the cell wall-plasma membrane interface is
maintained and remodeled during environmental stress, and how signals are perceived at this
interface in response to changes in internal (e.g., turgor pressure) or external cues (e.g.,
chemical or mechanical stimuli).
64
The plant cell wall is a polysaccharide-based extracellular matrix comprised of cellulose
microfibrils embedded in a molecular network composed of hemicelluloses, pectins, and
structural proteins.[5] Plant cell walls play key roles in defining cell shape, providing structural
support, and mediating cellular signaling.[6,7] The cell wall constrains the lateral diffusion of
plasma membrane proteins[8] and regulates the size and dynamics of membrane nanodomains,[9]
which are submicron molecular assemblies with diverse protein and lipid compositions.[10,11]
Much of our understanding of cell wall function has come from studies in *Arabidopsis thaliana*,
particularly through the characterization of mutants impaired in the synthesis, modification, or
degradation of specific wall components.[5] A number of studies have demonstrated the effects
that environmental stress has on cell wall properties, and the physiological impact that
disruptions in cell wall composition have on the ability of plants to acclimate to such stresses.[7,12]
However, whether specific wall constituents play a role in mediating wall-membrane
attachments has not been determined.
78
Under non-stress conditions, turgor pressure appresses the plasma membrane against the cell
wall (Figure 1). However, under hyperosmotic stress, cellular water loss can be so severe that
plasmolysis ensues, when the plasma membrane retracts from the cell wall (Figure 1C-1E,
Video S1). Notably, during plasmolysis, portions of the plasma membrane remain attached to
the cell wall, forming thin connecting membrane filaments known as Hechtian strands, first
described by Hecht over a century ago.[13] These strands are part of a larger network collectively
referred to as the Hechtian structure,[14] which includes Hechtian strands, the Hechtian reticulum,
and wall-membrane attachment sites. The attachment sites, located at the distal ends of
Hechtian structures (Figure 1D and 1E, Video S1) are reminiscent of focal adhesions in
mammalian cells,[15] yet their molecular identity and function remain largely unknown.[3,4] Some

Hechtian structures are likely attached to the wall via plasmodesmata,[14] membrane-lined
intercellular channels that mediate transport and communication between plant cells. However,
the presence of wall-membrane attachment sites at the outer surface of epidermal cells[3] and in
guard cells,[16] both of which lack plasmodesmata, suggests alternative mechanisms for
connecting the cell wall and the plasma membrane.
94
The plasma membrane is a lipid bilayer enriched with diverse lipid species and proteins, and it
exhibits heterogeneity in its physicochemical properties.[17] Importantly, many membrane proteins
are organized into nanodomains to fulfill their functions in plant growth, abiotic stress responses,
and biotic interactions.[10,11] For example, Cellulose Synthase Complexes (CSCs) form discrete
puncta at the plasma membrane[18] and are considered nanodomains by microscopy-based
definitions.[10] CSCs have been proposed to anchor the plasma membrane to the cell wall in
plasmolyzed plant cells through the nascently synthesized cellulose microfibrils that become
embedded into the wall matrix,[19] yet this idea remains experimentally untested. Hyperosmotic
stress can also induce the formation of nanodomains, which was recently demonstrated for
ROP6, a Rho GTPase that interacts with NADP oxidases to produce reactive oxygen species
important for osmotic signaling.[20] In the context of biotic interactions, REMORINs (REMs) are a
well-characterized protein family known to organize into nanodomains.[21] For instance, the
symbiosis-specific remorin SYMREM1 in *Medicago truncatula* is essential for stabilizing the
symbiotic receptor-like kinase LYK3[22] and maintaining membrane topology during infection.[23]
Interestingly, *in vitro* studies have shown that REMs can bind oligo- and polygalacturonides,[24]
components of pectins in the cell wall. However, whether REMs contribute to physical or
functional associations between the cell wall and plasma membrane is unknown.
112
Plant roots absorb water and nutrients from the environment, and their growth is highly sensitive
to environmental fluctuations. Here, we leveraged the effects of hyperosmotic stress to
investigate the structure and function of the cell wall-plasma membrane interface in roots. We
demonstrate that maintaining CSC density at the plasma membrane is essential for wall-
membrane attachments and sustained root growth under hyperosmotic stress conditions. We
show that wall-membrane attachment sites contain two populations of nanodomains, CSCs and
REMs, respectively. Genetic, biochemical, and co-localization studies identified SHOU4/4L,
known negative regulators of CSC exocytosis,[25] as being associated with REM nanodomains
that limit CSC density at the plasma membrane. Our data support a one-way antagonistic
mechanism where REM nanodomains restrict the abundance of CSC nanodomains. CSCs and
REMs function together at the plasma membrane to regulate wall-membrane attachments,
which are critical for maintaining cellular resilience under water deficit conditions.
125
126 **Results**
127
128 *Impaired cellulose and rhamnose levels have contrasting effects on root growth under*
129 *hyperosmotic stress*
130
To investigate the roles of cell wall polysaccharides in hyperosmotic stress response, we
surveyed a collection of previously published Arabidopsis mutants deficient in cellulose,

xyloglucan (a hemicellulose), pectic homogalacturonan (HG), rhamnogalacturonan-I (RG-I), or
rhamnogalacturonan-II (RG-II), respectively, for changes in root growth (Figure S1A, Figure 2).
Under hyperosmotic stress induced by 0.28 M sorbitol, wild-type roots exhibited a growth
reduction of 52-57% compared to the control conditions (Figure 2, Figure S1B). Mutants lacking
cellulose, including *cesa3^{je5}*, *cesa6^{prc1-1}*, and *cob-1*, all displayed a more severe growth
reduction (≥75%) (Figure 2A). Expression of *GFP-CESA3* under its native promoter in the
cesa3^{je5} mutant background restored root growth to wild-type levels (Figure 2A).
140
141 Mutants defective in xyloglucan (Figure 2B), HG (Figure 2C), RG-I acetylation (Figure 2D), or
142 RG-II crosslinking (Figure 2E) generally showed wild-type responses to stress. The only
143 exception was *qua2-1*, which exhibited a significantly greater reduction in root growth compared
144 to wild type under stress (Figure 2C), though this may be influenced by its known defects in cell-
145 cell adhesion.[26]
146
In contrast to the cellulose mutants, the *rhm1-1*, *rhm1-2*, and *rhm1-3* mutants, which are
deficient in rhamnose, a component of the RG-I backbone,[27] showed significantly less reduction
in root growth under hyperosmotic stress than the wild type (Figure 2D and Figure S1B). This
phenotype was reversed by expressing *RHM1-GFP* under its native promoter (Figure 2D).
Rhamnose in RG-I can be substituted with side chains, including α-1,5-arabinan and β-1,4-
galactan. While a deficiency in galactan, as in the *gals1/2/3* mutants,[28] partially mirrored the
rhm1 mutant's response, a deficiency in arabinan, as in the *mur4-1* mutant, did not show a
similar effect (Figure 2D). Taken together, our mutant survey suggests specific but contrasting
roles of cellulose and RG-I rhamnose in root growth responses to hyperosmotic stress.
156
157 *Deficiency in cellulose or rhamnose has opposite effects on plasmolysis and baseline plasma*
158 *membrane proteome*
159
To further understand the functions of cellulose and rhamnose in hyperosmotic stress response
at the cellular and molecular level, we performed live-cell imaging of five-day-old seedlings
expressing YFP-LTI6b, a classic plasma-membrane marker, after a brief hyperosmotic stress
treatment. The 0.28 M sorbitol concentration used in our root growth assay (Figure 2) did not
cause any noticeable changes in root cells compared to the control condition (Figure S1C).
However, treatment with 0.5 M sorbitol induced plasmolysis, with the plasma membrane
detaching from the cell wall at cell-cell junctions along the root's longitudinal axis (Figure S1C).
Increasing the sorbitol concentration to 1 M caused a more pronounced plasmolysis effect, with
the membrane also retracting from the outer periphery of root cells. Hechtian structures and cell
wall-plasma membrane attachment sites were clearly visible at these membrane retraction sites
(Figure S1C, Figure 1C-1E).
171
We selected 0.5 M sorbitol to compare plasmolysis in the wild type, cellulose-deficient, and
rhamnose-deficient mutants, as this concentration is less stressful than 1 M sorbitol and allows
for easier quantification of plasmolysis by area. The ratio of protoplast area (highlighted by YFP-
LTI6b) to cell area (outlined by the cell wall in the bright field channel) was used as a measure

of plasmolysis. Compared to wild-type cells, plasmolysis was more severe in the *cesa3^{je5}*
mutant, with a greater proportion of cells displaying lower ratios between 0.6-0.8 (26% in
cesa3^{je5} vs. 12% in wild type) (Figure 3A and 3B). In contrast, 70% of the *rhm1-1* cells exhibited
minimal plasmolysis at a ratio between 0.9-1.0, whereas this proportion in the wild type is 34%
(Figure 3A and 3B). These cellular-level plasmolysis results correlate with the severity with
which root growth was inhibited in these mutants (Figure 2A and 2D).
182
To identify membrane protein signatures underlying the distinctive hyperosmotic stress
responses of *cesa3^{je5}* and *rhm1-1* mutants, we used a biotin ligase-based proximity labeling
approach[29,30] by expressing a TurboID (TbID)-YFP-LTI6b fusion protein in the respective
genetic backgrounds. After confirming that TbID-YFP-LTI6b localized to the plasma membrane
(Figure S2A), we treated five-day-old seedlings with 50 μM biotin for 15 min (Figure 3C), which
was sufficient to biotinylate proteins beyond the endogenously biotinylated ones, as confirmed
by western blotting (Figure S2B and S2C). Following protein extraction, streptavidin pull-down of
the biotinylated proteins, tryptic digestion, and mass spectrometry, we conducted pairwise
protein enrichment analyses between each mutant and the wild type (Figure S2D and S2E).
192
In the *cesa3^{je5}* mutant, only one protein exhibited differential labeling: TOM1-LIKE1 (TOL1),[31,32]
which was significantly less enriched compared to wild type (Figure 3D). In contrast, over a
thousand proteins showed differential enrichment in the *rhm1-1* mutant relative to wild type
(Figure 3E). Among the most enriched proteins were CESA1, CESA5, CESA6, CELLULOSE
SYNTHASE-INTERACTIVE1 (CSI1),[33,34] CSI3,[35] PATROL1,[36] and CMU1[37] (Figure 3E), which
are involved in cellulose synthesis, CSC delivery to the plasma membrane, or microtubule
stability to guide CSC movement. These differences in enrichment were not caused by changes
in LTI6b levels, which were consistent across all transgenic lines. These data suggest that
defects in RG-I rhamnose biosynthesis lead to an increase in the abundance of CSC-associated
proteins at the plasma membrane.
203
To directly measure CSC density at the plasma membrane in root cells and assess its changes
due to deficiencies in cellulose or rhamnose, we introduced the GFP-CESA3 marker, driven by
its native promoter, into the wild-type, *cesa6^{prc1-1}*, and *rhm1-1* backgrounds. Similar to earlier
reports in etiolated hypocotyls,[18] CSCs were observed at the plasma membrane as particles
near or below the resolution limit of the spinning disk confocal microscope, and in Golgi as
bright "disks" larger than the membrane-localized particles (Figure 3F). Compared to the wild
type, GFP-CESA3-labeled CSC particles in *cesa6^{prc1-1}* roots were barely detectable at the
plasma membrane, though Golgi-localized CSCs remained visible (Figure 3F and 3G). In
contrast, *rhm1-1* roots exhibited a significantly higher CSC particle density at the plasma
membrane than wild-type roots (Figure 3F and 3G), consistent with our proteomic
characterization (Figure 3E).
215
216 *CSCs mediate cell wall-plasma membrane attachments under hyperosmotic stress*
217
Our results thus far point to a possible link between CSC density at the plasma membrane and
the severity of plasmolysis during hyperosmotic stress, suggesting that CSC may be directly or

indirectly involved in promoting wall-membrane attachments. To test this hypothesis, we
crossed the tdTomato-CESA6 marker with GFP-LTI6b and imaged root cells using spinning disk
confocal microscopy. Under control conditions, tdTomato-CESA6-labeled CSCs appeared as
particles that move along linear trajectories at the plasma membrane (Figure 4A, Figure S3A,
Video S2), consistent with previous observations in etiolated hypocotyls.[18] Treatment with 1 M
sorbitol immobilized CSCs into nanodomains at the plasma membrane (Figure S3A, Video S3),
which showed an increased degree of signal overlap with GFP-LTI6b-labeled wall-membrane
attachment sites compared to the control condition (Figure 4A and 4B, Video S4). Because
cortical microtubules guide the distribution of CSCs at the plasma membrane,[18,38] we also
examined how incipient hyperosmotic stress affects microtubules and their colocalization with
CSCs using the GFP-CESA3/mCherry-TUA5 double marker line.[38] In Hechtian structures where
the plasma membrane remained attached to the cell wall, mCherry-TUA5-labeled microtubules
appeared as strand-like structures, which were decorated by the immobilized population of
CSCs at the plasma membrane (Figure S3B).
234
To determine whether the plasma membrane-localized CSCs mediate wall-membrane
attachments, we treated YFP-LTI6b seedlings with the cellulose biosynthesis inhibitor isoxaben
(ISX), which has previously been shown to remove CSCs from the plasma membrane.[18] After
confirmation of this CSC clearance (Figure S3C and S3D), we induced plasmolysis with 0.5 M
sorbitol and quantified the degree of plasmolysis (Figure 4C). Compared to the mock condition,
ISX treatment resulted in a greater fraction of cells exhibiting increased plasmolysis (Figure 4D).
This trend is consistent with that observed in the cellulose-deficient mutants *cesa3^{je5}* (Figure 3A
and 3B) and *cesa6^{prc1-1}* (Figure 4E). Additionally, ISX caused changes in protoplast morphology
under 0.5 M sorbitol, specifically enhancing curvature at both ends (Figure 4C, Figure S3E). To
further assess the impact of ISX on wall-membrane attachments at the outer cell surface, we
induced plasmolysis using 1 M sorbitol. In mock-treated roots, wall-membrane attachment sites
were evident in regions where the plasma membrane had dissociated from the wall of the outer
cell surface. In contrast, in ISX-treated roots, protoplast detachment and shrinkage occurred
primarily at the cell-cell junctions (Figure 4F). These data confirm that wall-membrane
attachment sites contain CSCs, which are necessary for the physical stability of these
associations during hyperosmotic stress.
251
To further test this hypothesis, we analyzed another cell wall mutant, *shou4-3 4l-1*, which lacks
the negative regulators of CSC exocytosis, SHOU4 and SHOU4L, and was previously shown to
exhibit a higher CSC density at the plasma membrane relative to the wild type.[25] Root growth
reduction under hyperosmotic stress was significantly less severe in *shou4-3 4l-1* compared to
the wild type (Figure 4G). Moreover, the proportion of root cells in this mutant that exhibit no
measurable wall-membrane detachment was substantially higher than that in the wild type
(Figure 4H), resembling the phenotypes observed in the *rhm1-1* mutants (Figure 2D, Figure 3A
and 3B). The *cesa6^{prc1-1} shou4-1* double mutant,[25] however, displayed root growth phenotypes
similar to the *cesa6^{prc1-1}* single mutant (Figure S3F, Figure 2A), suggesting that the *cesa6^{prc1-1}*
mutation is epistatic to the *shou4-1* mutation. Similarly, the *cesa3^{je5}* mutation was epistatic to
rhm1-1 in root growth under both control and hyperosmotic stress conditions (Figure 4I, Figure
2A). As a complementary approach, we tested the effect of ISX on root growth in the *rhm1-1*

mutant or *shou4-3 4l-1* double mutant and found that this treatment similarly led to a more
severe degree of growth reduction than the non-ISX treated controls (Figure 4J). These results
indicate that CSCs act downstream of *RHM1* and *SHOU4/4L*.
267
268   Together, these data suggest that CSC density at the plasma membrane safeguards
269   hyperosmotic stress responses in roots by mediating wall-membrane attachments.
270
271   *REMs rapidly form nanodomains at wall-membrane attachment sites upon hyperosmotic stress*
272
CSCs are organized, by definition,[10] in nanodomains under both control and hyperosmotic
stress conditions (Figure 4A). To investigate how other plasma membrane nanodomains
respond to hyperosmotic stress, we focused on REMs, the best-characterized markers of
plasma membrane nanodomains[39] and known stabilizers of membrane topologies.[23] Using live-
cell laser scanning confocal microscopy in five-day-old seedling roots expressing YFP-REM1.2,
we observed a diffuse yet polarized localization, with stronger fluorescence signals detected at
the shootward regions of the plasma membrane (Figure 5A). Upon treatment with sorbitol for 5
minutes, YFP-REM1.2 coalesced into nanodomains that are approximately 0.4 µm in diameter,
and their density increased in a concentration-dependent manner with sorbitol (Figure 5A and
5B). This suggests that the density of YFP-REM1.2-positive nanodomains serves as a readout
for the severity of hyperosmotic stress. Similar nanodomain formation was observed in other
tissues, including cotyledons and hypocotyls, when treated with sorbitol (Figure S4A). Because
REMs are also components of plasmodesmata-localized nanodomains,[40] we crossed a
plasmodesmata marker, mCherry-PDCB1,[41] with YFP-REM1.2 to study their co-association. At
cell-cell junctions, a portion of YFP-REM1.2 nanodomains overlapped with mCherry-PDCB1
signal (Figure S4B), confirming their presence in plasmodesmata. However, YFP-REM1.2
nanodomains observed on the outer (periclinal) sides of cells were independent of
plasmodesmata (Figure S4B). Our study focused on this latter population of YFP-REM1.2
nanodomains. Time-lapse imaging revealed rapid formation of these immobile nanodomains,
likely instantaneously upon hyperosmotic stress (Figure S4C, Video S5). Notably, this process
was reversible: the nanodomains dispersed back into the plasma membrane when
hyperosmotic stress was alleviated (Video S6). Lastly, to evaluate the dependence of YFP-
REM1.2 nanodomain formation on sterols, a key membrane lipid species that facilitates REM
localization into highly ordered nanodomains[39], we treated YFP-REM1.2 seedlings with
fenpropimorph, a sterol biosynthesis inhibitor.[42] Unlike the mock-treated control, fenpropimorph-
treated seedlings did not exhibit YFP-REM1.2 nanodomains under hyperosmotic stress (Figure
S4D). These results suggest that the formation of YFP-REM1.2 nanodomains relies on the
sterol composition of the plasma membrane.
301
To test whether YFP-REM1.2 nanodomains are associated with wall-membrane attachment
sites, we performed live-cell imaging experiments using two approaches: staining five-day-old
YFP-REM1.2 seedlings with calcofluor white (Figure 5C) and generating a YFP-
REM1.2/mCherry-LTI6b double marker line (Figure 5D). Hyperosmotic stress-induced YFP-
REM1.2 nanodomains were attached to the wall (Figure 5C) and remained colocalized with
mCherry-LTI6b-labeled wall-membrane attachment sites at the outer surface of cells (Figure 5D

and 5E). To investigate the role of REMs in hyperosmotic stress responses, we analyzed the
rem1.2 1.3 double mutant[40,43] and the *rem1.2 1.3 1.4* triple mutant[43] by assessing root growth
and plasmolysis. *rem1.2 1.3* double mutants did not differ from the wild type (Figure S4E and
S4F), while *rem1.2 1.3 1.4* triple mutants displayed significantly reduced growth inhibition and
resistance to plasmolysis under stress conditions (Figure 5F-5H), phenotypes that were similar
to those of the *rhm1* and *shou4-3 4l-1* mutants (Figure 2D, Figure 3A and 3B, Figure 4G and
4H). Expression of YFP-REM1.2 under its native promoter in the *rem1.2 1.3 1.4* triple mutant
background restored the root growth phenotypes to wild-type levels (Figure 5F). Together, these
results suggest that the rapidly forming REM nanodomains function as negative regulators of
wall-membrane attachments.
318
319   *REMs antagonize CSCs to regulate hyperosmotic stress responses*
320
Given the observed formation of CSC nanodomains and REM nanodomains in root cells under
hyperosmotic stress, we next sought to investigate the relationship between these two
nanodomain populations. To visualize them, we crossed YFP-REM1.2 with tdTomato-CESA6
lines and imaged root cells using spinning disk confocal microscopy under control conditions or
in the presence of 1 M sorbitol. Under control conditions, tdTomato-CESA6-labeled CSC
particles were distributed across the plasma membrane, while YFP-REM1.2 signals were
predominantly localized to the shootward regions of the plasma membrane (Figure 6A). Upon
treatment with 1 M sorbitol, both markers formed distinct nanodomains (Figure 6A and 6B). To
quantify the spatial relationship of CSCs and REMs, we performed Manders' colocalization
analysis using the GFP-CESA3/tdTomato-CESA6 double marker line as a positive control
(Figure S5A). Under control conditions, YFP-REM1.2 signals showed minimal overlap with
tdTomato-CESA6-labeled CSC particles (Figure 6C). However, under hyperosmotic stress,
REM1.2 and CSC nanodomains exhibited partial colocalization, yet their Manders' overlap
coefficients remained significantly lower than those of the GFP-CESA3/tdTomato-CESA6
positive control (Figure 6C). These data suggest that CSCs and REMs function in spatially
distinct cellular domains under hyperosmotic stress, though partial colocalization suggests there
may be specific cellular contexts where their function may be coordinated.
338
To determine whether there is a functional relationship between the distribution of these two
nanodomains, we introduced the YFP-REM1.2 marker into the *cesa3^{je5}* and *rhm1-1* mutant
backgrounds and measured REM nanodomain density following 0.5 M sorbitol treatment. While
REM nanodomain density in the *cesa3^{je5}* mutant remained unchanged, it was significantly
reduced in the *rhm1-1* mutant compared to the wild type (Figure 6D and 6E), suggesting that
RHM1 promotes REM nanodomain formation. Additionally, we introduced the GFP-CESA3
marker into the *rem1.2 1.3 1.4* triple mutant and quantified CSC density at the plasma
membrane under control conditions. *rem1.2 1.3 1.4* triple mutants exhibited significantly higher
CSC density than wild type (Figure 6F and 6G). Given the importance of sterol biosynthesis for
REM1.2 nanodomain formation (Figure S4D), we tested whether it also influences CSC
patterning at the plasma membrane and affects plasmolysis. Compared to mock-treated roots,
treatment with fenpropimorph had no detectable effect on CSC density at the plasma membrane
or on protoplast behavior under 0.5 M sorbitol (Figure S5B-S5E). These data support the

hypothesis that CSC density at the plasma membrane is necessary for plasma membrane-cell
wall attachment, which plays a critical role in safeguarding root cells against the disruptive effect
of hyperosmotic stress. Furthermore, our data suggest that REMs inhibit the formation of these
attachments by limiting CSCs abundance at the plasma membrane, independently of REM
nanodomain formation.
357
To test this hypothesis, we analyzed the effect of ISX on hyperosmotic stress-induced growth
inhibition in *rem1.2 1.3 1.4* triple mutant roots. Under sorbitol treatment, mock-treated mutants
exhibited a 52% reduction in growth, whereas ISX treatment exacerbated the reduction to 83%
(Figure 6H), a level comparable to that observed in cellulose-deficient mutants (Figure 2A). We
also generated the *cesa3[ie5] rem1.2 1.3* triple mutant and found that this line phenocopied the
cesa3[ie5] single mutant under both control and hyperosmotic stress conditions (Figure 6I, Figure
2A). We further attempted to generate the *cesa3[ie5] rem1.2 1.3 1.4* quadruple mutant but were
unsuccessful because the *cesa3[ie5]* and *rem1.4* mutations are located on the same arm of
chromosome 5. Together, these results confirm that the reduced severity of hyperosmotic stress
responses in *rem1.2 1.3 1.4* triple mutants is a consequence of enhanced CSC density at the
plasma membrane, and further suggest an antagonistic relationship between REMs and CSCs.
369
370 *SHOU4/4L are components of REM nanodomains*
371
To uncover additional proteins present in REM nanodomains that could mediate the apparent
antagonism with CSCs, we employed proximity labeling and generated transgenic lines
expressing *TbID-YFP-REM1.2* driven by the *REM1.2* native promoter in the wild-type
background. After confirming that the TbID tag did not affect YFP-REM1.2 localization to the
plasma membrane or nanodomain formation (Figure S6A), we treated five-day-old seedlings
with 50 µM biotin for 15 minutes, in the presence or absence of 0.5 M sorbitol. In both
conditions, biotinylated proteins were detected in addition to those that are endogenously
biotinylated (Figure S6B and S6C). Overall, we identified 5,922 proteins, which were subjected
to two filtering steps (Figure 7A). To eliminate background proteins, we compared TbID-YFP-
REM1.2 samples with the wild-type control, identifying 30 and 111 proteins under the control
and sorbitol-treated conditions, respectively (Figure 7B and 7C). A subsequent comparison
between these conditions revealed 83 proteins enriched specifically under sorbitol treatment
(Figure 7D).
385
Our proteomic data revealed proteins known to interact with REM1.2, including REM1.2 itself,
REM1.3 and other REMs in the same family,[43] RPM1-INTERACTING PROTEIN4 (RIN4),[44] and
PLASMA-MEMBRANE ASSOCIATED CATION-BINDING PROTEIN1 (PCaP1)[45] (Figure 7B-
7D), confirming previous protein-protein interaction studies. Interestingly, some of these proteins
did not meet the statistical threshold across all three pairwise comparisons. For instance, RIN4
was enriched in TbID-YFP-REM1.2 samples compared to the wild type under both control and
sorbitol conditions, but not in the direct comparison between the two conditions (Figure 7B-7D).
Similarly, PCaP1 was enriched in TbID-YFP-REM1.2 samples compared to the wild type under
sorbitol conditions, but not under control conditions; however, it was enriched in the comparison
between the two conditions (Figure 7B-7D). These findings raise the possibility of false

negatives in the pairwise comparisons described above and led us to test whether these
proteins colocalize and interact with REM1.2 during hyperosmotic stress using an orthogonal
approach. To this end, we performed confocal microscopy and FRET-FLIM analyses by
transiently expressing YFP-REM1.2 in tobacco leaves, either alone or alongside an mRFP-
tagged protein of interest. mRFP-REM1.3 served as a positive control, while free mRFP was
used as a negative control. As anticipated, free mRFP localized to the cytosol did not colocalize
with or interact with YFP-REM1.2 under either control or sorbitol treatment conditions (Figure
7E-7H). mRFP-REM1.3 colocalized and interacted with YFP-REM1.2 at the plasma membrane
under control conditions and within nanodomains following sorbitol treatment (Figure 7E-7H).
Similarly, mRFP-RIN4 and PCaP1-mRFP also exhibited colocalization and interaction with YFP-
REM1.2 under both control and sorbitol conditions (Figure S7).
407
One group of proteins identified in the pairwise comparisons was the MECHANOSENSITIVE
CHANNEL OF SMALL CONDUCTANCE-LIKEs (MSLs), including MSL4, MSL6, and MSL9
(Figure 7B-7D). In tobacco infiltration assays, mRFP-MSL4 and mRFP-MSL6 colocalized with
YFP-REM1.2 at the plasma membrane under control conditions but not within nanodomains
following sorbitol treatment (Figure S8A and S8B). However, under prolonged sorbitol
treatment, colocalization was observed in the Hechtian reticulum (Figure S8A and S8B). In
contrast, the localization pattern of mRFP-MSL9 resembled that of the cortical endoplasmic
reticulum (ER) under both control and sorbitol conditions (Figure S8C), consistent with the close
proximity of the ER to the plasma membrane where YFP-REM1.2 is located. This observation
aligns with previous reports of MSL10, the closest homolog of MSL9 in the MSL family, being
localized at ER-plasma membrane contact sites.[46,47] To assess the functional role of MSLs in
Arabidopsis root growth responses to hyperosmotic stress, we compared the growth
phenotypes of the *msl4;5;6;9;10* quintuple mutant and the wild type. However, no significant
differences in growth were observed between the two genotypes under either control or sorbitol
treatment conditions (Figure S8D), suggesting that none of these MSLs are required for root
growth responses to hyperosmotic stress.
424
Another group of proteins identified in the pairwise comparisons was SHOU4/4L, known
inhibitors of CSC exocytosis.[25] Similar to the positive control mRFP-REM1.3, both SHOU4-
mRFP and SHOU4L-mRFP colocalized and interacted with YFP-REM1.2 at the plasma
membrane under control conditions and within nanodomains following sorbitol treatment (Figure
7E-7H, Figure S9). In Arabidopsis roots, the *shou4-3 4l-1* double mutants displayed reduced
root growth inhibition and increased resistance to plasmolysis than wild type under
hyperosmotic stress (Figure 4G and 4H). These phenotypes closely resembled those of the
rem1.2 1.3 1.4 triple mutants (Figure 5F-5H). Together, these findings suggest that SHOU4/4L
are components of the REM nanodomains that mediate the antagonism with CSC
nanodomains.
435
436   **Discussion**
437
In summary, we provided genetic, live-cell imaging, and proximity labeling-based proteomic
evidence that membrane nanodomains mediate the association and function of the cell wall-

plasma membrane interface. While these nanodomains are prominently observed at the outer
epidermal cell surface where they are in direct contact with the environment and function
independently of plasmodesmata, they are also likely present and functionally relevant at other
wall-membrane interfaces, such as in inner tissues where high-resolution live-cell imaging is
technically challenging. Maintaining CSC density at the plasma membrane is crucial for proper
cell wall-plasma membrane attachment and root responses to hyperosmotic stress. REMs
rapidly form nanodomains at the plasma membrane upon hyperosmotic stress and they harbor
many additional proteins in the vicinity, including the CSC exocytosis inhibitors SHOU4/4L.[25]
These findings reveal a molecular module in which CSCs are the core component that mediate
attachment between the cell wall and plasma membrane, acting downstream of and being
antagonized by REMs. Our findings provide new insights into the structural determinants of wall-
membrane attachment in plants and a physiological context in which their function supports the
resilience of plants to environmental stress.
453
Our survey of a panel of cell wall mutants in hyperosmotic stress-induced growth inhibition
revealed how different classes of cell wall polysaccharides are involved in this process. This
analysis led us to focus on two key proteins involved in cell wall biosynthesis: CSCs, and
RHM1. RHM1 encodes a UDP-L-rhamnose synthase, initially identified as a suppressor of
LEUCINE RICH REPEAT-EXTENSIN1 (LRX1), an extracellular protein that regulates cell wall
formation in root hairs.[48,49] RHM1 plays an important role in the morphogenesis of epidermal
cells, including root hairs,[48] cotyledon pavement cells,[50] and conical petal cells.[27,51] Although
rhamnose is required for both pectin biosynthesis and flavonol glycosylation, experimental
evidence suggests that rhamnose-containing RG-I directly contributes to epidermal
morphogenesis, whereas flavonol rhamnosylation has an indirect effect.[50,52] Our data reveal a
novel function of RHM1 in regulating root growth and responses to hyperosmotic stress (Figures
2 and 3). While we cannot definitively determine whether the observed *rhm1* mutant phenotypes
in our growth and plasmolysis assays result directly from RG-I rhamnose deficiency, we argue
that an indirect mechanism is more likely. This is supported by our findings that CSC machinery
components are enriched, and CSC density at the plasma membrane is increased, in the *rhm1-*
1 mutant (Figure 3). Additionally, the *cesa3* mutation is epistatic to *rhm1-1* (Figure 4). These
findings underscore the intricate interplay among cell wall polymers:[7] altering one wall
component can influence the synthesis, assembly, and/or organization of others.[53–55]
472
Environmental changes influence CSC dynamics and cellulose synthesis.[56] For example, salt
stress reduces cellulose production and leads to CSC depletion and reappearance at the
plasma membrane that coincides with cortical microtubule reorganization,[57] a process requiring
COMPANION OF CELLULOSE SYNTHASE proteins[58] and TETRATRICOPEPTIDE
THIOREDOXIN-LIKE (TTL) proteins.[59] Treating Arabidopsis seedlings with hyperosmotic stress
for several hours, whether mild (200 mM mannitol) or moderate (500 mM mannitol), triggers the
redistribution of CSCs from the plasma membrane into small CESA compartments (SmaCCs)[38]
or microtubule-associated cellulose synthase compartments (MASCs).[60] Prolonged severe
osmotic stress has also been shown to reduce cellulose production.[61] In alignment with these
previous observations, we demonstrate that acute, severe osmotic stress causes a subset of
CSCs to remain at the plasma membrane, decorating Hechtian structures at cell wall-plasma

membrane attachment sites (Figure 4, Video S4). These CSCs are immobile (Figure S3, Video
S3) and are likely inactive in cellulose synthesis, as CSC movement at the plasma membrane is
thought to be driven by cellulose polymerization.[18] Additionally, our study establishes a
functional link between CSC density at the plasma membrane and both root plasmolysis
response at the cellular level and growth response at the organ level. These findings provide
direct experimental evidence for the longstanding hypothesis that CSCs contribute to plasma
membrane anchoring to the cell wall, perhaps through the nascent cellulose microfibril that is
integrated into the cell wall.[19]
492
REMs, a plant-specific protein family,[62] are known to function in development,[40,43]
plasmodesmata aperture regulation,[40] membrane topology stabilization in root nodule
symbiosis,[23] and plant innate immunity.[63] Our work expands the understanding of REM function
in environmental stress responses by demonstrating that the sterol-dependent (Figure S4),
rapid formation of REM nanodomains at wall-membrane attachment sites (Figure 5, Video S5)
upon hyperosmotic stress serves as a hub for the aggregation of many other proteins (Figures
7, S7, S8, and S9). In addition, we show that REMs act upstream of CSCs, with SHOU4/4L as
their mediator, to regulate hyperosmotic stress responses in elongating roots (Figures 6 and 7).
The elevated CSC density observed in *rem1.2 1.3 1.4* triple mutants (Figure 6) and in *shou4-3
4l-1* double mutants under non-stress conditions,[25] the unchanged CSC density following
fenpropimorph treatment (Figure S5), along with the interaction between SHOU4/4L and
REM1.2 in the absence of stress (Figures 7 and S9), suggests that REM-CSC antagonism
occurs independently of REM nanodomain formation. This plasma membrane-localized
molecular module, comprising CSCs, SHOU4/4L, and REMs, functions both as a structural
anchor linking the cell wall to the plasma membrane and as a regulatory platform for
coordinating hyperosmotic stress responses.
509
510   **Limitations of the Study**
511
The ultrastructural and organizational details of how CSCs mediate the physical association
between the cell wall and plasma membrane remain to be elucidated. Although REM proteins
were initially characterized for their ability to bind both simple and complex galacturonides *in
vitro*,[24] it is still unclear whether such interactions occur *in vivo*. If they do, how REMs, peripheral
proteins anchored to the inner leaflet of the plasma membrane, engage with cell wall pectins
remains an open question. Our data revealed a correlation between plasmolysis upon
hyperosmotic shock and growth inhibition under hyperosmotic stress, but the relationship
between these two processes is not yet understood. In addition, proximity labeling identified
numerous proteins in the vicinity of REM1.2, yet how these interactions change in response to
hyperosmotic stress remains to be explored. These questions present exciting future directions
that we aim to pursue through interdisciplinary approaches to advance our understanding of
plant-environment interactions.
524
525   **Acknowledgements**
526

We thank past and present members of the Dinneny lab, Peter Dahlberg and his group at the
SLAC National Accelerator Laboratory, the Plant Biology community at Stanford University, and
Charles T. Anderson and his group at The Pennsylvania State University for insightful
discussions and critical feedback. We are especially grateful to Andrea Mair and Dominique
Bergmann for generously providing proximity labeling constructs and offering valuable advice on
proximity labeling experiments. We thank Renate Weizbauer, Andrey Malkovskiy, and the
Carnegie Advanced Imaging Facility for access to and assistance with the spinning disk
confocal microscope. We also thank Daniel Miranda and Olivia Creasey for their help with 3D
segmentation using AIVIA software. We gratefully acknowledge the following colleagues for
sharing genetic materials: Siamsa Doyle and Stéphanie Robert for *mur2-1*, *mur3-1,* and *mur4-1*
seeds; Olivier Hamant for *qua2-1* seeds; Henrik Scheller for *tbl10-1* and *gals1/2/3* seeds;
Christoph Ringli for *RHM1-GFP/rhm1-2* seeds; Adam Saffer and Vivian Irish for *rhm1-3* seeds;
Renee Weizbauer and David Ehrhardt for GFP-CESA3/mCherry-TUA5 seeds; Yvon Jaillais for
mCherry-LTI6b seeds; Yoselin Benitez-Alfonso for mCherry-PDCB1 seeds; Xu Chen for *rem1.2*
1.3c seeds; and Charles T. Anderson for GFP-CESA3/tdTomato-CESA6 seeds.
542
Y.R. is a 2020 Simons fellow of the Life Sciences Research Foundation. This project was
supported by grants awarded to J.R.D. from the Biological and Environmental Research (BER)
Program, US Department of Energy (DE-AC02-76SF00515), National Institutes of Health,
National Institute of General Medicine Sciences (R01 GM123259-01), the Stanford Precourt
Institute, and a Faculty Scholars grant from the Simons Foundation and Howard Hughes
Medical Institute (55108515), and a grant awarded to J.J.K. from the National Science
Foundation (IOS-1856431). J.R.D. is an investigator of the Howard Hughes Medical Institute.
N.B.A and T.O. were supported by the German Research Foundation (Deutsche
Forschungsgemeinschaft; DFG) in frame of the Collaborative Research Center 924 (project ID
170483403). S.-L.X. was supported by the National Institutes of Health (S10OD030441) and the
Carnegie endowment fund to the Carnegie mass spectrometry facility.

554
555 **Author contributions**
556
557 Conceptualization, Y.R. and J.R.D.; data acquisition and analysis, Y.R., A.V.R., T.S.G., N.B.A,
558 W.P.D., and S.-L.X.; data interpretation, all authors; funding, Y.R., T.O., J.J.K., S.-L.X., and
559 J.R.D; supervision, Y.R. and J.R.D; writing of the original draft, Y.R. and J.R.D; manuscript
560 editing, Y.R., N.B.A, W.P.D., T.O., J.J.K., S.-L.X., and J.R.D..
561
562 **Declaration of interests**
563
564 The authors declare no competing interests.
565
566 **Methods**
567
568 Plant Materials and Growth
569

Arabidopsis thaliana seeds were surface-sterilized using a solution of 30% bleach and 0.1%
SDS for 20 minutes, followed by four rinses with sterile water. The sterilized seeds were
stratified in the dark at 4°C for 2-3 days before being germinated on 1X Murashige and Skoog
(MS) plates supplemented with 1% sucrose and solidified with 0.7% gelzan (pH 5.7). Seedlings
were grown vertically in a growth chamber maintained at 22°C under a 16-hour light/8-hour dark
photoperiod.
576
577 Molecular Cloning and Plant Transformation
578
The *CESA3* coding sequence with attB sites was amplified from Arabidopsis Col-0 cDNA and
introduced into the pDONR221 vector (Thermo Fisher, catalog #12536017) using a Gateway BP
reaction. A 1831-bp promoter fragment upstream of the *CESA3* start codon was amplified from
Col-0 genomic DNA with SbfI and XbaI restriction sites. The destination vector pGWB506
(Addgene, catalog #74848) was digested with SbfI and XbaI to replace the 35S promoter with
the *CESA3* promoter. The *CESA3* coding sequence was subsequently inserted into the
modified pGWB506 vector via a Gateway LR reaction to generate the *ProCESA3:GFP-CESA3*
construct. This construct was transformed into Arabidopsis Col-0 plants, and transgenic lines
were selected on medium containing 25 µg/mL Hygromycin B.
588
TurboID and YFP were amplified from the plasmid pENTR_L1-YFP-Turbo-L2 (Addgene, catalog
#127350). *LTI6b* was amplified from Col-0 genomic DNA, including its exons, introns, and
3'UTR. *REM1.2* was amplified from Col-0 cDNA. To generate TbID-YFP-LTI6b and TbID-YFP-
REM1.2, overlapping PCR was performed using these amplified fragments, introducing AgeI
and EcoRI restriction sites at the N- and C-termini, respectively. A 1568-bp promoter fragment
upstream of the *REM1.2* start codon was amplified from Col-0 genomic DNA, incorporating
PmeI and AgeI sites. The pEGAD empty vector (ABRC stock #CD3-389) was digested with
PmeI and AgeI to replace the *35S* promoter with the *REM1.2* promoter. This modified pEGAD
vector was subsequently digested with AgeI and EcoRI to replace the GFP sequence with TbID-
YFP-LTI6b or TbID-YFP-REM1.2. Constructs were then transformed into Arabidopsis Col-0 or
mutant backgrounds to generate transgenic lines, which were then selected using 50 µM Basta.
600
The *35S:YFP-REM1.2* construct used in tobacco infiltration assays was generated following the
same restriction digestion strategy described above. For *35S* promoter-driven mRFP-tagged
protein constructs, the coding sequences of *REM1.3*, *RIN4*, *PCaP1*, *SHOU4*, and *SHOU4L*
were amplified from Col-0 cDNA with attB sites. The genomic sequences of *MSL4*, *MSL6*, and
MSL9, including exons and introns, were amplified from Col-0 genomic DNA with attB sites. All
amplified sequences were first cloned into the pDONR221 vector and subsequently introduced
into pGWB554 (Addgene, catalog #74884) or pGWB555 (Addgene, catalog #74885) to create
mRFP fusion constructs with C- or N-terminal tags, respectively.
609
610 Root Growth Assay
611
612 Four-day-old seedlings grown on standard 1X MS media were transferred to either fresh
613 standard media or media supplemented with 0.28 M sorbitol. At the time of transfer, root tips

614 were marked to track growth. Seedlings were then grown for an additional day, after which the
615 plates were scanned using a dual-light flatbed scanner (Epson V800). Root growth was
616 quantified in ImageJ by measuring the root length extending beyond the marked position.
617
618 Pharmacological Treatments
619
To inhibit cellulose biosynthesis, we used isoxaben (ISX) (Sigma, catalog #36138-100MG) in
two different experimental setups. For acute clearance of CSCs from the plasma membrane,
five-day-old seedlings expressing GFP-LTI6b and tdTomato-CESA6 were incubated in MS
liquid containing 100 nM ISX for 20 min prior to imaging for plasmolysis. MS liquid with 0.1%
DMSO served as the mock treatment. To assess the effect of ISX on root growth, four-day-old
seedlings were transferred from standard 1X MS plates to one of four treatment conditions:
standard plates with 0.1% DMSO, standard plates with 1 nM ISX, 0.28 M sorbitol plates with
0.1% DMSO, or 0.28 M sorbitol plates with 1 nM ISX. Root growth was subsequently measured
as described above.
629
630 To inhibit sterol biosynthesis, 100 µg/mL fenpropimorph (Sigma, catalog #40713-100MG) was
631 added to standard 1X MS media. 1X MS media containing 0.1% DMSO was used as the mock.
632 Four-day-old seedlings expressing YFP-REM1.2 were transferred from standard plate to either
633 the mock or fenpropimorph plate and let grow for an additional day.
634
635 Confocal Microscopy
636
To visualize plasmolysis in transgenic lines expressing YFP- or GFP-LTI6b, five-day-old
seedlings were treated with 0.5 M sorbitol for 5 min on a glass slide and imaged on a Leica TCS
SP8 microscope with an HC PL APO 93x/1.3 NA objective. YFP was excited with a white light
laser at 514 nm, with emission detected between 525-570 nm. GFP was excited at 488 nm, with
emission detected between 500-560 nm. Bright-field and fluorescence images were acquired
simultaneously. The plasmolysis ratio was quantified in ImageJ by dividing the protoplast area
(measured from the fluorescence image) by the total cell area (measured from the bright-field
image). Protoplast curvature at cell ends was measured following previously established
methods.[64]
646
For calcofluor white staining, five-day-old seedlings were stained with 0.1 µM (10^{-5}%, w/v)
calcofluor white (Santa Cruz Biotechnology, catalog #sc-218504) diluted in MS liquid in the dark
for 30 min, washed with MS liquid, treated with fresh MS liquid or 1 M sorbitol for 5 min on a
glass slide, and imaged on a Leica TCS SP8 microscope with an HC PL APO 93x/1.3 NA
objective. Calcofluor was excited by UV at 405 nm, with emission detected between 420-470
nm.
653
CESA imaging was done on a Leica DMI6000 inverted microscope attached to a Yokogawa
spinning disk confocal head with a 100x/1.4 NA objective. GFP was excited at 488 nm with a
525/50 nm emission filter. tdTomato was excited at 561 nm with a 593/40 nm emission filter.
Excitation energy was adjusted to 4 mW power and exposure time was 200 ms. Z-stacks were

obtained with a step size of 0.27 μm. Time-lapse images were recorded at a single focal plane
at the plasma membrane with an interval of 10 s and a duration of 5 min.
660
REM imaging was performed using a Leica TCS SP8 microscope with an HC PL APO 93x/1.3
NA objective. YFP excitation and emission parameters were consistent with those described
above. For time-lapse imaging of REM nanodomain formation, a five-day-old YFP-REM1.2
seedling was placed under a 1X MS + 1% sucrose, 0.7% gelzan pad in a Nunc Lab-Tek
chambered coverglass (Thermo Fisher, catalog #155360). A 0.5-cm diameter hole was created
adjacent to the root tip. Time-lapse z-stack images were taken at a step-size of 0.3 μm,
beginning at 0 min (pre-treatment) and then every minute for 10 min following the addition of
200 μL of 0.5 M sorbitol into the hole. To image REM nanodomain disappearance upon relief
from hyperosmotic stress, a five-day-old YFP-REM1.2 seedling was pre-treated with 0.5 M
sorbitol for 5 min and then placed under a 1X MS + 1% sucrose + 0.5 M sorbitol, 0.7% gelzan
pad in a Nunc Lab-Tek chambered coverglass. Time-lapse z-stack imaging was performed
under the same parameters as described above, except that 200 μL of 1X MS + 1% sucrose
liquid solution was added to the hole instead of 0.5 M sorbitol.
674
675   Image Analysis
676
677   3D renderings of confocal images were generated using Leica LAS X or AIVIA 15.0.0 software
678   (Leica Microsystems). Hechtian structures were segmented in AIVIA. Quantitative image
679   analyses were performed with ImageJ and Imaris 10.1.1 (Oxford Instruments).
680
CESA density was quantified in ImageJ. Z-stack images were first background-subtracted using
the sliding paraboloid algorithm with a rolling ball radius of 30 pixels, and contrast was
enhanced by setting saturated pixels to 0.35%. For each root cell, a substack containing CESA
particles at the plasma membrane was generated and a maximum intensity projection was
created. A region of interest (ROI) was defined using the polygon tool to exclude the Golgi, and
its area was measured. Within this ROI, CESA particles were identified using the Find Maxima
tool with a noise tolerance of 500.
688
689   Time-lapse images of REM nanodomain formation and disappearance were processed in
690   ImageJ by generating maximum projections of z-stacks at each time point using ImageJ. The
691   resulting projections were concatenated and aligned using the StackReg plugin to correct for
692   frame shifts.
693
To measure REM nanodomain density, z-stack images were similarly contrast-enhanced in
ImageJ with saturated pixels set at 0.35%. A substack containing REM nanodomains at the
plasma membrane was generated for each root cell, followed by maximum intensity projection
and ROI selection using the polygon tool. The projection was then cropped and imported into
Imaris, where REM nanodomains were detected using a 0.4-μm diameter threshold.

Colocalization analysis for dual marker lines was performed in ImageJ using the Coloc 2 plugin.
For each root cell, an ROI was added to the ROI Manager and selected. Manders' overlap

coefficient was calculated using the Costes threshold regression method with 100 Costes
randomizations.

Biotin Treatment
704
One gram of five-day-old whole seedlings grown on MS plates was harvested into a nylon
biopsy bag (Fisher Scientific, catalog #50-303-41). The bag was submerged in either 50 μM
biotin in MS liquid medium (control) or 50 μM biotin in MS liquid supplemented with 0.5 M
sorbitol for 15 minutes. After treatment, samples were washed with the corresponding solution
(MS or MS + 0.5 M sorbitol) and gently blotted on a paper towel to remove excess liquid.
Samples were then flash-frozen in liquid nitrogen, ground into fine powder using a mortar and a
pestle, and stored at -80°C for further use.
712
713   Immunoblotting
714
To confirm biotinylation, 100 μl of finely ground frozen tissue powder from the biotin treatment
was resuspended in 100 μl of 2× Laemmli buffer containing 120 mM Tris (pH 6.8), 4% SDS,
20% glycerol, 5% β-mercaptoethanol, and 0.05% bromophenol blue. Samples were boiled at
95°C for 5 minutes, and 10 μl of the supernatant was loaded onto a 10% SDS-PAGE gel.
Proteins were transferred to a 0.2 μm PVDF membrane (Bio-Rad, catalog #1704156) using the
Trans-Blot Turbo Transfer System (Bio-Rad). Membranes were either stained with Coomassie
Brilliant Blue R-250 (Bio-Rad, catalog #1610400) for 30 seconds and destained with a solution
containing 40% methanol and 10% acetic acid, or blocked in 5% BSA in TBS and probed with
HRP-conjugated streptavidin (Thermo Fisher, catalog #S911) diluted 1:5,000 in 5% BSA in
TBST for 1 hour at room temperature. Chemiluminescence was detected using the Clarity ECL
Western Substrate (Bio-Rad, catalog #1705061) and visualized with the ChemiDoc MP Imaging
System (Bio-Rad).
727
728   Protein Affinity Purification
729
One gram of the finely ground frozen tissue powder from the biotin treatment was resuspended
in 2 mL of protein extraction buffer containing 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.1%
SDS, 1% Triton X-100, 0.5% sodium deoxycholate, 1 mM EGTA, 1 mM DTT, 1× cOmplete
protease inhibitor cocktail (Roche, catalog #COUEDTAF-RO), and 1 mM PMSF. The
suspension was vortexed and incubated on a rotor wheel at 4°C for 10 min. Samples were then
sonicated in an ice bath using a Branson 450 sonifier at 20% amplitude with 10 s on/10 s off
pulses for a total of 2 min. Following sonication, samples were centrifuged at 12,000 g for 10
min at 4°C. A total of 2.5 mL of the resulting supernatant was loaded onto PD-10 desalting
columns (Cytiva, catalog #17-0851-01) pre-equilibrated with equilibration buffer composed of 50
mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.1% SDS, 1% Triton X-100, 0.5% sodium deoxycholate,
1 mM EGTA, and 1 mM DTT. Proteins were eluted with 3.5 mL of equilibration buffer into 5 mL
LoBind tubes (Eppendorf, catalog #0030108302). For streptavidin pulldown, 200 μL of
Dynabeads MyOne Streptavidin T1 beads (Thermo Fisher, catalog #65602) were washed with
extraction buffer and then incubated with the desalted protein extract supplemented with 1×

cOmplete protease inhibitor cocktail and 1 mM PMSF. The mixture was incubated overnight at
4°C on a rotor wheel. The following day, beads were separated using a magnetic rack,
transferred to 1.5 mL LoBind tubes (Eppendorf, catalog #022431081), and subjected to a series
of washes: twice with cold extraction buffer, twice with cold equilibration buffer, once with cold 1
M KCl, transferred to a fresh 1.5 mL LoBind tube, and washed twice more with cold extraction
buffer. After the final wash, beads were separated on a magnetic rack to remove residual buffer
and stored at -80°C for further use.
751
752 Mass Spectrometry Sample Preparation
753
Frozen streptavidin beads were thawed and washed once with PBS prepared in HPLC-grade
water (Thermo Fisher, catalog #022934-k7). The beads were transferred to a new 1.5 mL
LoBind tube and washed five additional times with PBS in HPLC-grade water. After the final
wash, beads were separated on a magnetic rack, and the buffer was carefully removed. Beads
were then incubated with 80 μL trypsin digestion buffer containing 50 mM Tris (pH 7.5), 1 M
Urea, 4 mM DTT, and 0.4 μg Trypsin at 25°C for 3 h with shaking at 1,200 rpm. Following
incubation, beads were magnetically separated and washed twice with 60 μL of 1 M urea in 50
mM Tris (pH 7.5). The resulting supernatants were pooled and treated with 10 mM
iodoacetamide at 25°C for 45 min in the dark with shaking at 1,200 rpm to alkylate cysteines.
Digestion was continued by adding 0.5 μg of additional trypsin, followed by overnight incubation
at 25°C with shaking at 1,200 rpm. The next day, a final 0.5 μg of trypsin was added, and
samples were incubated for an additional 4 h at 25°C with shaking. Following digestion, 1%
formic acid was added to acidify the samples, which were then vortexed and desalted using
OMIX C18 pipette tips (10-100 μL, Agilent, catalog #A57003100) according to the
manufacturer's instructions. Peptides were dried using a speed vacuum centrifuge and
resuspended in 0.1% formic acid for LC-MS/MS analysis. Samples were analyzed using a Q-
Exactive HF hybrid quadrupole-Orbitrap mass spectrometer (Thermo Fisher), coupled to an
Easy-nLC 1200 ultra-performance liquid chromatography system (Thermo Fisher).[29]
772
773 Mass Spectrometry Data Analysis
774
Mass spectrometry data were analyzed using MaxQuant (version 2.4.2.0)[65] and Perseus
(version 2.0.10.0)[66]. For plasma membrane proteome comparisons between wild type and
cesa3^{je5}, raw data from TbID-YFP-LTI6b/Col-0, Col-0, TbID-YFP-LTI6b/*cesa3^{je5}*, and *cesa3^{je5}*
were searched together in MaxQuant using default parameters with the following modifications.
The LFQ minimum ratio count was set to 2. Peptides were searched against the TAIR10 protein
database ("TAIR10_pep_20101214";
https://www.arabidopsis.org/download/list?dir=Proteins%2FTAIR10_protein_lists),
supplemented with a custom list of potential contaminants, such as trypsin, human keratin,
streptavidin, BSA, YFP, TbID-YFP, along with the standard MaxQuant contaminant list. "Match
between runs" was enabled with a match time window of 0.4 min and an alignment time window
of 20 min. The resulting "proteinGroups.txt" file was imported into Perseus with LFQ intensities
as the Main category. Proteins marked with reverse, contaminant, and only identified by site
identifications were removed. Data were log2-transformed, grouped, and filtered to retain

proteins with at least two valid values in at least one group. Data were imputed by replacing
missing LFQ values from normal distribution using the total matrix mode with a width of 0.3 and
down shift of 1.8. To remove all unspecific background labeling, each TbID-YFP-LTI6b-tagged
group was compared to its respective genetic background by a one-sided Welch's t-test with S0
of 0.5, FDR of 0.05, and 250 randomizations. The resulting filtered matrix was then used to
compare TbID-YFP-LTI6b/*cesa3^{je5}* and TbID-YFP-LTI6b/Col-0 using a two-sided Student's t-test
with S0 of 0.5, FDR of 0.05, and 250 randomizations.
795
796   The same workflow was applied to the plasma membrane proteome comparison between the
797   wild type and *rhm1-1*, using raw data of TbID-YFP-LTI6b/Col-0, Col-0, TbID-YFP-LTI6b/*rhm1-1*,
798   and *rhm1-1* samples.
799
For REM1.2 proximity labeling experiments, raw data from TbID-YFP-REM1.2/Col-0 (control
condition), TbID-YFP-LTI6b/Col-0 (control condition), Col-0 (control condition), TbID-YFP-
REM1.2/Col-0 (0.5 M sorbitol), TbID-YFP-LTI6b/Col-0 (0.5 M sorbitol), and Col-0 (0.5 M
sorbitol) were analyzed together using the same MaxQuant and Perseus workflow described
above. To remove background labeling, TbID-YFP-REM1.2/Col-0 was compared to Col-0 under
each condition by a one-sided Welch's t-test with S0 of 0.5, FDR of 0.05, and 250
randomizations. The resulting filtered matrix was then used to compare the 0.5 M sorbitol
treatment and the control condition in TbID-YFP-REM1.2/Col-0 using a two-sided Student's t-
test with S0 of 0.5, FDR of 0.05, and 250 randomizations.
809
810   The mass spectrometry proteomics data have been deposited to the ProteomeXchange
811   Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE[67] partner repository.
812
813   Transient Expression in Tobacco
814
Nicotiana benthamiana seeds were sown directly in PRO-MIX HP soil and grown in a growth
chamber at 25°C under a 12-hour light/12-hour dark photoperiod. DNA constructs were
transformed into *Agrobacterium tumefaciens* GV3101 (pMP90 and pSoup) strain using
electroporation. Transformed Agrobacterium were cultured in 5 mL LB medium supplemented
with antibiotics for two days, then pelleted by centrifugation at 3,000 g for 10 minutes. Pellets
were resuspended and diluted in an infiltration medium containing 10 mM MgCl$_2$, 10 mM MES to
an OD600 of 0.5. Cultures were incubated at room temperature for 3 h with 150 µM
acetosyringone. For infiltration, cultures were mixed with the Agrobacterium p19 strain at an
OD600 of 0.2 and infiltrated to the abaxial side of leaves from 5- to 6-week-old Nicotiana plants.
Infiltrated plants were kept in the growth chamber for an additional 3-5 days before imaging.
825
826   FRET-FLIM
827
FRET-FLIM experiments in Nicotiana leaves were performed using a Leica TCS SP8 confocal
microscope equipped with an HC PL APO 93x/1.3 NA objective. Leaves infiltrated with YFP-
REM1.2 alone served as donor-only controls, while those co-infiltrated with YFP-REM1.2 and
an mRFP-tagged construct served as donor + acceptor samples. Samples were incubated in

either MS liquid medium or 1 M sorbitol to induce hyperosmotic stress. The donor (YFP) was
excited at 514 nm using a pulsed white light laser operating at 80 MHz. Emission was collected
between 525–570 nm using a HyD SMD hybrid detector. Laser power was adjusted to ensure a
maximum of approximately one photon per laser pulse, and frame repetition was set to 20.
836
837 YFP fluorescence lifetimes were measured using the LAS X software. Under control conditions,
838 ROIs were defined as narrow rectangular areas along the plasma membrane at cell borders.
839 Under hyperosmotic stress, ROIs were selected based on individual YFP-REM1.2
840 nanodomains. Approximately 3-5 ROIs were analyzed per image. YFP lifetimes were calculated
841 using an n-Exponential Reconvolution model with one component.
842
843 Quantification and Statistical Analysis
844
845 Grouped and scattered column graphs displaying mean ± SD were generated using Prism 10
846 (GraphPad). Histograms were created in Microsoft Excel, and volcano plots were generated
847 using R. Statistical analyses were performed using two-way ANOVA in Prism 10. Details of
848 statistical tests, including sample sizes and significance levels, are provided in the figures and
849 corresponding legends.
850
References

1. Theocharis, A.D., Skandalis, S.S., Gialeli, C., and Karamanos, N.K. (2016). Extracellular
matrix structure. Adv. Drug Deliv. Rev. *97*, 4–27.

2. Humphries, J.D., Byron, A., and Humphries, M.J. (2006). Integrin ligands at a glance. J.
Cell Sci. *119*, 3901–3903.

3. Kohorn, B.D. (2000). Plasma membrane-cell wall contacts. Plant Physiol. *124*, 31–38.

4. Feng, W., Lindner, H., Robbins, N.E., 2nd, and Dinneny, J.R. (2016). Growing Out of
Stress: The Role of Cell- and Organ-Scale Growth Control in Plant Water-Stress
Responses. Plant Cell *28*, 1769–1782.

5. Lampugnani, E.R., Khan, G.A., Somssich, M., and Persson, S. (2018). Building a plant cell
wall at a glance. J. Cell Sci. *131*. https://doi.org/10.1242/jcs.207373.

6. Delmer, D., Dixon, R.A., Keegstra, K., and Mohnen, D. (2024). The plant cell wall-dynamic,
strong, and adaptable-is a natural shapeshifter. Plant Cell *36*, 1257–1311.

7. Rui, Y., and Dinneny, J.R. (2020). A wall with integrity: surveillance and maintenance of the
plant cell wall under stress. New Phytol. *225*, 1428–1439.

8. Martinière, A., Lavagi, I., Nageswaran, G., Rolfe, D.J., Maneta-Peyret, L., Luu, D.-T.,
Botchway, S.W., Webb, S.E.D., Mongrand, S., Maurel, C., et al. (2012). Cell wall constrains
lateral diffusion of plant plasma-membrane proteins. Proc. Natl. Acad. Sci. U. S. A. *109*,
12805–12810.

9. McKenna, J.F., Rolfe, D.J., Webb, S.E.D., Tolmie, A.F., Botchway, S.W., Martin-
Fernandez, M.L., Hawes, C., and Runions, J. (2019). The cell wall regulates dynamics and

size of plasma-membrane nanodomains in Arabidopsis. Proc. Natl. Acad. Sci. U. S. A. *116*,
12857–12862.

10. Jaillais, Y., Bayer, E., Bergmann, D.C., Botella, M.A., Boutté, Y., Bozkurt, T.O., Caillaud,
M.-C., Germain, V., Grossmann, G., Heilmann, I., et al. (2024). Guidelines for naming and
studying plasma membrane domains in plants. Nat. Plants *10*, 1172–1183.

11. Hdedeh, O., Mercier, C., Poitout, A., Martinière, A., and Zelazny, E. (2025). Membrane
nanodomains to shape plant cellular functions and signaling. New Phytol. *245*, 1369–1385.

12. Feng, W., Kita, D., Peaucelle, A., Cartwright, H.N., Doan, V., Duan, Q., Liu, M.-C., Maman,
J., Steinhorst, L., Schmitz-Thom, I., et al. (2018). The FERONIA Receptor Kinase Maintains
Cell-Wall Integrity during Salt Stress through Ca2+ Signaling. Curr. Biol. *28*, 666–675.e5.

13. Hecht., V.K. (1912). Studien über den Vorgang der Plasmolyse (Beitr. Biol. Pflanz.).

14. Arico, D.S., Dickmann, J.E.M., Hamant, O., and Canut, H. (2023). The plasma membrane -
cell wall nexus in plant cells: focus on the Hechtian structure. Cell Surf. *10*, 100115.

15. Haswell, E.S., and Verslues, P.E. (2015). The ongoing search for the molecular basis of
plant osmosensing. J. Gen. Physiol. *145*, 389–394.

16. Yu, Q., Ren, J.-J., Kong, L.-J., and Wang, X.-L. (2018). Actin filaments regulate the
adhesion between the plasma membrane and the cell wall of tobacco guard cells.
Protoplasma *255*, 235–245.

17. Gronnier, J., Gerbeau-Pissot, P., Germain, V., Mongrand, S., and Simon-Plas, F. (2018).
Divide and Rule: Plant Plasma Membrane Organization. Trends Plant Sci. *23*, 899–917.

18. Paredez, A.R., Somerville, C.R., and Ehrhardt, D.W. (2006). Visualization of cellulose
synthase demonstrates functional association with microtubules. Science *312*, 1491–1495.

19. Lang, I., Barton, D.A., and Overall, R.L. (2004). Membrane-wall attachments in
plasmolysed plant cells. Protoplasma *224*, 231–243.

20. Smokvarska, M., Francis, C., Platre, M.P., Fiche, J.-B., Alcon, C., Dumont, X., Nacry, P.,
Bayle, V., Nollmann, M., Maurel, C., et al. (2020). A plasma membrane nanodomain
ensures signal specificity during osmotic signaling in plants. Curr. Biol. *30*, 4654–4664.e4.

21. Gouguet, P., Gronnier, J., Legrand, A., Perraki, A., Jolivet, M.-D., Deroubaix, A.-F.,
German-Retana, S., Boudsocq, M., Habenstein, B., Mongrand, S., et al. (2021). Connecting
the dots: from nanodomains to physiological functions of REMORINs. Plant Physiol. *185*,
632–649.

22. Liang, P., Stratil, T.F., Popp, C., Marín, M., Folgmann, J., Mysore, K.S., Wen, J., and Ott, T.
(2018). Symbiotic root infections in Medicago truncatula require remorin-mediated receptor
stabilization in membrane nanodomains. Proc. Natl. Acad. Sci. U. S. A. *115*, 5289–5294.

23. Su, C., Rodriguez-Franco, M., Lace, B., Nebel, N., Hernandez-Reyes, C., Liang, P.,
Schulze, E., Mymrikov, E.V., Gross, N.M., Knerr, J., et al. (2023). Stabilization of
membrane topologies by proteinaceous remorin scaffolds. Nat. Commun. *14*, 323.

24. Reymond, P., Kunz, B., Paul-Pletzer, K., Grimm, R., Eckerskorn, C., and Farmer, E.E.

(1996). Cloning of a cDNA encoding a plasma membrane-associated, uronide binding
phosphoprotein with physical properties similar to viral movement proteins. Plant Cell *8*,
2265–2276.

25. Polko, J.K., Barnes, W.J., Voiniciuc, C., Doctor, S., Steinwand, B., Hill, J.L., Jr, Tien, M.,
Pauly, M., Anderson, C.T., and Kieber, J.J. (2018). SHOU4 proteins regulate trafficking of
cellulose synthase complexes to the plasma membrane. Curr. Biol. *28*, 3174–3182.e6.

26. Mouille, G., Ralet, M.-C., Cavelier, C., Eland, C., Effroy, D., Hématy, K., McCartney, L.,
Truong, H.N., Gaudon, V., Thibault, J.-F., et al. (2007). Homogalacturonan synthesis in
Arabidopsis thaliana requires a Golgi-localized protein with a putative methyltransferase
domain: Homogalacturonan synthesis in Arabidopsis: Plant J. *50*, 605–614.

27. Saffer, A.M., Carpita, N.C., and Irish, V.F. (2017). Rhamnose-Containing Cell Wall
Polymers Suppress Helical Plant Growth Independently of Microtubule Orientation. Curr.
Biol. *27*, 2248–2259.e4.

28. Ebert, B., Birdseye, D., Liwanag, A.J.M., Laursen, T., Rennie, E.A., Guo, X., Catena, M.,
Rautengarten, C., Stonebloom, S.H., Gluza, P., et al. (2018). The three members of the
Arabidopsis Glycosyltransferase Family 92 are functional β-1,4-galactan synthases. Plant
Cell Physiol. *59*, 2624–2636.

29. Mair, A., Xu, S.-L., Branon, T.C., Ting, A.Y., and Bergmann, D.C. (2019). Proximity labeling
of protein complexes and cell-type-specific organellar proteomes in Arabidopsis enabled by
TurboID. Elife *8*, e47864.

30. Xu, S.-L., Shrestha, R., Karunadasa, S.S., and Xie, P.-Q. (2023). Proximity labeling in
plants. Annu. Rev. Plant Biol. *74*, 285–312.

31. Moulinier-Anzola, J., De-Araujo, L., and Korbei, B. (2014). Expression of Arabidopsis TOL
genes. Plant Signal. Behav. *9*, e28667.

32. Korbei, B., Moulinier-Anzola, J., De-Araujo, L., Lucyshyn, D., Retzer, K., Khan, M.A., and
Luschnig, C. (2013). Arabidopsis TOL proteins act as gatekeepers for vacuolar sorting of
PIN2 plasma membrane protein. Curr. Biol. *23*, 2500–2505.

33. Li, S., Lei, L., Somerville, C.R., and Gu, Y. (2012). Cellulose synthase interactive protein 1
(CSI1) links microtubules and cellulose synthase complexes. Proc. Natl. Acad. Sci. U. S. A.
109, 185–190.

34. Bringmann, M., Li, E., Sampathkumar, A., Kocabek, T., Hauser, M.-T., and Persson, S.
(2012). POM-POM2/cellulose synthase interacting1 is essential for the functional
association of cellulose synthase and microtubules in Arabidopsis. Plant Cell *24*, 163–177.

35. Lei, L., Li, S., Du, J., Bashline, L., and Gu, Y. (2013). Cellulose synthase INTERACTIVE3
regulates cellulose biosynthesis in both a microtubule-dependent and microtubule-
independent manner in Arabidopsis. Plant Cell *25*, 4912–4923.

36. Zhu, X., Li, S., Pan, S., Xin, X., and Gu, Y. (2018). CSI1, PATROL1, and exocyst complex
cooperate in delivery of cellulose synthase complexes to the plasma membrane. Proc. Natl.
Acad. Sci. U. S. A. *115*, E3578–E3587.

37. Liu, Z., Schneider, R., Kesten, C., Zhang, Y., Somssich, M., Zhang, Y., Fernie, A.R., and
950   Persson, S. (2016). Cellulose-microtubule uncoupling proteins prevent lateral displacement
951   of microtubules during cellulose synthesis in Arabidopsis. Dev. Cell *38*, 305–315.

952 38. Gutierrez, R., Lindeboom, J.J., Paredez, A.R., Emons, A.M.C., and Ehrhardt, D.W. (2009).
953   Arabidopsis cortical microtubules position cellulose synthase delivery to the plasma
954   membrane and interact with cellulose synthase trafficking compartments. Nat. Cell Biol. *11*,
955   797–806.

956 39. Gronnier, J., Crowet, J.-M., Habenstein, B., Nasir, M.N., Bayle, V., Hosy, E., Platre, M.P.,
957   Gouguet, P., Raffaele, S., Martinez, D., et al. (2017). Structural basis for plant plasma
958   membrane protein dynamics and organization into functional nanodomains. Elife *6*, e26404.

959 40. Huang, D., Sun, Y., Ma, Z., Ke, M., Cui, Y., Chen, Z., Chen, C., Ji, C., Tran, T.M., Yang, L.,
960   et al. (2019). Salicylic acid-mediated plasmodesmal closure via Remorin-dependent lipid
961   organization. Preprint, https://doi.org/10.1073/pnas.1911892116
962   https://doi.org/10.1073/pnas.1911892116.

963 41. Benitez-Alfonso, Y., Faulkner, C., Pendle, A., Miyashima, S., Helariutta, Y., and Maule, A.
964   (2013). Symplastic intercellular connectivity regulates lateral root patterning. Dev. Cell *26*,
965   136–147.

966 42. Grison, M.S., Kirk, P., Brault, M.L., Wu, X.N., Schulze, W.X., Benitez-Alfonso, Y., Immel, F.,
967   and Bayer, E.M. (2019). Plasma Membrane-Associated Receptor-like Kinases Relocalize to
968   Plasmodesmata in Response to Osmotic Stress. Plant Physiol. *181*, 142–160.

969 43. Abel, N.B., Buschle, C.A., Hernandez-Ryes, C., Burkart, S.S., Deroubaix, A.-F., Mergner,
970   J., Gronnier, J., Jarsch, I.K., Folgmann, J., Braun, K.H., et al. (2021). A hetero-oligomeric
971   remorin-receptor complex regulates plant development. bioRxiv, 2021.01.28.428596.
972   https://doi.org/10.1101/2021.01.28.428596.

973 44. Liu, J., Elmore, J.M., and Coaker, G. (2009). Investigating the functions of the RIN4 protein
974   complex during plant innate immune responses. Plant Signal. Behav. *4*, 1107–1110.

975 45. Cheng, G., Yang, Z., Zhang, H., Zhang, J., and Xu, J. (2020). Remorin interacting with
976   PCaP1 impairs Turnip mosaic virus intercellular movement but is antagonised by VPg. New
977   Phytol. *225*, 2122–2139.

978 46. Codjoe, J.M., Richardson, R.A., McLoughlin, F., Vierstra, R.D., and Haswell, E.S. (2022).
979   Unbiased proteomic and forward genetic screens reveal that mechanosensitive ion channel
980   MSL10 functions at ER-plasma membrane contact sites in Arabidopsis thaliana. Elife *11*,
981   e80501.

982 47. Haswell, E.S., Peyronnet, R., Barbier-Brygoo, H., Meyerowitz, E.M., and Frachisse, J.-M.
983   (2008). Two MscS homologs provide mechanosensitive channel activities in the
984   Arabidopsis root. Curr. Biol. *18*, 730–734.

985 48. Diet, A., Link, B., Seifert, G.J., Schellenberg, B., Wagner, U., Pauly, M., Reiter, W.-D., and
986   Ringli, C. (2006). The Arabidopsis root hair cell wall formation mutant lrx1 is suppressed by
987   mutations in the RHM1 gene encoding a UDP-L-rhamnose synthase. Plant Cell *18*, 1630–
988   1641.

49. Baumberger, N., Ringli, C., and Keller, B. (2001). The chimeric leucine-rich repeat/extensin
cell wall protein LRX1 is required for root hair morphogenesis in Arabidopsis thaliana.
Genes Dev. *15*, 1128–1139.

50. Ringli, C., Bigler, L., Kuhn, B.M., Leiber, R.-M., Diet, A., Santelia, D., Frey, B., Pollmann, S.,
and Klein, M. (2008). The modified flavonol glycosylation profile in the Arabidopsis rol1
mutants results in alterations in plant growth and cell shape formation. Plant Cell *20*, 1470–
1481.

51. Field, S., Dorone, Y., Dwyer, W.P., Cox, J.A., Raba, D., Froehlich, J., Blea, M., Carmo,
O.M.S., Wallace, I.S., Hastings, R., et al. (2024). Arabidopsis thaliana RHAMNOSE 1
condensate formation drives UDP-rhamnose synthesis. bioRxivorg.
https://doi.org/10.1101/2024.02.15.580454.

52. Saffer, A.M., and Irish, V.F. (2018). Flavonol rhamnosylation indirectly modifies the cell wall
defects of RHAMNOSE BIOSYNTHESIS1 mutants by altering rhamnose flux. Plant J. *94*,
649–660.

53. Xiao, C., Zhang, T., Zheng, Y., Cosgrove, D.J., and Anderson, C.T. (2016). Xyloglucan
deficiency disrupts microtubule stability and cellulose biosynthesis in Arabidopsis, altering
cell growth and morphogenesis. Plant Physiol. *170*, 234–249.

54. Du, J., Kirui, A., Huang, S., Wang, L., Barnes, W.J., Kiemle, S.N., Zheng, Y., Rui, Y., Ruan,
M., Qi, S., et al. (2020). Mutations in the pectin methyltransferase QUASIMODO2 influence
cellulose biosynthesis and wall integrity in Arabidopsis. Plant Cell *32*, 3576–3597.

55. Saffer, A.M., Baskin, T.I., Verma, A., Stanislas, T., Oldenbourg, R., and Irish, V.F. (2023).
Cellulose assembles into helical bundles of uniform handedness in cell walls with abnormal
pectin composition. Plant J. *116*, 855–870.

56. Wang, T., McFarlane, H.E., and Persson, S. (2016). The impact of abiotic factors on
cellulose synthesis. J. Exp. Bot. *67*, 543–552.

57. Wang, C., Li, J., and Yuan, M. (2007). Salt tolerance requires cortical microtubule
reorganization in Arabidopsis. Plant Cell Physiol. *48*, 1534–1547.

58. Endler, A., Kesten, C., Schneider, R., Zhang, Y., Ivakov, A., Froehlich, A., Funke, N., and
Persson, S. (2015). A Mechanism for Sustained Cellulose Synthesis during Salt Stress.
Cell *162*, 1353–1364.

59. Kesten, C., García-Moreno, Á., Amorim-Silva, V., Menna, A., Castillo, A.G., Percio, F.,
Armengot, L., Ruiz-Lopez, N., Jaillais, Y., Sánchez-Rodríguez, C., et al. (2022). Peripheral
membrane proteins modulate stress tolerance by safeguarding cellulose synthases. Sci.
Adv. *8*, eabq6971.

60. Crowell, E.F., Bischoff, V., Desprez, T., Rolland, A., Stierhof, Y.-D., Schumacher, K.,
Gonneau, M., Höfte, H., and Vernhettes, S. (2009). Pausing of Golgi bodies on
microtubules regulates secretion of cellulose synthase complexes in Arabidopsis. Plant Cell
21, 1141–1154.

61. Iraki, N.M., Bressan, R.A., Hasegawa, P.M., and Carpita, N.C. (1989). Alteration of the
physical and chemical structure of the primary cell wall of growth-limited plant cells adapted

to osmotic stress. Plant Physiol. *91*, 39–47.

62. Raffaele, S., Mongrand, S., Gamas, P., Niebel, A., and Ott, T. (2007). Genome-wide
annotation of remorins, a plant-specific protein family: evolutionary and functional
perspectives. Plant Physiol. *145*, 593–600.

63. Ma, Z., Sun, Y., Zhu, X., Yang, L., Chen, X., and Miao, Y. (2022). Membrane nanodomains
modulate formin condensation for actin remodeling in Arabidopsis innate immune
responses. Plant Cell *34*, 374–394.

64. Segev-Zarko, L.-A., Dahlberg, P.D., Sun, S.Y., Pelt, D.M., Kim, C.Y., Egan, E.S., Sethian,
J.A., Chiu, W., and Boothroyd, J.C. (2022). Cryo-electron tomography with mixed-scale
dense neural networks reveals key steps in deployment of Toxoplasma invasion machinery.
PNAS Nexus *1*, gac183.

65. Tyanova, S., Temu, T., and Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nat. Protoc. *11*, 2301–2319.

66. Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M.Y., Geiger, T., Mann, M., and Cox,
J. (2016). The Perseus computational platform for comprehensive analysis of (prote)omics
data. Nat. Methods *13*, 731–740.

67. Vizcaíno, J.A., Côté, R.G., Csordas, A., Dianes, J.A., Fabregat, A., Foster, J.M., Griss, J.,
Alpi, E., Birim, M., Contell, J., et al. (2013). The PRoteomics IDEntifications (PRIDE)
database and associated tools: status in 2013. Nucleic Acids Res. *41*, D1063–D1069.

1048

A

B

C

Control   1 M sorbitol

D

Control

Plasma membrane   Cell wall   merge

XY   YZ   XY   YZ   YZ

1 M sorbitol

Plasma membrane   Cell wall   merge

XY   YZ   XY   YZ   YZ

E

Hechtian structure

3D   3D

cell surface

z

cell interior

Wall-membrane
attachment sites

3D
oblique slice

Figure 1. Hyperosmotic stress reveals cell wall-plasma membrane attachment sites in Arabidopsis roots.

(A) A 5-d-old Arabidopsis seedling grown on MS media. Box demarcates the root tip. Scale bar: 0.2 cm.

(B) Diagram of the longitudinal section of an Arabidopsis root tip in (A). Epidermal cells in the elongation zone are color coded in blue, which represent the regions where images in (C) were taken.

(C) 3D rendering of root epidermal cells in 5-d-old seedlings expressing a plasma membrane marker, YFP-LTI6b, under the control condition or treated with 1 M sorbitol for 5 min. Scale bar: 10 μm.

(D) Laser scanning confocal images of root epidermal cells from 5-day-old seedlings expressing the plasma membrane marker YFP-LTI6b and co-stained with the cell wall dye calcofluor white. Seedlings were imaged under control conditions or after treatment with 1 M sorbitol for 5 min. XY and YZ projections are shown. The positions where YZ projections were made are indicated by arrowheads. Scale bar: 5 μm.

(E) 3D segmentation of Hechtian structure and wall-membrane attachment sites from the dashed box region in (D). Segmented Hechtian structure is shown either in yellow, overlaid on YFP-LTI6b signals, or with depth-based color coding. Wall-membrane attachment sites (indicated by white dashed circles) are visualized by applying an oblique slicer (outlined in red) to the YZ projection. Scale bars: 2 μm. See also Video S1.

Figure 2. Reductions in cellulose and rhamnose impact root growth under hyperosmotic stress in opposing ways.

(A) Root growth of wild type, cellulose-deficient mutants, and a *CESA3* complementation line 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 27$ seedlings per genotype per treatment.

(B) Root growth of wild type and xyloglucan mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 20$ seedlings per genotype per treatment.

(C) Root growth of wild type and homogalacturonan mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 25$ seedlings per genotype per treatment.

(D) Root growth of wild type and rhamnogalacturonan-I mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 16$ seedlings per genotype per treatment.

(E) Root growth of wild type and rhamnogalacturonan-II mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 29$ seedlings per genotype per treatment.

Error bars indicate SD. ns, no significance; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by two-way ANOVA.

A

YFP-LTI6b | BF | merge | segmented (ratio)

0.5 M sorbitol

Col-0 — 0.88

cesa3^{ie5} — 0.73

rhm1-1 — 0.97

B

Plasmolysis ratio

Percent cells (%)

- Col-0 a
- *cesa3^{ie5}* b
- *rhm1-1* c

C

Biotin incubation → Pull-down → Tryptic digestion → Mass spec

Intensity / m/z

D

TbID-YFP-LTI6b/*cesa3^{ie5}* vs. TbID-YFP-LTI6b/Col-0

−log10 (P value)

log2 (fold change)

TbID-YFP-LTI6b/Col

TbID-YFP-LTI6b/*cesa3^{ie5}*

TOL1

E

TbID-YFP-LTI6b/*rhm1-1* vs. TbID-YFP-LTI6b/Col-0

−log10 (P value)

log2 (fold change)

TbID-YFP-LTI6b/Col

TbID-YFP-LTI6b/*rhm1-1*

CSI3
CMU1
CSI1
PATROL1
CESA6
CESA1
CESA5

F

GFP-CESA3

Col-0 | *cesa6^{prc1-1}* | *rhm1-1*

G

GFP-CESA3 particle density (cm^{−2})

Col-0 a | *cesa6^{prc1-1}* b | *rhm1-1* c

Figure 3. Deficiency in cellulose or rhamnose has opposite effects on plasmolysis and the plasma membrane proteome.

(A) Laser scanning confocal images and manual tracings of plasmolyzed root epidermal cells in 5-d-old seedlings expressing YFP-LTI6b in Col-0, $cesa3^{je5}$, or $rhm1-1$ background, respectively. Plasmolysis was induced by 0.5 M sorbitol for 5 min. Each displayed ratio corresponds to the same individual cell. Scale bar: 10 μm.

(B) Histogram showing plasmolysis ratio (protoplast area to cell area) in root epidermal cells of 5-d-old seedlings expressing YFP-LTI6b in Col-0, $cesa3^{je5}$, or $rhm1-1$ background, respectively. Different letters indicate significant differences by one-way ANOVA and Tukey's test. $n \geq 113$ cells from at least 10 seedlings per genotype.

(C) Schematic workflow of the proximity labeling-based mass spectroscopy.

(D) Volcano plot showing differentially enriched proteins in the plasma membrane proteome of $cesa3^{je5}$ compared to the wild type Col-0. Significantly enriched or depleted proteins are color coded in blue. Pre-filtering steps removed genetic background from each TbID-YFP-LTI6b-labeled samples using Welch t-test with an FDR of 0.05 and an S0 of 0.5. Pair-wise comparison was then conducted between TbID-YFP-LTI6b/$cesa3^{je5}$ and TbID-YFP-LTI6b/Col-0 using Student's t-test with an FDR of 0.05 and an S0 of 0.5. $n \geq 3$ biological replicates per genotype.

(E) Volcano plot showing differentially enriched proteins in the plasma membrane proteome of $rhm1-1$ compared to the wild type Col-0. Significantly enriched or depleted proteins are color coded in orange. Pre-filtering steps removed genetic background from each TbID-YFP-LTI6b-labeled samples using Welch t-test with an FDR of 0.05 and an S0 of 0.5. Pair-wise comparisons were then conducted between TbID-YFP-LTI6b/$rhm1-1$ and TbID-YFP-LTI6b/Col-0 using Student's t-test with an FDR of 0.05 and an S0 of 0.5. $n \geq 3$ biological replicates per genotype.

(F) Spinning disk confocal images of GFP-CESA3 in root epidermal cells of 5-d-old seedlings of Col-0, $cesa6^{prc1-1}$, and $rhm1-1$. The lower panels show magnified views of the regions outlined by white dashed boxes in the corresponding upper panels. Scale bars: 5 μm (upper panels), 1 μm (lower panels).

(G) Quantification of GFP-CESA3 particle density at the plasma membrane from (F). Error bars indicate SD. Different letters indicate significant differences by one-way ANOVA and Tukey's test. $n \geq 19$ cells from at least 10 seedlings per genotype.

Figure 4. CSC density at the plasma membrane safeguards wall-membrane attachments and plant growth under hyperosmotic stress.

(A) Spinning disk confocal images of root epidermal cells in 5-d-old seedlings expressing GFP-LTI6b and tdTomato-CESA6 under the control condition or treated with 1 M sorbitol for 5 min. XY and YZ projections are shown. The positions where YZ projections were made are indicated by arrowheads. Scale bar: 5 µm.

(B) Quantification of Manders overlap coeffiicient in 5-d-old seedlings of GFP-LTI6b/tdTomato-CESA6 under the control condition or treated with 1 M sorbitol for 5 min. Error bars indicate SD. $n \geq 10$ cells per genotype per treatment. *$p < 0.05$ by Student's t-test.

(C) Laser scanning confocal images and manual tracings of plasmolyzed root epidermal cells in 5-d-old Col-0 seedlings expressing YFP-LTI6b, treated with or without 100 nM isoxaben (ISX) for 20 minutes. Plasmolysis was subsequently induced by treatment with 0.5 M sorbitol for 5 minutes. Each displayed ratio corresponds to the same individual cell. Scale bar: 10 µm.

(D) Histogram showing plasmolysis ratio in root epidermal cells in 5-d-old Col-0 seedlings expressing YFP-LTI6b, treated with or without 100 nM ISX for 20 minutes. $n \geq 32$ cells from at least 5 seedlings per treatment. **$p < 0.01$ by Student's t-test.

(E) Histogram showing plasmolysis ratio induced by 0.5 M sorbitol in root epidermal cells of 5-d-old seedlings expressing YFP-LTI6b in Col-0 or $cesa6^{prc1-1}$ background, respectively. $n \geq 59$ cells from at least 10 seedlings per genotype. ****$p < 0.0001$ by Student's t-test.

(F) 3D renderings and YZ projections of root epidermal cells of 5-d-old YFP-LTI6b seedlings treated with or without 100 nM ISX for 20 minutes. Plasmolysis was subsequently induced by treatment with 1 M sorbitol for 5 minutes. Scale bar: 10 µm.

(G) Root growth of wild type and $shou4-3$ $4l-1$ mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 34$ seedlings per genotype per treatment.

(H) Histogram showing plasmolysis ratio induced by 0.5 M sorbitol in root epidermal cells of 5-d-old seedlings expressing YFP-LTI6b in Col-0 or $shou4-3$ $4l-1$ background, respectively. $n \geq 79$ cells from at least 9 seedlings per genotype. ****$p < 0.0001$ by Student's t-test.

(I) Root growth of wild type and $cesa3^{je5}$ $rhm1-1$ mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 42$ seedlings per genotype per treatment.

(J) Effects of ISX on root growth in wild type, $rhm1-1$, and $shou4-3$ $4l-1$ mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+) in the presence or absence of 1 nM ISX. $n \geq 19$ seedlings per genotype per treatment.

Error bars indicate SD. **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$ by two-way ANOVA.

Figure 5. The rapidly forming REM nanodomains function as negative regulators of wall-membrane attachments

(A) Laser scanning confocal images in root epidermal cells of 5-d-old seedlings expressing YFP-REM1.2 treated with 0 M, 0.28 M, 0.5 M, or 1 M sorbitol for 5 min, respectively. Scale bar: 10 μm.

(B) Bar graph showing YFP-REM1.2 nanodomain density at different sorbitol concentrations quantified from (A). $n \geq 21$ cells from at least 5 seedlings per treatment. Error bars indicate SD. Different letters indicate significant differences by one-way ANOVA and Tukey's test.

(C) Laser scanning confocal images of root epidermal cells in 5-d-old seedlings expressing YFP-REM1.2 co-stained with calcofluor white under the control condition or treated with 1 M sorbitol for 5 min. XY and YZ projections are shown. The positions where YZ projections were made are indicated by arrowheads. Scale bar: 10 μm.

(D) Laser scanning confocal images of root epidermal cells in 5-d-old seedlings expressing YFP-REM1.2 and mCherry-LTI6b under the control condition or treated with 1 M sorbitol for 5 min. XY and YZ projections are shown. The positions where YZ projections were made are indicated by arrowheads. Scale bar: 10 μm.

(E) Quantification of Manders overlap coeffiicient in 5-d-old seedlings of YFP-REM1.2/mCherry-LTI6b under the control condition or treated with 1 M sorbitol for 5 min. Error bars indicate SD. $n \geq 9$ cells per genotype per treatment. ns, no significance by Student's t-test.

(F) Root growth of wild type, *rem1.2 1.3 1.4* triple mutants, and *YFP-REM1.2/rem1.2 1.3 1.4* lines 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 27$ seedlings per genotype per treatment. Error bars indicate SD. ns, no significance; ***$p < 0.001$, ****$p < 0.0001$ by two-way ANOVA.

(G) Laser scanning confocal images and manual tracings of plasmolyzed root epidermal cells in 5-d-old seedlings expressing YFP-LTI6b in Col-0 or *rem1.2 1.3 1.4* background, respectively. Plasmolysis was induced by 0.5 M sorbitol for 5 min. Each displayed ratio corresponds to the same individual cell. Scale bar: 10 μm.

(H) Histogram showing plasmolysis ratio in root epidermal cells of 5-d-old seedlings expressing YFP-LTI6b in Col-0 or *rem1.2 1.3 1.4* background, respectively. $n \geq 102$ cells from at least 10 seedlings per genotype. ****$p < 0.0001$ by Student's t-test.

A

Control | 1 M sorbitol

YFP-REM1.2 | tdTomato-CESA6 | merge | YFP-REM1.2 | tdTomato-CESA6 | merge

XY YZ | XY YZ | XY YZ | XY YZ | XY YZ | XY YZ

B

Control | 1 M sorbitol

C

Manders' coefficient

**** ****

a a a a | a' b' c' c'

Control 1 M | Control 1 M

CESA3 ⇄ CESA6 | REM1.2 ⇄ CESA6

D

YFP-REM1.2

Col-0 | cesa3^je5 | rhm1-1

E

YFP-REM1.2 nanodomain density (μm⁻²)

a a b

Col-0 cesa3^je5 rhm1-1

F

GFP-CESA3

Col-0 | rem1.2 1.3 1.4

G

GFP-CESA3 particle density (μm⁻²)

***

Col-0 | rem 1.2 1.3 1.4

H

Col-0 | rem1.2 1.3 1.4

Root growth (cm)

**** **** | **** ****

57% | 84% | 52% | 83%

Mock 1 nM ISX | Mock 1 nM ISX
- + - + | - + - +

I

Root growth (cm)

**** ****

56% | 79%

Col-0 | cesa3^je5 rem1.2 1.3
- + | - +

Figure 6. REMs antagonize CSCs to regulate hyperosmotic stress responses.
(A) Spinning disk confocal images of root epidermal cells in 5-d-old seedlings expressing YFP-REM1.2 and tdTomato-CESA6 under the control condition or treated with 1 M sorbitol for 5 min. XY and YZ projections are shown. The positions where YZ projections were made are indicated by arrowheads. Scale bar: 5 μm.
(B) Magnified views of the regions outlined by white dashed boxes in (A). Scale bar: 2 μm.
(C) Quantification of Manders overlap coeffiicient in 5-d-old seedlings of GFP-CESA3/tdTomato-CESA6 or YFP-REM1.2/tdTomato-CESA6 under the control condition or treated with 1 M sorbitol for 5 min. Error bars indicate SD. $n \geq 6$ cells per genotype per treatment. In each genotype, different letters indicate significant differences by one-way ANOVA and Tukey's test. ****$p < 0.0001$ by Student's t-test when comparing the two genotypes under the same condition.
(D) Laser scanning confocal images of YFP-REM1.2 nanodomains in 5-d-old seedlings of the Col, *cesa3^{je5}*, or *rhm1-1* background, respectively when treated with 0.5 M sorbitol for 5 min. Scale bar: 10 μm.
(E) Quantification of YFP-REM1.2 nanodomain density from (C). $n \geq 54$ cells from at least 10 seedlings per genotype. Error bars indicate SD. Different letters indicate significant differences by one-way ANOVA and Tukey's test.
(F) Spinning disk confocal images of GFP-CESA3 in root epidermal cells of 5-d-old seedlings of Col and *rem1.2 1.3 1.4*. Scale bar: 5 μm.
(G) Quantification of GFP-CESA3 particle density at the plasma membrane from (E). $n = 27$ cells from at least 6 seedlings per genotype. Error bars indicate SD. ***$p < 0.001$ by Student's t-test.
(H) Effects of ISX on root growth in wild type and *rem1.2 1.3 1.4* mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+) in the presence or absence of 1 nM ISX. $n \geq 15$ seedlings per genotype per treatment.
(I) Root growth of wild type and *cesa3^{je5} rem1.2 1.3* mutants 1 day after transfer to either fresh MS media (-) or MS media supplemented with 0.28 M sorbitol (+). $n \geq 28$ seedlings per genotype per treatment.
Error bars indicate SD. ****$p < 0.0001$ by two-way ANOVA.

Figure 7. REM nanodomains harbor many other proteins in the vicinity, including SHOU4/4L.

(A) Workflow of filtering steps to identify proteins in the vicinity of TbID-YFP-REM1.2 in the presence or absence of 0.5 M sorbitol.

(B) and (C) Volcano plots showing differentially enriched proteins comparing TbID-YFP-REM1.2/Col-0 and Col-0 (WT) under the control (ctrl) condition (B) or treated with 0.5 M sorbitol for 15 min (C). Significantly enriched proteins are color coded using one-sided Welch's t-test with an FDR of 0.05 and an S0 of 0.5. $n \geq 3$ biological replicates.

(D) Volcano plot showing differentially enriched proteins comparing 0.5 M sorbitol treatment and the control condition in TbID-YFP-REM1.2/Col-0. Pair-wise comparison was conducted using Student's t-test with an FDR of 0.05 and an S0 of 0.5. $n \geq 3$ biological replicates.

(E) Laser scanning confocal images of tobacco leaf epidermal cells co-expressing YFP-REM1.2 with free mRFP, mRFP-REM1.3, SHOU4-mRFP, or SHOU4L-mRFP, respectively, under control conditions. Scale bar: 5 μm.

(F) Measurements of YFP lifetime from (E). Error bars indicate SD. Different letters indicate significant differences by one-way ANOVA and Tukey's test. $n \geq 23$ regions of interest from at least 3 infiltrated plants per construct combination.

(G) Laser scanning confocal images of tobacco leaf epidermal cells co-expressing YFP-REM1.2 with free mRFP, mRFP-REM1.3, SHOU4-mRFP, or SHOU4L-mRFP, respectively, upon 1 M sorbitol treatment for 10 min. Scale bar: 5 μm.

(H) Measurements of YFP lifetime from (E). Error bars indicate SD. Different letters indicate significant differences by one-way ANOVA and Tukey's test. $n \geq 20$ regions of interest from at least 3 infiltrated plants per construct combination.