Approaching an Error-Free Diploid Human Genome Assembly of East Asian Origin

Yanan Chu[1,6], Zhuo Huang[1,2,6], Changjun Shao[1,6], Shuming Guo[3,6], Yiji Yang[1,2], Xinyao Yu[1,2], Jian Wang[1], Yabin Tian[4], Jing Chen[1], Ran Li[5], Yukun He[5], Jun Yu[2*], Jie Huang[4*], Zhancheng Gao[3,5*], Yu Kang[1,2,7*].

1. Beijing Institute of Genomics, Chinese Academy of Sciences, and China National Center for Bioinformation, Beijing, 100101, China.

2. University of Chinese Academy of Sciences, Beijing, 100049, China.

3. Linfen Clinical Medicine Research Center, Institute of Chest and Lung Diseases, Shanxi Medical University, Linfen, 041000, China

4. State Key Laboratory of Drug Regulatory Science, National Institutes for Food and Drug Control, Beijing, 100050, China

5. Department of Respiratory and Critical Care Medicine, Human Genome Research Center, Peking University People's Hospital, Beijing 100044, China

6. These authors contributed equally

7. Lead contact

*Correspondence: Yu Kang, kangy@big.ac.cn, Zhancheng Gao, zcgao@bjmu.edu.cn, Jie Huang, jhuang5522@nifdc.org.cn, Jun Yu, junyu@big.ac.cn.

SUMMARY

Achieving an error-free diploid human genome remains a major challenge. We report T2T-YAO v2.0, a near-perfect assembly of a Han Chinese individual. To systematically evaluate assembly accuracy, we developed Sufficient Alignment Support (SAS), an alignment-based metric that detects structural and base-level errors in windows lacking adequate read support. Building on this, we established a "structural error–first" polishing strategy, correcting misassemblies with ultra-long ONT reads, then refining bases with PWC (Platform-integrated Window Consensus), a multi-platform, window-based polishing pipeline. Together, SAS and PWC provide a rigorous framework for

genome assessment and correction, resolving all detectable structure and non-homopolymer-related errors outside ribosomal DNA clusters. The resulting assembly contains no erroneous 21-mers absent from sequencing data, meeting k-mer–based criteria for an error-free genome. T2T-YAO v2.0 delivers the most perfect East Asian reference to date, enabling precise annotation, benchmarking, and variant discovery—laying a stronger foundation for human genomics and precision medicine.

KEYWORDS

Error-free genome, assembly quality assessment, near-perfect genome, SAS (Sufficient Alignment Support), erroneous k-mer, human diploid genome, Han Chinese, T2T-YAO, ultra-long ONT reads, homopolymer, PWC (Platform-integrated Window Consensus)

INTRODUCTION

Advancements in sequencing technologies and assembly algorithms have rapidly improved genome assembly quality, culminating in the release of the first telomere-to-telomere (T2T) human reference genome, T2T-CHM13, by the Telomere-to-Telomere (T2T) Consortium in 2022(Church, 2022; Nurk et al., 2022), which achieved unprecedent completeness and correctness. Nonetheless, assembly errors persist, arising from sources such as base-calling inaccuracies, incorrect haplotype assignment of contigs, imperfect read correction, and homopolymer compression during the assembly process(Mc Cartney et al., 2022; Rhie et al., 2021). As precision medicine advances, there is a growing demand for a diploid genome assembly with perfect base-level accuracy to serve as a benchmark for both sequencing and variant calling(Aganezov et al., 2022; Formenti and Howe, 2022; Jarvis et al., 2022; Olson et al., 2023), which is still a major challenge. While the last few years have seen dramatic improvements in base-calling accuracy across all major sequencing platforms of ONT, PacBio, and short-read sequencing, an interesting question arises: how close the current technologies can bring us to a perfect or error-free human genome assembly, whose accuracy ensures reliable downstream annotation and analysis?

The remaining errors in an initial assembly can be typically corrected *post hoc* by
realigning sequencing reads to the assembly and polishing regions of inconsistency. To
achieve this, a variety of structural and single-nucleotide variant (SNV) callers currently
available can be applied in parallel to identify and correct assembly errors based on
alternative allele detection. Then the accuracy of the resulting polished assembly is
typically evaluated using k-mer–based metrics such as Merqury, which quantifies
erroneous k-mers absent from the raw sequencing data(Rhie et al., 2020). However,
this polishing- evaluation strategy remains not efficient or accurate enough to achieve a
truly error-free human genome. Several limitations in current polishing strategies are
apparent. First, there is no comprehensive and reliable method to identify assembly
errors that can effectively guide correction. K-mer–based approaches are especially
inadequate for high-quality assemblies, as they tend to miss errors that do not introduce
novel k-mers, such as homopolymer length mismatches. Moreover, sequencing errors
can obscure genuine erroneous k-mers, making QV values highly sensitive to
sequencing depth and platform-specific error profiles. Second, uncorrected structural
errors can compromise read alignment, leading to failed corrections or overcorrection at
the base level. Third, no streamlined and accurate pipeline exists for correcting base-
level errors: false positives and false negatives are inevitable with current SNV callers.
These tools are typically optimized for haploid references and single-platform datasets,
often resulting in suboptimal performance on diploid genomes and posing challenges for
integrating calls across multiple sequencing platforms.

To address these issues, we developed an alignment-based quality assessment metric
called Sufficient Alignment Support (SAS), which directly identifies regions of the
assembly lacking adequate read support, an intrinsic indicator of assembly error.
Building on this, we implemented a "Structural Error-First" (SE-First) polishing strategy,
which prioritizes the correction of structural errors using ultra-long Oxford Nanopore
Technology (ONT) reads to improve the alignment of shorter, high-accuracy reads for
subsequent base-level corrections, then introduced PWC (Platform-integrated Window
Consensus), a multi-platform, window-based polishing pipeline to refine bases. Starting
from our previously published T2T-YAO v1.1, an initial complete diploid genome

assembly of a Han Chinese individual(He et al., 2023), we incorporated additional
sequencing data from high-accuracy platforms and iteratively applied SAS evaluation
and PWC polishing with SE-First principle. This approach yielded T2T-YAO v2.0, a
near-error-free diploid human genome, with the exception of residual issues in
ribosomal DNA (rDNA) clusters and a small number of homopolymer-associated errors
in repetitive regions. Based on Merqury analysis, T2T-YAO v2.0 contains no erroneous
21-mers absent from the raw sequencing data, yielding an "infinite" quality value (QV)
by k-mer standards, satisfying k-mer–based criteria for an error-free genome.

99
100 **RESULTS**
101

SAS: an alignment-based metric for assembly errors

To systematically evaluate the quality of genome assemblies, we developed a sensitive
alignment-based metric termed Sufficient Alignment Support (SAS). Rather than
detecting inconsistencies between the assembly and aligned sequencing reads, SAS
identifies assembly errors by flagging genomic windows that lack sufficient support from
mapped reads. In SAS, base-level errors (BEs; <50 bp) and structural errors (SEs; ≥50
bp) are quantified independently (**Figure 1A**).

For BEs, the diploid genome assembly is divided into 50 bp windows. Windows are
flagged as erroneous if they are not precisely supported by reads exceeding 10% of the
overall mean sequencing depth across any platform. This threshold is relaxed to 10% of
the local sequencing depth in specific low-coverage contexts: (1) regions within 10 kb of
chromosomal ends (*i.e.*, telomeres); (2) sequences containing degenerative GnA or
CnT repeats (GA/CT content ≥90% and GC content ≥60%); or (3) GC-rich regions (GC
content ≥90%), which are known to be underrepresented in polymerase-dependent,
high-accuracy sequencing platforms (*e.g.*, Element, Illumina, and PacBio platforms) and
rely predominantly on ONT reads for coverage.

SEs are assessed in 2 kb non-overlapping windows using ultra-long ONT reads for its
steady coverage without obvious sequencing bias. A window is considered structurally
erroneous if it meets one of the following conditions: (1) sequencing depth is outside the
20% to 200% range of the mean depth; or (2) fewer than 50% of local-depth reads span
the window without large indels (≥50 bp). This threshold is relaxed in chromosomal
termini (within 10 kb), where ≥10% of the local-depth reads must provide contiguous
support. Additionally, to improve detection of misjoin events, 30 kb sliding windows
(stepped every 2 kb) are evaluated. If no reads span a given window, the corresponding
15 consecutive 2 kb windows are flagged as SEs. Mitochondrial sequence, which is
small (~16 kb) and typically free of structural errors, are excluded from this analysis.

The overall SAS-based quality value (QV_{SAS}) is calculated by summing the BE and SE
windows and computing the predicted error rate (E) using the formula:

$$QV_{sas} = -10 \cdot \log E$$
$$E = \frac{BE + SE}{L}$$

Where E is the estimated error rate of the assessed assembly, BE and SE are the
number of non-redundant BE and SE windows, respectively, L is the base length of the
assembly.

To accurately evaluate the T2T-YAO assembly, we upgraded its raw data to high-
accuracy sequencing platforms: 260 Gb Q28 ultra-long ONT reads(Ni et al., 2023)
(filtered for >30 kb; N50 >120 kb), 172 Gb Pore-C(Deshpande et al., 2022), 306 Gb HiFi
reads (PacBio SPRQ; filtered >3 kb), and 320 Gb Element AVITI(Arslan et al., 2024)
Q50 reads (**Table S1**). To reduce ambiguity in haplotype assignment during mapping,
ONT and SPRQ reads were phased in advance using haplotype-specific k-mers
(hapmers) and mapped in a haplotype-aware manner (see Methods).

We applied the SAS metric to evaluate the previously released T2T-YAO v1.1 assembly
and compared the results to those obtained using Merqury, the most widely adopted k-

mer–based quality assessment tool (**Table 1**). SAS identified 21,511 base-level error (BE) windows and 4,438 structural error (SE) windows, yielding QV_{SAS} values of 53.12 and 54.13 for the maternal and paternal haplotypes, respectively. Chromosome-specific QV_{SAS} scores ranged from 45.36 to 61.37 (**Table S2**). In contrast, Merqury using 21-mers reported only 22,556 erroneous k-mers, corresponding to 1,074 estimated errors and Q67.42 across the diploid genome. This quality is notably lower than initially reported(He et al., 2023), as the use of more accurate Element and SPRQ reads revealed assembly errors that were previously obscured by sequencing errors, demonstrating the dependency of k-mer based QV on platform accuracy. Nonetheless, the much more errors identified by SAS than Merqury highlights the markedly greater sensitivity of SAS. Many BE windows detected by SAS lacked corresponding erroneous k-mers, even when more sensitive 31-mers were used, which identified 123,085 erroneous k-mers corresponding to 3,970 estimated errors and Q61.74 (**Table 1**). These results underscore the limitations of k-mer–based methods in capturing all types of assembly errors. As a representative example, we showcased the paternal chromosome 1 to illustrate the differences in error detection between SAS and other metrics (**Figure 2A**).

Closer inspection of BE windows revealed a frequent association with homopolymer sequences, where reads often disagreed on the number of repeated nucleotides. To classify these regions, we defined homopolymer sites as stretches of ≥10 A/T, ≥7 G/C mononucleotide repeats, or ≥10 dinucleotide repeats. Accordingly, BEs were divided into homopolymer-associated (h-BEs) and non-homopolymer-associated (nh-BEs), accounting for 17.17% (3,694) and 82.83% (17,817) of all BEs, respectively (**Table 1**). While nh-BEs were relatively enriched in centromeric regions and corresponded with k-mer–derived errors identified by Merqury, h-BEs were more broadly distributed outside centromeres and were largely undetected by k-mer-based methods (**Figure 2A**). Conversely, some erroneous k-mers reported by Merqury—using the Element-SPRQ hybrid k-mer set, following the protocol in T2T-CHM13(Mc Cartney et al., 2022)—were found outside SAS-flagged BE windows, clustering instead within degenerative GnA or CnT repeat regions. These loci typically had sufficient ONT coverage and were not

flagged as errors by SAS, but dropped out by SPRQ and Element reads due to
platform-specific sequencing bias. These discrepancies underscore the fundamental
limitations of k-mer–based assessment tools: their inability to detect errors that do not
introduce novel k-mers, such as homopolymer length discrepancies, and their
susceptibility to false positives in regions with biased or zero coverage. The latter arises
from Merqury's reliance on the hybrid k-mer sets derived exclusively from high-accuracy
Illumina (Element in this study) and PacBio platforms, while excluding ONT data to
avoid introducing large number of sequencing error-related k-mers given its relatively
higher sequencing error rate.

Given Merqury's limited ability to detect structural errors, we compared SAS with other
tools, including Sniffles2(Smolka et al., 2024) (ONT), the widely used structural variant
(SV) callers, and coverage-based structural error detectors— Flagger(Liao et al., 2023)
(ONT) and nucflag(Vollger et al., 2019) (SPRQ). Sniffles2 identified only 263 confident
SV-like errors (supported by ≥60% of local reads), demonstrating limited sensitivity,
particularly in repetitive regions (**Figure S1**), where it relies on repeat annotations often
unavailable in just completed assemblies. Flagger and nucflag are widely inconsistent
with each other, that they identify 4.03 Mb and 12.30 Mb of collapsed regions (>50 bp),
respectively, with only 1.08 Mb of overlap (**Table S3**). However, 83.63% of their
overlapping regions are either within or adjacent (<10 kb) to SE windows identified by
SAS. Furthermore, 78.50% of low-coverage regions jointly flagged by Flagger
("erroneous" and "duplication" regions) and nucflag ("misjoin" regions) are also
detectable by SAS. Notably, nucflag frequently misclassifies misjoins in GnA/CnT
repeats and other regions dropped out by SPRQ due to sequencing bias (**Figure 2A**),
which obvious damages the reliability of SE calling based on PacBio reads. Haplotypic
switch errors, which are flagged by nucflag as heterogeneous regions and by Merqury
as switch k-mers, are not treated as a separate error class in SAS. Depending on the
read evidence, such events are either classified as SEs if flanking clips are present or
as BEs if disagree with phased read but without a clear breakpoint.

Correction of Structural Errors

Using SAS and ultra-long ONT reads, we identified 4,438 non-redundant SE windows in the T2T-YAO v1.1 assembly. These windows were further categorized based on their underlying features: 1,295 were high-depth regions, 1,662 low-depth regions, 331 contained indels, 61 were marked by clipped alignments, and 1,929 were consecutive windows without spanning reads. Note that individual windows could be assigned to multiple categories. Errors related to clipped alignments and abnormal depth were particularly concentrated in the short arms of acrocentric chromosomes and the centromeres of maternal chromosomes 6 and 7, whereas indels were more evenly distributed across the genome (**Figure 3A**).

Given the critical importance of accurate read alignment for subsequent base-level error correction, we adopt a "Structural Error-first" (SE-first) strategy. This approach prioritizes correction of structural errors using ONT reads before applying PWC base-level polishing with high-accuracy sequencing data. By resolving large-scale mis-assemblies upfront, the SE-first strategy reduce the risk of overcorrection due to artificial alignments in shorter reads, thereby enhancing the overall fidelity of the assembly.

We employed a comprehensive correction pipeline combining alternate variant calls from Sniffles2, local reassemblies using ONT reads, and targeted telomere extension to eliminate all detectable SEs. After each correction round, reads were re-aligned, and SAS was reapplied to reassess error content (**Figure 1B**). Most SEs were resolved after several iterations of this correction–realignment–evaluation loop, though certain sites of clipped alignment required manual inspection and correction. When local assembly fails to resolve mis-assemblies, incorporation of Pore-C chromosomal configuration data often enables correction. The resulting contact map validates the structural integrity of the final assembly, confirming accurate haplotype phasing and the correct pairing of short and long arms across the ten acrocentric chromosomes, a particularly challenging structural feature to resolve (**Figure S2**). In telomeric and subtelomeric regions, ONT

reads often exhibited heterogeneity that impeded successful reassembly. In such cases,
stratifying ONT reads based on characteristic SNVs and assembling the dominant
haplotype group enabled resolution. Following these polishing efforts, all telomeres
were extended to an average length of 8.7 ± 2.6 kb, with all exceeding 5 kb except for
three in the q arms of maternal chromosome 12 and paternal chromosomes 21 and 22
(**Figure 4A**). The telomeres in v2.0 are closer to conventionally accepted lengths, and
far more complete than those in T2T-CHM13 (2.7 ± 0.7 kb).

Ultimately, T2T-YAO v2.0 becomes free of detectable SEs outside of ribosomal DNA
(rDNA) clusters (**Figure 2B, 3A**). Centromeric collapses, particularly in the higher-order
repeat (HOR) arrays of maternal chromosomes 6 and 7, are also resolved
(**Supplemental information file 1**, **Figure 4B, Figure S3A**). One potential exception is
observed in a 12 kb high-depth region upstream of the *TSPAN33* gene on maternal
chromosome 7q32. Although the local ONT depth (84–89×) marginally exceeds the
global cutoff (84×), the local SPRQ depth (45×) is quite near mean depth (51.2 ± 13.5),
and the sequences there and of flanks are fully supported by ONT and SPRQ reads
without signs of clip or misjoin. Thus, the reason for the elevated ONT depth in this
particular region is still unknown. Further inspection reveals that this region is composed
of L1M and Alu elements which sequences are widely present in human genome with
millions of copies(Belancio et al., 2009). We suspect that the repetitive nature of this
region may contribute to the elevated depth where 22.8% of mapped reads are shorter
than 60 kb and exhibiting MAPQ =1 (**Figure S4**).

Aside from this, ONT read depth is within the 20%~200% mean depth range and follows
a Gaussian distribution, with extremely few regions tallied by Flagger as false
duplication (2.81 Mb, 0.047%) or collapsed (1.55 Mb, 0.026%) (**Figure 5A**), with ~90%
present in rDNA clusters. All overlapping issue regions identified by Flagger and nucflag
are confined to rDNA arrays or telomeric regions, supporting the structural integrity of
T2T-YAO v2.0 (**Table S3**). To further validate continuity, we assessed the fraction of the
assembly covered by non-split ONT reads (NScov%). In T2T-YAO v1.1, NScov% is

99.98%, with ~0.91 Mb (excluding rDNA) left uncovered. In contrast, T2T-YAO v2.0

accomplishes 100% coverage by non-split ONT reads when rDNA is masked,

confirming the absence of unresolved SEs outside rDNA.

Depth fluctuations in v1.1 are much more pronounced in Element and SPRQ compared

to ONT (53.5 ± 42.0 and 51.2 ± 13.5 vs. 42.4 ± 8.1, averaged in 2 kb windows),

reflecting their greater susceptibility to artificial alignments due to shorter read lengths.

Such falsely collapsed alignments frequently introduce incorrect variant calls and

promote overcorrection during base-level polishing. In T2T-YAO v1.1, alignment of

SPRQ and Element reads revealed widespread regions of abnormally high depth (>2-

fold mean depth), totaling 4.71 Mb and 2.52 Mb, respectively, enriched for artificial

alignments (**Figure 5B**). These misaligned reads often contain consensus mismatches

to the reference and represent a major source of overcorrections. After SE correction,

the collapsed regions were reduced to 3.43 Mb (SPRQ) and 0.93 Mb (Element),

including those within rDNA (**Figure 5B**). Thus, SE correction reduced false alignments

in high-accuracy data, mitigating the risk of over-correction and improving reliability in

the polished assembly.

Correction of Base-level Errors

Using ONT, SPRQ, and Element sequencing data, we identified 21,511 base-level error

(BE) windows in T2T-YAO v1.1, comprising 3,694 homopolymer-associated BEs (h-

BEs) and 17,817 non-homopolymer-associated BEs (nh-BEs), broadly distributed

across all chromosomes (**Figure 3B**). Each flagged window is corrected using PWC

pipeline, which generates consensus sequence (supported by ≥60% reads) from

spanning reads (depth ≥10) for each flagged window. PWC follows an across-platform

prioritization scheme, where consensus supported by all platforms is directly applied. In

case where the three sequencing platforms disagree, corrections supported by two

platforms are adopted for nh-BEs, otherwise, platform-specific reliability is applied in the

order: Element > SPRQ > ONT. If no consensus is obtainable, ONT reads spanning the

region are locally reassembled to correct the sequence, which often enables consensus

generation by mitigating artificial alignments (**Figure 1C**). For h-BEs, local ONT-based

consensus or assembly is omitted due to the known limitations of ONT sequencing in

accurately resolving homopolymer lengths (**Figure 6A**). The stringent conditions for

generating consensus minimized overcorrection caused by alignment artifacts during

the polishing rounds. Read alignment and polishing are repeated iteratively until the

number of BEs is stabilized.

To correct additional base-level errors potentially missed by SAS, we applied multiple

variant callers: Deeppolisher(Mastoras et al., 2024b) (SPRQ), DeepVariant(Poplin et al.,

2018a; Shafin et al., 2021) (ONT and Element), and GATK HaplotypeCaller(Poplin et

al., 2018b) (Element) to call variants by mapping reads to the T2T-YAOv1.1 diploid

genome. These tools typically reported several thousand to over twenty thousand

homozygous variants (*i.e.* supported by the majority of aligned reads from the

corresponding platform), yielding a total of 63,398 non-redundant calls with genotype

quality (GQ) ≥20, with a little overlap among tools. Because obvious false positives

were present, we first filtered out variants whose reference sequence, along with 10 bp

flanking sequences, were supported by ≥60% local depth and ≥10 exact-matching

reads from any platforms. This filtering step yielded 43,444 potential SNV-like errors

across both haplotypes, of which 40.21% were associated with homopolymer regions,

an even higher proportion than the observed rate of h-BEs (**Figure 5C**). The elevated

frequency of homopolymer-associated errors likely stems from the homopolymer

compression step used in the initial T2T-YAO assembly with Verkko(Antipov et al.,

2025; Rautiainen et al., 2023). Of these SNV-like errors, 43.90% of non-homopolymer-

associated variants overlapped with SAS-detected nh-BEs, whereas only 7.08% of

homopolymer-related calls overlapped with SAS h-BEs. This discrepancy highlights the

challenge of identifying true errors via alignment, particularly those associated with

homopolymers. To avoid overcorrection from alternative sequences proposed by SNV

callers, we applied the same PWC pipeline to potential SNV-like errors outside BE

windows: replacing the variant site and its 10 bp flanking sequence with a read-derived

consensus sequence rather than the variant caller's prediction. In many cases, the

consensus sequence was identical to the reference, meaning that only a small
proportion of sites beyond BE windows are truly errors could be corrected.

After several rounds of iterative polishing, T2T-YAO v2.0 contains only 336 nh-BEs (all
within rDNA regions) and 837 h-BEs located in repetitive regions (including 489 in
rDNA) that remain unresolved due to the inherent limitations of current sequencing
technologies (**Table 1, Figure 3B**). When we reassess the v2.0 assembly using the
same SNV callers, the total number of filtered SNV-like potential errors (homozygous
SNV with GQ ≥20) drops significantly to 2,354 (**Figure 5C**). Of these, only 38 are non-
homopolymer-associated, with 36 (94.74%) located within rDNA units, consistent with
the distribution of nh-BEs identified by SAS. The remaining two non-homopolymer calls
outside rDNA are confirmed to be false positives supported by reads of other platforms.
The sharp drop in SNV-like errors without extensive corrections suggests that many of
them, particularly outside BE windows are likely false calls derived from artificially
aligned reads bearing variant sequences.

An illustrative example of unresolved h-BEs can be found in the 3.3 kb D4Z4 repeat
array on chromosome 4q35(Lemmers et al., 2018). In this region, the polyC tracts
between DUX4-like genes can not be confidently resolved: ONT and SPRQ reads
struggled with homopolymer sequencing, while Element reads suffered from
sequencing bias and mapping ambiguity (**Figure 6B**). Nonetheless, the remaining h-
BEs account for only 0.04% of the total 2,412,242 homopolymer sites genome-wide.
Although not completely resolved, T2T-YAOv2.0 provides an accurate count and
distribution of homopolymer sites in a human genome. Among these, polyA/polyT tracts
(≥10 bp) comprised 81.70%, while polyC/polyG tracts (≥7 bp) and poly-dinucleotide
repeats (≥10 bp) represented 9.20% and 9.10%, respectively. Overall, the distribution of
homopolymers clearly avoids centromeric regions and exhibit very similar pattern
across haplotypes (**Figure S5**). Notably, polyC/polyG and poly-dinucleotide repeats are
approximately 10-fold enriched in rDNA sequences (up to ~600/Mb), whereas

polyA/polyT tracts are much sparser in the short arms of acrocentric chromosomes,
encompassing rDNA.

Toward an Error-Free Human Genome Assembly

We re-evaluated the quality of the T2T-YAO v2.0 assembly using both SAS and
Merqury metrics, based on the same sequencing datasets (**Table 1**). The final SAS
assessment identified 1,773 structural error (SE) windows (all confined to rDNA regions
except for the six windows of high-depth in maternal chromosome 7 as described
above) and 1,175 base-level error (BE) windows, comprising 827 within rDNA and 348
homopolymer-associated BEs (h-BEs) outside rDNA. The resulting QV_{SAS} scores were
66.00 (maternal) and 61.25 (paternal). When rDNA regions were masked, leaving only
the 348 h-BEs, the QV_{SAS} values rose to 71.62 and 71.81, respectively. (**Table 1, Table**
S2, Supplemental information file 1).

T2T-YAO v2.0 contains tens of kilobases of GnA/CnT repeat sequences that are poorly
covered by Element and SPRQ reads but well supported by ONT data. Due to the
limited representation of these sequences in the k-mer set used by Merqury, a high
number of false positives are called in these loci. Specifically, Merqury identified 3,006
erroneous 21-mers (corresponding to a QV of 76.18) and 28,956 erroneous 31-mers
(QV 68.03) in the diploid assembly. Since all erroneous 21-mers are encompassed by
the erroneous 31-mers, we focused our inspection on the 31-mers. These were
consistently found in regions with near-zero coverage (depth ≤1) in Element and SPRQ
data but showed strong support from ONT reads. The 31-mers clustered into 1,127
discrete regions (after merging overlaps), collectively spanning 76,248 bp. Of these,
70% regions (totaling 53,307 bp in 674 regions) had extremely high GA/CT content
(>90%), while the rest were composed of non-canonical 6-mers found in telomeric
sequences or GC-rich tandem repeats, patterns consistent with known sequencing
biases(Freeman et al., 2020; Ross et al., 2013) (**Table S4**). Given the presence of these
sequences in human genome that are dropped out by PacBio and short-read

sequencing (SRS), achieving Q100 quality is unattainable with the current PacBio/SRS

hybrid k-mer set, even with the complete correction of all assembly errors.

The ONT data used in this study were generated using the R10 flow cell, known for high

base-calling accuracy (Q28; >99.8%). To leverage this accuracy, we incorporated k-

mers from the ONT dataset into our analysis, stringently filtering out k-mers of

occurrence ≤4 to exclude those of potential sequencing errors, compared to ≤1 for

Element and SPRQ reads. We first use this expanded trinary k-mer set to assess T2T-

YAO v1.1, which QV only shows a slight improvement over the binary set, indicating

minimal inclusion of k-mers caused by ONT sequencing errors. When we applied the

trinary k-mer set to assess T2T-YAO v2.0, the QV for 21-mers increased to "infinite,"

representing an Merqury-defined error-free diploid genome. The QV for 31-mers also

improved significantly, reaching 96.29 for the maternal haplotype and 88.73 for the

paternal, with only 143 erroneous 31-mers remaining (**Table 1**). All paternal erroneous

31-mers are confined to rDNA regions, while those of maternal are located in two h-BE

loci of GnA repeat regions containing several polyG tracts (≥8 bases), corresponding

precisely with the residual errors detected by the SAS metric.

Ensuring the accuracy of annotation

We generated the gene annotation of T2T-YAO v2.0 by transferring annotations from

T2T-CHM13 v2.0 using Liftoff(Shumate and Salzberg, 2021). rDNA clusters and Ig/TCR

VDJ segments, which are highly variable among individuals, were annotated separately

and excluded from gene count statistics. Of the 57,377 genes annotated in T2T-CHM13,

56,601 (98.65%) were successfully lifted to T2T-YAO, including most protein-coding

genes, lncRNAs, and pseudogenes. A total of 115 protein-coding genes (0.59%) failed

to lift over in either haplotype, while lncRNAs and pseudogenes showed higher loss

rates, 272 (1.66%) and 339 (1.99%), respectively (**Table 2, Table S5**). To recover the

missed protein-coding genes, we applied miniprot (Li, 2023) which enabled the

annotation of additional genes and extra gene copies based on MANE(Morales et al.,

2022)(v1.4) protein-to-genome alignments, adding 14 genes (11 absent in T2T-CHM13)

and 30 extra copies. In total, 773 genes, including 112 protein-coding genes, were
absent in T2T-YAO compared to T2T-CHM13. Notably, all missing protein-coding genes
were either uncharacterized ("LOC" genes) or members of gene families such as
FAM90 and *DEFB*, which are often confused among paralogues when lifting over due to
high sequence similarity. The overall lift-over rate was slightly lower than that of T2T-
HG002(Hansen and al, 2025), where only 39 distinct genes (3 protein-coding) were
missing across both haplotypes. Since both HG002 and CHM13 are of European origin,
this discrepancy may reflect population-related differences in gene content, particularly
in the number and sequence of paralogues. As expected, protein-coding genes are the
most conserved, whereas lncRNAs and pseudogenes exhibit greater variability between
any genomes (**Table 2**). Most lifted-over genes in T2T-YAO v2.0 retain synteny with
their counterparts in T2T-CHM13, with a few exceptions, particularly on chromosome Y
(**Figure S6**). In total, 4,143 extra copies of unique genes were identified in the diploid
T2T-YAO, of which only 9.5% are protein-coding (**Table 2**, **Table S5**). Notably, about
one-third of the extra lncRNA copies are located on chromosome Y, with their biological
significance largely unknown (**Figure S7**).

For the protein-coding genes successfully lifted in T2T-YAO, we identified 254,704
transcripts in both haplotypes. Among these, 194,042 (76.18%), including 374 corrected
by Lifton (Chao et al., 2025), encode protein sequences identical to their T2T-CHM13
counterparts (**Table S6**). Of the transcripts with altered protein sequences, 458 (0.75%)
showed markedly reduced sequence identity (<80%), suggesting potential impairment
of protein function. Considering both alleles, 363 non-redundant transcripts
corresponding to 200 genes exhibit either decreased identity (<80%) or absence in both
haplotypes. These findings highlight ancestry-related differences in gene content and
function, emphasizing the need to refine and complete the annotation of T2T-YAO v2.0
through additional strategies beyond liftover, such as *de novo* gene prediction,
transcriptome sequencing from diverse East Asian tissues and multiple lineages
differentiated from iPSC-YAO cells.

Comprehensive correction of SEs and nh-BEs in T2T-YAO v2.0 eliminated all
detectable errors in centromeres, which naturally lack homopolymer sites, providing
unprecedented resolution of centromeric sequences. We refined the annotation of
centromeres, higher-order repeats (HORs), and other satellite elements across all 46
chromosomes in T2T-YAO v2.0 (**Figure 4B, Supplement Information file 1**) (Altemose
et al., 2022; Logsdon et al., 2024). Compared to v1.1, subtle but significant differences
can be observed, exemplified by updates to the paternal chromosome 1 assembly
(**Figure S3B**), highlighting the importance of high-fidelity polishing for accurate
centromere characterization. In v2.0, we identified 536 active HOR monomer types,
comprising 320,424 sequences. These sequences showed limited within-type variance,
with 86.23% of sequences being 100% identical to their type consensus and 98.61%
showing <5 bp variance (identity >97%). Comparison of HOR variants between
maternal vs. paternal autosomes identified three major haplotype-specific patterns (>90
repeats), two in paternal chromosome 17 and one in maternal chromosome 6. Besides
these, T2T-YAO has five additional major patterns absent from both T2T-CHM13 and
CHM1, but lacks two compared to them (**Table S7**). The presence of these patterns
suggests population-specific differences in HOR variants, warranting further
investigation into centromere diversity across populations.

Using the recently developed Genomic Centromere Profiling (GCP) pipeline, a method
analogous to fragment length analysis for characterizing the position, orientation, and
organization of centromere protein B (CENP-B) binding motifs(Corda and Giunta, 2025),
we constructed structural models for centromeric repeat clusters in T2T-YAO. These
models confirmed the correctness of the v2.0 centromere assemblies and revealed
extensive variation in centromere size, structure, and repeat composition among
chromosomes(Logsdon et al., 2024). Although maternal and paternal centromeres of
the same chromosome generally exhibited similar organization, clear inter-haplotype
differences were observed in repeat size. For instance, the HORs of maternal
chromosomes 6 and 7 were substantially expanded compared to their paternal
counterparts in T2T-YAO (**Figure S3A**).

Implications in accurate benchmark and variant calling

High-quality, ancestry-matched references are known to enhance variant discovery by improving read mapping (Aganezov et al., 2022; Guo et al., 2024). With T2T-YAO, the completely assembled centromeres now enable systematic analysis of variation in these previously inaccessible regions. We first examined the small variants between T2T-YAO's paternal and maternal haplotypes in 100 kb windows of alignable regions across autosomes. Genome-wide, the density of small indels (1-3bp) was 0.14/kb and SNP density was 0.89/kb with a Ti/Tv ratio of 1.78. Centromeric regions showed strikingly different patterns, reflecting extensive haplotype divergence as illustrated in paternal chromosome 1(**Figure 7A**). The amount of alignable sequence dropped sharply, especially in satellite DNA outside HORs. Meanwhile SNP density was markedly elevated, 4.2-fold in HORs (3.77/kb) and 3.0-fold in other satellites (2.65/kb). Most variants in HORs arose from inactive or divergent HOR units, accompanied by a reduced Ti/Tv ratio of 0.72, consistent with previous report (Logsdon et al., 2024). Indels were rare, occurring at 0.08/kb in HORs and 0.06/kb in other satellites, compared with 0.17/kb in other repeat regions (**Table S8**).

When aligned T2T-YAO to the ancestry-unmatched T2T-CHM13, SNP and indel densities rose across the genome, with the strongest effects in centromeres, which showed further reduced alignability and higher variant density (**Figure 7B, Table S8**). While overall Ti/Tv kept relatively stable, the distribution of variant types across repeats shifted with the reference, highlighting the importance of ancestry-matched references for accurate variant calling in complex regions.

Coding regions were strongly constrained. SNP density in CDS was 0.46/kb—half the genome average—and indel density was 7.65/Mb, about one-twentieth. The Ti/Tv ratio rose to 2.65, reflecting purifying selection, and the missense-to-synonymous ratio (dN/dS) of SNPs was reduced, both annotated using VEP(McLaren et al., 2016).

However, using unmatched or lower-quality references of CHM13 and GRCh38, inflated variant density in CDS reduced dN/dS due to inflated synonymous calls from divergent alignments and selective depletion of missense variants (**Figure 7C**). Predictions with LOFTEE (Karczewski et al., 2020), which relies on population variant data coordinated in GRCh38, identified only 91 SNPs and 35 indels as high-confidence loss-of-function (LOF) variants, none in CDS regions (**Figure 7D**). Most LOF variants were homozygous and affected uncharacterized or gene family members, consistent with genes missed in liftover, implying potential confusions in variant calling among paralogs.

Finally, telomeres in T2T-YAO v2.0 were extended and corrected to error-free status across all chromosomes. Most contain non-canonical 6-mer repeats interspersed with canonical (TTAGGG)n motifs, especially at subtelomeric transitions (Lyčka et al., 2023) (**Figure 4B**). Motif composition and repeat length varied widely, without haplotype-, chromosome-, or arm-specific patterns, confirming earlier reports (Stephens and Kocher, 2024) but now with direct sequence evidence and strong read support, intriguing further investigation into telomere variations and their biological significance.

DISCUSSION

An error-free diploid human genome holds immense value for calibrating variant calling pipelines and enhancing the accuracy of clinical genetic diagnoses(Jarvis et al., 2022; Wagner et al., 2022). Following the completion of the T2T-CHM13 assembly, the "Q100" project was initiated with the ambitious goal of producing a diploid human genome of Phred quality score 100, equivalent to one error per 10 billion bases(Telomere-to-telomere-consortium, 2023). Given the approximate size of the human diploid genome (~6 billion bases), achieving Q100 implies constructing a truly error-free genome. The validation of T2T-YAO v2.0 by 100% coverage of non-split ONT reads except at rDNA regions and free of erroneous 21-mers absent from the trinary k-mer set, thus qualifying for an "infinite" QV under Merqury standards, further demonstrate the power of the SAS framework, PWC pipeline, and our "Structural Error-first" strategy. The resulting near perfect genome enables accurate annotation of genome components and benchmarks of variant calling. However, technical limitations

in both genome evaluation and error correction exist, preventing it from being deemed
absolutely error-free.

Achieving an error-free genome requires assembly quality metrics capable of evaluating
accuracy at such level. These metrics encompasses multiple dimensions, including
completeness, correctness, and continuity. Of these, completeness is relatively
straightforward to assess via k-mer-based approaches(Rhie et al., 2020). However,
accurate evaluation of correctness and continuity remains more challenging and
typically relies on alignment-based methods(Li et al., 2023; Mastoras et al., 2024a).
Most alignment-based tools identify discrepancies between the genome assembly and
realigned reads and confined to a single sequencing platform. These discrepancies
often include both false positives arising from artificial alignments of heterogeneous
reads, as well as false negatives missed due to sequencing dropout. Consequently,
integrating alignment discrepancies across platforms is not a trivial problem as simply
merging or intersecting them does not reliably identify true assembly errors.

To overcome these challenges, we developed SAS, which evaluate alignment support
rather than discrepancies with the assembly. This reverse strategy facilitates integration
of regions supported across multiple platforms, avoiding the need to set arbitrary cutoffs
to balance false positives and negatives. Assessing supportive alignments incurs a
greater computational burden than simply examining discrepancies embedded in the
output files of aligners. To improve efficiency, we first quickly filtered for regions
perfectly covered (depth ≥5) by Element reads without mismatches or clips, which
account for 99.2% of the T2T-YAO genome, thereby greatly reducing the overall
computational load. SAS implemented a window-based approach, which offers higher
sensitivity in identifying BEs than variation callers of single base resolution which read
support is occasionally unreliable particularly related to indels in repetitive regions. A
window (50-mer) is considered supported only if at least five reads (representing >10%
of the mean sequencing depth in our case) cover it exactly, a criterion more stringent
than k-mer-based methods like Merqury, which typically require only two occurrences of

21- or 31-mers. Structural errors can likewise be detected in the window-based strategy,

by flagging 2 kb windows lacking sufficient spanning reads, and scanning 30 kb

windows without spanning reads to verify continuity in broader genomic context.

Theoretically, an alignment-based approach can identify all assembly errors and correct

them according to the consensus of aligned reads—provided that the authentic

sequence is fully covered by reads and the reads are correctly mapped to their original

locations. However, artificial alignments are inevitable, especially with shorter reads,

compromising the accuracy of error detection and correction. Therefore, minimizing

alignment artifacts is crucial. Previously, MapQ (mapping quality, reflecting the

confidence that a read is placed in a unique place of the genome) was commonly used

to filter ambiguous alignments that are typically false alignments. However, MapQ is

ineffective in diploid genomes, as most reads receive a zero MapQ score due to the

high similarity between haplotypes as well as recent duplications within a haplotype.

This limitation explains why MapQ-dependent methods, such as GCI(Chen et al., 2024)

and DeepVariant(Shafin et al., 2021), often produce false calls. In this work, we apply

three strategies to reduce artificial alignments:1) A "SE–first" approach that leverages

ultra-long ONT reads to define genome structure, similar to the proprietary strategy

used in the chromosome X project(Miga et al., 2020). 2) Phasing ONT and SPRQ reads

into haplotypes based on the hapmers they contain prior to mapping. We use stringent

criteria—such as larger k-mers and significant hapmer differences—to avoid false

assignments caused by noise. 3)Applying filters to remove secondary alignments and

multi-split reads instead of MapQ. All together, these measures are more effective then

MapQ in reducing false alignment while preserving the correct ones.

Based on the near-perfect T2T-YAO v2.0, our results in variants between haplotypes

establish that ancestry-matched references are critical for accurate variant calling in

complex regions. When using unmatched genomes, SNP and indel densities were

artificially inflated, alignable fractions collapsed, and functional interpretations were

distorted. These effects were most pronounced in centromeres and paralogous genes,

where divergence across haplotypes exaggerated spurious variants and obscured true

biological differences. Even in coding regions, unmatched references skewed dN/dS by

inflating synonymous calls and underestimating missense variation, particularly in gene

family members. These findings underscore a broader principle: a single reference

genome cannot serve all populations. The ancestry-matched assemblies are more

powerful in reducing mapping artifacts, resolving repeats, and sharpening functional

inference than uniformed reference. As population-specific T2T references emerge,

they will provide not only more accurate baselines for variant discovery but also

essential resources for precise human genomics.

Limitations of the study

Despite its sensitivity, SAS still misses some base-level errors in repetitive regions

where misaligned short reads from other loci create false support. It also fails to detect

collapses or false duplications that fall within acceptable depth thresholds. These

problematic regions often show heterogeneous SNP patterns in the aligned

reads(Vollger et al., 2019), complicating distinction from sequencing noise, particularly

with ONT reads. A canonical example is the ribosomal DNA (rDNA) clusters, which

remain the final hurdle to a fully error-free human genome(Li and Durbin, 2024). In

these regions, ONT reads are long enough to span complete rDNA units but lack the

base-level accuracy to resolve unit-level SNVs, whereas SPRQ reads provide the

needed accuracy but are not long enough to span the entire unit. Thus, the primary

barriers to achieving a completely error-free diploid human genome are intrinsic to the

architecture of highly repetitive sequences, particularly their unit length and sequence

identity, and the current limitations of sequencing technologies in generating reads that

are both sufficiently long and accurate. Bridging this gap remains the final frontier in

complete human genome assembly.

RESOURCE AVAILABILITY

Lead contact

• Requests for further information and resources should be directed to and will be fulfilled by the
lead contact, Yu Kang (kangy@big.ac.cn).

Materials availability

- This study did not generate new unique reagents.

Data and code availability

ACKNOWLEDGMENTS

This study was supported by the grants of the National Key Research and Development Program of China (2024YFC3405701), the National Science Foundation of China (32371537).

AUTHOR CONTRIBUTIONS

Y.K. conceived and designed the project. Z.H. implemented the software SAS. Y.C. conducted the correction of T2T-YAO. S.G., C.S., J.W. performed sample preparation and sequencing. Y.Y.,X.Y., Y.T., J.C., R.L. and Y.H. participated in data analyses, and Y.K., Z.G., J.H., J.Y. wrote the manuscript. All the authors agree on the manuscript.

DECLARATION OF INTERESTS

All authors declare no competing interests.

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES

During the preparation of this work, the author(s) used Deepseek and chatGPT in order to improve language fluency, clarity, and readability. After using this tool or service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

FIGURE TITLES AND LEGENDS

Figure 1. Overview of the sufficient alignment support (SAS) metric for detecting assembly

errors and error correction pipeline. (**A**). SAS identifies structural errors (SEs) and base-level

errors (BEs) based on read alignment patterns across defined windows. Top panel (SE windows):

Structural errors are detected in 2 kb windows using ultra-long Oxford Nanopore (UL ONT) reads.

Windows are flagged based on five indicators: abnormally high or low depth, large insertions or

deletions (indels), clipped alignments, and lack of spanning reads. Telomeric regions with less

supportive reads are not flagged. Bottom panel (BE windows): Base-level errors are evaluated in 50

bp windows using reads from all high-accuracy sequencing platforms (Element, SPRQ, or UL ONT).

A window is considered error-free only if all bases are covered by perfectly matched spanning reads.

In regions not covered by Element reads, support from SPRQ or UL ONT reads is used to validate

sequence accuracy. SAS flags windows lacking sufficient read support (*i.e.*, <10% of average

sequencing depth), illustrating example regions containing various assembly errors. (**B**). Schematic

Overview of the Structural Error (SE) Correction Pipeline. (**C**). Schematic Overview of the Base-

Level Error (BE) Correction Pipeline.

Figure 2. Genomic distribution of assembly errors on paternal chromosome 1 before and

after polishing. Genomic features and assembly error profiles for T2T-YAO v1.1(**A**) compared to

the polished T2T-YAO v2.0 (**B**). Tracks from top to bottom: GC% and GA% content; normalized read

depth from Element, PacBio Revio (SPRQ), and ONT ultra-long reads; structural error (SE) windows

identified by SAS and Sniffles2; misjoins and collapses flagged by Nucflag and Flagger; base-level

errors (BEs) categorized as homopolymer-associated (h-BE) and non-homopolymer-associated (nh-

BE); erroneous 31-mers and 21-mers detected by Merqury with binary and trinary k-mer set.

Figure 3. Counts of base-level and structural errors in T2T-YAO before and after iterative

correction. Distribution of structural error (SE) windows (**A**) and base-level error (BE) windows (**B**)

across chromosomes in maternal and paternal haplotypes of T2T-YAO v1.1 compared to v2.0. SEs

were categorized into five types: low coverage, high coverage, indels (≥50 bp), clipped alignments,

and non-spanning regions. BEs are divided into homopolymer-associated BEs (h-BEs) and non-

homopolymer-associated BEs (nh-BEs). Bar plots show counts per chromosome, with the left bar in

each pair for v1.1 (light colors) and the right for v2.0 (dark colors). The rDNA-associated errors are

shown as hatched bars.

Figure 4. Structural characterization of telomeres and centromeres in the polished T2T-YAO

v2.0 assembly.

(**A**) Telomeric repeat content and extension across all 46 chromosomal termini. Each bar represents

the distance from the chromosomal terminus, with colored segments indicating specific 6-mer

tandem repeats. Canonical (TTAGGG)n (in long arm) and complementary (CCCTAA)n (in short arm)

motifs are shown in dark red and blue, along with shallow colors indicate non-canonical telomeric-

like 6-mer repeats. (**B**) Annotation of centromeric regions in all the 46 chromosomes. Repeat content

is visualized as stacked bars by annotated repeat family. Each pair of centromeres displays a unique

repeat composition and organization, with notable differences between haplotypes. The inset shows

a zoom-out view of the Y chromosome, which is full of repeat sequences.

Figure 5. Read depth distribution and SNV-like errors in T2T-YAO v1.1 and v2.0

(A) Depth distribution of ONT reads aligned to T2T-YAO v2.0, along with misassemblies predicted

by Flagger. (B) Comparison of read depth distributions for SPRQ and Element datasets in T2T-YAO

v1.1 and v2.0. (C) Quality score distribution of SNV-like errors identified by SNV callers in T2T-YAO

v1.1 and v2.0.

Figure 6. Unresolved homopolymer-related base-level errors

(**A**) Examples of discordance among ONT and SPRQ reads at homopolymer sites. (**B**) Example of

unresolved homopolymer sites in the D4Z4 repetitive region on chromosome 4q35. The sequencing

depth and read alignment from multiple sequencing platforms (ONT, SPRQ and Element) are

displayed with peaks above each track representing the sequencing depth at each position. The

gray horizontal bars indicate the sequencing reads, with colored vertical bars denoting mismatches

in these reads. The bottom tracks highlight the position of DUX4L genes and unresolved h-BE.

Figure 7. Impact of reference genome on variant calling

(A) Density of SNPs and small indels, and Ti/Tv ratio in 100 kb windows along paternal chromosome 1. Upper panel: whole chromosome; lower panel: centromere and flanking regions, with repeat annotations indicated in the background of the peak map. **(B)** Comparison of SNP and indel densities, and Ti/Tv ratios across different repeat classes using ancestry-matched (inter-haplotype) versus unmatched (T2T-CHM13) references. **(C)** Comparison of SNP and indel densities, Ti/Tv ratios, and missense-to-synonymous variant ratios in CDS regions across different references. **(D)** Counts of high-confidence loss-of-function variants per chromosome, shown separately for SNPs (upper panel) and small indels (lower panel). Heterozygous variants, present on a single haplotype, are indicated by hatched bars stacked above the homozygous variants.

Figure S1. Example of a structural error in a repetitive region missed by Sniffles2.

The vertical bar in top track indicates the structural error detected by Sniffles2 (missed in panels A and B, present in panel C but incomplete). The blue peaks above middle track represents the ONT sequencing depth at each position. The gray horizontal bars indicate the ONT sequencing reads, with colored vertical bars indicating mismatches. Black short lines within the reads indicate sequence deletions (i.e., base pair deletions relative to the reference genome). The purple vertical bars and the numbers above them indicate insertions and their length. The bottom SAS-SE track and dashed box highlight the location of structural errors detected by SAS pipeline.

Figure S2. Contact map of pore-c reads aligned to the T2T-YAO v2.0.

Figure S3. Comparison of centromere infrastructure of T2T-YAO v1.1 and v2.0.

(**A**) Comparison of centromere structures in T2T-YAO v1.1 and v2.0 via Genomic Centromere Profiling (GCP). (**B**) Patterns of higher-order repeats (HOR) and satellite sequences in the centromere of paternal chromosome 1 in T2T-YAO v1.1 and v2.0.

Figure S4. Repetitive elements in the SE (structural error) region on maternal chromosome 7 with abnormal ONT read depth.

The sequencing depth (blue peak for ONT and pink peak for SPRQ) and read alignment from ONT, SPRQ platforms are displayed across the SE region in maternal chromosome 7. The SAS-SE track

highlight the location of abnormal ONT read depth (dark red) detected by SAS pipeline. The bottom
tracks highlight the genomic features, including repeat motif and genes in this region.

Figure S5. Distribution of three major types of homopolymers in each chromosome of T2T-
YAO v2.0.

Figure S6. Synteny of the unique genes between T2T-CHM13 and T2T-YAO haplotype.

Synteny maps of (A) Maternal haplotype (MAT) and (B) paternal haplotype (PAT). Each dot
represents a gene shared between both genomes in the panel, colored according to gene transcript
sequence identity. The continuous diagonal line in panels indicates strong conservation of gene
order. Sparse off-diagonal dots suggest possible local rearrangements or transpositions, typically
accompanied by a slight reduction in sequence identity.

Figure S7. **Relative locations of extra gene copies in T2T-YAO v2.0 to their corresponding**
genes in T2T-CHM13.

Each line connects the position of an extra-copy gene in Maternal haplotype (MAT) **(A)** and
paternal haplotype (PAT) **(B)** of T2T-YAO to its corresponding gene position in T2T-CHM13, colored
with gene types.

Table S1. Sequencing datasets and phasing statistics for the evaluation of the T2T-YAO assembly.

Table S2. Chromosome-wise assembly quality metrics before and after polishing.

Table S3. Structural errors identified by multiple tools before and after correction.

Table S4. GC and GA/CT content of the remaining erroneous 31-mers in T2T-YAO v2.0.

Table S5. The copy number and location of each gene in T2T-CHM13, maternal and paternal
haplotype of T2T-YAO v2.0.

Table S6. Transcript variants and sequence identity between T2T-YAO v2.0 and T2T-CHM13

Table S7. Specific HOR units in T2T-YAO v2.0

Table S8. SNP and small indel in repetitive regions.

Supplemental Information file 1: S1_Figures 1-45. Genomic distribution of assembly errors on each
chromosome except paternal chromosome 1 before and after polishing

Supplemental Information file 2: S2_Figures 1-24. The composition of centromeric α-satellite

HOR units in each chromosome.

Table 1. Comparative quality metrics of the T2T-YAO assembly assessed by SAS and Merqury.

		v1.1			v2.0		
		mat	pat	Both	mat	pat	Both
SAS	QV (total)	53.12	54.13	53.59	66.00	61.25	63.04
	BE	11946	9565	21511	291	884	1175
	h-BE	1789	1905	3694	255	582	837
	nh-BE	10157	7660	17817	35	301	336
	SE	2752	1686	4438	469	1304	1773
	low_depth	920	742	1662	121	709	830
	high_depth	886	409	1295	302	376	678
	indel	141	190	331	9	49	58
	clip	39	22	61	6	24	30
	no spanning	1247	682	1929	99	469	568
	QV (rDNA masked)	53.65	55.39	54.42	71.62	71.81	71.72
	BE	10871	7863	18734	179	169	348
	h-BE	1426	1145	2571	179	169	348
	nh-BE	9445	6718	16163	0	0	0
	SE	2147	572	2719	6	0	6
	low_depth	730	57	787	0	0	0
	high_depth	643	115	758	6	0	6
	indel	116	125	241	0	0	0
	clip	23	8	31	0	0	0
	no spanning	1027	295	1322	0	0	0
Merqury-binary	QV-21mer	66.48	68.68	67.42	74.84	78.23	76.18
	err-kmer	14252	8304	22556	2085	921	3006
	total-kmer	$3.02{\times}10^{9}$	$2.92{\times}10^{9}$	$5.93{\times}10^{9}$	$3.02{\times}10^{9}$	$2.92{\times}10^{9}$	$5.93{\times}10^{9}$
	err rate	$2.25{\times}10^{7}$	$1.36{\times}10^{7}$	$1.81{\times}10^{7}$	$3.28{\times}10^{8}$	$1.50{\times}10^{8}$	$2.41{\times}10^{8}$
	switch err%	0.0107	0.0254	—	0.0017	0.0038	—
	Completeness%	97.770	93.708	99.839	97.777	93.710	99.850

	QV-31mer	**60.97**	**62.73**	**61.74**	**67.24**	**69.04**	**68.03**
	err-kmer	74864	48221	123085	17694	11262	28956
	total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
	err rate	8.00×10^7	5.34×10^7	6.69×10^7	1.89×10^7	1.25×10^7	1.57×10^7
	switch err%	0.0219	0.0413	—	0.0050	0.0072	—
	Completeness%	96.780	92.607	99.806	96.792	92.610	99.826
Merqury-trinary	**QV-21mer**	**67.7888**	**69.854**	**68.6823**	**inf**	**inf**	**inf**
	err-kmer	10543	6331	16874	0	0	0
	total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
	err rate	1.66×10^7	1.03×10^7	1.35×10^7	0	0	0
	switch err%	0.0107	0.0254	—	0.0017	0.0038	—
	Completeness%	97.7896	93.7264	99.8638	97.7966	93.7281	99.875
	QV-31mer	**62.4363**	**64.2025**	**63.2153**	**96.2941**	**88.7314**	**91.0965**
	err-kmer	53376	34337	87713	22	121	143
	total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
	err rate	5.71×10^7	3.80×10^7	4.77×10^7	2.35×10^7	1.34×10^7	7.77×10^7
	switch err%	0.0219	0.0413	—	0.0050	0.0072	—
	Completeness%	96.810	92.636	99.841	96.823	92.639	99.861

Table 2. Gene annotation of T2T-YAO v2.0 liftoff from T2T-CHM13

	Gene Type	CHM13	Unique	Success %	Extra copy	Failures	Failure %	Total	MANE Unique	MANE Extra
	Protein coding	19987	19842	99.27%	182	145	0.73%	20024	11	4
	LncRNA	15511	15291	98.58%	625	220	1.42%	15916	--	--
Maternal	Pseudogene	16599	16197	97.58%	674	402	2.42%	16871	--	--
	Others	3899	3806	97.61%	302	93	2.39%	4108	--	--
	Total	55996	55136	98.46%	1783	860	1.54%	56919	56930	56934
	Protein coding	19238	19087	99.22%	213	151	0.78%	19300	9	20
	LncRNA	16072	15773	98.14%	1258	299	1.86%	17031	--	--
Paternal	Pseudogene	16059	15651	97.46%	649	408	2.54%	16300	--	--
	Others	3733	3675	98.45%	240	58	1.55%	3915	--	--
	Total	55102	54186	98.34%	2360	916	1.66%	56546	56555	56575
	Protein coding	20090	19975	99.43%	--	115	0.57%	39324	14	30
	LncRNA	16388	16116	98.34%	--	272	1.66%	32947	--	--
Diploid	Pseudogene	16993	16654	98.01%	--	339	1.99%	33171	--	--
	Others	3906	3855	98.69%	--	51	1.31%	8023	--	--
	Total	57377	56600	98.65%	--	777	1.35%	113465	113479	113509

Box 1. Near-Perfect Genome

Thanks to advances in sequencing and computing technologies, human genomes can now be assembled almost exactly as they exist in nature. Apart from the challenging ribosomal DNA (rDNA) arrays—long stretches of nearly identical ~45 kb repeats—and the exact lengths of a few homopolymer runs, every other basepaire of the diploid human genome T2T-YAO can now be resolved from telomere to telomere. This achievement marks a major step beyond what was possible only a few years ago with the advent of T2T-CHM13.

Using diploid human genome as an example, we can see how close the current technologies bring us to an error-free or perfect eukaryotic genome:

1. **K-mer completeness**: >99.8%

2. **K-mer switch error rate** (diploid genome): <0.01%

In addition, the following criteria are met with rDNA excluded:

3.   **Coverage by non-split long reads** (N50 >100 kb): 100%

4.   **Number of 30 kb windows not spanned by long reads**: 0

5.   **Non-homopolymer sequences lacking sufficient read support**: 0

6.   **Homopolymer sites lacking sufficient read support**: <0.02% of total homopolymer
sites

Together, these metrics define the current benchmark for what can be considered a *near-*
perfect eukaryotic genome.

821
822

REFERENCES

Aganezov, S., Yan, S.M., Soto, D.C., Kirsche, M., Zarate, S., Avdeyev, P., Taylor, D.J., Shafin, K.,

Shumate, A., Xiao, C.*, et al.* (2022). A complete reference genome improves analysis of human genetic

variation. Science (New York, NY) *376*, eabl3533.

Altemose, N., Logsdon, G.A., Bzikadze, A.V., Sidhwani, P., Langley, S.A., Caldas, G.V., Hoyt, S.J.,

Uralsky, L., Ryabov, F.D., Shew, C.J.*, et al.* (2022). Complete genomic and epigenetic maps of human

centromeres. Science (New York, NY) *376*, eabl4178.

Antipov, D., Rautiainen, M., Nurk, S., Walenz, B.P., Solar, S.J., Phillippy, A.M., and Koren, S. (2025).

Verkko2 integrates proximity-ligation data with long-read De Bruijn graphs for efficient telomere-to-

telomere genome assembly, phasing, and scaffolding. Genome Res *35*, 1583–1594.

Arslan, S., Garcia, F.J., Guo, M., Kellinger, M.W., Kruglyak, S., LeVieux, J.A., Mah, A.H., Wang, H., Zhao,

J., Zhou, C.*, et al.* (2024). Sequencing by avidity enables high accuracy with low reagent consumption.

Nature Biotechnology *42*, 132–138.

Belancio, V.P., Deininger, P.L., and Roy-Engel, A.M. (2009). LINE dancing in the human genome:

transposable elements and disease. Genome Medicine *1*, 97.

Chao, K.H., Heinz, J.M., Hoh, C., Mao, A., Shumate, A., Pertea, M., and Salzberg, S.L. (2025).

Combining DNA and protein alignments to improve genome annotation with LiftOn. Genome Res *35*,

311–325.

Chen, Q., Yang, C., Zhang, G., and Wu, D. (2024). GCI: a continuity inspector for complete genome

assembly. bioRxiv, 2024.2004.2006.588431.

Church, D.M. (2022). A next-generation human genome sequence. Science (New York, NY) *376*, 34–35.

Corda, L., and Giunta, S. (2025). Chromosome-specific centromeric patterns define the centeny map of

the human genome. Science (New York, NY) *389*, eads3484.

Deshpande, A.S., Ulahannan, N., Pendleton, M., Dai, X., Ly, L., Behr, J.M., Schwenk, S., Liao, W.,

Augello, M.A., Tyer, C.*, et al.* (2022). Identifying synergistic high-order 3D chromatin conformations from

genome-scale nanopore concatemer sequencing. Nat Biotechnol *40*, 1488–1499.

Formenti, G., and Howe, K. (2022). An assembly line for an improved human reference genome. Nature

611, 519–+.

Freeman, T.M., Wang, D., and Harris, J. (2020). Genomic loci susceptible to systematic sequencing bias

in clinical whole genomes. Genome Res *30*, 415–426.

Guo, S., Huang, Z., Zhang, Y., He, Y., Chen, X., Wang, W., Li, L., Kang, Y., Gao, Z., Yu, J.*, et al.* (2024).

Enhancing Variant Calling in Whole-exome Sequencing Data Using Population-matched Reference

Genomes. Genomics Proteomics Bioinformatics *22*.

Hansen, N., and al, e. (2025). A complete diploid human genome benchmark for personalized genomics.

bioRxiv (to appear).

He, Y., Chu, Y., Guo, S., Hu, J., Li, R., Zheng, Y., Ma, X., Du, Z., Zhao, L., Yu, W.*, et al.* (2023). T2T-YAO:

A Telomere-to-telomere Assembled Diploid Reference Genome for Han Chinese. Genomics Proteomics

Bioinformatics *21*, 1085–1100.

Jarvis, E.D., Formenti, G., Rhie, A., Guarracino, A., Yang, C., Wood, J., Tracey, A., Thibaud-Nissen, F.,
Vollger, M.R., Porubsky, D.*, et al.* (2022). Semi-automated assembly of high-quality diploid human
reference genomes. Nature *611*, 519–531.

Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins, R.L., Laricchia,
K.M., Ganna, A., Birnbaum, D.P.*, et al.* (2020). The mutational constraint spectrum quantified from
variation in 141,456 humans. Nature *581*, 434–443.

Lemmers, R., van der Vliet, P.J., Vreijling, J.P., Henderson, D., van der Stoep, N., Voermans, N., van
Engelen, B., Baas, F., Sacconi, S., Tawil, R.*, et al.* (2018). Cis D4Z4 repeat duplications associated with
facioscapulohumeral muscular dystrophy type 2. Hum Mol Genet *27*, 3488–3497.

Li, H. (2023). Protein-to-genome alignment with miniprot. Bioinformatics (Oxford, England) *39*.

Li, H., and Durbin, R. (2024). Genome assembly in the telomere-to-telomere era. Nat Rev Genet *25*, 658–
670.

Li, K., Xu, P., Wang, J., Yi, X., and Jiao, Y. (2023). Identification of errors in draft genome assemblies at
single-nucleotide resolution for quality assessment and improvement. Nature Communications *14*, 6556.

Liao, W.W., Asri, M., Ebler, J., Doerr, D., Haukness, M., Hickey, G., Lu, S., Lucas, J.K., Monlong, J., Abel,
H.J.*, et al.* (2023). A draft human pangenome reference. Nature *617*, 312–324.

Logsdon, G.A., Rozanski, A.N., Ryabov, F., Potapova, T., Shepelev, V.A., Catacchio, C.R., Porubsky, D.,
Mao, Y., Yoo, D., Rautiainen, M.*, et al.* (2024). The variation and evolution of complete human
centromeres. Nature *629*, 136–145.

Lyčka, M., Bubeník, M., Závodník, M., Peska, V., Fajkus, P., Demko, M., Fajkus, J., and Fojtová, M.
(2023). TeloBase: a community-curated database of telomere sequences across the tree of life. Nucleic
Acids Research *52*, D311–D321.

Mastoras, M., Asri, M., Brambrink, L., Hebbar, P., Kolesnikov, A., Cook, D.E., Nattestad, M., Lucas, J.,
Won, T.S., Chang, P.-C.*, et al.* (2024a). Highly accurate assembly polishing with DeepPolisher. bioRxiv,
2024.2009.2017.613505.

Mastoras, M., Asri, M., Brambrink, L., Hebbar, P., Kolesnikov, A., Cook, D.E., Nattestad, M., Lucas, J.,
Won, T.S., Chang, P.C.*, et al.* (2024b). Highly accurate assembly polishing with DeepPolisher. bioRxiv.

Mc Cartney, A.M., Shafin, K., Alonge, M., Bzikadze, A.V., Formenti, G., Fungtammasan, A., Howe, K.,
Jain, C., Koren, S., Logsdon, G.A.*, et al.* (2022). Chasing perfection: validation and polishing strategies
for telomere-to-telomere genome assemblies. Nature methods *19*, 687–695.

McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P., and Cunningham, F.
(2016). The Ensembl Variant Effect Predictor. Genome biology *17*, 122.

Miga, K.H., Koren, S., Rhie, A., Vollger, M.R., Gershman, A., Bzikadze, A., Brooks, S., Howe, E.,
Porubsky, D., Logsdon, G.A.*, et al.* (2020). Telomere-to-telomere assembly of a complete human X
chromosome. Nature *585*, 79–84.

Morales, J., Pujar, S., Loveland, J.E., Astashyn, A., Bennett, R., Berry, A., Cox, E., Davidson, C.,

Ermolaeva, O., Farrell, C.M., *et al.* (2022). A joint NCBI and EMBL-EBI transcript set for clinical genomics
and research. Nature *604*, 310–315.

Ni, Y., Liu, X., Simeneh, Z.M., Yang, M., and Li, R. (2023). Benchmarking of Nanopore R10.4 and R9.4.1
flow cells in single-cell whole-genome amplification and whole-genome shotgun sequencing. Comput
Struct Biotechnol J *21*, 2352–2364.

Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., Vollger, M.R., Altemose, N.,
Uralsky, L., Gershman, A., *et al.* (2022). The complete sequence of a human genome. Science (New
York, NY) *376*, 44–53.

Olson, N.D., Wagner, J., Dwarshuis, N., Miga, K.H., Sedlazeck, F.J., Salit, M., and Zook, J.M. (2023).
Variant calling and benchmarking in an era of complete human genome sequences. Nature Reviews
Genetics *24*, 464–483.

Poplin, R., Chang, P.-C., Alexander, D., Schwartz, S., Colthurst, T., Ku, A., Newburger, D., Dijamco, J.,
Nguyen, N., Afshar, P.T., *et al.* (2018a). A universal SNP and small-indel variant caller using deep neural
networks. Nature Biotechnology *36*, 983–987.

Poplin, R., Ruano-Rubio, V., DePristo, M.A., Fennell, T.J., Carneiro, M.O., Van der Auwera, G.A., Kling,
D.E., Gauthier, L.D., Levy-Moonshine, A., Roazen, D., *et al.* (2018b). Scaling accurate genetic variant
discovery to tens of thousands of samples. bioRxiv, 201178.

Rautiainen, M., Nurk, S., Walenz, B.P., Logsdon, G.A., Porubsky, D., Rhie, A., Eichler, E.E., Phillippy,
A.M., and Koren, S. (2023). Telomere-to-telomere assembly of diploid chromosomes with Verkko. Nature
Biotechnology.

Rhie, A., McCarthy, S.A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W.,
Fungtammasan, A., Kim, J., *et al.* (2021). Towards complete and error-free genome assemblies of all
vertebrate species. Nature *592*, 737–746.

Rhie, A., Walenz, B.P., Koren, S., and Phillippy, A.M. (2020). Merqury: reference-free quality,
completeness, and phasing assessment for genome assemblies. Genome biology *21*, 245.

Ross, M.G., Russ, C., Costello, M., Hollinger, A., Lennon, N.J., Hegarty, R., Nusbaum, C., and Jaffe, D.B.
(2013). Characterizing and measuring bias in sequence data. Genome biology *14*, R51.

Shafin, K., Pesout, T., Chang, P.C., Nattestad, M., Kolesnikov, A., Goel, S., Baid, G., Kolmogorov, M.,
Eizenga, J.M., Miga, K.H., *et al.* (2021). Haplotype-aware variant calling with PEPPER-Margin-
DeepVariant enables high accuracy in nanopore long-reads. Nature methods *18*, 1322–1332.

Shumate, A., and Salzberg, S.L. (2021). Liftoff: accurate mapping of gene annotations. Bioinformatics
(Oxford, England) *37*, 1639–1643.

Smolka, M., Paulin, L.F., Grochowski, C.M., Horner, D.W., Mahmoud, M., Behera, S., Kalef-Ezra, E.,
Gandhi, M., Hong, K., Pehlivan, D., *et al.* (2024). Detection of mosaic and population-level structural
variants with Sniffles2. Nature Biotechnology *42*, 1571–1580.

Stephens, Z., and Kocher, J.-P. (2024). Characterization of telomere variant repeats using long reads
enables allele-specific telomere length estimation. BMC Bioinformatics *25*, 194.

Telomere-to-telomere-consortium (2023). HG002 Q100 project. In https://githubcom/marbl/hg002.

Vollger, M.R., Dishuck, P.C., Sorensen, M., Welch, A.E., Dang, V., Dougherty, M.L., Graves-Lindsay, T.A.,

Wilson, R.K., Chaisson, M.J.P., and Eichler, E.E. (2019). Long-read sequence and assembly of

segmental duplications. Nature methods *16*, 88–94.

Wagner, J., Olson, N.D., Harris, L., McDaniel, J., Cheng, H., Fungtammasan, A., Hwang, Y.C., Gupta, R.,

Wenger, A.M., Rowell, W.J.*, et al.* (2022). Curated variation benchmarks for challenging medically

relevant autosomal genes. Nat Biotechnol *40*, 672–680.

Figure 1

A

B

C

Figure 2

A

YAO v1.1 chr1_paternal

B

YAO v2.0 chr1_paternal

Figure 3

Figure 4

Figure 5

A

B

Chr4_paternal: 191,854,671-191,944,659

ONT

SPRQ

DUX4L
h-BE

0   5kb

Chr4_paternal: 191,860,741-191,880,744

ONT

SPRQ

Element

DUX4L
h-BE

0   2kb

Figure 7