Approaching an Error-Free Diploid Human Genome

Yanan Chu[1*], Zhuo Huang[1,2*], Changjun Shao[1*], Shuming Guo[3*], Xinyao Yu[1,2], Jian Wang[1], Yabin Tian[4], Jing Chen[1], Ran Li[5], Yukun He[5], Jun Yu[2#], Jie Huang[4#], Zhancheng Gao[3,5#], Yu Kang[1,2#].

1. Beijing Institute of Genomics, Chinese Academy of Sciences, and China National Center for Bioinformation, Beijing, 100101, China.

2. University of Chinese Academy of Sciences, Beijing, 100049, China.

3. Linfen Clinical Medicine Research Center, Institute of Chest and Lung Diseases, Shanxi Medical University, Linfen, 041000, China

4. State Key Laboratory of Drug Regulatory Science, National Institutes for Food and Drug Control, Beijing, 100050, China

5. Department of Respiratory and Critical Care Medicine, Human Genome Research Center, Peking University People's Hospital, Beijing 100044, China

Correspondence to: Yu Kang, kangy@big.ac.cn, Zhancheng Gao, zcgao@bjmu.edu.cn, Jie Huang, jhuang5522@nifdc.org.cn, Jun Yu, junyu@big.ac.cn.

Abstract

Achieving a truly error-free human diploid genome remains a central challenge in genomics. Here, we present T2T-YAO v2.0, a nearly perfect diploid genome assembly for a Han Chinese individual. We first developed Sufficient Alignment Support (SAS), a sensitive alignment-based metric that identifies both structural and base-level assembly erroneous windows by quantifying inadequate read support across sequencing platforms. Unlike conventional methods that detect erroneous k-mers (*e.g.*, Merqury) or inconsistencies between assemblies and aligned reads from single platform, SAS integrates supportive information from various platform and avoids false positives caused by sequencing bias and artificial alignments. Employing a "Structural Error-first" polishing strategy, prioritizing structural error correction using ultra-long Oxford Nanopore reads, we iteratively refined the assembly with high-accuracy PacBio and Element sequencing data. This approach eliminates all detectable structural and non-homopolymer-related base-level errors outside

ribosomal DNA clusters. The final T2T-YAO v2.0 assembly exhibits complete non-split ONT read coverage and free of erroneous 21-mers outside sequencing data, representing an error-free human genome by Merqury standards. Our results not only highlight the limitations of current genome quality assessment methods but also provide a roadmap for constructing high-fidelity diploid references essential for precision medicine and genome-based diagnostics.

Main Text

Introduction

Advancements in sequencing technologies and assembly algorithms have rapidly improved the quality of genome assemblies, culminating in the release of the first telomere-to-telomere (T2T) human reference genome, T2T-CHM13, by the Telomere-to-Telomere (T2T) Consortium in 2022[1,2]. As precision medicine advances, there is a growing demand for a diploid genome assembly with perfect base-level accuracy to serve as a benchmark for both sequencing and variant calling[3-6]. However, assembly errors can still arise from multiple sources, including base-calling inaccuracies, incorrect haplotype assignment of contigs, imperfect read correction, and homopolymer compression during the assembly process[7,8]. These errors must be corrected *post hoc* by realigning sequencing reads to the assembly and polishing regions of inconsistency.

Currently, a variety of structural and single-nucleotide variant (SNV) callers can be applied in parallel to identify and correct assembly errors based on alternative allele detection. The accuracy of the resulting polished assembly is typically evaluated using k-mer–based metrics such as Merqury, which quantifies erroneous k-mers absent from the raw sequencing data[9]. However, this polishing-evaluation strategy remains not efficient or accurate enough to achieve a truly error-free human genome. Several limitations in current polishing strategies are apparent. First, there is no comprehensive and reliable method to identify assembly errors that can effectively guide correction. K-mer–based approaches are especially inadequate for high-quality assemblies, as they tend to miss errors that do not introduce novel k-mers, such as homopolymer length mismatches. Moreover, sequencing errors can obscure genuine erroneous k-mers, making QV values highly sensitive to sequencing depth and platform-specific error profiles. Second, uncorrected structural errors can

compromise read alignment, leading to failed corrections or overcorrection at the base level. Third,

no streamlined and accurate pipeline exists for correcting base-level errors: false positives and false

negatives are inevitable with current SNV callers. These tools are typically optimized for haploid

references and single-platform datasets, often resulting in suboptimal performance on diploid

genomes and posing challenges for integrating calls across multiple sequencing platforms.

To address these issues, we developed an alignment-based quality assessment metric called

Sufficient Alignment Support (SAS), which directly identifies regions of the assembly lacking

adequate read support, an intrinsic indicator of assembly error. Building on this, we implemented a

"Structural Error-First" (SE-First) polishing strategy, which prioritizes the correction of structural

errors using ultra-long Oxford Nanopore Technology (ONT) reads to improve the alignment quality

for subsequent base-level corrections using shorter, high-accuracy reads. Starting from our

previously published T2T-YAO v1.1, an initial complete diploid genome assembly of a Han Chinese

individual[10], we incorporated additional sequencing data from high-accuracy platforms and

iteratively applied SAS evaluation and SE-First polishing. This approach yielded T2T-YAO v2.0, a

near-error-free diploid human genome, with the exception of residual issues in ribosomal DNA

(rDNA) clusters and a small number of homopolymer-associated errors in repetitive regions. Based

on Merqury analysis, T2T-YAO v2.0 contains no erroneous 21-mers absent from the raw sequencing

data, yielding an "infinite" quality value (QV) by k-mer standards, the first such result for a diploid

human genome.

Results

SAS: an alignment-based metric for assembly errors

To systematically evaluate the quality of genome assemblies, we developed a sensitive alignment-

based metric termed Sufficient Alignment Support (SAS). Rather than detecting inconsistencies

between the assembly and aligned sequencing reads, SAS identifies assembly errors by flagging

genomic windows that lack sufficient support from mapped reads. In SAS, base-level errors (BEs;

<50 bp) and structural errors (SEs; ≥50 bp) are quantified independently (**Fig. 1**).

For BEs, the diploid genome assembly is divided into 50 bp windows. Windows are flagged as

erroneous if they are not precisely supported by reads exceeding 10% of the overall mean

sequencing depth across any platform. This threshold is relaxed to 10% of the local sequencing

depth in specific low-coverage contexts: (1) regions within 10 kb of chromosomal ends (*i.e.*,

telomeres); (2) sequences containing degenerative GnA or CnT repeats (GA/CT content ≥90% and

GC content ≥60%); or (3) GC-rich regions (GC content ≥90%), which are known to be

underrepresented in polymerase-dependant, high-accuracy sequencing platforms (*e.g.*, Element,

Illumina, and PacBio platforms) and rely predominantly on ONT reads for coverage.

SEs are assessed in 2 kb non-overlapping windows using ultra-long ONT reads for its steady

coverage without obvious sequencing bias. A window is considered structurally erroneous if it meets

one of the following conditions: (1) sequencing depth is outside the 20% to 200% range of the global

average; or (2) fewer than 50% of local-depth reads span the window without large indels (≥50 bp).

This threshold is relaxed in chromosomal termini (within 10 kb), where ≥10% of the local-depth

reads must provide contiguous support. Additionally, to improve detection of misjoin events, 30 kb

sliding windows (stepped every 2 kb) are evaluated. If no reads span a given window, the

corresponding 15 consecutive 2 kb windows are flagged as SEs. Mitochondrial sequence, which is

small (~16 kb) and typically free of structural errors, are excluded from this analysis.

The overall SAS-based quality value (QV_{SAS}) is calculated by summing the BE and SE windows

and computing the predicted error rate (E) using the formula:

$$QV_{sas} = -10 \cdot \log E$$
$$E = \frac{BE + SE}{L}$$

Where E is the estimated error rate of the assessed assembly, BE and SE are the number of BE and

SE windows, respectively, L is the base length of the assembly.

To accurately evaluate the T2T-YAO assembly, we upgrated its raw data to high-accuracy

sequencing platforms: 260 Gb Q28 ultra-long ONT reads[11] (filtered for >30 kb; N50 >120 kb), ~172

Gb Pore-C[12], 306 Gb HiFi reads (PacBio SPRQ; filtered >3 kb), and 320 Gb Element AVITI[13] Q50

reads (**Table S1**). To reduce ambiguity in haplotype assignment during mapping, ONT and SPRQ

reads were phased in advance using haplotype-specific kmers (hapmers) and mapped in a haplotype-

118 aware manner.

120 We applied the SAS metric to evaluate the previously released T2T-YAO v1.1 assembly and

121 compared the results to those obtained using Merqury, a widely adopted k-mer–based quality

122 assessment tool (**Table 1**). SAS identified 21,511 base-level error (BE) windows and 4,438

123 structural error (SE) windows, yielding QV_{SAS} values of 53.12 and 54.13 for the maternal and

124 paternal haplotypes, respectively. Chromosome-specific QV_{SAS} scores ranged from 45.36 to 61.37

125 (**Table S2**). In contrast, Merqury using 21-mers reported only 22,556 erroneous k-mers,

126 corresponding to 1,074 estimated errors across the diploid genome. This number is notably higher

127 than initially reported[10], as the use of more accurate Element and SPRQ reads revealed assembly

128 errors that were previously obscured by sequencing errors. Nonetheless, the large discrepancy

129 between SAS and Merqury highlights the markedly greater sensitivity of SAS. Many BE windows

130 detected by SAS lacked corresponding erroneous k-mers, even when more sensitive 31-mers were

131 used, which identified 123,085 erroneous k-mers corresponding to 3,970 estimated errors (**Table 1**).

132 These results underscore the limitations of k-mer–based methods in capturing all types of assembly

133 errors. As a representative example, we showcased the paternal chromosome 1 to illustrate the

134 differences in error detection between SAS and other metrics (**Fig. 2a**).

136 Closer inspection of BE windows revealed a frequent association with homopolymer sequences,

137 where reads often disagreed on the number of repeated nucleotides. To classify these regions, we

138 defined homopolymer sites as stretches of ≥10 A/T, ≥7 G/C mononucleotide repeats, or ≥10

139 dinucleotide repeats. Accordingly, BEs were divided into homopolymer-associated (h-BEs) and

140 non-homopolymer-associated (nh-BEs), accounting for 17.17% (3,694) and 82.83% (17,817) of all

141 BEs, respectively (**Table 1**). While nh-BEs were relatively enriched in centromeric regions and

142 corresponded with k-mer–derived errors identified by Merqury, h-BEs were more broadly

143 distributed outside centromeres and were largely undetected by k-mer-based methods (**Fig.2**).

144 Conversely, some erroneous k-mers reported by Merqury were located outside SAS-flagged BE

145 windows and clustered within degenerative GnA or CnT repeat regions. These loci typically had

146 sufficient ONT coverage and were not flagged as errors by SAS, but lacked support from SPRQ and

147 Element reads due to platform-specific sequencing bias. This discrepancy underscores a

fundamental limitation of k-mer–based assessment tools: their inability to detect errors that do not

introduce novel k-mers, such as homopolymer length discrepancies, and their susceptibility to false

positives in regions with biased or zero coverage. The issue is particularly severe in the current

version of Merqury due to its reliance on the hybrid k-mer sets derived solely from high-accuracy

Illumina and PacBio platforms (Element in this study), excluding ONT data due to its relatively

higher sequencing error rate.

Given that Merqury cannot detect structural errors effectively, we compared SAS against several

structural variant (SV) callers: Sniffles2[14] (ONT), Flagger[15] (ONT), and nucflag[16] (SPRQ). Sniffles2

identified only 263 confident SV-like errors (supported by ≥60% of local reads), demonstrating

limited sensitivity, particularly in repetitive regions (**Extended Fig. 1**), where it relies on repeat

annotations often unavailable in *de novo* assemblies. Flagger and nucflag are widely inconsistent

with each other, that they identify 4.03 Mb and 12.30 Mb of collapsed regions (>50 bp), respectively,

with only 1.08 Mb of overlap (**Table S3**). Notably, 83.63% of their overlapping regions are either

within or adjacent (<10 kb) to SE windows identified by SAS. Furthermore, 78.50% of low-

coverage regions jointly flagged by Flagger ("erroneous" and "duplication" regions) and nucflag

("misjoin" regions) are also detectable by SAS. We also observed that nucflag frequently

misclassifies misjoins in GnA/CnT repeats and other regions with zero SPRQ coverage due to

sequencing bias (**Fig. 2**). The obvious bias of SPRQ in specific sequences damages the reliability

of SPRQ-based SE callers. Haplotypic switch errors, which are flagged by nucflag as heterogeneous

regions and by Merqury as switch k-mers, are not treated as a separate error class in SAS. Depending

on the read evidence, such events are classified as SEs if flanking clips are present or as BEs if

phased read support is available but without a clear breakpoint.

Correction of Structural Errors

Using SAS and ultra-long ONT reads, we identified 4,438 non-redundant SE windows in the T2T-

YAO v1.1 assembly. These windows were further categorized based on their underlying features:

1,295 were high-depth regions (29.18%), 1,662 low-depth regions (37.45%), 331 contained indels

(7.46%), 61 were marked by clipped alignments (1.37%), and 1,929 were consecutive windows

without spanning reads (43.46%). Note that individual windows could be assigned to multiple

categories. Errors related to clipped alignments and abnormal depth were particularly concentrated

in the short arms of acrocentric chromosomes and the centromeres of maternal chromosomes 6 and

7, whereas indels were more evenly distributed across the genome (**Fig. 3a**).

Given the critical importance of accurate read alignment for subsequent base-level error correction,

we adopt a "Structural Error-first" (SE-first) strategy. This approach prioritizes correction of

structural errors using ONT reads before applying base-level polishing with high-accuracy

sequencing data. By resolving large-scale misassemblies upfront, we reduce the risk of

overcorrection due to artificial alignments in shorter reads, thereby enhancing the overall fidelity of

the assembly.

We employed a comprehensive correction pipeline combining variant calls from Sniffles2, local

reassemblies using ONT reads, and targeted telomere extension to eliminate all detectable SEs. After

each correction round, reads were re-aligned, and SAS was reapplied to reassess error content

(**Extended Fig. 2**). Most SEs were resolved after several iterations of this correction–realignment–

evaluation loop, though certain clipped alignment sites required manual inspection and correction.

In telomeric and subtelomeric regions, ONT reads often exhibited heterogeneity that impeded

successful reassembly. In such cases, stratifying ONT reads based on characteristic SNVs and

assembling the dominant haplotype group enabled resolution. Following these polishing efforts, all

telomeres were extended to an average length of 8.7 ± 2.6 kb, with all exceeding 5 kb except for

three in the q arms of maternal chromosome 12 and paternal chromosomes 21 and 22 (**Fig. 4a**).

Notably, most telomeres contained some non-canonical tandem 6-mer repeats similar to

(TTAGGG)n[17], often interspersed with canonical motifs, particularly at the transition from

subtelomeric to telomeric regions, as previously described[18]. The structural integrity of the assembly

is further validated by the Pore-C chromosomal configuration data, with its contact map confirming

the haplotype phasing and the accurate pairing of short and long arms across the ten acrocentric

chromosomes, a particularly challenging structural feature to resolve (**Extended Fig. 3**).

Ultimately, T2T-YAO v2.0 becomes free of detectable SEs outside of ribosomal DNA (rDNA)

clusters (**Fig. 2b, 3b**). Centromeric collapses, particularly in maternal chromosomes 6 and 7, are

also resolved. Their corresponding higher-order repeat (HOR) arrays in T2T-YAO v2.0 are

successfully extended to near-native lengths (**Fig. 4b, Extended Fig. 4**). One potential exception of

high-depth is observed in a 12 kb region upstream of the *TSPAN33* gene on maternal chromosome

7q32. Although the local ONT depth (84–89×) marginally exceedes the global cutoff (84×), the

sequence there is fully supported by all platforms without signs of collapse or misjoin. Further

inspection reveals that this region is composed of LIM and Alu elements which sequences are

widely present in human genome with millions of copies[19]. The repetitive nature of this region may

be the source of the elevated depth where 22.8% of mapped reads are shorter than 60 kb and

exhibiting MAPQ scores of 0–1 (**Extended Fig. 5**).

Aside from this, ONT read depth followed a Gaussian distribution, with extremely few regions

tallied by Flagger as false duplication or collapsed (**Extended Fig. 6**). All overlapping issue regions

identified by Flagger and nucflag are confined to rDNA arrays or telomeric regions, supporting the

structural completeness of T2T-YAO v2.0 (**Table S3**). To further validate continuity, we assessed

the fraction of the assembly covered by non-split ONT reads (NScov%). In T2T-YAO v1.1, NScov%

is 99.98%, with ~0.91 Mb (excluding rDNA) left uncovered. In contrast, T2T-YAO v2.0

accomplishes 100% NScov when rDNA is masked, confirming the absence of unresolved SEs.

In T2T-YAO v1.1, alignment of SPRQ and Element reads revealed widespread artificial collapses,

totaling 4.71 Mb and 2.52 Mb, respectively, in regions with >2-fold mean depth (**Fig. 2a, Extended**

Fig. 7a). After correction, these figures were reduced to 3.43 Mb (SPRQ) and 0.93 Mb (Element),

with 22.99% and 35.37% of these remaining regions located within rDNA (**Fig. 2b, Extended Fig.**

7b). Reduced artifact-induced alignments in high-accuracy data minimized the risk of over-

correction of errors and improved reliability in the polished assembly.

Correction of Base-level Errors

Using ONT, SPRQ, and Element sequencing data, we identified 21,511 base-level error (BE)

windows in T2T-YAO v1.1, comprising 3,694 homopolymer-associated BEs (h-BEs) and 17,817

non-homopolymer-associated BEs (nh-BEs), broadly distributed across all chromosomes (**Fig. 3b**).

To avoid overcorrection, each flagged window is replaced with a consensus sequence derived from

reads spanning that window. In cases where the three sequencing platforms disagree, we prioritize

corrections supported by at least two platforms or follow the order of reliability: Element > SPRQ >

ONT. If no consensus is obtainable, ONT reads spanning the region are locally reassembled to

correct the sequence, provided the error is not homopolymer-related (**Extended Fig. 8**). For h-BEs,

local ONT-based consensus or assembly is omitted due to the known limitations of ONT sequencing

in accurately resolving homopolymers (**Extended Fig. 9**). This platform-aware consensus strategy

minimized alignment artifacts and reduced overcorrection during subsequent polishing rounds.

Read alignment and polishing are repeated iteratively until the number of BEs is stabilized.

To correct additional base-level errors, we applied multiple variant callers: Deeppolisher[20] (SPRQ),

DeepVariant[21] (ONT and Element), and GATK HaplotypeCaller[22] (Element) to call variants in the

diploid genome. These tools yielded a total of non-redundant 63,398 homozygous variants with

genotype quality (GQ) ≥20. To eliminate false positives, we filtered out variants which original

sequences as well as 10 bp flanking sequences were supported by ≥60% local depth and ≥10 exact-

matching reads. This filtering step yielded 43,444 potential SNV-like errors across both haplotypes,

of which 40.21% were associated with homopolymer regions, an even higher proportion than the

observed rate of h-BEs (**Extended Fig. 10a**). The elevated frequency of homopolymer-associated

errors likely stems from the homopolymer compression step used in the initial T2T-YAO assembly

with Verkko[23,24]. Of these SNV-like errors, 43.90% of non-homopolymer-associated variants

overlapped with SAS-detected nh-BEs, whereas only 7.08% of homopolymer-related calls

overlapped with SAS h-BEs. This discrepancy highlights the challenge of identifying true

homopolymer-associated errors via alignment. All variant callers detect discrepancies between the

assembly and aligned reads; however, many such discrepancies are false positives arising from read

misalignment or homopolymer inaccuracy. Accordingly, we treated SNV-like variants with the same

correction strategy as BEs: replacing both the variant site and its 10 bp flanking sequence with the

read-derived consensus sequence, rather than using the sequence predicted by the variant caller.

After several rounds of iterative polishing, T2T-YAO v2.0 containes only 336 nh-BEs (all within

rDNA regions) and 837 h-BEs located in repetitive regions (including 489 in rDNA) that remain

unresolved due to the inherent limitations of current sequencing technologies (**Table 1, Fig. 3b**).

When we reassess the v2.0 assembly using the same SNV callers, the total number of filtered SNV-

like potential errors (homozygous SNV with GQ ≥20) drops significantly to 2,354 (**Extended Fig.**

10b). Of these, only 38 are non-homopolymer-associated, with 36 (94.74%) located within rDNA

units, consistent with the distribution of nh-BEs identified by SAS. The remaining two non-

homopolymer calls outside rDNA are confirmed to be false positives.

An illustrative example of an unresolved h-BE can be found in the D4Z4 repeat array on

chromosome 4q35[25]. In this region, the polyC tracts within DUX4-like genes can not be confidently

resolved: ONT and SPRQ reads struggled with homopolymer sequencing, while Element reads

suffered from mapping ambiguity (**Extended Fig. 11**). Nonetheless, the remaining h-BEs account

for only 0.04% of the total 2,412,254 homopolymer sites genome-wide. Among these, polyA/polyT

tracts (≥10 bp) comprised 81.70%, while polyC/polyG tracts (≥7 bp) and poly-dinucleotide repeats

(≥10 bp) represented 9.20% and 9.10%, respectively. The distribution of these homopolymer types

is avoid of centromere regions and consistent across haplotypes (**Extended Fig. 12**).

Toward an Error-Free Human Genome Assembly

We re-evaluated the quality of the T2T-YAO v2.0 assembly using both SAS and Merqury metrics,

based on the same sequencing datasets (**Table 1**). The final SAS assessment identified 1,773

structural error (SE) windows (all confined to rDNA regions except for the six windows of high-

depth in maternal chromosome 7) and 1,175 base-level error (BE) windows, comprising 827 within

rDNA and 348 homopolymer-associated BEs (h-BEs) outside rDNA. The resulting QV_{SAS} scores

were 66.00 (maternal) and 61.25 (paternal). When rDNA regions were masked, leaving only the 348

h-BEs, the QV_{SAS} values rose to 71.62 and 71.81, respectively. (**Table S2, Supplementary Figs. 1–**

45).

T2T-YAO v2.0 contains tens of kilobases of GnA/CnT repeat sequences that are poorly supported

by Element and SPRQ reads but well supported by ONT data. Due to the limited representation of

these sequences in the k-mer set used by Merqury, a high number of false positives are called in

these loci. Specifically, Merqury identified 3,006 erroneous 21-mers (corresponding to a QV of

76.18) and 28,956 erroneous 31-mers (QV 68.03) in the diploid assembly. Since all erroneous 21-

mers are encompassed by the erroneous 31-mers, we focused our inspection on the 31-mers. These

were consistently found in regions with near-zero coverage in Element and SPRQ data but showed

strong support from ONT reads. The 31-mers clustered into 1,127 discrete regions (after merging

overlaps), collectively spanning 76,248 bp. Of these, 674 regions (totaling 53,307 bp) had extremely

high GA/CT content (>90%), while the rest were composed of non-canonical 6-mers found in

telomeric sequences or GC-rich tandem repeats, patterns consistent with known sequencing

biases[26,27] (**Table S4**).

The ONT data used in this study were generated using the R10 flow cell, known for high base-

calling accuracy (Q28; >99.8%). To leverage this accuracy, we incorporated k-mers from the ONT

dataset into our analysis, applying a stringent filter to exclude potential sequencing errors, setting

the minimum occurrence threshold at ≤4 for ONT data, compared to ≤1 for Element and SPRQ

reads. We first use this expanded trinary k-mer set to assess T2T-YAO v1.1, which QV only shows

a slight improvement over the binary set, indicating minimal inclusion of ONT sequencing errors.

When we applied the trinary k-mer set to assess T2T-YAO v2.0, the QV for 21-mers increased to

"infinite," representing the first Merqury-defined error-free diploid genome. The QV for 31-mers

also improved significantly, reaching 96.29 for the maternal haplotype and 88.73 for the paternal,

with only 143 erroneous 31-mers remaining. All paternal errors are confined to rDNA regions, while

the maternal errors are located in two h-BE loci of GnA repeat regions containing several polyG

tracts (≥8 bases), corresponding precisely with the residual errors detected by the SAS metric.

The comprehensive correction of SEs and non-homopolymer BEs outside rDNA enabled

unprecedented resolution of human centromeres, regions that remain among the most challenging

to assemble aside from rDNA clusters[28]. We refined the annotation of centromeres, higher-order

repeats (HORs), and other satellite elements across all 46 chromosomes (**Fig. 4b**). Subtle but

significant differences between T2T-YAO v1.1 and v2.0 were observed, exemplified by updates to

the paternal chromosome 1 assembly, highlighting the importance of high-fidelity polishing for

accurate centromere characterization (**Extended Fig. 13**). Using the recently developed Genomic

Centromere Profiling (GCP) pipeline, a method analogous to fragment length analysis for

characterizing the position, orientation, and organization of centromere protein B (CENP-B)

binding motifs[29], we constructed structural models for centromeric repeat clusters in T2T-YAO.

These models revealed that the sequence architecture of human centromeres varies substantially in

size, structure, and composition among chromosomes[30]. Although maternal and paternal

centromeres of the same chromosome exhibit similar patter, differences in the size of components

are still obvious between haplotypes, such as the HOR of maternal chromosomes 6 and 7 were

significantly expanded relative to their paternal counterparts in T2T-YAO (**Extended Fig. 4**).

Together, these advances, achieved through iterative structural and base-level polishing,

demonstrate the power of the SAS framework and our "Structural Error-first" strategy. The resulting

T2T-YAO v2.0 assembly represents one of the most accurate and complete diploid human genomes

to date, nearing the status of an error-free reference genome.

Discussion

An error-free diploid human genome holds immense value for calibrating variant calling pipelines

and enhancing the accuracy of clinical genetic diagnoses[5,31]. Following the completion of the T2T-

CHM13 assembly, the "Q100" project was initiated with the ambitious goal of producing a diploid

human genome of Phred quality score 100, equivalent to one error per 10 billion bases[32]. Given the

approximate size of the human diploid genome (~6 billion bases), achieving Q100 implies

constructing a truly error-free genome. While T2T-YAO v2.0 has been validated by non-split ONT

reads and is free of erroneous 21-mers, thus qualifying for an "infinite" QV under Merqury standards,

technical limitations in both genome evaluation and error correction prevent it from being deemed

absolutely error-free.

Genome assembly quality encompasses multiple dimensions, including completeness, correctness,

and continuity. Of these, completeness is relatively straightforward to assess via k-mer-based

approaches[9]. However, accurate evaluation of correctness and continuity remains more challenging

and often requires alignment-based methods[33,34]. Most alignment-based tools identify discrepancies

between the genome assembly and realigned reads. These discrepancies often include both false

positives, arising from artificial alignments of heterogeneous reads, and false negatives, which are

missed due to sequencing bias. Moreover, integrating alignment discrepancies across platforms

remains difficult.

To overcome these challenges, we developed SAS, a window-based alignment metric that flags 50

bp segments not sufficiently supported by reads from any platform. In our implementation, a 50-

mer is considered supported only if it is exactly supported by at least five reads (representing >10%

of the mean sequencing depth) in its context. This approach offers higher stringency and sensitivity

than k-mer-based methods such as Merqury, which typically examine 21- or 31-mers and require

only two occurrences. Similarly, structural errors are detected in windows lacking sufficient

spanning reads. The window-based design also improves computational efficiency: by initially

filtering for regions perfectly covered by Element reads without mismatches or clips, which

constitute 99.2% of the T2T-YAO genome, the overall computational burden is greatly reduced.

This enables SAS to complete genome-wide assessment in under two hours after alignment.

Despite its sensitivity, SAS still misses some base-level errors in repetitive regions where

misaligned reads from other loci create false support. It also fails to detect collapses or false

duplications that fall within acceptable depth thresholds. These problematic regions often show

heterogeneous SNP patterns in the aligned reads[16], complicating distinction from sequencing noise,

particularly with ONT reads. A canonical example is the ribosomal DNA (rDNA) clusters, which

remain the final hurdle to a fully error-free human genome[35]. In these regions, ONT reads are long

enough to span complete rDNA units but lack the base-level accuracy to resolve unit-level SNVs,

whereas SPRQ reads provide the needed accuracy but are not long enough to span the entire unit.

Thus, the primary barriers to achieving a completely error-free diploid human genome are intrinsic

to the architecture of highly repetitive sequences, particularly their unit length and sequence identity,

and the current limitations of sequencing technologies in generating reads that are both sufficiently

long and accurate. Bridging this gap remains the final frontier in complete human genome assembly.

Methods

iPSC Culture and DNA Sequencing

Induced pluripotent stem cells (iPSCs) used for the T2T-YAO assembly were cultured in mTeSR™ Plus medium on vitronectin-coated plates and passaged at a 1:10 ratio using 0.02% EDTA (Versene). High-molecular-weight genomic DNA for Oxford Nanopore Technologies (ONT) ultra-long read sequencing was extracted using the Grandomics BAC-long kit. DNA fragments >50 kb were enriched using SageHLS, and sequencing libraries were prepared with the SQK-LSK114 kit. Sequencing was performed on a PromethION platform with R10 flow cells. Basecalling was conducted using Dorado (v5.0.0, sup mode), and reads with quality scores <Q7 were excluded.

For PacBio Revio® sequencing, DNA was extracted via CTAB and purified using Grandomics kits. SMRTbell® libraries were prepared with the SMRTbell Prep Kit 3.0 and analyzed using SMRT Link v25.1. For Element AVITI sequencing, PCR-free libraries were generated using the Elevate Enzymatic Library Prep Kit (Cat. No. 830-00009) and Long UDI Adapter Kit Set A (Cat. No. 830-00010), following the manufacturer's protocols. Libraries were pooled, denatured, and sequenced using the AVITI 2×150 Sequencing Kit CB UltraQ (Cat. No. 860-00018) on the AVITI platform.

Detection of Structural Error (SE) Windows

ONT ultra-long reads were filtered to retain only those >30 kb and phased using haplotype-specific 31-mers (hapmers). A read was assigned to a haplotype if it contained >40 hapmers from one haplotype and at least three-fold more than from the opposite haplotype. Phased reads were aligned to the corresponding haplotype assemblies using Winnowmap2[36]; unassigned reads were aligned to the diploid reference. Only primary alignments were retained.

The assembly was scanned using non-overlapping 2 kb windows. A window was flagged as an SE if :

1. **INDELs and Clipping**: >40% of the local reads contained insertions or deletions ≥50 bp detected via CIGAR string analysis, or Presence of hard or soft clipping in aligned reads.

2. **Abnormal Depth**: Read depth <20% or >200% of the genome-wide mean (assessed using Mosdepth).

3. **Lack of Spanning Reads**: No spanning reads were observed within 15 consecutive 2 kb windows (*i.e.*, a 30 kb region).

417   Thresholds were relaxed near chromosome ends (<10 kb from termini), where a window was

418   flagged only if no or fewer than 10% of local reads spanned the region and lacked indels.

Detection of Base-Level Error (BE) Windows

Element reads were aligned to the diploid assembly using BWA-MEM. Alignments were filtered to

retain only perfect matches (CIGAR NM:i:0) without clipping. SPRQ reads were phased and

aligned similarly to ONT reads.

Base-level errors were evaluated in sliding 50 bp windows. Windows were flagged as BEs if the

exact sequence was supported by reads at <10% of the mean sequencing depth. Windows with

sufficient (>10% mean depth) perfect-match Element reads were excluded. Remaining regions were

further validated using SPRQ and ONT data. In low-coverage regions, thresholds were relaxed to

10% of **local** depth under the following conditions:

1.   Within 10 kb of chromosomal termini (*i.e.*, telomeric regions);

2.   Regions with degenerative GA/CT repeats (≥90% GA/CT and ≥60% GC content);

3.   GC-rich regions (≥90% GC content).

Merqury-Based Assembly Evaluation

435   Merqury and Meryl were used to assess k-mer-based quality metrics (QV), completeness, and

436   phasing accuracy using 21-mers and 31-mers. The evaluation followed the methods described by

437   McCartney et al[7] for the binary k-mer set using Element and SPRQ data, and for the trinary k-mer

438   set except combining the k-mer set of ONT reads of occurrence >4.

Variant Calling (SVs and SNVs)

441   Structural variants-like errors were called using Sniffles2 (ONT), Flagger (ONT), and NucFlag

442   (SPRQ), with default parameters. Base-level SNV-like errors were identified using DeepVariant

443   (ONT and Element), Deeppolisher (SPRQ), and GATK HaplotypeCaller (Element). Variants with

444   genotype quality (GQ) <20 were excluded. Heterozygous calls were also filtered out except for

445   haplotype-aware calls from Deeppolisher.

Error Correction

SE windows were first corrected using variant calls from Sniffles2 when available. In cases without confident calls, reads spanning the SE and 5–10 kb flanks were reassembled using Hifiasm or Flye. If required, flanking regions were extended to 50–100 kb for resolving large collapses or duplications. Reassembled contigs were aligned back to the reference, and the corrected sequences were substituted accordingly. Manual assessment and correction with IGV is often necessary for SEs of clipping or complex errors. Telomeric regions with misalignment or heterogeneity were extended using major-allele ONT reads. Corrections were followed by haplotype-aware re-alignment and re-evaluation with SAS. The process was repeated until no SEs were detected outside rDNA clusters.

BE windows were classified into homopolymer-associated (h-BE) and non-homopolymer-associated (nh-BE).

nh-BEs were corrected using consensus sequences from supporting reads in the order of platform priority: Element > SPRQ > ONT. Where consensus could not be reached, local ONT reassembly was performed.

h-BEs were polished using Element or SPRQ consensus only, due to ONT's known limitations in homopolymer resolution.

For SNV-like errors called by DeepVariant, Deeppolisher, and GATK HaplotypeCaller, false positives were removed if >55% and ≥10 reads from any platform exactly supported the reference allele sequence including ±10 bp flanks. Remaining SNV calls were corrected using consensus-based replacement as nh-BEs and h-BEs except within the ±10 bp flanks windows. Polishing iterations continued until the number of detected BEs plateaued.

Centromere and Telomere Annotation

Centromeric regions were annotated using RepeatMasker[37], with α-satellite arrays identified via HumAS-HMMER[30]. Telomeric repeats were detected and quantified using Telescope[38].

References

1   Church, D. M. A next-generation human genome sequence. *Science (New York, N.Y.)* **376**,
34-35 (2022). https://doi.org:10.1126/science.abo5367

2   Nurk, S. *et al.* The complete sequence of a human genome. *Science (New York, N.Y.)* **376**,
44-53 (2022). https://doi.org:10.1126/science.abj6987

3   Olson, N. D. *et al.* Variant calling and benchmarking in an era of complete human genome
sequences. *Nature Reviews Genetics* **24**, 464-483 (2023). https://doi.org:10.1038/s41576-
023-00590-0

4   Aganezov, S. *et al.* A complete reference genome improves analysis of human genetic
variation. *Science (New York, N.Y.)* **376**, eabl3533 (2022).
https://doi.org:doi:10.1126/science.abl3533

5   Jarvis, E. D. *et al.* Semi-automated assembly of high-quality diploid human reference
genomes. *Nature* **611**, 519-531 (2022). https://doi.org:10.1038/s41586-022-05325-5

6   An assembly line for an improved human reference genome. *Nature* (2022).
https://doi.org:10.1038/d41586-022-03151-3

7   Mc Cartney, A. M. *et al.* Chasing perfection: validation and polishing strategies for
telomere-to-telomere genome assemblies. *Nature methods* **19**, 687-695 (2022).
https://doi.org:10.1038/s41592-022-01440-3

8   Rhie, A. *et al.* Towards complete and error-free genome assemblies of all vertebrate
species. *Nature* **592**, 737-746 (2021). https://doi.org:10.1038/s41586-021-03451-0

9   Rhie, A., Walenz, B. P., Koren, S. & Phillippy, A. M. Merqury: reference-free quality,
completeness, and phasing assessment for genome assemblies. *Genome biology* **21**, 245
(2020). https://doi.org:10.1186/s13059-020-02134-9

10   He, Y. *et al.* T2T-YAO: A Telomere-to-telomere Assembled Diploid Reference Genome
for Han Chinese. *Genomics, proteomics & bioinformatics* **21**, 1085-1100 (2023).
https://doi.org:10.1016/j.gpb.2023.08.001

11   Ni, Y., Liu, X., Simeneh, Z. M., Yang, M. & Li, R. Benchmarking of Nanopore R10.4 and
R9.4.1 flow cells in single-cell whole-genome amplification and whole-genome shotgun
sequencing. *Comput Struct Biotechnol J* **21**, 2352-2364 (2023).
https://doi.org:10.1016/j.csbj.2023.03.038

12   Deshpande, A. S. *et al.* Identifying synergistic high-order 3D chromatin conformations
from genome-scale nanopore concatemer sequencing. *Nature biotechnology* **40**, 1488-
1499 (2022). https://doi.org:10.1038/s41587-022-01289-z

13   Arslan, S. *et al.* Sequencing by avidity enables high accuracy with low reagent
consumption. *Nature biotechnology* **42**, 132-138 (2024). https://doi.org:10.1038/s41587-
023-01750-7

14   Smolka, M. *et al.* Detection of mosaic and population-level structural variants with
Sniffles2. *Nature biotechnology* **42**, 1571-1580 (2024). https://doi.org:10.1038/s41587-
023-02024-y

15   Liao, W. W. *et al.* A draft human pangenome reference. *Nature* **617**, 312-324 (2023).
https://doi.org:10.1038/s41586-023-05896-x

16   Vollger, M. R. *et al.* Long-read sequence and assembly of segmental duplications. *Nature*
methods **16**, 88-94 (2019). https://doi.org:10.1038/s41592-018-0236-3

17   Lyčka, M. *et al.* TeloBase: a community-curated database of telomere sequences across
the tree of life. *Nucleic acids research* **52**, D311-D321 (2023).

https://doi.org:10.1093/nar/gkad672

18   Stephens, Z. & Kocher, J.-P. Characterization of telomere variant repeats using long reads
enables allele-specific telomere length estimation. *BMC Bioinformatics* **25**, 194 (2024).
https://doi.org:10.1186/s12859-024-05807-5

19   Belancio, V. P., Deininger, P. L. & Roy-Engel, A. M. LINE dancing in the human genome:
transposable elements and disease. *Genome Medicine* **1**, 97 (2009).
https://doi.org:10.1186/gm97

20   Mastoras, M. *et al.* Highly accurate assembly polishing with DeepPolisher. *bioRxiv* (2024).
https://doi.org:10.1101/2024.09.17.613505

21   Poplin, R. *et al.* A universal SNP and small-indel variant caller using deep neural networks.
Nature biotechnology **36**, 983-987 (2018). https://doi.org:10.1038/nbt.4235

22   Poplin, R. *et al.* Scaling accurate genetic variant discovery to tens of thousands of samples.
bioRxiv, 201178 (2018). https://doi.org:10.1101/201178

23   Rautiainen, M. *et al.* Telomere-to-telomere assembly of diploid chromosomes with
Verkko. *Nature biotechnology* (2023). https://doi.org:10.1038/s41587-023-01662-6

24   Antipov, D. *et al.* Verkko2 integrates proximity-ligation data with long-read De Bruijn
graphs for efficient telomere-to-telomere genome assembly, phasing, and scaffolding.
Genome research **35**, 1583-1594 (2025). https://doi.org:10.1101/gr.280383.124

25   Lemmers, R. *et al.* Cis D4Z4 repeat duplications associated with facioscapulohumeral
muscular dystrophy type 2. *Hum Mol Genet* **27**, 3488-3497 (2018).
https://doi.org:10.1093/hmg/ddy236

26   Ross, M. G. *et al.* Characterizing and measuring bias in sequence data. *Genome biology*
14, R51 (2013). https://doi.org:10.1186/gb-2013-14-5-r51

27   Freeman, T. M., Wang, D. & Harris, J. Genomic loci susceptible to systematic sequencing
bias in clinical whole genomes. *Genome research* **30**, 415-426 (2020).
https://doi.org:10.1101/gr.255349.119

28   Altemose, N. *et al.* Complete genomic and epigenetic maps of human centromeres.
Science (New York, N.Y.) **376**, eabl4178 (2022).
https://doi.org:doi:10.1126/science.abl4178

29   Corda, L. & Giunta, S. Chromosome-specific centromeric patterns define the centeny map
of the human genome. *Science (New York, N.Y.)* **389**, eads3484 (2025).
https://doi.org:10.1126/science.ads3484

30   Logsdon, G. A. *et al.* The variation and evolution of complete human centromeres. *Nature*
629, 136-145 (2024). https://doi.org:10.1038/s41586-024-07278-3

31   Wagner, J. *et al.* Curated variation benchmarks for challenging medically relevant
autosomal genes. *Nature biotechnology* **40**, 672-680 (2022).
https://doi.org:10.1038/s41587-021-01158-1

32   consortium, T.-t.-t. HG002 Q100 project. *https://github.com/marbl/hg002* (2023).

33   Li, K., Xu, P., Wang, J., Yi, X. & Jiao, Y. Identification of errors in draft genome assemblies
at single-nucleotide resolution for quality assessment and improvement. *Nature*
communications **14**, 6556 (2023). https://doi.org:10.1038/s41467-023-42336-w

34   Mastoras, M. *et al.* Highly accurate assembly polishing with DeepPolisher. *bioRxiv*,
2024.2009.2017.613505 (2024). https://doi.org:10.1101/2024.09.17.613505

35   Li, H. & Durbin, R. Genome assembly in the telomere-to-telomere era. *Nat Rev Genet* **25**,

658-670 (2024). https://doi.org/10.1038/s41576-024-00718-w

36   Jain, C., Rhie, A., Hansen, N. F., Koren, S. & Phillippy, A. M. Long-read mapping to repetitive reference sequences using Winnowmap2. *Nature methods* **19**, 705-710 (2022). https://doi.org/10.1038/s41592-022-01457-8

37   Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in genomic sequences. *Curr Protoc Bioinformatics* **Chapter 4**, 4.10.11-14.10.14 (2009). https://doi.org/10.1002/0471250953.bi0410s25

38   Formenti, G. *et al.* Gfastats: conversion, evaluation and manipulation of genome sequences using assembly graphs. *Bioinformatics (Oxford, England)* **38**, 4214-4216 (2022). https://doi.org/10.1093/bioinformatics/btac460

Data Availability

The raw sequencing data for T2T-YAO v2.0 generated in this study have been deposited in the GSA for Human database at the China National Center for Bioinformation (CNCB) under accession number HRA011075, publicly accessible at https://ngdc.cncb.ac.cn/gsa-human. The raw data for v1.1, released on August 16, 2023, are available from the same URL under accession number HRA004987. The T2T-YAO v1.1 and v2.0 genome assemblies are available in the Genome Warehouse at CNCB (https://ngdc.cncb.ac.cn/gwh) under the following accession numbers: v1.1: GWHDQZJ00000000 (maternal), GWHDOOG00000000 (paternal); v2.0: GWHGEYC00000000.1 (maternal), GWHGEYB00000000.1 (paternal). All genome sequences and associated datasets can be freely downloaded without application and are also publicly accessible at https://www.genomeyao.org.cn/v2.

Code Availability

Code for SAS, error correction, and all other workflows are available at https://github.com/KANGYUlab/sas-pipeline and https://github.com/KANGYUlab/genome-polish-pipeline.

Code for other software used in this work:

Meryl v1.3 (https://github.com/marbl/meryl) and Merqury v1.1 (https://github.com/marbl/merqury) for k-mer QV assessment. ONT and SPRQ reads were aligned to assembly using Winnowmap v2.03 (https://github.com/marbl/Winnowmap). Aligment between assemblies was using minimap2 2.27 (https://github.com/lh3/minimap2). Element reads were aligned with BWA-MEM 0.7.17

(https://github.com/lh3/bwa). SAMtools v1.19.2 (https://github.com/samtools/samtools/) and

bedtools v2.25.0 (https://github.com/arq5x/bedtools2) were used for BAM files and BED files

processing.   Reads   alignment   were   visualized   with   IGV   v2.17.3

(https://github.com/igvteam/igv.js#igvjs). Genome-wide alignment depth was calculated using

mosdepth (https://github.com/brentp/mosdepth). The SVs were detected using NucFlag v0.3.6

(https://github.com/logsdon-lab/NucFlag) from the phased alignmen of SPRQ, Flagger v0.3.2

(https://github.com/mobinasri/flagger/), Sniffles2 v2.4 (https://github.com/fritzsedlazeck/Sniffles)

from the phased alignment of ONT. The SNVs were detected by GATK v4.3.0.0

(https://github.com/broadinstitute/gatk),   Deepvariant   v1.9.0

(https://github.com/google/deepvariant),   Deepploisher   r0.1.0

(https://github.com/google/deeppolisher). Local reassemblies were performed with Flye v2.9.5

(https://github.com/mikolmogorov/Flye)   and   Hifiasm   v0.25.0

(https://github.com/chhylp123/hifiasm). Bcftools v1.20 (https://github.com/samtools/bcftools/)

was utilized for processing VCF files and incorporating polishing edits into the new assembly.

Centromere   regions   were   identified   using   RepeatMasker   v4.1.9

(https://www.repeatmasker.org/RepeatMasker), and the location of the α-satellite HOR arrays were

identified using HumAS-HMMER (https://github.com/enigene/HumAS-HMMER). The telomere

regions were identified using Telescope v0.1.1 (https://github.com/vgl-hub/teloscope). Haplotype

phasing   were   validated   with   Pore-C   contact   maps   using   wf-pore-C   v1.3.0

(https://github.com/epi2me-labs/wf-pore-c)   and   visualized   in   Juicebox   v2.20

(https://github.com/aidenlab/Juicebox). The organization patterns of centromeres and telomeres

were visualized using ggplot2 (v3.5.1) in R (v4.4.0). Statistical visualizations were generated using

Matplotlib (v3.7.1) with NumPy (v1.24.3) and Pandas (v2.0.3) in Python (v3.9.16).

Acknowledgments

This study was supported by the grants of the National Key Research and Development Program of

China (2024YFC3405701), the National Science Foundation of China (32371537).

Author contributions

Y.K. conceived and designed the project. Z.H. implemented the software SAS. Y.C. conducted the

correction of T2T-YAO. S.G., C.S., J.W. performed sample preparation and sequencing. X.Y., Y.T.,

J.C., R.L. and Y.H. participated in data analyses, and Y.K., Z.G., J.H., J.Y. wrote the manuscript. All

the authors agree on the manuscript.

Competing interest declaration

All authors declare no competing interests.

Table

Table 1. Comparative Quality Metrics of the T2T-YAO Assembly Assessed by SAS and Merqury.

		v1.1			v2.0		
		mat	pat	Both	mat	pat	Both
SAS	**QV (total)**	**53.12**	**54.13**	**53.59**	**66.00**	**61.25**	**63.04**
	BE	**11946**	**9565**	**21511**	**291**	**884**	**1175**
	h-BE	1789	1905	3694	255	582	837
	nh-BE	10157	7660	17817	35	301	336
	SE	**2752**	**1686**	**4438**	**469**	**1304**	**1773**
	low_depth	920	742	1662	121	709	830
	high_depth	886	409	1295	302	376	678
	indel	141	190	331	9	49	58
	clip	39	22	61	6	24	30
	no spanning	1247	682	1929	99	469	568
	QV (rDNA masked)	**53.65**	**55.39**	**54.42**	**71.62**	**71.81**	**71.72**
	BE	**10871**	**7863**	**18734**	**179**	**169**	**348**
	h-BE	1426	1145	2571	179	169	348
	nh-BE	9445	6718	16163	0	0	0
	SE	**2147**	**572**	**2719**	**6**	**0**	**6**
	low_depth	730	57	787	0	0	0
	high_depth	643	115	758	6	0	6
	indel	116	125	241	0	0	0
	clip	23	8	31	0	0	0
	no spanning	1027	295	1322	0	0	0
Merqury-binary	**QV-21mer**	**66.48**	**68.68**	**67.42**	**74.84**	**78.23**	**76.18**
	err-kmer	14252	8304	22556	2085	921	3006
	total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
	err rate	2.25×10^7	1.36×10^7	1.81×10^7	3.28×10^8	1.50×10^8	2.41×10^8
	switch err%	0.0107	0.0254	—	0.0017	0.0038	—
	Completeness%	97.770	93.708	99.839	97.777	93.710	99.850
	QV-31mer	**60.97**	**62.73**	**61.74**	**67.24**	**69.04**	**68.03**
	err-kmer	74864	48221	123085	17694	11262	28956
	total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
	err rate	8.00×10^7	5.34×10^7	6.69×10^7	1.89×10^7	1.25×10^7	1.57×10^7
	switch err%	0.0219	0.0413	—	0.0050	0.0072	—
	Completeness%	96.780	92.607	99.806	96.792	92.610	99.826
Merqury-trinary	**QV-21mer**	**67.7888**	**69.854**	**68.6823**	**inf**	**inf**	**inf**
	err-kmer	10543	6331	16874	0	0	0
	total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
	err rate	1.66×10^7	1.03×10^7	1.35×10^7	0	0	0
	switch err%	0.0107	0.0254	—	0.0017	0.0038	—
	Completeness%	97.7896	93.7264	99.8638	97.7966	93.7281	99.875
	QV-31mer	**62.4363**	**64.2025**	**63.2153**	**96.2941**	**88.7314**	**91.0965**

err-kmer	53376	34337	87713	22	121	143
total-kmer	3.02×10^9	2.92×10^9	5.93×10^9	3.02×10^9	2.92×10^9	5.93×10^9
err rate	5.71×10^7	3.80×10^7	4.77×10^7	2.35×10^7	1.34×10^7	7.77×10^7
switch err%	0.0219	0.0413	—	0.0050	0.0072	—
Completeness%	96.810	92.636	99.841	96.823	92.639	99.861

Figures

 Figure 1. Overview of the sufficient alignment support (SAS) metric for detecting assembly

 errors. SAS identifies structural errors (SEs) and base-level errors (BEs) based on read alignment

 patterns across defined windows. Top panel (SE windows): Structural errors are detected in 2 kb

 windows using ultra-long Oxford Nanopore (UL ONT) reads. Windows are flagged based on five

 indicators: abnormally high or low depth, large insertions or deletions (indels), clipped alignments,

 and lack of spanning reads. Telomeric regions with any supportive reads are not flagged. Bottom

 panel (BE windows): Base-level errors are evaluated in 50 bp windows using reads from high-

 accuracy sequencing platforms (Element, SPRQ, or UL ONT). A window is considered error-free

 only if all bases are covered by perfectly matched, non-clipped reads. In regions not covered by

 Element reads, support from SPRQ or UL ONT reads is used to validate sequence accuracy. SAS

 flags windows lacking sufficient read support (*i.e.*, <10% of average sequencing depth), illustrating

 example regions containing various assembly errors.

YAO v1.1 chr1_pat

YAO v2.0 chr1_pat

Figure 2. Genomic distribution of assembly errors on paternal chromosome 1 before and after

polishing. Genomic features and assembly error profiles for T2T-YAO v1.1(**a**) compared to the

polished T2T-YAO v2.0 (**b**). Tracks from top to bottom: GC% and GA% content; normalized read

depth from Element, PacBio Revio (SPRQ), and ONT ultra-long reads; structural error (SE)

windows identified by SAS and Sniffles2; misjoins and collapses flagged by Nucflag and Flagger;

base-level errors (BEs) categorized as homopolymer-associated (h-BE) and non-homopolymer-

associated (nh-BE); erroneous 31-mers and 21-mers detected by Merqury with binary and trinary

k-mer set.

Figure 3. Counts of base-level and structural errors in T2T-YAO before and after iterative correction. Distribution of structural error (SE) windows (**a**) and base-level error (BE) windows (**b**) across chromosomes in maternal and paternal haplotypes of T2T-YAO v1.1 compared to v2.0. SEs were categorized into five types: low coverage, high coverage, indels (≥50 bp), clipped alignments, and non-spanning regions. BEs are divided into homopolymer-associated BEs (h-BEs) and non-homopolymer-associated BEs (nh-BEs). Bar plots show counts per chromosome, with the left bar

675   in each pair for v1.1 (light colors) and the right for v2.0 (dark colors). The rDNA-associated errors

676   are shown as hatched bars.

a

b

Figure 4. Structural characterization of telomeres and centromeres in the polished T2T-YAO v2.0 assembly.

(**a**) Telomeric repeat content and extension across all 46 chromosomal termini. Each bar represents the distance from the chromosomal terminus, with colored segments indicating specific 6-mer tandem repeats. Canonical (TTAGGG)n (in long arm) and complementary (CCCTAA)n (in short arm) motifs are shown in dark red and blue, along with shallow colors indicate non-canonical telomeric-like 6-mer repeats. (**b**) Annotation of centromeric regions in all the 46 chromosomes. Repeat content was visualized as stacked bars by annotated repeat family. Each pair of centromeres displays a unique repeat composition and organization, with notable differences between haplotypes. The inset shows a zoom-out view of the Y chromosome centromere, which is full of repeat sequences.

Extended Figures

Extended Figure 1. Example of a Structural Error in a Repetitive Region Missed by Sniffles2

Extended Figure 2. Schematic Overview of the Structural Error (SE) Correction Pipeline

Extended Figure 3. Contact Map of Pore-C Reads Aligned to the T2T-YAO v2.0 Assembly

Extended Figure 4. Comparison of Centromere Structures in T2T-YAO v1.1 and v2.0 via Genomic Centromere Profiling (GCP)

Extended Figure 5. Repetitive Elements in Maternal Chromosome 7 with Abnormal ONT Read Depth

Extended Figure 6. Length Distributions of Collapsed and Duplicated Regions Predicted by Flagger

Extended Figure 7. Read Depth Distributions of SPRQ and Element Data in T2T-YAO v1.1 and v2.0

Extended Figure 8. Schematic Overview of the Base-Level Error (BE) Correction Pipeline

Extended Figure 9. Example of Discordance Between ONT and SPRQ Reads at Homopolymer Sites

Extended Figure 10. Quality Score Distribution of SNV-like Errors in T2T-YAO v1.1 and v2.0

Extended Figure 11. Homopolymer Sites in the D4Z4 Region on Chromosome 4q35

Extended Figure 12. Distribution of Homopolymers Across Chromosomes in T2T-YAO v2.0

Extended Figure 13. Patterns of Higher-Order Repeats and Satellite Sequences in the Paternal

Centromere of Chromosome 1 in T2T-YAO v1.1 and v2.0

Supplementary Tables

Table S1. Sequencing Datasets Utilized in This Study

Table S2. Chromosome-Wise Assembly Quality Metrics Before and After Polishing

Table S3. Structural Errors Identified by Multiple Tools Before and After Correction

Table S4. GC and GA/CT Content of Remaining Erroneous 31-mers in T2T-YAO v2.0

Supplement Methods

Supplement Figure 1-45 Distribution of assembly errors before and after correction in each

chromosome except paternal chromosome 1.

Extended Fig. 1

a

chrX_maternal: 70,977,242-70,982.071

Sniffles

ONT

SAS-SE

b

chr2_paternal: 131,255,507-121,280,700

Sniffles

ONT

SAS-SE

c

chr2_paternal: 87,402,206-87,496,228

Sniffles

ONT

SAS-SE

Extended Fig. 2

Extended Fig. 3

Extended Fig. 4

a. YAOv1.1

b. YAOv2.0

Extended Fig. 5

chr7_maternal: 133,025,150-133,364,953

Extended Fig. 6

Extended Fig. 7

a

b

Extended Fig. 8

Extended Fig. 9

chr1_maternal: 8,108,160-8,108,631

chr1_paternal: 15,216,801-15,216,840

chr1_maternal: 46,238,167-46,238,198

Extended Fig. 10

a   V1.1

b   V2.0

Extended Fig. 1

a

chr4_paternal: 191,675,743-191,983,215

b

chr4_paternal: 191,720,591-191,743,567

Extended Fig. 12

AT mononucleotide Distribution Across Chromosomes (500kb bins, >300=max color)

GC mononucleotide Distribution Across Chromosomes (500kb bins, >50=max color)

Dinucleotide Distribution Across Chromosomes (500kb bins, >50=max color)

Extended Fig. 13