1

Spatial Transcriptomics Reveals that the Local Immune Response to Placental Guinea Pig Cytomegalovirus Infection is Driven by Chemokine Signaling

Zachary W. Berkebile,[1] Colleen L. Forster,[2] Emma Stanley,[3] Fernanda M. Rodriguez,[3] Adam C. Herman[4], Micah D. Gearhart[5], Craig J. Bierle[1]*

[1.] Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Minnesota, Minneapolis, Minnesota, United States of America

[2.] Biorepository and Laboratory Services, Clinical and Translational Science Institute, University of Minnesota, Minneapolis, Minnesota, United States of America

[3.] University of Minnesota Genomics Center, Department of Pediatrics, University of Minnesota, Minneapolis, Minnesota, United States of America

[4.] Minnesota Supercomputing Institute, University of Minnesota, Minneapolis, Minnesota, United States of America

[5.] Division of Maternal-Fetal Medicine, Department of Obstetrics, Gynecology and Women's Health, University of Minnesota, Minneapolis, Minnesota, United States of America

* Corresponding Author

E-mail: cjbierle@umn.edu

Short title: Spatial Transcriptomics of GPCMV-infected placentas

Abstract

Cytomegalovirus infection can disrupt placental development and function either by directly infecting placental cells or by eliciting a pathogenic immune response. The relative contributions of these two mechanisms to adverse pregnancy outcomes remains poorly understood. In this study, we used spatial transcriptomics to quantify host and viral gene expression at the maternal-fetal interface at near single-cell resolution. Guinea pig cytomegalovirus (GPCMV) infection after mid-gestation causes focal infections at the base of the main placenta. Samples for spatial transcriptomics were collected from guinea pigs infected with GPCMV at 35 days gestation. Viral loads and the location of infected cells in placentas were assessed using virus-specific droplet digital PCR and *in situ* hybridization at 21 days post-infection. Representative placentas were sectioned onto Visium Spatial Gene Expression Slides and sequencing libraries were prepared from six infected and six uninfected tissue sections. Spatial transcriptomes from 33,687 55-μm spots were generated and used in subsequent analyses. To assess how infection affected gene expression at the maternal-fetal interface, a combination of graph-based clustering and manual classification was used to assign spatial transcriptomes to clusters representative of different anatomic regions. Infection dysregulated more transcripts in the decidua and junctional zone than in the labyrinth or non-capillarized syncytium. Notably, infection downregulated transcripts involved in lipid metabolism and upregulated transcripts involved in antiviral defense and chemokine signaling. A second analysis compared the spatial transcriptomes of GPCMV-infected cells and their immediate microenvironment with similar regions in uninfected placentas. A transcriptional signature indicative of immune activation was clear in this comparison, and the local placental response to cytomegalovirus infection was driven by upregulated chemokine signaling. Thus, spatial transcriptomics revealed regional patterns of gene expression in the guinea pig placenta and illuminated how the host response to GPCMV may compromise placental function.

Author Summary

The placenta supports fetal development while also acting as an immune barrier against bloodborne pathogens. Cytomegalovirus (CMV), the most common viral cause of congenital infections and preventable neurologic disability in children, evades host defenses to infect the placenta. How CMV affects the placenta and fetal health remains poorly understood. Using a guinea pig model of CMV infection during pregnancy and spatial transcriptomics, a recently-developed method that enables gene expression to be studied at near-cellular resolution, this study compared normal and infected placentas. The effects of infection on directly infected cells and their immediate environment and indirect effects that occur at more distant sites were revealed. This information may inform the development of therapies to improve placental function after CMV infection.

Introduction

Human cytomegalovirus (HCMV) is a ubiquitous herpesvirus and the most frequent cause of congenital viral infections [1]. Congenital cytomegalovirus (cCMV) infections occur in roughly 1 in 200 pregnancies and approximately 8000 children are disabled by cCMV each year in the United States [2]. Placental infection plays a central role in the pathogenesis of cCMV. The placenta is critical for supporting fetal development, facilitates the development and maintenance of maternal tolerance, and acts as an immunologic barrier protecting the developing offspring from bloodborne pathogens. A myriad of antiviral defenses, including the constitutive expression of pattern recognition receptors and type III interferon, prevent most viruses from replicating in the placenta and/or infecting the fetus [3]. HCMV can infect both the maternal decidua and the placenta, placental infection likely precedes congenital infection, and

some adverse pregnancy outcomes, including fetal growth restriction and stillbirth, can be

caused by placental HCMV infections that do not transmit to the fetus [4-7].

How HCMV infection affects placental development and function remains poorly

understood [8]. In clinical samples, HCMV antigens are most often detected in the decidua, in

cytotrophoblasts located in floating villi and cell columns, and in the villous stroma and fetal

capillaries [9, 10]. Villous malformations have been noted in cases of early fetal demise after

congenital HCMV infection, and it has been proposed that infection can interfere with

trophoblast proliferation and differentiation to compromise the early placental development [4].

Other less severe pathologies, including a hypoxia-like syndrome, are more frequent and

observed when placental infection occurs later in pregnancy [5, 11]. Isolated primary extravillous

trophoblasts, cytotrophoblasts, and trophoblast progenitor cells can be productively infected by

HCMV *in vitro* [12-15]. Infection reduces the capacity of cytotrophoblasts to proliferate, limits the

differentiation of trophoblast progenitor cells, and interferes with the anchoring of villous

explants *in vitro* [16, 17]. HCMV can infect trophoblast stem cells, where the virus dysregulates

the expression of transcripts involved in differentiation, but these infections are not productive,

have no apparent effect on the proliferation of the trophoblast stem cells, and viral genomes are

not maintained after passage [18, 19]. Similar observations have been made in trophoblast

organoids, which also do not support productive HCMV replication [20].

The inflammatory response to placental infection can cause pregnancy loss, growth

restriction, neurocognitive impairment, and developmental delay [21]. Proinflammatory cytokines

accumulate in amniotic fluid, the placenta, and maternal sera during congenital HCMV infections

[22-24]. Similar inflammatory responses occur when tissue explants and organoids are infected

with HCMV *in vitro* [20, 25, 26]. Therapeutic or genetic interference with inflammatory signaling

pathways can rescue murine pregnancies after infection, exposure to pathogen associated

molecular patterns, or cytokines treatments [27, 28]. The species-specificity of HCMV and

inability of mouse cytomegalovirus to cause placental and congenital infections has limited our

ability to experimentally study whether the inflammatory response to placental infection is a

cause of fetal harm [1, 29].

Guinea pigs are the most widely used animal model of cCMV infection. Guinea pigs

have ~65-day gestational periods and highly-developed, precocious offspring [30]. Major guinea

pig organ and immune systems develop *in utero* to an extent that is more like humans than the

mouse or rat. Guinea pig cytomegalovirus (GPCMV, also known as Caviid herpesvirus 2) was

isolated from the salivary gland of an infected guinea pig and primary GPCMV infection during

pregnancy leads to the sequential infection of the placenta and fetus [31-35]. Fetal outcomes

are affected by the timing of infection during pregnancy [36, 37]. Notably, maternal infection

after mid-gestation causes high rates of stillbirth and fetal growth restriction, and lesions,

including ischemic injury and areas of necrosis, are observed in GPCMV-infected placenta at 14

days post-infection (dpi) and later [33]. We recently compared how maternal GPCMV infection

either late in the embryonic period (21 days gestation [dGA]) or after mid-gestation (35 dGA)

affected the placenta [38]. While there were no significant differences in viral genome

abundance between these two groups at 21 dpi, bulk RNA-seq revealed that significant

changes in placental gene expression were largely restricted to the animals that had been

infected at 35 dGA. *In situ* hybridization further revealed differences in the location of infected

cells in older and younger placenta. These observations led us to hypothesize that the guinea

pig placenta becomes sensitized to infection late in gestation, when a population of GPCMV-

susceptible cells develop near the base of the main placenta, and that the immune response to

these infected cells compromises placental function.

In this study, we used spatial transcriptomics to compare gene expression in age-

matched GPCMV-infected and control placentas at a near-cellular resolution. We found that

infection affects gene expression in anatomic regions, such as the labyrinth, that are not directly

infected by the virus. When gene expression was compared between GPCMV-infected cells and

their immediate microenvironment and a comparable area in mock-infected placentas, we found

that the local inflammatory response to GPCMV was driven by upregulated chemokine gene

expression. Similarities in the placental response to CMV infection between humans and guinea

pigs indicate that the rodent could be a valuable model for testing immunomodulatory therapies

that aim to prevent fetal injury after chronic placental infections.

Results

Spatial transcriptomics of the guinea pig placenta

HCMV and GPCMV both cause focal infections in the placenta [10, 33, 38]. When virus-

specific immunohistochemistry or *in situ* hybridization is used to detect GPCMV, individual

infected cells or small groups of infected cells can be detected in both the placenta and decidua

but are rare [33, 38-40]. However, we have found that comparably large areas of infected cells

can be consistently detected at the base of the main placenta when dams are infected after mid-

gestation and euthanized near term [38]. These cells are often found adjacent to the

subplacenta or maternal arterial channels that perfuse the placenta. We utilized the Visium

Spatial Gene Expression platform (10x Genomics) to study transcription in the guinea pig

placenta and decidua at a near single-cell resolution and illuminate how infection directly and

indirectly affects gene expression. This next generation sequencing-based technology uses

glass slides tiled with 55 µm diameter, oligonucleotide-containing spots; each capture area

contains 5000 of these spots in a 6.5 x 6.5 mm^2 square. The oligonucleotides include location-

based barcodes, unique molecular identifiers, and poly-T sequences that are used to capture

spatial transcriptomes [41]. **Figure 1** summarizes the experimental approach used in this

project. Placentas were collected from guinea pigs that were bred during postpartum estrus,

either GPCMV- or mock-infected at 35 days gestation (dGA), and euthanized 21 days post-

infection (56 dGA) [42]. One half of each placenta was flash frozen in OCT compound for

histopathologic studies and spatial transcriptomics (ST). A portion of the remaining placenta

was collected for DNA extraction; extracted DNA was used for viral load quantification and fetal

sex screening by droplet digital PCR (ddPCR) and endpoint PCR [43, 44]. For ST, the flash

frozen tissue was sectioned onto Visium slides, hematoxylin and eosin (H&E) stained, and

imaged. The tissue was then permeabilized and an Illumina sequencing library was generated

from cDNA synthesized using the oligonucleotides that are tiled on the capture area.

Figure 1. Spatial transcriptomics workflow. Gestational age-matched placentas from

GPCMV- and mock-infected guinea pigs were divided and flash frozen in OCT

compound. Visium Tissue Optimization Slides were used to determine the ideal

incubation time for tissue permeabilization and on-slide cDNA synthesis. GPCMV-

specific RNAscope was used to screen placentas for GPCMV-infected cells and select

representative tissue to analyze on Visium Gene Expression Slides. Placentas were

sectioned onto these slides, H&E stained, and imaged. After on-slide cDNA synthesis

and Illumina sequencing, data was analyzed using Space Ranger and Partek Flow

software. Gene expression in GPCMV- and mock-infected tissue was compared after

graph based clustering and spatial transcriptomes that contained infected cells were

identified based on their viral read content.

On-slide cDNA synthesis was optimized using Visium Tissue Optimization Slides (10x

Genomics). Fluorescence microscopy revealed that more cDNA was produced from the main

placenta than either the decidua or the subplacenta, regardless of how long the tissue had been

incubated with permeabilization buffer (**Fig. 1**). This reflects the higher RNA content of the

placenta, and we have observed that an average of 2.5±0.71-fold more RNA is extracted per

mg of placenta than decidua in other unpublished experiments.

RNAscope specific to the viral transcript *gp3* was used to screen for infected cells and

identify representative GPCMV-infected samples (**S1 Fig.**); five placentas (three female and two

male) collected from three dams were selected for ST. GPCMV-infected cells were detected at

the base of the main placenta in all these placentas. Six age-matched placentas from three

mock-infected dams were used as controls. The guinea pig placenta is significantly larger than

6.5x6.5 mm^2 at 56 dGA and each placenta was trimmed to so that portions of the placenta,

decidua, and subplacenta would be included on each capture area. However, examination of

the H&E-stained gene expression slides revealed that the subplacenta was not present in

several tissue sections (**S2 Fig.**). To verify that infected cells were present in each capture area,

a serial section was collected from each trimmed placenta and used for a second round of

RNAscope (**S3 Fig.**). In the initial RNAscope analysis, one placenta (2395B) had a large area of

gp3$^+$ cells in the decidua; this placenta was divided and sectioned onto two capture areas.

Sample 2395B1, which should have corresponded to this area of infected decidua, did not

contain gp3$^+$ cells in the second round of RNAscope. GPCMV infected cells were detected

where expected in all other samples.

cDNA from the twelve capture areas was used to prepare sequencing libraries, pooled,

and sequenced. Apart from sample 735B, more than 5X10^4 reads were sequenced per spot and

a median of 3900 host transcripts were detected in each spatial transcriptome (**S1 Table**). In

GPCMV-infected placentas, a median of 1.45X10^5 (range 3X10^3 to 2.6X10^5) reads mapped to

the viral genome per sample. These reads mapped to the 3' end of viral transcripts and

spanned the entire GPCMV genome (**Fig. 2**). Fewer than 100 reads mapped to the viral

genome in mock-infected tissue sections and these reads were generally highly repetitive

sequences.

Figure 2. GPCMV sequences detected in spatial transcriptomes aligned to the viral

genome. Sequencing reads were aligned to the GPCMV genome (NC_020231.1) using

SpaceRanger and visualized with Geneious (v11.1.5). The total number of reads

mapping to the viral genome in each sample is shown. Coverage at each nucleotide

position is graphed in log scale. Regions with coverage <1 read are highlighted in blue

and >100 in orange.

GPCMV infection alters gene expression across the distinct anatomic regions of the guinea pig maternal-fetal interface

A combination of graph-based clustering and manual refinement was used to compare

gene expression across the different anatomic regions of the guinea pig placenta. The 3000

most variant guinea pig transcripts were used for graph-based clustering; viral transcripts were

excluded from this analysis. Ten clusters were produced when a resolution setting of 0.5 was

used for graph-based clustering (**S4 Fig.**). Cluster assignment generally reflected where spatial

transcriptomes originated at the maternal-fetal interface. The computationally generated clusters

were then manually refined. Firstly, data from spatial transcriptomes located in folded tissue or

near large blood spaces were excluded from later analyses. Three clusters that included the

interlobar or marginal syncytium were combined into a single cluster representative of all non-

capillarized syncytium and three clusters were combined and classified as parts of the placental

stem and junctional zone. Seven clusters remained after manual spot reassignment (**Fig. 3**).

While this underrepresents the complexity of the placenta, a conservative approach allowed us

to compare gene expression across many spatial transcriptomes per cluster. The capillarized

syncytium was represented by two clusters that comprise the core and periphery of the

labyrinth. The non-capillarized syncytium and subplacenta were each represented by a single

cluster. The remaining transcriptomes, located in the junctional zone, the stem of the placenta,

and the decidua, formed three clusters, the smallest of which was comprised of only 389

transcriptomes and was excluded from later analyses. Consistent with our observation that the

subplacenta was not sampled in several tissue sections, spots assigned to the subplacenta

cluster were not present in all twelve capture areas (**Fig. 4a**).

Figure 3. Gene expression in spatial transcriptomes reflects their location at the

maternal-fetal interface. Spatial transcriptomes were classified using a combination of

graph-based clustering and manual reassignment. (**A**) UMAP projection showing

transcriptomes colored by the cluster assignment used for differential expression

analyses. LP- labyrinth periphery, LC- labyrinth core, NS- non-capillarized syncytium,

DE- decidua, SP- subplacenta, N/A- not assigned, JZ-junctional zone and placental

stem. **(B)** UMAP projection showing spots colored by sample identifier, with GPCMV-

infected samples labeled in red, illustrating the integration of data from three Visium

slides and twelve capture areas. **(C)** Representative sections of mock- and GPCMV-

infected placenta. The H&E stained tissue sections are shown on the left. Spatial

transcriptomes were colored by cluster assignment in the panels on the right.

Transcriptomes that were excluded from subsequent analyses are not colored.

Figure 4. **GPCMV infection dysregulates gene expression in the placenta and**

decidua. Gene expression was compared between sections of GPCMV- and mock-

infected placenta (N=6/group). **(A)** The proportion of spatial transcriptomes in each

capture area that were assigned to a given cluster is illustrated by bar chart. **(B)**

Transcripts that were differentially regulated by GPCMV infection in each cluster were

identified by gene-specific analysis (two-fold or greater change, p ≤0.05). The total

number of spatial transcriptomes from GPCMV- and mock-infected capture areas

included in this analysis and number of genes that were up- or down-regulated in each

comparison are shown.

Cluster-specific biomarkers were identified by using Student's t-tests to compare gene

expression in the spatial transcriptomes assigned to one cluster with the combined data from all

other spatial transcriptomes. **S2 Table** ranks the top 20 biomarkers for each cluster by

significance and fold change. To illustrate how biomarker expression varies in the placenta and

decidua, expression data for select biomarkers was overlaid onto images of representative

control and GPCMV-infected placentas (**S5 Fig.**). The biomarkers for the two labyrinth clusters

were partially overlapping. Conversely, the decidua and subplacenta clusters both had a high

number of highly expressed biomarkers that were unique to the regions. Interestingly, several

prolactin-like transcripts were identified as biomarkers for the labyrinth and non-capillarized

syncytium. Guinea pigs, like murid rodents and ruminants, have an expanded prolactin locus

and two of these transcripts, *Prlrp1* and *Prlrp2,* were previously shown to be expressed in the

guinea pig placenta [45]. Our analysis found that *Prlrp1* was a biomarker for the labyrinth

clusters and *Prlrp2* was enriched in the non-capillarized syncytium but expressed at a high level

throughout the placenta and decidua. A third, previously uncharacterized prolactin-like

transcript, *Ghc2* (LOC111754958), was a biomarker of the non-capillarized syncytium. These

prolactin-like transcripts may have evolved to regulate placental development in guinea pigs.

To assess how GPCMV infection affected gene expression in the different regions of the

placenta, the spatial transcriptomes of infected and mock-infected tissues were compared on a

cluster-by-cluster basis. For this analysis, gene expression data was normalized to counts per

million plus one and $\log_2$ transformed. Differentially regulated transcripts were identified by

gene-specific analysis. Broadly, GPCMV infection downregulated more host transcripts than it

upregulated (**Fig. 4b**). Functional enrichment analyses of the differentially regulated transcripts

was done with g:Profiler [46]. Infection had a comparably modest effect on gene expression in

the labyrinth, the non-capillarized syncytium, and the subplacenta. Several terms related to an

antiviral response, including immune response (GO:0006955), type I interferon production

(GO:0032606), and chemokine activity (GO:0008009), were enriched in these three regions

after GPCMV infection (**S5 Fig.**). Terms related to metabolic functions such as peptidase

activity, sterol transport, and signaling receptor binding and activity also were enriched in these

clusters. GPCMV-infected cells were rarely detected in these regions by RNAscope and few of

the spatial transcriptomes assigned to these clusters contained viral reads. Thus, it can be

inferred that infection-associated changes in gene expression in the labyrinth, non-capillarized

syncytium, and subplacenta are largely indirect effects of viral infection elsewhere in the

placenta.

GPCMV infection affected markedly more transcripts in the two clusters that mapped to

the decidua or placental stem and junctional zone. Most GPCMV-infected cells were assigned

to the later cluster and 74 immune response (GO:0006955) transcripts, including many involved

in chemokine signaling, were dysregulated by infection (**Fig. S5**). Viral infection downregulated

transcripts with functions related to protein binding (GO:0005515), lipid transport activity

(GO:0005319), small molecule metabolic processes (GO:0044281), and developmental

processes (GO:0032502) in these tissue compartments. Taken together, this differential gene

expression analyses revealed how GPCMV infection affects regions of the placenta that are not

directly infected. While GPCMV infection dysregulates fewer transcripts in the main placenta

than in the decidua, placental stem, or junctional zone, the upregulation of *Irf7* and select

interferon stimulated genes (e.g. *Isg15*, *OasL*, and *Rsad2*) provides evidence that an antiviral

response occurs throughout the placenta. The dysregulation of transcripts involved in metabolic

or developmental processes suggests that GPCMV infection reprograms gene expression

throughout the decidua and placenta in ways that could be harmful to fetal development.

Gene expression changes in the GPCMV-infected microenvironment

GPCMV transcripts are polyadenylated and spatial transcriptomes that contained viral

reads were identified in all six samples of GPCMV-infected placenta (**Fig. 5**). To determine how

GPCMV affects transcription in infected cells and cells in their immediate vicinity, we compared

the results of *gp3*-specific RNAscope with the number of GPCMV reads that were detected per

transcriptome (**S2 and S6 Figs.**). Spots that contained more than one hundred viral reads

(N=99) colocalized with *gp3*+ cells in the serial tissue section and were annotated as GPCMV-

infected. Sample 2395B1 was excluded from this analysis because it lacked spatial

transcriptomes that contained more than one hundred viral reads and *gp3*-positive cells. Spots

that were located within 110 μm of GPCMV-infected cells were classified as part of the GPCMV-

infected microenvironment (N=285) for this differential gene expression analysis. In total, 384

spatial transcriptomes from GPCMV-infected placentas were compared with 1148 spatial

transcriptomes from the mock-infected tissue sections that were similarly located in the placenta

(Fig. S6). Most spatial transcriptomes included in this analysis had been assigned to either the

placental stem and junctional zone or the non-capillarized syncytium clusters (**Fig. 6a**).

Figure 5. GPCMV-infected cells are detected in spatial transcriptomes.

SpaceRanger was used to map reads to the GPCMV genome and spatial

transcriptomes that contained viral transcripts were identified. (**A**) The number of

GPCMV-specific reads in each transcriptome was plotted by sample identifier. GPCMV-

infected samples are noted in red, and the data was colored by the viral read count. (**B**)

Sequencing data was overlaid onto an image of a GPCMV-infected sample (2421A).

Spots were colored based on GPCMV-read count as in (A). (**C**) For differential gene

expression analyses, transcriptomes that contained >100 GPCMV reads were defined

as "infected" (blue) and any spot within 100 μM was classified as in the GPCMV-infected

microenvironment (red).

Figure 6. Chemokine and T cell-associated transcripts are upregulated in or near

GPCMV-infected cells. Gene expression was compared between spatial

transcriptomes that contained ≥100 GPCMV reads or were within 110 μm of infected

cells and a comparable area in mock-infected tissue. (**A**) Bar graph illustrating the

cluster assignments for the two groups of transcriptomes that were compared in this

analysis. (**B**) Transcript abundance illustrated by volcano plot, highlighting transcripts

with chemokine activity (GO:0008009). (**C**) Violin plots displaying the expression of

select chemokine transcripts or transcripts related to a cytotoxic immune response in

individual transcriptomes. (**D**) g:Profiler was used for functional enrichment analysis to

identify terms that are enriched among the transcripts that are differentially regulated by

GPCMV infection. Select gene ontology terms are highlighted on a modified Manhattan

plot.

448 guinea pig transcripts were upregulated in or near GPCMV-infected cells and 530

were downregulated (≥2-fold change, *p*≤0.05, **Fig. 6b**). Gene ontology analysis revealed a

transcriptional signature indicative of immune activation in the GPCMV-infected tissues (**Fig.**

6c). Notably, the inflammatory response to GPCMV infection appears to be predominantly

driven by chemokine signaling. 16 of 18 chemokine transcripts that were enriched in or near

GPCMV infected cells were upregulated, including multiple chemokines that signal through

Cxcr3 and several chemokines (*Ccl2*-like [LOC100717552], *Ccl3*-like [LOC100734903 and

LOC100735187], and *Ccl15*-like [LOC101788094]) that may be unique to guinea pigs (**Fig. 6b**).

67 transcripts involved in leukocyte activation (GO:0045321), including markers of a cytotoxic

immune response such as *Cd3d*, *Cd8a*, and granzyme transcripts, were also more highly

expressed at sites of GPCMV infection. Cytokines that are associated with acute placental

infection (e.g. IL-1β, TNFα, IL-6, and IL-8) were not differentially regulated by GPCMV.

Similarly, while numerous interferon stimulated gene transcripts (e.g. *Isg15*) were upregulated

by infection, transcripts encoding interferons were not affected by GPCMV infection.

Cumulatively, spatial transcriptomics resolved direct and indirect effects of GPCMV infection in

the placenta and has revealed the central role of chemokine signaling in the GPCMV-infected

placenta.

Discussion

Placental infection plays a central role in CMV pathogenesis during pregnancy. In this study, we used ST to investigate how GPCMV affects gene expression at the maternal-fetal interface. With its 55-μm spot resolution, the Visium Spatial Gene Expression Platform enabled the most detailed analysis of transcription in the guinea pig placenta and decidua to date, revealing how infection impacts transcription in infected cells and their immediate microenvironment and illuminating how regions of the placenta are affected by the virus when not directly infected.

Research in primary trophoblasts, tissue explants, and organoids has provided insights into how HCMV affects the maternal-fetal interface, but these models have notable limitations [18, 20, 25, 26, 47, 48]. During *in vitro* experiments, placental cells or tissue explants are generally exposed to high infectious doses of virus by non-physiologic routes. Samples are analyzed within days of the initial infection, and the resulting data may reflect features of the innate host response to HCMV. Bulk RNA seq experiments have found that HCMV elicits markedly different transcriptional responses from decidua and placenta [20, 26]. Moreover, first and third trimester trophoblasts and tissue explants are differentially susceptible and responsive to HCMV infection [5, 26, 49, 50]. Thus, how CMV infection can cause placental injury likely depends upon when infection occurs during gestation.

Clinically, HCMV causes chronic placental infections that are characterized by focal lesions and a heterogeneous population of maternal and fetal leukocytes respond to infection [4, 10]. This complexity cannot be easily modeled *in vitro*, but studying samples of infected placenta collected after congenital infections presents its own challenges. Unlike Zika or rubella viruses, primary HCMV infection does not present with diagnostic clinical signs (e.g. rash), molecular methods must be used to confirm infection during pregnancy, most cases are not

identified until after birth. Sample heterogeneity further complicates the analysis of clinical

samples. For example, ST was used to study placentas collected after SARS-Cov-2 infection

during pregnancy [51]. Most of nine samples analyzed did not contain virus that could be

detected by qRT-PCR, *in situ* hybridization, or ST. The highest viral loads and most significant

infection-associated changes in gene expression were found in two placentas recovered after

intrauterine fetal demise at significantly different times in pregnancy (22.3 versus 35.0 weeks).

Experimental studies in animal models thus play an important role in revealing how CMV and

other viral infections can affect the maternal-fetal interface.

Guinea pigs have long been used for preclinical CMV vaccine development, but the

limited availability of species-specific reagents has been a barrier to more sophisticated studies

of CMV pathogenesis [52]. Species-agnostic next generation sequencing technology has

revolutionized discovery science in guinea pigs and other unconventional animal models. For

example, scRNA-Seq revealed that naked mole-rats, a hystricomorph rodent like the guinea pig,

lack conventional natural killer cells [53]. We have previously used bulk RNA-seq to examine

how guinea pig amniotic epithelial cells respond to infection *in vitro* and assess how primary

GPCMV infection during the embryonic period or after mid-gestation differentially affect the

placenta [38, 43]. However, bulk RNA-seq and most other conventional gene expression

analyses quantify transcript abundance in tissue homogenates. This effectively averages

transcript abundance across numerous cell types and states. ST enables gene expression

analysis at a near single-cell resolution while preserving cellular location in the tissue of interest

[54]. While the fresh-frozen Visium Spatial Gene Expression workflow is not reliant on species

specific reagents, the 55-μm resolution of each spatial transcriptome is a limitation of the

technology. Depending on the cellular density in tissue being analyzed, each spatial

transcriptome reflects the RNA content of one to ten cells. Computational approaches have

been developed that can infer the cellular composition within spots, as demonstrated by recent

studies that used sc- or snRNA-Seq datasets to study the human maternal-fetal interface [51,

55]. While cell atlases have been developed for the human and mouse placenta, a similar

resource has yet to be developed for the guinea pig and an automated cell type classification

was not done in this study. [56-59]

This study revealed that GPCMV infection dysregulates more transcripts in the decidua

and junctional zone than in the labyrinth or non-capillarized syncytium. HCMV antigens and

DNA are more frequently detected in decidua than in placenta, but GPCMV is only sporadically

detected in the decidua by *in situ* hybridization at 21 dpi (**Fig. S1**) [38, 60]. Additional

experiments are necessary to assess whether GPCMV infects the decidua at earlier times in

infection and resolves. One observation from gene set enrichment analysis was that GPCMV

infection downregulates lipid metabolism and transport in the maternal-fetal interface. Fetal

development and growth are dependent upon the regulated transport of maternal nutrients,

including fatty acids, across the placenta and maternal hyperlipidemia is typical during a healthy

pregnancy [61, 62]. Abnormal maternal lipid levels are observed during pregnancy-induced

hypertension, preeclampsia, and preterm birth [63-65]. At a cellular level, DNA viruses hijack

lipid metabolism pathways to promote replication, and HCMV infection promotes fatty acid

synthesis and modifies cholesterol synthesis and trafficking *in vitro* [66-70]. Given that ZIKV

infection also perturbs lipid homeostasis in the placenta, altered lipid metabolism may a

mechanism by which viruses cause placental insufficiency [71].

Placental dysfunction and adverse fetal outcomes can be caused by interferon signaling

and other inflammatory responses to infection [8, 72]. Infection is a leading cause of preterm

birth, and the acute inflammatory responses to ascending bacterial infection involves cytokines,

including IL-1β, IL-6, IL-8, and TNF-α, that are normally upregulated during parturition [73].

Polyinosinic:polycytidylic acid (Poly I:C), a synthetic analog of the viral pathogen associated

molecular pattern double-stranded RNA, can affect placental and fetal development in mice and

rats depending upon the dose, route of administration, and the timing of treatment during

pregnancy [reviewed in [74]]. Experiments in *Ifnar*-transgenic mice have demonstrated that fetal

growth restriction and resorption can be caused by Type I interferon signaling that occurs in the

Zika virus (ZIKV)-infected fetus and placenta [27, 75, 76]. In humans, Aicardi–Goutières (AGS)

syndrome, a rare inherited disease characterized by dysregulated Type I interferon production,

can present during pregnancy with ultrasound findings reminiscent of congenital infections [77].

Multiple studies have found that cCMV causes proinflammatory cytokines to accumulate

in amniotic fluid, and elevated CXCL10 levels are a biomarker of congenital infection [22-24].

CMV infection also upregulates *CXCL10* transcription in tissue explants, amniotic epithelial

cells, and decidual organoids *in vitro* [20, 26, 43]. Elevated amniotic fluid concentrations of

CXCL10 also occurs during chronic chorioamnionitis and maternal anti-fetal rejection; the

chemokine may drive T cell recruitment to the placenta and/or fetal membranes [78-80]. Bulk

RNA-Seq found that *Cxcl10* is upregulated in the GPCMV-infected placenta, and ST confirmed

this finding while also revealing that a broad mix of chemokines are upregulated in the

microenvironment around GPCMV-infected cells [38]. Curiously, while individual interferon

stimulated genes are upregulated during placental GPCMV infection, interferon transcripts were

not regulated by infection. Interestingly, one study found no association between CXCL10

concentrations and the detection of HHV6, parvovirus B19, or Epstein-Barr virus in amniotic

fluid [23]. Nonhuman primate studies also indicate that the inflammatory response to CMV may

be distinct from other viral causes of placental and congenital infections such as ZIKV [81, 82].

Primates and rodents diverged 100 million years ago, and the cytomegaloviruses have

been cospeciating with their hosts ever since [83, 84]. The species-specificity of HCMV

precludes the direct study of the virus in animal models, and while *in vivo* experiments using

closely related primate and guinea pig cytomegaloviruses can reveal how infection affects the

placenta under controlled experimental conditions, virus and host-specific immune adaptations

could limit the translatability of these studies to humans. For example, GPCMV-specific

adaptations to its host's immune system include the convergent evolution of class I MHC mimics

and a unique viral chemokine, *gp1* [85, 86]. Guinea pigs and other caviomorph rodents have a

unique population of mononuclear cells, Kurloff cells, with natural killer (NK) cell activities [87].

While direct functional comparisons to human or mouse decidual NK cells have yet to be

completed, the abundance of circulating Kurloff cells is regulated by estrogen and the cells

localize to the placenta and uterus in pregnant animals [88, 89]. While this study and our

previous bulk RNA-seq of the guinea pig placenta provide valuable insights into the biology of

the maternal-fetal interface in guinea pigs, more extensive studies of placental gene expression

and guinea pig immunity are merited to support the use of the rodent as a comparative model

for human gestation [38].

Materials and Methods.

Ethics Statement.

All animal procedures were conducted in accordance with protocols approved by the

Institutional Animal Care and Use Committee (IACUC) at the University of Minnesota,

Minneapolis (Protocol IDs: 1810-36403A and 2106-39180A). Experimental protocols and

endpoints were developed in strict accordance to the National Institutes of Health Office of

Laboratory Animal Welfare (Animal Welfare Assurance #A3456-01), Public Health Service

Policy on Humane Care and Use of Laboratory Animals, and United States Department of

Agriculture Animal Welfare Act guidelines and regulations (USDA Registration # 41-R-0005)

with the oversight and approval of the IACUC. Guinea pigs were housed in a facility maintained

by the University of Minnesota Research Animal Resources, who were accredited through the

Association for Assessment and Accreditation of Laboratory Animal Care, International

(AAALAC). All procedures were conducted by trained personnel under the supervision of

veterinary staff.

Infection of Pregnant Guinea Pigs and Tissue Collection.

GPCMV seronegative, pregnant guinea pigs were mock- or GPCMV- infected as

previously described [38]. Briefly, two- to three-month-old strain 2 females were housed with

strain 2 males until pregnancy could be confirmed by palpation. At this point, the strain 2 male

was replaced with a strain 13 male, who remained with the female until three days postpartum

to establish a second, timed pregnancy [42]. Progesterone ELISA and/or palpation was used to

confirm the second pregnancy. At 35 days dGA, guinea pigs were injected subcutaneously into

the scruff of the neck with 0.5 mL of GPCMV-containing PBS (2×10^5 PFU, SG13J2) or PBS

alone [43]. Blood and plasma were collected from dams every seven days post-infection (dpi)

until the animals were euthanized at 21 dpi. Several dams also received intraperitoneal (IP)

injections of 1.5 mg of rat IgG (2 mg/mL in PBS) at days 0, 7, and 14 post-infection or an IP

injection of 80 mg of Hypoxyprobe 1 (pimonidazole HCl) dissolved in PBS that was administered

90 minutes before the animal was euthanized (Supplementary Table 1). During tissue collection,

the uterus was dissected, the decidua gently separated from the uterus, and each placenta was

halved. One half of each placenta was embedded in optimal cutting temperature (OCT)

compound and flash frozen in isopentane chilled with a dry ice ethanol bath. A portion of the

remaining tissue was frozen at -80ºC for DNA extraction and viral load quantification.

Screening Placentas for GPCMV Infection and Fetal Sex.

DNA was extracted from placentas using the DNeasy Blood and Tissue Kit (Qiagen).

The abundance of GPCMV genomes in the tissue was quantified by ddPCR using primers and

probes specific to *GP54* and the Bio-Rad QX200 system as previously described [43]. A

multiplexed endpoint PCR assay that targets *Sry* (Y-specific) and *Dmd* (X-specific) was used to

determine the sex of each placenta [44]. The location of GPCMV-infected cells in placenta was

determined using an RNAscope® assay specific to the viral transcript *gp3* (Probe V-CavHV-2-

gp3) [38]. Fresh-frozen tissue was prepared for single-plex RNAscope® 2.5 assays as

described in technical note 320536-TN Date 11112016. 10-µm sections of placenta were

mounted onto Superfrost Plus slides (ThermoFisher), fixed using 4% paraformaldehyde, and

dehydrated by sequential immersion in 50%, 70%, and 100% ethanol. After the slides were

dried, the tissue was treated with RNAscope® hydrogen peroxide and Protease IV. The slides

were then stained with RNAscope® 2.5 HD Detection Reagent – RED (322360-USM Rev. Date

11052015) as previously described (9).

Spatial Transcriptomics

To optimize on-slide cDNA synthesis for ST, mock-infected placentas collected at 42

and 56 dGA were analyzed using Tissue Optimization Slides and Reagents (10x Genomics,

PN-1000194). Blocks of frozen placenta were trimmed with the goal of including the main

placenta, placental stem, and decidua on each capture area. 10 µm tissue sections were placed

onto seven of the eight capture areas on the tissue optimization slide (CG000240 Rev B).

Tissue was fixed with methanol, H&E stained, and imaged with a Huron TissueScope LE

(CG000160 Rev A). For tissue permeabilization and fluorescent cDNA synthesis, capture areas

were incubated with Permeabilization Enzyme for times that ranged from 3 to 30 minutes

(CG000238 Rev C). After reverse transcription, slides were imaged with a Nikon Eclipse

fluorescent microscope. An 18-minute permeabilization time was selected for ST experiments

based on the quality of the fluorescent signal. Irrespective of the permeabilization time, more

signal was observed in the main placenta than in the subplacenta, placental stem, or decidua.

Five GPCMV-infected and six control placentas were sectioned onto the twelve capture

areas of three Visium Spatial Gene Expression Slides. These tissues originated from three

GPCMV- and three mock-infected dams and were a mixture of males and females. Blocks were

roughly trimmed to 6.5x6.5 mm^2 so that a similar region of interest would be included in each

capture area and, for GPCMV-infected samples, infected cells would be included in the capture

areas. One GPCMV-infected placenta (2395B) was divided and sectioned onto two capture

areas. The slides were fixed, H&E stained, and imaged with a Huron TissueScope LE. The

tissue was then permeabilized for 18 minutes and cDNA synthesis, amplification, and

sequencing library preparation was completed using the manufacturer recommended protocol

(CG000239 Rev C). Visium libraries were combined and sequenced on a single Illumina

NovaSeq S4 lane (2 x 150 PE run format). Sequencing results are summarized in

Supplementary Table 1. A median of 2.2X10^8 reads per capture area (range 1.0X10^8 to

6.9X10^8) and 7.3X10^4 reads per spatial transcriptome (range 3.3X10^4 to 1.8X10^5) were

sequenced. Sequencing data has been uploaded to the Gene Expression Omnibus (Series

GSE303764, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE303764).

522   Data analysis

To analyze ST data, a custom reference sequence that combined the guinea pig

genome (Cavpor3.0, GCF_000151735.1, Annotation Release 103), the GPCMV genome (strain

22122, NC_020231), and a list of Y chromosome transcripts, was prepared [90, 91]. Space

Ranger (10x Genomics, version 1.3.0) was used to process the raw Illumina sequencing data.

Space Ranger outputs were uploaded to *Partek® Flow®* software (v10.0). Reads that aligned to

the GPCMV genome from each sample of infected placenta were visualized with Geneious

(v11.1.5). Each capture area was manually examined and 927 spatial transcriptomes that were

located under folded or visibly damaged tissue were excluded from downstream analyses. For

graph-based clustering, data from the remaining 33,687 spots was filtered to the 3000 most

variable features, normalized with SCTransform, and batch corrected with Seurat3 integration

[92, 93]. 50 principal components were used for graph-based clustering and data presentation

by Uniform Manifold Approximation and Projection (UMAP) dimensional reduction. Sensitivities

from 0.3 to 1.0 were evaluated using the Louvain clustering algorithm. Higher resolution settings

predictably identified more clusters but also resulted in considerable variation in the number of

spatial transcriptomes assigned to each cluster across the twelve samples. A resolution setting

of 0.5 was selected for subsequent analyses. The data was further refined by excluding 1018

spatial transcriptomes that were located at the edges of tissue sections and combining several

clusters. Cluster-specific biomarkers were identified using Partek Flow's "compute biomarker"

tool. This algorithm compared the expression of features in each cluster with the expression

observed in all other clusters combined using student's t-tests.

To identify host transcripts that were differentially regulated by GPCMV infection, the

abundance of all transcripts that were detected in >0.1% of spatial transcriptomes was

normalized to counts per million plus one and $\log_2$ transformed. Differentially regulated

transcripts were identified by gene-specific analysis (GSA). In one analysis, transcripts that

were differentially regulated by infection were identified in each of the previously identified

clusters. In a second analysis, spatial transcriptomes that contained ≥100 GPCMV transcripts

and every spatial transcriptome within 110 µm were defined as the GPCMV-infected

microenvironment. Transcript abundance was compared with similar area that was manually

selected from mock-infected tissue sections, and 384 spatial transcriptomes from GPCMV-

infected placentas were compared with 1148 spatial transcriptomes from control tissue. For both

workflows, functional enrichment analyses on differentially regulated transcripts (twofold or

more, $p < 0.05$) were completed using g:Profiler [46].

Acknowledgements

556   This work was supported by the National Institutes of Health's National Center for Advancing

557   Translational Sciences, which provided funding for the Biorepository and Laboratory Services

558 Team, and the resources and staff at the University of Minnesota University Imaging Centers

559 (SCR_020997) and the Genomics Center (https://genomics.umn.edu). The content of this article

560 is solely the responsibility of the authors and does not necessarily represent the official views of

561 the National Institutes of Health's National Center for Advancing Translational Sciences.

562 **Competing interests.**

563 The authors have no competing interests to declare.

565   **Figure 1.**

570 **Figure 2.**

573   **Figure 3.**

577   **Figure 4.**

580 **Figure 5.**

583   **Figure 6.**

1. **Chemokine activity** (GO:0008009)
2. **Immune system process** (GO:0002376)
3. **Leukocyte activation** (GO:0045321)

Supporting Information.

S1 Figure. GPCMV-specific RNAscope.
Fresh-frozen, GPCMV-infected placentas were cross-sectioned and stained with a RNAscope assay that detects the viral transcript *gp3* (red). Scans of the five infected placentas that were selected for ST are shown.

S2 Figure. Hematoxylin and eosin-stained guinea pig placentas. GPCMV- (Red, N=5) and mock-infected placentas (N=6) were trimmed and sectioned onto twelve capture areas. The Visium Spatial Gene Expression Slides were H&E stained and imaged. The images were cropped to each capture area's fiducial frame and oriented so that the main placenta is up and decidua down.

602 **S3 Figure. GPCMV-specific RNAscope of trimmed placentas.** Placentas were trimmed to
603 accommodate the 6.5X6.5 mm^2 Visium capture area. After tissue was sectioned on the Visium
604 Gene Expression Slides, the serial section from each sample was used for gp3-specific
605 RNAscope (red) to confirm that infected cells had been placed within each capture area.

S4 Figure. Spatial transcriptomes cluster by anatomic region. Graph-based clustering
assigned spatial transcriptomes to ten groups. These were further manually refined to seven
clusters, which included excluding data under folded tissue sections, near large blood spaces,
and combining several smaller clusters into groups representing non-capillarized syncytium and
the junctional zone and placenta stem.

2396A **2395B2**

Prlp1

Ghc2 (LOC111754958)

Mfsd2a

Pvalb

Vcan

1. LP 2. LC 3. NS 4. DE 5. SP 7. JZ

S5 Figure. Biomarker expression in the anatomic regions of the guinea pig placenta. Transcripts were identified that were significantly enriched in each cluster. Gene expression data for select biomarkers was overlaid on two representative sections of placenta such that transcript abundance in each spot is represented by green intensity. Violin pots illustrate the expression of each biomarker in all samples with transcriptomes classified by cluster.

S6 Figure. Functional enrichment analyses reveal pathways dysregulated by GPCMV infection. For each anatomic region of the placenta, transcripts that were significantly up- or down-regulated in GPCMV-infected placenta were identified using GSA. Functional enrichment analyses were done using g:Profiler [46]. Enriched pathways are illustrated by Manhattan plot and the ten most significantly enriched molecular functions and biological processes for each cluster are indicated.

S7 Figure. Identification of GPCMV-infected cells and their microenvironment in spatial
transcriptomes. Transcriptomes that contained GPCMV reads were identified in each capture
area. In GPCMV-infected samples, spatial transcriptomes were either colored by the abundance
of viral reads or red for >100 reads or blue to designate the microenvironment near infected
cells. In mock-infected samples, transcriptomes colored in yellow were selected for the
differential gene expression analysis.

References

1.   Mocarski Jr ES, Shenk T, Griffiths PD, Pass RF. Cytomegaloviruses. In: Knipe DM,
Howley PM, editors. Fields virology. 2. 6th ed. Philadelphia, PA: Wolters Kluwer/Lippincott
Williams & Wilkins Health; 2013. p. 1960-2014.

2.   Kenneson A, Cannon MJ. Review and meta-analysis of the epidemiology of congenital
cytomegalovirus (CMV) infection. Rev Med Virol. 2007;17(4):253-76. Epub 2007/06/21. doi:
10.1002/rmv.535. PubMed PMID: 17579921.

3.   Megli CJ, Coyne CB. Infections at the maternal-fetal interface: an overview of
pathogenesis and defence. Nat Rev Microbiol. 2022;20(2):67-82. Epub 20210825. doi:
10.1038/s41579-021-00610-y. PubMed PMID: 34433930; PubMed Central PMCID:
PMCPMC8386341.

4.   Garcia AG, Fonseca EF, Marques RL, Lobato YY. Placental morphology in
cytomegalovirus infection. Placenta. 1989;10(1):1-18. Epub 1989/01/01. doi: 10.1016/0143-
4004(89)90002-7. PubMed PMID: 2541426.

5.   Maidji E, Nigro G, Tabata T, McDonagh S, Nozawa N, Shiboski S, et al. Antibody
treatment promotes compensation for human cytomegalovirus-induced pathogenesis and a
hypoxia-like condition in placentas with congenital infection. Am J Pathol. 2010;177(3):1298-
310. Epub 20100722. doi: 10.2353/ajpath.2010.091210. PubMed PMID: 20651234; PubMed
Central PMCID: PMCPMC2928963.

6.   Iwasenko JM, Howard J, Arbuckle S, Graf N, Hall B, Craig ME, Rawlinson WD. Human
cytomegalovirus infection is detected frequently in stillbirths and is associated with fetal
thrombotic vasculopathy. J Infect Dis. 2011;203(11):1526-33. Epub 2011/05/20. doi:
10.1093/infdis/jir121. PubMed PMID: 21592980.

7.   Pereira L, Petitt M, Fong A, Tsuge M, Tabata T, Fang-Hoover J, et al. Intrauterine
growth restriction caused by underlying congenital cytomegalovirus infection. J Infect Dis.

2014;209(10):1573-84. Epub 20140107. doi: 10.1093/infdis/jiu019. PubMed PMID: 24403553;

PubMed Central PMCID: PMCPMC3997585.

8.   Yockey LJ, Lucas C, Iwasaki A. Contributions of maternal and fetal antiviral immunity in

congenital disease. Science. 2020;368(6491):608-12. Epub 2020/05/10. doi:

10.1126/science.aaz1960. PubMed PMID: 32381717.

9.   Pereira L, Maidji E, McDonagh S, Tabata T. Insights into viral transmission at the

uterine-placental interface. Trends Microbiol. 2005;13(4):164-74. doi: 10.1016/j.tim.2005.02.009.

PubMed PMID: 15817386.

10.   Uenaka M, Morizane M, Tanimura K, Deguchi M, Kanzawa M, Itoh T, Yamada H.

Histopathological analysis of placentas with congenital cytomegalovirus infection. Placenta.

2019;75:62-7. Epub 20190104. doi: 10.1016/j.placenta.2019.01.003. PubMed PMID: 30712668.

11.   Faure-Bardon V, Magny JF, Parodi M, Couderc S, Garcia P, Maillotte AM, et al.

Sequelae of Congenital Cytomegalovirus Following Maternal Primary Infections Are Limited to

Those Acquired in the First Trimester of Pregnancy. Clin Infect Dis. 2019;69(9):1526-32. Epub

2019/01/01. doi: 10.1093/cid/ciy1128. PubMed PMID: 30596974.

12.   Fisher S, Genbacev O, Maidji E, Pereira L. Human cytomegalovirus infection of

placental cytotrophoblasts in vitro and in utero: implications for transmission and pathogenesis.

J Virol. 2000;74(15):6808-20. Epub 2000/07/11. doi: 10.1128/jvi.74.15.6808-6820.2000.

PubMed PMID: 10888620; PubMed Central PMCID: PMCPMC112198.

13.   Zydek M, Petitt M, Fang-Hoover J, Adler B, Kauvar LM, Pereira L, Tabata T. HCMV

infection of human trophoblast progenitor cells of the placenta is neutralized by a human

monoclonal antibody to glycoprotein B and not by antibodies to the pentamer complex. Viruses.

2014;6(3):1346-64. Epub 20140319. doi: 10.3390/v6031346. PubMed PMID: 24651029;

PubMed Central PMCID: PMCPMC3970154.

14.   Tabata T, Petitt M, Zydek M, Fang-Hoover J, Larocque N, Tsuge M, et al. Human

cytomegalovirus infection interferes with the maintenance and differentiation of trophoblast

progenitor cells of the human placenta. J Virol. 2015;89(9):5134-47. Epub 20150304. doi:

10.1128/JVI.03674-14. PubMed PMID: 25741001; PubMed Central PMCID: PMCPMC4403461.

15.　Halwachs-Baumann G, Wilders-Truschnig M, Desoye G, Hahn T, Kiesel L, Klingel K, et

al. Human trophoblast cells are permissive to the complete replicative cycle of human

cytomegalovirus. J Virol. 1998;72(9):7598-602. doi: 10.1128/JVI.72.9.7598-7602.1998. PubMed

PMID: 9696860; PubMed Central PMCID: PMCPMC110014.

16.　Pereira L, Tabata T, Petitt M, Fang-Hoover J. Congenital cytomegalovirus infection

undermines early development and functions of the human placenta. Placenta. 2017;59 Suppl

1:S8-S16. Epub 20170425. doi: 10.1016/j.placenta.2017.04.020. PubMed PMID: 28477968.

17.　Tabata T, Petitt M, Fang-Hoover J, Rivera J, Nozawa N, Shiboski S, et al.

Cytomegalovirus impairs cytotrophoblast-induced lymphangiogenesis and vascular remodeling

in an in vivo human placentation model. Am J Pathol. 2012;181(5):1540-59. Epub 20120907.

doi: 10.1016/j.ajpath.2012.08.003. PubMed PMID: 22959908; PubMed Central PMCID:

PMCPMC3483806.

18.　Rollman TB, Berkebile ZW, Okae H, Bardwell VJ, Gearhart MD, Bierle CJ. Human

Trophoblast Stem Cells Restrict Human Cytomegalovirus Replication. bioRxiv.

2023:2023.12.13.571456. Epub 20231214. doi: 10.1101/2023.12.13.571456. PubMed PMID:

38168202; PubMed Central PMCID: PMCPMC10760179.

19.　Okae H, Toh H, Sato T, Hiura H, Takahashi S, Shirane K, et al. Derivation of Human

Trophoblast Stem Cells. Cell Stem Cell. 2018;22(1):50-63 e6. Epub 20171214. doi:

10.1016/j.stem.2017.11.004. PubMed PMID: 29249463.

20.　Yang L, Semmes EC, Ovies C, Megli C, Permar S, Gilner JB, Coyne CB. Innate immune

signaling in trophoblast and decidua organoids defines differential antiviral defenses at the

maternal-fetal interface. Elife. 2022;11:e79794. Epub 20220817. doi: 10.7554/eLife.79794.

PubMed PMID: 35975985; PubMed Central PMCID: PMCPMC9470165.

21.   Goldstein JA, Gallagher K, Beck C, Kumar R, Gernand AD. Maternal-Fetal Inflammation

in the Placenta and the Developmental Origins of Health and Disease. Front Immunol.

2020;11:531543. Epub 20201113. doi: 10.3389/fimmu.2020.531543. PubMed PMID: 33281808;

PubMed Central PMCID: PMCPMC7691234.

22.   Scott GM, Chow SS, Craig ME, Pang CN, Hall B, Wilkins MR, et al. Cytomegalovirus

infection during pregnancy with maternofetal transmission induces a proinflammatory cytokine

bias in placenta and amniotic fluid. J Infect Dis. 2012;205(8):1305-10. Epub 20120301. doi:

10.1093/infdis/jis186. PubMed PMID: 22383678.

23.   Gervasi MT, Romero R, Bracalente G, Chaiworapongsa T, Erez O, Dong Z, et al. Viral

invasion of the amniotic cavity (VIAC) in the midtrimester of pregnancy. J Matern Fetal Neonatal

Med. 2012;25(10):2002-13. Epub 20120530. doi: 10.3109/14767058.2012.683899. PubMed

PMID: 22524157; PubMed Central PMCID: PMCPMC3498469.

24.   Bourgon N, Fitzgerald W, Aschard H, Magny JF, Guilleminot T, Stirnemann J, et al.

Cytokine Profiling of Amniotic Fluid from Congenital Cytomegalovirus Infection. Viruses.

2022;14(10). Epub 20220928. doi: 10.3390/v14102145. PubMed PMID: 36298700; PubMed

Central PMCID: PMCPMC9607316.

25.   Weisblum Y, Panet A, Zakay-Rones Z, Vitenshtein A, Haimov-Kochman R, Goldman-

Wohl D, et al. Human cytomegalovirus induces a distinct innate immune response in the

maternal-fetal interface. Virology. 2015;485:289-96. Epub 20150828. doi:

10.1016/j.virol.2015.06.023. PubMed PMID: 26318261.

26.   Weisblum Y, Oiknine-Djian E, Vorontsov OM, Haimov-Kochman R, Zakay-Rones Z, Meir

K, et al. Zika Virus Infects Early- and Midgestation Human Maternal Decidual Tissues, Inducing

Distinct Innate Tissue Responses in the Maternal-Fetal Interface. J Virol. 2017;91(4):e01905-16.

Epub 20170131. doi: 10.1128/JVI.01905-16. PubMed PMID: 27974560; PubMed Central

PMCID: PMCPMC5286880.

27.   Yockey LJ, Jurado KA, Arora N, Millet A, Rakib T, Milano KM, et al. Type I interferons

instigate fetal demise after Zika virus infection. Sci Immunol. 2018;3(19). Epub 2018/01/07. doi:

10.1126/sciimmunol.aao1680. PubMed PMID: 29305462; PubMed Central PMCID:

PMCPMC6049088.

28.   Rowe JH, Ertelt JM, Xin L, Way SS. Listeria monocytogenes cytoplasmic entry induces

fetal wastage by disrupting maternal Foxp3+ regulatory T cell-sustained fetal tolerance. PLoS

Pathog. 2012;8(8):e1002873. Epub 20120816. doi: 10.1371/journal.ppat.1002873. PubMed

PMID: 22916020; PubMed Central PMCID: PMCPMC3420962.

29.   Schleiss MR. Nonprimate models of congenital cytomegalovirus (CMV) infection: gaining

insight into pathogenesis and prevention of disease in newborns. ILAR J. 2006;47(1):65-72.

Epub 2006/01/05. doi: 10.1093/ilar.47.1.65. PubMed PMID: 16391432.

30.   Morrison JL, Botting KJ, Darby JRT, David AL, Dyson RM, Gatford KL, et al. Guinea pig

models for translation of the developmental origins of health and disease hypothesis into the

clinic. J Physiol. 2018;596(23):5535-69. Epub 20180530. doi: 10.1113/JP274948. PubMed

PMID: 29633280; PubMed Central PMCID: PMCPMC6265540.

31.   Hartley JW, Rowe WP, Huebner RJ. Serial propagation of the guinea pig salivary gland

virus in tissue culture. Proc Soc Exp Biol Med. 1957;96(2):281-5. Epub 1957/11/01. doi:

10.3181/00379727-96-23455. PubMed PMID: 13485080.

32.   Choi YC, Hsiung GD. Cytomegalovirus infection in guinea pigs. II. Transplacental and

horizontal transmission. J Infect Dis. 1978;138(2):197-202. Epub 1978/08/01. doi:

10.1093/infdis/138.2.197. PubMed PMID: 210239.

33.   Griffith BP, McCormick SR, Fong CK, Lavallee JT, Lucia HL, Goff E. The placenta as a

site of cytomegalovirus infection in guinea pigs. J Virol. 1985;55(2):402-9. Epub 1985/08/01. doi:

10.1128/JVI.55.2.402-409.1985. PubMed PMID: 2991565; PubMed Central PMCID:

PMCPMC254947.

34.   Auerbach MR, Yan D, Vij R, Hongo JA, Nakamura G, Vernes JM, et al. A neutralizing

anti-gH/gL monoclonal antibody is protective in the guinea pig model of congenital CMV

infection. PLoS Pathog. 2014;10(4):e1004060. Epub 20140410. doi:

10.1371/journal.ppat.1004060. PubMed PMID: 24722349; PubMed Central PMCID:

PMCPMC3983071.

35.   Griffith BP, Lucia HL, Hsiung GD. Brain and visceral involvement during congenital

cytomegalovirus infection of guinea pigs. Pediatr Res. 1982;16(6):455-9. Epub 1982/06/01. doi:

10.1203/00006450-198206000-00010. PubMed PMID: 6285263.

36.   Griffith BP, Hsiung GD. Cytomegalovirus infection in guinea pigs. IV. Maternal infection

at different stages of gestation. J Infect Dis. 1980;141(6):787-93. Epub 1980/06/01. doi:

10.1093/infdis/141.6.787. PubMed PMID: 6248603.

37.   Harrison CJ, Britt WJ, Chapman NM, Mullican J, Tracy S. Reduced congenital

cytomegalovirus (CMV) infection after maternal immunization with a guinea pig CMV

glycoprotein before gestational primary CMV infection in the guinea pig model. J Infect Dis.

1995;172(5):1212-20. Epub 1995/11/01. doi: 10.1093/infdis/172.5.1212. PubMed PMID:

7594656.

38.   Berkebile ZW, Putri DS, Abrahante JE, Seelig DM, Schleiss MR, Bierle CJ. The

Placental Response to Guinea Pig Cytomegalovirus Depends Upon the Timing of Maternal

Infection. Front Immunol. 2021;12:686415. Epub 20210615. doi: 10.3389/fimmu.2021.686415.

PubMed PMID: 34211475; PubMed Central PMCID: PMCPMC8239309.

39.   Coleman S, Choi KY, Root M, McGregor A. A Homolog Pentameric Complex Dictates

Viral Epithelial Tropism, Pathogenicity and Congenital Infection Rate in Guinea Pig

Cytomegalovirus. PLoS Pathog. 2016;12(7):e1005755. Epub 20160707. doi:

10.1371/journal.ppat.1005755. PubMed PMID: 27387220; PubMed Central PMCID:

PMCPMC4936736.

40.   Hashimoto K, Yamada S, Katano H, Fukuchi S, Sato Y, Kato M, et al. Effects of

immunization of pregnant guinea pigs with guinea pig cytomegalovirus glycoprotein B on viral

spread in the placenta. Vaccine. 2013;31(31):3199-205. Epub 20130515. doi:

10.1016/j.vaccine.2013.04.078. PubMed PMID: 23684839.

41.   Stahl PL, Salmen F, Vickovic S, Lundmark A, Navarro JF, Magnusson J, et al.

Visualization and analysis of gene expression in tissue sections by spatial transcriptomics.

Science. 2016;353(6294):78-82. Epub 2016/07/02. doi: 10.1126/science.aaf2403. PubMed

PMID: 27365449.

42.   Bierle CJ, Fernandez-Alarcon C, Hernandez-Alvarado N, Zabeli JC, Janus BC, Putri DS,

Schleiss MR. Assessing Zika virus replication and the development of Zika-specific antibodies

after a mid-gestation viral challenge in guinea pigs. PLoS One. 2017;12(11):e0187720. Epub

20171103. doi: 10.1371/journal.pone.0187720. PubMed PMID: 29099873; PubMed Central

PMCID: PMCPMC5669436.

43.   Putri DS, Berkebile ZW, Mustafa HJ, Fernandez-Alarcon C, Abrahante JE, Schleiss MR,

Bierle CJ. Cytomegalovirus infection elicits a conserved chemokine response from human and

guinea pig amnion cells. Virology. 2020;548:93-100. Epub 20200617. doi:

10.1016/j.virol.2020.06.005. PubMed PMID: 32838950; PubMed Central PMCID:

PMCPMC7447834.

44.   Depreux FF, Czech L, Whitlon DS. Sex Genotyping of Archival Fixed and

Immunolabeled Guinea Pig Cochleas. Sci Rep. 2018;8(1):5156. Epub 20180326. doi:

10.1038/s41598-018-23491-3. PubMed PMID: 29581456; PubMed Central PMCID:

PMCPMC5980087.

45.   Alam SM, Konno T, Rumi MA, Dong Y, Weiner CP, Soares MJ. Prolactin family of the

guinea pig, Cavia porcellus. Endocrinology. 2010;151(8):3918-28. Epub 20100609. doi:

10.1210/en.2010-0239. PubMed PMID: 20534723; PubMed Central PMCID:

PMCPMC2940522.

46.   Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J. g:Profiler: a web

server for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic

Acids Res. 2019;47(W1):W191-W8. Epub 2019/05/09. doi: 10.1093/nar/gkz369. PubMed PMID:

31066453; PubMed Central PMCID: PMCPMC6602461.

47.   Schleiss MR, Aronow BJ, Handwerger S. Cytomegalovirus infection of human

syncytiotrophoblast cells strongly interferes with expression of genes involved in placental

differentiation and tissue integrity. Pediatr Res. 2007;61(5 Pt 1):565-71. Epub 2007/04/07. doi:

10.1203/pdr.0b013e318045be6d. PubMed PMID: 17413859.

48.   Alfi O, Cohen M, Bar-On S, Hashimshony T, Levitt L, Raz Y, et al. Decidual-tissue-

resident memory T cells protect against nonprimary human cytomegalovirus infection at the

maternal-fetal interface. Cell Rep. 2024;43(2):113698. Epub 20240123. doi:

10.1016/j.celrep.2024.113698. PubMed PMID: 38265934.

49.   Lopez H, Benard M, Saint-Aubert E, Baron M, Martin H, Al Saati T, et al. Novel model of

placental tissue explants infected by cytomegalovirus reveals different permissiveness in early

and term placentae and inhibition of indoleamine 2,3-dioxygenase activity. Placenta.

2011;32(7):522-30. Epub 20110524. doi: 10.1016/j.placenta.2011.04.016. PubMed PMID:

21605903.

50.   Hemmings DG, Kilani R, Nykiforuk C, Preiksaitis J, Guilbert LJ. Permissive

cytomegalovirus infection of primary villous term and first trimester trophoblasts. J Virol.

1998;72(6):4970-9. Epub 1998/05/30. doi: 10.1128/JVI.72.6.4970-4979.1998. PubMed PMID:

9573266; PubMed Central PMCID: PMCPMC110059.

51.   Barrozo ER, Seferovic MD, Castro ECC, Major AM, Moorshead DN, Jochum MD, et al.

SARS-CoV-2 niches in human placenta revealed by spatial transcriptomics. Med.

2023;4(9):612-34 e4. Epub 20230708. doi: 10.1016/j.medj.2023.06.003. PubMed PMID:

37423216; PubMed Central PMCID: PMCPMC10527005.

52.   Schafer H, Burger R. Tools for cellular immunology and vaccine research the in the

guinea pig: monoclonal antibodies to cell surface antigens and cell lines. Vaccine.

2012;30(40):5804-11. Epub 20120720. doi: 10.1016/j.vaccine.2012.07.012. PubMed PMID:

22819989.

53.   Hilton HG, Rubinstein ND, Janki P, Ireland AT, Bernstein N, Fong NL, et al. Single-cell

transcriptomics of the naked mole-rat reveals unexpected features of mammalian immunity.

PLoS Biol. 2019;17(11):e3000528. Epub 20191121. doi: 10.1371/journal.pbio.3000528.

PubMed PMID: 31751331; PubMed Central PMCID: PMCPMC6894886.

54.   Moses L, Pachter L. Museum of spatial transcriptomics. Nat Methods. 2022;19(5):534-

46. Epub 20220310. doi: 10.1038/s41592-022-01409-2. PubMed PMID: 35273392.

55.   Arutyunyan A, Roberts K, Troule K, Wong FCK, Sheridan MA, Kats I, et al. Spatial

multiomics map of trophoblast development in early pregnancy. Nature. 2023;616(7955):143-

51. Epub 20230329. doi: 10.1038/s41586-023-05869-0. PubMed PMID: 36991123; PubMed

Central PMCID: PMCPMC10076224.

56.   Vento-Tormo R, Efremova M, Botting RA, Turco MY, Vento-Tormo M, Meyer KB, et al.

Single-cell reconstruction of the early maternal-fetal interface in humans. Nature.

2018;563(7731):347-53. Epub 20181114. doi: 10.1038/s41586-018-0698-6. PubMed PMID:

30429548; PubMed Central PMCID: PMCPMC7612850.

57.   Pique-Regi R, Romero R, Tarca AL, Sendler ED, Xu Y, Garcia-Flores V, et al. Single cell

transcriptional signatures of the human placenta in term and preterm parturition. Elife. 2019;8.

Epub 20191212. doi: 10.7554/eLife.52004. PubMed PMID: 31829938; PubMed Central PMCID:

PMCPMC6949028.

58.   Pique-Regi R, Romero R, Tarca AL, Luca F, Xu Y, Alazizi A, et al. Does the human

placenta express the canonical cell entry mediators for SARS-CoV-2? Elife. 2020;9:e58716.

Epub 20200714. doi: 10.7554/eLife.58716. PubMed PMID: 32662421; PubMed Central PMCID:

PMCPMC7367681.

59.   Marsh B, Blelloch R. Single nuclei RNA-seq of mouse placental labyrinth development.

Elife. 2020;9. Epub 20201103. doi: 10.7554/eLife.60266. PubMed PMID: 33141023; PubMed

Central PMCID: PMCPMC7669270.

60.   Pereira L, Maidji E, McDonagh S, Genbacev O, Fisher S. Human cytomegalovirus

transmission from the uterus to the placenta correlates with the presence of pathogenic bacteria

and maternal immunity. J Virol. 2003;77(24):13301-14. doi: 10.1128/jvi.77.24.13301-

13314.2003. PubMed PMID: 14645586; PubMed Central PMCID: PMCPMC296088.

61.   Gil-Sanchez A, Koletzko B, Larque E. Current understanding of placental fatty acid

transport. Curr Opin Clin Nutr Metab Care. 2012;15(3):265-72. Epub 2012/03/28. doi:

10.1097/MCO.0b013e3283523b6e. PubMed PMID: 22450774.

62.   Herrera E. Lipid metabolism in pregnancy and its consequences in the fetus and

newborn. Endocrine. 2002;19(1):43-55. Epub 2003/02/14. doi: 10.1385/ENDO:19:1:43. PubMed

PMID: 12583601.

63.   Ziaei S, Bonab KM, Kazemnejad A. Serum lipid levels at 28-32 weeks gestation and

hypertensive disorders. Hypertens Pregnancy. 2006;25(1):3-10. Epub 2006/04/15. doi:

10.1080/10641950500543756. PubMed PMID: 16613787.

64.   Ray JG, Diamond P, Singh G, Bell CM. Brief overview of maternal triglycerides as a risk

factor for pre-eclampsia. BJOG. 2006;113(4):379-86. Epub 2006/03/24. doi: 10.1111/j.1471-

0528.2006.00889.x. PubMed PMID: 16553649.

65.   Catov JM, Bodnar LM, Kip KE, Hubel C, Ness RB, Harger G, Roberts JM. Early

pregnancy lipid concentrations and spontaneous preterm birth. Am J Obstet Gynecol.

2007;197(6):610 e1-7. Epub 2007/12/07. doi: 10.1016/j.ajog.2007.04.024. PubMed PMID:

18060950.

66.   Lange PT, Lagunoff M, Tarakanova VL. Chewing the Fat: The Conserved Ability of DNA

Viruses to Hijack Cellular Lipid Metabolism. Viruses. 2019;11(2). Epub 20190129. doi:

10.3390/v11020119. PubMed PMID: 30699959; PubMed Central PMCID: PMCPMC6409581.

67.   Munger J, Bennett BD, Parikh A, Feng XJ, McArdle J, Rabitz HA, et al. Systems-level

metabolic flux profiling identifies fatty acid synthesis as a target for antiviral therapy. Nat

Biotechnol. 2008;26(10):1179-86. Epub 20080928. doi: 10.1038/nbt.1500. PubMed PMID:

18820684; PubMed Central PMCID: PMCPMC2825756.

68.   Abrahamsen LH, Clay MJ, Lyle JM, Zink JM, Fredrikson LJ, DeSiervo AJ, Jerkofsky MA.

The effects of cytomegalovirus infection on polar lipids and neutral lipids in cultured human

cells. Intervirology. 1996;39(4):223-9. Epub 1996/01/01. doi: 10.1159/000150521. PubMed

PMID: 9078462.

69.   Gudleski-O'Regan N, Greco TM, Cristea IM, Shenk T. Increased expression of LDL

receptor-related protein 1 during human cytomegalovirus infection reduces virion cholesterol

and infectivity. Cell Host Microbe. 2012;12(1):86-96. Epub 2012/07/24. doi:

10.1016/j.chom.2012.05.012. PubMed PMID: 22817990; PubMed Central PMCID:

PMCPMC3405916.

70.   Low H, Mukhamedova N, Cui HL, McSharry BP, Avdic S, Hoang A, et al.

Cytomegalovirus Restructures Lipid Rafts via a US28/CDC42-Mediated Pathway, Enhancing

Cholesterol Efflux from Host Cells. Cell Rep. 2016;16(1):186-200. Epub 20160616. doi:

10.1016/j.celrep.2016.05.070. PubMed PMID: 27320924; PubMed Central PMCID:

PMCPMC5389417.

71.   Chen Q, Gouilly J, Ferrat YJ, Espino A, Glaziou Q, Cartron G, et al. Metabolic

reprogramming by Zika virus provokes inflammation in human placenta. Nat Commun.

2020;11(1):2967. Epub 20200611. doi: 10.1038/s41467-020-16754-z. PubMed PMID:

32528049; PubMed Central PMCID: PMCPMC7290035.

72.   Maxwell AJ, You Y, Aldo PB, Zhang Y, Ding J, Mor G. The role of the immune system

during pregnancy: General concepts. In: Mor G, editor. Reproductive Immunology: Academic

Press; 2021. p. 1-21.

73. Gomez-Lopez N, Romero R. The nature of the immune response in microbial-associated

and sterile intraamniotic inflammation. In: Mor G, editor. Reproductive Immunology: Academic

Press; 2021. p. 207-37.

74. Bao M, Hofsink N, Plosch T. LPS versus Poly I:C model: comparison of long-term effects

of bacterial and viral maternal immune activation on the offspring. Am J Physiol Regul Integr

Comp Physiol. 2022;322(2):R99-R111. Epub 20211207. doi: 10.1152/ajpregu.00087.2021.

PubMed PMID: 34874190; PubMed Central PMCID: PMCPMC8782664.

75. Miner JJ, Cao B, Govero J, Smith AM, Fernandez E, Cabrera OH, et al. Zika Virus

Infection during Pregnancy in Mice Causes Placental Damage and Fetal Demise. Cell.

2016;165(5):1081-91. Epub 20160511. doi: 10.1016/j.cell.2016.05.008. PubMed PMID:

27180225; PubMed Central PMCID: PMCPMC4874881.

76. Buchrieser J, Degrelle SA, Couderc T, Nevers Q, Disson O, Manet C, et al. IFITM

proteins inhibit placental syncytiotrophoblast formation and promote fetal demise. Science.

2019;365(6449):176-80. Epub 2019/07/13. doi: 10.1126/science.aaw7733. PubMed PMID:

31296770.

77. Rice G, Patrick T, Parmar R, Taylor CF, Aeby A, Aicardi J, et al. Clinical and molecular

phenotype of Aicardi-Goutieres syndrome. Am J Hum Genet. 2007;81(4):713-25. Epub

20070904. doi: 10.1086/521373. PubMed PMID: 17846997; PubMed Central PMCID:

PMCPMC2227922.

78. Kim CJ, Romero R, Kusanovic JP, Yoo W, Dong Z, Topping V, et al. The frequency,

clinical significance, and pathological features of chronic chorioamnionitis: a lesion associated

with spontaneous preterm birth. Mod Pathol. 2010;23(7):1000-11. Epub 20100326. doi:

10.1038/modpathol.2010.73. PubMed PMID: 20348884; PubMed Central PMCID:

PMCPMC3096929.

79. Romero R, Chaemsaithong P, Chaiyasit N, Docheva N, Dong Z, Kim CJ, et al. CXCL10

and IL-6: Markers of two different forms of intra-amniotic inflammation in preterm labor. Am J

Reprod Immunol. 2017;78(1):e12685. Epub 20170519. doi: 10.1111/aji.12685. PubMed PMID:

28544362; PubMed Central PMCID: PMCPMC5488235.

80.   Maymon E, Romero R, Bhatti G, Chaemsaithong P, Gomez-Lopez N, Panaitescu B, et

al. Chronic inflammatory lesions of the placenta are associated with an up-regulation of amniotic

fluid CXCR3: A marker of allograft rejection. J Perinat Med. 2018;46(2):123-37. Epub

2017/08/23. doi: 10.1515/jpm-2017-0042. PubMed PMID: 28829757; PubMed Central PMCID:

PMCPMC5797487.

81.   Hirsch AJ, Roberts VHJ, Grigsby PL, Haese N, Schabel MC, Wang X, et al. Zika virus

infection in pregnant rhesus macaques causes placental dysfunction and immunopathology. Nat

Commun. 2018;9(1):263. Epub 20180117. doi: 10.1038/s41467-017-02499-9. PubMed PMID:

29343712; PubMed Central PMCID: PMCPMC5772047.

82.   Manuel T, Mostrom M, Crooks CM, Davalos A, Barfield R, Scheef E, et al. A nonhuman

primate model mirrors human congenital cytomegalovirus infection and reveals a spectrum of

vertical transmission outcomes. bioRxiv. 2025:2025.01.16.633406. Epub 20250121. doi:

10.1101/2025.01.16.633406. PubMed PMID: 40492197; PubMed Central PMCID:

PMCPMC12148051.

83.   Adkins RM, Walton AH, Honeycutt RL. Higher-level systematics of rodents and

divergence time estimates based on two congruent nuclear genes. Mol Phylogenet Evol.

2003;26(3):409-20. Epub 2003/03/20. doi: 10.1016/s1055-7903(02)00304-4. PubMed PMID:

12644400.

84.   Roberts RM, Green JA, Schulz LC. The evolution of the placenta. Reproduction.

2016;152(5):R179-89. Epub 20160802. doi: 10.1530/REP-16-0325. PubMed PMID: 27486265;

PubMed Central PMCID: PMCPMC5033709.

85.   Crumpler MM, Choi KY, McVoy MA, Schleiss MR. A live guinea pig cytomegalovirus

vaccine deleted of three putative immune evasion genes is highly attenuated but remains

immunogenic in a vaccine/challenge model of congenital cytomegalovirus infection. Vaccine.

2009;27(31):4209-18. Epub 20090503. doi: 10.1016/j.vaccine.2009.04.036. PubMed PMID:

19389443; PubMed Central PMCID: PMCPMC3539795.

86. Leviton MP, Lacayo JC, Choi KY, Hernandez-Alvarado N, Wey A, Schleiss MR. An

attenuated cytomegalovirus vaccine with a deletion of a viral chemokine gene is protective

against congenital CMV transmission in a guinea pig model. Clin Dev Immunol.

2013;2013:906948. Epub 20130820. doi: 10.1155/2013/906948. PubMed PMID: 24000289;

PubMed Central PMCID: PMCPMC3755440.

87. Pouliot N, Maghni K, Blanchette F, Cironi L, Sirois P, Stankova J, Rola-Pleszczynski M.

Natural killer and lectin-dependent cytotoxic activities of Kurloff cells: target cell selectivity,

conjugate formation, and Ca++ dependency. Inflammation. 1996;20(6):647-71. Epub

1996/12/01. doi: 10.1007/BF01488802. PubMed PMID: 8979153.

88. Marshall AH, Swettenham KV, Vernon-Roberts B, Revell PA. Studies on the function of

the Kurloff cell. Int Arch Allergy Appl Immunol. 1971;40(1):137-52. doi: 10.1159/000230401.

PubMed PMID: 4100635.

89. Revell PA, Vernon-Roberts B, Gray A. The distribution and ultrastructure of the Kurloff

cell in the guinea-pig. J Anat. 1971;109(Pt 2):187-99. Epub 1971/07/01. PubMed PMID:

5558229; PubMed Central PMCID: PMCPMC1270999.

90. Yang D, Tamburro K, Dittmer D, Cui X, McVoy MA, Hernandez-Alvarado N, Schleiss

MR. Complete genome sequence of pathogenic Guinea pig cytomegalovirus from salivary gland

homogenates of infected animals. Genome Announc. 2013;1(2):e0005413. Epub 20130314.

doi: 10.1128/genomeA.00054-13. PubMed PMID: 23516193; PubMed Central PMCID:

PMCPMC3622957.

91. Martinez-Pacheco M, Tenorio M, Almonte L, Fajardo V, Godinez A, Fernandez D, et al.

Expression Evolution of Ancestral XY Gametologs across All Major Groups of Placental

Mammals. Genome Biol Evol. 2020;12(11):2015-28. Epub 2020/08/14. doi:

10.1093/gbe/evaa173. PubMed PMID: 32790864; PubMed Central PMCID: PMCPMC7674692.

92.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq

data using regularized negative binomial regression. Genome Biol. 2019;20(1):296. Epub

20191223. doi: 10.1186/s13059-019-1874-1. PubMed PMID: 31870423; PubMed Central

PMCID: PMCPMC6927181.

93.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.

Comprehensive Integration of Single-Cell Data. Cell. 2019;177(7):1888-902 e21. Epub

20190606. doi: 10.1016/j.cell.2019.05.031. PubMed PMID: 31178118; PubMed Central PMCID:

PMCPMC6687398.

1026

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

1. Chemokine activity (GO:0008009)
2. Immune system process (GO:0002376)
3. Leukocyte activation (GO:0045321)

Figure 6