Escape from X inactivation drives sex differences and female trait variation

Carrie Zhu[1,2], Liaoyi Xu[1,2], Arbel Harpak[1,2,+]

[1]Department of Integrative Biology, University of Texas at Austin, Austin, TX, USA

[2]Department of Population Health, University of Texas at Austin, Austin, TX, USA

[+]To whom correspondence should be addressed: arbelharpak@utexas.edu

Abstract

X chromosome inactivation (XCI) partially balances gene dosage between sexes, yet expression from the inactive X (Xi) is variable across genes. In this study, we investigate whether gene-level Xi expression predicts transcriptional and phenotypic consequences of X-linked variation. We find that Xi expression levels are a strong linear predictor of female-male expression differences, suggesting that other compensatory or regulatory mechanisms play a more minor role in sex differences in X-linked gene expression. Among females, we identify three traits—lymphocyte percentage, HbA1c, and schizophrenia—for which higher Xi expression correlates with the strength of evidence for dominance effects. We hypothesize that an underappreciated mechanism could generate dominance effects of X-linked variants on a trait—specifically when the variant influences skew in X inactivation. This work establishes Xi expression as essential for understanding sex differences and the female-specific genetic basis of disease.

Introduction

X chromosome inactivation (XCI) in females is traditionally viewed as an equalizer, bringing X-linked gene dosage to be on par with dosage in males by silencing one X chromosome[1–3]. However, many genes "escape" this silencing and are expressed from the inactive X (Xi)[4,5]. Roughly 15–30% of X-linked genes show some degree of escape, with some escaping consistently across tissues and individuals, and others showing context-dependent escape[6,7]. The extent of escape can vary continuously, from low-level to equal expression between Xi and the active X (Xa). Understanding the consequences of expression from Xi is critical for interpreting both sex differences and individual variation among females[8–10].

An immediate expectation is that levels of Xi expression would impact dosage balance between the sexes. Genes in the pseudoautosomal regions (PAR) of the X chromosome, which recombine with the Y chromosome, are not subject to XCI and thus maintain expression from both sex chromosomes[11,12]. Yet, even PAR genes show unequal expression between the sexes, with higher expression in males reported in several tissues. This demonstrates that expression from Xi does not typically reach the full levels of the active X chromosome nor that of the Y chromosome[6,7,13]. In contrast, many non-PAR (NPX) genes have no Y homologs and exhibit higher expression in females due to unopposed Xi expression[6].

Beyond sex differences, Xi expression may also influence phenotypic variation within females[14,15]. Heterozygote loss of function mutations provide compelling case studies. For example, such mutations in DDX3X increase the risks of developmental delay and disability[16], and in KDM6A the risk of Kabuki syndrome[17–19]. These observations imply that Xi gene dosage can be impactful for development and disease. Further, Agarwal et al. recently showed that in a group of 17 NPX genes (those with Y homologs), the fitness cost of heterozygote loss of function mutations is higher than in any other group of genes genome-wide[20].

Here, we test whether gene-level Xi expression predicts transcriptional and phenotypic consequences of X-linked variation. We find that, across genes, Xi expression levels quantitatively predict both (i) the magnitude of female-male expression differences and (ii) the strength of genetic associations for disease risk and physiological traits in females. We also suggest a novel hypothesis for non-additive effects of X-linked variants on a trait, specifically when the variant also influences a skew in X inactivation. Testing for dominance effects in X-linked variants, we find weak evidence for these effects being mediated by genetically encoded skews at individual loci. However, integrating signals across Xi expressed genes, we identify three

54   traits—lymphocyte percentage, HbA1c, and schizophrenia—for which the strength of Xi

55   expression predicts the significance of dominance effects.

Results

Results

We used RNA-seq data from the Genotype-Tissue Expression (GTEx) v8 project[21] for 643 X-

linked genes across 14 tissue types in 635 XY and 312 XX adult individuals. While recognizing

that sex chromosome karyotype does not always align with biological sex, we henceforth label

XX individuals as female and XY individuals as male. We quantify Xi expression using allele-

specific expression data obtained from a single GTEx female donor whose X inactivation is non-

random across cells and tissues ("skewed XCI"), i.e. the inactivated chromosome is the one

inherited from the same parent in nearly all cells[6]. Importantly, our reliance on data from this single

GTEx donor implies that, throughout this work, we do not consider variation in XCI and escape

among females. Instead, our focus is on variation in Xi expression across genes and these genes'

contribution to sex differences and within-female variation.

Xi expression predicts sex differences in gene expression. Across genes in the non-

pseudoautosomal (NPX) region, expression from Xi is strongly correlated with sex differences in

median expression level at both the individual tissue level (e.g., liver and skin in **Fig. 1A**) and

when examining the per-gene medians across tissues (Pearson r = 0.77; p < 2.2e-16; **Fig. 1B**).

In contrast, in the pseudoautosomal region (PAR), 11 out of 11 genes had higher expression in

males (by 19% on average), and sex differences were not correlated with Xi expression (Pearson

p=0.95; **Fig. 1B**). These results are in line with prior work by Tukiainen et al.[6] and Tomofuji et al.[22]

who also report greater female expression in escape genes in the NPX region, and greater male

expression in the PAR1 region. This is, to our knowledge, the first illustration of the linear,

continuous and tight relationship between Xi expression and subsequent sex differences, which

suggests that other compensatory or regulatory mechanisms play a more minor role in sex

differences in X-linked gene expression.

Many Xi-expressed genes have a highly consistent expression levels across tissues—18 out of

43 (42%) per a previous report[6]. In such genes, e.g. in ZRSR2 and UBA1, we also see little

variance in sex differences in gene expression (**Fig. S1**). Genes such as ARSD and FHL1, which

have more variable Xi expression, tend to also exhibit greater variance in sex differences in

expression. In general, variance across tissues in sex differences is proportional to variance

across tissues in Xi expression (**Fig. S1**). One outlier to this trend is GYG2, which is involved in

glycogenesis[23]. GYG2 shows disproportionately larger variance in sex differences across tissues

and is in fact expressed from Xi only in the liver, pancreas, and stomach (between 27% and

Figure 1: Xi expression in a single female predicts sex differences in gene expression. (A) The x-

axis shows the ratio of expression from Xi and total expression in the tissue (liver on top, suprapubic skin

on the bottom) estimated in a single GTEx female donor. The y-axis displays the difference between median

female and male transcripts per million (TPM), standardized by the median male TPM for each gene. For

example, y=0 confers to equal medians and 0.5 confers to the median expression of the gene among

females being 50% higher than that of males. Gene expression data for the respective tissues was obtained

from GTEx v8. We estimate the Pearson correlation statistics independently for the non-PAR X (orange)

and PAR1 (gray) regions. Genes with homologs on the Y chromosome are indicated as dark red triangles.

(B) The x-axis shows the median proportion of expression from Xi across tissues. The y-axis shows the

median, across tissues, of standardized female-male differences.

111% higher median expression in females, across these tissues; **Figs. 1B, S2**). In other tissues

with no expression from Xi, sex differences still varied extensively. For example, in coronary

artery tissue, median expression was 83% higher in females while in the esophagus' muscularis

layer it was only 20% higher. Such tissue-specific sex differences may reflect additional regulatory

factors, such as epigenetic modifications or tissue-specific chromatin states, that are not

necessarily unique to Xi expression[24].

Beyond the impact of Xi expression on sex differences, it may also contribute uniquely to female

variation. We therefore also inquired into consequences for phenotypic variation among females.

Once again, we do so by examining, across genes, the relationship between Xi expression

(inferred from a single individual) and measures of gene-level contribution to variation. We

performed X-wide association studies (XWAS) for 13 diseases and continuous physiological

traits. We chose these traits based on either having relatively high SNP heritabilities or

documented phenotypic sex differences for greater statistical power[25–28]. We tested for standard,

additive effects. We discuss the relationship between additive effects and Xi expression later in

the section "**Xi expression predicts contribution to trait variation among females**".

A new hypothesis for dominance effects of X-linked variants. In our XWAS, we also tested

whether the heterozygote effect deviates from the average of homozygote effects (a "dominance-

deviation effect"). Hypothetically, dominance may be observed on the X chromosome for the

same reasons as they may exist in autosomes. However, dominance effects are thought to be

negligible among common variants in autosomes[29,30]. Should XCI itself, or perhaps escape from

XCI, alter our expectations for X-linked variants? We distinguish three biological scenarios for a

variant in an escape gene that yield different expectations for the nature of X-linked variants'

effects.

First, we consider a biallelic SNP with an allelic effect β when the effect allele is expressed from

Xa. If the locus completely escapes X inactivation, the expected effect of a heterozygote is equal

to the average of the two homozygotes ($\alpha = 1$ in **Table 1-i**) just like in autosomes. When

inactivation is complete and random, one of the two alleles' effects would be zeroed at random

and so the genotypic effect is halved ($\alpha = 0$ in **Table 1-i**). For some intermediate degree of Xi

expression (ranging from 0 to 1), the expected effect in homozygotes is $(1 + \alpha)\beta$, and in

heterozygotes it is $\frac{(1+\alpha)}{2}\beta$ (**Table 1-i**). In conclusion, the presence of a variant in an Xi-expressed

locus, in and within itself, does not lead us to expect dominance-deviation effects.

Second, we consider an additional effect of the variant—modulating Xi expression levels—when

it is present on the inactive chromosome. Still, in this scenario, the variant does not influence X

inactivation choice. While the effect size now depends on the allele-specific level of Xi expression,

the effect remains additive (**Table 1-ii**).

Case	Expected Effect of Genotype	Dominance Deviation Effect?
(i) Inactivation and expression from Xi (at level α) are independent of allele.	AA: 0 AT: $\frac{(1+\alpha)}{2}\beta$ TT: $(1+\alpha)\beta$	No
(ii) Inactivation is independent, but expression from Xi depends on allele (at level α_A for allele A on Xi and level α_T for allele T on Xi)	AA: 0 AT: $\frac{(1+\alpha_T)}{2}\beta$ TT: $(1+\alpha_T)\beta$	No
(iii) Inactivation is allele-dependent: In heterozygotes, an X chromosome with the T allele has probability δ of being the active X. (Expression from Xi is fixed at level α.)	AA: 0 AT: $\delta\beta + (1-\delta)\alpha\beta$ TT: $(1+\alpha)\beta$	Yes

Table 1. When an X-linked allele affects both a trait and the likelihood of inactivation, we expect it to have a dominance deviation effect on the trait.

A third case we consider is when the variant influences which parental X is inactivated. Heterozygotes will then exhibit skewing of XCI with unequal probabilities of each of the alleles being on Xa and, as a result, having the more pronounced effect. Therefore, under this scenario, we expect dominance effects in our XWAS (**Table 1-iii**). In other words, the allelic effect on the trait in this case is mediated by a pleiotropic effect on XCI.

Dominance effects may also arise for mechanisms that are not unique to the X or Xi. We emphasize, however, that the mechanism in case (iii) hinges on genetically encoded biases in XCI. While there is ample literature on skewed XCI in blood and other tissues[31–34], the majority of skewing is thought arise from stochastic processes, clonal expansion, or aging[35–37], rather than from heritable genetic variation. Nonetheless, some evidence supports the possibility of heritable components to skewed XCI, especially in extreme or familial cases[33,38–41]. These findings may also motivate the testing for dominance effects as a way to identify potential loci that influence XCI.

Across all 13 traits, we found only 17 SNPs with genome-wide significant (p<5e-8) dominance effects. All of the SNPs were in the NPX, but Xi expression data was not available for nearby

genes. Notably, the nearest genes also had no Y homologs. These effects were suggestive of complete dominance (**Fig. S3**). To illustrate, **Fig. 2** shows the mean phenotype across genotypes for three SNPs with significant dominance effects on HbA1c (**Fig. 2A-C**) and one with a significant effect on reported migraine cases (**Fig. 2D**). We caveat these observations in noting that sample sizes for the minor allele homozygotes were small, and the allelic effect of the SNPs among males (per a simple comparison of mean phenotypes conferring to hemizygote genotypes; **Fig. S3**) was often discordant. One intriguing case is a SNP close to ZFP92 (**Fig. 2A**), which encodes a zinc finger protein thought to regulate DNA transcription and transposable element silencing[42]. Therefore, a linked variant may influence X-inactivation, consistent with Case (iii) in **Table 1**.

Xi expression predicts contribution to trait variation among females. We next asked whether higher Xi expression suggests a greater role in explaining trait variation among females. We did so by testing for a correlation between Xi expression level (using the median across tissues) and the strength of a gene's association with the trait. To assign a strength of association, we annotated each gene with the minimal p-value association among SNPs for which the gene is the closest gene[43]. We estimated this correlation across 13 traits and two types of effects, additive and dominance deviation. For height, the significance of additive effects and expression from Xi were highly correlated (Pearson r = 0.45, p = 0.029; **Fig S4**).

Another intriguing correlation is observed between Xi expression levels and significance of dominance-deviation effects on percentage of lymphocytes in leukocytes (Pearson r = 0.52, p = 0.012; **Fig. S4**). If XCI skewness is heritable and more polygenic than previously appreciated, this correlation would fall in line with Case (iii) in **Table 1**. Overall, for both types of effects, we find that genes with higher tissue-averaged Xi expression tend to have stronger GWAS associations near them. For additive effects, there was a positive Pearson correlation for 11/13 traits (1 significant at p < 0.05). For dominance-deviation effects, this was true for 12/13 traits (1 significant at p < 0.05; **Fig. S4**).

We investigated this relationship further, in a tissue-specific manner, namely by examining correlations of XWAS significance with tissue-specific Xi expression. In some tissues, most traits we examined showed significant correlation between Xi expression and strength of XWAS association (p < 0.05). This includes stomach (7 traits for additive GWAS; 10 traits for dominance deviation), thyroid (4 additive, 7 dominance deviation), as well as in lymphoblastoid cell lines (LCLs) (4 additive, 8 dominance deviation) (**Fig. 3C**). These tissues were also highly correlated

Figure 2: Examples of NPX SNPs with significant dominance effects. Shown are four SNPs that

exhibited genome-wide significant dominance deviation effects in an XWAS for HbA1c (**A-C**) and reported

case of migraine (**D**). The y-values confer to the mean phenotype value (±SE), or proportion, in (D), for

females in the White British subset of the UKB with the respective genotype. Sample sizes (n) are printed

below each genotype.

with each other in terms of Xi expression (see dendrogram on the left hand side of **Fig. 3C**).

However, we were unable to discern whether this points to pleiotropic contributions from Xi

expressed genes. Other tissues such as esophagus mucosa, kidney cortex, suprapubic skin, and

lung, had no significant correlations. This may result from low variation in Xi expression among

genes or a lesser impact of Xi-expressed genes on variation in these traits. Since Xi expression

is generally similar across tissues (about 43% of genes with nonzero Xi expression were

characterized as having similar expression levels across tissues by Tukiainen et al.[6], the latter is

more likely.

For some traits, Xi expression and significance of allelic effects were significantly correlated

across multiple tissues for only one type of effect. Interestingly, lymphocyte percentage in

leukocytes, HbA1c, and schizophrenia were only significant for the dominance deviation model

across 5, 4, and 4 tissues respectively **(Fig. 3A, 3B)**. We therefore hypothesize that X inactivation

skew may more commonly mediate the effects of Xi expression in these tissues. Recognizing the

multiple hypothesis burden of testing 13 traits, 14 tissues and 2 effects types, we note that only 4

Figure 3: Correlation between Xi expression and gene-trait association across tissues. We estimate, for each tissue, trait and effect type (additive or dominance deviation) the Pearson correlation between the proportion of Xi expression out of total (Xi + Xa) expression and the maximum $-\log_{10}$(p-value) among variants overlapping the gene. Only genes with non-zero expression (blue points) were considered. **(A)** HbA1c in whole blood and dominance deviation effects. **(B)** Thyrotoxicosis in thyroid and dominance deviation effects. **(C)** Correlations across tissue, trait and effect type combinations. Only significant correlations are displayed, with blue hues conferring to the significance levels. The bottom-left triangles of each block show Pearson p-values for Xi expression and additive effect $-\log_{10}$(p-value), and the top-right triangles show the same for dominance-deviation effects. We cluster tissues by pairwise correlations of Xi expression, as shown by the dendrogram on the left. The length of each branch is inversely proportional to the correlation between tissues in Xi expression, For example, Xi expression is more correlated between stomach and lymphoblastoid cell lines (LCL) than it is between thyroid and kidney cortex.

tissue-trait-effect combinations are significant at an FDR significance level of 0.05 (bolded

triangles in **Fig. 3C**).

Discussion

Our study demonstrates that expression from Xi is a major, quantifiable contributor to

transcriptional sex differences and trait variation among females. First, we find that Xi expression,

as measured in a single individual with skewed inactivation, robustly predicts sex differences in

expression across non-pseudoautosomal (NPX) genes and tissues. While previous work has

documented sex differences in expression of escape genes[6], our findings go further by showing

that this relationship extends across the full range of Xi expression, even among genes not

formally categorized as 'escapees', supporting the view that escape from X inactivation acts along

a continuum, rather than as a binary on/off phenomenon (see relevant discussion in [7,44]). The

strong correlation (Pearson $r = 0.77$) between Xi expression and higher female expression in NPX

regions supports a model by which sex differences arise largely from unequal dosage, rather than

from differential regulation of the active X.

We also propose a simple model to delineate an X-chromosome specific expectation for

dominance-deviation effects. Namely, dominance-deviation effects will arise when allele-specific

skewing affects which haplotype is inactivated. By distinguishing between effects on expression

given that an allele is Xi-linked and effects on XCI, we provide a clear rationale for interpreting

dominance effects in the context of XCI. We identify several traits—lymphocyte percentage,

HbA1c, and schizophrenia—where dominance-deviation signals may point to genetically encoded

XCI bias. These traits are particularly compelling because they align with known biology of XCI

skew or tissue-specific gene regulation. Lymphocyte percentage reflects hematopoietic

processes, where XCI skew is well-documented due to clonal expansion and somatic

selection[33,37]. Some studies also suggest a heritable component to skew[33,38], which may explain

why dominance effects on the percentage of lymphocytes in leukocytes is strongly correlated with

Xi expression. Such skew may expose or silence functional alleles, helping explain the greater

incidence of autoimmune disorders in females. HbA1c, a marker of glucose control, could reflect

Xi expression effects in metabolic tissues like the liver or pancreas—tissues where genes like

GYG2 and PNPLA4 exhibit high Xi expression. Schizophrenia is a neurodevelopmental disorder

with known sex differences in presentation, as well as previous observations of an XCI skew in

affected females[45]. Together, these examples suggest that dominance effects on these traits may

reveal genetically encoded variation in XCI.

Finally, the correlation between Xi expression and GWAS association strength tends to cluster by
tissue type, suggesting that Xi expression acts through shared regulatory environments. Tissues
such as stomach, thyroid, and LCLs show the strongest and most consistent correlations across
traits. Given that a large proportion of Xi-expressed genes are similarly expressed across tissues,
these results could imply that Xi-driven trait effects may often arise from genes involved in
fundamental cellular processes such as metabolism or structure.

In sum, our study shows that escape from X inactivation—treated as a quantitative, gene-level
trait—has widespread implications for sex differences and trait variation in humans. Our findings
underscore the importance of Xi expression in shaping sex differences and influencing genetic
architecture of physiological and disease traits.

Methods

GTEx expression data. To estimate gene expression levels, we used bulk tissue RNA
sequencing data from the Genotype-Tissue Expression (GTEx) v8 project[46] for 727 genes across
14 tissue types in 635 males and 312 females. Gene expression levels are measured in
transcripts per million (TPM). The list of genes on the X chromosome was obtained from San
Roman et al.[7] which included 712 genes on the non-PAR (NPX) region and 15 genes on the
pseudo-autosomal 1 (PAR1) region of X chromosome. We removed sex-specific tissues and
genes with less than 10 individuals with TPM greater than 1. We also removed tissues, including
brain tissues, without corresponding Xi expression data described in "**Use of allele-specific
expression data to measure degree of escape from XCI**". Any gene-tissue pairs with 0 TPM
were removed.

Use of allele-specific expression data to measure degree of escape from XCI. We measured
the amount of expression from the inactive X chromosome using data obtained from Tukiainen et
al.[6] They use allele-specific expression data from a single female GTEx-UPIC donor who exhibits
a near complete degree of X inactivation skew, in which the same parental chromosome is
inactivated across nearly all cells. For each gene, one ASE site with the highest coverage in the
largest number of tissues is chosen. The allele with the lower combined expression across tissues
is labeled the inactive X (Xi) allele. XIST is the sole exception, with the Xi allele having higher
expression levels across tissues. We then quantify the degree of escape using the ratio of Xi

expression to total (inactive + active) X expression. In total, the combination of GTEx expression

data and allele-specific expression data is available for 190 NPX and 11 PAR1 genes.

UK Biobank. We used the UK Biobank, a large biomedical database of phenotypic and genotypic

information of around half a million UK participants[47,48]. In this study, we focused on female

participants, with female labeling based on the presence of an XX karyotype. Quality checks of

sample characteristics included removing individuals with sex chromosome aneuploidy, or a self-

reported sex other than female. We excluded all related individuals 3rd degree or closer, as listed

in data field 22020. We removed individuals who had withdrawn from the UK Biobank at any time

before we performed the analysis of UKB data (6/19/2024). Furthermore, we limited participants

to those who self-identified as White and as British and who were also clustered in the genetic

principal component space. After filtering, we had a total sample of 181012 female individuals.

UKB Genotype Data. We used X-chromosome, bi-allelic SNPs with INFO scores greater than

0.8 (Resource 1967 in UKB). Following a similar quality check method as used by Neale Lab[49],

we then filtered SNPs with call rate > 0.95, Hardy-Weinberg equilibrium test p-value larger than

1e-10 and minor allele frequency larger than 1e-3. Filtering was completed using *plink 2.0*

alpha[50,51].

Phenotype Data. In this study, we analyzed 13 phenotypes including 6 continuous and 7 binary

disease phenotypes. The continuous traits were chosen for their relatively high SNP heritability

estimates as provided by Neale Lab[49]. Disease phenotypes were classified using ICD10 coding

(data field 41270 in UK Biobank) and chosen based on known sex-specific differences in disease

incidence, course or prognosis.

XWAS. We conducted female-specific association analysis for the X chromosome using *plink 2.0*

alpha[50,51]. We included birth year and the first ten principal components (data field 22009 in UK

Biobank) as covariates. We applied two different models, additive and dominance-deviation, using

the --genotypic modifier.

Gene annotation. We used *Ensembl Variant Effect Predictor*[43] to annotate highly associated

SNPs ($p < 0.05$) obtained from XWAS. We found the closest gene within 5000 bp up- and down-

stream each SNP. For **Fig. 3**, we created a list of SNPs for each gene with available Xi expression

data using the *Ensembl REST API*[52] overlap endpoint, which retrieves variants that lie within the

gene region. We chose the SNP with the lowest p-value to represent the gene.

Summarizing the correlation distance between tissues in Xi expression using a tree. To

create the dendrogram shown on the left in **Fig. 3C**, we first estimated the pairwise Pearson

correlation of proportion of Xi expression levels for 14 tissues using the data from Tukiainen et

al[6]. We treated the inverse of these correlations as a distance measure between each pair of

tissues. Using the resulting pairwise distance matrix, we performed hierarchical clustering using

the function *hclust()* in *R version 4.3.1* using default settings. The length of (horizontal) branches

shown in **Fig. 3C** confers to the distance between each pair or cluster of tissues.

Acknowledgements

We thank Ipsita Agarwal, Andrés Bendesky, Jared Cole, Mark Kirkpatrick and Molly Przeworski

for helpful discussions and comments on the manuscript. This work was funded by NIH grant

R35GM151108 and a Pew Biomedical Scholarship to A.H. This study was conducted using the

UK Biobank resource under application 92741, as approved by the University of Texas at Austin

institutional review board (study protocol 00003287). The authors acknowledge the Texas

Advanced Computing Center (TACC) at The University of Texas at Austin for providing

computational resources that have contributed to the research results reported in this manuscript.

Literature Cited

1. Lyon, M. F. Gene Action in the X-chromosome of the Mouse (Mus musculus L.). *Nature* 190, 372–373 (1961).

2. Disteche, C. M. Dosage Compensation of the Sex Chromosomes. *Annu Rev Genet* 46, 537–560 (2012).

3. Disteche, C. M., Filippova, G. N. & Tsuchiya, K. D. Escape from X inactivation. *Cytogenet Genome Res* 99, 36–43 (2002).

4. Posynick, B. J. & Brown, C. J. Escape From X-Chromosome Inactivation: An Evolutionary Perspective. *Front Cell Dev Biol* 7, (2019).

5. Balaton, B. P., Dixon-McDougall, T., Peeters, S. B. & Brown, C. J. The eXceptional nature of the X chromosome. *Hum Mol Genet* 27, R242–R249 (2018).

6. Tukiainen, T. *et al.* Landscape of X chromosome inactivation across human tissues. *Nature* 550, 244–248 (2017).

7. San Roman, A. K. *et al.* The human inactive X chromosome modulates expression of the active X chromosome. *Cell Genomics* 3, 100259 (2023).

8. Khramtsova, E. A. *et al.* Quality control and analytic best practices for testing genetic models of sex differences in large populations. *Cell* 186, 2044–2061 (2023).

9. Fass, S. B. *et al.* Relationship between sex biases in gene expression and sex biases in autism and Alzheimer's disease. *Biol Sex Differ* 15, 47 (2024).

10. Ungar, R. A. *et al.* Transcriptomic signatures of rare variant impacts across sex and the X chromosome. *Human Genetics and Genomics Advances* 6, 100463 (2025).

11. Graves, J. The origin and evolution of the pseudoautosomal regions of human sex chromosomes. *Hum Mol Genet* 7, 1991–1996 (1998).

12. Blaschke, R. J. & Rappold, G. The pseudoautosomal regions, SHOX and disease. *Curr Opin Genet Dev* 16, 233–239 (2006).

13. Johnston, C. M. *et al.* Large-Scale Population Study of Human Cell Lines Indicates that Dosage Compensation Is Virtually Complete. *PLoS Genet* 4, e9 (2008).

14. Fu, Y., Kenttämies, A., Ruotsalainen, S., Pirinen, M. & Tukiainen, T. Role of X chromosome and dosage-compensation mechanisms in complex trait genetics. *The American Journal of Human Genetics* 112, 1330–1343 (2025).

15. Sidorenko, J. *et al.* The effect of X-linked dosage compensation on complex trait variation. *Nat Commun* 10, 3009 (2019).

16. Tang, L. *et al.* Prospective and detailed behavioral phenotyping in DDX3X syndrome. *Mol Autism* 12, 36 (2021).

17. Van Laarhoven, P. M. *et al.* Kabuki syndrome genes *KMT2D* and *KDM6A* : functional analyses demonstrate critical roles in craniofacial, heart and brain development. *Hum Mol Genet* 24, 4443–4453 (2015).

18. Lederer, D. *et al.* Deletion of KDM6A, a Histone Demethylase Interacting with MLL2, in Three Patients with Kabuki Syndrome. *The American Journal of Human Genetics* 90, 119–124 (2012).

19. Cheon, C. K. *et al.* Identification of KMT2D and KDM6A mutations by exome sequencing in Korean patients with Kabuki syndrome. *J Hum Genet* 59, 321–325 (2014).

20. Agarwal, I., Fuller, Z. L., Myers, S. R. & Przeworski, M. Relating pathogenic loss-of-function mutations in humans to their evolutionary fitness costs. *Elife* 12, (2023).

21.   Lonsdale, J. *et al.* The Genotype-Tissue Expression (GTEx) project. *Nat Genet* 45, 580–
585 (2013).

22.   Tomofuji, Y. *et al.* Quantification of escape from X chromosome inactivation with single-
cell omics data reveals heterogeneity across cell types and tissues. *Cell Genomics* 4,
100625 (2024).

23.   Mu, J. & Roach, P. J. Characterization of Human Glycogenin-2, a Self-glucosylating
Initiator of Liver Glycogen Metabolism. *Journal of Biological Chemistry* 273, 34850–34856
(1998).

24.   Qadir, M. M. F. *et al.* Single cell regulatory architecture of human pancreatic islets
suggests sex differences in β cell function and the pathogenesis of type 2 diabetes.
Preprint at https://doi.org/10.1101/2024.04.11.589096 (2024).

25.   Nyholt, D. *et al.* Evidence for an X-linked genetic component in familial typical migraine.
Hum Mol Genet 7, 459–463 (1998).

26.   Bauer, M., Glenn, T., Pilhatsch, M., Pfennig, A. & Whybrow, P. C. Gender differences in
thyroid system function: relevance to bipolar disorder and its treatment. *Bipolar Disord* 16,
58–71 (2014).

27.   Miquel, C. H., Faz-Lopez, B. & Guéry, J. C. Influence of X chromosome in sex-biased
autoimmune diseases. *J Autoimmun* 137, 102992 (2023).

28.   van Vollenhoven, R. F. Sex differences in rheumatoid arthritis: more than meets the eye...
BMC Med 7, 12 (2009).

29.   Visscher, P. M. *et al.* Genome Partitioning of Genetic Variation for Height from 11,214
Sibling Pairs. *The American Journal of Human Genetics* 81, 1104–1110 (2007).

30.   Palmer, D. S. *et al.* Analysis of genetic dominance in the UK Biobank. *Science (1979)*
379, 1341–1348 (2023).

31.   Minks, J., Robinson, W. P. & Brown, C. J. A skewed view of X chromosome inactivation.
Journal of Clinical Investigation 118, 20–23 (2008).

32.   Zito, A. *et al.* Escape from X-inactivation in twins exhibits intra- and inter-individual
variability across tissues and is heritable. *PLoS Genet* 19, e1010556 (2023).

33.   Bolduc, V. *et al.* No evidence that skewing of X chromosome inactivation patterns is
transmitted to offspring in humans. *Journal of Clinical Investigation* 118, 333–341 (2008).

34.   Sharp, A., Robinson, D. & Jacobs, P. Age- and tissue-specific variation of X chromosome
inactivation ratios in normal women. *Hum Genet* 107, 343–349 (2000).

35.   Newall, A. E. T. Primary non-random X inactivation associated with disruption of Xist
promoter regulation. *Hum Mol Genet* 10, 581–589 (2001).

36.   Busque, L. *et al.* Nonrandom X-inactivation patterns in normal females: lyonization ratios
vary with age. *Blood* 88, 59–65 (1996).

37.   Werner, J. M., Ballouz, S., Hover, J. & Gillis, J. Variability of cross-tissue X-chromosome
inactivation characterizes timing of human embryonic lineage specification events. *Dev*
Cell 57, 1995-2008.e5 (2022).

38.   Amos-Landgraf, J. M. *et al.* X Chromosome–Inactivation Patterns of 1,005 Phenotypically
Unaffected Females. *The American Journal of Human Genetics* 79, 493–499 (2006).

39.   Thorvaldsen, J. L., Krapp, C., Willard, H. F. & Bartolomei, M. S. Nonrandom X
Chromosome Inactivation Is Influenced by Multiple Regions on the Murine X
Chromosome. *Genetics* 192, 1095–1107 (2012).

40.   Chadwick, L. H. & Willard, H. F. Genetic and parent-of-origin influences on X
chromosome choice in Xce heterozygous mice. *Mammalian Genome* 16, 691–699
(2005).

41.   Chadwick, L. H., Pertz, L. M., Broman, K. W., Bartolomei, M. S. & Willard, H. F. Genetic
Control of X Chromosome Inactivation in Mice: Definition of the *Xce* Candidate Interval.
Genetics 173, 2103–2110 (2006).

42.   Osipovich, A. B. *et al.* ZFP92, a KRAB domain zinc finger protein enriched in pancreatic
islets, binds to B1/Alu SINE transposable elements and regulates retroelements and
genes. *PLoS Genet* 19, e1010729 (2023).

43.   McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol* 17, 122 (2016).

44.   San Roman, A. K. *et al.* The human Y and inactive X chromosomes similarly modulate
autosomal gene expression. *Cell Genomics* 4, 100462 (2024).

45.   Zhang, X. *et al.* A new sex-specific underlying mechanism for female schizophrenia:
accelerated skewed X chromosome inactivation. *Biol Sex Differ* 11, 39 (2020).

46.   Aguet, F. *et al.* The GTEx Consortium atlas of genetic regulatory effects across human
tissues. *Science (1979)* 369, 1318–1330 (2020).

47.   Sudlow, C. *et al.* UK Biobank: An Open Access Resource for Identifying the Causes of a
Wide Range of Complex Diseases of Middle and Old Age. *PLoS Med* 12, e1001779
(2015).

48.   Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data.
Nature 562, 203–209 (2018).

49.   Abbot, L. *et al.* GWAS Results. http://www.nealelab.is/uk-biobank/ (2017).

50.   Purcell, S. & Chang, C. PLINK 2.00 alpha. Preprint at www.cog-genomics.org/plink/2.0/
(2020).

51.   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* 4, 7 (2015).

52.   Yates, A. *et al.* The Ensembl REST API: Ensembl Data for Any Language. *Bioinformatics*
31, 143–145 (2015).

469
470