CCB79 is a primate-specific cilium initiation factor essential to maintain neural progenitor diversity in developing brain tissue.

Dhanasekaran Rathinam[1], Vaibhav Jadhav[1], Seema Wasim[2], Omkar Suhas Vinchure[1], Dina Elkahwagy[1], Tejas Borkar[1], Aruljothi Mariappan[1], Nazlican Altinisik[1], Samhan Alsolami[3, 4], Mo Li[3, 4], Johannes Ptok[5, 6], Ziliang Zhao[7,8], Christian Eggeling[7,8], Muhammad Tariq[9], Maria Giovanna Riparbelli[10], Giuliano Callaini[10], Peter Nürnberg[11], Elke Gabriel[12], and Jay Gopalakrishnan[1*]

[1]Institute of Human Genetics, University Hospital, Friedrich-Schiller-Universität Jena, 07740 Jena, Germany
[2]Institute of Human Genetics, Ulm University, Ulm University Medical Center, Ulm, Germany
[3]Bioscience Program, Biological and Environmental Science and Engineering Division (BESE), King Abdullah University of Science and Technology (KAUST), Thuwal, 23955, Saudi Arabia
[4]KAUST Center of Excellent for Smart Health (KCSH), Thuwal, 23955, Saudi Arabia
[5]Institute of Medical Microbiology and Hospital Hygiene, Heinrich Heine University Düsseldorf, Düsseldorf, Germany
[6]Institute of Virology, University Hospital Düsseldorf, Heinrich Heine University Düsseldorf, Düsseldorf, Germany
[7]Leibniz Institute of Photonic Technologies, Jena, Germany; member of the Leibniz Centre for Photonics in Infection Research (LPI), Jena, Germany
[8]Institute of Applied Optics and Biophysics, Friedrich-Schiller-Universität Jena, Jena, Germany
[9]National Institute for Biotechnology and Genetic Engineering College, Pakistan Institute of Engineering and Applied Sciences (NIBGE-C, PIEAS), Faisalabad, Pakistan
[10] Department of Life Sciences and Medical Biotechnology, University of Siena, Siena 53100, Italy
[11]ologne Center for Genomics (CCG), University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne, Germany
[12]Institute of Human Genetics, University Hospital, Heinrich-Heine-Universität, 40225 Düsseldorf, Germany

[*] **Correspondence:** jay.gopalakrishnan@med.uni-jena.de

Summary

Identifying the genes that regulate the accurate diversity of neural progenitor cells (NPCs) helps understand the mechanisms of human neocortex expansion. In primate brains, an additional intermediate progenitor layer, the outer subventricular zone (oSVZ) facilitates the expansion of the neocortex. Here, we identify an uncharacterized gene, KIAA0408, and show that its expression is enriched in intermediate progenitors. Removing KIAA0408 in human-induced pluripotent stem cell (iPSC)-derived brain organoids results in an impaired cortical organization characterized by abnormal cell fate and patterning defects, including the depletion of intermediate and ventral progenitors. Molecularly, KIAA0408 is a 79-kilodalton centriolar distal appendage protein (DAP) that controls cilium biogenesis (hereafter CCB79). CCB79 forms a complex with other DAP components and specifically localizes in the DAP at the onset of ciliogenesis, and its absence blocks ciliogenesis. Mechanistically, progenitors in 3D brain tissues are unable to form cilia, which induces aberrant hedgehog signaling and causes premature differentiation. Finally, human CCB79, rather than the mouse ortholog, rescues cilia defects, suggesting that CCB79 has undergone rapid evolution from rodents to primates to fine-tune ciliogenesis for proper brain development.

Introduction

Primate evolution is characterized by a significant expansion of the forebrain, resulting in a disproportionately large neocortex[1-5]. Emerging evidence indicates that these evolutionary differences originate at the NPC level—the fundamental unit of the developing brain[2,4-8]. NPC diversity is carefully coordinated to promote the proper development of the cerebral cortex. In the developing brain, NPCs are located in distinct layers of the ventricular zone (VZ) at the apical surface. Here, apical radial glia divide symmetrically to produce dorsal and ventral intermediate progenitors, which migrate to the subventricular zone (SVZ) and can proliferate further before symmetrically differentiating into migrating neurons across all cortical layers[9-12]. In species with more prominent, gyrencephalic brains (e.g., primates), an additional outer subventricular zone (oSVZ) contains a large number of outer radial glial cells and intermediate progenitors, a highly neurogenic NPC type that greatly promotes neocortex expansion[13-16].

Conversely, rodents, which typically have smooth-surfaced (lissencephalic) brains, have fewer outer radial glial cells, intermediate progenitors, and an indistinct oSVZ[13,16]. The surface area and thickness of the cerebral cortex in primates are much greater than in rodents due to the increased diversity of NPCs and neurons[1-5]. This wide range of NPC types and their developmental niches shows how the balance between self-renewing divisions and differentiation can fundamentally influence brain growth and function. Although this NPC diversity is critical for neocortex expansion, the molecular basis of NPC fate determination and diversity in the developing human brain remains unknown.

Primary cilia extend from the surface of most vertebrate cells, originating from the mother centriole, which acts as a basal body that docks to the plasma membrane[17]. The docking process is mediated

by a specialized, pinwheel-like structure called the distal appendages (DAPs) at the end of the mother centriole[18,19]. DAPs are essential for initiating ciliogenesis, and their molecular composition and biogenesis mechanisms are only beginning to emerge [20-22]. Primary cilia mediate several growth signaling pathways, including Sonic Hedgehog (SHH), which is vital for brain development[23,24]. Mutations in cilia-specific genes lead to ciliopathies, which are characterized by diverse phenotypes affecting multiple organs and are often accompanied by neurodevelopmental disorders [25,26]. Studies have shown that specific ciliary dysfunctions are linked to neurodevelopmental disorders, known as "non-classical ciliopathies," emphasizing the vital role of cilia in human brain development [17,27-29]. For instance, SHH signaling is crucial for controlling dorsal and ventral patterning during telencephalon development, and disruptions in this pathway can impair neural tube closure [30-32]. Mutations in SCLT1, a component of DAPs, are connected to neuroretinal degeneration[28]. Importantly, many genes related to basal body-cilia are enriched in the developing brain, and their regulated dynamics contribute to driving the lateral growth of the cerebral cortex [33,34]. This suggests that ciliary functions tailored to specific contexts, i.e., cell type-specific functions, may have evolved specifically to support brain development. This led us to search for unknown molecules that regulate ciliogenesis in the developing human brain.

In this study, we identify a previously uncharacterized gene, KIAA0408, and demonstrate that it encodes a 79-kilodalton (kDa) protein essential for regulating cilium formation, which we refer to as CCB79. CCB79 is mainly expressed in intermediate progenitors and both excitatory and inhibitory neurons located in the subventricular (SV) and outer subventricular (oSV) regions of the developing brain. Molecularly, CCB79 is a centriolar distal appendage protein (DAP) that is dynamically regulated during the cell cycle, appearing only at the start of ciliogenesis. Using proximity labeling and mass spectrometry, we show that CCB79 forms a complex with other DAP components. CRISPR/CAS9-mediated deletion of CCB79 disrupts the DAP and hinders cilium formation. Functional studies in genome-engineered brain organoids, combined with comparative single-cell (scRNA) and whole-transcriptome analyses, revealed abnormal ciliogenesis and misregulated SHH signaling, leading to altered NPC diversity and defects in brain tissue organization. We find that intermediate and ventral progenitors are particularly vulnerable when CCB79 is absent. Notably, the human gene can rescue cilia defects, while the mouse ortholog cannot. This suggests that, although CCB79 is a vertebrate gene, the primate variant has rapidly evolved to regulate specific aspects of ciliogenesis that are crucial for cortical expansion in primates.

Discovery of CCB79 as a controller of cilium biogenesis

To identify novel regulators of NPC diversity, we analyzed single-cell RNA sequencing (scRNA-seq) data from early-stage brain organoids, as described in Figures 5-6[35-37]. Uniform Manifold Approximation and Projection (UMAP) revealed six significant cell clusters, including telencephalic neural progenitors, dorsal and ventral progenitors, and excitatory and inhibitory neurons **(Figure 1A)**. Since intermediate progenitors are relatively abundant in gyrencephalic species with a high

neurogenic capacity for neocortex expansion [13-15], we calculated the differentially expressed genes in intermediate progenitor-containing clusters (C5 and C6 of SVZ and oSVZ region) compared to the telencephalic progenitor cluster (C2 of apical ventricular region) **(Figure 1B)**. We identified a group of genes that were significantly upregulated**.** We could assign known functions to all of these genes, except for the KIAA0408 mRNA **(Supplementary Table 1).** Computing the normalized expression of KIAA0408 across various human tissues revealed that its expression is enriched in fetal brain tissues, specifically in early neuronal cell types and at different stages of fetal development[38] **(Figure S1A-C).** This is corroborated by scRNA data derived from our brain organoids **(Figure 1C)**. Notably, in our UMAP, the expression of KIAA0408 coincides with an enriched expression of EOMES, INSM1, NKX2-2, GAD2, and SLC17A6, markers of intermediate progenitors and excitatory and inhibitory neurons emerging from dorsal and ventral progenitors, respectively **(Figure 1D and S1D)**. A pseudotime trajectory analysis confirms this, indicating that our brain organoids harbor cell lineages that follow sequential stages of brain development, including dorsal and ventral patterning events **(Figure 1D)**.

KIAA0408 is an uncharacterized gene initially identified as a novel human cDNA clone enriched in brain tissue, located on chromosome 6orf174[39,40]. It is considered a complex locus because it merges with another gene, MTCL3. Although their coding regions do not overlap, a read-through RNA product—likely a non-coding RNA predicted to be a candidate for nonsense-mediated mRNA decay—is produced[41]. KIAA0408 has six exons that encode a protein containing coiled-coil domains **(Figure 1E)**. To determine its subcellular localization, we transiently expressed the human KIAA0408 cDNA with a green fluorescent protein reporter tag at its N-terminus in human induced pluripotent stem cells (iPSCs). We observed the gene product as a distinct dot in each cell. To test whether these dots are basal bodies associated with primary cilia, we immunostained cilia using ARL13B antibodies and found that KIAA0408 localizes at the ciliary base **(Figure S1E)**. To label the endogenous protein, we generated polyclonal rat antibodies against the N-terminal peptide of KIAA0408 **(Figure 1E).** In these experiments, we used NPCs differentiated from iPSCs. Anti-KIAA0408 antibodies labeled KIAA0408 as a single, distinct dot at the distal end of the mother centriole, filling the space between the mother centriole (defined by γ-tubulin) and the cilium (defined by ARL13B) **(Figure 1F)**. We further immunostained the NPCs with antibodies against glutamylated tubulin GT335 and CP110 to confirm this localization pattern. GT335 antibodies label the mother and daughter centrioles as well as the cilium. CP110, a centriolar protein, specifically labels the daughter centriole when the mother centriole serves as a template for cilium assembly[42,43]. This analysis reaffirmed the endogenous localization of KIAA0408, showing it labels the distal end of the mother centriole and links it to the cilium **(Figure 1F-G).**

To validate the KIAA0408 antibody and determine the gene's function, we knocked out the KIAA0408 gene in iPSCs using the CRISPR/CAS9 method using two guide RNAs targeting exon 3 [44] **(Figure 1H).** Isolation of pure clones and Sanger sequencing of the PCR-amplified region revealed that the gene editing resulted in a 35-nucleotide deletion between the two guide RNA binding sites in the

targeted genomic region. We also observed the introduction of a stop codon within 22 bases of the deleted region, which stops translation after 90 amino acids **(Figure S2A)**. To detect the protein in Western blots, we raised polyclonal rabbit antibodies against the exon 3 peptides of KIAA0408 **(Figure 1E),** which recognized the protein bands only in unedited control iPSCs but not in knockout cells **(Figure 1I).** On the other hand, N-terminal antibodies did not show any immunoreactivity in KIAA0408-KO cells **(Figure 1J)**. If detected, there was only a faint signal, suggesting either reduced protein recruitment or that the residual signaling was due to the presence of the first 90 intact amino acids, which cover the epitope region for the antibody. As a result, unlike unedited cells, most KIAA0408-KO cells lacked primary cilia. When detected, the cilia were bulgy, short, thin, and long, indicating they are aberrant **(Figure 1J and S2B).** Ultrastructural investigation using transmission electron microscopy (EM) revealed that mother centrioles contained appendages in control cells that dock to the plasma membrane, form ciliary pockets, and emanate axonemal microtubules, and assemble cilia. In contrast, KIAA0408-KO cells lacked appendages in centrioles, forming a bulgy vesicle-like ciliary pocket, and failed to assemble structurally normal cilia **(Figure 1K)**. Finally, while the number of centrosomes remains unchanged in KIAA0408-KO cells, the frequency of ciliated cells and CCB79's recruitment to mother centrioles is significantly reduced compared to controls **(Figure 1L and S2C).** These results indicate that KIAA0408 localizes to the distal end of the mother centrioles and is critical for initiating ciliogenesis. For clarity, we refer to KIAA0408 as CCB79, designating it as a **C**ontroller of **C**ilium **B**iogenesis with a molecular weight of 79 kDa.

CCB79 is a cell cycle-regulated centriolar distal appendage protein

To map the precise localization of CCB79 within the mother centriole relative to centriolar appendage proteins, we used structured illumination microscopy (SIM) on NPCs. Compared to GT335 and CP110-specific staining (magenta), CCB79 (green) formed a distinct ring distal to the mother centriole at the base of the cilium **(Figure 2Ai-ii).** We then co-stained with ODF2 (magenta), which marks the subdistal appendages of the mother centriole [21,45]. While the top view showed that the CCB79 ring is clearly distal to ODF2 signals, the tilted view revealed that OFD2 forms a hollow cylinder surrounded by the CCB79 ring **(Figure 2Aiii).** To further confirm CCB79's position, we co-stained with CEP164 (magenta), a distal appendage protein. CEP164 formed a ring co-localizing with the CCB79 ring. Notably, the CCB79 ring displayed distinct periodic densities, suggesting they are distal appendages of the mother centriole **(Figure 2Aiv).** Finally, using super-resolution stimulated emission depletion (STED) microscopy, we visualized nine distinct densities of FBF1 (magenta), another distal appendage protein, which co-localized with CCB79 **(Figure 2Av-vi).** Importantly, imaging several centrioles showed that the diameter of CCB79's ring overlaps with the average diameter of CEP164 and FBF1 rings but not the OFD2 ring **(Figure 2B)**. These results indicate that CCB79 is a component of the centriolar distal appendage structure at the base of the cilium **(Figure 2C-D).**

The primary cilium is a dynamic organelle that assembles and disassembles, closely linked to the cell cycle[46,47]. It assembles when the cell exits mitosis and disassembles upon re-entry into the cycle. Our expanded analysis of CCB79's localization across different cell cycle stages revealed a single CCB79 dot in the G0/G1 phase, when cells build a cilium. Cells in G2, anaphase, and telophase lacked CCB79. Interestingly, CCB79 reappeared during late cytokinesis as a single dot in one of the dividing cells that begins to form a cilium **(Figure 2E).** This pattern shows that the mother centriole is preparing to assemble a cilium in G1 by recruiting CCB79.

Appendage proteins specify mother centrioles competent to template cilium biogenesis [19,20]. However, most appendage proteins not only localize to the mother centrioles at the onset of ciliogenesis but are present throughout the cell cycle. Comparing the localization behaviors of selected appendage proteins, such as CEP164 and FBF1, to CCB79 revealed that CCB79 intensity peaks at G0/G1, the stage when cells assemble cilia and rapidly declines as cells proceed to S, G2, and M phase, where cells disassemble cilia **(Figure 2F and S3A)**.

To further confirm that CCB79 appears only at the beginning of cilium assembly, we synchronized hTERT-immortalized retinal pigment epithelial cells (RPE-1) using serum starvation to reach G0 and serum stimulation to transition to G1-S. After 72 hours of serum starvation, cells assembled cilia with CCB79 prominently localized at the distal end of the mother centriole as a single dot. During subsequent hours of serum addition, CCB79 staining diminished, indicating cilium disassembly. CP110 is localized solely at the daughter centrioles at similar levels **(Figure S3B)**. We also examined CPAP, a direct interactor of CP110, in these assays. CPAP is a cilium disassembly factor whose levels gradually increased as cells disassembled the cilium[29,48] **(Figure S3C-D)**. These experiments show that CCB79, a cilium initiation factor, appears mainly in cells at the onset of cilium assembly.

CCB79 is a primate-specific cilium initiation factor

Since centriolar appendage proteins are essential regulators of cilium biogenesis, we conducted a phylogenetic analysis of centriolar DAPs with CCB79 to examine their conservation across different species. To achieve this, we employed clustering analysis to infer evolutionary models and identify gene expression patterns across various species[49]. The results showed that the CCB79 gene first appeared in frogs and then continued to evolve in higher vertebrates and terrestrial animals. Interestingly, the emergence of CCB79 expression coincided with other known DAP molecules such as TTBK2, CEP164, FBF1, CEP83, CEP89, C2CD3, and SCLT1 **(Figure 3A).** This finding prompted us to analyze its amino acid sequence. The basic pairwise alignment of sequences from different species relative to humans revealed significant changes in the CCB79 protein during evolution. For example, a mouse protein, which shares 55% identity with humans, rapidly evolved into a marmoset protein with 94% identity to the human protein within approximately 48 million years. Evolutionary pressure then slowed from marmosets to humans through macaques, bonobos, and chimpanzees over the past 43 million years **(Figure 3A)**. Additionally, multiple sequence alignment of CCB79 across various species uncovered notable insertions, substitutions, and deletions in their amino acid

sequences. Notably, primates, starting with Old World monkeys, gained a 43-amino-acid region in exon 5, which is absent in rodents **(Figure S4A).**

We therefore focused our analysis on exon 5. Alignment of the human CCB79 gene with its mouse counterpart shows that the human gene has gained an extension of 131 nucleotides at its C-terminus **(Figure S4B, green extension)**. This suggests that the transcript may have arisen due to small variations at the splice sites caused by point mutations. Analyzing the potential splice site revealed that there is a donor splice site in exon 5 of the mouse gene (we call this mSD5), which is suppressed in humans by a mutation—specifically, a nucleotide exchange (+3A>C), a point mutation three nucleotides downstream of the 5' splice site **(Figure S4B).** This is further supported by a decrease in the Hbond score (HBS), which measures complementarity to U1 snRNA of the spliceosome complex. Alternatively, this suppressed site is compensated by a novel splice site gained through another mutation further downstream at position +1 (SD5: +1T>G) in the human gene. This results in the exonization of intron 5 in the mouse transcript. HEXplorer plots in Figure S3B highlight the changes in the splicing environment caused by the mutations in human SD5 and mSD5. In summary, compared to the mouse gene, the human CCB79 gained 131 nucleotides at position 127,446,277 in the 5th exon, located on chromosome 6, leading to a new exonization and the addition of 43 amino acids to the translated protein.

To predict how the gain of 43 aa could alter the structure and folding characterization of the protein in primates, we utilized the AlphaFold protein structure prediction tool. This analysis revealed a gain of alpha-helices in primates, indicating that this alpha-helix can be functionally traced back to a common ancestor of Old-World monkeys but is absent in rodents **(Figure 3B).** Therefore, with the gain of a 43 aa containing alpha-helix, it is plausible that the CCB79 is a primate-specific transcript essential for ciliogenesis in primates and is functionally different from the rodent gene. Thus, although the CCB79 is a vertebrate gene, the human or primate transcript has undergone significant evolutionary pressure to acquire primate-specific functions **(Figure S4C-D)**.

To test this experimentally, we cloned the mouse ortholog and examined whether it could rescue the ciliogenesis defects of CCB79-KO iPSCs. First, we conducted our experiments in wild-type iPSCs. While the human gene CCB79 effectively targeted the ciliary base, the mouse gene did not do so efficiently. Furthermore, the mouse gene inhibited the cells from forming cilia **(Figure 3C)**. Accordingly, while the human CCB79 fully restored ciliogenesis in CCB79-KO iPSCs, the mouse gene neither targeted the ciliary base effectively nor rescued the loss of cilia. These experiments reveal functional differences between human and mouse transcripts, and CCB79 is a primate-specific factor involved in cilium initiation **(Figure 3D).** Figure 3E summarizes the findings. Obtaining stable clones of iPSCs expressing CCB79 was impossible, as overexpression did not produce healthy lines. Therefore, we expressed the gene in a doxycycline-inducible manner and analyzed a mixed population of cells.

CCB79 removal is sufficient to trigger NPCs to undergo premature differentiation and abolish cilia initiation.

We conducted a series of experiments to characterize the loss of function of CCB79 and its role in cilium initiation and NPC maintenance using iPSC-derived NPCs. While both variants (control and CCB79-KO) did not show a significant difference in EdU incorporation, unlike controls, CCB79-KO NPCs spontaneously differentiated into neurons at the expense of self-renewing NPCs. Notably, these experiments were performed under conditions that do not promote differentiation, indicating that CCB79 is essential for maintaining NPCs in their self-renewal state **(Figure S5A).**

Next, we aimed to dissect the underlying mechanisms of defective ciliogenesis. To achieve this, we conducted DAP recruitment assays across these variants. CCB79-KO NPCs revealed the following findings in comparison to healthy controls **(Figure S5B)**. *i)* Impaired CCB79 recruitment, meaning CCB79-KO NPCs recruited either no or significantly reduced levels of CCB79 (green, CCB79, panels B-F); *ii)* Aberrant cilia, wherein cilia were either absent or presented as a dot or bulging short structure (magenta, GT335 or ARL13B, panels B-F); *iii)* CP110 was never removed but remained on both daughter and mother centrioles (red, CP110, panel B); *iv)* Recruitment of γTubulin and DAPs CEP164 and FBF1 were unaffected (red, panels C, D, and E); and *v)* Strikingly, CCB79-KO NPCs failed to recruit TTBK2 (red, panel F). Panels on the right quantify the findings. In summary, these results indicate that CCB79 is a critical factor in cilium initiation, and its loss of function specifically inhibits TTBK2 recruitment, thereby impairing ciliogenesis. TTBK2 is an essential kinase that orchestrates cilia growth, and its absence abolishes cilium initiation[31]. Panel G presents a hypothetical topology of CCB79 as part of the DAP complex, initiating a cilium, and Panel H summarizes the findings.

CCB79 removal downregulates telencephalic progenitor genes, ciliopathy genes, and perturbs Hedgehog signaling in 3D human brain organoids

To gain mechanistic insights into how CCB79 can maintain NPC diversity in 3D brain tissues, we employed unbiased mRNA transcriptomics at both bulk and single-cell levels (scRNA) on 14-day-old brain organoids cultured for 14 days in spinner flasks (see Methods). We generated these organoids from control and CCB79-KO iPSCs using our previously described method [35-37], as detailed in Figure 6. Briefly, we directly induced the differentiation of iPSCs into neural epithelium. Early-stage brain organoids generated using this method expressed a diverse range of cell types, including telencephalic neural progenitors, as well as dorsal and ventral progenitors, and excitatory and inhibitory neurons. These organoids mirror the sequential stages of telencephalon development, exhibiting both dorsal and ventral patterning events, and could model developmental disorders such as microcephaly **(Figure 1A, C-D).**

The CCB79-KO organoids showed less growth compared to age-matched healthy controls, indicating developmental issues **(Figure S6A).** Thin sectioning and immunostaining for SOX2-positive progenitors, TUJ1-positive neurons, and ARL13-positive primary cilia revealed that control

organoids had well-organized, expanded ventricular zones (VZ) rich in SOX2-positive progenitors, ARL13B-primary cilia in their apical lumen of the VZ, and a thin layer of TUJ-positive primitive cortical plate at the basal side, mirroring the cytoarchitecture of developing brain tissue. In contrast, the CCB79-KO organoids showed disrupted cytoarchitecture, unrecognizable VZs, and barely visible cilia in their lumen, with a noticeable reduction in SOX2 intensities but an increase in TUJ1-positive neurons throughout the tissue, indicating early progenitor differentiation and resulting NPC loss **(Figure S6B).**

To examine the molecular changes caused by CCB79 removal, we conducted a differential gene expression (DGE) analysis comparing control and CCB79-KO brain organoids. We identified a down-regulation of pan-telencephalic genes (FOXG1), dorsal telencephalic progenitor genes specifying newly born cortical projection neurons (NEUROD6), ventral telencephalon progenitor genes (GSX2 and DLX family) crucial for the development of forebrain GABAergic interneurons, genes relevant to intermediate progenitors and early-born cortical (TBR1), inhibitory (GAD2 and GABRA1), excitatory (SLC6A17), and inter (Calbindin) neurons. Gene ontology ranking aligned with the DEG analysis **(Figure S6C).**

We also observed a decrease in the expression of known ciliopathy genes, which supports the finding that CCB79 loss prevents cilium formation **(Figure S6D).** Interestingly, GPR161 is among the most downregulated genes. GPR161 functions as a negative regulator of Sonic Hedgehog (SHH) signaling, and its loss in mice results in increased SHH activity in the neural tube[50,51]. Similarly, increased SHH signaling in cortical organoids caused by the loss of INPP5E, a ciliary gene, results in ventralization[52]. Therefore, we examined SHH target genes (PTCH1 and GLI1), ligands (DHH, SHH, and IHH), and receptors (PTCH1 and SMO), and found their levels elevated, indicating that SHH signaling is disrupted in organoid tissues after CCB79 removal **(Figure S6E).** To confirm the increase in SHH components and activity, we calculated a weighted SHH pathway activation score and combined it with the normalized expression of canonical SHH pathway genes[53]. This analysis showed a significant increase in SHH activity in CCB79-KO organoids **(Figure S6F)**, driven by higher levels of key SHH signaling molecules. Notably, SHH pathway scores were inversely related to the expression of forebrain patterning transcription factors, including FOXG1, LHX6, and TBR1, indicating a coordinated disruption of SHH signaling and neurodevelopmental programs, consistent with the known role of primary cilia in brain development **(Figure S6G)**[27,31,51,52]. Collectively, these findings suggest that CCB79 loss causes derepression of the SHH pathway and impairs forebrain gene expression, aligning with previously described ciliopathy phenotypes[54]. These changes in gene expression and ectopic SHH signaling could influence cell diversity in 3D brain tissue.

CCB79 removal derails progenitor fate determination, disrupts developmental patterning, and changes cell diversity in 3D human brain organoids.

To analyze changes in cell diversity, we examined the scRNA transcriptomes of approximately 18,000 control and 8,900 CCB79-KO cells isolated from 14-day-old brain organoids, tracking

developmental changes caused by the removal of CCB79. We assigned the sequenced cells to neural tube (WNT8B, RSPO2), neural crest (PAX3, COLEC12), progenitors of telencephalic (SOX2, FOXG1) and non-telencephalic regions (PAX3, HOXB2), proliferative (MKI67, AURKB), and immature neurons from both telencephalic (STMN, DLX5, GAD1) and non-telencephalic origins (STMN2, LHX1). We also identified oligodendrocyte progenitors (PLP1, SOX10) and microglia-like cells (AIF1) **(Figure 4A and S7-S8).** By evaluating similarities and differences, we found abnormal levels of oligodendrocyte precursors and neural crest lineages (clusters C2 and C9), as well as a reduction in telencephalic progenitors and immature neurons derived from ventral progenitors (clusters C3 and C6). The unusual appearance of oligodendrocyte precursors may be linked to increased SHH signaling in ventral progenitors, which is known to induce premature specification of oligodendrocyte precursors at the expense of inhibitory neurons. This aligns with the effects of cilia-mediated SHH enhancement observed in cortical organoids and the chick ventral spinal cord[52,55]. We then quantified the differential expression of SHH genes across our cell clusters and, notably, identified elevated levels of SHH signaling components in specific cell types within cluster 9, which contains oligodendrocyte progenitors, in CCB79-KO organoids (**Figure S9)**.

Since CCB79 is enriched in SVZ and oSVZ progenitors, as well as excitatory and inhibitory neurons derived from both dorsal and ventral progenitors, we aimed to map the changes in cell diversity in these regions precisely. First, we performed a detailed analysis of clusters C3, C4, C5, C6, and C7 **(Figures 4B and S10)**. Next, we calculated pseudotime to project all cortical lineage cells onto a continuous differentiation axis that begins from the ventricular zone (VZ), extends through the inner and outer subventricular zones (SVZ/oSVZ), and ends in postmitotic neurons. Control organoids follow a regulated trajectory with dorsal and ventral patterning, reflecting the typical sequence of corticogenesis seen in human organoids and fetal cortex[56,57]. The first trajectory (blue line), which expresses markers of ventral radial glia, leads to the formation of GABAergic inhibitory neurons. The second trajectory (magenta line) features markers of dorsal and outer radial glia, leading to excitatory neurons. Importantly, ventral patterning is disrupted in CCB79-KO organoids **(Figures 4C-D).**

In analyzing dorsal patterning events, we observed a clear, genotype-specific shift in the transcription factors of intermediate progenitors (IPs). In control samples, EOMES is a key transcription factor of IPs that promotes the specification of upper-layer excitatory neurons during forebrain development. In contrast, CCB79-KO cells expressed low levels of EOMES but an elevated level of ATOH1 along with its downstream regulator TLX3. ATOH1 is a transcription factor usually restricted to hindbrain neurogenesis, which enforces glutamatergic identity in dorsal spinal and hindbrain neurons **(Boxed line graphs in Figure 4D)** [58-60]. This signature suggests that the loss of CCB79 also disrupts dorsally specified IPs but induces an ATOH1-driven progenitor program, which interferes with the typical EOMES-driven expansion phase in the SVZ/oSVZ. This imbalance likely causes shifts in dorsal cell types by reducing the number of classical IPs and upper-layer neurons while increasing atypical progenitors that are stalled in dorsal identity.

Furthermore, measuring the relative expressions of identity markers in progenitors, precursors, and neurons showed that control organoids exhibit rostral patterning events (identified by FOXG1 and OTX2). In contrast, CCB79-KO organoids shifted from rostral to caudal patterning, accompanied by an increase in hindbrain progenitors (HOXB1 and MSX1) and a decrease in inhibitory interneurons (DLX1 and GAD2) **(Figure 4E)**. We suspected that the observed misdifferentiation could cause defects in the cytoarchitectural organization of the CCB79-KO organoids.

To map this, we calculated the relative proportion of cell types in specific brain regions during development **(Figure S11A)**. We observed a significant depletion of cells in the VZ of CCB79-KO organoids, along with a downregulation of early progenitor markers, including PAX6 and FOXG1. This suggests either premature differentiation or impaired maintenance of radial glial cells, leading to a reduced progenitor pool **(Figure S11B)**. The increased presence of cells in the intermediate zone (IZ) and migratory zone (MZ) coincides with a sharp decline in the tangential migratory stream, suggesting that migration has stalled. Differential gene expression analysis reveals a downregulation of key neurogenic transcription factors (SATB2, DCX, DLX1/2) and neuronal markers (NEUROD1/2), as well as interneuron-specific transcription factors (LHX6, ARX), indicating delayed or impaired neuronal migration to the cortical plate **(Figure S11C)**. Additionally, marker-based scoring of division modes of radial glial cells showed an increase in asymmetric divisions and a decrease in symmetric divisions in CCB79-KO, indicating that neuronal commitment expands at the expense of self-renewing progenitors **(Figure S11D).** Overall, these findings demonstrate that CCB79 deletion disrupts the neurogenesis program through combined defects in progenitor dynamics, transcriptional regulation, and regional signaling, emphasizing the essential role of primary cilia in coordinating human cortical development. The schematic summarizes potential events that occur when CCB79 is removed **(Figure S11E).**

CCB79 ablation perturbs brain tissue organization due to NPC depletion, and the mouse gene does not rescue the defects.

To verify the findings from our transcriptomic data and to characterize the effects of impaired ciliogenesis on 3D tissues, we performed imaging-based analyses of cell composition and cytoarchitecture. To visualize the entire brain organoid's cytoarchitecture in its intact form, we tissue-cleared them, immunostained with SOX2 (progenitors), ARL13B (cilia), and MAP2 (neurons), and imaged with a light-sheet microscope. Despite its low resolution, we were able to volumetrically visualize the cytoarchitecture and measure the VZ, including its diameter and lumen size. Compared to controls, CCB79-KO organoids showed disrupted organization, characterized by fewer but thinner rosettes and an irregular distribution of neurons **(Figure 5A-D)**. Since the primate CCB79 has undergone significant changes compared to the rodent gene, we tested whether the rodent gene could functionally rescue brain organization defects in 3D tissues. For this, we generated brain organoids using CCB79-KO iPSCs that express human CCB79 and mouse genes, as shown in Figure 3D. As stably expressing clones were not obtainable, we engineered them to express the

genes in an inducible manner using doxycycline. While the human gene significantly rescued the rosette defects, it did not rescue defects related to the VZ. Conversely, the mouse gene failed to rescue any of the defects and even worsened the phenotype of CCB79-KO organoids, reaffirming our results with 2D cells described in Figure 3D **(Figure 5E-F).**

We then analyzed the distribution of PAX6, SOX2, ARL13B, and TUJ1-positive cells in organoid slices and quantified their numbers. We found that CCB79-KO organoids exhibit perturbed ciliogenesis characterized by barely visible cilia in their lumen. Additionally, CCB79-KO organoids exhibited a disrupted and thinner VZ, with only a few layers of radial glial cells, and a noticeable decrease in PAX6/SOX2 intensities **(Figure 5G-H**).

We then investigated premature differentiation of NPCs as suggested by our scRNA analysis **(Figure 11D)**. We found that a TUJ1-positive region is confined to the basal side of control organoids, forming a primitive cortical plate (CP) separated from the VZ. In contrast, CCB79-KO organoids showed neurons throughout the organoids, including within the VZ, along with a decrease in radial glial progenitors **(Figure 6A).** Since CCB79 is enriched in intermediate progenitors, we examined the presence of these cells. In controls, TBR2-positive cells form a layer of the subventricular zone (SVZ) basal to apical progenitors lining the ventricular lumen. Conversely, CCB79-KO organoids did not display a TBR2-positive layer. Similarly, scoring for PTPRZ1-positive outer radial glial cells revealed a noticeably reduced population in both mutant organoids **(Figure 6B-D).** Therefore, the finding of increased neuron levels coupled with progenitor depletion indicates that premature neurogenesis occurs in organoids lacking functional CCB79 protein.

To explore the basis of premature differentiation, we studied the kinetics of apical progenitors whose division plane mostly orients horizontally toward the ventricular lumen. This orientation suggests a symmetric expansion of apical progenitors, leading to the formation of various NPC types critical for building a layered neocortex. Phospho-vimentin marks dividing apical progenitors, and by analyzing the division plane relative to the ventricular lumen, we determined the proportion of apical progenitors oriented horizontally (proliferating) versus vertically (differentiating) (Refs). While control organoids mainly exhibited horizontally oriented apical progenitors, CCB79-KO organoids contained cells oriented vertically, indicating a tendency to differentiate and be displaced from the apical side. This supports the finding that control organoids had significantly more Phospho-vimentin-positive cells than the CCB79-KO organoids **(Figure 6E-H)**. These results suggest that the loss of functional CCB79 impairs ciliogenesis, which subsequently triggers NPC differentiation, leading to NPC depletion and ultimately resulting in defective brain tissue organization, similar to that observed in microcephaly.

Discussion

Using a cell-type-specific gene enrichment analysis of human brain organoids, we identified KIAA0408 (CCB79), a previously uncharacterized vertebrate-specific gene. We determined its function as a primate-specific cilium initiation factor. Our approach demonstrates the effectiveness

of using human organoids to identify primate- and human-specific gene functions that contribute to the expansion of the human neocortex[61,62]. Molecularly, CCB79 is a centriolar DAP that anchors the mother centriole to the plasma membrane during ciliogenesis. In this process, CCB79 is a cell cycle-regulated molecule and appears only at the start of ciliogenesis (G1/G0 phase), where it recruits TTBK2, another DAP component, to initiate ciliogenesis **(Figures 2, S3, and S5).**

Even though centriolar DAPs and cilia are expected to be expressed in all neural cell types, CCB79's expression is enriched in intermediate progenitors, which are relatively abundant neurogenic cell types in gyrencephalic brain species critical for generating neurons in the evolutionary expansion of the human cortex[9,10]. This suggests that context-specific ciliary functions have evolved during brain development, warranting further research into the roles of cilia in human brain development and disease.

Notably, although CCB79 is conserved in vertebrates, the human gene has experienced significant changes and faced an evolutionary challenge in initiating ciliogenesis **(Figure 3).** Our rescue experiments show that the mouse gene neither targets the ciliary base nor restores ciliogenesis in human iPSCs lacking CCB79 **(Figure 3C-D)**. This suggests that human CCB79 has specifically evolved to promote ciliogenesis in human cells, leading us to hypothesize that human cilia may have recently evolved, at least in specific cell types of the developing brain, in terms of composition, length, and dynamics. This is surprising because centrioles containing DAPs are very ancient structures conserved from single-celled organisms to humans; thus, significant evolutionary changes in them are unexpected. However, it appears that CCB79 has undergone rapid evolution in terrestrial animals. It will be interesting to test the expression of human CCB79 in rodents and determine whether it can alter cilia, thereby affecting cell composition and inducing gyrification in lissencephalic rodent brains.

Our findings indicate that telencephalic and ventral progenitors, as well as the inhibitory interneurons they generate, are particularly susceptible to CCB79 disruptions **(Figure 4)**. We could precisely identify these defects because our organoids consistently mimic the in vivo cell type diversity of human brain development, including essential progenitor populations and both excitatory and inhibitory interneurons derived from the medial ganglionic eminence.

To date, no non-syndromic ciliopathy or neurodevelopmental disorder has been linked to CCB79. However, our findings suggest that the CCB79 gene might be a risk factor for microcephaly and autism-like disorders. This is because most affected cell types after CCB79 perturbation are ventral progenitors, which typically develop into GABAergic interneurons and oligodendrocyte precursors under normal conditions. When CCB79 is perturbed, the fate determination of ventral progenitors is disrupted, leading to the formation of abnormal oligodendrocyte precursors at the expense of ventral progenitors. Earlier studies have shown that prolonged or excessive SHH causes ventral progenitors to differentiate into oligodendrocytes, although the underlying mechanism remains unclear. Our data address this gap, as CCB79-KO cells lacking functional cilia display signs of excessive SHH activation, which may be a key factor in disrupting the balance.

Our study has limitations. First, we were unable to generate homogeneous populations of mouse transcript-expressing iPSCs with regulated expression of the gene product. This is primarily because expression of CCB79 is temporarily regulated and appears only at the onset of cilium initiation **(Figure 2).** Future studies should investigate the expression of the human transcript in the mouse endogenous locus or the mouse gene knocked into human iPSCs to dissect the specific aspects of cilia and progenitor biology affected by the mouse and human genes. Secondly, we do not know if CCB79 mutations have occurred in human patients. It would be interesting to screen patient cohorts for CCB79 mutations and determine if CCB79 is a risk factor for neurodevelopmental disorders.

Nevertheless, our findings provide new insights into cilia biology and its previously unknown connections to the maintenance of progenitors and human brain development mechanisms **(Figures 4-6).** Human cortical development involves intricate processes that generate a wide variety of cell types, with their diversity and connections carefully regulated through a complex interplay of transcription, epigenetics, cell adhesion, and polarity factors. CCB79 removal, however, primarily affects DAP assembly and ciliogenesis, yet it significantly disrupts cell diversity, indicating a crucial role for cilia in regulating brain organization. In terms of cilia biology, since CCB79 has never been studied and is a novel DAP component, our discoveries will improve the molecular understanding of DAP roles in cilium initiation. The results and resources presented lay the groundwork and open new avenues for cilia research to investigate CCB79's potential isoforms, interaction partners, and regulatory mechanisms in ciliogenesis. Additionally, these findings strengthen the underexplored connection between cilia dynamics and human brain development.

Acknowledgment

This work is supported by Fritz Thyssen Foundation (10.20.2.031MN), and Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 503306912 – FOR5547, DFG, GO 2301/5-2.

Author contributions statement

JG conceived the concept, and DR performed most of the experiments; SW performed the initial localization findings. EG reprogrammed iPSCs and generated NPC experiments; VJ performed transcriptomic data analysis, OSV conducted 3D tissue clearing and light sheet imaging; ZZ, AM, TB, and CG conducted SIM and STED-microscopy imaging; TB conducted Western blot experiments; NA edited the iPSCs and ablated CCB79; MR and GC conducted EM experiments. JG, EG, DR, and VJ wrote the manuscript. All authors have read the manuscript and approved it for publication.

Competing interest statement

The authors declare no competing interests.

METHOD DETAILS

Human iPS cells and culturing: Human iPS cell lines IMR90 and BJ151 were cultured in mTeSR™ plus medium (Stem Cell Technologies) on Matrigel-coated cell culture dishes (Corning) at 37°C and 5% CO_2 and were routinely checked for mycoplasma contamination using the MycoAlert Kit (Lonza). Cells were dissociated into small aggregates using ReLeSR (Stem cell Technologies) every 5-7 days and split 1:5 onto fresh Matrigel-coated dishes. Unless otherwise stated, most analyses shown were performed using organoids and NPCs derived from IMR90 (Coriell) or BJ151, which was reprogrammed in-house.

Generation of CCB79 (KIAA0408) knock-out iPSCs: Two single guide RNAs (sgRNAs) targeting exon 3 of the *KIAA0408* gene were chosen based on high predicted on-target activity and minimal off-target effects: sgRNA1: ATGTCGGTCGATGTGCAGCA, and sgRNA2: CATCGACCGACATCAGATCC. Each sgRNA was cloned into a lab-modified pX330A-derived plasmid encoding *S. pyogenes* Cas9, a U6 promoter-driven sgRNA cassette, and a GFP reporter to monitor transfection efficiency. Oligonucleotides were synthesized with BbsI-compatible overhangs as follows: sgRNA1-FP: CACCGATGTCGGTCGATGTGCAGCA, sgRNA1-RP: AAACTGCTGCACATCGACCGACATC; sgRNA2-FP: CACCGCATCGACCGACATCAGATCC, and sgRNA2-RP: AAACGGATCTGATGTCGGTCGATGC.
Golden Gate cloning was used to insert phosphorylated, annealed oligonucleotides into the BbsI-digested vector backbone. Briefly, oligonucleotides were phosphorylated using T4 Polynucleotide Kinase (NEB) at 37 °C for 30 minutes and annealed by gradual cooling from 95 °C to 25 °C. The pX330A plasmid was linearized by digestion with FastDigest BbsI (Thermo Fisher) and purified by column purification. Annealed oligos (1:200 dilution) were ligated into the digested vector using T4 DNA Ligase (NEB) in a Golden Gate assembly. Ligation products were treated with PlasmidSafe DNase (EpiCentre) to remove linear or un-ligated DNA and subsequently transformed. Positive clones were confirmed by Sanger sequencing, and validated constructs were used for downstream transfection.
hiPSC IMR90 cells were cultured on Matrigel-coated plates in mTeSR1 medium and dissociated using Accutase. For each sgRNA, 75,000 cells were seeded per well in a 12-well plate. Cells were transfected using GeneJuice (Sigma-Aldrich) with 1 μg of plasmid DNA per well. After 24 hours, cells were treated with 1 μg/mL puromycin for 24 hours. Transfection efficiency was assessed by monitoring GFP fluorescence.
Targeted mutations revealed by the mismatch-sensitive T7 endonuclease I (T7E1, NEB) assay. For clonal validation, single cells were seeded at low density to allow colony formation. Individual clones were manually picked, expanded, and screened by PCR and Sanger sequencing to confirm indels or frameshift mutations introduced by CRISPR-Cas9 editing.

NPC generation: NPCs were differentiated from iPSCs as previously described [38-40]. iPSCs were dissociated into single cells using Accutase. For the neurospheres, 45,000 iPSCs were seeded into a 96-well V-bottom plate and maintained in NIM for 5 days. Neurospheres were transferred into a PLO/Laminin-coated 35 mm dish using a 200 μL cut tip and evenly distributed. The dish was incubated at 37 °C with 5% CO_2 for 7 days, with a medium change performed daily. The adherent neurospheres with neural rosettes were washed once with DMEM-F12 and incubated for 1 hour in Neural Rosette Selection Medium at 37 °C and 5% CO_2. After incubation, DMEM-F12 was used to dislodge only the neurospheres from the dish. The neurospheres were gently resuspended using an uncut 1000 μL pipette tip to generate smaller aggregates. These aggregates were transferred to a Falcon tube, centrifuged at 700 × g for 3 minutes, and resuspended in 1 mL of NIM supplemented with 10 μM ROCK inhibitor. The cell suspension and small aggregates were evenly distributed in a PLO/Laminin-coated 35-mm dish and incubated at 37°C with 5% CO_2. For expanding the NPCs, medium changes were performed with STEMdiff™ Neural Progenitor Medium (STEMCELL Technologies, Cat #05833) daily or every other day until confluency was achieved.

Brain organoid generation: The brain organoids were generated by a non-directed differentiation method as previously described [35]. The iPSCs were cultured in mTeSR Plus culture medium and passaged using ReLeSR. The cells were regularly checked for mycoplasma contamination and their stem cell morphology. For initiating neural differentiation, cells were dissociated into single cells.

Therefore, iPS cells were carefully washed once with pre-warmed DMEM/F12 (GIBCO) and then treated for 5 minutes at 37°C with Accutase (Sigma-Aldrich). After cell collection, centrifugation, and counting, cells were diluted in neural induction medium (NIM, Stem Cell Technologies) plus Rock inhibitor (10 mM, Y-27632 2HCl, Selleck Chem) at a concentration of 3,5 × 10^5 cells/mL. 100 µL of this cell suspension was added to each well of a 96-well V-bottom low-adherent plate and incubated at 37°C and 5% CO2. Half of the medium was changed every day for 5 days until neurospheres were formed.

Neurospheres in Petri dishes and spinner flasks: At day 5 of neural induction, the neurospheres were carefully collected using a 200 µL pipette tip and embedded into Matrigel basement membrane extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma. The embedded neurospheres were cultured in a petri dish for another 4 days, containing neurosphere medium. On day 9, the neurospheres were transferred into a spinner flask containing brain organoid medium and were then cultured for an additional 14 days. Starting from first day of differentiation, the process takes a total of 24 days.

Immunofluorescence microscopy of 2D cells and 3D brain organoid tissues: Human iPSCs, NPCs, and brain organoids were immunostained using the following method. Materials were fixed in 4% PFA (Merck Millipore) for 30 min at room temperature (RT) and treated with PBS-Glycine (0.225 g Glycine per 100 mL PBS). For organoids, PBS-Glycine was removed, and 30% Sucrose (Merck Millipore) was added to dehydrate the tissues overnight at 4°C. The next day, organoids were embedded into Tissue-Tek® Cryomold® Cryomolds (Science Services) containing Tissue-Tek® O.C.T. Compound solution (Science Services) and immediately stored at 80°C for a minimum of 24 hours. Frozen specimens were embedded and cryo-sectioned at 12 µm thickness with Cryostat (Leica CM3050 S) and placed onto Poly-L-lysine (Sigma-Aldrich) pre-coated Superfrost glass slides (0.01% PLL diluted in distilled H2O and coated for 5 min at RT, subsequently dried for 3 hours at RT) (Thermo Scientific). Slides with organoid sections were dried for 1 hour at RT and then kept for long-term storage at 80°C.
Before immunofluorescent staining procedures, we thawed the slides with organoid sections for 30 min at room temperature, washed 2x 3 min with PBS-Glycine (0.225 g per 100 ml) and permeabilized for 15 min in 0.5% Triton X-100/0.1% Tween (Sigma Aldrich) in PBS solution at RT, and washed again 2x 3 min with PBS-Glycine. Subsequently, sections were blocked with blocking solution (0.5% Fish gelatin in PBS) for 1 hour at RT. Primary antibodies were diluted in blocking solution, and sections were incubated with primary antibodies for 2 hours at RT.
2D cells were treated on glass coverslips that were fixed with either 4% PFA or ice-cold methanol. Methanol fixation was used to stain CCB79. After 3x washing with blocking solution, corresponding secondary antibodies (goat-anti mouse Alexa 488 or 647; donkey-anti rabbit Alexa 488, 594 or 647; donkey-anti sheep 647 or donkey-anti goat Alexa 594) were diluted in blocking solution (1:800) and incubated for 1 hour at RT protected from light. Finally, sections were stained with DAPI, washed twice for 3 minutes with blocking solution, and after a final wash with distilled water, briefly air-dried, and then mounted with Mowiol (Sigma Aldrich) and glass coverslips. Slides were dried for one hour at RT and then stored at 4°C in the dark until imaging.

Confocal and light-sheet microscopy: Tissue sections or 2D cells on glass coverslips were imaged using a Leica SP8 confocal microscope. Our experiments utilized laser lines at 405, 488, 561, and 633 nm and were imaged using objective lenses with a 63x oil immersion magnification. We used 405, 488, 561, or 633 nm laser lines and imaged the samples using objective lenses with magnifications of 10x/0.3, 20x/0.8, and 40x/1.3 oil immersion objectives.
For light sheet imaging, the brain organoids were fixed in 4% warm PFA for 1 hour at room temperature on a gentle rocker. The samples were then washed twice in PBS before being stained with antibodies. Briefly, the organoids were permeabilized in a permeabilization buffer for 1 hour at room temperature, followed by blocking for 1.5 hours in a blocking solution with fish gelatin. The organoids were incubated overnight at 4 °C in the blocking solution with primary antibodies against SOX2 (Abcam, 1:400 dilution) and MAP2 (Proteintech, 1:400 dilution). The organoids were then washed three times in blocking solution for 30 minutes. The organoids were then incubated in a primary antibody against ARL13B (BiCell Scientific, 1:400 dilution) for 3 hours at room temperature. The organoids were again washed as indicated above, followed by incubation in blocking solution

with secondary antibodies (1:1000 dilution) overnight at room temperature. The other day, the organoids were washed as indicated above, followed by embedding the organoids in low-melting agar.

The embedded organoids were sequentially dehydrated using 30%, 50%, 70%, 90% and absolute ethanol. After the dehydration step, the blocks were treated with Ethyl Cinnamate overnight, followed by imaging.

Imaging data acquisition was done using the Light Sheet 7 system (Carl Zeiss) with a 5x objective at 1x zoom factor and a 20x objective at a 2x zoom factor. The same step-size (z step thickness) and imaging depth (number of z steps) were employed for computing the Maximum Intensity Projections, sections of the VZs, volumetric rendering, and image-based quantifications. Raw data were subjected to channel alignment (case-based) in Zen Black software (Zeiss) before analysis. Imaging data analysis was performed using Arivis Pro (Zeiss). Image-based machine learning and segmentation were employed for quantifying VZ volume distribution, lumen diameters, and the number of rosettes across samples. The blob finder module was used for performing segmentation and quantification of progenitor densities in a fixed volume in the rosettes across the samples. The data was exported in MS Excel. Statistical analysis and graph plotting were performed in GraphPad Prism 10. The raw images acquired were processed with ImageJ, Adobe Photoshop 2024, and Adobe Illustrator 2024. Movies of organoid Z-stack images were processed using ImageJ.

STED (Stimulated Emission Depletion) Super-Resolution Imaging: Two-color STED images were acquired using the Abberior Expert Line, equipped with pulsed excitation and depletion lasers, and an Olympus UPlanSApo 100x/1.4 oil immersion objective as described previously [63]. Proteins of interest were labeled with Abberior Star-Orange (Distal appendage protein FBF1 and Cilia marker ARL13B). Abberior Star-Red (rat CCB79) dyes via secondary antibodies and were excited by 561 nm and 640 nm pulsed diode lasers, respectively. The fluorescence signals were inhibited by a pulsed 775 nm depletion laser (total power = 3 W). DAPI was excited with 405 nm/488 nm pulsed diode lasers. Before image acquisition, the excitation and depletion beams were aligned for both 2D and 3D STED.1 Briefly, the center of the excitation and depletion beams was overlapped initially by scanning gold beads of 150 nm (BBI Solutions) in reflection mode. Afterwards, TetraSpeck beads of four colors (TetraSpeckTM Microspheres, 100 nm, fluorescent blue/green/orange/dark red) were used to correct mismatches between the scattering mode and the fluorescence mode. To ensure the correct and precise positioning of the same beads imaged by different laser lines, individual confocal and STED channels were compared and adjusted, respectively, as the last step on the Imspector SLM (spatial light modulator) adjustment panel. During STED sample imaging, sequential line-by-line scanning was applied to minimize photobleaching of the samples, starting with the Star-Red channel and then the Star-Orange channel. The images were acquired according to the Nyquist criterion for super-resolution, with a pixel size of 20 nm and a pixel dwell time of 10 µs; a distance of 100 nm was applied between each slice for xyZ stacks. The raw STED images were deconvolved using Huygens Professional version 24.10 (Scientific Volume Imaging, https://svi.nl/) with the classic maximum likelihood estimation (MLE) algorithm, incorporating lateral drift stabilization, and with 27 iterations.

Structured Illumination Microscopy (SIM) Super-Resolution Imaging: To provide complementary super-resolution imaging with lower phototoxicity and a broader field of view, SIM was performed to image the same protein-labeled samples used for STED. Images were acquired using a commercial Zeiss Elyra 7 Lattice SIM system, equipped with a Plan-Apochromat 100x/1.46 oil immersion objective, and sensitive sCMOS cameras. Structured illumination was applied in a 3D mode, and image reconstruction was performed using Zeiss ZEN Black, with default parameters for modulation contrast and noise filtering. SIM images were acquired at Nyquist sampling conditions, with voxel sizes ranging from 40 to 60 nm in the xy plane and from 120 to 150 nm in the z direction. Image post-processing was performed using Fiji.

Antibodies used: The following antibodies were used. The following antibodies were used. Rat anti-CCB79 (1:100, generated in house via BiCell Scientific), Rabbit anti-CCB79 (1:500, generated in-house via BiCell Scientific), Rabbit anti-CEP164 (1:200, Proteintech), Rabbit anti-TTBK2 (1: 100, Sigma Aldrich), Rabbit anti-SCLT1 (1: 25, Proteintech), Rabbit anti-FBF1 (1: 200, Proteintech), Rabbit anti-CEP89 (1: 500, Proteintech), Rabbit anti-CEP83 (1: 300, Sigma Aldrich), Rabbit anti-

ARL13B (1: 100, Proteintech), Rat anti-ARL13B (1: 100, BiCell Scientific), Mouse anti-ARL13B (1: 200, Proteintech), Mouse anti-GT335 (1: 2000, AdipoGen), Rab anti-CP110 (1: 200, Proteintech), Mouse anti-CPAP (1: 25, In house), Rabbit anti-ODF2 (1: 100, Proteintech), Mouse anti-SOX2 (1:500, Abcam), Rabbit anti-PAX6 (1:10 Proteintech), Rab anti-FoxG1 (1:300,), Mouse anti-Nestin (1:100, Novus Biologicals), Rabbit anti-TUJ1 (1:300, Sigma-Aldrich), Rabbit anti-pVIM (1:500, Abcam), Rabbit anti-TBR2 (1: 100, Abcam), Rabbit anti-PTPRZ1 (1: 200, Sigma-Aldrich), Rabbit anti-MAP2 (1: 200, Proteintech). We used Alexa Fluor Dyes conjugated with goat/donkey anti-mouse, anti-rabbit, or anti-rat antibodies (1:1000, Molecular Probes, Thermo Fisher, USA) as secondary antibodies. In addition, DAPI 1 µg/ml (Sigma Aldrich, USA) was used to stain the nucleus.

Tissue-specific expression analysis of CCB79 (KIAA0408): Tissue-specific expression data for CCB79 (KIAA0408) were obtained from the UCSC Genome Browser GTEx Gene V8 Track (Human genome GRCh38/hg38), summarizing mean expression across fetal and adult human tissues. Data were loaded using pandas and organized into fetal and adult expression matrices. To enable comparison, expression values were min–max normalized using the fetal expression range as the reference. Heatmaps were generated using seaborn, with tissues sorted by fetal KIAA0408 expression. A consistent viridis color scale was applied to both fetal and adult panels to highlight developmental differences. Single-cell RNA-seq data from human developing neocortex were obtained from a publicly available dataset[38] representing cells from 20 defined developmental stages. Expression of genes, including SOX2, PAX6, DCX, and CCB79 (KIAA0408), was visualized across these stages using a dotplot with Scanpy, with standardized expression to capture gene dynamics during brain development.

Organoid dissociation, single-cell preparation, and scRNA sequencing: Human iPSC-derived brain organoids at Day 14 from both control and CCB79-KO genotypes were processed for single-cell preparation using a modified protocol adapted from [56]. Briefly, 3–4 intact organoids were rinsed three times with 1× PBS to remove residual media. Organoids were then incubated in 2 mL Accutase (StemPro) supplemented with 0.2 U/µL DNase I (Roche, [Catalog #]) for approximately 45 minutes at 37°C, with gentle flicking every 10 minutes to facilitate dissociation into single cells. If visible aggregates or debris remained after dissociation, the suspension was sequentially passed through 40 µm strainers to ensure a uniform single-cell suspension. The dissociated cells were pelleted by centrifugation at 900 × g for 3 minutes, washed once with 1 mL of 1× PBS, and then recentrifuged under the same conditions. Cell viability and concentration were assessed using Trypan Blue exclusion on a Countess™ Automated Cell Counter (Thermo Fisher Scientific).
Single-cell fixation was performed according to the 10x Genomics Flex Gene Expression Profiling protocol. Briefly, up to 2 million cells were pelleted at 350 × g for 5 minutes at 4°C, resuspended in 1 mL Fixation Buffer, and incubated for 16–24 hours at 4°C. Fixation was quenched by centrifugation at 850 × g for 5 minutes at room temperature, followed by the addition of 1 mL Quenching Buffer (1× Conc. Quench Buffer; 10x Genomics PN-2000516). To enable long-term storage, 0.1 volumes of Enhancer (10x Genomics PN-2000482) and 10% glycerol were added, and samples were stored at -80°C. Before library preparation, samples were thawed at room temperature, centrifuged at 850 × g for 5 minutes, and resuspended in 1 mL of 0.5× PBS containing 0.02% BSA.

Hybridization of probe sets was carried out according to 10x Genomics recommendations. Up to 2 million fixed cells were used per hybridization reaction in a total volume of 80 µL, consisting of 60 µL hybridization buffer and 20 µL of either Human Transcriptome Probe Panel v1. Hybridizations were incubated at 42°C for 16–24 hours. After hybridization, samples were diluted in Post-Hybridization Wash Buffer and analyzed using an automated Cell Counter.
For standard pooling, equal cell numbers from each hybridization were combined; for differential pooling strategies, cell numbers were adjusted as needed. Cell pools underwent three washes in Post-Hybridization Wash Buffer at 42°C for 10 minutes each, followed by resuspension in Post-Hybridization Resuspension Buffer. Samples were filtered through a 30 µm filter to remove aggregates, then recounted before loading on the Chromium system.
NGS library preparation was performed using the 10x Genomics multiplexed Fixed RNA Profiling protocol, following the manufacturer's guidelines (CG000527 and CG000478 for sample fixation). Libraries were prepared with input ranging from approximately 140,000 to 2,000,000 cells per sample. Sixteen samples were pooled into a single batch, aiming for an average of 8,000 cells per

sample, and amplified through 20 PCR cycles. A post-PCR purification step was conducted to remove residual primers and adapter dimers. Library quality was assessed using a High Sensitivity DNA chip on a 2100 Bioanalyzer (Agilent Technologies) and quantified with the Qubit 1X dsDNA HS Assay Kit on a Qubit 4.0 Fluorometer (Invitrogen by Thermo Fisher Scientific). Sequencing was carried out on a NextSeq 2000 platform with a P3 flow cell, using 28 cycles for read 1, 90 cycles for read 2, and 10 cycles each for i7 and i5 indexes at the Core Facility Genomics, Institute of Molecular Biology GmbH (IMB), Mainz, Germany.

Data processing and analysis: Cell Ranger Analysis: Raw 10x Genomics Flex data (FASTQ files) were processed using Cell Ranger v7.1.0 (10x Genomics) and aligned to the human reference (GRCh38-2020-A, provided by 10x Genomics) using the Cell Ranger multi-pipeline with default parameters. The pipeline performed demultiplexing, read alignment, barcode processing, and UMI counting to generate feature-barcode matrices for downstream single-cell RNA-seq analysis. For downstream single-cell RNA-seq data processing, highly variable genes were identified, and the total number of counts per cell, along with the percentage of mitochondrial reads, were regressed out to normalize technical variation. Principal component analysis (PCA) was performed, followed by batch correction using the Harmony package (v1.2). To account for local neighborhood structure, a k-nearest neighbor graph was computed using the BBKNN algorithm, and a two-dimensional representation was generated with the UMAP and Force Atlas2 (FA) algorithm for visualization. Cell type annotation for the CCB79-KO and control was performed using canonical marker genes and top differentially expressed genes (DEGs) identified for each cluster **(Supplementary table 2)**. The proportion of each annotated cell type was used as input for principal coordinate analysis (PCA) to assess inter-sample relationships. Unsupervised clustering was performed using the Leiden algorithm, and marker genes for each cluster were identified through differential expression analysis using the Wilcoxon Rank Sum test and t-tests. Pseudotime trajectory inference was conducted using the diffusion pseudotime (DPT) algorithm, with proliferative radial glia cells designated as the root state. All analyses were implemented in Scanpy (version ≥ 1.10), and the whole pipeline was scripted and logged to ensure reproducibility.

Bulk RNA transcriptomic analysis: Bulk RNA sequencing analysis was conducted on six samples, representing two experimental groups (CCB79-KO and control), with three biological replicates each. Metadata detailing sample identifiers, condition labels, and replicate numbers was compiled into a structured tab-delimited file for downstream processing and analysis. Briefly, raw FASTQ files were aligned to the human reference genome (GRCh38) using the STAR aligner (Dobin et al., 2013) version 2.7. Gene-level quantification was then performed using feature counts from the Subread package version 2.0.2, using the corresponding annotation file. The resulting count files were parsed to extract gene-level expression values and merged into a unified expression matrix across all samples [64,65]. The matrix was subset to include only protein-coding genes, and genes with fewer than 10 total counts across all samples were excluded to reduce technical noise.

Differential expression analysis was conducted using the DESeq2 package ([66]version 1.44.0. All analyses compared CCB79-KO to control. Significantly differentially expressed genes (DEGs) were identified based on an adjusted p-value threshold of 0.05, corrected with the Benjamini-Hochberg method. For visualization, MA plots and volcano plots were created using DESeq2 and ggplot2 version 3.5.2, while heatmaps of the top 50 DEGs were generated with the pheatmap package version 1.0.13. Functional enrichment analyses used the clusterProfiler package [67], version 4.12.6, with a focus on Gene Ontology (GO) biological processes and KEGG pathways. Significant DEGs were mapped from gene symbols to Entrez Gene IDs using the org. Hs.eg.db annotation database version 3.19.1, and enrichment results were visualized as dot plots. All analyses were carried out in R (version 4.0 or later), with the complete pipeline scripted and logged for reproducibility.

Hedgehog signaling activity: To assess Hedgehog (HH) signaling activity, we implemented a weighted pathway activation scoring strategy using z-transformed expression values of genes associated with key functional roles, similar to previously described contexts that use gene expression of signaling components to infer pathway activation[68-71]. Briefly, genes were categorized into three main functional groups: activators, suppressors, and ligands. Activators (e.g., GLI1, GLI2, SMO, BCL2, WNT2B) were assigned a weight of +1, reflecting their positive regulatory role in HH signal transduction and transcriptional output [72,73]. Suppressors (e.g., PTCH1, PTCH2, SUFU, HHIP)

were given a weight of -1, as they inhibit pathway progression by sequestering or repressing activators [74]. Ligands (e.g., SHH, IHH, DHH) were assigned a moderate weight of +0.5 to reflect their role as extracellular initiators of the pathway without directly encoding for downstream transcriptional regulators. The weighted HH score was computed by first scaling gene expression across samples (z-transformation), then multiplying each gene's expression value by its assigned weight, and finally summing these weighted scores for each sample. This composite metric estimates the net activation state of the HH pathway in each biological replicate, capturing the balance between activation and repression signals at the transcriptomic level.

To explore the biological relevance of HH pathway modulation, we next investigated its relationship with the expression of genes involved in forebrain development. The HH signaling pathway is known to be crucial in early brain patterning, including the regionalization of the forebrain, midbrain, and hindbrain, and particularly in the specification of the ventral forebrain [75-77]. Genes such as FOXG1, PAX6, TBR1, DLX2, and EMX2 are canonical regulators of telencephalon development and neuronal differentiation. Additionally, DLX2 and GSX2 are also known transcriptional targets or downstream effectors of the HH pathway in neural progenitors [78]. To investigate this link, we correlated the weighted HH scores with the expression levels of selected forebrain marker genes across control and CCB79-KO samples using Pearson correlation. This correlation was visualized using both heatmaps and bar plots to identify candidate genes whose expression may be co-regulated or influenced by HH pathway activity. The results provide insight into whether alterations in HH signaling in the CCB79-KO condition may impact forebrain-related transcriptional programs, potentially reflecting neurodevelopmental dysregulation associated with genetic perturbations.

Transmission electron microscopy (TEM): iPSCs were washed once in PBS and then fixed in 2.5% Glutaraldehyde and 1.8% Sucrose in PBS overnight at 4°C. Samples were rinsed in PBS and then post-fixed with 1% osmium tetroxide for 1 hour at 4°C. After fixation, the samples were further dehydrated in a graded ethanol series and embedded in Epon-Araldite resin for 48 hr at 60°C. The Reichert Ultracut E ultramicrotome was used to obtain ultrathin sections (40-60 nm). These sections were further stained with uranyl acetate and lead citrate and then observed with a FEI Tecnai G2 Spirit 255 transmission electron microscope operating at 100 kV.

Molecular biology and cloning: The cDNA sequences of the human or mouse CCB79 gene were cloned into the PiggyBac plasmid pXLONE-MCS-GFP. Primers incorporating KpnI and NotI restriction sites were designed to amplify the target cDNA. Both the PCR product and the plasmid were digested with KpnI and NotI, followed by ligation. The ligation mix was transformed into TOP10 competent *E. coli* and plated on LB agar containing ampicillin. Positive clones were screened by restriction digestion and confirmed by Sanger sequencing to include the correct insert.

Western blot: Cell pellets were lysed in a standard lysis buffer on ice. 60-70ug of Protein samples were resolved in 10 or 12% Acrylamide gels and were transferred to nitrocellulose membrane. The membrane was then blocked with 5% skimmed milk at RT for 1 hr. All Primary antibodies were incubated overnight at 4 °C, and secondary antibodies were incubated at room temperature for 1 hour. Super signal Western Pico or Femto was used to detect the signal. Antibody dilutions are as follows: Anti-Rabbit CCB79 (C-terminal), anti-mouse beta-actin (1:3000), and peroxidase-conjugated secondary antibodies were used at 1:3000 (Life Technologies)

Statistical analysis: The statistical analyses were performed using GraphPad Prism (version 9). Experiments were carried out in triplicate, and all values were expressed as mean $\pm$ sem. Details of the statistical method applied and the number of data points (n = value, here "n" represents the number of organoids, slices, cells, or components) for each experiment are provided in the figure and its legend. Quantification graphs carry "n" and "p" values. In all cases, a p-value < 0.05 was considered statistically significant. Briefly, for two-group comparisons: Mann–Whitney U test, for three-group comparisons: Kruskal–Wallis test, followed by Dunn's post-hoc test with Bonferroni correction, and for categorical proportions: Fisher's exact test, was performed. Differences were considered significant at p < 0.05. All tests were two-tailed.

Main figure legends

Figure 1. Discovery of CCB79 as a controller of cilium biogenesis.
  A. UMAP plot showing significant cell clusters derived from Day 14 human brain organoids. Cluster description and markers used to annotate the clusters are given in the legend.
  B. Volcano plot of differentially expressed genes between cluster 2 (telencephalic NPCs) and cluster 5 (Intermediate progenitors) identified KIAA0408, which has unknown functions.
  C. A matrix plot to display the relative expression levels of the considered genes. Note that KIAA0408 is enriched in clusters 6 and 7, coinciding with intermediate progenitors, inhibitory, and excitatory neuronal markers.
  D. Pseudotime trajectory analysis of cell clusters showing cell lineages that follow sequential stages of brain development with dorsal and ventral patterning.
  E. Gene structure of KIAA0406. Predicted exons (green bar) are translated into a 79 kDa protein. The position and epitope sequences used to generate anti-KIAA0408 are given (blue line).
  F. Anti-KIAA0408 antibodies specifically recognize the endogenous protein located between the distal end of the mother centriole (γ-tubulin, red) and the ciliary (ARL13B, yellow) base. Scale bar 5 μm.
  G. Anti-KIAA0408 antibodies specifically recognize the mother centriole, which is also labelled by GT335 (glutamylated tubulin, yellow). The daughter centriole is specified by CP110 (red), which is absent from the mother centriole. Scale bar, 5 μm.
  H. Position of guide RNAs used to ablate CCB79 using CRISPR/CAS9 is shown on its 3^{rd} exon structure.
  I. Anti-KIAA0408 generated against the C-terminal epitope (shown in E) recognizes a 79 kDa band only in unedited iPSCs, not in KIAA0408-KO iPSCs that lack the translated protein.
  J. KIAA0408-KO iPSCs lack KIAA0408 immunoreactivity (green in control) and a fully assembled cilium (Yellow in control). Mother and daughter centrioles are stained with γ-tubulin, shown in red; scale bar, 5 μm.
  K. Serial sectioning EM analysis reveals that control iPSCs (top panel) typically exhibit structurally assembled primary cilia with a mother centriole at their base. Arrowheads indicate ciliary pockets, and the arrow points to an appendage structure. KIAA0408-KO iPSCs (bottom panel) lack both appendage structures (arrows) and the cilium. Instead of a typical ciliary pocket, the mother centriole is docked to a membrane bulge. Scale bar, 200 nm. A representative image from at least 15 cells is displayed.
  L. Bar graphs display the frequencies of ciliated cells and KIAA0408 intensity between control and KIAA0408-KO cells. Data are shown as mean ± SEM from at least three independent experiments. For KIAA0408 intensities (scatter plot at right), a non-parametric Mann-Whitney. The "n" numbers and p-values are provided in the graph.

Figure 2. CCB79 is a centriolar DAP.
  A. **(i)** SIM-based imaging of the mother (M) and daughter (D) centrioles specified by GT335 (magenta) and showing CCB79 (green) ring distal to the mother centriole, like a crown (Yellow arrowhead). Likewise, CP110 (magenta) labels only the daughter centriole that is distant from the CCB79 ring **(ii)**. OFD2, a subdistal appendage protein (magenta), is distinctly away from the CCB79 ring (green), forming a hollow tube projected into the CCB79 ring **(iii)**. CEP164, a distal appendage protein (magenta), forms a ring that co-localizes with the CCB79 ring (green). Both CEP164 and CC79 rings show periodic densities **(iv)**. STED-based imaging of FBF1, another centriolar DAP, forms a distinct nine-fold symmetrical structure of DAP that is co-localized with the CCB79 ring **(v-vi).** Top, side, and tilt views are shown for all examples. Scale bar 100 μm.

B. The scatter plot shows the mean diameter of the CCB79 ring from each centriole imaged, which is similar to DAP molecules and wider than OFD2, a subdistal appendage protein (sDAP).

C. A SIM image of CCB79 (green) and cilium (ARL13B, magenta)

D. Schematic diagram showing positions of DAP and sDAP rings at a mother centriole.

E. CCB79 staining across the cell cycle stages in NPCs. CCB79 (green) appears only at the start of ciliogenesis (ARL13B, red) at G0, disappears, and reappears in one of the late cytokinesis cells that is ciliated: γ-tubulin labels mother and daughter centrioles (yellow). The diagram of the mother (M) and daughter (D) centrioles and cilia is provided on the side to aid in interpreting the data. Scale bar 5 µm.

F. Relative intensities of DAP across the cell cycle. Note that CCB79 peaks only at G0/G1 and then declines.

Figure 3. CCB79 is a vertebrate protein and a primate-specific cilium initiation factor.

A. Phylogenetic conservation of CCB79 and other centriolar DAPs from bacteria, single-celled organisms, invertebrates, vertebrates, mammals, to humans, along with the known timeline of evolution. The last column shows the target protein (CCB79 from other species) and its percentage identity to the human CCB79 protein (represented by gradients of grey to black). Note, CCB79 appears only in amphibians and has evolved prominently in terrestrial animals.

B. AlphaFold prediction of secondary structure of CCB79 between rodents and primates. Compared to rodents, primate genes gain 43 amino acids, which enable the formation of an alpha helix in primates.

C. Inducible expression of GFP-CCB79 in wild-type iPSCs targets the ciliary base as a single dot (yellow arrowhead). The mouse gene (right panel) does not target the ciliary base efficiently and abolishes ciliogenesis. Both transfected and untransfected cells are shown, with the scale bar indicated in the panels (10 and 5 µm). Results are quantified in **C'**. Data presented as mean ± SEM from at least three independent experiments. For the frequencies of ciliated cells, a non-parametric T-test was used; for cells with ciliary GFP-CCB79 dots, a non-parametric Mann-Whitney. The "n" numbers and p-values are provided in the graph.

D. Human CCB97 expression in CCB79-KO rescues the cilia loss by targeting the mother centrioles and cilium initiation. The mouse gene (right panel) does not rescue the loss of cilia. Results are quantified in **D'** at right. Data presented as mean ± SEM from at least three independent experiments. For the frequencies of ciliated cells and cells with ciliary GFP-CCB79 dots, a non-parametric Mann-Whitney test was performed. The "n" numbers and p-values are provided in the graph.

Figure 4. CCB79 is crucial for maintaining telencephalic progenitors and ventral patterning.

A. Merged UMAP displaying various cell clusters. The legends below show the annotated cell clusters with the corresponding markers used. UMAP cell clusters of control and CCB-KO (middle panels) and proportions of various cell types between control and CCB79-KO organoids (right).

B. Sub-iteration analysis of clusters C3, C4, C5, C6, and C7 identifies 15 subclusters. The legends at right show the annotated cell clusters.

C. Pseudotime computation reveals regulatory trajectories exhibiting dorsal (magenta line) and ventral (blue line) patterning in the control. CCB79-KO shows a perturbed ventral patterning.

D. Pseudotime plot and line plots of pseudotime expression trends for individual markers using a rolling mean (window size = 500 cells), faceted by gene and colored by condition (control = blue, CCB79-KO = orange).

E. Stacked violin plots displaying expression distributions of the marker across annotated cell types in control (left) and CCB79-KO (right). Markers highlighting categories: caudal

patterning (steel blue), rostral TFs (light green), excitatory neurons (purple), inhibitory neurons (gray), and hindbrain EN (olive green). Interesting regions of discussion are boxed.

Figure 5. CCB79 ablation perturbs brain tissue organization, and the mouse gene does not rescue the defects.

   A. Tissue clearing and light sheet imaging of control and CCB79-KO brain organoids. Control organoids show defined VZs of several cell layers thick with a defined lumen at the apical side (SOX2, green). The VZ lumen (L) is defined by primary cilia (ARL13B, magenta). MAP2 defined the primitive cortical plate (CP) at the basal side (red). In contrast, CCB79-KO organoids do not exhibit a defined VZ, but rather rosette-like structures, which are significantly fewer in number than in the control: scale bar, 50 µm.

   B. Scattered plots quantify the number and dimensions of VZ between control and CCB79-KO. Data presented as mean ± SEM from at least three independent experiments. Unpaired Student's t-test was used. The "n" numbers and the "p" values are given in the graph.

   C. Volume rendered images of panel A in a 3D representation. Refer to the corresponding movies, 1 and 2

   D. Quantification of VZ volume and progenitor density between control and CCB79-KO after volume rendering. Data presented as mean ± SEM from at least three independent experiments. An unpaired Student's t-test was used. The "n" numbers and the "p" values are given in the graph.

   E. While the human CCB79 partially rescues the defects of CCB79-KO, the mouse gene fails to rescue these defects, including the number of neural rosettes, lumen diameter, and VZ diameter. Scale bar 20 µm.

   F. Scatter plots quantify the rescue effects. Data presented as mean ± SEM from at least three independent experiments. One-Way ANOVA with Multiple Comparisons test. The "n" numbers and the "p" values are given in the graph.

   G. Thin sectioning and immunostaining of control and CCB79-KO organoids. Control organoids show typical cytoarchitecture, but CCB79-KO organoids do not. The number of VZ and its thickness are reduced. Besides, both SOX2 (green) and PAX6 (magenta)-positive progenitors are weak in CCB79-KO sections. Scale bar 50 µm. Bar diagrams at right quantify the findings between control and CCB79-KO organoids. Data presented as mean ± SEM from at least three independent experiments. A nonparametric T-test, followed by a Mann-Whitney test, was performed. The "n" numbers and the "p" values are given in the graph.

   H. Zoomed-in images of VZ lumen show a well-defined VZ lumen whose apical region is lined by primary cilia (ARL13B, magenta) in control organoids (Upper panel). SOX2-positive (green) progenitors define VZ. In contrast, CCB79-KO organoids (Bottom panel) are not structurally normal, progenitors lack SOX2, lumens lack cilia, and TUJ1-positive neurons are all over. Scale bar 10 µm.

Figure 6. Progenitors in CCB79-KO brain organoids undergo premature differentiation.

   A. Control brain organoids show their typical cytoarchitecture with distinct VZ (SOX2, green) and TUJ1-positive (red) primitive cortical plate (CP). CCB79-KO organoids exhibit neurons throughout with a thinner VZ and a deficiency of SOX2-positive progenitors. Scale bar 50 µm.

   B. TBR2 (red) defines intermediate progenitors at the sVZ in controls. VZ is marked by SOX2-positive progenitors (green). Cilia (ARL13B, magenta) mark the VZ lumen. CCB79-KO organoids lack TBR2-positive cells.

   C. Control organoids display PTPRZ1, which is expressed in outer radial glial cells spanning from the VZ to the basal region (green). CCB79-KO organoids show a drastically reduced population of PTPRZ1 cells.

D. Bar diagrams quantify the findings from panels A, B, and C. Data are presented as mean ± SEM from at least three independent experiments. A non-parametric Mann-Whitney test was performed. The "n" numbers and p-values are provided in the graph.

E. Measuring premature differentiation by analyzing the kinetics of the apical progenitor division plane. Phospho-Vimentin (green) selectively labels the dividing apical progenitors at the VZ's apical side. Control organoids predominantly display dividing progenitors whose division plane is horizontal to the VZ lumen (Left panel). CCB79-KO brain organoids (Right panel) show apical progenitors whose division plane is mainly vertical to the VZ lumen. Scale bar 5 μm.

F. The bar diagram quantifies the distribution of the division plane. Horizontal and vertical. A schematic on the right shows horizontal and vertical division planes. Data are presented as proportion mean proportion ± SEM from at least three independent experiments. Fisher's exact test was performed. The "n" numbers and the "p" values are given in the graph.

G. Control organoids, in general, have numerous Phospho-Vimentin-positive (green) apical progenitors, whereas CCB79-KO do not. Scale bar 50 μm.

H. A bar diagram quantifies the distribution of Phospho-Vimentin-positive apical progenitors. Data are presented as mean ± SEM from at least three independent experiments. A non-parametric Mann-Whitney test was performed. The "n" numbers and p-values are provided in the graph.

Supplementary Figure 1 (Related to main Figure 1).

A. Heatmaps display the minimum–maximum normalized expression of CCB79 (KIAA0408) across multiple types of fetal (left) and adult (right) tissues, using GTEx V8 data.

B. Expression of CCB79 (KIAA0408) across fetal and adult brain cell types, with all values normalized to the fetal expression range to emphasize developmental differences.

C. The dot plot shows the standardized expression of key developmental and neuronal marker genes across 20 defined stages of human neocortical development. Dot size reflects the proportion of expressing cells in each group, while color indicates the normalized expression level. CCB79 (KIAA0408) expression is highlighted with a green box, indicating a similar expression profile to that of early neurons expressing DCX.

D. UMAP showing distribution of SOX2, DCX, EOMES, INSM1, NKX2-2, GAD2, SLC17A6 and KIAA0408.

E. GFP-KIAA0408 (green) expression in iPSCs showing a single dot indicated by yellow arrowheads. Scale bar 10 μm. Left panel; GFP-KIAA0408 (green) is at the base of the cilium (ARL13B, yellow). γ-tubulin labels the mother and daughter centrioles (red). Scale bar 5 μm.

Supplementary Figure 2 (Related to main Figure 1).

F. Exon structure of KIAA0408, positions of guide RNA to ablate the gene, and the results of gene ablation. The region deleted after CRISPR editing is shown. Agarose gel shows the fragment after the deletion. The predicted peptide with an induced stop codon is shown in the bottom panel.

G. **(i)** A pool of control NPCs shows normally ciliated cells (ARL13B, green). G-tubulin labels the mother and daughter centrioles (red). **(ii)** A pool of CCB79-KO NPCs shows no ciliated cells. If cilia are observed, the following categories apply. No cilia, cilia as a dot, bulgy, short, or thin, long. (ARL13B, green). γ-tubulin labels the mother and daughter centrioles (red): scale bars 20 μm and 5 μm. The bar diagram at the bottom quantifies the various categories of cilia in CCB79-KO cells. Data are presented as mean ± SEM from at least three independent experiments. The "n" numbers are given in the graph.

H. Centrosome numbers are unchanged in CCB79-KO NPCs. A bar diagram quantifies centrosome numbers between control and CCB79-KO cells. Data are presented as mean ± SEM from at least three independent experiments. A two-way ordinary ANOVA, followed by Sidak's multiple comparisons test, was used. The "n" numbers and the "p" values are given in the graph.

Supplementary Figure 3 (Related to main Figure 2).

A. Localization behavior of DAP components, CEP164, and FBF1 (yellow) across various cell cycle stages. Cilia are labelled by GT335 (red). Note, in contrast to CCB79, which localizes only at the onset of cilium initiation (G0), these DAPs localize throughout the cell cycle stages. Scale bar 5 μm.

B. Serum starvation and serum stimulation experiments using hTERT inactivated RPE1 cells. After 72 hrs of serum starvation, all cells are ciliated. CCB79 (green) is at the distal site of the mother centriole (M). Centrioles and cilia are labeled by GT335 (yellow). CP110 (red) labels only daughter centrioles (D). Cells begin to disassemble cilia after three hours of serum addition. Note that the CCB79 intensity is reduced (middle panel). The same observation is at six hours after serum addition. Dotted boxes indicate the cells that are zoomed in on the panel. Cilia are shown in the magnified panel. Asterik is not the cilium from the same cell. Scale bar 10 μm.

C. The same experiment, which includes CPAP (red), a cilium disassembly factor, which, in contrast to CCB79, gains intensity. Scale bar 10 μm.

D. Graphs that quantify the findings, such as intensity of CPAP (blue), CP110 (red), and CCB79 (green). A scatter plot displays the cilia length at various time points following serum addition. Data are presented as mean ± SEM from at least three independent experiments, and a non-parametric Mann-Whitney test was performed. The "n" numbers and p-values are provided in the graph.

Supplementary Figure 4 (Related to main Figure 3).

A. Pairwise alignment of amino acid sequences from different species is shown on the left. The boxed area is the epitope used to generate anti-CCB79 antibodies. The red boxed area, positioned at the end of exon five, indicates the region gained in primates, which comprises 43 amino acids. (Describe different species as abbreviations)

B. Sequence alignment of the region, including exon 5, indicates changes in alternatively spliced exons between the mouse and human genomes. Bases in red with an arrow indicate the area where a primate-specific point mutation likely occurred in the evolutionary time between man and mouse. Asterisks indicate conserved positions between both species. Splice site strength and splicing regulatory landscape predictions using scoring algorithms (MaxEnt, H-bond score) were analyzed by Hexplorer. The HEXplorer score profiles represent splicing regulatory potential across the exon-intron boundary. The splicing region modified by the point mutation in humans exhibits a pronounced enhancer regulatory region and a more favorable splice site (splicing domain five or SD5), as indicated by a high HBS (15.9). This occurs downstream of the mouse splice site (mouse splicing domain five or mSD5), which is further suppressed by a second point mutation in humans (HBS score changes from 14.4 to -7.3), thus enabling exonization in humans and other primates.

C. Gene structure and the coding regions of the mouse gene compared to human CCB79. The area gained in primates is highlighted.

D. Mouse and human protein sequences are given. The red bar underlines the region of 43 amino acids that is unique to primates but is absent in the mouse protein.

Supplementary Figure 5.

A. CCB79 is critical for maintaining NPCs in their self-renewal state. Under non-differentiating conditions, control NPCs self-renew. CCB79-KO NPCs undergo spontaneous differentiation into neurons (TUJ1, magenta), leading to the loss of NPCs (Nestin, green). Edu (yellow) incorporation does not differ between control and CCB79-KO NPCs. Bar graphs at right quantify the findings. Data presented as mean ± SEM from at least three independent experiments. For quantifying Nestin versus TUJ-1 positive cells, a non-parametric Mann-Whitney test was performed. The "n" numbers and p-values are provided in the graph.

B-F. DAP recruitment assay. Control NPCs recruit CP110, a centriolar protein **(red, B),** γ-tubulin **(red, C),** CEP164 **(red, D),** FBF1 **(red, E)**, and TTBK2 **(red, F)** (red) to the base of the cilia (ARL13B or GT335, magenta). CP110 labels only the daughter centrioles when the mother centriole templates ciliogenesis. In all cases, CCB79 is recruited to the mother centrioles at the ciliary base (green). The right panels show CCB79-KO NPCs, which either fail or faintly recruit CCB79 (green), and cells have either no cilia or aberrant cilia (magenta). Except for TTBK2 **(red, F),** the recruitment of other components is not significantly perturbed. Note that CP110 remains persistent in the mother centrioles that do not assemble cilia. Scale bar 5 μm.

Bar diagrams on the right quantify the findings. Data presented as mean ± SEM from at least three independent experiments in each case. For intensity measurements (CCB79, CEP164, FBF1, and TTBK2), a non-parametric Mann-Whitney test was performed. The "n" numbers and p-values are provided in the graph.

G. The drawing shows a mother centriole templating a cilium with the hypothetical topology of various DAP components, as indicated by CCB79. Legends are provided to the right to aid in understanding the experimental findings.

H. The schematic summarizes the experimental results where CCB79-KO NPCs specifically fail to recruit TTBK2, and thus, no cilium is initiated.

Supplementary Figure 6.

A. Macroscopic images of control and CCB79-KO brain organoids. Scale bar 200 µm. The scatter plot below quantifies individual brain organoids and shows that the CCB79-KO brain organoids are significantly smaller in size.

B. Thin-sectioning and imaging of whole surface brain organoids. Control shows typical architecture with VZ, whose lumen is lined by cilia (ARL13B, yellow). SOX2 (green) shows NPCs. At this magnification, there are only barely visible TUJ1-positive neurons at the basal side (left panel). CCB79-KO organoids in the right panels show a disorganized cytoskeleton characterized by the loss of VS, SOX2-positive progenitors with a gain of TUJ1-positive neurons all over the place. Scale bar 200 µm.

C. Left panel: Volcano plot depicting differential gene expression with log2 fold-change on the x-axis and –log10 p-value on the y-axis. Significantly upregulated and downregulated genes ($log_2FC > 2$, FDR < 0.05) are highlighted between the CCB79-KO and control samples. Right panel: Dot plots of enriched considerably Gene Ontology Biological Processes (GO: BP) for the top 200 upregulated and downregulated genes in CCB79-KO.

D. Left panel: Volcano plot illustrating differential gene expression, highlighting confirmed ciliopathy genes—right panel: Heatmap of the top 50 most variable ciliopathy genes, scaled by z-score. Columns represent samples grouped by control (purple) and CCB79-KO (green).

E. Categorical heatmap of HH-related gene groups (Ligands, Receptors, GLI factors, Ciliary components, Suppressors, Targets). Columns represent samples grouped by control (purple) and CCB79-KO (green).

F. Weighted HH signaling activity score computed by integrating activator, suppressor, and receptor gene expression z-scores shown as a boxplot comparing control and CCB79-KO.

G. Correlation heatmap showing Pearson correlation coefficients between HH signaling activity scores and forebrain developmental markers (FOXG1, EMX1, PAX6, TBR1, ARX, NKX2-1, and NEUROD6), color-coded from blue (positive) to red (negative) correlation.

Supplementary Figure 7 (related to the main Figure 4).

A. UMAP plots showing the distribution of various gene set markers used to annotate cell cluster types observed in control and CCB79-KO brain organoids.

Supplementary Figure 8 (related to the main Figure 4).

A. Dot plot showing expression of lineage- and region-specific markers across annotated cell types. Dot size represents the fraction of expressing cells; reversed viridis color intensity indicates the average scaled expression, ranging from 0 to 3.

B. Hierarchical clustering represented as heatmaps showing the top 5 differentially expressed genes per cell type identified in control (left) and CCB79-KO (right) conditions. Color intensity reflects scaled expression, with gene names and corresponding cell types annotated.

Supplementary Figure 9 (related to the main Figure 4).

A. The UMAP showing the spatial distribution and normalized expression levels of the selected HH pathway in control (left) and CCB79-KO (right) conditions reveals striking differences in distribution, particularly in the oligodendrocyte precursor-containing cluster (dotted circle). Color intensity (gray-to-brown colormap) represents a scaled expression of 0–5.

B. Matrix plots display the average expression (z-score) of HH pathway genes grouped by functional categories across annotated cell types. Rows represent HH pathway gene sets

(Ciliary Complex, GLI Factors, Ligands, Receptors, Suppressors, Targets), and columns represent cell clusters. Dendrograms display the hierarchical clustering of genes and cell types based on their expression similarity. Color scale ranges from low (light pink) to high (dark red) expression. The most affected cluster 9 is highlighted with a box.

Supplementary Figure 10 (related to the main Figure 4).
  A. UMAP embedding of all cells colored by 15 major developmental cell type clusters after sub-clustering in control (left) and CCB79-KO (right) organoids. Legends represent colors that correspond to annotated cell types.
  B. UMAP plots showing cell type-specific expression of selected markers used to assign cell types in control (left) and CCB79-KO (right).

Supplementary Figure 11 (related to the main Figure 4).
  A. Right panel, Force-directed graph (FA layout) showing combined developmental trajectories, control = green, CCB79-KO = purple. Left panel, annotated regions Hem (lavender), VZ (red), SVZ (orange), oSVZ (mustard yellow), IZ (beige), MZ (light pink), Migratory stream (pink), CP (sky blue), comparing control vs CCB79-KO.
  B. Stacked bar plot of proportional representation of cells across ordered cortical layers comparing control vs CCB79-KO.
  C. Dot plot quantifies the relative expression of markers (left side) in progenitors, precursors, and neurons between control and CCB-KO brain organoids.
  D. Classification of progenitor division modes using gene module scoring for symmetric and asymmetric division markers. The bar plot displays the stacked percentages for symmetric, asymmetric, and ambiguous cases. Each panel illustrates distinct aspects of cortical development and perturbations in CCB79-KO organoids, including regional mispatterning, altered layer composition, and division-type imbalance.

Supplementary Figure 12 (related to the main Figure 4)

Supplementary Movie 1 and Movie 2 (related to the main Figure 6)

References

Lui, J. H., Hansen, D. V. & Kriegstein, A. R. Development and evolution of the human neocortex. *Cell* **146**, 18-36 (2011). https://doi.org/10.1016/j.cell.2011.06.030

Melchionna, M. *et al.* Cortical areas associated to higher cognition drove primate brain evolution. *Commun Biol* **8**, 80 (2025). https://doi.org/10.1038/s42003-025-07505-1

Rakic, P. Specification of cerebral cortical areas. *Science* **241**, 170-176 (1988). https://doi.org/10.1126/science.3291116

Kriegstein, A. R. & Noctor, S. C. Patterns of neuronal migration in the embryonic cortex. *Trends in neurosciences* **27**, 392-399 (2004). https://doi.org/10.1016/j.tins.2004.05.001

Ball, G. *et al.* Molecular signatures of cortical expansion in the human foetal brain. *Nature communications* **15**, 9685 (2024). https://doi.org/10.1038/s41467-024-54034-2

Shao, W. *et al.* Centrosome anchoring regulates progenitor properties and cortical formation. *Nature* **580**, 106-112 (2020). https://doi.org/10.1038/s41586-020-2139-6

Nowakowski, T. J. *et al.* Spatiotemporal gene expression trajectories reveal developmental hierarchies of the human cortex. *Science* **358**, 1318-1323 (2017). https://doi.org/10.1126/science.aap8809

Bhaduri, A. *et al.* An atlas of cortical arealization identifies dynamic molecular signatures. *Nature* **598**, 200-204 (2021). https://doi.org/10.1038/s41586-021-03910-8

Pebworth, M. P., Ross, J., Andrews, M., Bhaduri, A. & Kriegstein, A. R. Human intermediate progenitor diversity during cortical development. *Proceedings of the National Academy of Sciences of the United States of America* **118** (2021). https://doi.org/10.1073/pnas.2019415118

Martínez-Cerdeño, V., Noctor, S. C. & Kriegstein, A. R. The role of intermediate progenitor cells in the evolutionary expansion of the cerebral cortex. *Cerebral cortex* **16 Suppl 1**, i152-161 (2006). https://doi.org/10.1093/cercor/bhk017

Florio, M. & Huttner, W. B. Neural progenitors, neurogenesis and the evolution of the neocortex. *Development* **141**, 2182-2194 (2014). https://doi.org/10.1242/dev.090571

Haubensak, W., Attardo, A., Denk, W. & Huttner, W. B. Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 3196-3201 (2004). https://doi.org/10.1073/pnas.0308600100

Shitamukai, A., Konno, D. & Matsuzaki, F. Oblique radial glial divisions in the developing mouse neocortex induce self-renewing progenitors outside the germinal zone that resemble primate outer subventricular zone progenitors. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 3683-3695 (2011). https://doi.org/10.1523/jneurosci.4773-10.2011

Reillo, I. & Borrell, V. Germinal Zones in the Developing Cerebral Cortex of Ferret: Ontogeny, Cell Cycle Kinetics, and Diversity of Progenitors. *Cerebral cortex* **22**, 2039-2054 (2011). https://doi.org/10.1093/cercor/bhr284

Smart, I. H. M., Dehay, C., Giroud, P., Berland, M. & Kennedy, H. Unique Morphological Features of the Proliferative Zones and Postmitotic Compartments of the Neural Epithelium Giving Rise to Striate and Extrastriate Cortex in the Monkey. *Cerebral cortex* **12**, 37-53 (2002). https://doi.org/10.1093/cercor/12.1.37

Kriegstein, A., Noctor, S. & Martínez-Cerdeño, V. Patterns of neural stem and progenitor cell division may underlie evolutionary cortical expansion. *Nature reviews. Neuroscience* **7**, 883-890 (2006). https://doi.org/10.1038/nrn2008

Gopalakrishnan, J. *et al.* Emerging principles of primary cilia dynamics in controlling tissue organization and function. *The EMBO journal* **42**, e113891 (2023). https://doi.org/10.15252/embj.2023113891

Anderson, R. G. The three-dimensional structure of the basal body from the rhesus monkey oviduct. *The Journal of cell biology* **54**, 246-265 (1972). https://doi.org/10.1083/jcb.54.2.246

Tanos, B. E. *et al.* Centriole distal appendages promote membrane docking, leading to cilia initiation. *Genes & development* **27**, 163-168 (2013). https://doi.org/10.1101/gad.207043.112

Kumar, D. *et al.* A ciliopathy complex builds distal appendages to initiate ciliogenesis. *The Journal of cell biology* **220** (2021). https://doi.org/10.1083/jcb.202011133

Chong, W. M. *et al.* Super-resolution microscopy reveals coupling between mammalian centriole subdistal appendages and distal appendages. *eLife* **9** (2020). https://doi.org/10.7554/eLife.53580

Bowler, M. *et al.* High-resolution characterization of centriole distal appendage morphology and dynamics by correlative STORM and electron microscopy. *Nature communications* **10**, 993 (2019). https://doi.org/10.1038/s41467-018-08216-4

Goetz, S. C. & Anderson, K. V. The primary cilium: a signalling centre during vertebrate development. *Nature reviews. Genetics* **11**, 331-344 (2010). https://doi.org/10.1038/nrg2774

Sasai, N., Toriyama, M. & Kondo, T. Hedgehog Signal and Genetic Disorders. *Frontiers in genetics* **10**, 1103 (2019). https://doi.org/10.3389/fgene.2019.01103

Bachmann-Gagescu, R. *et al.* Healthcare recommendations for Joubert syndrome. *American journal of medical genetics. Part A* **182**, 229-249 (2020). https://doi.org/10.1002/ajmg.a.61399

Reiter, J. F. & Leroux, M. R. Genes and molecular pathways underpinning ciliopathies. *Nature reviews. Molecular cell biology* **18**, 533-547 (2017). https://doi.org/10.1038/nrm.2017.60

Guemez-Gamboa, A., Coufal, N. G. & Gleeson, J. G. Primary cilia in the developing and mature brain. *Neuron* **82**, 511-521 (2014). https://doi.org/10.1016/j.neuron.2014.04.024

Sangermano, R. *et al.* Variants in the ciliopathy gene SCLT1 are associated with non-syndromic retinal degeneration. *Res Sq* (2025). https://doi.org/10.21203/rs.3.rs-6507107/v1

Gabriel, E. *et al.* CPAP promotes timely cilium disassembly to maintain neural progenitor pool. *The EMBO journal* **35**, 803-819 (2016). https://doi.org/10.15252/embj.201593679

Andreu-Cervera, A., Catala, M. & Schneider-Maunoury, S. Cilia, ciliopathies and hedgehog-related forebrain developmental disorders. *Neurobiology of disease* **150**, 105236 (2021). https://doi.org/10.1016/j.nbd.2020.105236

Goetz, S. C., Liem, K. F., Jr. & Anderson, K. V. The spinocerebellar ataxia-associated gene Tau tubulin kinase 2 controls the initiation of ciliogenesis. *Cell* **151**, 847-858 (2012). https://doi.org/10.1016/j.cell.2012.10.010

Jackson, P. K. TTBK2 kinase: linking primary cilia and cerebellar ataxias. *Cell* **151**, 697-699 (2012). https://doi.org/10.1016/j.cell.2012.10.027

Morozov, Y. M. & Rakic, P. Lateral expansion of the mammalian cerebral cortex is related to anchorage of centrosomes in apical neural progenitors. *Cerebral cortex* **34** (2024). https://doi.org/10.1093/cercor/bhae293

O'Neill, A. C. *et al.* Spatial centrosome proteome of human neural cells uncovers disease-relevant heterogeneity. *Science* **376**, eabf9088 (2022). https://doi.org/10.1126/science.abf9088

Gabriel, E. *et al.* Recent Zika Virus Isolates Induce Premature Differentiation of Neural Progenitors in Human Brain Organoids. *Cell stem cell* **20**, 397-406.e395 (2017). https://doi.org/10.1016/j.stem.2016.12.005

Goranci-Buzhala, G. *et al.* Rapid and Efficient Invasion Assay of Glioblastoma in Human Brain Organoids. *Cell reports* **31**, 107738 (2020). https://doi.org/10.1016/j.celrep.2020.107738

Ramani, A. *et al.* SARS-CoV-2 targets neurons of 3D human brain organoids. *The EMBO journal*, e106230 (2020). https://doi.org/10.15252/embj.2020106230

Wang, L. *et al.* Molecular and cellular dynamics of the developing human neocortex. *Nature* (2025). https://doi.org/10.1038/s41586-024-08351-7

Nomura, N. *et al.* Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Res* **1**, 27-35 (1994). https://doi.org/10.1093/dnares/1.1.27

Kikuno, R., Nagase, T., Waki, M. & Ohara, O. HUGE: a database for human large proteins identified in the Kazusa cDNA sequencing project. *Nucleic acids research* **30**, 166-168 (2002). https://doi.org/10.1093/nar/30.1.166

Ishikawa, K. *et al.* Prediction of the coding sequences of unidentified human genes. VIII. 78 new cDNA clones from brain which code for large proteins in vitro. *DNA Res* **4**, 307-313 (1997). https://doi.org/10.1093/dnares/4.5.307

Kanie, T. *et al.* A hierarchical pathway for assembly of the distal appendages that organize primary cilia. *eLife* **14** (2025). https://doi.org/10.7554/eLife.85999

Spektor, A., Tsang, W. Y., Khoo, D. & Dynlacht, B. D. Cep97 and CP110 suppress a cilia assembly program. *Cell* **130**, 678-690 (2007). https://doi.org/10.1016/j.cell.2007.06.027

Hsu, P. D., Lander, E. S. & Zhang, F. Development and applications of CRISPR-Cas9 for genome engineering. *Cell* **157**, 1262-1278 (2014). https://doi.org/10.1016/j.cell.2014.05.010

Huang, N. *et al.* Hierarchical assembly of centriole subdistal appendages via centrosome binding proteins CCDC120 and CCDC68. *Nature communications* **8**, 15057 (2017). https://doi.org/10.1038/ncomms15057

Kim, S. & Tsiokas, L. Cilia and cell cycle re-entry: more than a coincidence. *Cell cycle* **10**, 2683-2690 (2011). https://doi.org/10.4161/cc.10.16.17009

Sánchez, I. & Dynlacht, B. D. Cilium assembly and disassembly. *Nature cell biology* **18**, 711-717 (2016). https://doi.org/10.1038/ncb3370

Iyer, S. S. *et al.* Centriolar cap proteins CP110 and CPAP control slow elongation of microtubule plus ends. *The Journal of cell biology* **224** (2025). https://doi.org/10.1083/jcb.202406061

Liu, C., Wright, B., Allen-Vercoe, E., Gu, H. & Beiko, R. Phylogenetic Clustering of Genes Reveals Shared Evolutionary Trajectories and Putative Gene Functions. *Genome Biol Evol* **10**, 2255-2265 (2018). https://doi.org/10.1093/gbe/evy178

Shimada, I. S. *et al.* Derepression of sonic hedgehog signaling upon Gpr161 deletion unravels forebrain and ventricular abnormalities. *Developmental biology* **450**, 47-62 (2019). https://doi.org/10.1016/j.ydbio.2019.03.011

Mukhopadhyay, S. *et al.* The ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic hedgehog pathway via cAMP signaling. *Cell* **152**, 210-223 (2013). https://doi.org/10.1016/j.cell.2012.12.026

Schembs, L. *et al.* The ciliary gene INPP5E confers dorsal telencephalic identity to human cortical organoids by negatively regulating Sonic hedgehog signaling. *Cell reports* **39**, 110811 (2022). https://doi.org/10.1016/j.celrep.2022.110811

Bangs, F. & Anderson, K. V. Primary Cilia and Mammalian Hedgehog Signaling. *Cold Spring Harbor perspectives in biology* **9** (2017). https://doi.org/10.1101/cshperspect.a028175

Valente, E. M., Rosti, R. O., Gibbs, E. & Gleeson, J. G. Primary cilia in neurodevelopmental disorders. *Nat Rev Neurol* **10**, 27-36 (2014). https://doi.org/10.1038/nrneurol.2013.247

Danesin, C. *et al.* Ventral neural progenitors switch toward an oligodendroglial fate in response to increased Sonic hedgehog (Shh) activity: involvement of Sulfatase 1 in modulating Shh signaling in the ventral spinal cord. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **26**, 5037-5048 (2006). https://doi.org/10.1523/jneurosci.0715-06.2006

Camp, J. G. *et al.* Human cerebral organoids recapitulate gene expression programs of fetal neocortex development. *Proceedings of the National Academy of Sciences of the United States of America* **112**, 15672-15677 (2015). https://doi.org/10.1073/pnas.1520760112

Zhang, Z., O'Laughlin, R., Song, H. & Ming, G. L. Patterning of brain organoids derived from human pluripotent stem cells. *Current opinion in neurobiology* **74**, 102536 (2022). https://doi.org/10.1016/j.conb.2022.102536

Parvathy, S. *et al.* TLX3 regulates CGN progenitor proliferation during cerebellum development and its dysfunction can lead to autism. *iScience* **27**, 111260 (2024). https://doi.org/10.1016/j.isci.2024.111260

Butts, J. C. *et al.* A single-cell transcriptomic map of the developing Atoh1 lineage identifies neural fate decisions and neuronal diversity in the hindbrain. *Developmental cell* **59**, 2171-2188.e2177 (2024). https://doi.org/10.1016/j.devcel.2024.07.007

Yamada, M. *et al.* Specification of spatial identities of cerebellar neuron progenitors by ptf1a and atoh1 for proper production of GABAergic and glutamatergic neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **34**, 4786-4800 (2014). https://doi.org/10.1523/jneurosci.2722-13.2014

Trujillo, C. A. *et al.* Reintroduction of the archaic variant of NOVA1 in cortical organoids alters neurodevelopment. *Science* **371** (2021). https://doi.org/10.1126/science.aax2537

Benito-Kwiecinski, S. *et al.* An early cell shape transition drives evolutionary expansion of the human forebrain. *Cell* **184**, 2084-2102.e2019 (2021). https://doi.org/10.1016/j.cell.2021.02.050

Zhao, Z. *et al.* Super-Resolution Imaging of Highly Curved Membrane Structures in Giant Vesicles Encapsulating Molecular Condensates. *Advanced materials* **34**, e2106633 (2022). https://doi.org/10.1002/adma.202106633

Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014). https://doi.org/10.1093/bioinformatics/btt656

Liao, Y., Smyth, G. K. & Shi, W. The Subread aligner: fast, accurate and scalable read mapping by seed-and-vote. *Nucleic acids research* **41**, e108 (2013). https://doi.org/10.1093/nar/gkt214

Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology* **15**, 550 (2014). https://doi.org/10.1186/s13059-014-0550-8

Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics : a journal of integrative biology* **16**, 284-287 (2012). https://doi.org/10.1089/omi.2011.0118

Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-seq data. *BMC bioinformatics* **14**, 7 (2013). https://doi.org/10.1186/1471-2105-14-7

Hynes, A. M. *et al.* Murine Joubert syndrome reveals Hedgehog signaling defects as a potential therapeutic target for nephronophthisis. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 9893-9898 (2014). https://doi.org/10.1073/pnas.1322373111

Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 15545-15550 (2005). https://doi.org/10.1073/pnas.0506580102

Yin, W. C. *et al.* Dual Regulatory Functions of SUFU and Targetome of GLI2 in SHH Subgroup Medulloblastoma. *Developmental cell* **48**, 167-183.e165 (2019). https://doi.org/10.1016/j.devcel.2018.11.015

Hui, C. C. & Angers, S. Gli proteins in development and disease. *Annu Rev Cell Dev Biol* **27**, 513-537 (2011). https://doi.org/10.1146/annurev-cellbio-092910-154048

Ingham, P. W. & McMahon, A. P. Hedgehog signaling in animal development: paradigms and principles. *Genes & development* **15**, 3059-3087 (2001). https://doi.org/10.1101/gad.938601

Rohatgi, R., Milenkovic, L. & Scott, M. P. Patched1 regulates hedgehog signaling at the primary cilium. *Science* **317**, 372-376 (2007). https://doi.org/10.1126/science.1139740

Chiang, C. *et al.* Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* **383**, 407-413 (1996). https://doi.org/10.1038/383407a0

Kohtz, J. D., Baker, D. P., Corte, G. & Fishell, G. Regionalization within the mammalian telencephalon is mediated by changes in responsiveness to Sonic Hedgehog. *Development* **125**, 5079-5089 (1998). https://doi.org/10.1242/dev.125.24.5079

Rallu, M., Corbin, J. G. & Fishell, G. Parsing the prosencephalon. *Nature reviews. Neuroscience* **3**, 943-951 (2002). https://doi.org/10.1038/nrn989

Fuccillo, M., Joyner, A. L. & Fishell, G. Morphogen to mitogen: the multiple roles of hedgehog signalling in vertebrate neural development. *Nature reviews. Neuroscience* **7**, 772-783 (2006). https://doi.org/10.1038/nrn1990

Fig. 1

Fig. 2

Fig 3

Fig. 4

A

B

- (SC1) Cycling RG
- (SC2) Quiescent / Apical RG
- (SC3) Dorsal VZ RG
- (SC4) Outer RG
- (SC5) Transitional oRG
- (SC6) IPC (NPAS3+)
- (SC7) IPC (INSM1+)
- (SC8) Neuroblasts
- (SC9) Immature excitatory neurons
- (SC10) Maturing excitatory neurons
- (SC11) Immature GABAergic INs
- (SC12) Cajal-Retzius neurons
- (SC13) Reactive / IEG RG
- (SC14) Ventral RG
- (SC15) Roof-plate / neural-crest-like

- (C1) Neural tube (*WNT8B, RSPO2*)
- (C2) Neural crest (*PAX3,COLEC12*)
- (C3) Telencephalic NPC (*SOX2,FOXG1*)
- (C4) Non-telencephalic NPC (*PAX3,HOXB2*)
- (C5) Proliferative NPC (*MKI67, AURKB*)
- (C6) Telencephalic immature Neuron (*STMN2,DLX5,GAD1*)
- (C7) Non telencephalic immature Neuron (*STMN2,LHX1*)
- (C8) Mesenchymal/Meningeal (*DCN,TWIST1, MEIS2*)
- (C9) Oligodendrocyte lineage (*PLP1, SOX10*)
- (C10) Microglia (Ambiguous) (*AIF1*)

C

D

E

Fig. 5

Fig. 6

Fig. S1 (related to main figure 1)

A CCB79 (KIAA0408)

B CCB79 (KIAA0408) in cell types

C Developmental stages of the human embryo

D

E

Fig. S2 (related to main figure 1)

A

KIAA0408

WT   TTAATGGGGCTAAATCCTGTTTCACTTAAATATTCCTGCTTAGTGTTTCTACCAGACT
KIAA0408-KO   TTAATGGGGCTA..........deleted region by CRSPR..........TCTACCAGACT
                                        Δ35

WT

KIAA0408 KO

Deleted region

Predicted mRNA transcript and peptide sequence

612bp

WT   TCCCCAAATTACCCCGATTTAGGACAAAGTGAATTTATAAGGACGAATCAC
KIAA0408 KO   TCCCCAAATTACCCCGATAGATGGTCTGAGAAAAGAAAATAAAAGAGAGC
                                                              *(stop codon)

        80   90   100   110
WT   S S P N Y P D L G Q S E F I R T N H K D G L R K E N K R E Q S L V
KIAA0408 KO   S S P N Y P ----------------------------------- E R W S E K R K *(stop codon)

1kb plus ladder | WT | KIAA0408-KO

612bp
577bp

Bi Control NPCs

ARL13B/γ-Tub

ARL13B/γ-Tub

Merge | ARL13B
γ-Tub

20µm

Bii KIAA0408-KO NPCs

ARL13B/γ-Tub

20µm

Cilia as a dot
ARL13B/γ-Tub

Merge | ARL13B
γ-Tub

Bulgy short Cilia
ARL13B/γ-Tub

5µm

Merge | ARL13B
γ-Tub

Thin long cilia

5µm

Merge | ARL13B
γ-Tub

C

n=336 Control   n=334 CCB79-KO

<0.0001   <0.0001
<0.0001   <0.0001

Aberrant cilia (%)

Cilia as a dot n=108
Bulgy short Cilia n=102
Thin long cilia n=102

Cells (%)

Centrosomes/cell
2   >2   <2   2   >2   <2

Fig. S3 (related to main figure 2)

Fig S4 (related to main figure 3)

Fig. S5

Fig. S6

Fig S7 (related to main figure 4)

A

Fig S8 (related to main figure 4)

Fig. S9 (related to main figure 4)

Fig. S10 (related to main figure 4)

Control

CCB79-KO

- (SC1) Cycling RG
- (SC2) Quiescent / Apical RG
- (SC3) Dorsal VZ RG
- (SC4) Outer RG
- (SC5) Transitional oRG
- (SC6) IPC (NPAS3+)
- (SC7) IPC (INSM1+)
- (SC8) Neuroblasts
- (SC9) Immature excitatory neurons
- (SC10) Maturing excitatory neurons
- (SC11) Immature GABAergic INs
- (SC12) Cajal-Retzius neurons
- (SC13) Reactive RG
- (SC14) Ventral RG
- (SC15) Roof-plate / neural-crest-like

Control

CCB79-KO

Fig. S11 (related to main figure 4)

Fig. S12

Control

CCB79-KO

MZ
ULN
CP
IZ
SVZ/oSVZ
VZ

RGC

NEC

IPC

oRG

DLN

EN

IPC-D

IN

IPC-V

EN

IPC-D*

IPC-V

Neuroepithelial cells(NEC)

Radial glia cells (RG)

Outer radial glia cells(oRG)

Intermediate progenitor - Dorsal (IPC-D)

Intermediate progenitor - Ventral (IPC-V)

Atypical intermediate progenitor - Dorsal

Atypical Excitatory neuron- Dorsal

Excitatory neuron (EN)

Inhibitory neuron (IN)

Interneuron

Upper layer neuron (ULN)

Deep layer neuron (DLN)

Cajal-Retzius

Primary cilia