Title: Long-term forest-sector mitigation and radiative forcing under contrasting

management, climate, and substitution pathways

I. Boukhris[1,2]*, F. Cherubini[3], A. Collalti[4,5], D. Dalmonech[4,5], C. Vonderach[6],

G. Marano[7], F. Giannetti[8], S. Lahssini[9], M. Santini[2], R. Valentini[1,2]

1. Department for Innovation in Biological, Agri-Food and Forest Systems (DIBAF),

University of Tuscia, 01100, Viterbo, Italy (issamboukhris@gmail.com; rik@unitus.it)

2. CMCC Foundation — Euro-Mediterranean Center on Climate Change, Lecce, Italy

(monia.santini@cmcc.it)

3. Industrial Ecology Programme, Department of Energy and Process Engineering, Norwegian

University of Science and Technology (NTNU), 7034 Trandheim, Norway

(francesco.cherubini.ntnu.no)

4. Forest modeling Lab, Institute for Agriculture and Forest Systems in the Mediterranean,

National Research Council of Italy (CNR-ISAFOM), 06128, Perugia, Italy

(alessio.collalti@cnr.it; daniela.dalmonech@cnr.it)

5. National Biodiversity Future Center (NBFC), 90133, Palermo, Italy

6. Forest Research Institute Baden-Wurttemberg, 79100, Freiburg, Germany

(christian.vonderach@forst.bwl.de)

7. Swiss Federal Institute for Forest, Snow and Landscape Research, WSL, 8903 Birmensdorf,

Switzerland (gina.marano@wsl.ch)

8. Department of Agriculture, Food, Environment and Forestry (DAGRI), University of

Florence, Florence, Italy

9. Department of Forest Development, National School of Forestry Engineering, 11000, Salé,

Morocco (marghadi@gmail.com)

*issamboukhris@gmail.com

phone: +39 327 328 2507

Abstract

Forests are central to the EU's climate neutrality strategy, currently offsetting ~9% of total

greenhouse gas emissions and offering further mitigation potential through harvested wood

products and the substitution effect. However, the climate benefit of the forest sector is

influenced by multiple interacting factors, including forest management, climate change,

wood-use strategies, and assumptions about substitution benefits, as well as the timing and fate

of carbon across the forest ecosystem and technosphere. To evaluate these drivers, we used a

coupled forest growth and a wood products model to simulate five different silvicultural

strategies under three climate change scenarios, four wood use schemes, and five displacement

factor decay pathways over a 285-year period (2015–2300), applied to a *Pinus nigra* forest in

Italy, questioning the impact of these factors on climate mitigation potential of the forest sector.

We assessed forest sector balance (FSB, net carbon exchange between forest system and

atmosphere), radiative forcing from biogenic CO_2 (RF_{bio}), and mitigation efficiency (ME) –

the proportion of sequestered carbon contributing to net climate benefit. Results showed that

FSB and RF_{bio} were "broadly" aligned, but ME varied with the magnitude and duration of

biogenic emissions. The scenarios BIOE (bioenergy) and TM (modular cutting) achieved high

FSB but showed lower ME due to concentrated or sustained emissions. WOOD (promotion of

long-lived wood) and ADAPT (adaptation management) yielded higher ME under SSP1-2.6,

while several strategies (WOOD, ADAPT, TRANS) became net sources under SSP5-8.5 after

2200. Substitution benefits declined under degressive assumptions, reducing mitigation by up

to 53% especially for high-harvest scenarios. FSB was primarily shaped by climate and

management, secondly by substitution, however; wood-use strategies had no significant long-

term effect provided they did not impact resource availability. Together, these findings

underscore that effective forest-sector mitigation requires not only maximizing cumulative

carbon stocks, but also minimizing the magnitude, timing, and atmospheric residence time of

emissions while carefully considering the role of substitution benefits.

Keywords

Forest management, Climate mitigation, Wood substitution, Carbon balance, Mitigation

efficiency, Climate change, Process-based model, Wood products model

Highlights

• The long-term forest sector carbon balance is mainly governed by active management,

climate conditions, and wood substitution pathways.

• Forest Sector Balance as a metric is "broadly" aligned with the radiative forcing from

biogenic emissions

• The mitigation efficiency of forest sector options depends on emissions timing, duration,

and amplitude

• Substitution factors also known as displacement factors need to be considered with greater

caution

1. Introduction

Forests are key to the EU's climate mitigation framework for achieving carbon neutrality by 2050. Currently they remove approximately 9% of total greenhouse gas (GHG) emissions from the atmosphere (EEA, 2022). Forest ecosystems serve as both carbon sinks, sequestering additional carbon, and reservoirs, storing it across several pools. When managed sustainably, forest carbon can be transferred laterally to harvested wood products (HWPs), establishing an external carbon pool and contributing to the substitution effect, as HWPs replace energy-intensive materials and fossil fuels, avoiding emissions (Rüter and Fulvio, 2025). In the EU, forests and HWPs together sequester approximately 400 Mt CO_2-eq.yr^{-1}, with HWPs contributing for about 10% to this total. Substitution provides a similar level of contribution with values ranging from 18 to 43 Mt CO_2-eq.yr^{-1}, further enhancing the forest sector's overall mitigation potential (Jonsson et al., 2021; Nabuurs et al., 2017; Rüter S. et al., 2016).To assess forest sector mitigation, a system-wide perspective is needed to account for the distinct roles of carbon pools. This approach ensures that all relevant mitigation pathways, sequestration, storage, and substitution, are critically evaluated (Grassi et al., 2021; Lemprière et al., 2013). Forest-sector mitigation outcomes are shaped by multiple interacting factors. Understanding these drivers is essential to accurately assess the effectiveness and trade-offs among strategies. The forest carbon sink is primarily determined by both forest management and climate conditions (Ameray et al., 2021; Dye et al., 2024; Gregor et al., 2024; Liu et al., 2025). Similarly, the HWPs sink is affected by these factors but also depends on wood-use characteristics and the carbon footprint of the displaced material, which ultimately determines the substitution effect (Bozzolan et al., 2024; Brunet-Navarro et al., 2017; Budzinski et al., 2020). Beyond biophysical and technological factors, temporal aspects of carbon fluxes also play a critical role.

Among these factors, forest management has received particular attention (Valentini and

Miglietta, 2015). Practices range from passive conservation to intensive silvicultural

interventions (Barredo et al., 2024; Dalmonech et al., 2022; Pilli et al., 2022). An ongoing

debate in the EU revolves on whether active or reduced management contributes more

effectively to climate mitigation (Luyssaert et al., 2007; Noormets et al., 2015). In managed

forests, early interventions such as tending and thinning are used to improve wood quality,

promote tree growth, and reduce the likelihood of natural mortality. As a result, more wood

can be harvested for use in HWPs instead of being lost to natural decay. At the same time, these

operations enhance forest resilience to disturbances, which are becoming more frequent due to

climate change (Barrette et al., 2023; Kilpeläinen and Peltola, 2022; Moreau et al., 2022). In

the long term, while unmanaged forests tend to maintain higher carbon stocks, especially in

the dead wood pool (Nagel et al., 2023), studies suggest that managed forests can achieve a

superior overall carbon balance when accounting for the substitution effect (Krug et al., 2012;

Schulze et al., 2020).

Differences in the mitigation potential of multiple active management strategies exist and are

primarily driven by factors such as structure complexity, rotation length, thinning intensity,

and thinning intervals. In parallel, at least two key factors influence the HWPs sink: the

recycling rate and the lifespan of HWPs, both of which delay carbon emissions to the

atmosphere through combustion or decomposition in disposal sites (Brunet-Navarro et al.,

2017; Pingoud et al., 2003; Wei et al., 2023). Finally, one of the most critical assumptions in

estimating the forest sector's substitution effect is the assumption of stationary displacement

factors (DF). Assuming fixed DF fails to account for potential improvements in energy

efficiency and reductions in the carbon footprint of competing sectors (e.g., steel, concrete, and

energy) that may lead to reductions in GHG emissions. Any advancements in these sectors

would consequently diminish the substitution effect of wood (Brunet-Navarro et al., 2021;

Harmon, 2019; Howard et al., 2021). Therefore, assessing alternative strategies with a fixed

DF reflects only the maximum substitution potential, while the actual effect is likely lower and

often overestimated in global reports.

A critical yet often overlooked aspect of climate mitigation assessments is the timing of carbon

fluxes and how this aligns with the time horizon of mitigation projects. As Levasseur (2010)

highlighted, "*carbon neutrality does not equate climate neutrality*", as carbon persists in the

atmosphere before being reabsorbed, contributing in the meantime to the climate forcing

(Levasseur et al., 2010). Forest-Technosphere systems that release carbon rapidly but sequester

it slowly can have a stronger short-term climate impact than those with faster sequestration

rates and slower carbon release (Holtsmark, 2012; Laganière et al., 2017). These short-term

emissions influence radiative forcing, potentially intensifying near-term warming and affecting

long-term climate stability. Similarly, the time horizon, the timeframe over which forest

systems are evaluated, significantly shapes their assessed climate impact. Over short time

frames, immediate emissions from harvesting may outweigh sequestration benefits, resulting

in a higher net climate impact. In contrast, longer horizons allow forest regrowth along with

the other climate system components to offset initial emissions, improving mitigation

outcomes (Cherubini et al., 2011; Holtsmark, 2013; Sierra et al., 2021). Given this, assessing

the mitigation potential of forests based on carbon balances alone is insufficient; it must also

account for their contribution to radiative forcing over time. To capture these temporal

dynamics, we first quantified climate impact using a timing-sensitive indicator, RF_{bio}, which

integrates the net carbon fluxes over time through a global impulse response function.

However, while RF_{bio} describes the magnitude of climate impact, it does not indicate how

efficiently carbon is processed to achieve that outcome. To address this limitation, we introduce

the concept of mitigation efficiency (ME), a metric reflecting the climate benefit delivered per

unit of carbon cycled through the forest sector. Together, RF_{bio} and ME offer a more nuanced

basis for evaluating forest-sector strategies, emphasizing not just how much carbon is removed,

but how effectively and durably that removal contributes to climate mitigation.

Despite the well-established role of forests and HWPs in climate mitigation, significant

uncertainties remain regarding the long-term effectiveness of different forest management

strategies. These uncertainties arise from the complex interactions among forest growth,

harvest intensity, product use, and substitution dynamics, as well as from simplifying

assumptions, such as stationary DF. Moreover, most studies focus on short-term carbon

balances, overlooking the long temporal cycles of forest systems and the fact that mitigation

potential may emerge at different timescales depending on the applied management (Metzler

et al., 2024; Skytt et al., 2021).

The main aim of this study is to quantify the long-term climate-mitigation performance of the

forest sector by jointly accounting for forest carbon dynamics, HWPs, and time-dependent

substitution effects under climate change. To achieve this, we compare five forest management

strategies under three climate change scenarios over a 285-year period, evaluate how

alternative displacement pathways and the decay of displacement factors influence mitigation

outcomes, assess whether high carbon balances consistently correspond to stronger climate

benefits across different time horizons, and determine how the mitigation efficiency of the

forest-sector options emerges and evolves over time.

We addressed these questions using an integrated modeling framework coupling a process-

based forest model with a wood product model (WPM) in a factorial experimental design. The

framework was applied to a *Pinus nigra* J.F. Arnold stand in the Vallombrosa Forest (Tuscany,

Italy), representative of temperate European forests. *P. nigra* was selected due to its wide

ecological adaptability, increasing use in European reforestation programs, and relevance for

climate-resilient forestry (EU council, 2025; Vacek et al., 2023).

2. Materials and Methods

2.1. Study area

The study was conducted in the state forest of Vallombrosa (43° 44' N, 11° 34'), a biogenetic

reserve of approximately 1000 ha in Tuscany, Italy. The forest experiences a mean annual

precipitation of about 1195 mm and a mean annual temperature slightly higher than 10 °C.

January is the coldest month (1.9 °C), July the warmest (19.6 °C). All the conifer plantations

were traditionally managed as even-aged. *Pinus nigra* was introduced in the forest (Bottalico

et al., 2012) as part of the main reforestation program that occurred after the second World

War. These stands cover 11% of the total forest area, with ages ranging from 60 to 95 years.

From 2005, as reported in the Vallombrosa forest management plan (*Piano di gestione e*

silvomuseo) (Travaglini, 2009). *Pinus nigra* stands are managed under close-to-nature

silviculture principles following the systemic silviculture approach (Ciancio and Nocentini,

2011), with the principles of modular/selective continuous cutting. This approach rooted in

Continuous Cover Forestry (CCF) and prescribed by Ciancio (2004) promotes resource

sustainability and natural regeneration by dynamically reassessing the harvestable volume

(Ciancio and Nocentini, 2004). The silvicultural systemic method for growing stock harvesting

calculates the real wood provision over a 5-year period. Thinning is allowed only if the

remaining standing volume is kept above the minimum threshold (typically 100-150 m^3 ha^{-1}

for *Pinus nigra*). If the condition is not met, only the portion of increment that preserves the

minimum stock is thinned. Thinning is also carried out to remove trees at high risk of falling.

When harvesting is conducted, the resulting residues are left on site. To monitor forest yield

and health, and in the context of the forest management plan, local permanent forest inventory

plots were established. The permanent circular plots (530 m²) are identical to those of the

Italian National Forest Inventory (NFI; https://www.inventarioforestale.org) and were

surveyed in 2005. Several of these plots are located within *Pinus nigra* stands and were

resurveyed again in 2017, 2018, and 2023.

2.2. Study key silvicultural and modeling assumptions

The following assumptions are considered in this study: the selected forest stand is assumed to

be representative of the average conditions across the broader *Pinus nigra* landscape. The same

forest management practices and wood use behavior are consistently applied/maintained

throughout the study period. Additionally, species composition is assumed to remain

unchanged, meaning no species conversion occurs over the time horizon. Finally, each

management scenario assumes a starting point of a recent plantation established in 2015

following a clear-cut, with a planting density of 2500 seedlings per hectare. Tree dimensions

at establishment followed species-specific default parameters of the growth model (initial

height = 1.3 m). Biomass and soil carbon pools were initialized using a 2500-year spin-up to

equilibrium under local climate conditions prior to the start of simulations. All management

scenarios began from identical stand conditions, ensuring that the differences among scenarios

arise solely from management, climate forcing, and wood-use assumptions.

In addition, landfilling activities were assumed to gradually phase out by the year 2060,

following an exponential decay model calibrated on national data. This estimate is based on

the observed reduction in the share of Italian municipalities utilizing landfill, which declined

from 46% in 2010 to 18% in 2022 (EEA, 2024). By assuming a constant relative rate of

decrease, landfilling is projected to approach zero use by 2060, and this cutoff year is adopted

as the upper limit for landfill emissions modeling.

2.3. Definition of scenarios

2.3.1. Climate change pathways

We investigated two coupled representative concentration pathways (RCPs) and shared

socioeconomic pathways (SSPs) based on distinct emission scenarios to assess potential future

climate trajectories and their socioeconomic implications. The SSP1-RCP2.6 scenario (SSP1-

2.6 hereafter) represents a world of sustainability-focused growth and equality, and it assumes

that strong climate protection measures are implemented, aiming to limit global warming to

well below 2°C above pre-industrial levels (O'Neill et al., 2014; Riahi et al., 2017; van Vuuren

et al., 2011). In contrast, SSP5-RCP8.5 (SSP5-8.5 hereafter) reflects a world of rapid and

unconstrained growth in economic output, energy demand and use, as well as limited to no

mitigation actions and a persistent reliance on fossil fuels (Calvin et al., 2017; Meinshausen et

al., 2011; Rogelj et al., 2018). Climate projections were obtained from the Coupled Model

Intercomparison (CMIP 6) project (Eyring et al., 2016), covering the period from 2015 to 2300.

Data were sourced from the MRI-ESM2-0 global climate model, as it was the only model

available that simulated the variables of interest over the long term, extending to the year 2300.

The outputs were downscaled to a 0.25° resolution using the perfect prognosis downscaling

method. A set of statistical models and algorithms was initially calibrated using historical

(observed) data from both coarse predictors (ERA5 reanalysis; (Hersbach et al., 2020) and local

predictand data (E-OBS reanalysis, including precipitation, maximum, and minimum

temperature; Cornes et al., 2018) over a representative 30-year climate period (1991–2020).

The methods were then validated and evaluated, with the best-performing models selected to

downscale daily precipitation, maximum, and minimum daily temperatures. Moreover,

atmospheric CO_2 concentrations for different climate trajectories were obtained from

(Meinshausen et al., 2020). Additionally, we simulated the current climate (CUR) for the

period 2020–2300 by randomly sampling each day of the year from the local station records

(Vallombrosa, locality of Reggello) over the period 2005–2023 (see **Figure 1**).

Figure 1: Climate projections from the downscaled MRI-ESM2-0 model under the SSP1-2.6 and SSP5-8.5 scenarios, driving the simulation for the period 2015–2300. The top panel shows annual precipitation (mm), the middle panel displays the annual mean of daily maximum and minimum temperatures (°C), and the bottom panel shows atmospheric CO_2 concentration (ppm), labeled as Ca.

2.3.2. Forest management scenarios

Five management scenarios were developed, grounded in five policies outlined in the EU Forest Strategy (European Commision, 2021), the EU Bioeconomy Strategy (European Commision, 2018), and the EU Forest Biodiversity Strategy (European Commission, 2021). These scenarios were designed to reflect the specific characteristics of the target species and the study area. In this process, the regulations for *P. nigra* management in the Tuscany region were also considered (refer to (Cantiani P et al., 2018) for further details). Each policy was

subsequently translated into stand-based forest management plans with the expert input of local

stakeholders, including forest managers and academic scholars.

The silvicultural interventions applied across the scenarios include pre-thinning and thinning

operations, implemented with varying intensities and intervals. In shelterwood-based

scenarios, additional interventions such as preparatory cuts and seedling cuts are employed to

facilitate natural regeneration under partial canopy cover. Final cuts are applied in both

shelterwood and clearcut systems, serving to remove the remaining overstory once

regeneration is established in shelterwood, or to harvest the entire stand at once in clearcutting

scenarios. In the latter case, artificial regeneration follows harvest operations, with planting

carried out at a density of 2500 seedlings per hectare (See **Figure 2** for detailed silvicultural

intervention schedules).

The first scenario is known as '*Tagli Modulari*' which represents modular cutting (hereafter

TM), and reflecting the typical practices applied in the Vallombrosa forest and described in

detail in Section 1. The "Wood-Based Bioenergy Production" (BIOE) assumes that the EU is

transitioning to a more sustainable energy future by gradually promoting the use of wood

bioenergy as a replacement for fossil fuel energy. While direct use of wood logs for energy is

discouraged under the EU's Renewable Energy Directive (RED II and RED III), only the

portion of trees unsuitable for higher-value uses are directly allocated to bioenergy. Practically,

this management plan consists of a short rotation (70-80 yrs) and the application of 1-3 thinning

treatments. The "Promotion of long-lived HWPs" (WOOD) explores how forest management

can support climate mitigation by increasing the supply of high-quality timber suitable for

durable HWPs. The scenario assumes that domestic wood use in the EU shifts increasingly

toward long-lived applications such as construction materials, which can store carbon for

extended periods and substitute for more carbon-intensive materials. While the lifespan of

HWPs also depends on post-harvest factors, such as recycling systems and end-use, this

scenario emphasizes the forest management component that facilitates the availability of

timber suitable for such uses. On the forest side, this objective is supported by extending

rotation lengths (e.g., 120-150 yrs) aiming to harvest trees with larger diameters and higher

wood quality through stronger thinning. The "Enhancing Forest Adaptation and Resilience"

(ADAPT) aims to strengthen the resilience of forests, particularly Mediterranean forests, which

are already experiencing severe drought conditions. Practically, this management plan will

consist of applying intensive thinning to alleviate competition on site resources and adopting

short rotation cycles to alleviate disturbance risks, in addition to promoting artificial

regeneration. The last scenario, "Transition to passive rewilding" (TRANS) is designed to

facilitate the shift from structured stand management toward a self-regulating forest system

with minimal long-term disturbance. Initial interventions are intended to accelerate natural

processes such as gap creation, early mortality, and canopy opening, which commonly occur

in the late stages of forest succession and structural diversification (Bauhus et al., 2009). These

actions aim to promote structural diversification and regeneration. Thinning is applied at

regular intervals between 2030 and 2150, with intensities ranging from 30% to 50% of the

stand basal area. After 2150, no further interventions were applied, allowing the forest to

develop autonomously under passive conditions.

Figure 2: Overview of the five forest management strategies evaluated in this study, arranged along a gradient of management intensity from high (top) to low (bottom). Strategies include: BIOE, TM, WOOD, ADAPT, and TRANS. Each strategy is defined by its objective, silvicultural approach, and target stand characteristics, progressing from strategic objectives to operational implementation. The right-hand panels show the corresponding management plans for each strategy, including timing and type of intervention (i.e., Prth: pre-thinning, Thin: thinning, Prep: preparatory cut, SeedC: Seeding cut, and FinalC: final cut), percentage of trees removed (%N), percentage of basal area removed (%BA) scheduled at specific ages.

2.3.3. Wood use scenarios

To explore the potential implications of improved cascading through an increase in product

lifespan and recycling rates, four different wood use scenarios were developed. The first, the

BAU as "Baseline Scenario", maintained constant recycling rate and lifespan values. In the

LS20 as "Longevity Scenario", the lifespan of products was increased by 20%. The RR20 as

"Reuse Scenario" focused on increasing the recycling rate of products by 20%. Finally, the

RRLS20 "Sustainability Scenario" increased both the lifespan of products and the recycling

rate by 20%. In BAU, the recycling rate of wood products was fixed at 10% through the

simulation period while product lifespans were parametrized from published studies (please

refer to **Table S1.5**).

2.3.4. Substitution scenarios

In addition to a baseline scenario (SUB0), which assumes that the DF remains constant over

time, we considered four decarbonization pathways with varying linear rates of DF reduction.

We adopt either a gradual (25% DF reduction by 2050, called SUB25), moderate (50%,

SUB50), significant (75%, SUB75) or a rapid (100%, SUB100), the latter being in line with

the EU's climate neutrality targets. DF values were parametrized based on literature sources

(Sathre and O'Connor, 2010; Suter et al., 2017) (please refer to Table S1.6).

2.4. Modeling framework

2.4.1. Modelling system and key concepts

The system considered in this study includes both forest and HWPs pools, collectively

representing the forest sector. It accounts for incoming and outgoing carbon fluxes while

disregarding internal exchanges, as these remain within the system and do not affect its net

balance. The forest sector system consists of five key components: (1) Aboveground biomass,

comprising stem wood, branches, stumps, and foliage of live trees, where the focus is on CO_2 fixation through photosynthesis and biomass accumulation; (2) Belowground biomass, including fine and coarse roots of live trees, also assessed for CO_2 fixation; (3) Aboveground Dead Organic Matter (ADOM), consisting of standing dead trees, downed woody debris, litter, and humus, where carbon is gradually released through decomposition; (4) Belowground Dead Organic Matter (BDOM), encompassing soil organic matter and dead roots, with carbon emissions primarily occurring through microbial respiration; and (5) HWPs, representing carbon stored in HWPs during their use phase, with CO_2 emissions occurring over time through disposal in landfills or combustion for bioenergy (see **Figure 3**). Beyond these carbon pools, the substitution effect plays a critical role in the forest sector's overall climate mitigation.

The annual forest ecosystem carbon balance (FEB; tC ha^{-1} yr^{-1}) is defined as:

$$FEB = (GPP - Ra - Rh_{onsite}) - Harvest = GPP - R_{ECO} - Harvest = -NEE - Harvest \qquad (Eq.\ 1)$$

where GPP represents the gross primary production; Ra, Rh_{onsite}, and R_{ECO} correspond to autotrophic, heterotrophic, and ecosystem respirations occurring within the forest, respectively, Harvest accounts for biomass removal through forestry operations, and NEE represents the net ecosystem exchange within the forest. All terms are expressed in units of tC ha^{-1} yr^{-1}.

Harvested biomass enters the wood sector, where it is temporarily stored in HWPs before eventually being released as emissions through disposal, or combustion for bioenergy. Therefore, the wood sector carbon balance (WSB; tC ha^{-1} yr^{-1}) is expressed as:

$$WSB = Harvest + Substitution - Rh_{offsite} = Harvest + Wbalance \qquad (Eq.\ 2)$$

where $Substitution$ represents avoided emissions resulting from the replacement of fossil fuels and energy-intensive materials with HWPs, $Rh_{offsite}$ (tC ha^{-1} yr^{-1}) refers to the heterotrophic respiration occurring outside the forest, mainly from HWPs decomposition and combustion, and $Wbalance$ refers to the balance between $Substitution$ and $Rh_{offsite}$, both of which are expressed in units of tC ha^{-1} yr^{-1}.

368   The annual overall forest sector balance (FSB; tC ha^{-1} yr^{-1}) integrates both forest and wood

369   sector dynamics, incorporating the mitigation benefits of substitution:

$$FSB = FEB + WSB = -NEE - Harvest + Harvest + Wbalance = Wbalance - NEE \quad \text{(Eq. 3)}$$

The terms Rh_{onsite}, and $Rh_{offsite}$ follow the terminology established by (Schulze et al., 2021).

Figure 3: Conceptual representation of carbon flows and balances with the forest sector,

divided into the forest ecosystem (green box) and wood sector (brown box). Gross Primary

Productivity (GPP) constitutes the carbon inflow to the forest ecosystem while both Ra and

Rh$_{onsite}$ constitute the carbon outflow from the forest ecosystem. Harvested biomass is

transferred from the forest ecosystem to the wood sector where it contributes to substitution

benefits and eventually to Rh$_{offsite}$.

2.4.2. Modelling framework and simulation setup

A forest sector modeling framework was developed by coupling 3D-CMCC-FEM (v5.7 BGC)

(Collalti et al., 2014; Dalmonech et al., 2024; Puchi et al., 2026; Saponaro et al., 2025; Vangi

et al., 2026), a stand-level, climate-sensitive, process-based model, with TimberTracer

(Boukhris et al., 2025) a WPM that tracks carbon stocks in HWPs, substitution effects, and

emissions from disposal sites (see **Figure 4**). The choice of 3D-CMCC-FEM was motivated

by its suitability for long-term climate-mitigation analyses. The model explicitly represents

photosynthesis, respiration, and carbon allocation, allowing sensitivity to changing climate

conditions and capturing climate-carbon feedback affecting growth and mortality. Its process-

based stress-response formulation enables realistic simulation of drought induced decline,

while the structural representation of stand dynamics provides a mechanistic description of

mortality and competition at the species scale. At the same time, the model remains

computationally efficient, which is essential for factorial multi-century simulations and

sensitivity analyses. Its relatively transparent parametrization further facilitates uncertainty

assessment and interpretation. Complementarily, TimberTracer was selected to represent the

technosphere component of the forest sector. Based on a material-flow approach, the model

simulates the temporal dynamics of carbon storage and emissions across all HWP pools and

disposal pathways, while explicitly accounting for material and energy substitution effects. Its

compatibility stand-level inputs allow seamless coupling with forest growth models such as

3D-CMCC-FEM, enabling consistent carbon accounting from forest ecosystem processes to

post-harvest product use. The modular structure of the model further supports scenario analysis

and sensitivity testing, making it suitable for evaluating long-term mitigation pathways.

Given the complexity and level of detail associated with each model, including their structure,

parametrization, input data, and validation procedures, full technical descriptions have been

moved to the Supplementary Materials (**Tables S1.1-S1.7**) to maintain clarity and focus on the

main text.

Figure 4: Carbon modeling framework coupling a forest process-based model (3D-CMCC-

FEM) and TimberTracer, a WPM. DBH is the mean diameter at breast height, N is the stand

density (ha^{-1}), BA is the basal area (m^2 ha^{-1}), H is the mean tree height (m), SA is sapwood area

of the mean tree (cm^2), HA is the heartwood area of the mean tree (cm^2), and BT is the bark

thickness of the mean tree (cm).

2.4.3. Modeling the climate impact of the forest sector

To assess the climate impact of the forest sector, we model atmospheric CO_2 emissions decay

using an Impulse Response Function (IRF), which quantifies the temporal reaction of a

dynamic system in response to some external change (Joos et al., 2013). The multi-model mean

from the carbon cycle model intercomparison in Joos et al. (2013) is used to simulate this decay

process, incorporating a multi-exponential decay function to simulate the response of different

carbon pools, accurately reflecting the transient and long-term behavior of carbon in the

atmosphere, land, and ocean reservoirs (Cherubini et al., 2011; Joos et al., 2013; Strassmann

and Joos, 2018). Its associated IRF is mathematically expressed as:

$IRF(t) = A_0 + \sum_1^3 A_i\, e^{-t/\tau_i}$ *(Eq. 4)*

where $A_0 = 0.2173$, $A_1 = 0.2240$, $A_2 = 0.2824$, $A_3 = 0.2763$, $\tau_1 = 394.4$, $\tau_2 = 36.54$, $\tau_3 = 4.304$

The amplitude A_0 represents the asymptotic airborne fraction of CO_2, which remains in the

atmosphere in the long term due to the equilibrium response of the ocean–atmosphere system.

Each A_i is associated with a characteristic residence time τ_i, which describes how long the

corresponding fraction of carbon remains in the atmosphere before being taken up by the

combined action of terrestrial and oceanic processes. Together, the coefficients $(A_0 + A_1 +$

$A_2 + A_3)$ sum to 1, and thus each A_i can be interpreted as the fractional contribution of a given

decay component to the overall atmospheric response. It is important to note that these

components do not correspond to individual, physically distinct carbon sinks, but rather to a

mathematical approximation of the combined response of the global carbon cycle.

The atmospheric concentration of CO_2 over time can be described as the convolution of the

IRF with the net carbon flux. It is expressed as the integral of past net emissions at time t',

weighted by the fraction of CO_2 remaining in the atmosphere after a time interval t -t', where t

is the reference time point.

To compare the climate impact of biogenic CO_2 emissions across different forest-sector

scenarios, a consistent metric is required. In this paper, we developed an approach based on

the integrated radiative forcing concept adapted for biological systems, referred to here as

RFbio. While originally formulated for pulse emissions in Life Cycle Assessment contexts (e.g.,

GWPbio; see Cherubini et al., 2011), here we extend the concept to account for net carbon fluxes

over time, making it applicable to dynamic forest-sector modeling. RFbio is derived from the

time-integrated radiative forcing caused by emissions, quantifying the perturbation of the

Earth's energy balance. It is calculated as follows:

$RFbio(t) = \int_0^t -FSB(t') \times IRF(t - t')\, dt'$ *(Eq. 5)*

A positive RF_{bio} indicates a net warming effect. In contrast, an RF_{bio} equals to or less than zero

implies that the system has no net contribution to atmospheric warming. This condition can

therefore be interpreted as the forest-sector system being climate neutral ($RF_{bio} \approx 0$), adds to

climate warming ($RF_{bio} > 0$), or climate cooling ($RF_{bio} < 0$) over the assessed time horizon.

Figure 5 provides an illustration of these concepts.

The modeling approach described in Sections 4.2 and 4.3 is operationalized through an

integrated methodological workflow, illustrated in Figure A.1.

Figure 5: CO_2 dynamics across Earth system compartments and the carbon removal dynamics.

(a) Schematic representation of CO_2 exchanges between the atmosphere, ocean and biosphere.

Red arrows indicate CO_2 emissions to the atmosphere; blue arrows indicate CO_2 uptake from

the atmosphere. (b) Net atmospheric CO_2 change (%) over time following a pulse emission,

with equivalent carbon removal occurring at year 60. For analogy, assume that by the starting

year, harvested wood is burned for bioenergy and the resulting emissions are fully captured by

year 60. Response is simulated using the IRF Bern 2.5 CC model. The shaded area represents

the RF$_{bio}$, highlighting temporary overshoot and long-term legacy effect. Positive values,

denote a forcing effect while negative values denote a cooling effect.

2.5. Assessment of carbon and climate mitigation potentials

Carbon balance (CB) was estimated by tracking net changes in forest carbon pools and HWPs,

while also accounting for the generated substitution effects. CB was determined by summing

the FSB (see Eq. 3) over a time horizon T:

$CB(T) = -\sum_{t=1}^{T} FSB(t)$ *(Eq. 6)*

where $FSB(t)$ represents the annual forest sector balance at time t. The total time horizon T

was considered over 85, 185, or 285 years. The minus sign in Equation 6 reflects the convention

that a negative CB indicates a net carbon sink (i.e., net removal of CO_2 from the atmosphere),

while a positive value indicates a net source (i.e., net emissions).

Climate mitigation potential, expressed as radiative forcing due to forest management, was

assessed using the RF$_{bio}$ metric (see Eq. 5), which accounts for the temporary nature of biogenic

CO_2 emissions and their atmospheric decay over time.

Furthermore, the concept of ME is introduced to measure the effectiveness of mitigation

efforts, is defined as the ratio between the negative RF$_{bio}$ and the corresponding carbon

balance. The negative sign ensures that a climate-cooling effect (negative RF$_{bio}$) yields a

positive ME, aligning higher ME values with greater mitigation benefit. This makes ME a

consistent and intuitive indicator of strategy performance across time horizons.

$ME(T) = \frac{RFbio(T)}{CB(T)} \cdot \begin{cases} -1, & RFbio(T) > 0 \text{ and } CB(T) > 0 \\ 1, & otherwise \end{cases}$ (Eq.7)

ME is calculated across different time horizons (85, 185, and 285 years), providing a metric

that relates the remaining or effectively removed biogenic emissions from the atmosphere to

the total carbon balance. The higher the ME value, the greater the climate benefit delivered per

unit of carbon captured, making it a useful comparative indicator of mitigation strategy

performance.

2.5.1 Statistical Analysis of Mitigation Potential

The effects of forest management, climate scenarios, wood use, and substitution factors were

statistically evaluated across a range of carbon-related response variables in the forest sector.

Carbon ecosystem fluxes (i.e., GPP, R_{ECO}, and NEE) were analyzed with respect to forest

management and climate, while wood emissions, substitution effects, wood balance, and FSB

were evaluated in relation to all four factors.

We used linear mixed-effects models (LMEs) to regress GPP, RECO, NEE, and FSB against

five key drivers: Management, Wood Use (WU), Substitution (SUB), Climate, and Century.

These models were used to assess the influence of each factor and their interactions on

ecosystem fluxes and forest sector balances. For wood emissions, substitution effects, and

wood balance, we applied generalized additive mixed models (GAMMs) to account for

potential nonlinear relationships. Model selection was guided by residual diagnostics,

including assessments of linearity, normality, and homoscedasticity. The factors and their

acronyms, used consistently throughout the manuscript, are summarized in **Table 1**. Full model

specifications, transformations, and diagnostic results are provided in **S2**.

Table 1: Overview of the factorial design used in the simulation experiment, showing the five

key factors and their respective levels along with their corresponding modality acronyms.

Factor	Levels	Modality acronyms
Management	5	TM, WOOD, BIOE, ADAPT, TRANS
Wood use	4	BAU, RR20, LS20, RRLS20
Substitution	5	SUB0, SUB25, SUB50, SUB75, SUB100
Climate change scenario	3	CUR, SSP1-2.6, SSP5-8.5
Century	3	C1 (2015-2099), C2 (2100-2199), C3 (2200-2300), ALL (2015-2300)

2.5.2. Consistency between carbon and climate mitigation rankings and the concept of

mitigation efficiency

To evaluate the relationship between CB and RF_{bio} outcomes, a linear regression analysis was

conducted, modeling RF_{bio} as a function of CB across all studied scenarios. Model fit was

assessed using the coefficient of determination (R^2) and statistical significance (p-value).

ME was calculated for each combination of climate scenario, management scheme, substitution

pathway, and time horizon (H1: 2015–2100; H2: 2015–2200; H3: 2015–2300). To assess the

sensitivity of ME to scenario context and temporal scope, we conducted a descriptive statistical

comparison across each factor. Patterns in ME and CB were summarized using means, with

scenario-specific anomalies (e.g., negative ME values) noted.

Additionally, an in-depth analysis of two forest mitigation options that exhibited the two most

divergent ME was conducted. This comparison focused on long-term FSB dynamics, RF_{bio}

patterns, time in a net emissions regime, and disturbance-recovery cycles, to elucidate the

underlying trade-offs between cumulative carbon benefits and the timing and distribution of

emissions.

3. Results

3.1. Forest sector balance (FSB)

FSB, defined as the sum of Net Ecosystem Exchange (NEE) and the wood-product balance

(Wbalance; i.e., substitution effects minus emissions from wood decay in disposal sites), varied

significantly with climate scenarios, management strategies, and across centuries. Climate

effects on FSB were not consistent across forest management types, and both drivers showed

strong century-dependent patterns (**see Table S2.4**). Although the displacement factors (DF)

had a low overall effect size ($\eta^2 = 0.03$, p<0.001), its influence on FSB was substantial in

specific scenarios, particularly those representing rapid decarbonization pathways. Over the

full simulation period, SSP5-8.5 yielded the lowest FSB (0.07 ± 0.01 tC ha^{-1} year^{-1}), while

SSP1-2.6 and current climate (CUR) resulted in significantly higher values (0.75 ± 0.01 and

0.64 ± 0.01, respectively). The ranking of scenarios changed over time: SSP5-8.5 initially

outperformed others but declined in the 22nd and 23rd centuries, ultimately ranking lowest

(**see Figure 6**). By contrast, SSP1-2.6 and CUR remained more stable across centuries with

lower inter-temporal variability.

Management strategies showed clear performance gradients mainly driven by NEE. Modular

harvesting (TM) and bioenergy (BIOE) produced the highest forest sector balance (FSB;

0.76 ± 0.02 and 0.59 ± 0.04), supported by relatively strong sinks (NEE: -1.25 ± 0.05 and $-$

1.10 ± 0.08). Promotion of long-lived wood products (WOOD) and forest adaptation

management (ADAPT) showed intermediate performance (FSB: 0.39 ± 0.02 and 0.26 ± 0.02)

associated with weaker sinks (NEE: -0.78 ± 0.05 and -0.54 ± 0.05). In contrast, transition to

passive rewilding (TRANS) consistently produced the lowest FSB (0.05 ± 0.03), declining to

negative values under SSP5-8.5 (-0.47 ± 0.03) due to forest ecosystem turning into carbon

source (NEE: $+0.83 \pm 0.01$) (**see Figure 6**). Decreases were most severe in TRANS and

ADAPT, where net carbon source behavior emerged in the third century.

DF, which modulates substitution levels over time, had a major influence on Wbalance and

thus on FSB. Under the baseline pathway (SUB0), the substitution effect was maintained

throughout all centuries, while decarbonization pathways (SUB25–SUB100) showed a

progressive decline in substitution, reaching zero by 2155 (SUB25), 2085 (SUB50), 2062

(SUB75), and 2050 (SUB100), respectively. These trajectories form three distinct temporal

patterns: persistent (SUB0), medium-lived (SUB25), and short-lived (SUB50–SUB100),

which resulted in markedly different Wbalance outcomes.

Component analysis indicated that NEE accounted for ~53–65% of FSB, particularly high

under TM. Substitution effect was lower in terms of contribution to the FSB but variable,

peaking under the SUB0 displacement pathway (18%), decreasing under SUB25 (7%), and

falling below 3% for SUB50–SUB100, which caused a (0.28 ± 0.07 tC ha^{-1} year^{-1}; $53 \pm 25\%$)

reduction in FSB compared to SUB0 (**see Table S2.5**).

Figure 6: Mean FSB, in tC ha^{-1} year^{-1} for five forest management strategies (TM, ADAPT, BIOE, TRANS, and WOOD), averaged across two climate scenarios (SSP1-2.6 and SSP5-8.5), shown for the 21st, 22nd, and 23rd centuries, as well as for the full period combined. Red error bars indicate the 95% confidence interval of FSB across climate scenarios. The upper panel shows FSB values by substitution scenarios, while the lower panel shows FSB values by climate scenario.

3.2. Radiative forcing & mitigation efficiency (RF$_{bio}$ and ME)

Forest management scenarios with a more negative carbon balance (CB) also tended to yield the greatest climate benefits. A linear regression, performed using outcomes from all studied scenarios, confirmed that CB is a strong predictor of the radiative forcing from biogenic emissions (RF$_{bio}$; $R^2 = 0.95$; $p < 2.2e{-}16$; see **Table S2.9**). Thus, the results and analyses previously conducted on CB are also applicable to RF$_{bio}$. However, this relationship is not always strictly linear. In some exceptional cases where CB values are close, a mitigation option with a slightly less negative CB produced a more favorable RF$_{bio}$. This was observed, for instance, under the SUB25 decarbonization pathway at horizon H2 (2015-2200): the TRANS

scenario had a slightly higher CB (-128 ± 0.91 tC ha^{-1}) than WOOD (-122 ± 2.75 tC ha^{-1}), yet

WOOD achieved a more favorable RF$_{bio}$ (-66 ± 1.33 vs. -60 ± 0.13 tC ha^{-1}) and a higher ME

(0.54 ± 0.00 vs. 0.47 ± 0.002). At horizon H3 (2015-2399), this contrast became even more

pronounced: although TRANS reached a more negative CB (-162 ± 0.92 tC/ha) compared to

WOOD (-155 ± 2.05 tC ha^{-1}), it resulted in a substantially less favorable RF$_{bio}$ (-66 ± 0.70 vs.

-73 ± 0.20 tC ha^{-1}) and a lower mitigation efficiency (ME; 0.41 ± 0.002 vs. 0.57 ± 0.002). ME

showed strong sensitivity to both climate conditions and forest management practices.

Under CUR and SSP1-2.6, ME remained relatively stable across management schemes,

typically ranging from 0.4 to 0.7 (**Figure 7** and **Table S2.10**). In contrast, under the SSP5-8.5

climate scenario, the ME of different management scenarios was more variable and often with

lower efficiencies, with the efficiency of several scenarios falling below zero, particularly

ADAPT and TRANS under H2 and H3 (**Figure 7** and **Table S2.10**), indicating that these

scenarios were contributing to climate warming rather than cooling it. The time horizon (H1,

H2, and H3) also influenced ME: across all climate scenarios, the short-term horizon (H1)

consistently resulted in higher efficiency values (>0.6) compared to the longer-term time

horizon (H2 and H3).

Interestingly, a more negative CB did not always equate to high efficiency. In CUR and SSP1-

2.6, forest projects with stronger (more negative) CB often showed lower or slightly lower

efficiency, whereas the opposite was true under SSP5-8.5, where stronger CB tended to support

higher ME. For example, ADAPT (0.53, SSP1-2.6, H2) and WOOD (0.54, SSP1-2.6, H2)

sometimes outperformed TM (0.49, SSP1-2.6, H2) or BIOE (0.45, SSP1-2.6, H2) in efficiency

despite having less negative CB values.

These dynamics were further supported and extended by generalized additive models (GAMs;

Figures S2.1-S2.2), which demonstrated that, in addition to CB, both the magnitude and

duration of emissions play dominant roles in determining ME. Under the SSP5-8.5 scenario,

emissions magnitude and CB were strongly positively correlated (r = 0.85), indicating that

scenarios with more negative CB generally exhibited low emissions. In contrast, a weak

negative correlation (r = –0.28) was observed under CUR and SSP1-2.6, suggesting no clear

relationship between CB and emissions magnitude. In the ADAPT and TRANS scenarios,

under the SSP5-8.5 climate scenario, positive CB values resulted in sustained net emissions

and positive contributions to radiative forcing.

Figure 7: CB and ME under various forest management and climate scenarios. Circle size

scales with CB via $-0.9 \times CB^{1.3}$; ME is quantile-transformed, both for improved visibility and

comparability across scenarios. Red borders indicate negative ME; hexagons show positive CB

(carbon source sector). Subplots represent climate scenarios, with circles colored by

management schemes.

3.3. In depth-insights from BIOE and WOOD strategies

Across the simulation period (2015–2300), the BIOE management scenario consistently

achieved a more negative CB than WOOD. By 2300, BIOE reached –337 tC ha^{-1}, while WOOD

reached –206 tC ha^{-1}. ME, however, remained higher in WOOD throughout. Both scenarios

began with comparable ME in 2100 (0.62 for BIOE, 0.63 for WOOD). By 2200, WOOD

retained a ME of 0.54, while BIOE declined to 0.46. In 2300, ME values were 0.47 for WOOD

and 0.44 for BIOE (see **Figure 8**).

BIOE followed an 80-year shelterwood rotation, resulting in three major final harvests

occurring in 2095, 2175, and 2265. WOOD used a 130-year rotation, with final cuts in 2145

and 2260. The first BIOE final harvest in 2095 led to emissions of approximately 17.3 tC ha^{-1},

with RF$_{bio}$ returning to pre-disturbance levels by 2135, indicating a 40-year payback period.

For WOOD, the first final cut in 2145 resulted in emissions of 5.6 tC ha^{-1}, with RF$_{bio}$ returning

to net cooling within 15 years. Following these major harvests, NEE in WOOD returned to

negative values (net sink) within 4 years, whereas BIOE remained a net carbon source (NEE >

0) for approximately 10 years.

In the three years immediately following each final cut, RF$_{bio}$ increased sharply in BIOE, while

the increase in WOOD was more moderate. This early post-harvest rise in RF$_{bio}$ coincided with

higher volumes of short-lived HWPs (i.e., firewood and paper) in BIOE; 37.25 m³ ha^{-1},

38.51 m³ ha^{-1}, and 37.88 m³ ha^{-1} after its respective final harvests, compared to 23.70 m³ ha^{-1}

and 14.82 m³ ha^{-1} in WOOD.

Emission timing and magnitude differed between the two scenarios. In BIOE, emissions were

concentrated in a limited number of high-intensity events, associated with initial establishment

and final harvests. These events corresponded to periods where GPP was low and ecosystem

respiration (R$_{ECO}$) remained high although the carbon outflow from the forest ecosystem

(Rh$_{onsite}$) decreased proportionally to GPP. Additional immediate emissions from firewood

combustion or from products at their end-life incineration (i.e., paper) were also observed

during these intervals. In WOOD, emissions were more evenly distributed and of lower

magnitude, arising from thinning, regeneration phases, and smaller final cuts. RF$_{bio}$ patterns in

WOOD were more stable, with shorter durations of net positive forcing following harvest

events.

Figure 8: Temporal dynamics of RF$_{bio}$, CB, and ME under BIOE (panel a) and WOOD (panel

b) management scenarios from 2015 to 2300. Shaded red bands indicate periods when the

forest sector is a net source (FSB<0). Green horizontal bands denote the number of years

required for RF$_{bio}$ to return to its pre-harvest level.

4. Discussions

Our results suggest that effective forest-sector climate mitigation requires more than

maximizing carbon balances. It demands a systems perspective integrating forest dynamics,

wood use, and evolving socio-technical conditions. Over a multi-century assessment, we found

that while the forest sector balance (FSB) is strongly influenced by climate and management,

mitigation efficiency (ME) depends more on emission timing and scale. Substitution effects,

particularly the contrast between the baseline pathway (SUB0) and the degressive pathways

(SUB25-SUB100) proved substantial, while HWP lifespan and recycling rate had negligible

long-term impact. Notably, scenarios with the lowest carbon balance (CB) did not always

deliver the greatest climate efficiency. ME declined over time, emphasizing the need to

consider temporal dynamics. All management options eventually showed declining carbon sink

capacity under SSP5-8.5, with TRANS, ADAPT, and WOOD becoming net sources.

4.1. Management strategies, wood use, and substitution pathways

Forest management has a strong influence on FSB. Intensive silvicultural practices, such as

TM and BIOE, consistently maintained high FSB across climate projections. (Giasson et al.,

2023) showed that harvest scenarios with high intervention and regular regrowth cycles sustain

stronger carbon sink than low-harvest or unmanaged systems. Furthermore, our results showed

that shifting from active to passive management under the TRANS scenario significantly

reduced FSB in the long term, especially under contrasted climate conditions where forest

carbon dynamics approached equilibrium. These findings align with (Pukkala, 2017), who

reported that unmanaged forests exhibit a long-term decline, nearing zero around 200 years

post-establishment. Beyond carbon storage, management also influences wood volume and

characteristics, affecting substitution potential.

Recent studies increasingly challenge static DFs, advocating for dynamic approaches that

reflect evolving technological, industrial, and policy contexts (I. Boukhris et al., 2025; Brunet-

Navarro et al., 2021; Hurmekoski et al., 2023). Our dynamic modeling confirms these

concerns. In scenarios where DFs reaches zero by 2050, substitution benefits disappear

entirely, resulting in a 53% lower FSB by 2300 compared to the SUB0 trajectory. This mirrors

projections by Brunet-Navarro et al. (2021), who estimated substitution benefits may decrease

by approximately 30% by 2030 and up to 96% by 2100, in line with the decarbonization goals

of the Paris Agreement.

Conversely, wood-use parameters like product lifespan and recycling rate had no discernable

long-term impact on FSB. Over 285 years, stored carbon in HWPs is eventually emitted

through decay, regardless of whether turnover occurs at 40 or 60 years. Although extended

lifespans and higher recycling rates may reduce emissions in the short- to medium-term, as

shown by (Brunet-Navarro et al., 2016), who projected up to 5 Mt CO_2 yr^{-1} savings in the EU-

28 with lifespan increases of 19.54% or a 20.92% recycling boost by 2030, these effects

diminish across multi-century timescales. Such results highlight the need to align interventions

with the analytical timeframe. Short-term gains may appear significant but are absorbed over

longer horizons when delayed emissions fall within the evaluation period.

4.2. Carbon sink instability under high-emission scenarios

The shift to net-carbon neutrality or a carbon source state by the last considered century under

SSP5-8.5 marks a critical loss of ecosystem resilience, driven by prolonged structural

degradation and physiological stress from high temperatures. Although precipitation increases

during this period, and vapor pressure deficit stays moderate (~0.825 ± 0.1 kPa), GPP declines

sharply across all strategies. This drop is more likely due to heat-induced impacts on

photosynthetic processes, such as enzyme deactivation, increased photorespiration, and

reduced carboxylation efficiency (Oliver et al., 2022; Qu Y et al., 2023). As biomass and

metabolic activity decline, both Ra and Rh fall due to limited substrate and structural loss, but

GPP declines faster, leading to sustained positive NEE in the final simulation century. This

aligns with Earth System Model outputs under SSP5-8.5 which show that warming suppresses

photosynthesis as temperatures exceed physiological thresholds, particularly in temperate and

boreal forests. The summer GPP-temperature correlation weakens over the 21st century,

turning significantly negative by 2070 (Chen and Peng, 2025; Zhang et al., 2022) also found

that 33% to 63% of global vegetated areas show declining productivity sensitivity to

temperature, with the most severe declines under SSP5-8.5.

4.3. Climate-Carbon dynamics and mitigation efficiency

The strong correlation observed between carbon balance (CB) and radiative forcing from

biogenic emissions (RF_{bio}) reflects the key role of long-lived CO_2 removals in determining

climate impacts of forest-based mitigation. This aligns with carbon cycle theory and is

consistent with impulse response functions (IRFs) such as the Bern model (Joos et al., 2013),

where most climate benefit arises from sustained atmospheric CO_2 reduction (Archer et al.,

2009; Masson-Delmotte et al., 2021). However, the CB-RF_{bio} relationship is not strictly

proportional. Our results show that under the SUB25 decarbonization pathway at both H2 and

H3, the WOOD scenario achieved a lower (better) RF_{bio} than TRANS, despite similar or less

negative CB. This suggests CB alone is insufficient basis for assessing climate performance,

and other factors also influence radiative forcing.

A key reason for this explanation for the discrepancy is the timing of carbon fluxes. CB

summarizes total emissions and removals but masks their temporal distribution. Many

scenarios involve non-monotonic dynamics, with initial emissions from establishment,

thinning, or harvest followed by delayed regrowth. Early emissions exert greater radiative

forcing due to the CO_2's long atmospheric lifetime cumulative warming (Allen et al., 2010;

Matthews and Caldeira, 2008). Although IRF is linear for pulses, time-distributed fluxes

interact non-linearly with the climate system. RF_{bio}, which captures these dynamics, more

accurately reflects climate impact than CB alone (Sierra et al., 2021; Tanaka and O'Neill,

2018).

Nevertheless, neither CB nor RFbio capture the carbon "cost" of achieving mitigation. We

therefore introduced the mitigation efficiency (ME; RFbio divided by CB). ME indicates how

effectively carbon removal translates into avoided radiative forcing. For instance, BIOE had a

more negative CB (-339 tC ha^{-1}) than ADAPT (-162 tC ha^{-1}), but a lower ME (0.44 vs. 0.49),

meaning less efficient climate benefit. BIOE also required 17 silvicultural interventions and 3

final cuts over 300 years, compared to 14 interventions and 2 final cuts in ADAPT. Thus, more

intensive management does not always yield proportionally greater mitigation and may entail

higher ecological or operational costs.

ME declined consistently as the time horizon extended. Initially, forests sequester carbon in

biomass and soils, but later interventions transfer carbon to HWPs, which eventually decay and

emit CO_2. Over time, emissions from earlier cycles accumulate, reducing net benefit. This

dynamic, "sequester now, emit later" (Guest et al., 2013), underscores the value of ME as a

complementary metric that integrates both climate effectiveness and implementation

sustainability.

4.4. Uncertainty, Model assumptions, and Long-Term Implications

Uncertainties within the modeling framework stem from initialization, process representation,

and parameterization ((Hamish) Kimmins et al., 2008; Reyer et al., 2020). The 3D-CMCC-

FEM forest growth model, while effective for stand-level analyses, simplifies spatial

heterogeneity and omits landscape-scale dynamics, potentially underrepresenting key

interactions such as seed dispersal and spatially explicit disturbance spread (Patacca et al.,

2023; Seidl et al., 2011). It also omits processes like tree establishment, disturbance-induced

mortality, and species composition are not represented, limiting its ability to capture ecological

feedback under climate change. Adaptive responses, including species acclimation, migration,

and phenological change, are excluded or overly simplified, despite their growing relevance

over multi-century timescales (Issam Boukhris et al., 2025; Nicotra et al., 2010).

On the HWPs side, the TimberTracer model uses fixed bucking rules and constant product

allocations, without accounting for future technological shifts, circular economy

developments, or changing consumer demand (Harju, 2022; Olšiaková et al., 2016). This

restricts its ability to reflect how innovation or policy changes could influence product

lifespans, recycling rates, or end-of-life treatment.

Additional uncertainty comes from external drivers. Climate projections diverge significantly

after 2100 (Hausfather and Peters, 2020), and future land-use and forest management pathways

depend on socio-economic and policy trends that are inherently hard to predict (Popp et al.,

2017). DF estimates used to quantify substitution benefits are also highly uncertain and

context-specific (Leskinen et al., 2018; Pingoud et al., 2010). Their magnitude will likely

evolve as energy systems decarbonize and industry shifts to lower-carbon alternatives

(Gustavsson et al., 2017).

Although the Vallombrosa site represents a productive temperate European forest, this study

aims to identify general mechanisms rather than universal prescriptions. The relative ranking

of management strategies observed here should be interpreted as site-dependent. In highly

productive temperate systems, rapid regrowth can offset harvest emissions, whereas in boreal

forests slower recovery may extend emissions payback times and favor lower harvest

intensities. Conversely, in drought-prone Mediterranean regions or disturbance-dominated

mountain forests, mortality dynamics may outweigh planned interventions. Nevertheless, the

structural relationships identified in this study appear robust: carbon balance alone does not

determine climate benefit, mitigation efficiency depends on emission timing and magnitude,

and declining substitution factors systematically reduce long-term climate mitigation. These

mechanisms arise from carbon-cycle dynamics rather than site-specific properties and are

therefore expected to apply broadly across forest contexts. Considering this, EU-level

mitigation assessments based solely on carbon stock indicators may overlook important

climate-impact differences among strategies. Rather than prescribing uniform management

targets, policy frameworks may benefit from approaches that evaluate mitigation performance

using temporally explicit metrics and that allow regional adaptation to ecological conditions

and evolving substitution effectiveness.

Conclusions

This study critically examined the long-term climate mitigation potential of the forest sector

by integrating forest growth, wood-use, and substitution dynamics within a unified modeling

framework. Our aim was not only to assess carbon balances but to evaluate whether commonly

used indicators accurately reflect true climate benefit over multi-century timescales.

Using a coupled modeling approach applied on a *P. nigra* forest in Italy, we simulated a broad

range of futures across five forest management strategies, four wood-use patterns, and five

substitution decay scenarios under three contrasting climate pathways. While CB and RF_{bio}

were generally aligned, they captured different aspects of climate impact. CB reflects net

carbon flows, but not their atmospheric effect; RF_{bio} represents the warming or cooling

influence overtime. To bridge this gap, we introduced ME, a metric that relates carbon flows

to actual contribution to actual mitigation effectiveness.

Our findings show that scenarios with the highest CB (e.g., BIOE and TM) do not always

deliver the greatest climate benefit, especially when emissions are front-loaded or persistent.

Meanwhile, strategies like WOOD and ADAPT, which extended carbon residence times in

both forest and technosphere, achieved higher ME under SSP1-2.6. However, under the SSP5-

8.5 scenario, even these strategies became net radiative sources after 2200 due to declining

sink capacity.

Substitution benefits, often assumed constant, declined significantly under degressive

assumptions, reducing the overall forest sector mitigation by up to 53%, especially in high

harvest volumes. While substitution played a secondary role compared to climate and

management, its sensitivity to decarbonization pathways highlights the risk of overreliance on

this lever.

Taken together, these findings highlight that effective forest-sector mitigation requires more

than achieving high carbon balances. It also depends on minimizing the magnitude, timing, and

persistence of emissions, and on reevaluating the assumptions behind substitution effects.

Metrics like RF_{bio} and ME provide valuable tools to capture these temporal and systemic

dynamics, offering a more nuanced, climate-relevant basis for evaluating forest-sector

strategies.

Ultimately, meaningful forest-sector contributions to climate neutrality require assessment not

just of carbon flows, but of their actual atmospheric impact over time.

Appendix

Figure A.1: Integrated modeling framework used to evaluate the long-term climate mitigation potential of a managed *Pinus nigra* forest (2015-2300). The framework combines forest dynamics (3D-CMCC-FEM), wood lifecycle modeling (TimberTracer), and an impulse response function (Bern 2.5 CC IRF). Simulations cover multiple scenario combinations (for management, climate, wood-use, and substitution). Outputs include temporal net ecosystem exchange, substitution effect, and emissions from wood combustion and decay, which are used to compute forest sector balance and radiative forcing from biogenic CO_2 emissions.

References

Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.H. (Ted), Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S.W., Semerci, A., Cobb, N., 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. For. Ecol. Manag., Adaptation of Forests and Forest Management to Changing Climate 259, 660–684. https://doi.org/10.1016/j.foreco.2009.09.001

Ameray, A., Bergeron, Y., Valeria, O., Montoro Girona, M., Cavard, X., 2021. Forest Carbon Management: a Review of Silvicultural Practices and Management Strategies Across Boreal, Temperate and Tropical Forests. Curr. For. Rep. 7, 245–266. https://doi.org/10.1007/s40725-021-00151-w

Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K., Matsumoto, K., Munhoven, G., Montenegro, A., Tokos, K., 2009. Atmospheric Lifetime of Fossil Fuel Carbon Dioxide. Annu. Rev. Earth Planet. Sci. 37, 117–134. https://doi.org/10.1146/annurev.earth.031208.100206

Barredo, J.I., Vizzarri, M., Kuželová, K., 2024. Archetypal typology of European forest ecosystems integrating management intensity and naturalness. Ambio 53, 1587–1598. https://doi.org/10.1007/s13280-024-02050-3

Barrette, J., Achim, A., Auty, D., 2023. Impact of Intensive Forest Management Practices on Wood Quality from Conifers: Literature Review and Reflection on Future Challenges. Curr. For. Rep. 9, 101–130. https://doi.org/10.1007/s40725-023-00181-6

Bauhus, J., Puettmann, K., Messier, C., 2009. Silviculture for old-growth attributes. For. Ecol. Manag., Old forests, new management: the conservation and use of old-growth forests in the 21st century 258, 525–537. https://doi.org/10.1016/j.foreco.2009.01.053

Bottalico, F., Chirici, G., Travaglini, D., 2012. La gestione della foresta di Vallombrosa dal 1876 al 2006: analisi delle cartografie storiche. Ital. For. E Mont. 449–458. https://doi.org/10.4129/ifm.2012.6.01

Boukhris, I., Collalti, A., Lahssini, S., Dalmonech, D., Nakhle, F., Testolin, R., Chiriacò, M.V., Santini, M., Valentini, R., 2025. TimberTracer: a comprehensive framework for the evaluation of carbon sequestration by forest management and substitution of harvested wood products. Carbon Balance Manag. 20, 12. https://doi.org/10.1186/s13021-025-00296-2

Boukhris, Issam, Marano, G., Dalmonech, D., Valentini, R., Collalti, A., 2025. Modeling Forest Growth Under Current and Future Climate. Curr. For. Rep. 11, 17. https://doi.org/10.1007/s40725-025-00249-5

Bozzolan, N., Grassi, G., Mohren, F., Nabuurs, G.-J., 2024. Options to improve the carbon balance of the harvested wood products sector in four EU countries. GCB Bioenergy 16, e13104. https://doi.org/10.1111/gcbb.13104

Brunet-Navarro, P., Jochheim, H., Cardellini, G., Richter, K., Muys, B., 2021. Climate mitigation by energy and material substitution of wood products has an expiry date. J. Clean. Prod. 303, 127026. https://doi.org/10.1016/j.jclepro.2021.127026

Brunet-Navarro, P., Jochheim, H., Muys, B., 2017. The effect of increasing lifespan and recycling rate on carbon storage in wood products from theoretical model to application for the European wood

877   sector. Mitig. Adapt. Strateg. Glob. Change 22, 1193–1205. https://doi.org/10.1007/s11027-016-
878   9722-z
879
880   Brunet-Navarro, P., Jochheim, H., Muys, B., 2016. Modelling carbon stocks and fluxes in the wood
881   product sector: a comparative review. Glob. Change Biol. 22, 2555–2569.
882   https://doi.org/10.1111/gcb.13235
883
884   Budzinski, M., Bezama, A., Thrän, D., 2020. Estimating the potentials for reducing the impacts on
885   climate change by increasing the cascade use and extending the lifetime of wood products in
886   Germany. Resour. Conserv. Recycl. X 6, 100034. https://doi.org/10.1016/j.rcrx.2020.100034
887
888   Calvin, K., Bond-Lamberty, B., Clarke, L., Edmonds, J., Eom, J., Hartin, C., Kim, S., Kyle, P., Link,
889   R., Moss, R., McJeon, H., Patel, P., Smith, S., Waldhoff, S., Wise, M., 2017. The SSP4: A world of
890   deepening inequality. Glob. Environ. Change 42, 284–296.
891   https://doi.org/10.1016/j.gloenvcha.2016.06.010
892
893   Cantiani P, Di Salvatore U, Romano R, 2018. La selvicoltura delle pinete artificiali di pino nero:
894   analisi delle legislazioni regionali italiane. For. - J. Silvic. For. Ecol. 16, 99.
895   https://doi.org/10.3832/efor2985-015
896
897   Chen, T., Peng, L., 2025. Global GPP Changes and Its Sensitivity to Temperature, in: Chen, T., Peng,
898   L. (Eds.), Multi-Scale Understanding of Vegetation Ecosystems: Changes, Drivers and Inspirations.
899   Springer Nature Switzerland, Cham, pp. 145–169. https://doi.org/10.1007/978-3-031-90163-8_8
900
901   Cherubini, F., Peters, G.P., Berntsen, T., Strømman, A.H., Hertwich, E., 2011. CO2 emissions from
902   biomass combustion for bioenergy: atmospheric decay and contribution to global warming. GCB
903   Bioenergy 3, 413–426. https://doi.org/10.1111/j.1757-1707.2011.01102.x
904
905   Ciancio, O., Nocentini, S., 2011. Biodiversity conservation and systemic silviculture: Concepts and
906   applications. Plant Biosyst. - Int. J. Deal. Asp. Plant Biol. 145, 411–418.
907   https://doi.org/10.1080/11263504.2011.558705
908
909   Ciancio, O., Nocentini, S., 2004. Il bosco ceduo. Selvicoltura, assestamento, gestione. Firenze:
910   Academia Italiana di Scienze Forestali, 721 pp. ISBN 978-88-87553-06-2.
911
912   Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., Valentini, R., 2014. A
913   process-based model to simulate growth in forests with complex structure: Evaluation and use of 3D-
914   CMCC Forest Ecosystem Model in a deciduous forest in Central Italy. Ecol. Model. 272, 362–378.
915   https://doi.org/10.1016/j.ecolmodel.2013.09.016
916
917   Dalmonech, D., Marano, G., Amthor, J.S., Cescatti, A., Lindner, M., Trotta, C., Collalti, A., 2022.
918   Feasibility of enhancing carbon sequestration and stock capacity in temperate and boreal European
919   forests via changes to management regimes. Agric. For. Meteorol. 327, 109203.
920   https://doi.org/10.1016/j.agrformet.2022.109203
921
922   Dalmonech, D., Vangi, E., Chiesi, M., Chirici, G., Fibbi, L., Giannetti, F., Marano, G., Massari, C.,
923   Nolè, A., Xiao, J., Collalti, A., 2024. Regional estimates of gross primary production applying the
924   Process-Based Model 3D-CMCC-FEM vs. Remote-Sensing multiple datasets. Eur. J. Remote Sens.
925   57, 2301657. https://doi.org/10.1080/22797254.2023.2301657
926
927   Dye, A.W., Houtman, R.M., Gao, P., Anderegg, W.R.L., Fettig, C.J., Hicke, J.A., Kim, J.B., Still,
928   C.J., Young, K., Riley, K.L., 2024. Carbon, climate, and natural disturbance: a review of mechanisms,
929   challenges, and tools for understanding forest carbon stability in an uncertain future. Carbon Balance
930   Manag. 19, 35. https://doi.org/10.1186/s13021-024-00282-0
931

European Environment Agency (EEA), 2024. Diversion of waste from landfill in Europe. EEA indicator assessment. Copenhagen: European Environment Agency. Available at: https://www.eea.europa.eu/en/analysis/indicators/diversion-of-waste-from-landfill (accessed 25 Feb 2026).

European Environment Agency (EEA), 2022. Annual European Union greenhouse gas inventory 1990-2020 and inventory report 2022. EEA Report No 02/2022. Copenhagen: European Environment Agency. https://www.eea.europa.eu/publications/annual-european-union-greenhouse-gas-inventory-2022 (Accessed 25 February 2026).

Council of the European Union, 2025. Improved EU rules for forest reproductive material: Council approves negotiating position. Press release, 13 June 2025. Available at: https://www.consilium.europa.eu/en/press/press-releases/2025/06/13/improved-eu-rules-for-forest-reproductive-material-council-approves-negotiating-position/(accessed 27 July 2025).

European Commission, 2021. New EU forest strategy for 2030. European Commission, Brussels. Available at: https://environment.ec.europa.eu/strategy/forest-strategy_en (accessed 27 July 2025).

European Commission, 2018. A sustainable bioeconomy for Europe: strengthening the connection between economy, society and the environment – Updated bioeconomy strategy. Brussels: European Commission. Available at: https://environment.ec.europa.eu/strategy/bioeconomy-strategy_en (accessed 27 July 2025).

European Commission, 2021. EU biodiversity strategy for 2030: Bringing nature back into our lives. Brussels: European Commission. Available at: https://environment.ec.europa.eu/strategy/biodiversity-strategy-2030_en (accessed 27 July 2025).

Eyring, V., Bony, S., Meehl, G.A., Senior, C.A., Stevens, B., Stouffer, R.J., Taylor, K.E., 2016. Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. Geosci. Model Dev. 9, 1937–1958. https://doi.org/10.5194/gmd-9-1937-2016

Giasson, L.-A., Thiffault, E., Lebel, L., Carle, J.-F., 2023. Carbon balance of forest management and wood production in the boreal forest of Quebec (Canada). Front. For. Glob. Change 6. https://doi.org/10.3389/ffgc.2023.1242218

Grassi, G., Fiorese, G., Pilli, R., Jonsson, K., Blujdea, V., Korosuo, A., Vizzarri, M., 2021. Brief on the role of the forest-based bioeconomy in mitigating climate change through carbon storage and material substitution. EUR 30763 EN. Luxembourg: Publications Office of the European Union, European Commission Joint Research Centre. Available at: https://publications.jrc.ec.europa.eu/repository/handle/JRC124374 (accessed 18 December 2023).

Gregor, K., Krause, A., Reyer, C.P.O., Knoke, T., Meyer, B.F., Suvanto, S., Rammig, A., 2024. Quantifying the impact of key factors on the carbon mitigation potential of managed temperate forests. Carbon Balance Manag. 19, 10. https://doi.org/10.1186/s13021-023-00247-9

Guest, G., Cherubini, F., Strømman, A.H., 2013. The role of forest residues in the accounting for the global warming potential of bioenergy. GCB Bioenergy 5, 459–466. https://doi.org/10.1111/gcbb.12014

984
985 Gustavsson, L., Haus, S., Lundblad, M., Lundström, A., Ortiz, C.A., Sathre, R., Truong, N.L.,
986 Wikberg, P.-E., 2017. Climate change effects of forestry and substitution of carbon-intensive
987 materials and fossil fuels. Renew. Sustain. Energy Rev. 67, 612–624.
988 https://doi.org/10.1016/j.rser.2016.09.056
989
990 (Hamish) Kimmins, J.P., Blanco, J.A., Seely, B., Welham, C., Scoullar, K., 2008. Complexity in
991 modelling forest ecosystems: How much is enough? For. Ecol. Manag., 6th North American Forest
992 Ecology Workshop: From science to sustainability 256, 1646–1658.
993 https://doi.org/10.1016/j.foreco.2008.03.011
994
995 Harju, C., 2022. The perceived quality of wooden building materials—A systematic literature review
996 and future research agenda. Int. J. Consum. Stud. 46, 29–55. https://doi.org/10.1111/ijcs.12764
997 Harmon, M.E., 2019. Have product substitution carbon benefits been overestimated? A sensitivity
998 analysis of key assumptions. Environ. Res. Lett. 14, 065008. https://doi.org/10.1088/1748-
999 9326/ab1e95
1000
1001 Hausfather, Z., Peters, G.P., 2020. Emissions – the 'business as usual' story is misleading. Nature
1002 577, 618–620. https://doi.org/10.1038/d41586-020-00177-3
1003
Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J.,
Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G.,
Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis,
M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan,
R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay,
P., Rozum, I., Vamborg, F., Villaume, S., Thépaut, J.-N., 2020. The ERA5 global reanalysis. Q. J. R.
Meteorol. Soc. 146, 1999–2049. https://doi.org/10.1002/qj.3803
1011
1012 Holtsmark, B., 2013. The outcome is in the assumptions: analyzing the effects on atmospheric CO2
1013 levels of increased use of bioenergy from forest biomass. GCB Bioenergy 5, 467–473.
1014 https://doi.org/10.1111/gcbb.12015
1015
1016 Holtsmark, B., 2012. Harvesting in boreal forests and the biofuel carbon debt. Clim. Change 112,
1017 415–428. https://doi.org/10.1007/s10584-011-0222-6
1018
1019 Howard, C., Dymond, C.C., Griess, V.C., Tolkien-Spurr, D., van Kooten, G.C., 2021. Wood product
1020 carbon substitution benefits: a critical review of assumptions. Carbon Balance Manag. 16, 9.
1021 https://doi.org/10.1186/s13021-021-00171-w
1022
1023 Hurmekoski, E., Kunttu, J., Heinonen, T., Pukkala, T., Peltola, H., 2023. Does expanding wood use in
1024 construction and textile markets contribute to climate change mitigation? Renew. Sustain. Energy
1025 Rev. 174, 113152. https://doi.org/10.1016/j.rser.2023.113152
1026
1027 Jonsson, R., Rinaldi, F., Pilli, R., Fiorese, G., Hurmekoski, E., Cazzaniga, N., Robert, N., Camia, A.,
1028 2021. Boosting the EU forest-based bioeconomy: Market, climate, and employment impacts. Technol.
1029 Forecast. Soc. Change 163, 120478. https://doi.org/10.1016/j.techfore.2020.120478
1030
Joos, F., Roth, R., Fuglestvedt, J.S., Peters, G.P., Enting, I.G., von Bloh, W., Brovkin, V., Burke, E.J.,
Eby, M., Edwards, N.R., Friedrich, T., Frölicher, T.L., Halloran, P.R., Holden, P.B., Jones, C.,
Kleinen, T., Mackenzie, F.T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A.,
Segschneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., Weaver,
A.J., 2013. Carbon dioxide and climate impulse response functions for the computation of greenhouse
gas metrics: a multi-model analysis. Atmospheric Chem. Phys. 13, 2793–2825.
https://doi.org/10.5194/acp-13-2793-2013
1038

Kilpeläinen, A., Peltola, H., 2022. Carbon Sequestration and Storage in European Forests, in: Hetemäki, L., Kangas, J., Peltola, H. (Eds.), Forest Bioeconomy and Climate Change. Springer International Publishing, Cham, pp. 113–128. https://doi.org/10.1007/978-3-030-99206-4_6

Krug, J., Koehl, M., Kownatzki, D., 2012. Revaluing unmanaged forests for climate change mitigation. Carbon Balance Manag. 7, 11. https://doi.org/10.1186/1750-0680-7-11

Laganière, J., Paré, D., Thiffault, E., Bernier, P.Y., 2017. Range and uncertainties in estimating delays in greenhouse gas mitigation potential of forest bioenergy sourced from Canadian forests. GCB Bioenergy 9, 358–369. https://doi.org/10.1111/gcbb.12327

Lemprière, T.C., Kurz, W.A., Hogg, E.H., Schmoll, C., Rampley, G.J., Yemshanov, D., McKenney, D.W., Gilsenan, R., Beatch, A., Blain, D., Bhatti, J.S., Krcmar, E., 2013. Canadian boreal forests and climate change mitigation. Environ. Rev. 21, 293–321. https://doi.org/10.1139/er-2013-0039

Leskinen, P., Cardellini, G., González-García, S., Hurmekoski, E., Sathre, R., Seppälä, J., Smyth, C., Stern, T., Verkerk, P. J. and European Forest Institute, 2018. Substitution effects of wood-based products in climate change mitigation (From Science to Policy). European Forest Institute. Available at: https://doi.org/10.36333/fs07.

Levasseur, A., Lesage, P., Margni, M., Deschênes, L., Samson, R., 2010. Considering Time in LCA: Dynamic LCA and Its Application to Global Warming Impact Assessments. Environ. Sci. Technol. 44, 3169–3174. https://doi.org/10.1021/es9030003

Liu, P., Zha, T., Black, T.A., Jassal, R.S., Jia, X., Noormets, A., Ouimette, A., Tian, Y., Li, X., 2025. Seasonal warming responses of the carbon dioxide sink from northern forests are sensitive to stand age. Commun. Earth Environ. 6, 43. https://doi.org/10.1038/s43247-025-02008-7

Luyssaert, S., Inglima, I., Jung, M., Richardson, A.D., Reichstein, M., Papale, D., Piao, S.L., Schulze, E.-D., Wingate, L., Matteucci, G., Aragao, L., Aubinet, M., Beer, C., Bernhofer, C., Black, K.G., Bonal, D., Bonnefond, J.-M., Chambers, J., Ciais, P., Cook, B., Davis, K.J., Dolman, A.J., Gielen, B., Goulden, M., Grace, J., Granier, A., Grelle, A., Griffis, T., Grünwald, T., Guidolotti, G., Hanson, P.J., Harding, R., Hollinger, D.Y., Hutyra, L.R., Kolari, P., Kruijt, B., Kutsch, W., Lagergren, F., Laurila, T., Law, B.E., Le Maire, G., Lindroth, A., Loustau, D., Malhi, Y., Mateus, J., Migliavacca, M., Misson, L., Montagnani, L., Moncrieff, J., Moors, E., Munger, J.W., Nikinmaa, E., Ollinger, S.V., Pita, G., Rebmann, C., Roupsard, O., Saigusa, N., Sanz, M.J., Seufert, G., Sierra, C., Smith, M.-L., Tang, J., Valentini, R., Vesala, T., Janssens, I.A., 2007. CO2 balance of boreal, temperate, and tropical forests derived from a global database. Glob. Change Biol. 13, 2509–2537. https://doi.org/10.1111/j.1365-2486.2007.01439.x

IPCC, 2021. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K., Waterfield, T., Yelekçi, Ö., Yu, R., Zhou, B. (eds.). Cambridge University Press, Cambridge, UK & New York, USA. https://doi.org/10.1017/9781009157896

Matthews, H.D., Caldeira, K., 2008. Stabilizing climate requires near-zero emissions. Geophys. Res. Lett. 35. https://doi.org/10.1029/2007GL032388

Meinshausen, M., Nicholls, Z.R.J., Lewis, J., Gidden, M.J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J.G., Daniel, J.S., John, A., Krummel, P.B., Luderer, G., Meinshausen, N., Montzka, S.A., Rayner, P.J., Reimann, S., Smith, S.J., van den Berg, M., Velders, G.J.M., Vollmer, M.K., Wang, R.H.J., 2020. The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. Geosci. Model Dev. 13, 3571–3605. https://doi.org/10.5194/gmd-13-3571-2020

Meinshausen, M., Raper, S.C.B., Wigley, T.M.L., 2011. Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1: Model description and calibration. Atmospheric Chem. Phys. 11, 1417–1456. https://doi.org/10.5194/acp-11-1417-2011

Metzler, H., Launiainen, S., Vico, G., 2024. Amount of carbon fixed, transit time and fate of harvested wood products define the climate change mitigation potential of boreal forest management—A model analysis. Ecol. Model. 491, 110694. https://doi.org/10.1016/j.ecolmodel.2024.110694

Moreau, G., Chagnon, C., Achim, A., Caspersen, J., D'Orangeville, L., Sánchez-Pinillos, M., Thiffault, N., 2022. Opportunities and limitations of thinning to increase resistance and resilience of trees and forests to global change. For. Int. J. For. Res. 95, 595–615. https://doi.org/10.1093/forestry/cpac010

Nabuurs, G.-J., Delacote, P., Ellison, D., Hanewinkel, M., Hetemäki, L., Lindner, M., 2017. By 2050 the Mitigation Effects of EU Forests Could Nearly Double through Climate Smart Forestry. Forests 8, 484. https://doi.org/10.3390/f8120484

Nagel, R., Meyer, P., Blaschke, M., Feldmann, E., 2023. Strict forest protection: A meaningful contribution to Climate-Smart Forestry? An evaluation of temporal trends in the carbon balance of unmanaged forests in Germany. Front. For. Glob. Change 6. https://doi.org/10.3389/ffgc.2023.1099558

Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E.J., Mathesius, U., Poot, P., Purugganan, M.D., Richards, C.L., Valladares, F., van Kleunen, M., 2010. Plant phenotypic plasticity in a changing climate. Trends Plant Sci. 15, 684–692. https://doi.org/10.1016/j.tplants.2010.09.008

Noormets, A., Epron, D., Domec, J.C., McNulty, S.G., Fox, T., Sun, G., King, J.S., 2015. Effects of forest management on productivity and carbon sequestration: A review and hypothesis. For. Ecol. Manag., Carbon, water and nutrient cycling in managed forests 355, 124–140. https://doi.org/10.1016/j.foreco.2015.05.019

Oliver, R.J., Mercado, L.M., Clark, D.B., Huntingford, C., Taylor, C.M., Vidale, P.L., McGuire, P.C., Todt, M., Folwell, S., Shamsudheen Semeena, V., Medlyn, B.E., 2022. Improved representation of plant physiology in the JULES-vn5.6 land surface model: photosynthesis, stomatal conductance and thermal acclimation. Geosci. Model Dev. 15, 5567–5592. https://doi.org/10.5194/gmd-15-5567-2022

Olšiaková, M., Loučanová, E., Paluš, H., 2016. Monitoring changes in consumer requirements for wood products in terms of consumer behavior. Acta Facultatis Xylologiae Zvolen 58(1), 15‑25. https://doi.org/10.17423/afx.2016.58.1.15

O'Neill, B.C., Kriegler, E., Riahi, K., Ebi, K.L., Hallegatte, S., Carter, T.R., Mathur, R., van Vuuren, D.P., 2014. A new scenario framework for climate change research: the concept of shared socioeconomic pathways. Clim. Change 122, 387–400. https://doi.org/10.1007/s10584-013-0905-2

Patacca, M., Lindner, M., Lucas-Borja, M.E., Cordonnier, T., Fidej, G., Gardiner, B., Hauf, Y., Jasinevičius, G., Labonne, S., Linkevičius, E., Mahnken, M., Milanovic, S., Nabuurs, G.-J., Nagel, T.A., Nikinmaa, L., Panyatov, M., Bercak, R., Seidl, R., Ostrogović Sever, M.Z., Socha, J., Thom, D., Vuletic, D., Zudin, S., Schelhaas, M.-J., 2023. Significant increase in natural disturbance impacts on European forests since 1950. Glob. Change Biol. 29, 1359–1376. https://doi.org/10.1111/gcb.16531

Pilli, R., Alkama, R., Cescatti, A., Kurz, W.A., Grassi, G., 2022. The European forest carbon budget under future climate conditions and current management practices. Biogeosciences 19, 3263–3284. https://doi.org/10.5194/bg-19-3263-2022

Pingoud, K., Perälä, A.-L., Soimakallio, S., Pussinen, A., 2003. Greenhouse gas impacts of harvested

1149 wood products: Evaluation and development of methods, RESEARCH NOTES. VTT
1150 TIEDOTTEITA.
1151

1152 Pingoud, K., Perälä, A.-L., Soimakallio, S., Pussinen, A., 2003. Greenhouse gas impacts of harvested
1153 wood products: Evaluation and development of methods. VTT Tiedotteita – Research Notes 2189.
1154 Espoo: VTT Technical Research Centre of Finland, 79 pp.
1155

1156 Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B.L., Dietrich,
1157 J.P., Doelmann, J.C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabeau, A., Takahashi, K.,
1158 Valin, H., Waldhoff, S., Weindl, I., Wise, M., Kriegler, E., Lotze-Campen, H., Fricko, O., Riahi, K.,
1159 Vuuren, D.P. van, 2017. Land-use futures in the shared socio-economic pathways. Glob. Environ.
1160 Change 42, 331–345. https://doi.org/10.1016/j.gloenvcha.2016.10.002
1161

1162 Puchi, P.F., Dalmonech, D., Castagneri, D., Genovese, G., Helgason, W., Khomik, M., Brilli, L.,
1163 Collalti, A., 2026. Decoding carbon allocation in boreal forests: Integrating multi-proxy observations
1164 and process-based modelling. Agric. For. Meteorol. 378, 110923.
1165 https://doi.org/10.1016/j.agrformet.2025.110923
1166

1167 Pukkala, T., 2017. Does management improve the carbon balance of forestry? For. Int. J. For. Res. 90,
1168 125–135. https://doi.org/10.1093/forestry/cpw043
1169

1170 Qu Y, Mueller-Cajar O, Yamori W, 2023. Improving plant heat tolerance through modification of
1171 Rubisco activase in C3 plants to secure crop yield and food security in a future warming world. J.
1172 Exp. Bot. 74. https://doi.org/10.1093/jxb/erac340
1173

Reyer, C.P.O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-Born, P.,
Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders, T.G.M., Kolari,
P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., Collalti, A., Trotta, C., Matteucci, G.,
D'Andrea, E., Foltýnová, L., Krejza, J., Ibrom, A., Pilegaard, K., Loustau, D., Bonnefond, J.-M.,
Berbigier, P., Picart, D., Lafont, S., Dietze, M., Cameron, D., Vieno, M., Tian, H., Palacios-Orueta,
A., Cicuendez, V., Recuero, L., Wiese, K., Büchner, M., Lange, S., Volkholz, J., Kim, H., Horemans,
J.A., Bohn, F., Steinkamp, J., Chikalanov, A., Weedon, G.P., Sheffield, J., Babst, F., Vega del Valle,
I., Suckow, F., Martel, S., Mahnken, M., Gutsch, M., Frieler, K., 2020. The PROFOUND Database
for evaluating vegetation models and simulating climate impacts on European forests. Earth Syst. Sci.
Data 12, 1295–1320. https://doi.org/10.5194/essd-12-1295-2020
1184

Riahi, K., van Vuuren, D.P., Kriegler, E., Edmonds, J., O'Neill, B.C., Fujimori, S., Bauer, N., Calvin,
K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J.C., Kc, S., Leimbach, M., Jiang, L.,
Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L.A.,
Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L.,
Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J.C., Kainuma, M.,
Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., Tavoni, M., 2017. The
Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions
implications: An overview. Glob. Environ. Change 42, 153–168.
https://doi.org/10.1016/j.gloenvcha.2016.05.009
1194

1195 Rogelj, J., Popp, A., Calvin, K.V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S., Strefler, J.,
1196 Hasegawa, T., Marangoni, G., Krey, V., Kriegler, E., Riahi, K., van Vuuren, D.P., Doelman, J.,
1197 Drouet, L., Edmonds, J., Fricko, O., Harmsen, M., Havlík, P., Humpenöder, F., Stehfest, E., Tavoni,
1198 M., 2018. Scenarios towards limiting global mean temperature increase below 1.5 °C. Nat. Clim.
1199 Change 8, 325–332. https://doi.org/10.1038/s41558-018-0091-3
1200

1201 Rüter, S., Di Fulvio, F., 2025. Report on HWP contribution potentials to EU forest sector mitigation
1202 (Deliverable D5.2). ForestNavigator Project (Horizon Europe, Grant Agreement No. 101056875).

1203 Available at: https://www.forestnavigator.eu/wp-content/uploads/FN_D5.2_final.pdf (accessed 18
1204 December 2023).

1205

1206 Rüter, S., Werner, F., Forsell, N., Prins, C., Vial, E., Levet, A.-L., 2016. *ClimWood2030 –*
1207 *Climate benefits of material substitution by forest biomass and harvested wood products:*
1208 *Perspective 2030.* Thünen Report 42. Braunschweig: Johann Heinrich von Thünen
1209 Institute. https://doi.org/10.3220/REP1480332042000

1210

1211 Saponaro, V., Dalmonech, D., Vangi, E., Puchi, P.F., Rezaie, N., D'Andrea, E., Tomelleri, E.,
1212 Collalti, A., 2025. Climate change, more than management, drives short- and long-term changes in
1213 iWUE in a sub-Alpine beech forest. J. For. Res. 37, 16. https://doi.org/10.1007/s11676-025-01942-8

1214

1215 Sathre, R., O'Connor, J., 2010. Meta-analysis of greenhouse gas displacement factors of wood
1216 product substitution. Environ. Sci. Policy 13, 104–114. https://doi.org/10.1016/j.envsci.2009.12.005

1217

1218 Schulze, E.D., Sierra, C.A., Egenolf, V., Woerdehoff, R., Irslinger, R., Baldamus, C., Stupak, I.,
1219 Spellmann, H., 2020. The climate change mitigation effect of bioenergy from sustainably managed
1220 forests in Central Europe. GCB Bioenergy 12, 186–197. https://doi.org/10.1111/gcbb.12672

1221

1222 Schulze, E.D., Valentini, R., Bouriaud, O., 2021. The role of net ecosystem productivity and of
1223 inventories in climate change research: the need for "net ecosystem productivity with harvest",
1224 NEPH. For. Ecosyst. 8, 15. https://doi.org/10.1186/s40663-021-00294-z

1225

1226 Seidl, R., Fernandes, P.M., Fonseca, T.F., Gillet, F., Jönsson, A.M., Merganičová, K., Netherer, S.,
1227 Arpaci, A., Bontemps, J.-D., Bugmann, H., González-Olabarria, J.R., Lasch, P., Meredieu, C.,
1228 Moreira, F., Schelhaas, M.-J., Mohren, F., 2011. Modelling natural disturbances in forest ecosystems:
1229 a review. Ecol. Model. 222, 903–924. https://doi.org/10.1016/j.ecolmodel.2010.09.040

1230

1231 Sierra, C.A., Crow, S.E., Heimann, M., Metzler, H., Schulze, E.-D., 2021. The climate benefit of
1232 carbon sequestration. Biogeosciences 18, 1029–1048. https://doi.org/10.5194/bg-18-1029-2021
1233 Skytt, T., Englund, G., Jonsson, B.-G., 2021. Climate mitigation forestry—temporal trade-offs.
1234 Environ. Res. Lett. 16, 114037. https://doi.org/10.1088/1748-9326/ac30fa

1235

1236 Strassmann, K.M., Joos, F., 2018. The Bern Simple Climate Model (BernSCM) v1.0: an extensible
1237 and fully documented open-source re-implementation of the Bern reduced-form model for global
1238 carbon cycle–climate simulations. Geosci. Model Dev. 11, 1887–1908. https://doi.org/10.5194/gmd-
1239 11-1887-2018

1240

1241 Suter, F., Steubing, B., Hellweg, S., 2017. Life Cycle Impacts and Benefits of Wood along the Value
1242 Chain: The Case of Switzerland. J. Ind. Ecol. 21, 874–886. https://doi.org/10.1111/jiec.12486
1243 Tanaka, K., O'Neill, B.C., 2018. The Paris Agreement zero-emissions goal is not always consistent
1244 with the 1.5 °C and 2 °C temperature targets. Nat. Clim. Change 8, 319–324.
1245 https://doi.org/10.1038/s41558-018-0097-x

1246

1247 Travaglini, D., 2009. Piano di gestione e Silvomuseo 2006–2025 della Foresta di Vallombrosa:
1248 compartimentazione e cartografia. Technical report, Università degli Studi di Firenze - Dipartimento
1249 di Gestione dei Sistemi Agrari, Alimentari e Forestali, Firenze, Italy.

1250

1251 Vacek, Z., Cukor, J., Vacek, S., Gallo, J., Bažant, V., Zeidler, A., 2023. Role of black pine (Pinus
1252 nigra J. F. Arnold) in European forests modified by climate change. Eur. J. For. Res. 142, 1239–1258.
1253 https://doi.org/10.1007/s10342-023-01605-5

1254

1255 Valentini, R., Miglietta, F., 2015. The greenhouse gas balance of Italy. An insight on managed and
1256 natural terrestrial ecosystems. https://doi.org/10.1007/978-3-642-32424-6

van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S.J., Rose, S.K., 2011. The representative concentration pathways: an overview. Clim. Change 109, 5. https://doi.org/10.1007/s10584-011-0148-z

Vangi, E., Sacchelli, S., Nocentini, S., Plutino, M., Dalmonech, D., Collalti, A., Travaglini, D., Corona, P., 2026. Enhancing ecosystem service provision through the silvicultural management of European black pine stands from afforestation and reforestation in Italy. J. For. Res. 37, 44. https://doi.org/10.1007/s11676-025-01975-z

Wei, X., Zhao, J., Hayes, D.J., Daigneault, A., Zhu, H., 2023. A life cycle and product type based estimator for quantifying the carbon stored in wood products. Carbon Balance Manag. 18, 1. https://doi.org/10.1186/s13021-022-00220-y

Zhang, Y., Piao, S., Sun, Y., Rogers, B.M., Li, X., Lian, X., Liu, Z., Chen, A., Peñuelas, J., 2022. Future reversal of warming-enhanced vegetation productivity in the Northern Hemisphere. Nat. Clim. Change 12, 581–586. https://doi.org/10.1038/s41558-022-01374-w

Credit authorship contribution statement

Issam Boukhris: Writing - original draft**,** Writing - review & editing, Methodology, Visualization, Validation, Investigation, Data curation, Conceptualization, Project administration. **Francesco Cherubini:** Writing - review & editing, Methodology, Resources, Formal analysis, Resources. **Alessio Collalti:** Writing - review & editing, Validation, Software, Formal analysis, Resources. **Daniela Dalmonech:** Validation, Software, Resources. **Christian Vonderach:** Writing - review & editing, Investigation, Validation, Formal analysis. **Francesca Giannetti**: Writing - review & editing, Resources, Formal analysis. **Gina Marano:** Writing - review & editing, Methodology, Formal analysis **Said Lahssini:** Writing - review & editing, Formal analysis, Software. **Monia Santini:** Writing - review & editing, Software, Funding acquisition. **Riccardo Valentini:** Formal analysis, Supervision, Funding acquisition.

Funding sources

Issam Boukhris was supported by the National Research Center for Agricultural Technologies —AGRITECH— PNRR (Italian National Plan of Recovery and Resilience), CN00000022 in WP 4.3 Task 3 Risk management strategies and policies in the context of climate change to carry out multiple field missions, during which he collected essential data in the Vallombrosa Forest in Tuscany, Italy. Alessio Collalti and Daniela Dalmonech acknowledge funding by the European Union – NextGenerationEU under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of December 16, 2021, rectified by Decree n.3175 of December 18, 2021 of the Italian Ministry of University and Research under award Number: Project code CN_00000033, Concession Decree No. 1034 of June 17, 2022 adopted by the Italian Ministry of University and Research, CUP B83C22002930006, Project title "National Biodiversity Future Centre - NBFC".

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability Statement

Data are available from the corresponding author upon reasonable request.

Acknowledgment

Issam Boukhris would like to thank the University of Tuscia and the CMCC Foundation, both based in Italy, for supporting his Ph.D. thesis, of which this work is a part. He also extends his gratitude to Professor Gherardo Chirici and Dr. Elia Vangi from the University of Florence, Vincenzo Saponaro from the CNR of Italy, Colonel Giuliano Savelli and Maréchal Giovanni Galipò from the local forest services of Vallombrosa for their technical and logistical support. Special thanks are also due to Dr. Khalil Misbah from the IRD of Toulouse, as well as to Adnan Yousaf and Renato Zompanti from the University of Tuscia, for their invaluable assistance during fieldwork.