Title: Long-term forest-sector mitigation and radiative forcing under

contrasting management, climate, and substitution pathways

Running Title: Forest sector mitigation pathways

I. Boukhris[1,2*], F. Cherubini[3], A. Collalti[4,5], D. Dalmonech[4,5], C.

Vonderach[6], G. Marano[7], F. Giannetti[8], S. Lahssini[9], M. Santini[2], R.

Valentini[1,2]

1. Department for Innovation in Biological, Agri-Food and Forest Systems (DIBAF),

University of Tuscia, 01100, Viterbo, Italy (issam.boukhris@unitus.it; rik@unitus.it)

2. CMCC Foundation — Euro-Mediterranean Center on Climate Change, Lecce, Italy

(monia.santini@cmcc.it)

3. Industrial Ecology Programme, Department of Energy and Process Engineering,

Norwegian University of Science and Technology (NTNU), 7034 Trandheim, Norway

(francesco.cherubini.ntnu.no)

4. Forest modeling Lab, Institute for Agriculture and Forest Systems in the Mediterranean,

National Research Council of Italy (CNR-ISAFOM), 06128, Perugia, Italy

(alessio.collalti@cnr.it; daniela.dalmonech@cnr.it)

5. National Biodiversity Future Center (NBFC), 90133, Palermo, Italy

6. Forest Research Institute Baden-Wurttemberg, 79100, Freiburg, Germany

(christian.vonderach@forst.bwl.de)

7. Department of Environmental Systems Science, Forest Ecology, Institute of Terrestrial

Ecosystems, ETH Zurich, Zurich, Switzerland (gina.marano@usys.ethz.ch)

8. Department of Agriculture, Food, Environment and Forestry (DAGRI), University of

Florence, Florence, Italy

9. Department of Forest Development, National School of Forestry Engineering, 11000, Salé,

Morocco (marghadi@gmail.com)

*issam.boukhris@unitus.it

phone: +39 327 328 2507

30
Abstract

Forests are central to the EU's climate neutrality strategy, currently offsetting ~9% of total

greenhouse gas emissions and offering further mitigation potential through harvested wood

products and the substitution effect. However, the climate benefit of the forest sector is

influenced by multiple interacting factors, including forest management, climate change,

wood-use strategies, and assumptions about substitution benefits, as well as the timing and

fate of carbon across the forest ecosystem and technosphere. To evaluate these drivers, we

used a coupled forest growth and a wood products model to simulate five different

silvicultural strategies under three climate change scenarios, four wood use schemes, and five

displacement factor decay pathways over a 285-year period (2015–2300), applied to a *Pinus

nigra* forest in Italy, questioning the impact of these factors on climate mitigation potential of

the forest sector. We assessed forest sector balance (FSB, net carbon exchange between forest

system and atmosphere), radiative forcing from biogenic CO_2 (RF_{bio}), and mitigation

efficiency (ME) – the proportion of sequestered carbon contributing to net climate benefit.

Results showed that FSB and RF_{bio} were "broadly" aligned, but ME varied with the

magnitude and duration of biogenic emissions. The scenarios BIOE (bioenergy) and TM

(modular cutting) achieved high FSB but showed lower ME due to concentrated or sustained

emissions. WOOD and ADAPT yielded higher ME under SSP1-2.6, while several strategies

(WOOD, ADAPT, TRANS) became net sources under SSP5-8.5 after 2200. Substitution

benefits declined under degressive assumptions, reducing mitigation by up to 53% especially

for high-harvest scenarios. FSB was primarily shaped by climate and management, secondly

by substitution, however; wood-use strategies had no significant long-term effect provided they did not impact resource availability. Together, these findings underscore that effective forest-sector mitigation requires not only maximizing cumulative carbon stocks, but also minimizing the magnitude, timing, and atmospheric residence time of emissions while carefully considering the role of substitution benefits.

Keywords

Forest management, Climate mitigation, Wood substitution, Carbon balance, Radiative Forcing, Mitigation efficiency, Climate change, Process-based model, Wood products model

Background

Forests are a key component of the EU's climate mitigation framework for achieving carbon

neutrality by 2050. Currently they remove approximately 9% of total greenhouse gas (GHG)

emissions from the atmosphere (EEA, 2022). Forest ecosystems serve as both carbon sinks,

sequestering additional carbon, and reservoirs, storing it across several pools. When managed

sustainably, forest carbon can be transferred laterally to harvested wood products (HWPs),

establishing an external carbon pool and contributing to the substitution effect, as HWPs

replace energy-intensive materials and fossil fuels, avoiding emissions (Rüter & Fulvio,

2025). In the EU, forests and HWPs together sequester approximately 400 Mt CO_2-eq.yr^{-1},

with HWPs contributing for about 10% to this total. Substitution provides a similar level of

contribution with values ranging from 18 to 43 Mt CO_2-eq.yr^{-1}, further enhancing the forest

sector's overall mitigation potential (Jonsson et al., 2021; Nabuurs et al., 2017; Rüter S. et al.,

2016).To assess forest sector mitigation, a system-wide perspective is needed to account for

the distinct roles of carbon pools. This approach ensures that all relevant mitigation

pathways, sequestration, storage, and substitution, are critically evaluated (Grassi et al., 2021;

Lemprière et al., 2013).

Forest-sector mitigation outcomes are shaped by multiple interacting factors. Understanding

these drivers is essential to accurately assess the effectiveness and trade-offs among

strategies. The forest carbon sink is primarily determined by both forest management and

climate conditions (Ameray et al., 2021; Dye et al., 2024; Gregor et al., 2024; Liu et al.,

2025). Similarly, the HWPs sink is affected by these factors but also depends on wood-use

characteristics and the carbon footprint of the displaced material, which ultimately

determines the substitution effect (Bozzolan et al., 2024; Brunet-Navarro et al., 2017;

Budzinski et al., 2020). Beyond biophysical and technological factors, temporal aspects of

carbon fluxes also play a critical role, a point further developed below.

Among these factors, forest management has received particular attention (Valentini &

Miglietta, 2015). Practices range from passive conservation to intensive silvicultural

interventions (Barredo et al., 2024; Dalmonech et al., 2022; Pilli et al., 2022). An ongoing

debate in the EU revolves on whether active or reduced management contributes more

effectively to climate mitigation (Luyssaert et al., 2007; Noormets et al., 2015). In managed

forests, early interventions such as tending and thinning are used to improve wood quality,

promote tree growth, and reduce the likelihood of natural mortality. As a result, more wood

can be harvested for use in HWPs instead of being lost to natural decay. At the same time,

these operations enhance forest resilience to disturbances, which are becoming more frequent

due to climate change (Barrette et al., 2023; Kilpeläinen & Peltola, 2022; Moreau et al.,

2022). In the long term, while unmanaged forests tend to maintain higher carbon stocks,

especially in the dead wood pool (Nagel et al., 2023), studies suggest that managed forests

can achieve a superior overall carbon balance when accounting for the substitution effect

(Krug et al., 2012; Schulze et al., 2020).

Differences in the mitigation potential of multiple active management strategies exist and are

primarily driven by factors such as structure complexity, rotation length, thinning intensity,

and thinning intervals. In parallel, at least two key factors influence the HWPs sink: the

recycling rate and the lifespan of HWPs, both of which delay carbon emissions to the

atmosphere through combustion or decomposition in disposal sites (Brunet-Navarro et al.,

2017; Pingoud et al., 2003; Wei et al., 2023). Finally, one of the most critical assumptions in

estimating the forest sector's substitution effect is the assumption of stationary displacement

factors (DF). Assuming fixed DF fails to account for potential improvements in energy

efficiency and reductions in the carbon footprint of competing sectors (e.g., steel, concrete,

and energy) that may lead to reductions in GHG emissions. Any advancements in these

sectors would consequently diminish the substitution effect of wood (Brunet-Navarro et al.,

2021; Harmon, 2019; Howard et al., 2021). Therefore, assessing alternative strategies with a

fixed DF reflects only the maximum substitution potential, while the actual effect is likely

lower and often overestimated in global reports.

A critical yet often overlooked aspect of climate mitigation assessments is the timing of

carbon fluxes and how this aligns with the time horizon of mitigation projects. As Levasseur

(2010) highlighted, "*carbon neutrality does not equate climate neutrality*", as carbon persists

in the atmosphere before being reabsorbed, contributing in the meantime to the climate

forcing (Levasseur et al., 2010). Forest-Technosphere systems that release carbon rapidly but

sequester it slowly can have a stronger short-term climate impact than those with faster

sequestration rates and slower carbon release (Holtsmark, 2012; Laganière et al., 2017).

These short-term emissions influence radiative forcing, potentially intensifying near-term

warming and affecting long-term climate stability. Similarly, the time horizon; the timeframe

over which forest systems are evaluated, significantly shapes their assessed climate impact.

Over short time frames, immediate emissions from harvesting may outweigh sequestration

benefits, resulting in a higher net climate impact. In contrast, longer horizons allow forest

regrowth along with the other climate system components to offset initial emissions,

improving mitigation outcomes (Cherubini et al., 2011; Holtsmark, 2013; Sierra et al., 2021).

Given this, assessing the mitigation potential of forests based on carbon balances alone is

insufficient; it must also account for their contribution to radiative forcing over time. To

capture these temporal dynamics, we first applied a timing-sensitive indicator (RF_{bio}) which

integrates the net carbon fluxes over time using a global impulse response function. Yet,

while RF_{bio} captures the climate impact, it does not indicate how efficiently carbon is

processed to achieve that outcome. To address this, we introduce the concept of mitigation

efficiency (ME), a metric that reflects the climate benefit delivered per unit of carbon cycled

through the forest sector. Together, RF_{bio} and ME offer a more nuanced basis for evaluating

forest-sector strategies, emphasizing not just how much carbon is removed, but how

effectively and durably that removal contributes to climate mitigation.

Despite the well-established role of forests and HWPs in climate mitigation, significant

uncertainties remain regarding the long-term effectiveness of different forest management

strategies. These uncertainties arise from the complex interactions among forest growth,

harvest intensity, product use, and substitution dynamics, as well as from simplifying

assumptions, such as stationary DF. Moreover, most studies focus on short-term carbon

balances, overlooking the long temporal cycles of forest systems and the fact that mitigation

potential may emerge at different timescales depending on the applied management (Metzler

et al., 2024; Skytt et al., 2021). To address these gaps, we developed an integrated modeling

framework combining a process-based forest model with a wood product model (WPM). We

evaluated five different forest management strategies under three climate change scenarios

and five displacement pathways over a 285-year simulation period, using a factorial design.

The analysis also examines whether high carbon balances consistently correspond to strong

climate outcomes, and how time horizons, climate trajectories, and decay of DF influence

long-term forest-sector mitigation performance. We applied the modeling framework to a

Pinus nigra J.F. Arnold stand in the Vallombrosa Forest, Tuscany, Italy, representative of

temperate European forest ecosystems. *P. nigra* was selected due to its wide ecological

adaptability, increasing use in European reforestation efforts, and its relevance to climate-

resilient forestry (EU council, 2025; Vacek et al., 2023).

Materials and Methods

1. Study area

The study was conducted in the state forest of Vallombrosa (43° 44' N, 11° 34'), a biogenetic

reserve of approximately 1000 ha in Tuscany, Italy. The forest experiences a mean annual

precipitation of about 1195 mm and a mean annual temperature slightly higher than 10 °C.

January is the coldest month (1.9 °C), July the warmest (19.6 °C). All the conifer plantations

were traditionally managed as even-aged. *Pinus nigra* was introduced in the forest (Bottalico

et al., 2012) as part of the main reforestation program that occurred after the second World

War. These stands cover 11% of the total forest area, with ages ranging from 60 to 95 years.

From 2005, as reported in the Vallombrosa forest management plan (*Piano di gestione e*

silvomuseo) (Travaglini, 2009). *Pinus nigra* stands are managed under close-to-nature

silviculture principles following the systemic silviculture approach (Ciancio & Nocentini,

2011), with the principles of modular/selective continuous cutting. This approach rooted in

Continuous Cover Forestry (CCF) and prescribed by Ciancio (2004) promotes resource

sustainability and natural regeneration by dynamically reassessing the harvestable volume

(Ciancio & Nocentini, 2004). The silvicultural systemic method for growing stock harvesting

calculates the real wood provision over a 5-year period. Thinning is allowed only if the

remaining standing volume is kept above the minimum threshold (typically 100-150 m^3 ha^{-1}

for *Pinus nigra*). If the condition is not met, only the portion of increment that preserves the

minimum stock is thinned. Thinning is also carried out to remove trees at high risk of falling.

When harvesting is conducted, the resulting residues are left on site. To monitor forest yield

and health, and in the context of the forest management plan, local permanent forest

inventory plots were established. The permanent circular plots (530 m²) are identical to those

of the Italian National Forest Inventory (NFI; https://www.inventarioforestale.org) and were

surveyed in 2005. Several of these plots are located within *Pinus nigra* stands and were

resurveyed again in 2017, 2018, and 2023.

2. Study key silvicultural and modeling assumptions

The following assumptions are considered in this study: the selected forest stand is assumed

to be representative of the average conditions across the broader *Pinus nigra* landscape. The

same forest management practices and wood use behavior are consistently

applied/maintained throughout the study period. Additionally, species composition is

assumed to remain unchanged, meaning no species conversion occurs over the time horizon.

Finally, each management scenario assumes a starting point of a recent plantation established

in 2015 following a clear-cut, with a planting density of 2500 trees per hectare.

In addition, landfilling activities were assumed to gradually phase out by the year 2060,

following an exponential decay model calibrated on national data. This estimate is based on

the observed reduction in the share of Italian municipalities utilizing landfill, which declined

from 46% in 2010 to 18% in 2022 (EEA, 2024). By assuming a constant relative rate of

decrease, landfilling is projected to approach zero use by 2060, and this cutoff year is adopted

as the upper limit for landfill emissions modeling.

3. Definition of scenarios

3.1. Climate change pathways

We investigated two coupled representative concentration pathways (RCPs) and shared

socioeconomic pathways (SSPs) based on distinct emission scenarios to assess potential

future climate trajectories and their socioeconomic implications. The SSP1-RCP2.6 scenario

(SSP1-2.6 hereafter) represents a world of sustainability-focused growth and equality, and it

assumes that strong climate protection measures are implemented, aiming to limit global

warming to well below 2°C above pre-industrial levels (O'Neill et al., 2014; Riahi et al.,

2017; van Vuuren et al., 2011). In contrast, SSP5-RCP8.5 (SSP5-8.5 hereafter) reflects a

world of rapid and unconstrained growth in economic output, energy demand and use, as well

as limited to no mitigation actions and a persistent reliance on fossil fuels (Calvin et al.,

2017; Meinshausen et al., 2011; Rogelj et al., 2018). Climate projections were obtained from

the Coupled Model Intercomparison (CMIP 6) project (Eyring et al., 2016), covering the

period from 2015 to 2300. Data were sourced from the MRI-ESM2-0 global climate model,

as it was the only model available that simulated the variables of interest over the long term,

extending to the year 2300. The outputs were downscaled to a 0.25° resolution using the

perfect prognosis downscaling method. A set of statistical models and algorithms was

initially calibrated using historical (observed) data from both coarse predictors (ERA5

reanalysis; Hersbach et al., 2020) and local predictand data (E-OBS reanalysis, including

precipitation, maximum, and minimum temperature; Cornes et al., 2018) over a

representative 30-year climate period (1991–2020). The methods were then validated and

evaluated, with the best-performing models selected to downscale daily precipitation,

maximum, and minimum daily temperatures. Moreover, atmospheric CO_2 concentrations for

different climate trajectories were obtained from (Meinshausen et al., 2020). Additionally, we

simulated the current climate (CUR) for the period 2020–2300 by randomly sampling each

day of the year from the local station records (Vallombrosa, locality of Reggello) over the

period 2005–2023 (see **Figure 1**).

Figure 1: Climate projections from the downscaled MRI-ESM2-0 model under the SSP1-2.6

and SSP5-8.5 scenarios, driving the simulation for the period 2015–2300. The top panel

shows annual precipitation (mm), the middle panel displays the annual mean of daily

maximum and minimum temperatures (°C), and the bottom panel shows atmospheric CO_2

concentration (ppm), labeled as Ca.

3.2. Forest management scenarios

Five management scenarios were developed, grounded in five policies outlined in the EU

Forest Strategy (European Commision, 2021), the EU Bioeconomy Strategy (European

Commision, 2018), and the EU Forest Biodiversity Strategy (European Commission, 2021).

These scenarios were designed to reflect the specific characteristics of the target species and

the study area. In this process, the regulations for *P. nigra* management in the Tuscany region

were also considered (refer to Cantiani P et al., 2018 for further details). Each policy was

subsequently translated into stand-based forest management plans with the expert input of

local stakeholders, including forest managers and academic scholars.

The silvicultural interventions applied across the scenarios include pre-thinning and thinning

operations, implemented with varying intensities and intervals. In shelterwood-based

scenarios, additional interventions such as preparatory cuts and seedling cuts are employed to

facilitate natural regeneration under partial canopy cover. Final cuts are applied in both

shelterwood and clearcut systems, serving to remove the remaining overstory once

regeneration is established in shelterwood, or to harvest the entire stand at once in

clearcutting scenarios. In the latter case, artificial regeneration follows harvest operations,

with planting carried out at a density of 2500 seedlings per hectare (See **Figure 2** for detailed

silvicultural intervention schedules).

The first scenario is known as '*Tagli Modulari'* which represents modular cutting (hereafter

TM), and reflecting the typical practices applied in the Vallombrosa forest and described in

detail in Section 1. The "Wood-Based Bioenergy Production" (BIOE) assumes that the EU is

transitioning to a more sustainable energy future by gradually promoting the use of wood

bioenergy as a replacement for fossil fuel energy. While direct use of wood logs for energy is

discouraged under the EU's Renewable Energy Directive (RED II and RED III), only the

portion of trees unsuitable for higher-value uses are directly allocated to bioenergy.

Practically, this management plan consists of a short rotation (70-80 yrs) and the application

of 1-3 thinning treatments. The "Promotion of long-lived HWPs" (WOOD) explores how

forest management can support climate mitigation by increasing the supply of high-quality

timber suitable for durable HWPs. The scenario assumes that domestic wood use in the EU

shifts increasingly toward long-lived applications such as construction materials, which can

store carbon for extended periods and substitute for more carbon-intensive materials. While

289 the lifespan of HWPs also depends on post-harvest factors, such as recycling systems and

290 end-use, this scenario emphasizes the forest management component that facilitates the

291 availability of timber suitable for such uses. On the forest side, this objective is supported by

292 extending rotation lengths (e.g., 120-150 yrs) aiming to harvest trees with larger diameters

293 and higher wood quality through stronger thinning. The "Enhancing Forest Adaptation and

294 Resilience" (ADAPT) aims to strengthen the resilience of forests, particularly Mediterranean

295 forests, which are already experiencing severe drought conditions. Practically, this

296 management plan will consist of applying intensive thinning to alleviate competition on site

297 resources and adopting short rotation cycles to alleviate disturbance risks, in addition to

298 promoting artificial regeneration. The last scenario, "Transition to passive rewilding"

299 (TRANS) is designed to facilitate the shift from structured stand management toward a self-

300 regulating forest system with minimal long-term disturbance. Initial interventions are

301 intended to accelerate natural processes such as gap creation, early mortality, and canopy

302 opening, which commonly occur in the late stages of forest succession and structural

303 diversification (Bauhus et al., 2009). These actions aim to promote structural diversification

304 and regeneration. Thinning is applied at regular intervals between 2030 and 2150, with

305 intensities ranging from 30% to 50% of the stand basal area. After 2150, no further

306 interventions were applied, allowing the forest to develop autonomously under passive

307 conditions.

Figure 2: Overview of the five forest management strategies evaluated in this study, arranged along a gradient of management intensity from high (top) to low (bottom). Strategies include: BIOE, TM, WOOD, ADAPT, and TRANS. Each strategy is defined by its objective, silvicultural approach, and target stand characteristics, progressing from strategic objectives to operational implementation. The right-hand panels show the corresponding management plans for each strategy, including timing and type of intervention (i.e., Prth: pre-thinning, Thin: thinning, Prep: preparatory cut, SeedC: Seeding cut, and FinalC: final cut), percentage of trees removed (%N), percentage of basal area removed (%BA) scheduled at specific ages.

3.3. Wood use scenarios

To explore the potential implications of improved cascading through an increase in product

lifespan and recycling rates, four different wood use scenarios were developed. The first, the

BAU as "Baseline Scenario", maintained constant recycling rate and lifespan values. In the

LS20 as "Longevity Scenario", the lifespan of products was increased by 20%. The RR20 as

"Reuse Scenario" focused on increasing the recycling rate of products by 20%. Finally, the

RRLS20 "Sustainability Scenario" increased both the lifespan of products and the recycling

rate by 20% (please refer to **Table S1.5**).

3.4. Substitution scenarios

In addition to a baseline scenario (SUB0), which assumes that the DF remains constant over

time, we considered four decarbonization pathways with varying linear rates of DF reduction.

We adopt either a gradual (25% DF reduction by 2050, called SUB25), moderate (50%,

SUB50), significant (75%, SUB75) or a rapid (100%, SUB100), the latter being in line with

the EU's climate neutrality targets.

4. Modeling framework

4.1. Modelling system and key concepts

The system considered in this study includes both forest and HWPs pools, collectively

representing the forest sector. It accounts for incoming and outgoing carbon fluxes while

disregarding internal exchanges, as these remain within the system and do not affect its net

balance. The forest sector system consists of five key components: (1) Aboveground

biomass, comprising stem wood, branches, stumps, and foliage of live trees, where the focus

is on CO_2 fixation through photosynthesis and biomass accumulation; (2) Belowground

biomass, including fine and coarse roots of live trees, also assessed for CO_2 fixation; (3)

Aboveground Dead Organic Matter (ADOM), consisting of standing dead trees, downed

woody debris, litter, and humus, where carbon is gradually released through decomposition;

(4) Belowground Dead Organic Matter (BDOM), encompassing soil organic matter and dead

roots, with carbon emissions primarily occurring through microbial respiration; and (5)

HWPs, representing carbon stored in HWPs during their use phase, with CO_2 emissions

occurring over time through disposal in landfills or combustion for bioenergy (see **Figure 3**).

Beyond these carbon pools, the substitution effect plays a critical role in the forest sector's

overall climate mitigation.

The annual forest ecosystem carbon balance (FEB; tC ha^{-1} yr^{-1}) is defined as:

$$□□□ = (□□□ − □□ − □□_{□□□□□□}) − □□□□□□□□$$
$$= □□□ − □_{□□□} − □□□□□□□□$$

$$= −□□□ − □□□□□□□□ \quad (Eq.\ 1)$$

where GPP represents the gross primary production; Ra, $□□_{□□□□□□□}$, and $□_{□□□}$ correspond

to autotrophic, heterotrophic, and ecosystem respirations occurring within the forest,

respectively, Harvest accounts for biomass removal through forestry operations, and NEE

represents the net ecosystem exchange within the forest. All terms are expressed in units of

tC ha^{-1} yr^{-1}.

Harvested biomass enters the wood sector, where it is temporarily stored in HWPs before

eventually being released as emissions through disposal, or combustion for bioenergy.

Therefore, the wood sector carbon balance (WSB; tC ha^{-1} yr^{-1}) is expressed as:

$$□□□ = □□□□□□□□ + □□□□□□□□□□□□□ − □□_{□□□□□□□} = □□□□□□□□ +$$

$$□□□□□□□□□ \quad (Eq.\ 2)$$

369 where Substitution represents avoided emissions resulting from the replacement of fossil

370 fuels and energy-intensive materials with HWPs, $R_{h_{wood}}$ (tC ha^{-1} yr^{-1}) refers to the

371 heterotrophic respiration occurring outside the forest, mainly from HWPs decomposition and

372 combustion, and ΔC_{wood} refers to the balance between $NPP_{to\ wood}$ and

373 $R_{h_{wood}}$, both of which are expressed in units of tC ha^{-1} yr^{-1}.

375 The annual overall forest sector balance (FSB; tC ha^{-1} yr^{-1}) integrates both forest and wood

376 sector dynamics, incorporating the mitigation benefits of substitution:

$$FSB = NBP + NWB = -R_h - R_{h_{wood}} + NPP_{to\ wood} + Substitution$$

378 $$= NPP_{to\ wood} - R_e \quad (Eq.\ 3)$$

380 The terms $R_{h_{wood}}$, and $NPP_{to\ wood}$ follow the terminology established by (Schulze et

381 al., 2021).

Figure 3: Conceptual representation of carbon flows and balances with the forest sector,

divided into the forest ecosystem (green box) and wood sector (brown box). GPP constitutes

the carbon inflow to the forest ecosystem while both Ra and Rh_{onsite} constitute the carbon

outflow from the forest ecosystem. Harvested biomass is transferred from the forest

ecosystem to the wood sector where it contributes to substitution benefits and eventually to

$Rh_{offsite}$.

4.2. Modelling framework and simulation setup

A forest sector modeling framework was developed by coupling 3D-CMCC-FEM (v5.7

BGC) (Collalti et al., 2014; Dalmonech et al., 2024), a stand-level, climate-sensitive, process-

based model, with TimberTracer (Boukhris et al., 2025) a WPM that tracks carbon stocks in

HWPs, substitution effects, and emissions from disposal sites (see **Figure 4**). Given the

complexity and level of detail associated with each model, including their structure,

parametrization, input data, and validation procedures, full technical descriptions have been

moved to the Supplementary Materials (**Tables S1.1-S1.7**) to maintain clarity and focus on

the main text.

400
Figure 4: Carbon modeling framework coupling a forest process-based model (3D-CMCC-

FEM) and TimberTracer, a WPM. DBH is the mean diameter at breast height, N is the stand

density (ha^{-1}), BA is the basal area (m^2 ha^{-1}), H is the mean tree height (m), SA is sapwood

area of the mean tree (cm^2), HA is the heartwood area of the mean tree (cm^2), and BT is the

bark thickness of the mean tree (cm).

4.3. Modeling the climate impact of the forest sector

To assess the climate impact of the forest sector, we model atmospheric CO_2 emissions decay

using an Impulse Response Function (IRF), which quantifies the temporal reaction of a

dynamic system in response to some external change (Joos et al., 2013). The multi-model

mean from the carbon cycle model intercomparison in Joos et al. (2013) is used to simulate

this decay process, incorporating a multi-exponential decay function to simulate the response

of different carbon pools, accurately reflecting the transient and long-term behavior of carbon

in the atmosphere, land, and ocean reservoirs (Cherubini et al., 2011; Joos et al., 2013;

Strassmann & Joos, 2018). Its associated IRF is mathematically expressed as:

$$IRF(t) = A_0 + \sum_1^3 A_i \, e^{-t/\tau_i} \qquad (Eq. 4)$$

where $A_0 = 0.2173$, $A_1 = 0.2240$, $A_2 = 0.2824$, $A_3 = 0.2763$, $\tau_1 = 394.4$, $\tau_2 = 36.54$, $\tau_3 = 4.304$

The amplitude A_0 represents the asymptotic airborne fraction of CO_2, which remains in the

atmosphere in the long term due to the equilibrium response of the ocean–atmosphere

system. Each A_i is associated with a characteristic residence time τ_i, which describes how

long the corresponding fraction of carbon remains in the atmosphere before being taken up

by the combined action of terrestrial and oceanic processes. Together, the coefficients

($A_0 + A_1 + A_2 + A_3$) sum to 1, and thus each A_i can be interpreted as the fractional

contribution of a given decay component to the overall atmospheric response. It is important

to note that these components do not correspond to individual, physically distinct carbon

sinks, but rather to a mathematical approximation of the combined response of the global

carbon cycle.

The atmospheric concentration of CO_2 over time can be described as the convolution of the

IRF with the net carbon flux. It is expressed as the integral of past net emissions at time t',

weighted by the fraction of CO_2 remaining in the atmosphere after a time interval t -t', where

t is the reference time point.

To compare the climate impact of biogenic CO_2 emissions across different forest-sector

scenarios, a consistent metric is required. In this paper, we developed an approach based on

the integrated radiative forcing concept adapted for biological systems, referred to here as

RF_{bio}. While originally formulated for pulse emissions in Life Cycle Assessment contexts

(e.g., GWP_{bio}; see Cherubini et al., 2011), here we extend the concept to account for net

carbon fluxes over time, making it applicable to dynamic forest-sector modeling. RF_{bio} is

derived from the time-integrated radiative forcing caused by emissions, quantifying the

perturbation of the Earth's energy balance. It is calculated as follows:

$RFbio(t) \ = \ \int_0^t -FSB(t') \ IRF(t-t') \, dt'$ (Eq. 5)

A positive RF_{bio} indicates a net warming effect. In contrast, an RF_{bio} equals to or less than

zero implies that the system has no net contribution to atmospheric warming. This condition

can therefore be interpreted as the forest-sector system being climate neutral ($RF_{bio} \approx 0$),

adds to climate warming ($RF_{bio} > 0$), or climate cooling ($RF_{bio} < 0$) over the assessed time

horizon. Figure 5 provides an illustration of these concepts.

The modeling approach described in Sections 4.2 and 4.3 is operationalized through an

integrated methodological workflow, illustrated in Figure A.1.

Figure 5: CO_2 dynamics across Earth system compartments and the carbon removal dynamics. (a) Schematic representation of CO_2 exchanges between the atmosphere, ocean and biosphere. Red arrows indicate CO_2 emissions to the atmosphere; blue arrows indicate CO_2 uptake from the atmosphere. (b) Net atmospheric CO_2 change (%) over time following a pulse emission, with equivalent carbon removal occurring at year 60. For analogy, assume that by the starting year, harvested wood is burned for bioenergy and the resulting emissions are fully captured by year 60. Response is simulated using the IRF Bern 2.5 CC model. The shaded area represents the RF_{bio}, highlighting temporary overshoot and long-term legacy effect. Positive values, denote a forcing effect while negative values denote a cooling effect.

5. Assessment of carbon and climate mitigation potentials

Carbon balance (CB) was estimated by tracking net changes in forest carbon pools and

HWPs, while also accounting for the generated substitution effects. CB was determined by

summing the FSB (see Eq. 3) over a time horizon T:

$CB(T) = -\sum_{t=1}^{T} FSB(t)$ (Eq. 6)

where $FSB(t)$ represents the annual forest sector balance at time t. The total time horizon

T was considered over 85, 185, or 285 years. The minus sign in Equation 6 reflects the

convention that a negative CB indicates a net carbon sink (i.e., net removal of CO_2 from the

atmosphere), while a positive value indicates a net source (i.e., net emissions).

Climate mitigation potential, expressed as radiative forcing due to forest management, was

assessed using the RF_{bio} metric (see Eq. 5), which accounts for the temporary nature of

biogenic CO_2 emissions and their atmospheric decay over time.

Furthermore, the concept of ME is introduced to measure the effectiveness of mitigation

efforts, is defined as the ratio between the negative RF_{bio} and the corresponding carbon

balance. The negative sign ensures that a climate-cooling effect (negative RFbio) yields a

positive ME, aligning higher ME values with greater mitigation benefit. This makes ME a

consistent and intuitive indicator of strategy performance across time horizons.

ME(T) = {

$\frac{-RFbio(T)}{CB(T)}$, *if RFbio(T) > 0 and CB(T) > 0*

$\frac{RFbio(T)}{CB(T)}$, otherwise } (Eq. 7)

ME is calculated across different time horizons (85, 185, and 285 years), providing a metric

that relates the remaining or effectively removed biogenic emissions from the atmosphere to

the total carbon balance. The higher the ME value, the greater the climate benefit delivered

per unit of carbon captured, making it a useful comparative indicator of mitigation strategy

performance.

5.1 Statistical Analysis of Mitigation Potential

The effects of forest management, climate scenarios, wood use, and substitution factors were

statistically evaluated across a range of carbon-related response variables in the forest sector.

Carbon ecosystem fluxes (i.e., GPP, R_{ECO}, and NEE) were analyzed with respect to forest

management and climate, while wood emissions, substitution effects, wood balance, and FSB

were evaluated in relation to all four factors.

We used linear mixed-effects models (LMEs) to regress GPP, RECO, NEE, and FSB against

five key drivers: Management, Wood Use (WU), Substitution (SUB), Climate, and Century.

These models were used to assess the influence of each factor and their interactions on

ecosystem fluxes and forest sector balances. For wood emissions, substitution effects, and

wood balance, we applied generalized additive mixed models (GAMMs) to account for

potential nonlinear relationships. Model selection was guided by residual diagnostics,

including assessments of linearity, normality, and homoscedasticity. The factors and their

acronyms, used consistently throughout the manuscript, are summarized in **Table 1**. Full

model specifications, transformations, and diagnostic results are provided in **S2**.

Table 1: Overview of the factorial design used in the simulation experiment, showing the five

key factors and their respective levels along with their corresponding modality acronyms.

Factor	Levels	Modality acronyms
Management	5	TM, WOOD, BIOE, ADAPT, TRANS
Wood use	4	BAU, RR20, LS20, RRLS20
Substitution	5	SUB0, SUB25, SUB50, SUB75, SUB100
Climate change scenario	3	CUR, SSP1-2.6, SSP5-8.5
Century	3	C1 (2015-2099), C2 (2100-2199), C3 (2200-2300), ALL (2015-2300)

5.2. Consistency between carbon and climate mitigation rankings and the concept of mitigation efficiency

To evaluate the relationship between CB and RF_{bio} outcomes, a linear regression analysis was conducted, modeling RF_{bio} as a function of CB across all studied scenarios. Model fit was assessed using the coefficient of determination (R^2) and statistical significance (p-value).

ME was calculated for each combination of climate scenario, management scheme, substitution pathway, and time horizon (H1: 2015–2100; H2: 2015–2200; H3: 2015–2300). To assess the sensitivity of ME to scenario context and temporal scope, we conducted a descriptive statistical comparison across each factor. Patterns in ME and CB were summarized using means, with scenario-specific anomalies (e.g., negative ME values) noted.

Additionally, an in-depth analysis of two forest mitigation options that exhibited the two most two divergent ME was conducted. This comparison focused on long-term FSB dynamics, RF_{bio} patterns, time in a net emissions regime, and disturbance-recovery cycles, to

527 elucidate the underlying trade-offs between cumulative carbon benefits and the timing and

528 distribution of emissions.

529 **Results**

530 **Forest sector balance (FSB)**

531 FSB, defined as the sum of NEE and Wbalance, varied significantly with climate scenarios,

532 management strategies, and across centuries. Climate effects on FSB were not consistent

533 across forest management types, and both drivers showed strong century-dependent patterns

534 (**see Table S2.4).** Although the DF had a low overall effect size ($\eta^2 = 0.03$, $p<0.001$), its

535 influence on FSB was substantial in specific scenarios, particularly those representing rapid

536 decarbonization pathways. Over the full simulation period, SSP5-8.5 yielded the lowest FSB

537 (0.07 ± 0.01 tC ha^{-1} year^{-1}), while SSP1-2.6 and CUR resulted in significantly higher

538 values (0.75 ± 0.01 and 0.64 ± 0.01, respectively). The ranking of scenarios changed

539 over time: SSP5-8.5 initially outperformed others but declined in the 22nd and 23rd

540 centuries, ultimately ranking lowest (**see Figure 6**). By contrast, SSP1-2.6 and CUR

541 remained more stable across centuries with lower inter-temporal variability.

542 Management strategies showed clear performance gradients mainly driven by NEE: TM and

543 BIOE produced the highest overall FSB (0.76 ± 0.02 and 0.59 ± 0.04), supported by

544 relatively strong sinks (NEE: -1.25 ± 0.05 and -1.10 ± 0.08). WOOD and ADAPT followed

545 with moderate FSB (0.39 ± 0.02 and 0.26 ± 0.02) with moderate sinks (NEE: $-0.78 \pm$

546 0.05 and -0.54 ± 0.05), while TRANS consistently showed the lowest values (0.05 ± 0.03),

547 declining to negative values under SSP5-8.5 (-0.47 ± 0.03) due to forest ecosystem turning

548 into carbon source (NEE: $+0.83 \pm 0.01$) (**see Figure 6**). Decreases were most severe in

549 TRANS and ADAPT, where net carbon source behavior emerged in the third century.

DF, which modulates substitution levels over time, had a major influence on Wbalance and

thus on FSB. Under SUB0, the substitution effect was maintained throughout all centuries,

while SUB25–SUB100 declined progressively, reaching zero by 2155 (SUB25), 2085

(SUB50), 2062 (SUB75), and 2050 (SUB100), respectively. These dynamics divide

pathways into three distinct temporal patterns: persistent (SUB0), medium-lived (SUB25),

and short-lived (SUB50–SUB100), which also significantly influence the Wbalance value.

Component analysis showed that NEE accounted for ~53–65% of FSB, particularly high

under TM. Substitution effect was lower in terms of contribution to the FSB but variable,

peaking under the SUB0 displacement pathway (18%), decreasing under SUB25 (7%), and

falling below 3% for SUB50–SUB100, which caused a (0.28 $\pm$ 0.07 tC ha^{-1} year^{-1};

53 $\pm$ 25%) reduction in FSB compared to SUB0 (**see Table S2.5**).

Figure 6: Mean FSB, in tC ha^{-1} year^{-1} for five forest management strategies (TM, ADAPT,

BIOE, TRANS, and WOOD), averaged across two climate scenarios (SSP1-2.6 and SSP5-

8.5), shown for the 21st, 22nd, and 23rd centuries, as well as for the full period combined.

Red error bars indicate the 95% confidence interval of FSB across climate scenarios. The

upper panel shows FSB values by substitution scenarios, while the lower panel shows FSB

values by climate scenario.

Radiative forcing & mitigation efficiency (RF_{bio} and ME)

Forest management scenarios with a more negative CB also tended to yield the greatest

climate benefits. A linear regression, performed using outcomes from all studied scenarios,

confirmed that CB is a strong predictor of RF_{bio} ($R^2 = 0.95$; $p < 2.2e{-}16$; see **Table S2.9**).

Thus, the results and analyses previously conducted on CB are also applicable to RF_{bio}.

However, this relationship is not always strictly linear. In some exceptional cases where CB

values are close, a mitigation option with a slightly less negative CB produced a more

favorable RF_{bio}. This was observed, for instance, under the SUB25 decarbonization pathway

at horizon H2: the TRANS scenario had a slightly higher CB (-128 ± 0.91 tC ha^{-1}) than

WOOD (-122 ± 2.75 tC ha^{-1}), yet WOOD achieved a more favorable RF_{bio} (-66 ± 1.33 vs. $-$

60 ± 0.13 tC ha^{-1}) and a higher ME (0.54 ± 0.00 vs. 0.47 ± 0.002). At horizon H3, this

contrast became even more pronounced: although TRANS reached a more negative CB ($-$

162 ± 0.92 tC/ha) compared to WOOD (-155 ± 2.05 tC ha^{-1}), it resulted in a substantially less

favorable RF_{bio} (-66 ± 0.70 vs. -73 ± 0.20 tC ha^{-1}) and a lower ME (0.41 ± 0.002 vs. $0.57 \pm$

0.002). ME showed strong sensitivity to both climate conditions and forest management

practices.

Under CUR and SSP1-2.6, ME remained relatively stable across management schemes,

typically ranging from 0.4 to 0.7 (**Figure 7** and **Table S2.10**). In contrast, under the SSP5-8.5

climate scenario, the ME of different management scenarios was more variable and often

with lower efficiencies, with the efficiency of several scenarios falling below zero,

particularly ADAPT and TRANS under H2 and H3 (**Figure 7** and **Table S2.10**), indicating

that these scenarios were contributing to climate warming rather than cooling it. The time

horizon (H1, H2, and H3) also influenced ME: across all climate scenarios, the short-term

horizon (H1) consistently resulted in higher efficiency values (>0.6) compared to the longer-

term time horizon (H2 and H3).

Interestingly, a more negative CB did not always equate to high efficiency. In CUR and

SSP1-2.6, forest projects with stronger (more negative) CB often showed lower or slightly

lower efficiency, whereas the opposite was true under SSP5-8.5, where stronger CB tended to

support higher ME. For example, ADAPT (0.53, SSP1-2.6, H2) and WOOD (0.54, SSP1-2.6,

H2) sometimes outperformed TM (0.49, SSP1-2.6, H2) or BIOE (0.45, SSP1-2.6, H2) in

efficiency despite having less negative CB values.

These dynamics were further supported and extended by generalized additive models

(GAMs; **Figures S2.1-S2.2**), which demonstrated that, in addition to CB, both the magnitude

and duration of emissions play dominant roles in determining ME. Under the SSP5-8.5

scenario, emissions magnitude and CB were strongly positively correlated (r = 0.85),

indicating that scenarios with more negative CB generally exhibited low emissions. In

contrast, a weak negative correlation (r = –0.28) was observed under CUR and SSP1-2.6,

suggesting no clear relationship between CB and emissions magnitude. In the ADAPT and

TRANS scenarios, under the SSP5-8.5 climate scenario, positive CB values resulted in

sustained net emissions and positive contributions to radiative forcing.

Figure 7: CB and ME under various forest management and climate scenarios. Circle size scales with CB via   ; ME is quantile-transformed, both for improved visibility and comparability across scenarios. Red borders indicate negative ME; hexagons show positive CB (carbon source sector). Subplots represent climate scenarios, with circles colored by management schemes.

In depth-insights from BIOE and WOOD strategies

Across the simulation period (2015–2300), the BIOE management scenario consistently achieved a more negative CB than WOOD. By 2300, BIOE reached –337 tC ha^{-1}, while WOOD reached –206 tC ha^{-1}. ME, however, remained higher in WOOD throughout. Both scenarios began with comparable ME in 2100 (0.62 for BIOE, 0.63 for WOOD). By 2200, WOOD retained a ME of 0.54, while BIOE declined to 0.46. In 2300, ME values were 0.47 for WOOD and 0.44 for BIOE (see **Figure 8**).

BIOE followed an 80-year shelterwood rotation, resulting in three major final harvests occurring in 2095, 2175, and 2265. WOOD used a 130-year rotation, with final cuts in 2145 and 2260. The first BIOE final harvest in 2095 led to emissions of approximately 17.3 tC ha^{-1}, with RF$_{bio}$ returning to pre-disturbance levels by 2135, indicating a 40-year

payback period. For WOOD, the first final cut in 2145 resulted in emissions of $5.6\,tC\,ha^{-1}$,

with RF_{bio} returning to net cooling within 15 years. Following these major harvests, NEE in

WOOD returned to negative values (net sink) within 4 years, whereas BIOE remained a net

carbon source (NEE > 0) for approximately 10 years.

In the three years immediately following each final cut, RF_{bio} increased sharply in BIOE,

while the increase in WOOD was more moderate. This early post-harvest rise in RF_{bio}

coincided with higher volumes of short-lived HWPs (i.e., firewood and paper) in BIOE;

$37.25\,m^3\,ha^{-1}$, $38.51\,m^3\,ha^{-1}$, and $37.88\,m^3\,ha^{-1}$ after its respective final harvests,

compared to $23.70\,m^3\,ha^{-1}$ and $14.82\,m^3\,ha^{-1}$ in WOOD.

Emission timing and magnitude differed between the two scenarios. In BIOE, emissions were

concentrated in a limited number of high-intensity events, associated with initial

establishment and final harvests. These events corresponded to periods where GPP was low

and R_{ECO} remained high although Rh_{onsite} decreased proportionally to GPP. Additional

immediate emissions from firewood combustion or from products at their end-life

incineration (i.e., paper) were also observed during these intervals. In WOOD, emissions

were more evenly distributed and of lower magnitude, arising from thinning, regeneration

phases, and smaller final cuts. RF_{bio} patterns in WOOD were more stable, with shorter

durations of net positive forcing following harvest events.

Figure 8: Temporal dynamics of RF_{bio}, CB, and ME under BIOE (panel a) and WOOD (panel b) management scenarios from 2015 to 2300. Shaded red bands indicate periods when the forest sector is a net source (FSB<0). Green horizontal bands denote the number of years required for RF_{bio} to return to its pre-harvest level.

Discussions

Our results suggest that effective forest-sector climate mitigation requires more than maximizing carbon balances. It demands a systems perspective integrating forest dynamics, wood use, and evolving socio-technical conditions. Over a multi-century assessment, we found that while the FSB is strongly influenced by climate and management, ME depends more on emission timing and scale. Substitution effects, particularly the contrast between SUB0 and SUB25-SUB100 proved substantial, while HWP lifespan and recycling rate had negligible long-term impact. Notably, scenarios with the lowest CB did not always deliver the greatest climate efficiency. ME declined over time, emphasizing the need to consider

temporal dynamics. All management options eventually showed declining carbon sink

capacity under SSP5-8.5, with TRANS, ADAPT, and WOOD becoming net sources.

Management strategies, wood use, and substitution pathways

Forest management has a strong influence on FSB. Intensive silvicultural practices, such as

TM and BIOE, consistently maintained high FSB across climate projections. Giasson et al.,

(2023) showed that harvest scenarios with high intervention and regular regrowth cycles

sustain stronger carbon sink than low-harvest or unmanaged systems. Furthermore, our

results showed that shifting from active to passive management under the TRANS scenario

significantly reduced FSB in the long term, especially under contrasted climate conditions

where forest carbon dynamics approached equilibrium. These findings align with Pukkala,

(2017), who reported that unmanaged forests exhibit a long-term decline, nearing zero around

200 years post-establishment. Beyond carbon storage, management also influences wood

volume and characteristics, affecting substitution potential.

Recent studies increasingly challenge static DF, advocating for dynamic approaches that

reflect evolving technological, industrial, and policy contexts (Boukhris, Collalti, et al., 2025;

Brunet-Navarro et al., 2021; Hurmekoski et al., 2023). Our dynamic modeling confirms these

concerns. In scenarios where DF reaches zero by 2050, substitution benefits disappear

entirely, resulting in a 53% lower FSB by 2300 compared to the SUB0 trajectory. This

mirrors projections by Brunet-Navarro et al. (2021), who estimated substitution benefits

could by $\cong$ 30% by 2030 and as much as by 96% by 2100, in line with the decarbonization

goals of the Paris Agreement.

Conversely, wood-use parameters like product lifespan and recycling rate had no discernable

long-term impact on FSB. Over 285 years, stored carbon in HWPs is eventually emitted

through decay, regardless of whether turnover occurs at 40 or 60 years. Although extended

lifespans and higher recycling rates may reduce emissions in the short- to medium-term, as

shown by (Brunet-Navarro et al., 2016), who projected up to 5 Mt CO_2 yr^{-1} savings in the EU-

28 with lifespan increases of 19.54% or a 20.92% recycling boost by 2030, these effects

diminish across multi-century timescales. Such results highlight the need to align

interventions with the analytical timeframe. Short-term gains may appear significant but are

absorbed over longer horizons when delayed emissions fall within the evaluation period.

Carbon sink instability under high-emission scenarios

The shift to net-carbon neutrality or a carbon source state by the last considered century under

SSP5-8.5 marks a critical loss of ecosystem resilience, driven by prolonged structural

degradation and physiological stress from high temperatures. Although precipitation increases

during this period, and VPD stays moderate (~0.825 ± 0.1 kPa), GPP declines sharply across

all strategies. This drop is more likely due to heat-induced impacts on photosynthetic

processes, such as enzyme deactivation, increased photorespiration, and reduced

carboxylation efficiency (Oliver et al., 2022; Qu Y et al., 2023). As biomass and metabolic

activity decline, both Ra and Rh fall due to limited substrate and structural loss, but GPP

declines faster, leading to sustained positive NEE in the final simulation century. This aligns

with Earth System Model outputs under SSP5-8.5 which show that warming suppresses

photosynthesis as temperatures exceed physiological thresholds, particularly in temperate and

boreal forests. The summer GPP-temperature correlation weakens over the 21st century,

turning significantly negative by 2070 (Chen & Peng, 2025; Zhang et al., 2022) also found

that 33% to 63% of global vegetated areas show declining productivity sensitivity to

temperature, with the most severe declines under SSP5□8.5.

Climate-Carbon dynamics and mitigation efficiency

The strong correlation observed between CB and RF_{bio} reflects the key role of long-lived CO_2 removals in determining climate impacts of forest-based mitigation. This aligns with carbon cycle theory and is consistent with IRFs such as the Bern model (Joos et al., 2013), where most climate benefit arises from sustained atmospheric CO_2 reduction (Archer et al., 2009; Masson-Delmotte et al., 2021). However, the CB-RF_{bio} relationship is not strictly proportional. Our results show that under the SUB25 decarbonization pathway at both H2 and H3, the WOOD scenario achieved a lower (better) RF_{bio} than TRANS, despite similar or less negative CB. This suggests CB alone is insufficient basis for assessing climate performance, and other factors also influence radiative forcing.

A key reason for this explanation for the discrepancy is the timing of carbon fluxes. CB summarizes total emissions and removals but masks their temporal distribution. Many scenarios involve non-monotonic dynamics, with initial emissions from establishment, thinning, or harvest followed by delayed regrowth. Early emissions exert greater radiative forcing due to the CO_2's long atmospheric lifetime cumulative warming (Allen et al., 2010; Matthews & Caldeira, 2008). Although IRF is linear for pulses, time-distributed fluxes interact non-linearly with the climate system. RF_{bio}, which captures these dynamics, more accurately reflects climate impact than CB alone (Sierra et al., 2021; Tanaka & O'Neill, 2018).

Nevertheless, neither CB nor RFbio capture the carbon "cost" of achieving mitigation. We therefore introduced ME, RFbio divided by CB. ME indicates how effectively carbon removal translates into avoided radiative forcing. For instance, BIOE had a more negative CB (−339 tC ha-1) than ADAPT (−162 tC ha-1), but a lower ME (0.44 vs. 0.49), meaning less efficient climate benefit. BIOE also required 17 silvicultural interventions and 3 final cuts

over 300 years, compared to 14 interventions and 2 final cuts in ADAPT. Thus, more

intensive management does not always yield proportionally greater mitigation and may entail

higher ecological or operational costs.

ME declined consistently as the time horizon extended. Initially, forests sequester carbon in

biomass and soils, but later interventions transfer carbon to HWPs, which eventually decay

and emit CO_2. Over time, emissions from earlier cycles accumulate, reducing net benefit.

This dynamic, "sequester now, emit later" (Guest et al., 2013), underscores the value of ME

as a complementary metric that integrates both climate effectiveness and implementation

sustainability.

Uncertainty, Model assumptions, and Long-Term Implications

Uncertainties within the modeling framework stem from initialization, process representation,

and parameterization ((Hamish) Kimmins et al., 2008; Reyer et al., 2020). The 3D-CMCC-

FEM forest growth model, while effective for stand-level analyses, simplifies spatial

heterogeneity and omits landscape-scale dynamics, potentially underrepresenting key

interactions such as seed dispersal and spatially explicit disturbance spread (Patacca et al.,

2023; Seidl et al., 2011). It also omits processes like tree establishment, disturbance-induced

mortality, and species composition are not represented, limiting its ability to capture

ecological feedback under climate change. Adaptive responses, including species

acclimation, migration, and phenological change, are excluded or overly simplified, despite

their growing relevance over multi-century timescales (Boukhris, Marano, et al., 2025;

Nicotra et al., 2010).

On the HWPs side, the TimberTracer model uses fixed bucking rules and constant product

allocations, without accounting for future technological shifts, circular economy

developments, or changing consumer demand (Harju, 2022; Olšiaková et al., 2016). This

restricts its ability to reflect how innovation or policy changes could influence product

lifespans, recycling rates, or end-of-life treatment.

Additional uncertainty comes from external drivers. Climate projections diverge significantly

after 2100 (Hausfather & Peters, 2020), and future land-use and forest management pathways

depend on socio-economic and policy trends that are inherently hard to predict (Popp et al.,

2017). DF estimates used to quantify substitution benefits are also highly uncertain and

context-specific (Leskinen et al., 2018; Pingoud et al., 2010). Their magnitude will likely

evolve as energy systems decarbonize and industry shifts to lower-carbon alternatives

(Gustavsson et al., 2017).

Conclusions

This study critically examined the long-term climate mitigation potential of the forest sector

by integrating forest growth, wood-use, and substitution dynamics within a unified modeling

framework. Our aim was not only to assess carbon balances but to evaluate whether

commonly used indicators accurately reflect true climate benefit over multi-century

timescales.

Using a coupled modeling approach applied on a *P. nigra* forest in Italy, we simulated a

broad range of futures across five forest management strategies, four wood-use patterns, and

five substitution decay scenarios under three contrasting climate pathways. While CB and

RF_{bio} were generally aligned, they captured different aspects of climate impact. CB reflects

net carbon flows, but not their atmospheric effect; RF_{bio} represents the warming or cooling

influence overtime. To bridge this gap, we introduced ME, a metric that relates carbon flows

to actual contribution to actual mitigation effectiveness.

Our findings show that scenarios with the highest CB (e.g., BIOE and TM) do not always

deliver the greatest climate benefit, especially when emissions are front-loaded or persistent.

Meanwhile, strategies like WOOD and ADAPT, which extended carbon residence times in

both forest and technosphere, achieved higher ME under SSP1-2.6. However, under the

SSP5-8.5 scenario, even these strategies became net radiative sources after 2200 due to

declining sink capacity.

Substitution benefits, often assumed constant, declined significantly under degressive

assumptions, reducing the overall forest sector mitigation by up to 53%, especially in high

harvest volumes. While substitution played a secondary role compared to climate and

management, its sensitivity to decarbonization pathways highlights the risk of overreliance

on this lever.

Taken together, these findings highlight that effective forest-sector mitigation requires more

than achieving high carbon balances. It also depends on minimizing the magnitude, timing,

and persistence of emissions, and on reevaluating the assumptions behind substitution effects.

Metrics like RF_{bio} and ME provide valuable tools to capture these temporal and systemic

dynamics, offering a more nuanced, climate-relevant basis for evaluating forest-sector

strategies.

Ultimately, meaningful forest-sector contributions to climate neutrality require assessment

not just of carbon flows, but of their actual atmospheric impact over time.

Appendix

Figure A.1: Integrated modeling framework used to evaluate the long-term climate mitigation

potential of a managed *Pinus nigra* forest (2015-2300). The framework combines forest

dynamics (3D-CMCC-FEM), wood lifecycle modeling (TimberTracer), and an impulse

response function (Bern 2.5 CC IRF). Simulations cover multiple scenario combinations (for

management, climate, wood-use, and substitution). Outputs include temporal net ecosystem

exchange, substitution effect, and emissions from wood combustion and decay, which are

used to compute forest sector balance and radiative forcing from biogenic CO_2 emissions.

References

Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M.,
Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H. (Ted), Gonzalez, P., Fensham, R.,
Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A., &
Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals
emerging climate change risks for forests. *Forest Ecology and Management*, *259*(4), 660–
684. https://doi.org/10.1016/j.foreco.2009.09.001
820

821 Ameray, A., Bergeron, Y., Valeria, O., Montoro Girona, M., & Cavard, X. (2021). Forest
822 Carbon Management: A Review of Silvicultural Practices and Management Strategies Across
823 Boreal, Temperate and Tropical Forests. *Current Forestry Reports*, *7*(4), 245–266.
824 https://doi.org/10.1007/s40725-021-00151-w
825

826 Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K.,
827 Matsumoto, K., Munhoven, G., Montenegro, A., & Tokos, K. (2009). Atmospheric Lifetime
828 of Fossil Fuel Carbon Dioxide. *Annual Review of Earth and Planetary Sciences*, *37*(Volume
829 37, 2009), 117–134. https://doi.org/10.1146/annurev.earth.031208.100206
830

831 Barredo, J. I., Vizzarri, M., & Kuželová, K. (2024). Archetypal typology of European forest
832 ecosystems integrating management intensity and naturalness. *Ambio*, *53*(11), 1587–1598.
833 https://doi.org/10.1007/s13280-024-02050-3
834

835 Barrette, J., Achim, A., & Auty, D. (2023). Impact of Intensive Forest Management Practices
836 on Wood Quality from Conifers: Literature Review and Reflection on Future Challenges.
837 *Current Forestry Reports*, *9*(2), 101–130. https://doi.org/10.1007/s40725-023-00181-6
838

839 Bauhus, J., Puettmann, K., & Messier, C. (2009). Silviculture for old-growth attributes.
840 *Forest Ecology and Management*, *258*(4), 525–537.
841 https://doi.org/10.1016/j.foreco.2009.01.053
842

843 Bottalico, F., Chirici, G., & Travaglini, D. (2012). La gestione della foresta di Vallombrosa
844 dal 1876 al 2006: Analisi delle cartografie storiche. *L'Italia Forestale e Montana*, 449–458.
845 https://doi.org/10.4129/ifm.2012.6.01
846

Boukhris, I., Collalti, A., Lahssini, S., Dalmonech, D., Nakhle, F., Testolin, R., Chiriacò, M. V., Santini, M., & Valentini, R. (2025). TimberTracer: A comprehensive framework for the evaluation of carbon sequestration by forest management and substitution of harvested wood products. *Carbon Balance and Management*, *20*(1), 12. https://doi.org/10.1186/s13021-025-00296-2

Boukhris, I., Marano, G., Dalmonech, D., Valentini, R., & Collalti, A. (2025). Modeling Forest Growth Under Current and Future Climate. *Current Forestry Reports*, *11*(1), 17. https://doi.org/10.1007/s40725-025-00249-5

Bozzolan, N., Grassi, G., Mohren, F., & Nabuurs, G.-J. (2024). Options to improve the carbon balance of the harvested wood products sector in four EU countries. *GCB Bioenergy*, *16*(1), e13104. https://doi.org/10.1111/gcbb.13104

Brunet-Navarro, P., Jochheim, H., Cardellini, G., Richter, K., & Muys, B. (2021). Climate mitigation by energy and material substitution of wood products has an expiry date. *Journal of Cleaner Production*, *303*, 127026. https://doi.org/10.1016/j.jclepro.2021.127026

Brunet-Navarro, P., Jochheim, H., & Muys, B. (2016). Modelling carbon stocks and fluxes in the wood product sector: A comparative review. *Global Change Biology*, *22*(7), 2555–2569. https://doi.org/10.1111/gcb.13235

Brunet-Navarro, P., Jochheim, H., & Muys, B. (2017). The effect of increasing lifespan and recycling rate on carbon storage in wood products from theoretical model to application for the European wood sector. *Mitigation and Adaptation Strategies for Global Change*, *22*(8), 1193–1205. https://doi.org/10.1007/s11027-016-9722-z

Budzinski, M., Bezama, A., & Thrän, D. (2020). Estimating the potentials for reducing the impacts on climate change by increasing the cascade use and extending the lifetime of wood products in Germany. *Resources, Conservation & Recycling: X*, *6*, 100034. https://doi.org/10.1016/j.rcrx.2020.100034

Calvin, K., Bond-Lamberty, B., Clarke, L., Edmonds, J., Eom, J., Hartin, C., Kim, S., Kyle, P., Link, R., Moss, R., McJeon, H., Patel, P., Smith, S., Waldhoff, S., & Wise, M. (2017). The SSP4: A world of deepening inequality. *Global Environmental Change*, *42*, 284–296. https://doi.org/10.1016/j.gloenvcha.2016.06.010

Cantiani P, Di Salvatore U, & Romano R. (2018). La selvicoltura delle pinete artificiali di pino nero: Analisi delle legislazioni regionali italiane. *Forest@ - Journal of Silviculture and Forest Ecology*, *16*(1), 99. https://doi.org/10.3832/efor2985-015

Chen, T., & Peng, L. (2025). Global GPP Changes and Its Sensitivity to Temperature. In T. Chen & L. Peng (Eds.), *Multi-scale Understanding of Vegetation Ecosystems: Changes, Drivers and Inspirations* (pp. 145–169). Springer Nature Switzerland. https://doi.org/10.1007/978-3-031-90163-8_8

Cherubini, F., Peters, G. P., Berntsen, T., Strømman, A. H., & Hertwich, E. (2011). CO2
emissions from biomass combustion for bioenergy: Atmospheric decay and contribution to
global warming. *GCB Bioenergy*, *3*(5), 413–426. https://doi.org/10.1111/j.1757-
1707.2011.01102.x

Ciancio, O., & Nocentini, S. (2004). *Il bosco ceduo. selvicoltura, assestamento, gestione.* ita.
https://flore.unifi.it/handle/2158/25946

Ciancio, O., & Nocentini, S. (2011). Biodiversity conservation and systemic silviculture:
Concepts and applications. *Plant Biosystems - An International Journal Dealing with All*
Aspects of Plant Biology, *145*(2), 411–418. https://doi.org/10.1080/11263504.2011.558705

Collalti, A., Perugini, L., Santini, M., Chiti, T., Nolè, A., Matteucci, G., & Valentini, R.
(2014). A process-based model to simulate growth in forests with complex structure:
Evaluation and use of 3D-CMCC Forest Ecosystem Model in a deciduous forest in Central
Italy. *Ecological Modelling*, *272*, 362–378. https://doi.org/10.1016/j.ecolmodel.2013.09.016

Dalmonech, D., Marano, G., Amthor, J. S., Cescatti, A., Lindner, M., Trotta, C., & Collalti,
A. (2022). Feasibility of enhancing carbon sequestration and stock capacity in temperate and
boreal European forests via changes to management regimes. *Agricultural and Forest*
Meteorology, *327*, 109203. https://doi.org/10.1016/j.agrformet.2022.109203

Dalmonech, D., Vangi, E., Chiesi, M., Chirici, G., Fibbi, L., Giannetti, F., Marano, G.,
Massari, C., Nolè, A., Xiao, J., & Collalti, A. (2024). Regional estimates of gross primary
production applying the Process-Based Model 3D-CMCC-FEM vs. Remote-Sensing multiple
datasets. *European Journal of Remote Sensing*, *57*(1), 2301657.
https://doi.org/10.1080/22797254.2023.2301657

Dye, A. W., Houtman, R. M., Gao, P., Anderegg, W. R. L., Fettig, C. J., Hicke, J. A., Kim, J.
B., Still, C. J., Young, K., & Riley, K. L. (2024). Carbon, climate, and natural disturbance: A
review of mechanisms, challenges, and tools for understanding forest carbon stability in an
uncertain future. *Carbon Balance and Management*, *19*(1), 35.
https://doi.org/10.1186/s13021-024-00282-0

EEA. (2022). *Annual European Union greenhouse gas inventory 1990-2020 and inventory*
report 2022. https://www.eea.europa.eu/en/analysis/publications/annual-european-union-
greenhouse-gas-1

EEA. (2024). *Diversion of waste from landfill in Europe*.
https://www.eea.europa.eu/en/analysis/indicators/diversion-of-waste-from-landfill

EU council. (2025). *Improved EU rules for forest reproductive material: Council approves*
negotiating position. Consilium. https://www.consilium.europa.eu/en/press/press-

releases/2025/06/13/improved-eu-rules-for-forest-reproductive-material-council-approves-negotiating-position/

European Commision. (2018). *Bioeconomy Strategy*. https://environment.ec.europa.eu/strategy/bioeconomy-strategy_en

European Commision. (2021). *New EU forest strategy for 2030*. https://environment.ec.europa.eu/strategy/forest-strategy_en

European Commission. (2021). *Biodiversity strategy for 2030*. https://environment.ec.europa.eu/strategy/biodiversity-strategy-2030_en

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, *9*(5), 1937–1958. https://doi.org/10.5194/gmd-9-1937-2016

Giasson, L.-A., Thiffault, E., Lebel, L., & Carle, J.-F. (2023). Carbon balance of forest management and wood production in the boreal forest of Quebec (Canada). *Frontiers in Forests and Global Change*, *6*. https://doi.org/10.3389/ffgc.2023.1242218

Grassi, G., Fiorese, G., Pilli, R., Jonsson, K., Blujdea, V., Korosuo, A., & Vizzarri, M. (2021, May 6). *Brief on the role of the forest-based bioeconomy in mitigating climate change through carbon storage and material substitution*. JRC Publications Repository. https://publications.jrc.ec.europa.eu/repository/handle/JRC124374

Gregor, K., Krause, A., Reyer, C. P. O., Knoke, T., Meyer, B. F., Suvanto, S., & Rammig, A. (2024). Quantifying the impact of key factors on the carbon mitigation potential of managed temperate forests. *Carbon Balance and Management*, *19*(1), 10. https://doi.org/10.1186/s13021-023-00247-9

Guest, G., Cherubini, F., & Strømman, A. H. (2013). The role of forest residues in the accounting for the global warming potential of bioenergy. *GCB Bioenergy*, *5*(4), 459–466. https://doi.org/10.1111/gcbb.12014

Gustavsson, L., Haus, S., Lundblad, M., Lundström, A., Ortiz, C. A., Sathre, R., Truong, N. L., & Wikberg, P.-E. (2017). Climate change effects of forestry and substitution of carbon-intensive materials and fossil fuels. *Renewable and Sustainable Energy Reviews*, *67*, 612–624. https://doi.org/10.1016/j.rser.2016.09.056

(Hamish) Kimmins, J. P., Blanco, J. A., Seely, B., Welham, C., & Scoullar, K. (2008). Complexity in modelling forest ecosystems: How much is enough? *Forest Ecology and Management*, *256*(10), 1646–1658. https://doi.org/10.1016/j.foreco.2008.03.011

Harju, C. (2022). The perceived quality of wooden building materials—A systematic
literature review and future research agenda. *International Journal of Consumer Studies*,
46(1), 29–55. https://doi.org/10.1111/ijcs.12764

Harmon, M. E. (2019). Have product substitution carbon benefits been overestimated? A
sensitivity analysis of key assumptions. *Environmental Research Letters*, *14*(6), 065008.
https://doi.org/10.1088/1748-9326/ab1e95

Hausfather, Z., & Peters, G. P. (2020). Emissions – the 'business as usual' story is
misleading. *Nature*, *577*(7792), 618–620. https://doi.org/10.1038/d41586-020-00177-3

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas,
J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X.,
Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., … Thépaut, J.-N. (2020). The
ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, *146*(730),
1999–2049. https://doi.org/10.1002/qj.3803

Holtsmark, B. (2012). Harvesting in boreal forests and the biofuel carbon debt. *Climatic*
Change, *112*(2), 415–428. https://doi.org/10.1007/s10584-011-0222-6

Holtsmark, B. (2013). The outcome is in the assumptions: Analyzing the effects on
atmospheric CO2 levels of increased use of bioenergy from forest biomass. *GCB Bioenergy*,
5(4), 467–473. https://doi.org/10.1111/gcbb.12015

1000

1001 Howard, C., Dymond, C. C., Griess, V. C., Tolkien-Spurr, D., & van Kooten, G. C. (2021).
1002 Wood product carbon substitution benefits: A critical review of assumptions. *Carbon*
1003 *Balance and Management*, *16*(1), 9. https://doi.org/10.1186/s13021-021-00171-w

1004

1005 Hurmekoski, E., Kunttu, J., Heinonen, T., Pukkala, T., & Peltola, H. (2023). Does expanding
1006 wood use in construction and textile markets contribute to climate change mitigation?
1007 *Renewable   and   Sustainable   Energy   Reviews*,   *174*,   113152.
1008 https://doi.org/10.1016/j.rser.2023.113152

1009

1010 Jonsson, R., Rinaldi, F., Pilli, R., Fiorese, G., Hurmekoski, E., Cazzaniga, N., Robert, N., &
1011 Camia, A. (2021). Boosting the EU forest-based bioeconomy: Market, climate, and
1012 employment impacts. *Technological Forecasting and Social Change*, *163*, 120478.
1013 https://doi.org/10.1016/j.techfore.2020.120478

1014

Joos, F., Roth, R., Fuglestvedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V.,
Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden,
P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., … Weaver,
A. J. (2013). Carbon dioxide and climate impulse response functions for the computation of
greenhouse gas metrics: A multi-model analysis. *Atmospheric Chemistry and Physics*, *13*(5),
2793–2825. https://doi.org/10.5194/acp-13-2793-2013

1021

Kilpeläinen, A., & Peltola, H. (2022). Carbon Sequestration and Storage in European Forests.
In L. Hetemäki, J. Kangas, & H. Peltola (Eds.), *Forest Bioeconomy and Climate Change* (pp.
113–128). Springer International Publishing. https://doi.org/10.1007/978-3-030-99206-4_6

Krug, J., Koehl, M., & Kownatzki, D. (2012). Revaluing unmanaged forests for climate
change mitigation. *Carbon Balance and Management*, 7(1), 11. https://doi.org/10.1186/1750-
0680-7-11

Laganière, J., Paré, D., Thiffault, E., & Bernier, P. Y. (2017). Range and uncertainties in
estimating delays in greenhouse gas mitigation potential of forest bioenergy sourced from
Canadian forests. *GCB Bioenergy*, 9(2), 358–369. https://doi.org/10.1111/gcbb.12327

1031

1032 Lemprière, T. C., Kurz, W. A., Hogg, E. H., Schmoll, C., Rampley, G. J., Yemshanov, D.,
1033 McKenney, D. W., Gilsenan, R., Beatch, A., Blain, D., Bhatti, J. S., & Krcmar, E. (2013).
1034 Canadian boreal forests and climate change mitigation. *Environmental Reviews*, 21(4), 293–
1035 321. https://doi.org/10.1139/er-2013-0039

1036

1037 Leskinen, P., Cardellini, G., González-García, S., Hurmekoski, E., Sathre, R., Seppälä, J.,
1038 Smyth, C., Stern, T., Verkerk, P. J., & European Forest Institute. (2018). *Substitution effects*
1039 *of wood-based products in climate change mitigation* (From Science to Policy) [From
1040 Science to Policy]. European Forest Institute. https://doi.org/10.36333/fs07

1041

1042 Levasseur, A., Lesage, P., Margni, M., Deschênes, L., & Samson, R. (2010). Considering
1043 Time in LCA: Dynamic LCA and Its Application to Global Warming Impact Assessments.
1044 *Environmental Science & Technology*, 44(8), 3169–3174. https://doi.org/10.1021/es9030003

1045

1046 Liu, P., Zha, T., Black, T. A., Jassal, R. S., Jia, X., Noormets, A., Ouimette, A., Tian, Y., &
1047 Li, X. (2025). Seasonal warming responses of the carbon dioxide sink from northern forests
1048 are sensitive to stand age. *Communications Earth & Environment*, 6(1), 43.
1049 https://doi.org/10.1038/s43247-025-02008-7

1050

Luyssaert, S., Inglima, I., Jung, M., Richardson, A. D., Reichstein, M., Papale, D., Piao, S.
L., Schulze, E.-D., Wingate, L., Matteucci, G., Aragao, L., Aubinet, M., Beer, C., Bernhofer,
C., Black, K. G., Bonal, D., Bonnefond, J.-M., Chambers, J., Ciais, P., … Janssens, I. A.
(2007). CO2 balance of boreal, temperate, and tropical forests derived from a global
database. *Global Change Biology*, 13(12), 2509–2537. https://doi.org/10.1111/j.1365-
2486.2007.01439.x

1057

Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N.,
Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B.
R., Maycock, T. K., Waterfield, T., Yelekçi, Ö., Yu, R., & Zhou, B. (Eds.). (2021). *Climate*
Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth
Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
University Press. https://doi.org/10.1017/9781009157896

1064

1065 Matthews, H. D., & Caldeira, K. (2008). Stabilizing climate requires near-zero emissions.
1066 *Geophysical Research Letters*, *35*(4). https://doi.org/10.1029/2007GL032388
1067

Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle,
U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P.
B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., … Wang, R.
H. J. (2020). The shared socio-economic pathway (SSP) greenhouse gas concentrations and
their extensions to 2500. *Geoscientific Model Development*, *13*(8), 3571–3605.
https://doi.org/10.5194/gmd-13-3571-2020
1074

1075 Meinshausen, M., Raper, S. C. B., & Wigley, T. M. L. (2011). Emulating coupled
1076 atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 – Part 1:
1077 Model description and calibration. *Atmospheric Chemistry and Physics*, *11*(4), 1417–1456.
1078 https://doi.org/10.5194/acp-11-1417-2011
1079

1080 Metzler, H., Launiainen, S., & Vico, G. (2024). Amount of carbon fixed, transit time and fate
1081 of harvested wood products define the climate change mitigation potential of boreal forest
1082 management—A model analysis. *Ecological Modelling*, *491*, 110694.
1083 https://doi.org/10.1016/j.ecolmodel.2024.110694
1084

1085 Moreau, G., Chagnon, C., Achim, A., Caspersen, J., D'Orangeville, L., Sánchez-Pinillos, M.,
1086 & Thiffault, N. (2022). Opportunities and limitations of thinning to increase resistance and
1087 resilience of trees and forests to global change. *Forestry: An International Journal of Forest
1088 Research*, *95*(5), 595–615. https://doi.org/10.1093/forestry/cpac010
1089

1090 Nabuurs, G.-J., Delacote, P., Ellison, D., Hanewinkel, M., Hetemäki, L., & Lindner, M.
1091 (2017). By 2050 the Mitigation Effects of EU Forests Could Nearly Double through Climate
1092 Smart Forestry. *Forests*, *8*(12), Article 12. https://doi.org/10.3390/f8120484
1093

1094 Nagel, R., Meyer, P., Blaschke, M., & Feldmann, E. (2023). Strict forest protection: A
1095 meaningful contribution to Climate-Smart Forestry? An evaluation of temporal trends in the
1096 carbon balance of unmanaged forests in Germany. *Frontiers in Forests and Global Change*,
1097 *6*. https://doi.org/10.3389/ffgc.2023.1099558
1098

1099 Nicotra, A. B., Atkin, O. K., Bonser, S. P., Davidson, A. M., Finnegan, E. J., Mathesius, U.,
1100 Poot, P., Purugganan, M. D., Richards, C. L., Valladares, F., & van Kleunen, M. (2010).
1101 Plant phenotypic plasticity in a changing climate. *Trends in Plant Science*, *15*(12), 684–692.
1102 https://doi.org/10.1016/j.tplants.2010.09.008
1103

1104 Noormets, A., Epron, D., Domec, J. C., McNulty, S. G., Fox, T., Sun, G., & King, J. S.
1105 (2015). Effects of forest management on productivity and carbon sequestration: A review and
1106 hypothesis. *Forest Ecology and Management*, *355*, 124–140.
1107 https://doi.org/10.1016/j.foreco.2015.05.019
1108

1109   Oliver, R. J., Mercado, L. M., Clark, D. B., Huntingford, C., Taylor, C. M., Vidale, P. L.,
1110   McGuire, P. C., Todt, M., Folwell, S., Shamsudheen Semeena, V., & Medlyn, B. E. (2022).
1111   Improved representation of plant physiology in the JULES-vn5.6 land surface model:
1112   Photosynthesis, stomatal conductance and thermal acclimation. *Geoscientific Model*
1113   *Development*, *15*(14), 5567–5592. https://doi.org/10.5194/gmd-15-5567-2022

1114

1115   Olšiaková, M., Loučanová, E., & Paluš, H. (2016). Monitoring changes in consumer
1116   requirements for wood products in terms of consumer behavior. *Acta Facultatis Xylologiae*
1117   *Zvolen*, *58(1)*. https://doi.org/10.17423/afx.2016.58.1.15

1118

1119   O'Neill, B. C., Kriegler, E., Riahi, K., Ebi, K. L., Hallegatte, S., Carter, T. R., Mathur, R., &
1120   van Vuuren, D. P. (2014). A new scenario framework for climate change research: The
1121   concept of shared socioeconomic pathways. *Climatic Change*, *122*(3), 387–400.
1122   https://doi.org/10.1007/s10584-013-0905-2

1123

Patacca, M., Lindner, M., Lucas-Borja, M. E., Cordonnier, T., Fidej, G., Gardiner, B., Hauf,
Y., Jasinevičius, G., Labonne, S., Linkevičius, E., Mahnken, M., Milanovic, S., Nabuurs, G.-
J., Nagel, T. A., Nikinmaa, L., Panyatov, M., Bercak, R., Seidl, R., Ostrogović Sever, M. Z.,
… Schelhaas, M.-J. (2023). Significant increase in natural disturbance impacts on European
forests since 1950. *Global Change Biology*, *29*(5), 1359–1376.
https://doi.org/10.1111/gcb.16531

1130

1131   Pilli, R., Alkama, R., Cescatti, A., Kurz, W. A., & Grassi, G. (2022). The European forest
1132   carbon budget under future climate conditions and current management practices.
1133   *Biogeosciences*, *19*(13), 3263–3284. https://doi.org/10.5194/bg-19-3263-2022

1134

1135   Pingoud, K., Perälä, A.-L., Soimakallio, S., & Pussinen, A. (2003). *Greenhouse gas impacts*
1136   *of harvested wood products: Evaluation and development of methods* (RESEARCH NOTES).
1137   VTT TIEDOTTEITA.

1138

1139   Pingoud, K., Pohjola, J., & Valsta, L. (2010). Assessing the integrated climatic impacts of
1140   forestry and wood products. *Silva Fennica*, *44*(1). https://www.silvafennica.fi/article/166

1141

1142   Popp, A., Calvin, K., Fujimori, S., Havlik, P., Humpenöder, F., Stehfest, E., Bodirsky, B. L.,
1143   Dietrich, J. P., Doelmann, J. C., Gusti, M., Hasegawa, T., Kyle, P., Obersteiner, M., Tabeau,
1144   A., Takahashi, K., Valin, H., Waldhoff, S., Weindl, I., Wise, M., … Vuuren, D. P. van.
1145   (2017). Land-use futures in the shared socio-economic pathways. *Global Environmental*
1146   *Change*, *42*, 331–345. https://doi.org/10.1016/j.gloenvcha.2016.10.002

1147

1148   Pukkala, T. (2017). Does management improve the carbon balance of forestry? *Forestry: An*
1149   *International Journal of Forest Research*, *90*(1), 125–135.
1150   https://doi.org/10.1093/forestry/cpw043

1151

Qu Y, Mueller-Cajar O, & Yamori W. (2023). Improving plant heat tolerance through
modification of Rubisco activase in C3 plants to secure crop yield and food security in a
future warming world. *Journal of Experimental Botany*, *74*(2).
https://doi.org/10.1093/jxb/erac340

Reyer, C. P. O., Silveyra Gonzalez, R., Dolos, K., Hartig, F., Hauf, Y., Noack, M., Lasch-
Born, P., Rötzer, T., Pretzsch, H., Meesenburg, H., Fleck, S., Wagner, M., Bolte, A., Sanders,
T. G. M., Kolari, P., Mäkelä, A., Vesala, T., Mammarella, I., Pumpanen, J., … Frieler, K.
(2020). The PROFOUND Database for evaluating vegetation models and simulating climate
impacts on European forests. *Earth System Science Data*, *12*(2), 1295–1320.
https://doi.org/10.5194/essd-12-1295-2020

Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer,
N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S.,
Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., … Tavoni, M. (2017). The Shared
Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions
implications: An overview. *Global Environmental Change*, *42*, 153–168.
https://doi.org/10.1016/j.gloenvcha.2016.05.009

Rogelj, J., Popp, A., Calvin, K. V., Luderer, G., Emmerling, J., Gernaat, D., Fujimori, S.,
Strefler, J., Hasegawa, T., Marangoni, G., Krey, V., Kriegler, E., Riahi, K., van Vuuren, D.
P., Doelman, J., Drouet, L., Edmonds, J., Fricko, O., Harmsen, M., … Tavoni, M. (2018).
Scenarios towards limiting global mean temperature increase below 1.5 °C. *Nature Climate*
Change, *8*(4), 325–332. https://doi.org/10.1038/s41558-018-0091-3

Rüter, S., & Fulvio, F. D. (2025). *ForestNavigator: Navigating European forests and forest*
bioeconomy sustainably to EU climate neutrality.
https://www.openagrar.de/servlets/MCRFileNodeServlet/openagrar_derivate_00065387/dn06
9634.pdf

Rüter S., Werner F., Forsell, N., Prins C., Vial, E., & Levet A.L. (2016). *ClimWood2030—*
Climate benefitsof material substitution by forest biomass and harvested wood products:
Perspective 2030 (Thünen Report 42; p. 148). Thünen Institute.
https://op.europa.eu/en/publication-detail/-/publication/993ecb81-2e54-11e8-b5fe-
01aa75ed71a1/language-en.

Schulze, E. D., Sierra, C. A., Egenolf, V., Woerdehoff, R., Irslinger, R., Baldamus, C.,
Stupak, I., & Spellmann, H. (2020). The climate change mitigation effect of bioenergy from
sustainably managed forests in Central Europe. *GCB Bioenergy*, *12*(3), 186–197.
https://doi.org/10.1111/gcbb.12672

Schulze, E. D., Valentini, R., & Bouriaud, O. (2021). The role of net ecosystem productivity
and of inventories in climate change research: The need for "net ecosystem productivity with
harvest", NEPH. *Forest Ecosystems*, *8*(1), 15. https://doi.org/10.1186/s40663-021-00294-z

Seidl, R., Fernandes, P. M., Fonseca, T. F., Gillet, F., Jönsson, A. M., Merganičová, K., Netherer, S., Arpaci, A., Bontemps, J.-D., Bugmann, H., González-Olabarria, J. R., Lasch, P., Meredieu, C., Moreira, F., Schelhaas, M.-J., & Mohren, F. (2011). Modelling natural disturbances in forest ecosystems: A review. *Ecological Modelling*, *222*(4), 903–924. https://doi.org/10.1016/j.ecolmodel.2010.09.040

Sierra, C. A., Crow, S. E., Heimann, M., Metzler, H., & Schulze, E.-D. (2021). The climate benefit of carbon sequestration. *Biogeosciences*, *18*(3), 1029–1048. https://doi.org/10.5194/bg-18-1029-2021

Skytt, T., Englund, G., & Jonsson, B.-G. (2021). Climate mitigation forestry—Temporal trade-offs. *Environmental Research Letters*, *16*(11), 114037. https://doi.org/10.1088/1748-9326/ac30fa

Strassmann, K. M., & Joos, F. (2018). The Bern Simple Climate Model (BernSCM) v1.0: An extensible and fully documented open-source re-implementation of the Bern reduced-form model for global carbon cycle–climate simulations. *Geoscientific Model Development*, *11*(5), 1887–1908. https://doi.org/10.5194/gmd-11-1887-2018

Tanaka, K., & O'Neill, B. C. (2018). The Paris Agreement zero-emissions goal is not always consistent with the 1.5 °C and 2 °C temperature targets. *Nature Climate Change*, *8*(4), 319–324. https://doi.org/10.1038/s41558-018-0097-x

Travaglini, D. (2009). *Piano di Gestione e Silvomuseo 2006-2025 della Foresta di Vallombrosa: Compartimentazione e cartografia*. https://flore.unifi.it/handle/2158/393724

Vacek, Z., Cukor, J., Vacek, S., Gallo, J., Bažant, V., & Zeidler, A. (2023). Role of black pine (Pinus nigra J. F. Arnold) in European forests modified by climate change. *European Journal of Forest Research*, *142*(6), 1239–1258. https://doi.org/10.1007/s10342-023-01605-5

Valentini, R., & Miglietta, F. (2015). *The greenhouse gas balance of Italy. An insight on managed and natural terrestrial ecosystems*. https://doi.org/10.1007/978-3-642-32424-6

van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., & Rose, S. K. (2011). The representative concentration pathways: An overview. *Climatic Change*, *109*(1), 5. https://doi.org/10.1007/s10584-011-0148-z

Wei, X., Zhao, J., Hayes, D. J., Daigneault, A., & Zhu, H. (2023). A life cycle and product type based estimator for quantifying the carbon stored in wood products. *Carbon Balance and Management*, *18*(1), 1. https://doi.org/10.1186/s13021-022-00220-y

1239 Zhang, Y., Piao, S., Sun, Y., Rogers, B. M., Li, X., Lian, X., Liu, Z., Chen, A., & Peñuelas, J.
1240 (2022). Future reversal of warming-enhanced vegetation productivity in the Northern
1241 Hemisphere. *Nature Climate Change*, *12*(6), 581–586. https://doi.org/10.1038/s41558-022-
1242 01374-w

1243

1244

Credit authorship contribution statement

Issam Boukhris: Writing - original draft**,** Writing - review & editing, Methodology, Visualization, Validation, Investigation, Data curation, Conceptualization, Project administration. **Francesco Cherubini:** Writing - review & editing, Methodology, Resources, Formal analysis, Resources. **Alessio Collalti:** Writing - review & editing, Validation, Software, Formal analysis, Resources. **Daniela Dalmonech:** Validation, Software, Resources. **Christian Vonderach:** Writing - review & editing, Investigation, Validation, Formal analysis. **Francesca Giannetti**: Writing - review & editing, Resources, Formal analysis. **Gina Marano:** Writing - review & editing, Methodology, Formal analysis **Said Lahssini:** Writing - review & editing, Formal analysis, Software. **Monia Santini:** Writing - review & editing, Software, Funding acquisition. **Riccardo Valentini:** Formal analysis, Supervision, Funding acquisition.

Funding sources

Issam Boukhris was supported by the National Research Center for Agricultural Technologies —AGRITECH— PNRR (Italian National Plan of Recovery and Resilience), CN00000022 in WP 4.3 Task 3 Risk management strategies and policies in the context of climate change to carry out multiple field missions, during which he collected essential data in the Vallombrosa Forest in Tuscany, Italy. Alessio Collalti and Daniela Dalmonech acknowledge funding by the European Union – NextGenerationEU under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of December 16, 2021, rectified by Decree n.3175 of December 18, 2021 of the Italian Ministry of University and Research under award Number: Project code CN_00000033, Concession Decree No. 1034 of June 17, 2022 adopted by the Italian Ministry of University and Research, CUP B83C22002930006, Project title "National Biodiversity Future Centre - NBFC".

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability Statement

Data are available from the corresponding author upon reasonable request.

Acknowledgment

Issam Boukhris would like to thank the University of Tuscia and the CMCC Foundation, both based in Italy, for supporting his Ph.D. thesis, of which this work is a part. He also extends his gratitude to Professor Gherardo Chirici and Dr. Elia Vangi from the University of Florence, Vincenzo Saponaro from the CNR of Italy, Colonel Giuliano Savelli and Maréchal Giovanni Galipò from the local forest services of Vallombrosa for their technical and logistical support. Special thanks are also due to Dr. Khalil Misbah from the IRD of Toulouse, as well as to Adnan Yousaf and Renato Zompanti from the University of Tuscia, for their invaluable assistance during fieldwork.