Distinguishing cortical indexes of arousal and awareness in sleeping patients with unresponsive wakefulness syndrome

Man Li[a,b], Jiahui Pan[b,c], Junrong Han[d], Xiaoyu Bao[a,b], Hang Wu[d], Fei Wang[b,c], Yangzuyi Yu[c], Aoqi Li[c], Yuhao Liu[c], Di Chen[a,b], Xiaochun Yang[e], Yanbin He[e,*], Pengmin Qin[b,d,*], and Yuanqing Li[a,b,f,*]

Consciousness is assumed to be defined by two components: awareness and arousal. However, the cortical indexes for these two components remain unclear. Since patients with unresponsive wakefulness syndrome (UWS) are supposed to have no awareness but arousal, sleep-wake cycle in patients with UWS may reflect pure arousal fluctuations, and then could distinguish the cortical indexes for arousal from ones for awareness. This study recorded nighttime polysomnography in patients with UWS, patients with minimally conscious state (MCS) and healthy controls, showing that spectral slope could index arousal, fluctuating among sleep stages in all three groups. Both spectral entropy and Lempel-Ziv complexity can index awareness, showing a significant difference between conscious and unconscious states. These findings provide fundamental evidence for the two-component hypothesis of consciousness.

N3-sleep | REM-sleep | neural correlates of consciousness | vegetative state | alpha band power

The debate on consciousness has attracted worldwide attention. In one popular theory, consciousness could be defined by two components: awareness (subjective experience, the content of consciousness) and arousal (the levels of wakefulness)[1–3]. Most states of consciousness can be described from these two components (Fig.1A). Different from awake healthy people who have high awareness and arousal, all anesthesia, N3-sleep and coma have low awareness and low arousal[2]. Rapid eye movement sleep (REM-sleep) has relatively high awareness but low arousal as the dreams commonly happen in REM-sleep and reflect rich subjective experience[4]. Patients with unresponsive wakefulness syndrome (UWS, also known as vegetative state) have no awareness but arousal as they have sleep-wake cycle, reflex response and high neural activity in brain stem ascending reticular activating system[5]. Although the majority of studies on consciousness (especially, disorders of consciousness) follow this two-component hypothesis of consciousness, the cortical indexes that can distinguish awareness and arousal are not found yet.

In the most states of consciousness, awareness fluctuates along with arousal. For example, in sleep-wake cycle of healthy person, both awareness and arousal keep decreasing from awake state to N3-sleep[6]. Although REM-sleep was usually adopted to investigate the biomarker for awareness by comparing with N3-sleep, some previous studies suggested that REM-sleep also had higher arousal than N3-sleep[7, 8] (Fig.1A). Thus, the covariation of arousal and awareness generates huge obstacles for disclosing the independent cortical indexes for awareness and arousal, and even undermines the foundation of the two-component hypothesis of consciousness (there might not be independent arousal or awareness in consciousness at all). Facing this dilemma, sleep-wake cycle in patient with UWS provides a solution because patient with UWS is supposed to have no awareness, and then could present the pure arousal fluctuation among the sleep stages. Most importantly, this solution is feasible now. As recent studies had demonstrate that polysomnography (PSG) could successfully discriminate sleep stages in UWS[9–12]), and could acquire electroencephalography (EEG) signals simultaneously which has been commonly used in the investigations on neural substrate of consciousness [9, 13–17] .

Several measurements on EEG signals could be candidates for cortical indexes for arousal or awareness. The spectral slope of the electrophysiological power spectrum, representing the scale-free component of cortical activity, could reflect the levels of consciousness according to the results from sleep and anesthesia states[18–22]. Additionally, there is one recent study which suggested spectral slope as a potential candidate as an arousal index[23]. The spectral entropy and Lempel- Ziv complexity,

Author affiliations: [a]School of Automation Science and Engineering, South China University of Technology, Guangzhou 510641, China.; [b]Research Center for Brain–Computer Interface, Pazhou Laboratory, Guangzhou 510330, China.; [c]School of Artificial Intelligence, South China Normal University, Foshan 528225, China.; [d]School of Psychology, South China Normal University, Guangzhou 510631, China.; [e]Guangdong Provincial Work Injury Rehabilitation Hospital, Guangzhou 510450, China.; [f]South China Brain–Computer Interface Technology Co., Ltd., Guangzhou 510320, China.

Author contributions: Y.Q.L. conceptualized the study. P.M.Q., M.L., and Y.Q.L. designed the methodology. M.L., P.M.Q., J.H.P., and Y.Q.L. conducted the investigation. M.L., Y.B.H., J.H.P., X.Y.B., Y.Z.Y.Y., A.Q.L., Y.H.L., X.C.Y., and D.C. curated the data. M.L. performed the formal analysis. M.L., J.R.H., H.W., and F.W. created the visualizations. Y.Q.L., P.M.Q., and J.H.P. acquired funding. Project administration was managed by P.M.Q. and Y.Q.L. Supervision was provided by Y.Q.L., P.M.Q., and Y.B.H. M.L. and P.M.Q. drafted the manuscript, with critical revisions from P.M.Q., J.H.P., M.L., and Y.Q.L.

The authors declare no competing interests.

*To whom correspondence should be addressed. E-mail: auyqli@scut.edu.cn or qin.pengmin@m.scnu.edu.cn or heyb_17@sina.com

Table 1. Dataset Information.

Dataset	State	Subjects	Gender (M/F)	Age (Mean ± SD)	ABI (%)	TBI (%)	CVD/ID (%)
Sleep dataset1	UWS	16	9/7	51.8 ± 11.43	62.5%	31.25%	6.25%
	MCS	18	15/3	50.47 ± 9.28	55.55%	38.89%	5.55%
	HC	18	8/10	41.7 ± 9.84	–	–	–
Sleep dataset2	HC	37	20/17	26.58 ± 8.19	–	–	–
BCI task dataset1	UWS	45	31/14	39.45 ± 14.09	26.66%	42.22%	31.11%
	MCS	33	20/13	35.9 ± 14.58	15.15%	66.66%	18.18%
BCI task dataset2	UWS	6	3/3	48.8 ± 9.39	50%	50%	0
	MCS	4	3/1	50 ± 23.3	75%	25%	0

Dataset specifies the data collection used in this study, including two Sleep datasets and two BCI task datasets. State represents the clinical state of the participants (UWS, MCS, or HC). Gender is reported as male/female, and Age as mean ± standard deviation. ABI, TBI, and CVD/ID (Other Disorders) indicate the percentages of participants diagnosed with anoxic brain injury, traumatic brain injury, and combined cerebrovascular/inflammatory diseases, respectively. "-" denotes unavailable data.

reflecting the richness and diversity of neural dynamics, have been proposed as potential markers of consciousness levels in many studies(24, 25). The Perturbational Complexity Index, combining TMS-EEG and Lempel-Ziv complexity, has been used to distinguish UWS from minimally conscious state (MCS) who have weak and unstable consciousness(24, 26). Additionally, alpha power is also a usually mentioned EEG measurement which was suggested to be related to levels of consciousness, showing significant reduction in UWS in some previous studies(13, 25, 27, 28). Based on these EEG measurements in the previous studies, we assumed that it should be achievable to disclose the distinct cortical indexes for awareness and arousal by adopting the sleep-wake cycle in both healthy and UWS participants.

To achieve the above aim, the current study included 67 UWS, 55 MCS and 55 healthy controls (HC) in total. See Table 1 for detailed information. The whole study included two parts (Fig.1). In the first part, PSG was used to record nighttime EEG, along with other physiological signals (EOG and EMG) in 16 UWS, 18 MCS and 18 HC. The EEG signals within Wake, N3-sleep, and REM-sleep stages were used to compute spectral slope, spectral entropy, and Lempel-Ziv complexity for each group (Fig.1B). Secondly, the findings from the first part were further validated using an independent sleep EEG dataset in healthy participants (n=37) and two independent BCI task-related EEG datasets (the first including 45 UWS, 33 MCS, the second including 6 UWS, 4 MCS) (Fig.1C). Following two-components hypothesis of consciousness, UWS would have lower awareness than HC and MCS during both Wake state and REM-sleep, but have the similar arousal as the HC and MCS among Wake state, REM-sleep and N3-sleep (Fig.1A). Our results showed that the spectral slope can work as cortical index for arousal while the spectral entropy and Lempel-Ziv complexity can work as the indexes for awareness. These findings demonstrate the two-component theory of consciousness, and shed light on the biological substrate of consciousness.

Result

Altered Sleep Pattern in UWS. The sleep pattern of patients with UWS appeared to be characterized by frequent changes between sleep and wakefulness(29). To evaluate group differences, several sleep architecture metrics including total sleep time (TST), wake time after sleep onset (WASO), sleep efficiency (SE), sleep maintenance efficiency (SME), night wake dominance (NWD), and sleep architecture completeness (SAC) were quantified. Figure.2 shows expert-scored hypnograms for the HC, MCS, and UWS groups.

One-way ANCOVAs were performed to compare these standard sleep metrics from hypnogram between groups (HC, MCS, UWS), respectively, with age as a covariate to control for potential age-related effects. Post-hoc pairwise comparisons with Bonferroni correction were conducted to assess differences between groups. All the sleep metrics showed significant differences among groups (TST: $F(2,47) = 13.971$, $p < 0.001$, $\eta_p^2 = 0.373$; WASO: $F(2,48) = 14.401$, $p < 0.001$, $\eta_p^2 = 0.375$; SE: $F(2,47) = 13.728$, $p < 0.001$, $\eta_p^2 = 0.369$; SME: $F(2,48) = 17.104$, $p < 0.001$, $\eta_p^2 = 0.416$). TST was significantly higher in the HC group compared with the MCS ($p = 0.03$) and UWS groups ($p < 0.001$). MCS group also exhibited higher values than the UWS group ($p = 0.016$). For WASO, UWS and MCS patients exhibited more frequent sleep-wake transitions than HC group (UWS vs. HC: $p < 0.001$; MCS vs. HC: $p = 0.003$). For SE, HC group had higher values than both the MCS ($p = 0.022$) and UWS groups ($p < 0.001$), as well as the MCS group had higher value than the UWS group (p=0.025). Similarly, for SME, the HC group had significantly higher values compared to the MCS ($p = 0.01$) and UWS groups ($p < 0.001$), with the MCS group also exhibiting higher values than the UWS group ($p = 0.01$) (Fig.2B-E). Please see SI Appendix Table S1 for detailed statistical information. Notable differences were observed in SAC and NWD among the three groups, with the UWS group exhibiting the lowest SAC and the most pronounced NWD. As illustrated in Fig.2F-G, only 62.5% of UWS patients exhibited the presence of all standard sleep stages. Furthermore, NWD was absent in HC participants but present in 22.2% of MCS patients and 50% of UWS patients. These results highlight the distinct sleep architecture of each group, characterized by impaired sleep efficiency and maintenance in UWS patients, in contrast to the more stable sleep patterns observed in the HC and MCS groups.

Spectral slope as the cortical index for Arousal. Among the EEG measurements mentioned above, only the spectral slope fits the hypothesis on the arousal. The spectral slope reliably tracked arousal variations across sleep stages in all three

A Theory and Hypothesis

Content of consciousness

Level of arousal

Locked in
Conscious Wakefulness
REM sleep
Drowsiness
Light sleep
Deep sleep
Minimally Conscious state
General anesthesia
Vegetative state
Coma

Cortical index of awareness ?

Cortical index of arousal ?

HC
MCS
UWS

Wake
REM
N3
N3
REM
Wake
N3
REM
Wake

B Feature Extraction and Comparative Analysis

Discovery dataset

HC
MCS
UWS

Overnight recording

Feature extracts

Arousal → Spectral Slope
channels, epochs

Awareness → Spectral Entropy
Lempel-Ziv Complexity
channels, epochs

Comparison within participant

Wake
VS.
N3 VS. REM

Comparison within sleep stage

HC
VS.
MCS VS. UWS

C Independent Dataset Validation

Sleep dataset

Fea-Arousal & Fea-Awareness
Spectral Slope
Spectral Entropy
Lempel-Ziv Complexity
channels, epochs

SVM Classifier
Lemple-Ziv complexity
high / low
Spectral Entropy
high / low
REM Wake
N3
Spectral Slope
low / high

Wake
REM
N3

BCI task dataset1
BCI task dataset2

Fea-Awareness
Spectral Entropy
Lempel-Ziv Complexity
channels, epochs

Fea-Arousal
Spectral Slope
channels, epochs

SVM Classifier
MCS
UWS

MCS
UWS

Fig. 1. Overview of the study design. (A) Theory and Hypothesis. (B) Feature Extraction and Comparative Analysis: Detailing the EEG feature computation and the comparative analysis across different groups and sleep stages. (C) Independent Dataset Validation: Sleep staging and consciousness state diagnosis based on arousal and awareness-related features.

Fig. 2. Sleep Patterns. (A) Exemplary hypnograms: Upper panel represents healthy controls (HC), middle panel depicts minimally conscious state (MCS), and lower panel shows unresponsive wakefulness syndrome (UWS). (B) Total sleep time (TST) for each group, defined as the cumulative duration of N1, N2, N3, and REM sleep stages within the Sleep Period Time (SPT; interval from first to last sleep epoch). (C) Wake after sleep onset (WASO) for each group, representing total wake duration within SPT. (D) Sleep efficiency (SE), calculated as TST / Time in Bed (TIB)× 100 (%), where TIB reflects the total recording duration. (E) Sleep maintenance efficiency (SME), calculated as TST / SPT × 100 (%). (F) Sleep architecture completeness (SAC): Percentage of participants in each group exhibiting all standard sleep stages (N1, N2, N3, and REM) during the recorded sleep period. (G) Night wake dominance (NWD): Percentage of participants with greater wake duration than sleep time during nocturnal hours. The white dot indicates the mean, and the box boundaries indicate the upper and lower quartiles.$*p < 0.05, **p < 0.01, ***p < 0.001$.

groups, especially in UWS. Additionally, the spectral slope in HC shows no difference with MCS and UWS during Wake state. See the following for the detailed information.

For the HC and MCS groups, a one-way repeated measures ANOVA was used to assess the effects of sleep stage. The results showed significant differences in spectral slope between sleep stages within each group (HC: $F(1.404, 23.860) = 242.340$, $p < 0.001$, $\eta_p^2 = 0.934$; MCS: $F(2, 34) = 27.661$, $p < 0.001$, $\eta_p^2 = 0.619$). In post-hoc comparisons with Bonferroni correction, the HC group showed significantly higher spectral slope during Wake state compared to REM-sleep ($p < 0.001$) and N3-sleep ($p < 0.001$), as well as significantly higher spectral slope during REM compared to N3 ($p < 0.001$). Similarly, in the MCS group, spectral slope was highest in Wake state, followed by REM-sleep and N3-sleep, with significant differences between Wake and REM ($p < 0.001$), Wake and N3 ($p < 0.001$), and REM and N3 ($p = 0.011$). For the UWS group, due to missing data for certain sleep stages in several patients (SI Appendix, Table S2), a one-way ANCOVA was performed for comparison between sleep stages, with age included as a covariate. The results showed significant difference among Wake, REM-sleep and N3-sleep ($F(2, 37) = 9.259$, $p < 0.001$, $\eta_p^2 = 0.334$). Post-hoc analysis with Bonferroni correction showed that

the spectral slope was also significantly higher during Wake compared to REM-sleep ($p = 0.039$) and N3-sleep ($p < 0.001$) (Fig.3A).

Furthermore, one-way ANCOVA was performed to compare spectral slope between groups (HC, MCS, UWS) for each sleep stage with age as a covariate, respectively. There were no significant differences among groups in Wake stage, while there were significant differences between among in REM and N3 stage (REM: $F(2, 45) = 4.02$, $p = 0.025$, $\eta_p^2 = 0.152$; N3: $F(2, 44) = 8.271$, $p < 0.001$, $\eta_p^2 = 0.273$). For post-hoc comparisons with Bonferroni corrections, during REM stage, spectral slope in HC group was significantly higher than that in the UWS group ($p = 0.04$), a marginally significant differences were observed between HC and MCS ($p = 0.059$), while no difference between MCS and UWS. During N3-sleep, HC group exhibited significantly lower spectral slope compared to both MCS ($p < 0.001$) and UWS ($p = 0.014$), no significant difference was found between MCS and UWS (Fig.3B). Please see SI Appendix Tables S3-5 for detailed statistical information. These results demonstrated that the spectral slope primarily reflects the arousal dimension of consciousness, rather than the awareness dimension.

The measurements of complexity as cortical index for Awareness. Different from spectral slope, the measurements of

Fig. 3. Arousal-Related Spectral Slope Trends. (A) Violin plots of spectral slope distributions for each sleep stage within HC, MCS, and UWS groups. (B) Violin plots of spectral slope distributions for individual sleep stages across HC, MCS, and UWS groups. The white dot indicates the mean, and the box boundaries represent the upper and lower quartiles. $*p < 0.05, **p < 0.01, ***p < 0.001$.

complexity (spectral entropy and Lempel-Ziv complexity) fit the hypothesis on awareness. HC and MCS showed higher complexity during Wake state and REM-sleep than during N3-sleep, while UWS showed no fluctuation among the sleep stages. Additionally, HC and MCS showed higher complexity than UWS in both Wake state and REM-sleep.

Figure 4A illustrates the variations in spectral entropy across the sleep stages within each group. One-way repeated-measures ANOVA revealed significant differences in spectral entropy across sleep stages in the HC and MCS groups (HC: $F(1.283, 21.803) = 66.073$, $p < 0.001$, $\eta_p^2 = 0.795$; MCS: $F(1.490, 25.328) = 21.773$, $p < 0.001$, $\eta_p^2 = 0.562$). Post-hoc comparisons with Bonferroni correction showed that, in the HC group, spectral entropy was significantly higher during Wake state compared to N3-sleep ($p < 0.001$), higher during REM-sleep compared to N3-sleep ($p < 0.001$). In the MCS group, post-hoc tests with Bonferroni correction revealed significant higher spectral entropy being higher in Wake state compared to N3-sleep ($p < 0.001$), as well as in REM-sleep compared to N3-sleep ($p < 0.001$). Most importantly, in the UWS group, no significant differences were found among sleep stages, indicating the absence of awareness modulation

among the wake-sleep cycle while the arousal still fluctuated independently.

To further validate the role of spectral entropy as a marker of awareness, one-way ANCOVA was performed to compare spectral entropy between groups (HC, MCS, UWS) for each sleep stage, with age as a covariate to control for potential age-related effects (Fig.4B). The results of one-way ANCOVA showed significant difference between groups in all Wake state, REM-sleep and N3-sleep (Wake state: $F(2, 48) = 26.139$, $p < 0.001$, $\eta_p^2 = 0.521$; REM-sleep: $F(2, 45) = 22.422$, $p < 0.001$, $\eta_p^2 = 0.499$; N3-sleep: $F(2, 44) = 8.008$, $p = 0.001$, $\eta_p^2 = 0.267$). In post-hoc comparisons with Bonferroni correction, in the Wake state, spectral entropy was significantly higher in the HC group compared to both the MCS ($p < 0.001$) and UWS groups ($p < 0.001$), with the MCS group also exhibiting significantly higher spectral entropy than the UWS group ($p = 0.006$). In the REM-sleep and N3-sleep stages, the HC group exhibited significantly higher spectral entropy compared to the MCS group (REM-sleep: HC vs. MCS, $p < 0.001$; N3-sleep: HC vs. MCS, $p = 0.008$) and UWS group (REM: HC vs. UWS, $p < 0.001$, N3: HC vs. UWS, $p = 0.002$). No significant differences were found between the MCS and UWS groups.

Fig. 4. Spectral Entropy and Lempel-Ziv Complexity Trends. (A) Mean spectral entropy variations across Wake, N3, and REM stages, combining data from HC, MCS, and UWS groups. (B) Violin plots of spectral entropy distributions for each sleep stage within HC, MCS, and UWS groups. (C) Mean Lempel-Ziv complexity variations across Wake, N3, and REM stages, combining data from HC, MCS, and UWS groups. (D) Violin plots of Lempel-Ziv complexity distributions for individual sleep stages across HC, MCS, and UWS groups. The white dot indicates the mean, and the box boundaries represent the upper and lower quartiles. $*p < 0.05, **p < 0.01, ***p < 0.001$.

Lempel-Ziv complexity as another popular measurement for complexity, a one-way repeated measures ANOVA show significant differences of Lempel-Ziv complexity between sleep stages in HC and MCS group (HC: $F(1.455, 24.743) = 127.414$, $p < 0.001$, $\eta_p^2 = 0.882$; MCS: $F(2, 34) = 27.550$, $p < 0.001$, $\eta_p^2 = 0.618$). As illustrated in Fig.4C, post-hoc comparisons with Bonferroni- correction showed that, in the HC group, both Wake and REM-sleep showed higher Lempel-Ziv complexity than in N3-sleep (for Wake vs. N3-sleep: $p < 0.001$; for REM-sleep vs. N3-sleep: $p < 0.001$). In the MCS group, the post-hoc comparisons with Bonferroni-correction also showed that both Wake and REM-sleep showed higher Lempel-Ziv complexity than in N3-sleep (for Wake vs. N3-sleep: $p < 0.001$; for REM-sleep vs. N3-sleep: $p < 0.001$). However, in the UWS group, one-way ANCOVA (with age as a covariate) showed no significant differences were observed across sleep stages, indicating the absence of awareness modulation during sleep-wake cycle.

As illustrated in Fig.4D, the result of one-way ANCOVA (with age included as a covariate) showed significant difference between groups in Wake, REM and N3 stage (Wake: $F(2, 48) = 27.608$, $p < 0.001$, $\eta_p^2 = 0.535$; REM: $F(2, 45) = 25.246$, $p < 0.001$, $\eta_p^2 = 0.529$; N3: $F(2, 44) = 4.154$, $p = 0.022$, $\eta_p^2 = 0.159$). For post-hoc comparisons with Bonferroni-correction, during Wake and REM, Lempel-Ziv complexity in the HC group was significantly higher than in both MCS (Wake: $p < 0.001$; REM: $p < 0.001$) and UWS (Wake: $p < 0.001$; REM: $p < 0.001$), as well as Lempel-Ziv complexity in the MCS group was significantly higher than in the UWS group (Wake: $p = 0.045$; REM: $p = 0.026$). In the N3 stage, only the HC group exhibited significantly higher Lempel-Ziv complexity compared to the UWS group ($p = 0.021$). Please see SI Appendix Tables S3-5 for detailed statistical information.

Sleep Staging based on cortical index of arousal and awareness. To confirm the effectiveness of arousal and awareness-related features, we evaluated their contributions in sleep stage classification using three analytical approaches on an independent sleep dataset in healthy participants. Fig.5A presents the confusion matrices from the leave-one-subject-out cross-validation (LOSO-CV) analysis, illustrating three classification performances: combining both arousal and awareness-related features, using arousal-related features, and using awareness-related features. Specifically, the combined feature set achieved the highest overall accuracy ($ACC = 0.80$), macro-average F1 score ($MF1 = 0.7745$), and Cohen's Kappa ($K = 0.6864$). In contrast, using only arousal-related features resulted in the lowest classification performance ($ACC = 0.6481$, $MF1 = 0.6481$, $K = 0.4695$), while the awareness-only feature set achieved intermediate performance ($ACC = 0.7617$, $MF1 = 0.7344$, $K = 0.6316$).

Detecting consciousness based on cortical index of awareness. To further confirm the relationship between cortical indexes (awareness and arousal) and consciousness, we conducted classification analyses across two independent DOC groups (both including UWS and MCS). Both groups had the EEG datasets during brain-computer interface tasks (see Materials and Methods, Participants and Data Acquisition for details). The awareness-related features (spectral entropy and Lempel-Ziv complexity) achieved significantly higher classification performance (on UWS and MCS) compared to arousal-related features (spectral slope) which had no ability to discriminate UWS from MCS (see Fig.5B-C). Specifically, for the BCI task dataset1, the accuracy for awareness-related features was 82.05%, with an AUC of 0.84; and for the BCI task dataset2, the accuracy was 90%, with an AUC of 0.92. In contrast, arousal-related feature (spectral slope) resulted in much worse performance, with an accuracy of 51.28% and an AUC of 0.52 on the BCI task dataset1, and 40% accuracy with an AUC of 0.46 on the BCI task dataset2. These findings validate that spectral entropy and Lempel-Ziv complexity could index awareness while spectral slope, as the index of arousal, cannot index awareness.

Reduced alpha power in healthy controls' REM-sleep and UWS. For the HC and MCS groups, a one-way repeated measures ANOVA was used to assess the effects of sleep stage. The results showed significant differences in alpha power across sleep stages within each group (HC: $F(1.458, 24.789) = 20.006$, $p < 0.001$, $\eta_p^2 = 0.541$, MCS: $F(1.372) = 9.420$, $p = 0.003$, $\eta_p^2 = 0.357$). Post-hoc comparisons with Bonferroni correction showed that in the HC group, alpha power during REM-sleep was significantly lower than during Wake state ($p < 0.001$) and N3-sleep ($p < 0.001$). In the MCS group, both REM-sleep and N3-sleep exhibited significantly lower alpha power compared to Wake state (REM vs. Wake: $p < 0.001$; N3 vs. Wake: $p = 0.017$). However, in the UWS group, a one-way ANCOVA (with age as a covariate) revealed no significant differences in alpha power across sleep stages.

Furthermore, one-way ANCOVA was performed to compare alpha power between groups (HC, MCS, UWS) for each sleep stage with age as a covariate, respectively. The result showed significant difference between groups in Wake state, REM-sleep and N3-sleep (Wake: $F(2, 48) = 26.848$, $p < 0.001$, $\eta_p^2 = 0.528$; REM: $F(2, 45) = 17.463$, $p < 0.001$, $\eta_p^2 = 0.437$; N3: $F(2, 44) = 20.700$, $p < 0.001$, $\eta_p^2 = 0.485$). Post-hoc comparisons with Bonferroni correction showed that alpha power in the UWS group was significantly lower than in both the HC and MCS groups during Wake (vs. HC: $p < 0.001$; vs. MCS: $p = 0.003$), REM (vs. HC: $p < 0.001$; vs. MCS: $p < 0.001$). In addition, the MCS group exhibited significantly lower alpha power than the HC group during Wake ($p < 0.001$) and N3 ($p < 0.001$). For detailed statistical results, see SI Appendix Tables S3–S5 and fig. S1.

Discussion

The current study distinguishes the awareness from arousal with independent cortical indexes for each other. In specific, the spectral slope can work as a cortical index of arousal while the measurements of complexity (spectral entropy and Lempel-Ziv complexity) work as the cortical indexes of awareness. It is the first time to use cortical index describes the pure arousal fluctuation during sleep-wake cycle in patients with UWS, and demonstrates the hypothesis that the UWS has the similar arousal level as the healthy control. These findings strongly support the two-components hypothesis of consciousness.

The spectral slope works as an index for arousal. In the last two decades, most studies took the opinion that the patients with UWS have similar arousal level as healthy participants during wakefulness(30, 31). However, the evidence supporting

Fig. 5. Evaluation of Sleep Staging Performance and Consciousness Diagnosis. (A) Cross-validated classification performance under different feature combinations. Left: Joint use of arousal-related (spectral slope) and awareness-related features (spectral entropy, Lempel-Ziv complexity) with leave-one-subject-out cross-validation (LOSO-CV). Middle: Arousal-related features (spectral slope). Right: Awareness-related features (spectral entropy, Lempel-Ziv complexity). (B) Awareness driven diagnostic performance: Top row, Confusion matrices for BCI task dataset1 (left) and BCI task dataset2 (right); Bottom row, ROC curves (BCI task dataset1 AUC: 0.84, BCI task dataset2 AUC: 0.92) confirm robust MCS-UWS spectral entropy arability. (C) Arousal driven diagnostic performance: Top row, Confusion matrices for identical datasets; Bottom row, ROC curves (dataset1 AUC: 0.52, dataset2 AUC: 0.46) approaching chance-level discrimination.

this assumption is only from sleep-wake cycle in behavioral level(5, 32, 33) and intact brain stem activity shown by PET(3, 34)in patients with UWS. Since there was no any evidence from cortical activity, these findings cannot eliminate the suspicion whether the patients with UWS have the similar brain arousal level as healthy individuals or not. In order to discover the cortical index for arousal, recent studies adopted fMRI or EEG, and acquired the signals from multiple altered conscious states (i.e., N3-sleep, deep anesthesia, REM-sleep and UWS)(20, 23, 31). Unfortunately, in these studies, all the contrasts based on these states cannot rule out the mutual influence of the two components. For example, although REM-sleep have relative high awareness and low arousal(23, 24), it still shows higher value in both arousal and awareness compared to the N3-sleep. In the current study, following the hypothesis that patients with UWS have no awareness, their sleep-wake cycle could reflect the fluctuation of arousal purely, which could eliminate the mutual influence between awareness and arousal. Standing on this point, our results showed the spectral slope varied among the sleep stages in

UWS in the same way as in MCS and healthy controls, could work as the cortical index for arousal.

The spectral slope reflects a 1/f dynamic in EEG signal, i.e. higher frequency activity exhibits reduced power compared with lower frequency activity, which has been highlighted its importance in consciousness(35–37). Previous studies indicated that the general anesthesia had more negative spectral slope values compared with awake states(18, 38). In the literature on sleep, within a brand frequency band (0.5-45Hz), N3-sleep showed the most negative spectral slope values compared with REM-sleep and Wake state while the REM-sleep had more negative spectral slopes value than Wake state(21, 39–41). The current study showed consistent results with the previous studies in healthy participants. Most importantly, the current study showed that the fluctuation of spectral slope during sleep stages in UWS patients was similar as healthy participants. Additionally, spectral slope showed no difference among UWS, MCS, HC during Wake state, and could not discriminate the MCS from UWS in the two independent datasets. These demonstrate that the

spectral slope does reflect the arousal level, but not the awareness. Two issues need to be noticed. First, although the most previous studies indicated that the REM-sleep might have higher arousal than N3-sleep as it has more brain activities(7, 8), some other studies showed REM-sleep had more negative spectral slope values compared with NREM-sleep and wake state within a narrow frequency band (30-45Hz or 30-50Hz)(21, 23). The current study reproduced this finding in healthy participants but not in UWS (SI Appendix Fig. S3 and S4). This narrow band spectral slope might not work as a biomarker of arousal. Second, N3-sleep showed more reduced spectral slope in healthy controls compared with UWS and MCS, which may indicate the intact function of brainstem and basal forebrain(42).

The complexity of cortical activity indexes awareness. The links between complexity and awareness have been the focus of attention(43). According to the integrated information theory (IIT) and similar ones(44, 45), a minimal level of complexity of cortical activities is needed to support conscious experience (e.g., awareness). In fact, some previous studies reported that complexity of cortical activities diminishes in the absence of awareness(41, 46–49). The spectral entropy and Lempel-Ziv complexity, two popular measurements of complexity, showed reduced values in unconscious states (UWS, N3-sleep, and anesthesia) compared with conscious states (awake, REM-sleep)(24, 27, 50–53). Although the previous studies intended to highlight the relationship between complexity and awareness by showing the higher complexity during REM-sleep(24), there is still a suspicion that REM-sleep in healthy participants may have a higher arousal than unconscious states (i.e., N3-sleep or UWS), in other words, the difference of complexity between REM-sleep and unconscious states might reflect the arousal difference, but not the awareness difference. Beyond these previous studies, our results specify the relationship between complexity and awareness by demonstrating patients with UWS had similar arousal as healthy controls in Wake state (higher than REM-sleep). Furthermore, spectral entropy and Lempel-Ziv complexity did not fluctuate with sleep-wake cycle in UWS where the arousal (e.g. spectral slope) did, and could discriminate MCS from UWS patients in the two independent datasets. These further clarify the relationship between complexity and awareness. Finally, both spectral entropy and Lempel-Ziv complexity showed higher value in healthy controls than UWS during N3-sleep. This may indicate that there is more information processing in unconscious healthy brain than UWS, which is supported by the previous studies on the cognitive processing during N3-sleep(54–56).

Both UWS and MCS had detectable sleep stages during night-time. However, they showed altered sleep structure compared with healthy controls. Compared with MCS patients, UWS patients had shorter sleep duration. Additionally, there were 62.5% of the UWS patients had all sleep stages while all MCS patients had all sleep stages (Fig.2, SI Appendix, Table S2). These results were consistent with the previous studies(12, 29, 33, 57–61). Besides the difference of sleep structure between the levels of consciousness (UWS and MCS), the previous studies also intended to investigate the relationship between sleep elements and the outcome of the patients with UWS and MCS(9, 42, 62–65), and found the positive relationship between sleep structure and better outcome(10, 66, 67). For example, the recent studies showed that the sleep spindles, K-complex, and REM-sleep could predict the favorable outcome of patients(9, 64, 65). The well-formed sleep spindles could even predict the detection of patients with cognitive-motor dissociation (CMD)(9). All the above findings indicated that the sleep structure could reflect the remaining brain function and have important clinical values in patients with UWS and MCS. The current study extended the current knowledge, for the first time describing the pure arousal fluctuation during sleep-wake cycle using cortical index (e.g. spectral slope).

The relationship between alpha power and awareness was also tested as some previous studies suggested that alpha power could be a biomarker of awareness(13, 25, 27, 28, 68–71). Consistent with the studies, our results showed that healthy controls and MCS had stronger alpha power than UWS during Wake state. Furthermore, the current study showed that there was no difference among Wake state, REM-sleep and N3-sleep in UWS. However, our results also showed that both Wake state and N3-sleep had stronger alpha power than REM-sleep in healthy controls (SI Appendix, Fig. S1 and S2), which was also supported by previous study(72). Considering that one in REM-sleep might have more conscious experience (e.g., awareness) than N3-sleep, this result might indicate that alpha could not reflect the level of awareness. Since the locked-in syndrome, who had lesions in brain stem, also showed reduced alpha power during awake states(73), the reduced alpha power in REM-sleep might due to the reduced functional integrity between brain stem and cortices(73, 74). Furthermore, one recent study found that alpha power was suppressed only in UWS with severe post-anoxia but failed to discriminate un/consciousness in other aetiologies(75). They supposed that the alpha power reduction might be related to the diffuse cortical damage. Taken together, the alpha power might not index awareness.

In conclusion, combining the brain signal during sleep-wake cycle in UWS and healthy controls, we found the cortical indexes for arousal (e.g., spectral slope), as well as for awareness (the measurements of complexity, e.g., spectral entropy and Lempel-Ziv complexity), supported the two-components theory of consciousness. The cortical indexes for awareness may be a useful tool for the diagnosis of patients with disorders of consciousness, or for monitoring the levels of sedation. Furthermore, it will be important for understanding the neural substrate of consciousness to investigate mechanism for the interaction between arousal and awareness in future.

Materials and Methods

Participants and data acquisition.

Sleep dataset1. The discovery dataset comprised 52 participants, including 34 patients with chronic disorders of consciousness (DOC) and 18 HC. Data were collected between February 2023 and January 2024 at the Guangdong Work Injury Rehabilitation Hospital. Among the DOC patients, 16 were diagnosed with UWS (mean age: 51.8 ± 11.43 years; male/female: 9/7), and 18 with MCS (mean age: 50.47 ± 9.28 years; male/female: 15/3). The HC group (mean age: 41.7 ± 9.84 years; male/female: 8/10). Clinical diagnosis was confirmed using the Coma Recovery Scale-Revised (CRS-R), which was administered one week before and on the day of the experiment, with a follow-up assessment conducted three months later. In each assessment period, a minimum of two

CRS-R assessments were performed conducted by two experienced physicians.

Patients were included according to the following criteria: a diagnosis of UWS or MCS and age between 18 and 70 years (to minimize age dependent effects on sleep). Patients with severe or unstable clinical conditions, pre-existing central nervous system impairments unrelated to disorders of consciousness, or recent use of barbiturates, antipsychotics, or antidepressants within 24 hours were excluded.

Sleep data was collected during overnight using Compumedics Grael Profusion PSG system (version 4.5, build 500, Compumedics Limited). The average recording duration was 8.402 ± 0.54 hours. Recordings included EEG, EOG, and EMG signals. EEG data were recorded according to the international 10–20 system (F3, F4, C3, C4, O1, O2) with a sampling frequency of 256 Hz. EOG and EMG signals were simultaneously captured to provide comprehensive sleep monitoring. Video recordings were synchronized with EEG to annotate key behavioral events (eye-opening moments, rest and agitation for endogenous movements and nursing for exogenous movements). Electrode impedance was maintained below 5 kΩ.

Ethical approval was obtained (approval number: AF/SC-07/2020.03), and informed consent was acquired from legal guardians in accordance with the Declaration of Helsinki. Detailed information can be found in SI Appendix, Table S6.

Sleep dataset2. This independent sleep dataset comprised recordings collected from 37 healthy volunteers (mean age: 26.58 ± 8.19 years, male/female ratio: 20/17), recruited from South China University of Technology between June 2018 and January 2023. Sleep data were collected during overnight polysomnography session using the SynAmps2 amplifier (Compumedics, Neuroscan, Inc., Australia) and a 32-channel EEG cap according to the international 10-20 system. The average recording duration was 7.02 ± 1.34 hours. EEG signals were sampled at 250 Hz, and electrode impedance was maintained below 5 kΩ.

BCI task dataset1. The dataset used in this study originates from a brain-computer interface (BCI) experiment conducted with DOC patients performing an audiovisual recognition task. This dataset, previously published in a study investigating the prognostic implications of BCI-based assessments in patients with DOC[76], comprises EEG recordings from 78 DOC patients , including 45 UWS (mean age: 39.45 ± 14.09 years, male/female: 31/14), and 33 MCS (mean age: 35.9 ± 14.58 years, male/female: 20/13). The data were collected during item-selection tasks aimed at detecting conscious awareness at the General Hospital of Guangzhou Military Command of People's Liberation Army, China, between October 2014 and August 2018. EEG signals were recorded using a 32-channel EEG cap positioned according to the international 10–20 system, with a sampling frequency of 250 Hz. Three BCIs were used, including a hybrid P300 and SSVEP BCI based on photograph stimuli (photograph paradigm), a hybrid P300 and SSVEP BCI based on visual number stimuli (number paradigm), and an audiovisual BCI based on audiovisual number stimuli (audiovisual paradigm). Thirty-five patients participated in the experiment based on the photograph paradigm, 13 in the experiment based on the number paradigm, and 30 in the experiment based on the audiovisual paradigm. In this study, we repurposed this dataset to validate cortical indexes for awareness.

BCI task dataset2. The BCI task dataset 2 was collected from an audiovisual multimodal BCI experiment, designed to integrate P300 and auditory steady-state response (ASSR) for assessing awareness in DOC patients. The data were acquired at the Guangdong Work Injury Rehabilitation Hospital, Guangzhou, China, between October 2022 and April 2023, using a SynAmps2 amplifier (Compumedics, Neuroscan, Inc., Australia). Although this dataset was obtained from the same hospital as the sleep dataset1, it was collected from a different cohort of patients with DOC patients.

The dataset included 10 DOC patients, comprising 6 UWS patients (mean age: 48.8 ± 9.39 years, male/female: 3/3) and 4 MCS patients (mean age: 50 ± 23.3 years, male/female: 3/1). EEG recordings were conducted over a 5-minute period using a 32-channel EEG cap, following the international 10–20 system, with a sampling frequency of 250 Hz. To ensure high signal fidelity,

electrode impedance was maintained below 5 kΩ During each trial, the two facial photographs flashed in a random order and the same side's earphone played the corresponding sound when each photo flashed. As with the BCI task dataset1, we repurposed this dataset to validate cortical indexes for awareness. Detailed information can be found in SI Appendix, Table S7.

Sleep Scoring. Polysomnography recordings (EEG, EOG, EMG) were segmented into 30-s epochs for sleep scoring. Given the abnormal sleep patterns in UWS and MCS patients, the standard sleep scoring criteria used in healthy individuals (77) cannot be applied directly, but have to be adjusted to UWS and MCS patients sleep patterns(12, 29, 57). In this study, we applied these adjusted criteria for sleep scoring in UWS and MCS patients.

Specifically, given that UWS and MCS patients may show local intermittent rhythmic delta activity that might be confused with sleep related delta waves, all six EEG electrodes (F3, F4, C3, C4, O1, O2) were utilized for visual sleep scoring to ensure accurate interpretation. Wake state was defined by relatively high EEG frequency, the presence of eye blinks (from video), and high EMG. N1-sleep was scored when the EEG showed a general slowing, accompanied by the disappearance of blinks and decreased muscle activity (rolling eye movements were considered optional indicators). N2-sleep was identified based on the presence of K-complexes and, occasionally, typical sleep spindles (0.5–3 s duration, 20–100 μV amplitude, 11–16 Hz frequency) or prolonged "atypical" slow spindles (10–12 Hz), along with EEG slowing, the emergence of slow-wave activity, and a further reduction in muscle tone, provided that slow waves < 20% of the epoch. N3-sleep was scored when > 20% of the epoch consist of slow-wave occurring synchronously across intact cortical regions. Given that EEG amplitude in UWS and MCS patients may be affected by various extracerebral factors, both standard high-amplitude slow waves (> 75 μV) and dominant low-voltage slow waves (< 75 μV) were considered indicative of slow-wave sleep (SWS). REM-sleep was defined as fast, low-voltage EEG activity with minimal muscular tone and the presence of muscle atonia and EEG characteristics were prioritized over EOG features(78).

All sleep data were independently scored by two expert clinical neurophysiologists. Discrepancies in sleep scoring and arousal recognition were reviewed and resolved through consensus. Uncertain epochs were excluded from statistical analyses to maintain data integrity.

Standard Sleep Architecture Metrics. Total sleep time (TST) was defined as the cumulative duration of all sleep stages (N1, N2, N3, and REM) within the sleep period. Wake time after sleep onset (WASO) was measured as the total duration of wake episodes occurring after the initial sleep onset. Sleep efficiency (SE) is defined as the percentage of total sleep time relative to the total time in bed (TIB, the total duration of the recording period) and was computed as follows:

$$SE(\%) = \frac{TST}{TIB} \times 100 \qquad [1]$$

Sleep maintenance efficiency (SME) was defined as the percentage of total sleep time relative to the total sleep period time (SPT):

$$SME(\%) = \frac{TST}{SPT} \times 100 \qquad [2]$$

Sleep architecture completeness (SAC) was defined as the proportion of individuals exhibiting a complete sleep architecture, characterized by the presence of all major sleep stages (N1, N2, N3, and REM) during the sleep period. It was computed as:

$$SAC(\%) = \frac{N_{\text{complete sleep architecture}}}{N_{\text{total}}} \times 100 \qquad [3]$$

where $N_{\text{complete sleep architecture}}$ represents the number of individuals experiencing all major sleep stages, and N_{total} is the total number of individuals analyzed.

Night wake dominance (NWD) was used to assess the proportion of individuals whose total wake time exceeded total sleep time during the recording period. The calculation was performed as:

$$NWD(\%) = \frac{N_{\text{wake > sleep}}}{N_{\text{total}}} \times 100 \qquad [4]$$

where $N_{\text{wake} > \text{sleep}}$ represents the number of individuals whose total wake time was greater than their TST.

EEG Preprocessing. A unified preprocessing pipeline was rigorously applied to both sleep EEG datasets (the primary discovery dataset, Sleep dataset1, and the independent sleep dataset, Sleep dataset2) and BCI task datasets (dataset1 and dataset2) using MNE-Python (v1.6.0) integrated with NumPy (v1.22.4) in Python 3.9.6. To ensure methodological consistency across modalities, six electrodes (F3, F4, C3, C4, O1, O2) following the international 10–20 system were selected, encompassing frontal, central, and occipital regions relevant to both sleep and task-related neural dynamics.

The EEG signals were initially processed with a 50 Hz notch filter (4th-order Butterworth) to remove power-line interference, followed by band-pass filtering in the range of 0.5–45 Hz using a finite impulse response (FIR) Hamming window filter. The signals were then re-referenced to the global average of all non-excluded channels to minimize artifacts from electrode displacement during prolonged recordings. Artifact rejection was performed manually by expert clinical neurophysiologists through visual inspection, using criteria such as an amplitude threshold of ± 100 μV to identify and exclude segments contaminated by noise. In order to reduce inter-channel variability, z-score normalization was applied independently to each channel. Finally, the signals were down sampled to 100 Hz and segmented into non-overlapping 30-second epochs.

Feature Extraction. For the discovery dataset (i.e., Sleep dataset 1; comprising UWS, MCS, and HC groups), which was primarily used for statistical analysis, feature extraction was restricted to three sleep stages: Wake-state (high arousal/high awareness), REM-sleep (low arousal/high awareness), and N3-sleep (low arousal/low awareness). Features were initially computed at the epoch level and then averaged for each stage within each subject. For each subject, all three sleep stages (Wake, REM, and N3) included more than 7 epochs (i.e., at least 3.5 minutes of data per stage). Additionally, to obtain a more generalized representation of neural activity, features were further averaged across EEG channels before statistical comparisons. In contrast, for the independent sleep and BCI task datasets (used for sleep staging and consciousness diagnosis, respectively), each epoch was treated as an independent sample.

Spectral Slope. The spectral slope is a quantitative feature characterizing the power distribution across frequency components, derived from the power spectral density (PSD) analysis of neural signals(19, 21, 23). In this study, spectral slope was computed for each channel and epoch following the methods used in previous studies(23, 79). To obtain the spectral slope, we fitted a linear regression line to the power spectrum in log-log space between 0.5 and 45 Hz.

Formally, for a given temporal signal $x(t)$, the PSD of the signal $x(t)$ is computed using the "welch" method to ensure robust and smoothed spectral estimation. The PSD is denoted as $P(f)$ where f represents the frequency. The PSD is then transformed into the logarithmic domain:

$$(\log f, \log P(f)) \quad [5]$$

this transformation linearizes the relationship between frequency and power in the log-log space, enabling a more interpretable slope calculation.

A linear regression is performed on the log-transformed PSD values within a predefined frequency range (0.5–45 Hz). The spectral slope is defined as the slope of the regression line:

$$Slope = \frac{\Delta(\log P(f))}{\Delta(\log f)} \quad [6]$$

Spectral Entropy. Spectral entropy(SEp) quantifies the uniformity or peakedness of the EEG power spectral distribution within a specific frequency range(80). It reflects the degree of randomness or predictability in the frequency components of the signal and provides an indicator of its sinusoidal characteristics. A lower SEp suggests that the signal is dominated by fewer frequency components, with the spectral power concentrated around one or several discrete frequencies, thereby exhibiting a more pronounced

sinusoidal nature. In contrast, a higher SEp value indicates a greater number of frequency components with a broader spectral distribution, signifying reduced sinusoidal characteristics and increased complexity of the signal. Building upon established methodologies in neurophysiological signal analysis(25), we implemented the calculation of spectral entropy within the 0.5-45 Hz bandwidth through three systematic phases for each channel and epoch.

For a continuous signal $x(t)$, the formula can be defined as follows:

$$X(f) = \int_{-\infty}^{\infty} x(t)e^{-j2\pi ft}dt \quad [7]$$

The power spectral density of each frequency component is calculated as follows:

$$\text{PSD}(f) = |X(f)|^2 \quad [8]$$

where $|X(f)|^2$ represents the power at frequency, which is the energy of the signal at that frequency. P denotes is the relative power of frequency bands f :

$$P(f) = \frac{\text{PSD}(f)}{\sum_f \text{PSD}(f)} \quad [9]$$

The spectral entropy (SEp) is defined as:

$$SE\text{p} = -\sum_f P(f)\log(P(f)) \quad [10]$$

Lempel-Ziv complexity. Lempel-Ziv Complexity (LZC) was used to quantify the complexity of EEG signals by assessing their temporal dynamics and unpredictability. In this study, LZC was computed for each channel and epoch following the calculation methods used in previous studies(81–83). We employed a mean symbolic encoding method to compute LZC, where the continuous EEG signal was discretized by mapping it to symbolic representations based on its mean value within a sliding window. To ensure that LZC was computed within the same frequency range as the spectral slope, a 0.5 Hz high-pass filter and a 45 Hz low-pass filter were applied to the EEG signals before computation.

The computation of LZC begins with the preprocessing of the continuous signal $x(t)$, which is divided into non-overlapping windows of size. For each window $X_i = \{x_1, x_2, \ldots, x_L\}$, the mean value of the window μ_i is calculated as:

$$\mu_i = \frac{1}{L}\sum_{j=1}^{L} x_j \quad [11]$$

This mean value μ_i is then used to symbolize the window. Specifically, the symbol for each window is determined by comparing the mean value μ_i to predefined thresholds or bins. The EEG signal was transformed into a binary sequence . The Lempel-Ziv parsing algorithm sequentially scans , identifying unique subsequences $S = \{s_1, s_2, \ldots, s_N\}$. The complexity count $C(N)$ increases whenever a new, non-redundant subsequence is encountered. To account for sequence length dependency, the raw complexity count $C(N)$ was normalized by the theoretical upper bound of a random binary sequence:

$$LZC = \frac{C(N)}{N/\log_2(N)} \quad [12]$$

where N is the length of the binary sequence. This normalized LZC provides a robust measure of EEG complexity, facilitating comparisons across different conditions and subjects.

Machine learning-based classification modelling. To evaluate the effectiveness of the identified arousal- and awareness-related features, we constructed a classification framework designed for two tasks: sleep staging and consciousness state diagnosis. The primary motivation was to validate the effectiveness of the cortical index of arousal and awareness.

Sleep staging. In this study, three separate Support Vector Machine (SVM) classifiers with radial basis function (RBF) kernels were constructed to evaluate the contribution of distinct feature domains to sleep staging. Specifically, the classifiers were trained using arousal-related features (spectral slope), awareness-related features (spectral entropy and Lempel-Ziv complexity), and a combination of both feature types (cortical index of arousal and awareness). All features were extracted from preprocessed EEG data during Wake, N3, and REM stages, as described in the Feature Extraction section.

To account for class imbalance across sleep stages, all classifiers were implemented using scikit-learn (v1.1.3) with class weight = "balanced". Sleep staging performance was evaluated using leave-one-subject-out cross-validation (LOSO-CV), where each subject was iteratively held out for testing while the remaining data were used for training.Performance metrics included overall classification accuracy (ACC), macro-averaged F1 score (MF1), and Cohen's Kappa coefficient (K).

Consciousness state diagnosis. In this study, two separate Support Vector Machine (SVM) classifiers with radial basis function (RBF) kernels were constructed using wake-stage EEG data from the discovery dataset (comprising UWS and MCS groups). One classifier was trained exclusively on the arousal-related feature (spectral slope), while the other utilized awareness-related features (spectral entropy and Lempel-Ziv complexity). Feature extraction procedures followed the same pipeline described in the Feature Extraction section.

To validate the effectiveness of the awareness-related cortical index in distinguishing state of consciousness, both classifiers were independently validated on two external BCI task datasets (dataset 1 and dataset 2), with strict alignment between feature types during training and testing (i.e., the arousal classifier was tested only on spectral slope, and the awareness classifier on spectral entropy and Lempel-Ziv complexity).

For each epoch, the classifiers generated probabilistic outputs indicating the likelihood of the predicted consciousness state (MCS or UWS). To obtain a subject-level diagnosis, a majority-vote strategy was applied across all epochs: if the aggregated probability of being classified as MCS exceeded 0.5, the subject was labeled as MCS; otherwise, UWS. This approach accounted for the dynamic and fluctuating nature of consciousness, particularly in patients with severe brain injuries. The classification performance was assessed using accuracy and the area under the receiver operating characteristic curve (AUC) as primary evaluation metrics.

Statistical analysis. All statistical analyses were conducted using SPSS (v27.0, IBM Corp., Armonk, NY, USA). For the within-group comparisons (HC and MCS groups), a one-way repeated measures ANOVA was used to assess the effects of sleep stage (Wake, N3, REM). Post-hoc pairwise comparisons were conducted with Bonferroni correction to control for multiple comparisons and identify significant differences between sleep stages within each group. For the UWS group, a one-way ANCOVA was performed separately for each sleep stage, with age as a covariate, due to missing data for certain sleep stages. Post-hoc pairwise comparisons were also conducted for the UWS group, with Bonferroni correction applied to control for multiple comparisons.

For between-group comparisons (HC, MCS, UWS groups), a one-way ANCOVA was performed separately for each sleep stage (Wake, REM, N3), with age as a covariate to control for potential age-related effects. Post-hoc pairwise comparisons with Bonferroni correction were conducted to assess differences between the three groups.The assumptions of sphericity were assessed using Mauchly's test, and when violated, the Greenhouse-Geisser correction was applied to adjust the degrees of freedom. Outliers beyond $\pm 2SD$ from the mean were excluded.

ACKNOWLEDGMENTS. We thank all participants for their contribution to this study. This work was supported by the STI 2030 Major Projects (Grant 2022ZD0208900), the Key Research and Development Program of Guangdong Province (Grant 2018B030339001), the Guangdong Basic and Applied Basic Research Foundation (Grants 2024A1515011429 and 2023A1515140100), the National Natural Science Foundation of China (Grant 32371098), the Research Center for Brain Cognition and Human Development, Guangdong (Grant 2024B0303390003), and the Guangzhou Talent Plan (Grant 2024D02J0008).

1. S Laureys, AM Owen, ND Schiff, Brain function in coma, vegetative state, and related disorders. *The Lancet Neurol.* **3**, 537–546 (2004).
2. V Cologan, et al., Sleep in disorders of consciousness. *Sleep medicine reviews* **14**, 97–105 (2010).
3. S Laureys, The neural correlate of (un) awareness: lessons from the vegetative state. *Trends cognitive sciences* **9**, 556–559 (2005).
4. JM Siegel, Rem sleep: a biological and psychological paradox. *Sleep medicine reviews* **15**, 139–142 (2011).
5. P Avantaggiato, et al., Polysomnographic sleep patterns in children and adolescents in unresponsive wakefulness syndrome. *The J. Head Trauma Rehabil.* **30**, 334–346 (2015).
6. P Halász, M Terzano, L Parrino, R Bódizs, The nature of arousal in sleep. *J. sleep research* **13**, 1–23 (2004).
7. L Parrino, et al., The 70 years since the discovery of rapid eye movement sleep: history, electroencephalographic features and unsolved issues. *J. Sleep Res.* p. e14342 (2024).
8. B Baird, SA Mota-Rolim, M Dresler, The cognitive neuroscience of lucid dreaming. *Neurosci. & Biobehav. Rev.* **100**, 305–323 (2019).
9. EE Carroll, et al., Sleep spindles as a predictor of cognitive motor dissociation and recovery of consciousness after acute brain injury. *Nat. Medicine* pp. 1–8 (2025).
10. J Nekrasova, et al., Retrospective analysis of sleep patterns in patients with chronic disorders of consciousness. *Sleep Medicine: X* **2**, 100024 (2020).
11. T Wielek, et al., Sleep in patients with disorders of consciousness characterized by means of machine learning. *PloS one* **13**, e0190458 (2018).
12. DR Sebastiano, et al., Sleep patterns associated with the severity of impairment in a large cohort of patients with chronic disorders of consciousness. *Clin. Neurophysiol.* **129**, 687–693 (2018).
13. X Ma, et al., How well do neural signatures of resting-state eeg detect consciousness? a large-scale clinical study. *Hum. Brain Mapp.* **45**, e26586 (2024).
14. DA Engemann, et al., Robust eeg-based cross-site and cross-protocol classification of states of consciousness. *Brain* **141**, 3179–3192 (2018).
15. JL Kästle, B Anvari, J Krol, HA Wurdemann, Correlation between situational awareness and eeg signals. *Neurocomputing* **432**, 70–79 (2021).
16. O Dressler, G Schneider, G Stockmanns, E Kochs, Awareness and the eeg power spectrum: analysis of frequencies. *Br. journal anaesthesia* **93**, 806–809 (2004).
17. S Stefan, et al., Consciousness indexing and outcome prediction with resting-state eeg in severe disorders of consciousness. *Brain Topogr.* **31**, 848–862 (2018).
18. MA Colombo, et al., The spectral exponent of the resting eeg indexes the presence of consciousness during unresponsiveness induced by propofol, xenon, and ketamine. *NeuroImage* **189**, 631–644 (2019).
19. Y Zhang, et al., Eeg spectral slope: A reliable indicator for continuous evaluation of consciousness levels during propofol anesthesia. *NeuroImage* **283**, 120426 (2023).
20. S Widmann, et al., Aperiodic component of the electroencephalogram power spectrum reflects the hypnotic level of anaesthesia. *Br. J. Anaesth.* **134**, 392–401 (2025).
21. C Höhn, MA Hahn, JD Lendner, K Hoedlmoser, Spectral slope and lempel–ziv complexity as robust markers of brain states during sleep and wakefulness. *eneuro* **11** (2024).
22. F Zilio, et al., Are intrinsic neural timescales related to sensory processing? evidence from abnormal behavioral states. *NeuroImage* **226**, 117579 (2021).
23. JD Lendner, et al., An electrophysiological marker of arousal level in humans. *elife* **9**, e55092 (2020).
24. AG Casali, et al., A theoretically based index of consciousness independent of sensory processing and behavior. *Sci. translational medicine* **5**, 198ra105–198ra105 (2013).
25. A Piarulli, et al., Eeg ultradian rhythmicity differences in disorders of consciousness during wakefulness. *J. neurology* **263**, 1746–1760 (2016).
26. JT Giacino, et al., Comprehensive systematic review update summary: disorders of consciousness: report of the guideline development, dissemination, and implementation subcommittee of the american academy of neurology; the american congress of rehabilitation medicine; and the national institute on disability, independent living, and rehabilitation research. *Neurology* **91**, 461–470 (2018).
27. JD Sitt, et al., Large scale screening of neural signatures of consciousness in patients in a vegetative or minimally conscious state. *Brain* **137**, 2258–2270 (2014).
28. ES Lutkenhoff, et al., Eeg power spectra and subcortical pathology in chronic disorders of consciousness. *Psychol. Medicine* **52**, 1491–1500 (2022).
29. I Mertel, et al., Sleep in disorders of consciousness: behavioral and polysomnographic recording. *Bmc Medicine* **18**, 1–14 (2020).
30. M Lee, et al., Quantifying arousal and awareness in altered states of consciousness using interpretable deep learning. *Nat. communications* **13**, 1064 (2022).
31. J Han, et al., The neural correlates of arousal: Ventral posterolateral nucleus-global transient co-activation. *Cell Reports* **43** (2024).
32. C Blume, R Del Giudice, M Wislowska, J Lechinger, M Schabus, Across the consciousness continuum—from unresponsive wakefulness to sleep. *Front. human neuroscience* **9**, 105 (2015).
33. E Landsness, et al., Electrophysiological correlates of behavioural changes in vigilance in vegetative state and minimally conscious state. *Brain* **134**, 2222–2232 (2011).
34. S Laureys, ND Schiff, Coma and consciousness: paradigms (re) framed by neuroimaging. *Neuroimage* **61**, 478–491 (2012).

35. L Wojtecki, et al., Modulation of central thalamic oscillations during emotional-cognitive processing in chronic disorder of consciousness. *Cortex* **60**, 94–102 (2014).

36. Y Huang, et al., Spectral and phase-amplitude coupling signatures in human deep brain oscillations during propofol-induced anaesthesia. *Br. J. Anaesth.* **121**, 303–313 (2018).

37. Y Huang, et al., Characteristics of thalamic local field potentials in patients with disorders of consciousness in *2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*. (IEEE), pp. 3779–3782 (2015).

38. R Gao, Interpreting the electrophysiological power spectrum. *J. neurophysiology* **115**, 628–630 (2016).

39. Y Shen, E Olbrich, P Achermann, P Meier, Dimensional complexity and spectral properties of the human sleep eeg. *Clin. Neurophysiol.* **114**, 199–209 (2003).

40. E Pereda, A Gamundi, R Rial, J González, Non-linear behaviour of human eeg: fractal exponent versus correlation dimension in awake and sleep stages. *Neurosci. letters* **250**, 91–94 (1998).

41. V Miskovic, KJ MacDonald, LJ Rhodes, KA Cote, Changes in eeg multiscale entropy and power-law frequency scaling during the human sleep cycle. *Hum. brain mapping* **40**, 538–551 (2019).

42. F Gobert, et al., Twenty-four-hour rhythmicities in disorders of consciousness are associated with a favourable outcome. *Commun. Biol.* **6**, 1213 (2023).

43. G Tononi, GM Edelman, Consciousness and complexity. *science* **282**, 1846–1851 (1998).

44. G Tononi, Consciousness as integrated information: a provisional manifesto. *The Biol. Bull.* **215**, 216–242 (2008).

45. JR King, et al., Information sharing in the brain indexes consciousness in noncommunicative patients. *Curr. biology* **23**, 1914–1919 (2013).

46. AI Luppi, et al., Consciousness-specific dynamic interactions of brain integration and functional diversity. *Nat. communications* **10**, 4616 (2019).

47. V Medel, M Irani, N Crossley, T Ossandón, G Boncompte, Complexity and 1/f slope jointly reflect brain states. *Sci. reports* **13**, 21700 (2023).

48. M Schartner, et al., Complexity of multi-dimensional spontaneous eeg decreases during propofol induced general anaesthesia. *PloS one* **10**, e0133532 (2015).

49. M Massimini, et al., Breakdown of cortical effective connectivity during sleep. *Science* **309**, 2228–2232 (2005).

50. S Sarasso, et al., Consciousness and complexity during unresponsiveness induced by propofol, xenon, and ketamine. *Curr. Biol.* **25**, 3099–3105 (2015).

51. P Coppola, et al., The complexity of the stream of consciousness. *Commun. biology* **5**, 1173 (2022).

52. M Massimini, et al., Hemispherotomy: cortical islands of deep sleep in awake humans (Research Square Preprint) (2024).

53. R Comolatti, et al., A fast and general method to empirically estimate the complexity of brain responses to transcranial and intracranial stimulations. *Brain stimulation* **12**, 1280–1289 (2019).

54. X Liu, BD Ward, JR Binder, SJ Li, AG Hudetz, Scale-free functional connectivity of the brain is maintained in anesthetized healthy participants but not in patients with unresponsive wakefulness syndrome. *PLoS One* **9**, e92182 (2014).

55. S Chen, et al., Disrupted interactions between arousal and cortical awareness networks in mcs and vs/uws patients: Evidence from resting-state functional imaging connectivity. *Neuroscience* **382**, 115–124 (2018).

56. A Markl, et al., Brain processing of pain in patients with unresponsive wakefulness syndrome. *Brain behavior* **3**, 95–103 (2013).

57. YG Pavlov, et al., Night sleep in patients with vegetative state. *J. Sleep Res.* **26**, 629–640 (2017).

58. V Cologan, et al., Sleep in the unresponsive wakefulness syndrome and minimally conscious state. *J. neurotrauma* **30**, 339–346 (2013).

59. U Malinowska, et al., Electroencephalographic profiles for differentiation of disorders of consciousness. *Biomed. engineering online* **12**, 1–18 (2013).

60. I Aricò, et al., Could combined sleep and pain evaluation be useful in the diagnosis of disorders of consciousness (doc)? preliminary findings. *Brain injury* **30**, 159–163 (2016).

61. M Wislowska, et al., Night and day variations of sleep in patients with disorders of consciousness. *Sci. reports* **7**, 266 (2017).

62. J Pan, J Wu, J Liu, J Wu, F Wang, A systematic review of sleep in patients with disorders of consciousness: from diagnosis to prognosis. *Brain sciences* **11**, 1072 (2021).

63. J Yu, et al., Sleep patterns correlates with the efficacy of tdcs on post-stroke patients with prolonged disorders of consciousness. *J. Transl. Medicine* **20**, 601 (2022).

64. JL Gottshall, DR Sebastiano, Sleep in disorders of consciousness: diagnostic, prognostic, and therapeutic considerations. *Curr. opinion neurology* **33**, 684–690 (2020).

65. MM Grigg-Damberger, O Hussein, T Kulik, Sleep spindles and k-complexes are favorable prognostic biomarkers in critically ill patients. *J. Clin. Neurophysiol.* **39**, 372–382 (2022).

66. D Arnaldi, et al., The prognostic value of sleep patterns in disorders of consciousness in the sub-acute phase. *Clin. Neurophysiol.* **127**, 1445–1451 (2016).

67. M Valente, et al., Sleep organization pattern as a prognostic marker at the subacute stage of post-traumatic coma. *Clin. Neurophysiol.* **113**, 1798–1805 (2002).

68. R Lehembre, et al., Resting-state eeg study of comatose patients: a connectivity and frequency analysis to find differences between vegetative and minimally conscious states. *Funct. neurology* **27**, 41 (2012).

69. S Chennu, et al., Spectral signatures of reorganised brain networks in disorders of consciousness. *PLoS computational biology* **10**, e1003887 (2014).

70. DR Sebastiano, et al., Significance of multiple neurophysiological measures in patients with chronic disorders of consciousness. *Clin. Neurophysiol.* **126**, 558–564 (2015).

71. A Naro, et al., Towards a method to differentiate chronic disorder of consciousness patients' awareness: The low-resolution brain electromagnetic tomography analysis. *J. Neurol. Sci.* **368**, 178–183 (2016).

72. RM Benca, et al., Eeg alpha power and alpha power asymmetry in sleep and wakefulness. *Psychophysiology* **36**, 430–436 (1999).

73. C Babiloni, et al., Resting state eyes-closed cortical rhythms in patients with locked-in-syndrome: an eeg study. *Clin. Neurophysiol.* **121**, 1816–1824 (2010).

74. Z Liu, et al., Finding thalamic bold correlates to posterior alpha eeg. *Neuroimage* **63**, 1060–1069 (2012).

75. MA Colombo, et al., Beyond alpha power: Eeg spatial and spectral gradients robustly stratify disorders of consciousness. *Cereb. cortex* **33**, 7193–7210 (2023).

76. J Pan, et al., Prognosis for patients with cognitive motor dissociation identified by brain-computer interface. *Brain* **143**, 1177–1189 (2020).

77. RB Berry, et al., The aasm manual for the scoring of sleep and associated events. *Rules, Terminol. Tech. Specif. Darien, Illinois, Am. Acad. Sleep Medicine* **176**, 7 (2012).

78. PB Forgacs, et al., Preservation of electroencephalographic organization in patients with impaired consciousness and imaging-based evidence of command-following. *Annals neurology* **76**, 869–879 (2014).

79. C Demanuele, CJ James, EJ Sonuga-Barke, Distinguishing low frequency oscillations within the 1/f spectral behaviour of electromagnetic brain signals. *Behav. Brain Funct.* **3**, 1–14 (2007).

80. T Inouye, et al., Quantification of eeg irregularity by use of the entropy of the power spectrum. *Electroencephalogr. clinical neurophysiology* **79**, 204–210 (1991).

81. A Lempel, J Ziv, On the complexity of finite sequences. *IEEE Transactions on information theory* **22**, 75–81 (1976).

82. F Kaspar, H Schuster, Easily calculable measure for the complexity of spatiotemporal patterns. *Phys. review A* **36**, 842 (1987).

83. R Nagarajan, Quantifying physiological data with lempel-ziv complexity-certain issues. *IEEE Transactions on Biomed. Eng.* **49**, 1371–1373 (2002).