**Terminal Loop Sequences in Viral Double-Stranded RNAs Modulate RIG-I Signaling**

Matthew Hackbart[1], Patrick Wang[1], Victoria Gnazzo[1], and Carolina B. López[1,2*]

[1]Department of Molecular Microbiology, Washington University School of Medicine, St. MO

[2]Center for Women Infectious Disease Research, Washington University School of Medicine, St. MO

*Corresponding Author, clopezzalaquett@wustl.edu

**ABSTRACT**

Detection of foreign RNAs is a crucial activation step for innate immunity pathways in response to viral infections. Retinoic acid-inducible gene I (RIG-I) is a cytoplasmic RNA sensor that triggers type I and III interferon (IFN) expression and activates the antiviral response. The activating ligand for RIG-I has been shown to be 5'-triphosphated blunt-ended double stranded(ds) RNA, but questions remain on the molecular mechanisms for RIG-I activation during viral infections. Here we show that immune-activating copy-back viral genomes (cbVGs) contain RNA stem loops away from the 5' end of the RNA that enhance RIG-I signaling and IFN expression. Importantly, the sequence of the terminal loops of the activating motifs impacts the strength of IFN expression. Additionally, we show that synthetic versions of these cbVG-derived stem loops trigger innate immune responses in mice demonstrating their potential as immunostimulants *in vivo*.

**INTRODUCTION**

Retinoic acid-inducible gene I (RIG-I) is an important cytoplasmic sensor of viral
infections that initiate the antiviral immune response. The RIG-I-like receptor (RLR) family
includes RIG-I, melanoma differentiation-associated protein 5 (MDA5), and laboratory of
genetics and physiology 2 (LGP2), which are host proteins that bind to RNAs and initiate
the antiviral response[1,2]. Binding of RLRs by foreign RNA triggers a signaling cascade
through the mitochondrial antiviral signaling (MAVS) protein that ends in the expression
of antiviral proteins, including type I and III interferons (IFNs)[3-7]. RIG-I contains a C-
terminal regulatory domain that recognizes the foreign RNA and drives conformational
changes in the protein helicase domain to wrap around the RNA[8-13]. These
conformational changes also release caspase activation and recruitment domains
(CARDs) that promote oligomerization of the RIG-I proteins and signaling through
MAVS[3,14,15]. RLRs recognize unique pathogen associated molecular patterns (PAMPs)
found in foreign RNA but not in the host cell RNAs.

The most well-studied of the RLR PAMPs are within the RNAs that activate RIG-I.
The regulatory domain of RIG-I detects terminal 5' triphosphates on RNA[16-19] and the
detection of this motif can be inhibited by methylation of the 5' end of the RNA[20,21]. While
ssRNA can activate RIG-I, the highest activation occurs when blunt ended dsRNA[22]
contains the 5' triphosphate[17]. The addition of overhangs to the end of the dsRNA limits
activation of RIG-I[18,19]. RIG-I is activated best by short dsRNA while longer synthetic
dsRNAs such as poly I:C (> 500nt in length) activate MDA5[23-25]. With these guidelines in
place, minimal ligands for activating RIG-I that contain a short 10 base pair dsRNA stem
with a 5' triphosphate have been synthetically generated[26,27]. While these studies have
provided critical data to identify the motifs that RIG-I will bind to in the RNA, in general,

they are not based on RNAs found naturally during viral infections. Further work is thus

needed to confirm and identify the characteristics of viral RNAs that activate RIG-I

signaling during infection.

Previously, we and others determined that the RNA that most robustly activates

RIG-I signaling during Sendai Virus (SeV) infection is a copy-back viral genome (cbVG)[28-

30]. CbVGs contain complementary ends predicted to form a blunt-ended dsRNA PAMP

with a 5' triphosphate, characteristic of the canonical RIG-I ligand[31,32]. However, detailed

folding analyses of the most prominent cbVG from SeV (cbVG 546) revealed a much

more complex folding of the molecule, including a stem loop that is necessary for strong

activation of the RIG-I pathway to stimulate IFN expression. This stem loop is comprised

of nucleotides 70-114 of cbVG 546, and when transferred to inert RNAs, it significantly

enhanced the immunostimulatory capability of the RNA[33]. Interestingly, the removal of

stem loop 70-114 from cbVG 546 severely decreased RIG-I activation even though the

cbVG still contained the 5' triphosphate and blunt-end dsRNA.

It is unknown how SeV 70-114 enhanced RIG-I activity or whether other RIG-I

stimulatory cbVGs contain stem loops similar to SeV 70-114. We therefore sought to

determine if other viral cbVGs contain RNA stem loops that can transfer RIG-I stimulatory

activity to otherwise inert RNAs and identify features of these RNA stem loops that are

necessary for robust RIG-I activation. Here, we identified multiple sequences in cbVGs

from human respiratory syncytial virus (RSV) and Nipah virus (NiV) infections that can

transfer RIG-I stimulatory activity to the inert X RNA from the hepatitis C virus. Through

mutation of the terminal loop sequences of the cbVG-derived RNAs, we discovered that

specific nucleotides in the terminal loop of the RNA are necessary for strong activation of

RIG-I signaling in cells. Lastly, *in vivo* administration of the cbVG-derived RNA loops

activated innate immune signaling pathways, suggesting their potential use as

immunostimulants for vaccines and immunotherapies.

**RESULTS**

**RSV and NiV cbVGs stimulate interferon expression**

To identify additional natural RNA stem loops that could activate antiviral innate

immune pathways, we selected previously identified cbVGs from either RSV (RSV cbVG

236)[34] and NiV (DI15, here called NiV cbVG 378)[35]. We first determined if these cbVGs

can activate the IFN signaling pathway in human A549 lung cells and found that *in vitro*

transcribed RNAs of both RSV cbVG 236 and NiV cbVG 378 could stimulate the

transcription of type I (*IFNB1*) and type III (*IFNL1)* interferons at 6 hours post transfection,

but not the hepatitis C virus X region (Fig 1A and 1B). This IFN activation was largely

MAVS-dependent, confirming activation of the RLR signaling, similar to what we had

reported for SeV cbVG 268, a shortened version of cbVG 546 (Fig 1C)[33].

**RSV cbVGs contain immunostimulatory stem loops**

To identify which specific RSV cbVG 236 sequence confers a strong RIG-I

stimulatory activity to the molecule, we first made a series of deletions in cbVG 236 that

either removed internal sequences disrupting the stem loops or removed a

complementary end of the cbVG (Fig 2A). We found that nucleotides 1-48 and 189-236

in the complementary ends of cbVG 236 contribute to RIG-I activation, as deletion

mutants del 1-48 and 189-236 lost immunostimulatory activity (Fig 2B). Interestingly, we

also found that nucleotides 49-100 are necessary for maximum IFN expression. Mutants

with deletions in this range (del 49-187 and del 64-187) show reduced stimulation of the

IFN response, but del 101-147 and del 137-187 induced strong IFN expression.

To determine if these RNA sequences can enhance expression of IFNs to an

otherwise inert RNA molecule, we transferred the RSV cbVG sequences to the inert X

RNA motif from the hepatitis C virus, ensuring that the overall structures of the final

molecules were similar (Fig S2)[36]. We found that X-region chimeras carrying RSV cbVG

sequences 1-48 or 39-68 induced the highest IFN expression when transfected into A549

cells, but RSV cbVG sequence 189-236 did not significantly enhance IFN expression in

this context (Fig 2C), suggesting that additional features of the 189-236 region participate

in inducing IFN expression.

To determine if specific immunostimulatory stem loops were present within the

immunostimulatory sequences, we then performed *in silico* prediction of the RNA stem

loops using RNAfold[37]. Since the complement ends of the cbVGs interfere with the *in*

*silico* folding predictions based on minimal free energy, and it is likely that favorable

secondary structures form during RNA synthesis rather than once the entire molecule has

been produced, we determined all possible RNA structures on partial cbVGs lacking

either the 3' or 5' complementary sequences. RSV cbVG 236 was predicted to form five

RNA stem loops (Fig 2D and Fig S1) in the absence of 3'-5' complementation with two of

the stem loops (1-34 and 39-68) present within the immunostimulatory sequences. RNA

folding of X region + RSV 1-48 predicted an immunostimulatory stem loop of sequence

1-34 similar to the full length cbVG (Fig S2).

**NiV cbVGs contain immunostimulatory stem loops**

Similar to RSV, we made a series of deletions to remove specific RNA sequences

from NiV cbVG 378 and determined which portions of the cbVG are necessary for strong

IFN expression. We found that nucleotides within the complementary ends of the cbVG

were necessary for strong expression of IFNs and removal of nucleotides 325-360 also

decreased IFN expression (Fig 3A and B). When attached to the X RNA, we found that

while NiV 325-360 induced *IFNB1* and *IFNL1* expression, NiV sequences 44-78 and 85-

136 induced greater IFN expression (Fig 3C). NiV cbVG 378 is predicted to have eight

RNA stem loops (Fig 3D and Fig S3) with stem loops of NiV 44-78 and NiV 85-136

enhancing IFN stimulation when transferred to the inert X RNA. Unexpectedly, deletion

of 44-78 or 85-136 sequences did not significantly reduce *IFNL1* expression from upon

RNA transfection, suggesting that the RNA folds differently in these regions of the

complementary ends than what was predicted *in silico*.

**Terminal loop sequences are critical for IFN expression**

We   next   investigated   the   characteristics   conserved   among   high

immunostimulatory RNA sequences versus lower immunostimulatory sequences (Fig

4A). The cbVG RNAs are predicted to form stem loops that are diverse in their size and

shape. However, two complementary sequences that are predicted to form loops with the

same structure (RSV 1-34 and RSV 203-236) exhibit different immunostimulatory

capabilities (Fig 2C). We did not observe significant differences in the minimum free

energy (MFE) between the stem loops, most of which had a predicted MFE between -6

and -10 kcal/mol. Upon observation, we found that the terminal loops with the higher

immunostimulatory activity contain either a CAA motif or poly-A stretch at the tip of the

loop (Fig 4A). In contrast, the low immunostimulatory loops are instead U-rich at the

terminal loop residues, except for NiV 324-361 which contains an A-rich terminal loop.

We then hypothesized that the terminal loop adenines played a key role in

triggering RIG-I signaling. To test this hypothesis, we mutated the terminal loops of the

X-RNA attached sequences and tested their ability to induce IFN expression. Mutation of

the terminal loop sequence did not alter the overall structure of the stem loops (Fig S4-

7). For X + SeV 70-114, we found that mutation of the A89 or A90 terminal residues

greatly reduced IFN activation by the RNA construct, however mutation of the C88

residue did not (Fig 4B). Mutations in X + RSV 1-48 showed that A18, but not C17 or A19,

was necessary for full IFN activation (Fig 4C). Additionally, mutation of X + NiV 85-136

showed that both terminal adenines, A111 and A112, were necessary for IFN signaling

(Fig 4D). However, this trend was not observed for the low immunostimulatory X + NiV

324-361, which did not show a reduction in IFN stimulation upon mutation of the terminal

adenines (Fig 3E). These data suggest that terminal adenines are critical to trigger robust

IFN expression and that additional sequence-specific factors impact their stimulatory

activity.

**RSV 1-48 activates innate immune signaling *in vivo***

Previously, we found that the SeV cbVG 546 stem loop 70-114 activates a strong

type I immune responses in mice when administered alone or within a lipid nanoparticle

(LNP)[38-40]. Here, we aimed to determine whether RSV sequence 1-48 similarly activates

innate sensors and induces an immune response *in vivo*. To ensure the RNA constructs

were delivered intracellularly to trigger the RIG-I pathway, we packaged the X + RSV 1-

48 and the control X + RSV 189-236 chimera constructs into LNPs (Fig 5A) and confirmed

the homogeneity of the formulations before *in vivo* administration (Fig 5B). These

formulations were then administered subcutaneously in the footpad of C57BL/6 mice.

After 16 hours, we found that LNPs containing X + RSV 1-48 induced significantly higher

expression of *IFNB1*, *MX1*, *IL1B*, and *CXCL10* compared to empty LNPs and X + RSV

189-236 (Fig 5C). These data demonstrate that the RSV cbVG-derived 1-48 sequence

retains its immunostimulatory activity *in vivo* when packaged into LNPs and together with

data from SeV 70-114 demonstrate that small RNA motifs with terminal loop adenines

can serve as potent immunostimulators in mice.

**DISCUSSION**

We have identified multiple RNA stem loops derived from cbVGs of RSV and NiV

that can transfer robust IFN stimulatory capabilities to an inert RNA. One unique feature

of the higher immunostimulatory RNA stem loops is a richness for adenines at the tip of

the loop. When these sequences were mutated, the mutant RNAs stimulated significantly

less IFN upon transfection into cells. Upon packaging into lipid nanoparticles, the RSV

cbVG-derived stem loop was able to activate innate immune signaling in mice.

While the enhanced stimulation appears to occur with adenine-rich sequences, we

also observe that a stem loop with 5 adenines (NiV 325-360) did not have decreased IFN

activation when the adenines were mutated (Fig. 4E). Since an adenine-rich loop does

not always enhance IFN expression, more work is necessary to determine the exact motif

or motifs that make up an optimal terminal loop PAMP. Some possibilities are that

sequences in the stems of the RNA interact with the loop to alter binding or that host

proteins are recognizing a specific RNA motif that is more than adenines.

One major question that remains is what step of the RIG-I pathway is impacted by

these terminal adenine-rich sequence. We propose the following possibilities for how the

terminal loop sequences may alter IFN expression: 1) RIG-I is heavily modulated by

phosphorylation and ubiquitination by host proteins[41], so the terminal loop may recruit or

inhibit additional host factors from interacting and altering RIG-I function in cells. 2) MAVS

has recently been shown to interact with host RNAs[42]. The terminal sequence may

stabilize RIG-I binding to MAVS through RNA/MAVS interactions and enhance

downstream signaling to boost IFN expression. 3) The terminal loop adenines may be

prone to RNA modifications such as N6 methylation to generate N6 methyladenosine

(m6A)[43]. m6A RNA modification can alter the stability of the RNA in a cell and modulate

the binding of RNA binding proteins. 4) Other unknown host RNA-binding proteins support

robust RIG-I signaling. Further work is necessary to parse out what interactions occur

with this terminal loop sequence and why adenine-rich sequences enhance IFN

expression from the RIG-I pathway.

With the advent of mRNA vaccines and the usage of synthetic RNAs as

therapeutics, understanding how RNAs can activate innate immunity is crucial to

optimizing RNA-based adjuvants and treatments. Synthetic RNAs can act as immune

adjuvants during vaccination[27,38-40,44,45], and the addition of immunostimulatory stem

loops to mRNA vaccines may bypass the need for additional adjuvants such as alum.

Although additional work is necessary to assess the immunostimulatory properties of

RNA stem loops *in vivo*, our discovery that the terminal loop sequence modulates immune

activation can lead to the development of strategies to tailor the immune response not

only based on dosage of stimulatory RNA, but on the sequence of attached RNA stem

loops.

**MATERIALS AND METHODS**

**Cells and Reagents.** A549 human type II alveolar cells (ATCC, CCL-185) and MAVS-

KO A549 cells (gifted from S. Weiss) were cultured at 37°C and 5% $CO_2$ with Dulbecco's

modified Eagle's media (Thermofisher, 11995065) supplemented with 10% fetal bovine

serum (FBS), 1 mM sodium pyruvate, 2 mM L-Glutamine, and 50 mg/mL gentamicin. All

cell lines were treated with mycoplasma removal agent (MP Biomedicals, 093050044)

and routinely tested for mycoplasma before use.

*In Vitro* **Transcription of RNA.** RNAs were cloned into a pSL1180-T7 plasmid that

contains T7 promoter at the 5' end of the RNA sequence and hepatitis D virus ribozyme

after the RNA sequence. *In vitro* transcription was performed with the Megascript T7

Transcription Kit (Thermofisher, AM1334) and RNA was isolated with LiCl precipitation

according to manufacturer's protocols. Purified RNA was measured by Qubit and tested

for quality by Bioanalyzer (Agilent). Full protocol available Protocols.io:

dx.doi.org/10.17504/protocols.io.3byl49qprgo5/v2

**RT-PCR and qPCR.** A549 cells were transfected with 5 pmol of indicated RNA and

Lipofectamine 2000 (Thermofisher, 11668027). After 6 hours, RNA was isolated with

Trizol (Thermofisher, 15596026). cDNA was generated with random hexamers using the

High-Capacity RNA to cDNA kit (Thermofisher, 4387406) according to the manufacturer's

protocols. For qPCR quantification, cDNA was quantified with Power SYBR Green Mix

(Thermofisher, 4367660). *IFNL1, IFNB1, IL1B, MX1, and CXCL10* were normalized to a

house keeping index calculated from *ACTB* and *GAPDH* genes. Sequences of primers

are given in Table S1. Full protocol available at Protocols.io:

dx.doi.org/10.17504/protocols.io.8epv5rxw6g1b/v1

**Mice**

C57BL/6 mice (The Jackson Laboratory) were bred in house. All mice were sex

and age matched. Both male and female mice were included in the experiments.

**Lipid Nanoparticle formulation**

X + RSV 1-48 and X + RSV 189-236 were encapsulated in the GenVoy Ionizable

Lipid Mix (ILM) using a NanoAssemblr Ignite machine (Precision Nanosystems) following

the manufacturer's instructions. Empty LNPs were performed and used as controls.

Encapsulation efficiency and RNA concentration were tested by Ribogreen Assay

(ThermoFisher). Final particle size was measure by Dynamic Light Scattering. Full

protocol available at Protocols.io: dx.doi.org/10.17504/protocols.io.36wgqnn4kgk5/v1

**Mice immunization**

Mice were anesthetized with isoflurane and injected subcutaneously (s.c.) into a

rear footpad. For immune response studies mice were inoculated with 0.5 µg X 1-48; 0.5

µg X 189-236; Empty LNPs or PBS at a final volume of 30 µl per dose. RNA was extracted

from footpad 16 hours after inoculation using TRIzol (Ambion Inc.).

***In Silico* RNA Folding Prediction.** All RNA folding predictions were performed using the

University of Vienna RNAfold website with the ViennaRNA Package (Version 2.6.3).

Structures shown are the minimum free energy structures. All sequences used for folding

are given in Table S2.

Website: http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi

**Statistical Analysis and Reproducibility**

All statistics were calculated using GraphPad Prism. Version 9. Specific tests and

significance values are indicated in each figure legend. Data reported are biological

triplicates unless otherwise stated.

**Ethics statement**

All described studies adhered to the Guide for the Care and Use of Laboratory

Animal of the National Institute of Health. Institutional Animal Care and Use Committee,

Washington University in St. Louis approved protocol 23-0083.

**ACKNOWLEDGEMENTS**

The authors acknowledge the use of the Chemical and Environmental Analysis Facility at

Washington University in Saint Louis for instrumentation and analytical support.

**Funding:**

NIH T32 HL007317-44 (MH)

NIH T32 AI007163-45 (MH)

NIH A137062 (CBL)

WashU BJC Investigator Program (CBL)

## 282 BIBLIOGRAPHY

1 Kato, H., Takahasi, K. & Fujita, T. RIG-I-like receptors: cytoplasmic sensors for non-self
RNA. *Immunol Rev* **243**, 91-98 (2011). https://doi.org:10.1111/j.1600-065X.2011.01052.x

2 Thoresen, D. *et al.* The molecular mechanism of RIG-I activation and signaling. *Immunol*
*Rev* **304**, 154-168 (2021). https://doi.org:10.1111/imr.13022

3 Yoneyama, M. *et al.* The RNA helicase RIG-I has an essential function in double-stranded
RNA-induced innate antiviral responses. *Nat Immunol* **5**, 730-737 (2004).
https://doi.org:10.1038/ni1087

4 Tanner, N. K. & Linder, P. DExD/H box RNA helicases: from generic motors to specific
dissociation functions. *Mol Cell* **8**, 251-262 (2001). https://doi.org:10.1016/s1097-
2765(01)00329-x

5 Kawai, T. *et al.* IPS-1, an adaptor triggering RIG-I- and Mda5-mediated type I interferon
induction. *Nat Immunol* **6**, 981-988 (2005). https://doi.org:10.1038/ni1243

6 Seth, R. B., Sun, L., Ea, C. K. & Chen, Z. J. Identification and characterization of MAVS, a
mitochondrial antiviral signaling protein that activates NF-kappaB and IRF 3. *Cell* **122**, 669-
682 (2005). https://doi.org:10.1016/j.cell.2005.08.012

7 Xu, L. G. *et al.* VISA is an adapter protein required for virus-triggered IFN-beta signaling.
*Mol Cell* **19**, 727-740 (2005). https://doi.org:10.1016/j.molcel.2005.08.014

8 Cui, S. *et al.* The C-terminal regulatory domain is the RNA 5'-triphosphate sensor of RIG-I.
*Mol Cell* **29**, 169-179 (2008). https://doi.org:10.1016/j.molcel.2007.10.032

9 Kowalinski, E. *et al.* Structural basis for the activation of innate immune pattern-
recognition receptor RIG-I by viral RNA. *Cell* **147**, 423-435 (2011).
https://doi.org:10.1016/j.cell.2011.09.039

10 Wang, Y. *et al.* Structural and functional insights into 5'-ppp RNA pattern recognition by
the innate immune receptor RIG-I. *Nat Struct Mol Biol* **17**, 781-787 (2010).
https://doi.org:10.1038/nsmb.1863

11 Lu, C., Ranjith-Kumar, C. T., Hao, L., Kao, C. C. & Li, P. Crystal structure of RIG-I C-terminal
domain bound to blunt-ended double-strand RNA without 5' triphosphate. *Nucleic Acids*
*Res* **39**, 1565-1575 (2011). https://doi.org:10.1093/nar/gkq974

12 Lu, C. *et al.* The structural basis of 5' triphosphate double-stranded RNA recognition by
RIG-I C-terminal domain. *Structure* **18**, 1032-1043 (2010).
https://doi.org:10.1016/j.str.2010.05.007

13 Luo, D. *et al.* Structural insights into RNA recognition by RIG-I. *Cell* **147**, 409-422 (2011).
https://doi.org:10.1016/j.cell.2011.09.023

14  Peisley, A., Wu, B., Yao, H., Walz, T. & Hur, S. RIG-I forms signaling-competent filaments
in an ATP-dependent, ubiquitin-independent manner. *Mol Cell* **51**, 573-583 (2013).
https://doi.org:10.1016/j.molcel.2013.07.024
15  Ramanathan, A. *et al.* The autoinhibitory CARD2-Hel2i Interface of RIG-I governs RNA
selection. *Nucleic Acids Res* **44**, 896-909 (2016). https://doi.org:10.1093/nar/gkv1299
16  Kim, D. H. *et al.* Interferon induction by siRNAs and ssRNAs synthesized by phage
polymerase. *Nat Biotechnol* **22**, 321-325 (2004). https://doi.org:10.1038/nbt940
17  Pichlmair, A. *et al.* RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-
phosphates. *Science* **314**, 997-1001 (2006). https://doi.org:10.1126/science.1132998
18  Schlee, M. *et al.* Recognition of 5' triphosphate by RIG-I helicase requires short blunt
double-stranded RNA as contained in panhandle of negative-strand virus. *Immunity* **31**,
25-34 (2009). https://doi.org:10.1016/j.immuni.2009.05.008
19  Schmidt, A. *et al.* 5'-triphosphate RNA requires base-paired structures to activate antiviral
signaling via RIG-I. *Proc Natl Acad Sci U S A* **106**, 12067-12072 (2009).
https://doi.org:10.1073/pnas.0900971106
20  Devarkar, S. C. *et al.* Structural basis for m7G recognition and 2'-O-methyl discrimination
in capped RNAs by the innate immune receptor RIG-I. *Proc Natl Acad Sci U S A* **113**, 596-
601 (2016). https://doi.org:10.1073/pnas.1515152113
21  Schuberth-Wagner, C. *et al.* A Conserved Histidine in the RNA Sensor RIG-I Controls
Immune Tolerance to N1-2'O-Methylated Self RNA. *Immunity* **43**, 41-51 (2015).
https://doi.org:10.1016/j.immuni.2015.06.015
22  Marques, J. T. *et al.* A structural basis for discriminating between self and nonself double-
stranded RNAs in mammalian cells. *Nat Biotechnol* **24**, 559-565 (2006).
https://doi.org:10.1038/nbt1205
23  Kato, H. *et al.* Length-dependent recognition of double-stranded ribonucleic acids by
retinoic acid-inducible gene-I and melanoma differentiation-associated gene 5. *J Exp Med*
**205**, 1601-1610 (2008). https://doi.org:10.1084/jem.20080091
24  Kato, H. *et al.* Differential roles of MDA5 and RIG-I helicases in the recognition of RNA
viruses. *Nature* **441**, 101-105 (2006). https://doi.org:10.1038/nature04734
25  Binder, M. *et al.* Molecular mechanism of signal perception and integration by the innate
immune sensor retinoic acid-inducible gene-I (RIG-I). *J Biol Chem* **286**, 27278-27287
(2011). https://doi.org:10.1074/jbc.M111.256974
26  Kohlway, A., Luo, D., Rawling, D. C., Ding, S. C. & Pyle, A. M. Defining the functional
determinants for RNA surveillance by RIG-I. *EMBO Rep* **14**, 772-779 (2013).
https://doi.org:10.1038/embor.2013.108
27  Linehan, M. M. *et al.* A minimal RNA ligand for potent RIG-I activation in living mice. *Sci
Adv* **4**, e1701854 (2018). https://doi.org:10.1126/sciadv.1701854
28  Baum, A., Sachidanandam, R. & García-Sastre, A. Preference of RIG-I for short viral RNA
molecules in infected cells revealed by next-generation sequencing. *Proc Natl Acad Sci U
S A* **107**, 16303-16308 (2010). https://doi.org:10.1073/pnas.1005077107
29  Mercado-López, X. *et al.* Highly immunostimulatory RNA derived from a Sendai virus
defective viral genome. *Vaccine* **31**, 5713-5721 (2013).
https://doi.org:10.1016/j.vaccine.2013.09.040

30    Sun, Y. *et al.* Immunostimulatory Defective Viral Genomes from Respiratory Syncytial
Virus Promote a Strong Innate Antiviral Response during Infection in Mice and Humans.
*PLoS Pathog* **11**, e1005122 (2015). https://doi.org:10.1371/journal.ppat.1005122
31    Kolakofsky, D. Isolation and characterization of Sendai virus DI-RNAs. *Cell* **8**, 547-555
(1976). https://doi.org:10.1016/0092-8674(76)90223-3
32    Lazzarini, R. A., Keene, J. D. & Schubert, M. The origins of defective interfering particles of
the negative-strand RNA viruses. *Cell* **26**, 145-154 (1981). https://doi.org:10.1016/0092-
8674(81)90298-1
33    Xu, J. *et al.* Identification of a Natural Viral RNA Motif That Optimizes Sensing of Viral RNA
by RIG-I. *mBio* **6**, e01265-01215 (2015). https://doi.org:10.1128/mBio.01265-15
34    Sun, Y. *et al.* A specific sequence in the genome of respiratory syncytial virus regulates the
generation of copy-back defective viral genomes. *PLoS Pathog* **15**, e1007707 (2019).
https://doi.org:10.1371/journal.ppat.1007707
35    Welch, S. R. *et al.* Inhibition of Nipah Virus by Defective Interfering Particles. *J Infect Dis*
**221**, S460-s470 (2020). https://doi.org:10.1093/infdis/jiz564
36    Stone, A. E. *et al.* Hepatitis C virus pathogen associated molecular pattern (PAMP) triggers
production of lambda-interferons by human plasmacytoid dendritic cells. *PLoS Pathog* **9**,
e1003316 (2013). https://doi.org:10.1371/journal.ppat.1003316
37    Zuker, M. & Stiegler, P. Optimal computer folding of large RNA sequences using
thermodynamics and auxiliary information. *Nucleic Acids Res* **9**, 133-148 (1981).
https://doi.org:10.1093/nar/9.1.133
38    Fisher, D. G., Gnazzo, V., Holthausen, D. J. & López, C. B. Non-standard viral genome-
derived RNA activates TLR3 and type I IFN signaling to induce cDC1-dependent CD8+ T-
cell responses during vaccination in mice. *Vaccine* **40**, 7270-7279 (2022).
https://doi.org:10.1016/j.vaccine.2022.10.052
39    Gnazzo, V. *et al.* DDO-adjuvanted influenza A virus nucleoprotein mRNA vaccine induces
robust humoral and cellular type 1 immune responses and protects mice from challenge.
*mBio*, e0358924 (2024). https://doi.org:10.1128/mbio.03589-24
40    Fisher, D. G., Coppock, G. M. & López, C. B. Virus-derived immunostimulatory RNA induces
type I IFN-dependent antibodies and T-cell responses during vaccination. *Vaccine* **36**,
4039-4045 (2018). https://doi.org:10.1016/j.vaccine.2018.05.100
41    Rehwinkel, J. & Gack, M. U. RIG-I-like receptors: their regulation and roles in RNA sensing.
*Nat Rev Immunol* **20**, 537-551 (2020). https://doi.org:10.1038/s41577-020-0288-3
42    Gokhale, N. S. *et al.* Cellular RNA interacts with MAVS to promote antiviral signaling.
*Science* **386**, eadl0429 (2024). https://doi.org:10.1126/science.adl0429
43    Boulias, K. & Greer, E. L. Biological roles of adenine methylation in RNA. *Nat Rev Genet*
**24**, 143-160 (2023). https://doi.org:10.1038/s41576-022-00534-0
44    Jiang, X. *et al.* Intratumoral delivery of RIG-I agonist SLR14 induces robust antitumor
responses. *J Exp Med* **216**, 2854-2868 (2019). https://doi.org:10.1084/jem.20190801
45    Mao, T. *et al.* A stem-loop RNA RIG-I agonist protects against acute and chronic SARS-CoV-
2 infection in mice. *J Exp Med* **219** (2022). https://doi.org:10.1084/jem.20211818
400
401

**FIGURE LEGENDS**

Figure 1. CbVG IFN Activation

**Figure 1. CbVG from RSV and NiV induce MAVS-dependent IFN expression. (A+B)**

Quantitative PCR (qPCR) for (A) type I interferon (*IFNB1*) and (B) type III interferon

(*IFNL1*) expression by A549 cells at 6 hours post transfection (hpt) with 5 pmol of HCV X

RNA, SeV cbVG 268, RSV cbVG 236, or NiV cbVG 378. One-way ANOVA was performed

for statistical analysis. ns: p>0.05; *: p<0.05, ***: p<0.001; ****: p<0.0001. Data are

represented as fold change over Mock. (C) qPCR for *IFNL1* of control or MAVS-KO A549

cells at 6 hpt with 5 pmol of SeV cbVG 268, RSV cbVG 236, or NiV cbVG 378. One-way

ANOVA was performed for statistical analysis. ****: p<0.0001. Data are represented as

fold change over Mock. Data reported are biological triplicates.

**Figure 2. RSV cbVG sequences 1-48 and 39-68 are immunostimulatory.** (A) Indicated

deletions of RSV cbVG 236 were generated by *in vitro* transcription then 1 pmol of IVT

RNA was transfected into A549 cells. (B) Expression of *IFNL1* was measured by qPCR

at 6 hpt. One-way ANOVA was performed for statistical analysis. ns: $p > 0.05$; ***: $p < 0.001$,

****: $p < 0.0001$. Data are represented as fold change over cbVG 236. (C) Sequences from

RSV cbVG 236 were attached to X RNA then *in vitro* transcribed. 1 pmol of RNA was

transfected into A549 cells and expression of *IFNB1* and *IFNL1* was measured by qPCR

at 6 hpt. One-way ANOVA was performed for statistical analysis. ns: $p > 0.05$; *: $p < 0.05$,

****: $p < 0.0001$. Data are represented as fold change over the X RNA control. (D)

Predicted structure of RNA stem loops from RSV cbVG 236. Data reported are biological

triplicates.

**Figure 3. NiV cbVG sequences 44-78 and 85-136 are immunostimulatory.** (A)

Indicated deletions of NiV cbVG 378 were generated by *in vitro* transcription then 1 pmol

of IVT RNA was transfected into A549 cells. (B) Expression of *IFNL1* was measured by

qPCR at 6 hpt. One-way ANOVA was performed for statistical analysis. ns: $p > 0.05$; ***:

$p < 0.001$, ****: $p < 0.0001$. Data are represented as fold change over cbVG 378. (C)

Sequences from NiV cbVG 378 were attached to X RNA then *in vitro* transcribed. 1 pmol

of IVT RNA was transfected into A549 cells and expression of *IFNB1* and *IFNL1* was

measured by qPCR at 6 hpt.. One-way ANOVA was performed for statistical analysis. ns:

$p > 0.05$; *: $p < 0.05$, **: $p < 0.01$; ***: $P < 0.001$; ****: $p < 0.0001$. Data are represented as fold

change over the X RNA control. (D) Predicted structure of RNA stem loops from NiV cbVG

378. Data reported are biological triplicates.

**Figure 4. Terminal loop adenines are necessary for high IFN expression.** (A) Predicted RNA structures of high immunostimulatory and low immunostimulatory stem loops derived from cbVGs. (B-E) Indicated mutations were generated for cbVG sequences attached to the X RNA. The following sequences were mutated: (B) SeV 70-

114, (C) RSV 1-48, (D) NiV 85-136, and (E) NiV 324-361. RNAs were *in vitro* transcribed

and 1 pmol of IVT RNA was transfected into A549 cells. Expression of *IFNL1* was

measured by qPCR at 6 hpt. One-way ANOVAs were performed for statistical analysis.

ns: $p > 0.05$; *: $p < 0.05$, **: $p < 0.01$; ***: $P < 0.001$; ****: $p < 0.0001$. Data are represented as

fold change over the wild-type stem loops. Data reported are biological triplicates.

**Figure 5. X + RSV 1-48 activates innate immune signaling in mice. (**A) Schematic of

lipid nanoparticles (LNPs) containing X + RSV 1-48 (X 1-48) or X + RSV 189-236 (X 189-

236). (B) Dynamic light scattering measurement of LNP sizes of empty LNPs, X 1-48, and

X 189-236. (C) LNPs were injected subcutaneously into the footpad of mice. Sixteen

hours post injection, RNA was collected and the RNA expression of *IFNB1*, *MX1*,

*CXCL10*, and *IL1B* was quantified by qPCR. One-way ANOVAs were performed for

statistical analysis. ns: $p > 0.05$; **: $p < 0.01$; ***: $P < 0.001$; ****: $p < 0.0001$. Data reported are

5 individual mice.