Retrograde Optogenetics Reveals Functional Convergence within a Corticotectal Pathway of Non-Human Primates

Xuefei Yu[1+], Atul Gopal[1+], Ken-ichi Inoue[2], Martin Bohlen[3], Genevieve Kuczewski[3], Marc A. Sommer[3], Hendrikje Nienborg[1], Masahiko Takada[2], Okihide Hikosaka[1]

1. Laboratory of Sensorimotor Research, National Eye Institute, National Institutes of Health, Bethesda, Maryland 20892, United States of America

2. Center for the Evolutionary Origins of Human Behavior, Kyoto University, Inuyama, Aichi 484-8506, Japan

3. Department of Biomedical Engineering, Duke University, Durham, North Carolina, 27708, United States of America

+ Corresponding author

Abstract

Retrograde optogenetics enables tagging and manipulation of pathway-defined neurons, although its application to the non-human primate brain remains challenging. We applied the technique to identify, record, and activate neurons forming the frontal eye field – superior colliculus projection in behaving macaques. Optical stimulation evoked robust contralateral saccades and pathway-selective opto-tagging revealed the functional visuo-motor convergences within this pathway. Our findings resolve a longstanding controversy about the frontal oculomotor outputs and highlight retrograde optogenetics as a powerful tool for primate circuit-level studies.

Main

Complex and flexible cognitive behaviors critically depend on the efficient flow of information across distributed neural circuits[1,2]. While system neuroscience has made significant advances in understanding information processing within individual brain regions, the mechanism by which information is transmitted between areas — particularly through specific anatomical pathways —remains poorly understood. One powerful approach to address this challenge is retrograde optogenetics, which enables selective labeling of neurons based on their projection targets and allows precise manipulation of activity to assess their causal contributions to behavior[3-6]. This method employs viral vectors transported retrogradely from axon terminals to somata, thereby allowing selective stimulation of pathway-defined neurons via light[6]. Although successfully applied in rodents, the use of retrograde optogenetics for projection-specific recording and behavioral manipulation in non-human primates has remained technically challenging and largely unexplored[7-9].

In the present study, we evaluated the efficacy of retrograde optogenetics for neural identification and behavioral manipulation in non-human primates by tagging the oculomotor pathway from the frontal eye field (FEF) to the superior colliculus (SC), two key nodes essential for saccade generation[10,11], and use this comprehensive approach to resolve a longstanding question about the nature of the FEF's output to the SC. The FEF-SC pathway offers an ideal

testbed for retrograde optogenetics for several reasons. First, both regions are known to integrate
diverse sensory inputs and play a causal role in saccade execution[12-14]. This provides a strong
foundation for behaviorally relevant pathway manipulation. Second, the nature of the FEF's
output to the SC in sensorimotor transformation in saccade generation remains unsolved. While
early studies suggested a motor-dominant signal in the FEF-SC pathway[15], subsequent research
indicated a richer signal composition that encompasses visual, visuomotor, and motor signals[16,17].
Third, although previous studies indicated that SC neurons receive FEF input[18], how these
signals are transformed between the regions is still unclear. To date, no single study has causally
tested the role of the FEF-SC projection in saccade behavior, directly characterized the
functional content of signals in the projection, and quantified the impact of the projection on SC-
recipient neurons, all within the same animals. Addressing all these questions together would
offer critical insight into the circuit-level and behavioral roles of this pathway.
59
To achieve such a cohesive study, we injected AAV2retro-hSyn-ChR2-EYFP unilaterally into
the SC of two monkeys (Fig.1a). To confirm gene expression in the FEF-SC projection, we
successfully manipulated neural activity and saccadic behavior using fiber optic illumination of
the SC (Extended Data Fig.1a-h) and showed post-mortem expression of the ChR2-EYFP
complex via histological analysis of the SC (Fig.2a) and FEF. Expression was profuse in the
ipsilateral FEF (Fig.2b) but negligible in the contralateral FEF (Fig.2c).
66
Our first main experimental goal was to optically stimulate the FEF to establish that selective
activation of retrogradely labeled FEF-to-SC neurons could evoke saccades. To test this, the
monkeys freely viewed a movie (Fig.1b), while 0.1s optical stimulation pulses (470nm,
50mW/mm^2) were delivered intermittently to the FEF (Fig.1b,c). Saccades during the 0.1 sec
stimulation period were compared between randomly interleaved "Stim Trials" (Fig.1c, top
center) and non-stimulated "Control Trials" (Fig.1c, bottom center) for each session.
73
Two key observations emerged. First, across 110 sessions, saccade frequency significantly
increased during the optical stimulation period (4.24 saccades/s, N = 2443 saccades) relative to
the control period (2.41 saccades/s, N = 1408 saccades, p = 4.28e-11, two-tailed paired t-test).
Second, evoked saccades were consistently directed toward the left visual field (Fig.1d shows
saccades for an example session). Since optical stimulation was applied to the right FEF, these
leftward saccades were contralateral to the site of stimulation (see also Extended Data Fig.2a-c),
resembling the effects of electrical stimulation at the same sites (Extended Data Fig.2b-d). The
onset latency of these example optically evoked saccades from FEF was 35ms post-stimulation
(Fig.1e), about 12ms longer than the latency observed from optical stimulation in SC in the same
monkey (22.5ms, Extended Data Fig.1f). Saccadic vectors evoked from optical stimulation in
FEF and the SC were generally well aligned (Fig.1f compared with Extended Data Fig.1g). The
FEF optical stimulation results were highly reproducible across sessions and animals (Fig.1f,
Extended Data Fig.2a), with a consistent latency peak ranging from 27.5 ms to 32.5ms following
laser onset (Fig. 1g), compared with 22.5ms latency from SC (Extended Data Fig.1h). This 5 -
10ms latency difference is consistent with axonal conduction and synaptic delay through the
FEF-SC pathway[19,20]. Optical stimulation of the FEF in fixation tasks also induced consistent
contralateral saccades, though with smaller amplitudes and reduced frequency (Extended Data
Fig.2e-j). Importantly, no such effects were observed prior to the viral vector injection (Extended

Data Fig.2k) or during red light stimulation, ineffective to activate ChR2 (Extended Data Fig.2l),
confirming the specificity of the ChR2-mediated effects.
94
95
Next, we investigated whether subthreshold optical stimulation in the FEF could modulate the
planning of saccades without directly evoking them, as found in electrical stimulation studies of
the FEF and SC[21,22]. To do this, we applied low-intensity optical stimulation ($< 20mW/mm^2$)
during a delayed saccade task (Fig.1h) targeting the final phase of saccade preparation (200ms
after fixation offset). For contralateral saccades, reaction times were significantly faster with
stimulation (Opto: 158.23 ± 7.81ms, [Mean $\pm$ S.E.M], N = 20 sessions. Control: 171.44 ± 6.89ms;
N = 20 sessions. p = 0.0068, two-tailed paired t-test, Fig.1i,j). The facilitation effect was
strongest when the target location was aligned with the direction of the evoked saccade vector
(Extended Data Fig.2m). By contrast, ipsilateral saccades were delayed (Opto: 176.43 ± 6.54ms;
Control: 166.04 ± 6.69ms, N = 20 sessions; p = 0.0084, two-tailed paired t-test; Fig.1i,k). This
effect was most pronounced when the target was positioned opposite to the sites' preferred
direction (Extended Data Fig.2m). During this manipulation, optical stimulation did not evoke
any additional saccades, confirming that the stimulation remained subthreshold and acted
specifically to modulate the reaction time. Taken together, these findings demonstrated the
efficacy of our approach, in that retrograde optogenetic manipulation of the FEF-SC pathway can
both trigger saccades when suprathreshold (Fig.1d-g) and modulate the saccade timing when
subthreshold depending on the stimulation strength (Fig.1h-k).
113
114   We then sought to identify and characterize the FEF neurons projecting directly to the SC using
115   pathway-selective targeting with opto-tagging to determine whether the transmitted signals span
116   the full sensorimotor continuum (visual to visuomotor to motor) or are preferentially motor-
117   related. To accomplish this, we recorded neuronal activity in the FEF using a customized optrode
118   (Fig.2d), permitting simultaneous optical stimulation and electrophysiological recording[23].
119
Across recording sessions, a substantial proportion of FEF neurons (34.4%, 246/716) showed
robust modulation in response to optical stimulation (examples in Fig.2e, f), most commonly
with increased firing rates. A subset of neurons exhibited rapid and pronounced activation, with
response latencies of less than 10ms following laser onset (Fig.2e and g; Extended Data Fig.3a).
These short latency responses—observed in 20.4% of the recorded units (146/716)—were
presumed to reflect direct activation of SC-projecting FEF neurons expressing ChR2. (Note that
the results were robust by using a shorter-latency cutoff, Extended Fig.3g left). In addition to
direct activation, we also observed neurons exhibiting either inhibition (Fig.2f, upper panel) or
delayed activation (Fig.2f, lower panel), consistent with indirect modulation via presumed local
circuit interactions[24] or reciprocal connections with other brain areas, e.g. the mediodorsal
thalamus[25]. Beyond the different latencies, the presumed indirectly and directly modulated units
also showed different strengths of optical response modulation (Extended Data Fig.3b,
Modulation strength: Fast Ex, 65.54 ± 7.14, N = 146; Slow Ex, 16.32 ± 3.03, N = 32; Inh, 12.44
± 0.87, N = 68; p = 7.84e-36, Kruskal-Wallis Test). None of these effects were observed prior to
the viral vector injection (Extended Data Fig.3c,e) or during red-light stimulation which is
ineffective for ChR2 activation (Extended Data Fig.3d,f), ruling out nonspecific thermal effects.
Subsequent analyses were focused on this putatively directly activated SC-projecting FEF

neuronal population (the N = 146 "Fast Ex" FEF neurons, Fig.2g), to investigate their functional
properties in greater detail.
139
140
The FEF-SC pathway can be conceptualized in three alternative organizational schemes, based
on the functional properties of FEF output neurons and the properties of FEF-recipient SC
neurons (Fig.3a). One possibility, consistent with motor-enriched output of FEF[15], is the *motor*
dominance model. It posits that the FEF outputs are primarily motor-related signals that target
visuo-movement and movement neurons in the SC (Fig.3a, left). On the contrary, two other
possibilities align with a broad spectrum of signals leaving the FEF[16,17]. The *no convergence*
model proposes that the FEF transmits the full spectrum of signals, from purely visual to purely
motor, broadly targeting all SC neuron types (Fig.3a, right). The *visuomotor convergence model*
is an intermediate model and proposes a full-spectrum output from the FEF, but emphasizes
convergence onto movement-related neurons in the SC (Fig.3a, middle). Discriminating among
these models requires examination of both the functional properties of SC-projecting FEF
neurons and their targets within the SC.
153
To this end, we defined three subtypes of SC-projecting FEF neurons using a memory-guided
saccade task, which was designed to dissociate sensory responses from motor-related activity[16]
(Fig.3b,c). "Visual neurons" responded exclusively following target onset, without activation
preceding the saccade (example in Fig.3d top; subtype average in Fig.3e top). "Visuo-movement
neurons" showed both visual responses after target onset and motor-related activity prior to
saccade initiation (Fig.3d,e middle). By contrast, "movement neurons" exhibited activation only
before saccade onset, with no responses to the target presentation (Fig.3d, e bottom). The
optically labeled SC-projecting FEF neurons were distributed in approximately equal proportions
across all the three functional subtypes (Fig.3f, p = 0.86, Chi-Squared Test), indicating that a
diverse range of signal types is transmitted from the FEF to the SC. This distribution remained
qualitatively similar when applying a shorter-latency cutoff of < 5ms to define SC-projecting
FEF neurons (Extended Data Fig.3g right). These findings are therefore inconsistent with the
motor dominance model (Fig.3a left).
167
To further discriminate between the no convergence and visuomotor convergence models (Fig.3a,
right vs. middle), we next mapped the corresponding postsynaptic targets of the FEF projection
neurons within the SC. For this purpose, we performed laminar recordings in the SC while
optically stimulating the FEF (Fig.3g). Out of 163 SC neurons recorded across 10 sessions, FEF
stimulation modulated a substantial number (N = 64, 39.26%), with excitation being the
dominant effect (81%, 52/64; Extended Data Fig.4a). To isolate putative monosynaptic FEF
targets, we focused on SC neurons displaying rapid activation (Latency < 10ms; Extended Data
Fig.4b,). These units were primarily recorded from the middle channels of the SC probe (Fig.3g),
consistent with the SC intermediate layers. To quantify their functional properties, we again used
the memory-guided saccade task and applied the same classification as described above
(example raw data across channels in Fig.3h). Within each class, we examined the proportion of
rapidly activated units. The vast majority of activated SC neurons were either visuomotor (N =
22/52, 42.3%) or motor (N = 15/45, 33.3%) in nature, whereas visual-only neurons were rare
among the FEF-recipient population (N = 5/52, 9.6%; Fig.3i, p = 9.86e-06, Chi-Squared Test).
This functional bias could not be explained by sampling artifacts, because our recordings

sampled comparable numbers of visual (N = 52), visuo-movement (N = 52) and movement
neurons (N = 45), nor could it be attributed to the injection variability, as direct optical
stimulation of the SC produced a balanced distribution of all three subtypes (Extended Data
Fig.1d).
187
Hence using retrograde optogenetics, we conclude that FEF projections preferentially target
visuo-movement and movement neurons in the SC. This finding aligns with a prior study
employing anterograde activation via electrical stimulation[18]. But at the source of the projection,
the SC-projecting FEF neurons, we found that the FEF-SC pathway transmits a continuous
spectrum of sensorimotor signals ranging from visual, visuomotor and motor. This result helps
reconcile previous conflicting findings derived from electrical antidromic stimulation
approaches[15,17]. A clarifying feature of our study lies in its integrated approach: We
simultaneously examined both the source and the target regions within the same animals using
consistent methodologies. Thus this pathway-selective strategy provides a clearer view of the
underlying neural mechanisms supporting cognitive processes and reduces the variability
associated with studies conducted across different laboratories, species, or time periods.
199
Taken together, our findings support the visuomotor convergence model, which posits that
projections from the diverse signals in the FEF converge primarily onto motor-related neurons
within the SC. This functional convergence favors a view of sensorimotor processing as a
continuous, multistage transformation, where activity at any stage along the pathway can
advance the computation toward action[26]. Thus, the present results are also consistent with
cognitive models proposing graded transitions between perception and action[27,28]. Future work
will be needed to determine whether this organizational principle—convergence of diverse
cortical output signals onto motor representations in subcortical structures—can be generalized
across cortico-subcortical circuits in other sensorimotor systems.
209
Using the primate oculomotor system as a model, our study also highlights the utility of
retrograde optogenetics as a powerful tool for simultaneously manipulating behavior and
labeling pathway-defined neuronal populations. The selective manipulation of behavior is a
critical foundation for probing higher cognitive functions[29]. While prior optogenetic studies in
non-human primates have established causal links between neural activity and behaviors, these
efforts have largely relied on local stimulation at injection sites[30] or anterograde manipulation[31,32].
Our study fills a critical gap by achieving robust and selective manipulation of behavior via
retrograde, pathway-selective optogenetics. This approach yields behavioral effects comparable
in strength to classical methods including electrical stimulation, but with far greater specificity
by targeting pathway-defined neuronal subpopulations. Recent histological studies in macaques
have shown that AAV2-retro enables reliable retrograde labeling in several other cortical areas
such as the anterior cingulate cortex, primary motor cortex, and ventrolateral prefrontal cortex[6,33].
These cortical long-range projections are all likely glutamatergic, but further studies could
explore whether this approach is equally effective in targeting subcortical structures having
GABAergic projections, such as the substantia nigra and zona incerta[22,34,35]. Together, the
projection specificity, labeling precision, and behavioral efficacy of retrograde optogenetics
make it a highly promising tool for dissecting circuit-level computations underlying cognitive
processes in non-human primates, especially where functional heterogeneity and pathway
specificity are essential for understanding the neural basis of behavior.

Methods

Subjects

Two adult male rhesus monkeys (*Macaca mulatta*, monkeys R and A), weighing 9-12kg, participated in the present study. All experimental protocols and procedurals were approved by the National Eye Institute Animal Care and Use Committee and complied with the United States Public Health Service Policy on the Humane Care and Use of Laboratory Animals.

Surgery

We implanted a plastic head holder and two recording chambers for accessing the SC and FEF in the skull under general anesthesia and sterile surgical conditions for each monkey. The SC recording chamber was positioned on the midline and angled 38 degrees posterior to vertical on the sagittal plane, allowing electrode penetrations to approach the SC surface approximately orthogonally in both monkeys. The FEF recording chamber was placed over the frontal lobe and tilted laterally by 20 degrees. The head holder and recording chambers were secured with ceramic screws and embedded in dental acrylic resin, which covered the top of the skull. After confirming the position of recording chambers using MRI (4.7T, Bruker), a craniotomy was performed for each chamber during another surgery for future recordings.

Behavioral tasks

All behavioral task events and data acquisition were conducted using custom-made C++ based software (Blip, http://www.robilis.com/blip/). The monkeys were seated in a primate chair with their heads restrained, facing a fronto-parallel screen in a sound-attenuated and electrically shielded room. Visual stimuli were rear projected onto the screen using a digital light processing projector (PJD8353s, ViewSonic). Eye position was monitored at a sampling rate of 1KHz using an infrared eye-tracking system (EyeLink 1000 Plus, SR Research). Water rewards were delivered through a computer-controlled electromagnetic solenoid valve (Parker-Hannifin). The timing of all task-related visual events was recorded relative to the timing of the evoked signal using a photodiode positioned at the corner of the screen.

Behavior tasks for stimulation

We trained monkeys to perform four different tasks for evaluating the effect of optical or electrical stimulation.

Movie free viewing task

The subjects freely watched randomly selected natural history movies (15-25 minutes long) continuously with occasional random water rewards to keep them engaged. Unbeknownst to the subject, the entire viewing period was artificially divided into trials 2-3 seconds long separated by inter-trial intervals of 1-2 seconds. Optical (4.5mW, equals 50mW/mm^2, 100ms) or electrical (50uA, 100Hz, 100ms) stimulation was applied in half of the total trials, at a random time point between 500ms and 1s after the start of the trials. To ensure wakefulness and engagement, the

eye movements were monitored, and the subjects were required to keep their eyes open and
within the designated 40° × 40° window throughout the trial. Optical or electrical stimulation
was applied only when the eyes were detected within this window. If the animals moved their
eyes outside the window or closed them, the current trial was aborted and repeated later. No
movement constraints were imposed during the inter-trial intervals. If a subject successfully
completed a trial, a small amount of water reward was delivered at the end of the trial. A typical
session consisted of 80 or 160 successful trials.

Fixation Task

In the fixation task, subjects were required to maintain fixation throughout each trial. At the
beginning of a trial, a central fixation point (1° × 1° degree) was illuminated, and the subjects
had to fixate on the point within 500ms. Between 500ms and 1000ms after fixation onset, the
optical stimulation was applied for 100ms in half of the trials (4.5mW, equivalent to 50mW/mm^2,
100ms; stimulated trials), while no stimulation was applied in the other half (control trials).
Stimulated and control trials were randomly interleaved. Following the stimulation period,
subjects continued fixating for an additional 500ms to 1000ms, making the total fixation duration
for each trial 2s. The fixation point remained visible throughout the entire fixation period. The
fixation window was ±5°. Subjects were allowed to break fixation during the stimulation period
(100ms) and 200ms after the stimulated period, as well as during the corresponding 200ms
period in control trials. If fixation was broken during this permissive window, subjects had
500ms to re-fixate on the fixation point and complete the remaining fixation period to
successfully finish the trial. Successfully maintaining fixation for the required duration resulted
in a water reward. A typical session consisted of 80 or 160 trials.

Gap Fixation task

The procedure for the gap fixation task was nearly identical to that of the fixation task, except
that the fixation point disappeared for 500ms in the middle of the trial. During training, subjects
were required to maintain fixation in each trial even when the fixation point was absent. In the
stimulation experiment, optical stimulation was applied 200ms after the fixation disappeared,
and ended 200ms before the end of the fixation-off period. Eye movements during the stimulated
period were compared with those during the corresponding fixation-off period in control trials.

Delayed saccade task

In the delayed saccade task, subjects were required to make a saccadic eye movement to a visual
target that appeared in the periphery after a random delay. Each trial began with the presentation
of a central fixation point. After subjects maintained fixation for 500ms to 1000ms, a saccade
target was displayed either contralateral or ipsilateral to the stimulated hemifield. The target
location was aligned with the saccade vectors elicited by stimulating during the free-viewing task.
Following a delay period of 800ms to 1200ms, the fixation point was turned off, serving as the
go signal instructing the subject to make a saccadic eye movement toward the target location to
receive a liquid reward. Optical stimulation was applied using blue laser light for 200ms, starting
with the go signal. Trials with and without stimulation were randomly interleaved. To ensure
subthreshold stimulation, the laser power was reduced to less than half (< 2mW, equivalent to
20mW/mm^2). An initial set of 200 trials was first conducted before the formal experimental

session, to confirm that the stimulation did not directly trigger saccades (i.e. fewer than 1% of
cases, i.e. fewer than one saccade per 100 trials). A typical recording session consisted of 160 -
320 trials in total.

Behavior tasks for neural recordings

312 Once units were identified, we first manually mapped their receptive field approximately. We
313 then fixed the target presentation eccentricity and varied the target presentation angle to
314 determine the preferred vector using a receptive field mapping task. Following this, we
315 employed a memory saccade task to classify the functional subtype of each unit.

Receptive field mapping task

317 The procedure of the receptive field mapping task was identical to that of the delayed saccade
318 task except that no stimulation was applied and the target appeared at one of 8 possible locations:
319 0 degree (up), 45 (left - up), 90 degree (left), 135 degree (left - down), 180 degree (down), 225
320 degree (right - down), 270 degree (right), or 315 degree (right - up).

Memory-guided saccade task

The task began with the appearance of a fixation point. The subject was instructed to acquire and
maintain fixation within a $5° \times 5°$ window. After holding fixation for 800ms, a target was briefly
flashed for 200ms, then the fixation point remained for an additional 800ms –1200ms before
disappearing. The offset of the fixation point served as the go signal, prompting the subject to
saccade to the remembered target location within 500ms. To successfully trigger the
reappearance of the target stimulus, the saccade had to land within 5 degrees of the target
location. The subject then had to maintain fixation on the target for another 400ms to receive a
reward.

Viral vector preparation

Adeno-associated viral vectors with retrograde infection expressing the channelrhodopsin gene
(AAV2-Retro hSyn-ChR2-EYFP) were produced by the helper-free triple transfection procedure
and purified by affinity chromatography (GE Healthcare). Viral titer was determined by
quantitative PCR using Taq-Man technology (Life Technologies). The transfer plasmid (pAAV-
hSyn-ChR2-EYFP-WPRE) was constructed by replacing the CMV promoter of the pAAV-
CMV-ChR2-EYFP-WPRE plasmid[32] with the human Synapsin (hSyn) promoter.

Viral vector injection

We first identified the SC using a combination of magnetic resonance imaging and
electrophysiological properties. After the identification, AAV2retro-hSyn-ChR2-EYFP (2.5 x
10^{13} genome copies/mL) was injected into the SC of one hemisphere in both monkeys (monkey
R: left SC, monkey A: right SC) using a custom-made injectrode. For each monkey, two
penetrations were made at least 1.4mm apart. In each penetration, 1.0ul of the vector was
injected at two different depths, separated by at least 500um. The injection process began with an
initial infusion of 0.2uL at a rate of 0.4uL/min, followed by a slower infusion rate 0.02uL/min

for the remaining volume. This process was controlled using a 10uL Hamilton syringe and a
motorized infusion pump (Harvard Apparatus, Holliston, MA, USA). After each injection, the
injectrode was left at the site for 1 hour before proceeding to the next site to ensure absorption of
viral solution into the surrounding neural tissue. Recording and testing experiments were
conducted 2 - 3 month after injection.

Laser light stimulation

We used custom-made single channel or multi-channel optrodes for optical stimulation and
electrophysiology recordings. The single channel optrode had an optic fiber (Doric Lenses,
200um diameter, 0.5NA) attached to a tungsten recording electrode (FHC, 0.5-2MOhm
impedance, 125um diameter). The tip of the electrode typically extended 200um beyond the fiber
tip. The multi-channel optrodes had the same optic fiber attached to a 24-channel laminar probe
(V probe, Plexon INC,100um inter-contact- distance). The tip of the fiber was placed close to the
third and fourth channels on the probe. The light source was a 473nm blue diode laser with a
maximum power up to 300mW (Topica photonics, iBeam smart). Before the optrode was
inserted into the brain, we measured the light intensity at the tip of the optrode using an optical
power meter (Newport Corporation, Centauri). During the movie free viewing task, we typically
fixed the output power to be 4.5mW (~50mW/mm^2) in FEF and 1.5mW (~16mW/mm^2) in SC,
for significant behavior and neuronal modulation.

Electrophysiology

Electrophysiological recordings were performed using either single channel or 24-channel V
probe optrodes (Plexon). Electrodes were inserted into the brain through a stainless-steel guide
tube and advanced using oil-driven micromanipulator (MO-97A, Narishige). Recording sites
were targeted using a grid system that enable electrode placement at 1-mm intervals in x and y
directions, orthogonal to the guide tube. Extracellular signals were amplified and band-pass-
filtered (100☐Hz to 8kHz) using a multichannel processor (Plexon). Spike sorting was
conducted offline using Kilosort[36] and manually curated by a human expert using Phy2
(www.github.com/cortex-lab/phy) to ensure that the sorted units exhibited physiologically
plausible inter-spike interval distributions and waveform shapes consistent with action potentials.
Neurons were excluded from analysis if they had low signal-to-noise ratio, a low average trial
firing rate (<1spikes/s), or lacked a clear visual or motor responses. The FEF was identified in
the anterior bank of the arcuate sulcus, where saccades could be evoked via low threshold
electrical stimulation (Threshold < 50uA, 100Hz) and neurons showed clear visual and/or
saccade related responses. The SC was identified as showing clear visual and/or motor responses
and clear transition from visual neurons to visual-movement neurons and to movement units
during the penetration.

General Experimental Procedures

During a typical recording session, once neurons were stabilized and well-isolated, the receptive
field of a randomly selected neuron was initially mapped by manually placing a square target at
various screen locations to identify the position that elicited the strongest neural responses. At
this stage, the receptive field locations were roughly determined by subjectively evaluating the
strength of the neural responses through auditory monitoring, and those responses were

confirmed as either visual or saccade related. Next, the movie free viewing task was
administered to assess the optical modulation of the recorded neurons. The isolated units were
then tested on a subset or all of the following tasks: fixation stimulation task, gap fixation task,
delayed saccade task, receptive mapping task, and the memory saccade task, among others.
Neurons were recorded across all subsequent tasks regardless of whether they exhibited optical
modulation, in order to avoid sampling bias.

Dataset

In total, we recorded 716 neurons from the FEF, comprising 562 neurons from Monkey R across
82 sessions and 154 neurons from Monkey A across 28 sessions, to evaluate the effects of optical
modulation during the movie free-viewing task. Of these, 612 neurons were also recorded during
the memory saccade task for functional subtype characterization. We excluded 30 neurons due to
a mean firing rate of less than 1 spike/s during the task period, and 77 neurons due to the absence
of significant visual or saccade-related responses. This resulted in 505 neurons being included
for further functional subtype characterization. To assess optical modulation effects in the SC,
we recorded 368 neurons (285 from Monkey R in 37 sessions and 83 from Monkey A in 8
sessions). Among these, 323 neurons were also recorded during the memory saccade task. We
excluded 23 neurons for low firing rate (< 1 spike/s) and 59 neurons for lacking significant visual
or saccade related activity, resulting in 241 SC neurons retained for functional subtype
characterization. In addition, we recorded 163 extra SC neurons in 10 sessions while conducting
optical stimulation in the FEF, of which 149 were also recorded during the memory saccade task
and used for functional subtype characterization.

Data analysis

Saccade analysis

In the free-viewing task, eye traces spanning 300ms—comprising 100ms before optical
stimulation onset, the 100ms optical stimulation period or un-stimulated control period, and the
100ms after stimulation offset—were extracted for saccade analysis. Raw eye position traces
were differentiated to obtain velocity and acceleration profile. Saccade detection was based on a
modified version of the algorithm described previously[37]. Briefly, candidate saccades were
identified when eye velocity exceeded 200 degrees/s and acceleration exceeded 1000 degree/s^2.
Saccades were then detected as the time point when velocity surpassed a threshold of 40
degrees/s. Saccade onset was defined as the time point when velocity first exceeded 40 degrees/s
and offset was defined as the time point when velocity fell below this threshold. Sequential
saccades were considered valid only if the inter-saccade interval was at least 20ms. For each
detected saccade, direction was calculated as the vector from the saccade start to end point, and
amplitude was defined as the Euclidean distance between the saccade start to the end point. To
minimize noise, only saccades with amplitudes between 5 to 40 degrees were included in the
analysis for this task. In the delayed saccade task, reaction time was defined as the interval
between the offset of the fixation point and the onset of the saccade towards the visual target.

Optical modulation analysis

For optical stimulation experiments, the neural responses of FEF during the 100ms stimulation
period were compared to the corresponding 100ms period in the non-stimulated (control) trials.
Neurons were labeled as optically modulated if their average responses during stimulation
differed significantly from those in control trials, tested via a two-sample t-test using a criterion
of $p<.05$. Neurons showing higher activity during stimulation compared to control were
classified as excitatory modulated, while those with lower activity were classified as inhibitory
modulated. For significantly modulated neurons, we then measured the modulation latency. Raw
spike rasters were divided into two segments: pre-stimulation and stimulation (from onset to
offset). Within the stimulation segment, we compared the spike counts in each 10ms bin to the
10ms bin immediately preceding stimulation onset using a paired t-test. The 10ms window was
advanced in 1ms steps for successive comparisons. If 5 consecutive windows showed significant
differences, the latency was defined as the onset time of the first significant window. The latency
was marked as NaN and excluded from the significantly modulated group if no significant
window was detected by the end of the stimulation period. Based on the latency, the excitatory
modulated population was further divided into 2 groups: (1) fast excitatory neurons with latency
<10ms; (2) slow excitatory neurons with latency >=10ms.

Functional subtype characterization

We used spike rates from three time windows in the memory saccade task to categorize neurons
in the FEF, following the criteria defined previously[38]: a baseline window (−100ms to -50ms
relative to target onset), a visual response window (+60ms to +100ms after target onset), and a
pre-saccadic window (−100ms from saccade onset up to saccade offset). For each neuron, a one-
way Kruskal-Wallis nonparametric ANOVA was performed across trials using spike rates from
the three time windows. Post hoc comparisons were then used to assess whether the visual and/or
saccadic activity differed significantly from baseline. Neurons showing significant visual but not
saccadic modulation were labeled visual neurons (N = 177). Neurons with significant activity in
both visual and saccade windows were classified as visual-movement neurons (N = 227).
Neurons with significant activity only in the saccade window were labeled movement neurons
(N = 103). The characterization for SC neurons used the same approach but with a time window
adjusted according to the response properties of SC neurons: baseline window: -100 to -50ms
relative to target onset; visual response window: +30 to +100ms after target onset, and the pre-
saccadic window: -50ms from saccade onset up to saccade offset.

Histology

After completing all experiments, monkey R was sedated with ketamine HCl and then deeply
anesthetized with an overdose of sodium pentobarbital (50 mg/kg, intravenous). Once the animal
reached a fully areflexic state, the transcranial perfusion was performed, beginning with 6 liters
of 0.1 M phosphate-buffered saline (PBS, pH7.2), followed by 4L of 4% paraformaldehyde in
0.1M PBS (pH7.2). After perfusion, the brain was fixed to the stereotaxic frame and cut into
blocks in the coronal plane including the midbrain regions. These blocks were postfixed in 4%
paraformaldehyde overnight at 4°C, then cryoprotected in a 30% sucrose solution in 0.1 m PBS
at 4°C. Once fully saturated and sunk, the tissue was frozen and serially sectioned into 50um
slices using a freezing stage sliding microtome (American Optical Company). The tissue sections
were then stored in PBS at 4°C until processed for immunofluorescence.

Sections were divided into six evenly spaced series. One series was reserved for assessing endogenous fluorescence while the other five were subjected to immunofluorescent amplification. Detailed procedures for immunofluorescent amplification have been previously published[39]. Briefly, serial free-floating sections were rinsed in PBS, permeated with 0.5% Triton-X 100 in PBS, blocked with 1% Bovine Serum Albumin and Triton-X 100 solution, and incubated in Goat Anti-GFP (1:800, Rockland Cat#: 600-101-215) antibody solution at 4°C on a rocker for 48 hours. Next, the tissue was rinsed and incubated in a secondary antibody solution of 0.5% Triton-X 100, 1% Normal Donkey Serum, and Donkey anti-Goat Alexa Fluor 488 (1:200, ThermoFisher Cat#: A-11055) at room temperature. After three hours, the sections were rinsed in PBS then incubated in Hoechst 33342 (1:1000, ThermoFisher Cat#: 62249) for 10 minutes. Finally, the tissue was rinsed in PBS and mounted on 1% gelatinized microscope slides. Mounted tissue was then dried at room temperature overnight, then dehydrated in graded ethanols (70%, 95%, 100%) and cleared with xylene. The tissue was coverslipped using Cytoseal 60. The sections were then digitized using an Axioscan.Z1 (Carl Zeiss) at 5x using an Axiocam 506 and HXP 120V as a light source using the DAPI, Green Fluorescent, and Texas Red filters.

Acknowledgements

We thank Nick Nichols, Daniel Yochelson, Denise Parker, and Hayden Warnock for technical support and Carlos Mejias-Apont for assistance in histological preparation of the material. We are also grateful to Anthony Movshon and Mike Hasse from NYU for their insightful discussions and valuable suggestions. This research was supported by the Intramural Research Program of the National Institutes of Health (NIH), United States and by award number R01NS125843 to M.A.S. from the National Institute Of Neurological Disorders And Stroke of the National Institutes of Health. The contributions of the NIH authors were made as part of their official duties as NIH federal employees, are in compliance with agency policy requirements, and are considered Works of the United States Government. However, the findings and conclusions presented in this paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human Services.

Author Contributions

X.Y., A.G. and O.H. contributed to the conceptualization and experimental design. X.Y and A.G. performed the experiments and analyzed the data. K.I. and M.T. produced the viral vector. M.B. and G.K conducted the histology and immunofluorescence histochemistry. X.Y. wrote the original draft of the manuscript. A.G., K.I, M.B., H.N, M.A.S, and M.T. contributed to manuscript editing. All authors contributed to data interpretation and reviewed and approved the final manuscript.

Declaration of interests

The authors declare no competing financial interests.

Figure Legends

Fig1. Robust saccade modulation induced by optical stimulation of the FEF-SC pathway.
a, Schematic illustration of the experimental design. The retrograde viral vector was injected into the SC, from where it was transported retrogradely to the somata of neurons in the FEF, thereby tagging the projection neurons to the SC. Optical stimulation was applied in the FEF.
b, Illustration of the movie free-viewing task. Macaques viewed movies and were rewarded for keeping their gaze within the display.
c, Optical stimulation protocol during the movie free-viewing task. Each free-viewing session consisted of two randomly interleaved trial types: stimulation trials, which included a 0.1s optical pulse from a blue laser (left red panel), and control trials, which did not include stimulation (right black panel). Each optical stimulation was embedded within two non-stimulated periods lasting 2-3s.
d, An example session showing saccades that occurred in stimulation trials (left) and control (right) trials during optical stimulation in the FEF.
e, Example peri-stimulus time histogram (PSTH) of saccades in the stimulation (red) and control trials (black) during optical stimulation in the FEF (N = 40 trials per condition). Upper raster shows time of saccade onsets in individual trials (both spontaneous and putatively stimulation-evoked). Lower histogram shows distributions of those onset times binned at a 5ms intervals. The triangle indicates the time point of peak saccade occurrence (Peak time: 35ms).
f, Polar histograms of saccade directions in stimulation and control trials for two monkeys during optical stimulation in the FEF. The histogram is binned in 22.5° intervals. The radial (ρ) axis represents the proportion of saccades within each angular bin, normalized to the total number of saccades recorded during the 100ms interval under each condition for each monkey.
g, Population PSTH of saccades in stimulation and control trials for two monkeys during optical stimulation in the FEF. To generate the PSTH, saccades were first counted and averaged across trials in 5ms bins from -100ms to 200ms relative to stimulation onset for each session. The average counts were then divided by the 5ms bin duration to yield the saccade frequency. The triangle indicates the time point of peak saccade occurrence for monkey R (red: 32.5ms) and monkey A. (magenta: 27.5ms)
h, Schematic of the delayed saccade paradigm. In each trial, a target appeared in either the left or right visual field, followed by a delay period of 0.8-1.2s. After the delay indicated by disappearance of the fixation point, the animal was required to make a saccadic eye movement to the target location.
i, An example session showing eye movement traces for contralateral (top) and ipsilateral saccades (bottom) during optical stimulation (red) and control (black) trials. Contralateral: Reaction Time (RT) for Optical stim: 122.9 ± 8.1ms [Mean ± S.E.M] N = 20 trials; RT for Control: 154.4 ± 3.3ms, N = 20 trials; p = 7.4516e-04, two sample two-tailed t-test. Ipsilateral: RT for Optical stim: 170.4 ± 5.7ms, N = 20 trials. RT for Control: 157.6 ± 5.7ms, N = 20 trials; p = 0.06, two sample two-tailed t-test.
j, Scatter plot of reaction times (RTs) for contralateral saccades in the optogenetic stimulation versus the control condition. N=20 sessions. The inset displays a bar plot for the mean RT for each condition. Mean RT during Optical stim: 158.23 ± 7.81ms, [Mean ± S.E.M]; Mean RT during Control: 171.44 ± 6.89ms; p = 0.0068, two-tailed paired t-test. Filled Symbols: Sessions

with a significant difference in RT between control and optogenetic stimulation (p < 0.05, one-tailed two sample t-test); Open Symbol: No significant difference (p > 0.05). Triangle: Monkey R; Square: Monkey A.

k, Scatter plot of reaction times for ipsilateral saccades in the optogenetic stimulation versus the control condition. N = 20 sessions. The inset displays a bar plot for the mean RT for each condition. Mean RT during Optical stim: 176.43 ± 6.54ms; Mean RT during Control: 166.04 ± 6.69ms; p = 0.0084, two-tailed paired t-test.

Fig2. AAV2Retro efficiently labeled neurons within the Frontal Eye Field (FEF)-Superior Colliculus (SC) pathway.

a, Immunofluorescence showing ChR2-EYFP expression in the injected hemifield of SC.

b, Immunofluorescence showing retrograde labeling of ipsilateral FEF neurons by ChR2-EYFP. The right panel is the magnified window(100X) showing the labeled somata. The triangles indicate two examples.

c, Lack of immunofluorescence in the contralateral FEF. Weak, expected labeling was observed in the adjacent area, ventrolateral prefrontal cortex (area 45), which is known to project to the SC[40].

d, A schematic illustrating the experiment involving optical stimulation and simultaneous electrical recording in the FEF.

e, Two examples of putative direct activation by optical stimulation, in which neurons showed rapid (Latency: 0ms) and robust activation. The top panels show the spike waveforms for each unit: red traces represent waveforms recorded during optical stimulation, while grey traces represent those recorded outside the stimulation period. The action potential waveforms remained consistent during optical stimulation, indicating stable unit identity.

f, Two examples of putative indirect activation by optical stimulation. The top example shows inhibition (top, Latency 15ms) and the bottom one shows delayed activation (Latency: 32ms). The top panels show the waveforms for each unit. Similar to panel **e,** the waveforms remained consistent during optical stimulation.

g, Population average across three groups of neurons in the FEF based on optical stimulation effects and modulation latency. The raw peristimulus histogram (PSTH) for each neuron was normalized by dividing by its peak firing rate. Subsequently, the average baseline activity (-100ms to 0ms relative to laser onset) was subtracted, resulting in a normalized firing rate ranging approximately from -1 to 1. The pie chart in the bottom-right corner shows the proportion of FEF neurons in each group. Fast Excitation—Fast Ex (Red): N = 146; Slow Excitation—Slow Ex (Orange): N = 32; Inhibition—Inh (Blue): N = 68; Not Effective—No effect (Grey): N = 470.

Fig 3. Combining optogenetic identification and behavior reveals the functional organization of the FEF-SC pathway for saccade generation.

a, Schematic illustrating possible models of information flow from the FEF to the SC. Both the FEF and SC contain visual (V), visuomotor (VM) and motor (M) related signals, but the nature of information transmitted from FEF and its target within SC remains controversial. Motor dominant model (left): FEF transmits primarily motor signals that target visuo-movement and movement neurons. Visuomotor convergence model (middle): FEF transmits a full spectrum of signals from visual to motor, and it converges onto movement related neurons in the SC. No

convergence model (right): the output from FEF is also a full-spectrum, but it broadly targets all
signal types in the SC. Data presented in panels **d-i** support the Visuomotor convergence model.
b, A schematic showing the behavioral paradigm for the memory saccade task. A target was
briefly presented for 100ms, and then disappeared. After a variable delay of 800-1200ms, the
fixation spot disappeared, instructing the monkey to make a saccade to the remembered target
location.
c, Time course of the memory saccade task.
d, Representative responses of three example SC-projecting neurons in the FEF during the
memory saccade task: a visual neuron (top), a visual-movement neuron (mid), and a movement
neuron (bottom).
e, Population PSTH for SC-projecting FEF neurons: visual neurons (top, N = 42/177, 23.73%),
visual-movement neurons (mid, N = 49/227, 21.59%), and movement neurons (bottom, N =
22/101, 21.78%). For each neuron, the raw PSTH was normalized by z-scoring, whereby the
mean firing rate was subtracted from each time point and the result was divided by the standard
deviation.
f, Proportion of SC-projecting FEF neurons that were fast-excited during optical stimulation,
across the three functional subtypes of recorded FEF neurons: visual neurons, visual-movement
neurons, and movement neurons. The proportion for each subtype was calculated as the number
of SC-projecting FEF neurons identified in each subtype (numerator above each bar) divided by
the total number of FEF neurons recorded for that subtype (the denominator above each bar).
This proportion definition provides an unbiased estimate of the likelihood of identifying SC-
projecting neurons within each subtype, minimizing potential bias from under- or oversampling
of specific subtypes. n.s.: no significant difference in the proportions among the three groups, p =
0.86, tested by chi-squared test.
g, An example session of laminar recordings in the SC during optical stimulation of the FEF,
shown as a heatmap (left). Modulated units were further classified as fast excitatory units based
on excitatory responses with latencies below 10ms. These units are considered putative SC
neurons directly activated by FEF stimulation, representing SC targets receiving direct
projections from the FEF. They are highlighted in green in the line plot (right).
h, A line plot showing the firing rate of the same SC population showing in panel **g** during the
memory saccade task, aligned on target onset (left) and saccade onset (right). Green thick lines
indicate channels on the laminar probe with the fast-excitation (latency < 10ms) optically
modulated units.
i, Proportion of FEF-recipient SC neurons across the three functional subtypes in the SC. The
definition of proportion is the same as in panel **f**. ***, p < 0.001, chi-squared test.
633

References

Gou, L. *et al.* Single-neuron projectomes of macaque prefrontal cortex reveal refined axon targeting and arborization. *Cell* **188**, 3806–3822 e3824 (2025). https://doi.org/10.1016/j.cell.2025.06.005

Miller, E. K. & Cohen, J. D. An integrative theory of prefrontal cortex function. *Annu Rev Neurosci* **24**, 167–202 (2001). https://doi.org/10.1146/annurev.neuro.24.1.167

San Sebastian, W. *et al.* Adeno-associated virus type 6 is retrogradely transported in the non-human primate brain. *Gene Ther* **20**, 1178–1183 (2013). https://doi.org/10.1038/gt.2013.48

Tervo, D. G. *et al.* A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons. *Neuron* **92**, 372–382 (2016). https://doi.org/10.1016/j.neuron.2016.09.021

Chen, Y. *et al.* A Novel Retrograde AAV Variant for Functional Manipulation of Cortical Projection Neurons in Mice and Monkeys. *Neurosci Bull* **40**, 90–102 (2024). https://doi.org/10.1007/s12264-023-01091-0

Cushnie, A. K. *et al.* Using rAAV2-retro in rhesus macaques: Promise and caveats for circuit manipulation. *J Neurosci Methods* **345**, 108859 (2020). https://doi.org/10.1016/j.jneumeth.2020.108859

Galvan, A. *et al.* Nonhuman Primate Optogenetics: Recent Advances and Future Directions. *J Neurosci* **37**, 10894–10903 (2017). https://doi.org/10.1523/JNEUROSCI.1839-17.2017

Tremblay, S. *et al.* An Open Resource for Non-human Primate Optogenetics. *Neuron* **108**, 1075–1090 e1076 (2020). https://doi.org/10.1016/j.neuron.2020.09.027

Shaw, L., Padmanabhan, K., Buckleaw, A., Mitchell F, J. & Wang K, H. Projection-Specific Intersectional Optogenetics for Precise Excitation and Inhibition in the Marmoset Brain. *bioRxiv* (2025). https://doi.org/10.1101/2025.06.18.660378

Schiller, P. H., True, S. D. & Conway, J. L. Effects of frontal eye field and superior colliculus ablations on eye movements. *Science* **206**, 590–592 (1979). https://doi.org/10.1126/science.115091

Tehovnik, E. J., Sommer, M. A., Chou, I. H., Slocum, W. M. & Schiller, P. H. Eye fields in the frontal lobes of primates. *Brain Res Brain Res Rev* **32**, 413–448 (2000). https://doi.org/10.1016/s0165-0173(99)00092-2

Sparks, D. L. Translation of sensory signals into commands for control of saccadic eye movements: role of primate superior colliculus. *Physiol Rev* **66**, 118–171 (1986). https://doi.org/10.1152/physrev.1986.66.1.118

Schall, J. D. The neural selection and control of saccades by the frontal eye field. *Philos Trans R Soc Lond B Biol Sci* **357**, 1073–1082 (2002). https://doi.org/10.1098/rstb.2002.1098

Everling, S. & Munoz, D. P. Neuronal correlates for preparatory set associated with pro-saccades and anti-saccades in the primate frontal eye field. *J Neurosci* **20**, 387–400 (2000). https://doi.org/10.1523/JNEUROSCI.20-01-00387.2000

15   Segraves, M. A. & Goldberg, M. E. Functional properties of corticotectal neurons in the
monkey's frontal eye field. *J Neurophysiol* **58**, 1387–1419 (1987).
https://doi.org/10.1152/jn.1987.58.6.1387

16   Sommer, M. A. & Wurtz, R. H. Frontal eye field sends delay activity related to
movement, memory, and vision to the superior colliculus. *J Neurophysiol* **85**, 1673–1685
(2001). https://doi.org/10.1152/jn.2001.85.4.1673

17   Sommer, M. A. & Wurtz, R. H. Composition and topographic organization of signals
sent from the frontal eye field to the superior colliculus. *J Neurophysiol* **83**, 1979–2001
(2000). https://doi.org/10.1152/jn.2000.83.4.1979

18   Helminski, J. O. & Segraves, M. A. Macaque frontal eye field input to saccade-related
neurons in the superior colliculus. *J Neurophysiol* **90**, 1046–1062 (2003).
https://doi.org/10.1152/jn.00072.2003

19   Robinson, D. A. Eye movements evoked by collicular stimulation in the alert monkey.
Vision Res **12**, 1795–1808 (1972). https://doi.org/10.1016/0042-6989(72)90070-3

20   Bruce, C. J., Goldberg, M. E., Bushnell, M. C. & Stanton, G. B. Primate frontal eye fields.
II. Physiological and anatomical correlates of electrically evoked eye movements. *J*
Neurophysiol **54**, 714–734 (1985). https://doi.org/10.1152/jn.1985.54.3.714

21   White, R. L., 3rd & Snyder, L. H. Subthreshold microstimulation in frontal eye fields
updates spatial memories. *Exp Brain Res* **181**, 477–492 (2007).
https://doi.org/10.1007/s00221-007-0947-7

22   Carello, C. D. & Krauzlis, R. J. Manipulating intent: evidence for a causal role of the
superior colliculus in target selection. *Neuron* **43**, 575–583 (2004).
https://doi.org/10.1016/j.neuron.2004.07.026

23   Acker, L., Pino, E. N., Boyden, E. S. & Desimone, R. FEF inactivation with improved
optogenetic methods. *Proc Natl Acad Sci U S A* **113**, E7297–E7306 (2016).
https://doi.org/10.1073/pnas.1610784113

24   Liske, H., Qian, X., Anikeeva, P., Deisseroth, K. & Delp, S. Optical control of neuronal
excitation and inhibition using a single opsin protein, ChR2. *Sci Rep* **3**, 3110 (2013).
https://doi.org/10.1038/srep03110

25   Sommer, M. A. & Wurtz, R. H. Frontal eye field neurons orthodromically activated from
the superior colliculus. *J Neurophysiol* **80**, 3331–3335 (1998).
https://doi.org/10.1152/jn.1998.80.6.3331

26   Wurtz, R. H., Sommer, M. A., Paré, M. & Ferraina, S. Signal transformations from
cerebral cortex to superior colliculus for the generation of saccades. *Vision Res* **41**, 3399–
3412 (2001). https://doi.org/10.1016/s0042-6989(01)00066-9

27   Miller, J. Discrete and continuous models of human information processing: theoretical
distinctions and empirical results. *Acta Psychol (Amst)* **67**, 191–257 (1988).
https://doi.org/10.1016/0001-6918(88)90013-3

28   Taylor, D. A. Stage analysis of reaction time. *Psychological Bulletin* **83**, 161–101 (1976).
https://doi.org/ https://doi.org/10.1037/0033-2909.83.2.161

29   Jazayeri, M. & Afraz, A. Navigating the Neural Space in Search of the Neural Code.
Neuron **93**, 1003–1014 (2017). https://doi.org/10.1016/j.neuron.2017.02.019

30   Katz, L. N. *et al.* Optogenetic Manipulation of Covert Attention in the Nonhuman
Primate. *J Cogn Neurosci* **37**, 266–285 (2025). https://doi.org/10.1162/jocn_a_02274

Amita, H., Kim, H. F., Inoue, K. I., Takada, M. & Hikosaka, O. Optogenetic manipulation of a value-coding pathway from the primate caudate tail facilitates saccadic gaze shift. *Nat Commun* **11**, 1876 (2020). https://doi.org/10.1038/s41467-020-15802-y

Inoue, K. I., Takada, M. & Matsumoto, M. Neuronal and behavioural modulations by pathway-selective optogenetic stimulation of the primate oculomotor system. *Nat Commun* **6**, 8378 (2015). https://doi.org/10.1038/ncomms9378

Weiss, A. R., Liguore, W. A., Domire, J. S., Button, D. & McBride, J. L. Intra-striatal AAV2.retro administration leads to extensive retrograde transport in the rhesus macaque brain: implications for disease modeling and therapeutic development. *Sci Rep* **10**, 6970 (2020). https://doi.org/10.1038/s41598-020-63559-7

May, P. J. & Basso, M. A. Connections between the zona incerta and superior colliculus in the monkey and squirrel. *Brain Struct Funct* **223**, 371–390 (2018). https://doi.org/10.1007/s00429-017-1503-2

Wurtz, R. H. & Hikosaka, O. Role of the Basal Ganglia in the Initiation of Saccadic Eye-Movements. *Prog Brain Res* **64**, 175–190 (1986).

Pachitariu, M., Sridhar, S., Pennington, J. & Stringer, C. Spike sorting with Kilosort4. *Nat Methods* **21**, 914–921 (2024). https://doi.org/10.1038/s41592-024-02232-7

Maeda, K., Inoue, K. I., Kunimatsu, J., Takada, M. & Hikosaka, O. Primate Amygdalo-Nigral Pathway for Boosting Oculomotor Action in Motivating Situations. *iScience* **23**, 101194 (2020). https://doi.org/10.1016/j.isci.2020.101194

McPeek, R. M. & Keller, E. L. Saccade target selection in the superior colliculus during a visual search task. *J Neurophysiol* **88**, 2019–2034 (2002). https://doi.org/10.1152/jn.2002.88.4.2019

Daw, T. B. *et al.* Direct Comparison of Epifluorescence and Immunostaining for Assessing Viral Mediated Gene Expression in the Primate Brain. *Hum Gene Ther* **34**, 228–246 (2023). https://doi.org/10.1089/hum.2022.194

Borra, E., Gerbella, M., Rozzi, S. & Luppino, G. Projections from caudal ventrolateral prefrontal areas to brainstem preoculomotor structures and to Basal Ganglia and cerebellar oculomotor loops in the macaque. *Cereb Cortex* **25**, 748–764 (2015). https://doi.org/10.1093/cercor/bht265

Figure 1

Figure 2

Figure 3