Drought and methyl jasmonate memory interact to determine plant functioning under

current drought stress in a perennial grass

Tarun Bhatt[1*], Nikita Rathore[2], Jaroslav Semerad[3], Tomas Cajthaml[3], Dinesh Thakur[2],

Zuzana Münzbergová[1,2]

[1]Department of Botany, Faculty of Sciences, Charles University, Benatska 2, 12800, Prague,

Czech Republic, 12800

[2]Institute of Botany, Czech Academy of Sciences, Zamek 1,25243, Pruhonice, Czech

Republic

[3]Institute of Microbiology of the Czech Academy of Sciences, Videnska 1083, 142 20 Prague,

Czech Republic

*Corresponding author e-mail: bhatttarun954@gmail.com

Abstract

Climate-driven droughts necessitate deeper insights into plant resilience, particularly in clonal

perennials. This study presents the first integrative analysis of how recurrent drought (D2),

drought memory (D1), and methyl jasmonate-induced memory (M1) independently and

interactively shape metabolomic diversity, composition, and morpho-physiological traits.

Using *Festuca rubra* as a model species and using a full-factorial design, we found that current

drought (D2) was the primary driver of metabolic reprogramming, reducing metabolite

richness but enhancing uniqueness, suggesting a streamlined, stress-optimized strategy. D1 and

M1 had significant, context-dependent impacts: D1 increased richness under well-watered

conditions but reversed under drought, while M1 stabilized both metabolomic and trait

variability. Importantly, D1 and M1 effects were non-additive, producing emergent

biochemical states and morpho-physiology. A significant D1 × D2 × M1 interaction revealed

complex integration of environmental and hormonal legacies. The metabolomic shifts closely

aligned with traits such as specific leaf area and relative water content, indicating a coordinated

adaptation at both metabolomic and morpho-physiological levels through memory-driven

mechanisms. Our findings confirm that both drought and hormone induced memories are

transgenerational and compositional, with synergistic roles in plant stress adaptation. This

suggests that combining hormonal and environmental priming strategies may be a viable

approach to enhancing plant resilience under increasing climate variability.

Keywords: Metabolomics, methyl jasmonate, MeJA memory, drought resilience, clonal plants,

transgenerational effects, phenotypic plasticity, drought memory, *Festuca rubra*.

1 Introduction

Plants are experiencing rapidly changing environments and face various stress factors, which

can impact their growth and reproduction (Crisp et al., 2016). Drought stress is among the most

significant stresses affecting plant growth and productivity worldwide (Michaletti et al., 2018).

The 2023 IPCC report indicates that under projected climate change, drought events will occur

more frequently and become more severe, threatening the structure, function, and productivity

of ecosystems. Therefore, understanding the effects of drought on plants is crucial for

developing effective strategies to mitigate damage and enhance plant resilience and adaptation

to these changing conditions.

To cope with drought, plants employ various strategies involving numerous biological

mechanisms activated during different growth stages at the cellular, organ, and whole plant

levels (Kumar et al., 2018). While plant responses to drought stress are intensively investigated

from cellular to community levels (e.g., Kumar et al., 2018; Ploughe et al., 2019), plant drought

responses at the metabolite level are still limited to investigating a few cellular processes (e.g.,

Almeida et al., 2020; Fàbregas & Fernie, 2019; You et al., 2019), and general metabolic profile

changes are poorly understood. Metabolic profiles provide an understanding of whole cellular

functioning and may thus help in developing strategies to enhance plant resilience to

drought(Bowne et al., 2012; Calleja-Satrustegui et al., 2024). Untargeted metabolomics also

provides a broad, unbiased view of metabolic shifts, allowing identification of novel

metabolites and pathways that may be involved in drought responses.

Most research on the drought effects on plants looks at the responses to single drought

events (e.g., Gupta et al., 2020; Ploughe et al., 2019), and far less attention has been given to

how plants cope with recurrent drought episodes (but see e.g., Virlouvet & Fromm, 2015;

Kambona et al., 2023). Understanding the effect of recurrent drought is, however, becoming

increasingly important as drought events are becoming more frequent (IPCC 2023), requiring

plants to not just survive individual stress events but also develop adaptive strategies to deal

with repeated drought via stress memory (Crisp et al., 2016).

Stress memory involves structural changes, epigenetic reprogramming, and

biochemical adjustments during initial drought exposure, priming plants for more efficient

future responses through sustained gene expression changes, protective protein accumulation,

and ongoing metabolic reprogramming (Crisp et al., 2016; Fleta-Soriano & Munné-Bosch,

2016; Sharma et al., 2022). Exploring metabolic changes linked to stress memory may broaden

our understanding of how plants adapt to environmental challenges by identifying key

metabolites, revealing diverse cellular pathways and structural responses involved in long term

stress adaptation (Aina et al., 2024). Studying the metabolic changes will thus help us to

understand how plants 'remember' stress at the molecular level and reorganize their

metabolome to become more resilient to recurrent drought. Even stronger inferences can be

made by linking the metabolic changes to morphophysiological changes in the same

individuals(Walker et al., 2023), but such comparisons are extremely rare.

Plant hormones play a crucial role in modifying plant performance under drought

conditions (Salvi et al., 2021a; Waadt et al., 2022). Methyl jasmonate (MeJA) enhances plant

resilience to drought by regulating key developmental processes such as seed germination, root

growth, flowering, and senescence (Hewedy, Elsheery, Karkour, Elhamouly, Arafa, Mahmoud,

Dawood, Hussein, Mansour, Amin, et al., 2023; Yu et al., 2018). It also influences root and

shoot growth, promoting water uptake and plant health during drought (Lee & Zwiazek, 2019;

Wei et al., 2025). Despite growing evidence for MeJA's effectiveness in enhancing drought

tolerance(Balbontín et al., 2024; Đurić et al., 2024; Khan et al., 2024; Wei et al., 2024), its

potential to induce transgenerational stress memory carried through metabolic or epigenetic

changes remains largely unexplored.

This study uniquely combines untargeted metabolomics with morpho-physiological

analyses to investigate whether MeJA priming imprints a stress memory that benefits

subsequent generations under recurrent drought conditions. It further explores the role of

drought induced memory and how MeJA and drought related memory interact to shape plant

performance under water stress. At the metabolomic level, we assessed changes in metabolite

diversity and composition of the entire metabolome. At the morpho-physiological level, we

evaluated specific leaf area, relative water content, leaf dry matter content, chlorophyll content,

and chlorophyll fluorescence. Finally, we investigated whether metabolomic profiles and plant

morpho-physiology are related, revealing predictive power from each other and indicating

integrated, memory driven strategies for enhanced stress resilience. We used *F. rubra*, a

dominant grass species of alpine grasslands, as a model. Thanks to its frequent vegetative

reproduction, this species is a perfect model for studying the transmission of stress memory

among clonal generations.

We hypothesize the following: **H1** – Exposure to current drought (D2) strongly

influence the metabolomic and morpho-physiological traits of *F. rubra*, representing immediate

stress induced changes. **H2**-Prior exposure to drought (D1) and methyl jasmonate (MeJA, M1)

induces transgenerational memory in clonal offspring, leading to persistent reprogramming of

the metabolome and modulation of key morpho-physiological traits such as specific leaf area

(SLA), relative water content (RWC), and chlorophyll dynamics thereby enhancing plant responses to recurrent drought stress even in the absence of current drought. **H3** – Drought and methyl jasmonate memories will interact with current drought such that combinations of past and present stress exposures will produce synergistic or compensatory effects on plant metabolome and morpho-physiology. **H4**- Finally, we hypothesize that metabolomic profiles and morphophysiological traits are tightly linked, reflecting coordinated memory-driven responses across molecular and phenotypic levels that enhance drought resilience.

2. Materials and Methods

2.1 Study system

We used ramets of *F. rubra* from our previous experiment (Bhatt et al. subm.) for this study. The previous experiment involved a factorial combination of drought (yes/no) and MeJA (yes/no) treatments and 10 replicates per treatment. *F. rubra* ramets were grown in pots with natural grassland soil under controlled conditions. Drought stress was applied in two phases, with recovery phases in between, while MeJA was applied to the soil during drought phases and not during the recovery phases. Control plants received regular watering. The MeJA soil application was chosen to ensure consistent distribution and minimize environmental loss. More details from the previous experiment are given in (Text S1).

In this experiment, ramets previously subjected to different drought and MeJA treatments (referred to as parental treatment) were grown in pots containing soil exposed to either drought or MeJA + drought in the previous experiment. Six replicates of ramets from the four parental treatments were grown in each soil with different memories (i.e., 2 parental drought treatments × 2 parental MeJA treatments × 2 soil memories × 2 current drought × 6 replicates). Although soil memory (SM) was included in the experimental design to account for potential legacy effects from the previous generation, it was not a primary focus of this

study. SM showed statistically significant effects only in the metabolomic composition

analyses and was not significant in any other aspect of the study. Consequently, SM is not

presented in the main text, and its results are presented only in (Table S1). We focus only on

intrinsic plant memory mechanisms, specifically current drought (D2), drought memory (D1),

and MeJA memory (M1).

Three of these replicates were well watered, and the remaining three were subjected to

recurrent drought cycles followed by recovery phases. This included 30 day cycles of drought

and recovery (Drought stage 1 - Recovery stage 1 - Drought stage 2 - Recovery stage 2). No

MeJA treatment was applied, as the focus was on investigating the legacy effects of prior MeJA

treatment on the metabolome of next-generation ramets under recurrent drought conditions.

The plants were grown in 200 ml pots. One *F. rubra* ramet was planted into each pot.

All planted ramets were of similar age and size (3 weeks old, 3.5-4 cm tall). They were grown

under long-day conditions (16 hours of daylight) in a growth chamber (17–20 °C Day and 7–

10 °C Night), reflecting peak growing season conditions at the original locality. The plants

were allowed to establish for 21 days before starting the experiment.

2.2 Plant performance

Plant physiology was characterized by chlorophyll content, chlorophyll fluorescence (Fv/Fm),

specific leaf area (SLA), leaf dry matter content (LDMC), and leaf relative water content

(RWC). These traits provide critical information about how plants structurally and

physiologically adapt to drought stress(Maxwell & Johnson, 2000; Perez-Harguindeguy et al.,

2013a).

Chlorophyll content and Chlorophyll fluorescence (Fv/Fm) were measured at the end

of each stress and recovery phase. SLA, LDMC, and RWC were measured only once, at the

end of the second recovery phase, due to their destructive nature, which prevented repeated

measurements.

2.2.1 Chlorophyll Content and Chlorophyll Fluorescence

Chlorophyll is an essential pigment crucial for photosynthesis, and its detection in plants is a

significant indicator of their chlorophyll fluorescence and overall well-being (Muller et al.,

2011). In our study, we utilized a chlorophyll CCM 300 meter to quantify the chlorophyll

content in plant leaves. Three randomly selected leaves were chosen from each pot, and the

chlorophyll content was measured according to the instructions provided by the manufacturer

once per leaf. The mean of these 3 values has been used as the dependent variable in the

subsequent analyses.

Chlorophyll degradation can decrease the maximum efficiency of Photosystem II (PSII)

known as the Fv/Fm ratios, decreasing the net photosynthetic rate (Chungloo et al., 2023).

Photon System Instrument FluorPen FP-100 MAX determined the chlorophyll fluorescence of

PSII. Healthy plants adapted to darkness have chlorophyll fluorescence ratios ranging from

0.75 to 0.85 (Björkman & Demmig, 1987). Lower values indicate damage to the system.

chlorophyll fluorescence was measured in the afternoon, and a 1-hour dark acclimation was

achieved by using the manufacturer provided clips. Chlorophyll content and chlorophyll

fluorescence were measured across four key phases: stress phase 1, recovery phase 1, stress

phase 2, and recovery phase 2. Based on the correlation analysis (Figure S1), two representative

variables were selected to reduce redundancy and multicollinearity: CCS2 (Chlorophyll

Content during Stress Phase 2) and CFR1 (Chlorophyll Fluorescence during Recovery Phase

1).

CCS2 showed strong correlations ($r \geq 0.7$) with other chlorophyll content traits from

stress phase 2 and both recovery phases, as well as with chlorophyll fluorescence traits from

both stress phases. This suggests that CCS2 effectively captures broader chlorophyll related

variation across phases. Similarly, CFR1 was highly correlated with CFR2 (Chlorophyll

Fluorescence during Recovery Phase 2), making it a suitable representative of the recovery-

phase fluorescence response.

2.2.2 SLA, LDMC, and RWC

SLA (Specific leaf area) and LDMC (Leaf dry matter content) are frequently used to predict

drought resistance because they essentially represent a trade-off between conservation and

rapid resource uptake (Blumenthal et al., 2020; Perez-Harguindeguy et al., 2013b). RWC

(Relative water content) serves as the most appropriate measure for assessing a plant's water

status, providing insight into cellular water deficit and the impact of Osmotic Adjustment on

maintaining cellular hydration during drought stress (Barrs & Weatherley, 1962).

Due to the extremely narrow and naturally folded morphology of *F. rubra* leaves,

accurately determining leaf area from scanned images proved unreliable. To standardize

measurements, we selected approximately 4 cm-long segments from individual leaves per pot.

Fresh weight was recorded for each segment. Because the leaves could not be fully unfolded

without damage, we measured both the length and width of each segment using a magnifying

glass. Leaf area was then estimated using the formula: length × width × 2, accounting for the

folding of the leaf blade. Subsequently, the leaf segment was placed in a Petri plate filled with

water overnight in a refrigerator to rehydrate. After the leaves had been rehydrated, any extra

water on the surface was carefully removed, and the leaves were weighed to determine their

turgid weight. The leaves were then dried in a 65°C oven for two days to determine their dry

weight. Using these measurements, we calculated the SLA by determining the ratio of leaf area

(mm^2) to dry mass (mg), LDMC by dividing the dry leaf mass (mg) by the turgid mass (g), and

Relative water content RWC as RWC (%) = [(Fresh weight - Dry weight) / (Turgid weight –

Dry weight)] x 100.

2.2.3 **Metabolite Profiling**

2.2.3.1 Plant sampling

Samples were collected at the end of the second drought phase. Approximately 20 leaves per

individual from multiple ramets per individual were sampled and pooled. Leaves were carefully

excised using sterile scissors, then wrapped in aluminium foil and immediately placed in liquid

nitrogen for snap freezing and temporary storage. After sampling all the plants, the samples

stored in liquid nitrogen were transferred to a -80°C freezer for storage until further analyses.

2.2.3.2 Methanol extraction

We took ~100 mg of each sample. We placed it into 1.5 ml microcentrifuge tubes (MCT)

containing 3 metal beads and 500 µL of an extraction solvent composed of methanol and water

(80:20). The samples were homogenized using a homogenizer. After homogenization, samples

were thoroughly vortexed for 10 seconds, followed by a 24-hour incubation at -20 °C. After

incubation, the tubes were vortexed again and centrifuged at 13000 rpm for 15 minutes. The

supernatant was carefully transferred into new tubes. The same steps of the extraction process,

from adding the solvent, were repeated once to ensure complete metabolite extraction. The

supernatants from both extraction rounds were pooled together. From this combined pool, 320

µL was transferred to a new tube, and 80 µL of a water/formic acid solution (99.9/0.1 v/v,

respectively) was added to the tube. The samples were then stored at -20 °C for at least 48

hours. Three blanks were processed alongside the samples to detect solvent-related

contamination, following all extraction steps but excluding leaf tissue. Additionally, quality

controls were created by pooling 20 µL from each sample extract.

2.2.3.3 Liquid Chromatography Data Acquisition

Chromatographic separation was performed at 40 °C using a 6546 LC/Q-TOF (Agilent) system

equipped with an Acclaim® RSLC 120 column (150 × 2.1 mm, 2.2 µm particle size; Thermo

Fisher Scientific). The chromatographic run began with an initial mobile-phase composition

of 95 % A, maintained for the first 3 minutes. From 0–3 minutes, the proportion of A was

linearly reduced to 82.7 %, then decreased to 76 % by the 10-minute mark. A steep linear

gradient followed between 10 and 17 minutes, dropping sharply from 76 % to 5 %, which was

held constant for one minute. A rapid re-equilibration occurred between 18 and 18.1 minutes,

raising A back to 95 %, and this composition was maintained through 20 minutes. The entire

sequence was executed for both positive and negative ionization modes. MS/MS analysis was

also performed in the same settings in a data-dependent mode, with 4 analyses conducted from

QC samples.

2.3.3.4 Spectral processing

Spectral processing of the LC-MS data was performed to extract relevant features for

metabolite identification and quantification. The raw data, which contained metabolites and

their concentrations across multiple time points, was processed using the MetaboAnalystR

Package of R (Pang et al., 2024). Several preprocessing steps were further applied to ensure

the quality and reliability of the acquired data. First, data were filtered to remove features with

more than 85% missing values, which helps eliminate unreliable or incomplete data. Missing

values were then imputed using the $1/5^{th}$ of the lowest value (Pang et al., 2024). This step is

essential to minimize the impact of missing values, which may arise due to biological

variability or technical limitations in detection sensitivity. Subsequently, the data were quantile

normalized, log transformed and scaled to stabilize variance and ensure comparable

distributions across samples. However, for estimating metabolite diversity metrics (richness,

Shannon diversity, and evenness), the processed data were deemed unsuitable due to

transformations that alter the original distribution of metabolite presence. Therefore, diversity

estimates were calculated using the unprocessed data to preserve the integrity of presence-

absence and abundance patterns.

2.3 Data analysis

All the metabolome analyses have been done separately for positive and negative ionization

modes as in previous studies (e.g. Calderón-Santiago et al., 2016; Nordström et al., 2008; Tian

et al., 2013).

2.3.1 Metabolite diversity and Uniqueness

To evaluate the effects of transgenerational memory of drought and Methyl Jasmonate (MeJA)

on metabolite diversity, we analyzed three key diversity indices, i.e., metabolite richness,

Shannon diversity, and Hill evenness (HillEven). Richness was estimated as the total number

of detected metabolites in each of the samples. Shannon diversity and HillEven were estimated

using the Chemodiv Package of R (Petrén et al., 2023). These indices were chosen because

they provide complementary insights into the metabolome: Shannon diversity captures both

the number and relative abundance of metabolites, reflecting metabolic complexity, while Hill

evenness quantifies how evenly metabolite abundances are distributed, which is critical for

understanding dominance patterns in untargeted metabolomics datasets. Additionally,

metabolomic uniqueness was evaluated by calculating the average Euclidean distance of each

sample from all others based on standardized metabolite intensities (Ren et al., 2015) .

Linear models were fitted using a full factorial design incorporating all treatments: D1, D2,

M1, and their two-way interactions. Before analyses, Richness, Shannon Diversity, and Hill

Evenness were log-transformed to fulfil the data normality assumption. Because the results of

Hill evenness largely matched the results of Shannon diversity in our analyses ($r = 0.96$), they

are not reported further. They are only shown in supporting information (Table S2 and Figure

S2, S3).

2.3.2 Multivariate analyses

Multivariate analyses were applied to examine the interactive effects of treatments and morpho-

physiological traits on the metabolomic composition. Redundancy Analysis (RDA) as

implemented in the vegan package in R was used throughout to explore these relationships,

using standardized data for all dependent variables (i.e., metabolites). A full model including

all main effects of D1, D2, M1, and their interactions was constructed. The significance of each

predictor was determined using permutation tests (n = 999), and variance partitioning was used

to quantify their contribution to overall variation. A similar RDA approach was used to

investigate how morpho-physiological traits influence metabolomic composition and how the

treatments affected plant morpho-physiological traits.

2.3.3 Effect of treatments on each metabolite

To visualize the overlap and specificity of metabolite responses across treatment groups, we

quantified the number of significantly affected metabolites (FDR-adjusted $p < 0.05$) for each

main effect (D1, M1, D2) and their pairwise interactions (D1×D2, D1×M1, D2×M1).

Significant metabolites were extracted from the linear model output, and treatment-specific

metabolite sets were defined. A binary presence/absence matrix was constructed across all

groups, enabling classification of each metabolite as either unique (significant in one group

only) or shared (significant in multiple groups). The resulting counts were visualized using bar

plots in R (version 4.2.1). To visualize changes in metabolite content, a heatmap was

constructed based on group-level averages of significant metabolites, with hierarchical

clustering. Also, due to the huge number of significant metabolites, only the top 100 were used

in the heatmap.

2.3.4 Annotation of significantly affected metabolites

We annotated the metabolites significantly affected by climate. For this, data-dependent LC-

MS/MS analysis was performed on a pooled sample (used as a quality control in LC-MS

analysis) to collect the necessary data. Subsequently, we used MS-DIAL (Tsugawa et al., 2015)

software to putatively annotate the metabolites detected in LC-MS/MS mode data. LC-MS/MS

data was processed by aligning retention times, detecting peaks, and deconvoluting spectra.

The annotation was then done by matching detected compounds against freely available

libraries at https://systemsomicslab.github.io/compms/msdial/main.html. A mass accuracy

threshold of 10 ppm was used in the annotation process. After annotation, the metabolites were

assigned to their chemical classes.

3 Results

3.1 Metabolite Diversity

The effect of current drought (D2) and its interaction with drought memory (D1) was

statistically significant (Table 1) in the positive ionization mode. Metabolite richness declined

under D2, indicating a narrowing of the metabolomic profile in response to stress. The

significant interaction between D1 and D2 suggests that the benefits of drought memory on

metabolomic diversification are conditional upon subsequent drought exposure. Specifically,

drought memory (D1) increased metabolite richness under non stress conditions, but had the

opposite effect under current drought, where richness was lowest in plants that had been

previously exposed to drought (Figure 1a).

For Shannon diversity, both D1 and M1 had statistically significant effects in the

positive ionization mode (Table 1), indicating that each memory treatment independently

affects metabolite diversity. D1 individually caused a decline in Shannon diversity, while M1

caused an increase. Their interaction (D1 × M1) was also significant (Table 1), indicating that

the effect of M1 on Shannon diversity depends on D1. Specifically, M1 caused an increase in

Shannon diversity in the absence of D1 memory, but the combined presence of D1 and M1

memories shaped Shannon diversity by negating the effects of each other and bringing the

Shannon diversity to control levels (both D1 and M1 are absent) (Figure 1b). In contrast, D2

did not significantly influence Shannon diversity in the positive mode but showed a strong

effect in the negative ionization mode (Table 1), where it significantly decreased Shannon

diversity, reflecting a more uneven distribution of metabolites under stress.

3.2 Metabolomic Uniqueness

The current drought (D2) was a significant predictor of metabolomic uniqueness in both

positive and negative ionization modes (Table 1), while M1 and D1 were not. A significant

interaction between D1 and D2 was also detected (Table 1), showing that metabolite uniqueness

declined under D1 alone but increased notably when drought memory plants were exposed to

current drought (Figure 1c). Also, the interaction between D2 and M1 was statistically

significant (Table 1). Plants without MeJa memory without current drought showed the lowest

uniqueness, plants without MeJa memory with current drought showed the highest uniqueness

and plants with MeJa memory, independent of current drought being in between (Figure 1d).

While the figures show results for the positive mode only, the results for the negative mode are

largely similar (Figure S4 a)

The triple interaction (D1 × D2 × M1) was significant for metabolic uniqueness in

negative mode. Plants with both drought and MeJa memory grown under drought exhibited the

highest metabolic uniqueness compared to all the other treatments (Figure S5).

3.3 Determinants of metabolome composition

D2 was the strongest predictor of metabolome composition in both positive and negative

ionization modes (Table 2). M1 and D1 (Table 2) also had significant and independent impacts

on metabolite profiles in both modes, though their effects were comparatively lower than D2.

The interactions D1×M1 and D2×M1 were also statistically significant (Table 2), indicating

that the combination of stress memory with current drought or between both memory types

altered metabolite composition. In contrast, the interaction between D1 and D2 did not show a

significant effect in either mode.

3.4 Effect of MeJA memory and its interaction with drought on specific metabolites

In the positive ionization mode, D2 induced the most significant metabolic changes, uniquely

affecting 475 metabolites and sharing 935 with other treatments, underscoring its central role

in metabolomic reprogramming (Figure S7a). M1 had the second largest impact, with 90

unique metabolites and 389 shared, while D1 had the smallest individual effect with 19 unique,

249 shared metabolites. Interaction terms also demonstrated pronounced metabolic responses.

The D2 × M1 interaction influenced the most metabolites among combinations (39 unique, 205

shared), followed by D1 × M1 (19 unique, 192 shared), while D1 × D2 showed minimal effect

(0 unique, 1 shared).

In the negative ionization mode (Figure S7b), D2 again triggered the strongest

metabolic shifts (462 unique, 952 shared), with M1 (73 unique, 396 shared) and D1 (30 unique,

311 shared) trailing behind. Similarly, interactions revealed substantial effects: D2 × M1 (54

unique, 170 shared), D1 × M1 (28 unique, 245 shared), and again, D1 × D2 (0 unique, 1 shared).

These findings highlight D2 as the primary driver of metabolic reprogramming, while

memory effects from M1 and D1 also shape distinct biochemical landscapes. The notably high

number of unique metabolites in D2 × M1 and D1 × M1 interactions indicates a non-additive,

context-dependent integration of stress memory with ongoing drought stress.

The heatmap of the top 200 differentially accumulated under different conditions and

annotated metabolites in both positive and negative mode (Figure 3). Clusters 1-4 in the

positive ion mode heatmap illustrate distinct metabolic strategies shaped by MeJA memory in

F. rubra. Cluster 1 metabolites, including phenolic compounds, flavonoids, curcuminoids,

saccharolipids, and linear diarylheptanoids, are upregulated in M1D2-treated plants, revealing

a jasmonic acid (JA) specific metabolic imprint, while D2, D1D2, and M1D1D2 treatments

show downregulation, suggesting that drought memory alone or in combination interferes with

this JA-associated response. Cluster 2 metabolites are selectively upregulated in both M1 and

M1D2, including alkaloids, terpenoids, isoflavonoids, lignans, and oligopeptides, indicating

that this set is a robust MeJA memory marker that does not require current stress for activation.

This exclusive response in MeJA primed plants makes Cluster 2 a clear biomarker for JA-

driven metabolic readiness. In contrast, Cluster 3 metabolites such as phosphocholines,

salicylamides, diacylglycerols, ether lipids, and flavonoid derivatives are selectively

downregulated only in M1 and M1D2, whereas all other treatments, regardless of memory,

show upregulation. This suggests that MeJA priming deliberately suppresses this non-primed,

general stress response pathway, likely to conserve energy and direct metabolism toward more

specialized defenses. Lastly, Cluster 4 is uniquely downregulated in M1D2, whereas D2,

D1D2, and M1D1D2 exhibit strong upregulation of their lipid and redox related metabolites

including ceramides (Cer_NDS, CerP), fatty alcohol esters, gamma amino acids, and

pyridoxines. This pattern indicates that MeJA memory strategically suppresses broad stress

induced metabolic activation in favor of targeted defense, reinforcing its role in refining and

focusing the plant's metabolic response under recurring drought conditions.

In negative ion mode, Cluster 1 is selectively upregulated in M1, and M1D2 includes

metabolites such as flavonoids, coumarins, limonoids, glycerophospholipids, oligosaccharides,

and rotanones, establishing it as a MeJA memory-specific metabolic signature. In contrast, D2,

D1D2, and M1D1D2 show clear suppression of this cluster, indicating that drought memory

alone or even in combination with MeJA fails to activate, and may actively repress, these

pathways. This underscores the specificity and strength of MeJA driven priming in shaping

targeted metabolic defense readiness. Conversely, Cluster 2 displays the opposite trend M1D2

exhibits strong downregulation of broad stress response metabolites such as alkaloids,

phosphatidic acid, alpha and N-acyl amino acids, methoxyphenols, and dihexosylceramides,

while D2, D1D2, and M1D1D2 show robust upregulation. This contrast highlights a MeJA

memory induced suppression of diffuse metabolic activation, reinforcing a strategic shift in *F.*

rubra toward focused, energy efficient defense responses during drought stress.

3.5 Relationship between plant morpho-physiological traits and metabolite profiles

394 The redundancy analysis (RDA) shows that SLA and RWC were the plant traits having the

395 strongest effects on metabolite profiles under both positive and negative ionization modes

396 (Table 3). Chlorophyll content (CCS2) also contributed to explaining metabolomic variation,

397 but to a lesser extent and LDMC had no significant effect on metabolite patterns.

3.6 Determinants of plant morpho-physiological traits

The redundancy analysis (RDA) showed that M1 was the strongest predictor of morpho-

physiological traits in *F. rubra,* followed by D2 and D1 (Table 2). The interaction between D2

and M1 was also statistically significant, while other interactions (D1×D2 and D1×M1) were

not significant. The triple interaction between D1×D2×M1 was also significant (Table 2). This

suggests that the way D1 affects dependence on M1 changes depending on whether the plants

are experiencing drought (D2).

LDMC, RWC, and CCS2 increased with M1 and D2×M1, indicating positive

associations with these treatments. In contrast, SLA pointed in the opposite direction,

suggesting a likely trade-off under those same treatments. CFR1 shows a positive association

with M1, as both vectors point in the same leftward direction on the RDA biplot. This suggests

higher CFR1 values under M1 treatment (Figure 2).

4. Discussion

This study provides one of the first comprehensive multi-trait and metabolomic analyses of

transgenerational plant memory in response to drought and methyl jasmonate (MeJA) priming

and interactions among different memories. Using *F. rubra* as a clonal model system, it reveals

how recurrent drought (D2), drought memory (D1), and MeJA memory (M1) individually and

interactively modulate metabolomic diversity and composition, along with morpho-

physiological traits. A key novelty lies in the demonstration that memory effects persist and

influence not only morphological adjustments but also the plant's systemic metabolic

reprogramming. The study uniquely identifies that memory effects are interaction dependent,

revealing conditional metabolomic shifts shaped by environmental context. It also shows that

MeJA memory can compensate for or stabilize drought induced metabolic alterations,

suggesting a promising tool for enhancing resilience in clonal or perennial species. This work

advances understanding of plant stress memory mechanisms, particularly highlighting

metabolomic memory as a transgenerational trait. We also demonstrated plant responses at

morpho-physiological levels and how strongly the metabolomic and morpho-physiological

changes relate to each other.

4.1 Current drought (D2) is the most important factor affecting metabolomic profiles and

morpho-physiology

Our results strongly support Hypothesis 1, demonstrating that current drought (D2) is the most

influential factor affecting both the metabolomic architecture and morpho-physiological traits

of *F. rubra*. This aligns with previous studies showing that plants facing acute water deficit

rapidly engage systemic metabolic and structural adjustments to maintain cellular integrity and

ensure survival (Fàbregas & Fernie, 2019; Gupta et al., 2020). However, our study extends this

link to the whole metabolome, showing that cellular processes are strongly affected by drought.

A key finding is the reduced metabolite richness under current drought (D2) in positive

ionization mode, suggesting a metabolic streamlining strategy. Rather than initiating a broad

response, plants appear to conserve resources by prioritizing metabolites critical for drought

tolerance, supporting the idea of targeted metabolic adjustment under stress. In contrast, the

negative ionization mode showed stable richness but increased Shannon diversity, indicating a

shift toward even metabolite distribution in negative mode. This pattern likely reflects the

prioritization of compounds such as lipids, amino acids, phenolics, lactones, and terpenoids,

which are metabolites known for their roles in membrane stability, antioxidant defense,

osmoprotection, and signaling under drought (Bowne et al., 2012; Fàbregas & Fernie, 2019).

We also observed a marked increase in metabolite uniqueness under drought,

suggesting individualized metabolic trajectories. This aligns with findings in *Arabidopsis*

thaliana and maize, where genotypes and tissues exhibited distinct metabolic responses to

drought (Lozano-Elena et al., 2022)highlighting metabolic plasticity as a key resilience

mechanism.

Morpho-physiological traits also responded strongly to drought. Notably, specific leaf

area (SLA) and relative water content (RWC) were closely associated with D2 conditions. A

higher SLA under drought reflects a strategy to maximize light capture with reduced carbon

cost, although it comes with trade-offs, namely, thinner leaves with lower water retention and

photosynthetic capacity (Gonzalez-Paleo & Ravetta, 2018; Sun et al., 2025; Zhou et al.,

2020).These changes underscore that drought resilience involves adjustments at both

metabolomic and morpho-physiological levels.

4.2 Transgenerational memory reprograms plant metabolome and morpho-physiology

A key finding of our study is the context-dependent manner in which D1 and M1 affect

metabolomic richness and Shannon diversity. Both D1 and M1 significantly increased

metabolite diversity under non-stressed conditions, reflecting a primed state where plants maintain a wider array of metabolites in anticipation of future drought stress. This aligns with earlier findings that priming via prior stress exposure or hormone signaling can enhance metabolic readiness and resilience(Hilker & Schmülling, 2019a; Liu et al., 2022a).

The independent effect of M1 uncovered in this study highlights its powerful role in transgenerational metabolomic and morpho-physiological reprogramming. Its effects on metabolomic composition and morpho-physiological traits were stronger than the effect of drought memory (D1). Importantly, M1 significantly enhanced metabolomic Shannon diversity without increasing richness. This suggests that M1 enhances not the quantity, but the balance and distribution of metabolites, possibly preparing the plant for a more versatile response to various stressors. This aligns with the known role of jasmonates in mediating systemic defense signaling and metabolic priming (Avramova, 2019). MeJA memory plants also maintained higher relative water content (RWC) and chlorophyll content under stress (CCS2), reflecting enhanced drought preparedness through improved water retention and photosynthetic integrity (Maxwell & Johnson, 2000; Qiu et al., 2020). To our knowledge, no previous study has explored MeJa memory in non-model, clonally reproducing species integrating untargeted metabolome profiling and morpho-physiology. By linking MeJA memory to stable changes in metabolic architecture and drought relevant traits, this work provides a novel framework for understanding non-genetic inheritance of stress resilience and offers practical insights for enhancing plant performance in increasingly unpredictable climates (Latzel & Münzbergová, 2018; Liu et al., 2022b).

Our study shows that drought memory (D1) alone maintained high metabolite richness, comparable to control plants, consistent with prior studies showing that stress primed plants often exhibit enhanced metabolic capacity due to epigenetic remodeling of stress responsive pathways (Nguyen et al., 2022). Drought memory (D1) also played a pivotal role in

reprogramming plant morpho-physiology. In D1 plants, we observed a marked increase in

specific leaf area (SLA), indicative of a shift toward a resource acquisitive strategy. This

morphological adjustment, however, was accompanied by lower relative water content (RWC)

and reduced leaf structural density, suggesting a trade-off between growth efficiency and water

conservation(Blumenthal et al., 2020; Pérez-Ramos et al., 2013). The fact that drought memory

(D1) triggers plant responses even in the absence of current drought suggests it acts as a priming

mechanism, keeping plants in a preconditioned state(Crisp et al., 2016; Fleta-Soriano &

Munné-Bosch, 2016). This form of anticipatory plasticity allows plants to respond more

quickly and efficiently when drought returns. In unpredictable environments, such stress

memory enhances survival by proactively shaping plant metabolism and morphology, rather

than relying solely on reactive responses(Liu et al., 2022b).

Crucially, drought memory (D1) not only influenced morpho-physiological traits and

metabolomic diversity but also altered metabolome composition, supporting the idea that plants

retain a biochemical memory of past stress(Sharma et al., 2022). This memory appears to guide

transgenerational metabolomic strategies, exerting effects on both structural and biochemical

components. While earlier studies have shown that drought stressed plants can transmit

morpho-physiological traits to clonal offspring(Hilker & Schmülling, 2019b; Kambona et al.,

2023b; Latzel & Münzbergová, 2018; Liu et al., 2022b), our study is the first to extend these

inherited traits to metabolic reprogramming, demonstrating that drought memory modulates

both structural and biochemical systems across generations.

4.3 Interactive effects of MeJA memory, drought memory, and current drought shape

plant responses

In line with H3, the different types of memory interacted with the current drought. Specifically,

drought memory (D1) increased metabolomic richness in plants not exposed to recurrent

drought (D2), and reduced it in plants exposed to current drought. This suggests that drought

memory is not expressed uniformly but is modulated by current environmental conditions.

Under recurrent drought, drought memory may streamline the metabolic response, focusing on

a core set of drought tolerance metabolites, thereby reducing overall richness. In contrast, in

the absence of current drought, drought memory appears to promote a broader metabolic

profile, possibly as a generalized preparedness strategy. This finding extends previous work in

Leymus chinensis and rice, which showed that drought memory alters physiological states

(Kambona et al., 2023b), by demonstrating that it also shapes whole metabolome richness in a

context-dependent manner.

Similarly, MeJA induced memory (M1) influenced metabolism through interactions

with D1 and D2, highlighting its context-sensitive regulatory role. These interactions reinforce

the idea that MeJA establishes a stable and heritable priming state, enhancing the plant's

capacity to respond to future stress. While MeJA is known to modulate antioxidant pathways,

secondary metabolism, and hormonal signaling (Chungloo et al., 2023; Hewedy, Elsheery,

Karkour, Elhamouly, Arafa, Mahmoud, Dawood, Hussein, Mansour, & Amin, 2023; Tayyab et

al., 2020; Yu et al., 2018), our study is the first to demonstrate that MeJA priming induces

transgenerational metabolic memory, affecting both metabolomic diversity and morpho-

physiological traits, as well as interactions with other types of memory.

The D1 × M1 interaction revealed that MeJA memory restored metabolomic diversity

in plants with drought memory to levels comparable to controls, suggesting a compensatory

mechanism. This implies that MeJA memory may reactivate or stabilize biosynthetic pathways

that are otherwise suppressed by drought memory. Such a capacity to "reset" the metabolic

state across generations positions MeJA priming as a promising strategy to mitigate the long-

term effects of drought, particularly in clonal and perennial species. While prior studies report

individual effects of memory in plants (Aswathi et al., 2025), the interaction among memory

types is rarely studied (Aswathi et al., 2025; Kambona et al., 2023b).

Another key and novel finding of this study is a significant three way interaction

between drought memory (D1), current drought (D2), and MeJA memory (M1), which strongly

influenced metabolomic uniqueness. This interaction reveals that multiple stress memories do

not act independently or additively, but instead interact in complex, emergent ways to shape

the plant metabolome. The resulting metabolomic profiles were distinct and variable among all

treatment groups, suggesting that plants integrate diverse stress legacies to generate unique,

context-specific metabolic responses. To our knowledge, this is the first study to demonstrate

such a three way interaction affecting the plant metabolome, highlighting a previously

unrecognized layer of complexity in stress memory research. These findings underscore the

importance of considering multi-factorial and transgenerational stress interactions when

studying plant adaptation. Our results also point to the potential of leveraging combined

priming strategies such as MeJA application alongside drought conditioning to enhance

adaptive plasticity and biochemical resilience in clonal species under increasingly

unpredictable environmental conditions.

4.4 Metabolome and physiology relate strongly to each other

In line with Hypothesis 4 (H4), we found that metabolomic profiles and morpho-physiological

traits are tightly linked. Redundancy analysis demonstrated that SLA and RWC were the

strongest predictors of metabolomic variation, followed by Chlorophyll Content (CCS2). These

traits, which are key indicators of plant water use strategy and photosynthetic function(Barrs

& Weatherley, 1962; Blumenthal et al., 2020; Maxwell & Johnson, 2000; Pérez-Ramos et al.,

2013) explained substantial portions of the variation in both positive and negative ionization

modes. The tight coupling between these physiological traits and metabolome composition

strongly supports the perception that plants coordinate biochemical and phenotypic traits to

adapt to both current drought and stress memory.

Cluster heatmap analysis revealed distinct regulation patterns among memory

treatments, showing how specific metabolite families support morpho-physiological resilience.

In positive ionization mode, Cluster 1 (Figure S8a)—including phenolics, flavonoids, lipids,

and curcuminoids—was strongly upregulated only in M1D2. These compounds, known for

their antioxidant, membrane stabilizing, and osmoprotective roles (Almeida et al., 2020b;

Fàbregas & Fernie, 2019; Falcone Ferreyra et al., 2012) , help maintain high relative water

content (RWC) and reduce oxidative stress. This aligns with lower specific leaf area (SLA) and

greater chlorophyll retention in MeJA memory plants, indicating efficient structural and

biochemical defense. Cluster 2 (Figure S8b), enriched in alkaloids, lignans, oligopeptides,

amino acids, and terpenoids, was activated in M1 and M1D2 signatures of jasmonate regulated

metabolism that preemptively bolster stress tolerance via ROS scavenging, structural

fortification, and signaling (Avramova, 2019; Khan et al., 2024; Yu et al., 2018) . In contrast,

Clusters 3 and 4 (Figures S8c–d), upregulated in non-MeJA treatments (D2, D1D2, M1D1D2)

but suppressed in M1D2, contained phosphocholines, salicylamides, ceramides, pyridoxines,

and other broad stress related metabolites. These are energetically costly and potentially

disruptive to specific JA responses; their repression in M1D2 suggests metabolic economy and

response specificity (Crisp et al., 2016; Liu et al., 2022b; Salvi et al., 2021b; Sharma et al.,

2022).

In negative ionization mode, Cluster 1 (Figure S8e), selectively upregulated in M1D2,

comprised flavonoids, coumarins, phosphatidic acids, macrolides, and limonoids—key agents

in membrane repair, antioxidative buffering, and defense mobilization (Chungloo et al., 2023;

You et al., 2019; Zhang et al., 2023) . This activation strengthens the link between metabolomic

priming and stable RWC and photosynthesis. Conversely, Cluster 2 (Figure S8f), with amino

acid derivatives, purines, lipid messengers, and phenylpropanes, was elevated in D2, D1D2, and M1D1D2 but repressed in M1D2, indicating avoidance of energetically expensive signaling in MeJA memory plants (Aswathi et al., 2025; Hilker & Schmülling, 2019b). Altogether, the data show that MeJA memory orchestrates the upregulation of protective metabolites and suppression of redundant or costly pathways, tightly linking metabolomic profiles with enhanced physiological outcomes highlighting the mechanistic basis of drought resilience in clonal plants (Aina et al., 2024; Balbontín et al., 2024).

5 Conclusion

Our study demonstrates that both drought memory and methyl jasmonate-induced memory play critical roles in shaping plant responses to current drought, not only through individual effects but also via complex interactions. We show for the first time that these memories influence whole metabolome diversity and uniqueness in a context-dependent and transgenerational manner. Drought memory appears to streamline metabolic responses under recurring drought, while methyl jasmonate induced memory fine tunes and stabilizes them, even compensating for the legacy of past stress. These memory driven adjustments extend beyond metabolism to influence morpho-physiological traits, enhancing water retention and structural resilience.

Importantly, our findings highlight that stress memory is not static but dynamically expressed depending on environmental context. The ability of methyl jasmonate induced memory to restore diversity and promote protective traits across generations underscores its potential as a biotechnological tool for enhancing resilience in clonally reproducing perennial species. Together, these insights advance our understanding of plant stress memory as a multi-layered, heritable system, offering promising avenues for developing crops and wild species better adapted to increasingly variable climates.

Conflict of interest: Nothing to declare

Acknowledgements: GAUK project number 171724 supported the study. It was also partly

supported by GAČR 22-00761S, a long-term research development project no. RVO 67985939

of the Czech Academy of Sciences and institutional support for science and research of the

Ministry of Education, Youth, and Sports of the Czech Republic.

Author contributions: TB, DT, and ZM conceptualized the study; TB, ZM, and DT did the

experimental setup; NR and TB meticulously prepared the samples for UPLC analysis, while

NR, DT, and JS optimized the UPLC-MS method; JS and TC performed UPLC-MS analysis;

The raw data were processed to obtain metabolite feature data by TB, DT, and JS; TB carried

out data analysis with the help of DT and ZM. TB led the writing, with in-depth editing and

suggestions from DT and ZM; DT and ZM are the senior authors; All authors approved the

submission and declare no conflict of interest.

References

Aina, O., Bakare, O. O., Fadaka, A. O., Keyster, M., & Klein, A. (2024). Plant biomarkers as
early detection tools in stress management in food crops: a review. *Planta*, *259*(3), 60.
https://doi.org/10.1007/s00425-024-04333-1

Almeida, T., Pinto, G., Correia, B., Gonçalves, S., Meijón, M., & Escandón, M. (2020a). In-
depth analysis of the Quercus suber metabolome under drought stress and recovery
reveals potential key metabolic players. *Plant Science*, *299*, 110606.
https://doi.org/https://doi.org/10.1016/j.plantsci.2020.110606

Almeida, T., Pinto, G., Correia, B., Gonçalves, S., Meijón, M., & Escandón, M. (2020b). In-
depth analysis of the Quercus suber metabolome under drought stress and recovery
reveals potential key metabolic players. *Plant Science*, *299*, 110606.
https://doi.org/https://doi.org/10.1016/j.plantsci.2020.110606

Aswathi, K. P. R., Ul-Allah, S., Puthur, J. T., Siddique, K. H. M., Frei, M., & Farooq, M.
(2025). The Plant Mind: Unraveling Abiotic Stress Priming, Memory, and Adaptation.
Physiologia Plantarum, *177*(4), e70372.

Avramova, Z. (2019). Defence-related priming and responses to recurring drought: Two
manifestations of plant transcriptional memory mediated by the ABA and JA signalling
pathways. *Plant, Cell & Environment*, *42*(3), 983–997.

Balbontín, C., Reyes, M., Yáñez, M. A., Parra-Palma, C., Morales-Quintana, L., & Ramos, P.
(2024). Enhancing blueberry drought resilience: ABA and MeJA hormonal formulations

unveil water-saving strategies. *Horticulture, Environment, and Biotechnology*. https://doi.org/10.1007/s13580-024-00640-4

Barrs, H. D., & Weatherley, P. E. (1962). A re-examination of the relative turgidity technique for estimating water deficits in leaves. *Australian Journal of Biological Sciences*, *15*(3), 413–428.

Björkman, O., & Demmig, B. (1987). Photon yield of O 2 evolution and chlorophyll fluorescence characteristics at 77 K among vascular plants of diverse origins. *Planta*, *170*, 489–504.

Blumenthal, D. M., Mueller, K. E., Kray, J. A., Ocheltree, T. W., Augustine, D. J., & Wilcox, K. R. (2020). Traits link drought resistance with herbivore defence and plant economics in semi-arid grasslands: The central roles of phenology and leaf dry matter content. *Journal of Ecology*, *108*(6), 2336–2351.

Bowne, J. B., Erwin, T. A., Juttner, J., Schnurbusch, T., Langridge, P., Bacic, A., & Roessner, U. (2012). Drought responses of leaf tissues from wheat cultivars of differing drought tolerance at the metabolite level. *Molecular Plant*, *5*(2), 418–429.

Calderón-Santiago, M., Fernández-Peralbo, M. A., Priego-Capote, F., & Luque de Castro, M. D. (2016). MSCombine: a tool for merging untargeted metabolomic data from high-resolution mass spectrometry in the positive and negative ionization modes. *Metabolomics*, *12*(3), 43. https://doi.org/10.1007/s11306-016-0970-4

Calleja-Satrustegui, A., Echeverría, A., Ariz, I., Peralta de Andrés, J., & González, E. M. (2024). Unlocking nature's drought resilience: a focus on the parsimonious root phenotype and specialised root metabolism in wild Medicago populations. *Plant and Soil*, 1–17.

Chungloo, D., Tisarum, R., Sotesaritkul, T., Praseartkul, P., Himanshu, S. K., Datta, A., & Cha-um, S. (2023). Exogenous Foliar Application of Methyl Jasmonate Alleviates Water-Deficit Stress in Andrographis paniculata. *Journal of Soil Science and Plant Nutrition*. https://doi.org/10.1007/s42729-023-01414-0

Crisp, P. A., Ganguly, D., Eichten, S. R., Borevitz, J. O., & Pogson, B. J. (2016). Reconsidering plant memory: Intersections between stress recovery, RNA turnover, and epigenetics. *Science Advances*, *2*(2), e1501340.

Đurić, M., Motyka, V., Dobrev, P. I., Nedvěd, D., Pokorná, E., Subotić, A., Trifunović-Momčilov, M., & Milošević, S. (2024). Changes in the Endogenous Phytohormone Status of Drought-Stressed Impatiens walleriana Hook. f. Plants Treated with Methyl Jasmonate. *Journal of Plant Growth Regulation*, 1–18.

Fàbregas, N., & Fernie, A. R. (2019). The metabolic response to drought. *Journal of Experimental Botany*, *70*(4), 1077–1085. https://doi.org/10.1093/jxb/ery437

Falcone Ferreyra, M. L., Rius, S. P., & Casati, P. (2012). Flavonoids: biosynthesis, biological functions, and biotechnological applications. *Frontiers in Plant Science*, *3*, 222.

Fiehn, O. (2002). Metabolomics—the link between genotypes and phenotypes. *Functional Genomics*, 155–171.

Fleta-Soriano, E., & Munné-Bosch, S. (2016). Stress memory and the inevitable effects of
drought: a physiological perspective. *Frontiers in Plant Science*, *7*, 143.

Gonzalez-Paleo, L., & Ravetta, D. A. (2018). Relationship between photosynthetic rate, water
use and leaf structure in desert annual and perennial forbs differing in their growth.
Photosynthetica, *56*(4), 1177–1187. https://doi.org/10.1007/s11099-018-0810-z

Gupta, A., Rico-Medina, A., & Caño-Delgado, A. I. (2020). The physiology of plant
responses to drought. *Science*, *368*(6488), 266–269.
https://doi.org/10.1126/science.aaz7614

Hewedy, O. A., Elsheery, N. I., Karkour, A. M., Elhamouly, N., Arafa, R. A., Mahmoud, G.
A.-E., Dawood, M. F.-A., Hussein, W. E., Mansour, A., & Amin, D. H. (2023). Jasmonic
acid regulates plant development and orchestrates stress response during tough times.
Environmental and Experimental Botany, *208*, 105260.

Hewedy, O. A., Elsheery, N. I., Karkour, A. M., Elhamouly, N., Arafa, R. A., Mahmoud, G.
A.-E., Dawood, M. F.-A., Hussein, W. E., Mansour, A., Amin, D. H., Allakhverdiev, S.
I., Zivcak, M., & Brestic, M. (2023). Jasmonic acid regulates plant development and
orchestrates stress response during tough times. *Environmental and Experimental
Botany*, *208*, 105260. https://doi.org/https://doi.org/10.1016/j.envexpbot.2023.105260

Hilker, M., & Schmülling, T. (2019a). Stress priming, memory, and signalling in plants.
Plant, Cell & Environment, *42*(3), 753–761.
https://doi.org/https://doi.org/10.1111/pce.13526

Hilker, M., & Schmülling, T. (2019b). Stress priming, memory, and signalling in plants. In
Plant, Cell & Environment (Vol. 42, Issue 3, pp. 753–761). Wiley Online Library.

Kambona, C. M., Koua, P. A., Léon, J., & Ballvora, A. (2023a). Stress memory and its
regulation in plants experiencing recurrent drought conditions. *Theoretical and Applied
Genetics*, *136*(2), 26.

Kambona, C. M., Koua, P. A., Léon, J., & Ballvora, A. (2023b). Stress memory and its
regulation in plants experiencing recurrent drought conditions. *Theoretical and Applied
Genetics*, *136*(2), 26. https://doi.org/10.1007/s00122-023-04313-1

Khan, V., Princi, Mubashshir, M., Umar, S., & Iqbal, N. (2024). Methyl Jasmonate and
Pseudomonas fluorescens Synergistically Boost Antioxidative Defense, Secondary
Metabolites, and Osmolyte Production to Enhance Drought Resilience in Mustard.
Journal of Plant Growth Regulation. https://doi.org/10.1007/s00344-024-11310-1

Kumar, S., Sachdeva, S., Bhat, K. V, & Vats, S. (2018). Plant responses to drought stress:
physiological, biochemical and molecular basis. *Biotic and Abiotic Stress Tolerance in
Plants*, 1–25.

Latzel, V., & Münzbergová, Z. (2018). Anticipatory Behavior of the Clonal Plant Fragaria
vesca. *Frontiers in Plant Science*, *9*.
https://www.frontiersin.org/articles/10.3389/fpls.2018.01847

Lee, S. H., & Zwiazek, J. J. (2019). Regulation of water transport in Arabidopsis by methyl
jasmonate. *Plant Physiology and Biochemistry*, *139*, 540–547.
https://doi.org/https://doi.org/10.1016/j.plaphy.2019.04.023

Liu, H., Able, A. J., & Able, J. A. (2022a). Priming crops for the future: rewiring stress
memory. *Trends in Plant Science*, *27*(7), 699–716.
https://doi.org/https://doi.org/10.1016/j.tplants.2021.11.015

Liu, H., Able, A. J., & Able, J. A. (2022b). Priming crops for the future: rewiring stress
memory. *Trends in Plant Science*, *27*(7), 699–716.

Lozano-Elena, F., Fàbregas, N., Coleto-Alcudia, V., & Caño-Delgado, A. I. (2022). Analysis
of metabolic dynamics during drought stress in Arabidopsis plants. *Scientific Data*, *9*(1),
90. https://doi.org/10.1038/s41597-022-01161-4

Maxwell, K., & Johnson, G. N. (2000). Chlorophyll fluorescence—a practical guide. *Journal
of Experimental Botany*, *51*(345), 659–668.

Michaletti, A., Naghavi, M. R., Toorchi, M., Zolla, L., & Rinalducci, S. (2018).
Metabolomics and proteomics reveal drought-stress responses of leaf tissues from
spring-wheat. *Scientific Reports*, *8*(1), 5710.

Nguyen, N. H., Vu, N. T., & Cheong, J.-J. (2022). Transcriptional stress memory and
transgenerational inheritance of drought tolerance in plants. *International Journal of
Molecular Sciences*, *23*(21), 12918.

Nordström, A., Want, E., Northen, T., Lehtiö, J., & Siuzdak, G. (2008). Multiple Ionization
Mass Spectrometry Strategy Used To Reveal the Complexity of Metabolomics.
Analytical Chemistry, *80*(2), 421–429. https://doi.org/10.1021/ac701982e

Pang, Z., Lu, Y., Zhou, G., Hui, F., Xu, L., Viau, C., Spigelman, A. F., MacDonald, P. E.,
Wishart, D. S., Li, S., & Xia, J. (2024). MetaboAnalyst 6.0: towards a unified platform
for metabolomics data processing, analysis and interpretation. *Nucleic Acids Research*,
52(W1), W398–W406. https://doi.org/10.1093/nar/gkae253

Perez-Harguindeguy, N., Diaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P.,
Bret-Harte, M. S., Cornwell, W. K., Craine, J. M., & Gurvich, D. E. (2013a). *New
handbook for standardised measurement of plant functional traits worldwide. Aust. Bot.
61, 167–234.*

Perez-Harguindeguy, N., Diaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P.,
Bret-Harte, M. S., Cornwell, W. K., Craine, J. M., & Gurvich, D. E. (2013b). *New
handbook for standardised measurement of plant functional traits worldwide. Aust. Bot.
61, 167–234.*

Pérez-Ramos, I. M., Volaire, F., Fattet, M., Blanchard, A., & Roumet, C. (2013). Tradeoffs
between functional strategies for resource-use and drought-survival in Mediterranean
rangeland species. *Environmental and Experimental Botany*, *87*, 126–136.

Petrén, H., Köllner, T. G., & Junker, R. R. (2023). Quantifying chemodiversity considering
biochemical and structural properties of compounds with the R package chemodiv. *New
Phytologist*, *237*(6), 2478–2492. https://doi.org/https://doi.org/10.1111/nph.18685

Ploughe, L. W., Jacobs, E. M., Frank, G. S., Greenler, S. M., Smith, M. D., & Dukes, J. S.
(2019). Community Response to Extreme Drought (CRED): a framework for drought-
induced shifts in plant–plant interactions. *New Phytologist*, *222*(1), 52–69.

Ren, S., Hinzman, A. A., Kang, E. L., Szczesniak, R. D., & Lu, L. J. (2015). Computational
and statistical analysis of metabolomics data. *Metabolomics*, *11*(6), 1492–1513.
https://doi.org/10.1007/s11306-015-0823-6

Salvi, P., Manna, M., Kaur, H., Thakur, T., Gandass, N., Bhatt, D., & Muthamilarasan, M.
(2021a). Phytohormone signaling and crosstalk in regulating drought stress response in
plants. *Plant Cell Reports*, *40*(8), 1305–1329. https://doi.org/10.1007/s00299-021-
02683-8

Salvi, P., Manna, M., Kaur, H., Thakur, T., Gandass, N., Bhatt, D., & Muthamilarasan, M.
(2021b). Phytohormone signaling and crosstalk in regulating drought stress response in
plants. *Plant Cell Reports*, *40*, 1305–1329.

Sharma, M., Kumar, P., Verma, V., Sharma, R., Bhargava, B., & Irfan, M. (2022).
Understanding plant stress memory response for abiotic stress resilience: Molecular
insights and prospects. *Plant Physiology and Biochemistry*, *179*, 10–24.
https://doi.org/https://doi.org/10.1016/j.plaphy.2022.03.004

Sun, D., Lei, Z., Carriquí, M., Zhang, Y., Liu, T., Wang, S., Song, K., Zhu, L., Zhang, W., &
Zhang, Y. (2025). Reductions in mesophyll conductance under drought stress are
influenced by increases in cell wall chelator-soluble pectin content and denser
microfibril alignment in cotton. *Journal of Experimental Botany*, *76*(4), 1116–1130.

Tayyab, N., Naz, R., Yasmin, H., Nosheen, A., Keyani, R., Sajjad, M., Hassan, M. N., &
Roberts, T. H. (2020). Combined seed and foliar pre-treatments with exogenous methyl
jasmonate and salicylic acid mitigate drought-induced stress in maize. *PLOS ONE*,
15(5), e0232269-. https://doi.org/10.1371/journal.pone.0232269

Tian, H., Bai, J., An, Z., Chen, Y., Zhang, R., He, J., Bi, X., Song, Y., & Abliz, Z. (2013).
Plasma metabolome analysis by integrated ionization rapid-resolution liquid
chromatography/tandem mass spectrometry. *Rapid Communications in Mass
Spectrometry*, *27*(18), 2071–2080. https://doi.org/https://doi.org/10.1002/rcm.6666

Tsugawa, H., Cajka, T., Kind, T., Ma, Y., Higgins, B., Ikeda, K., Kanazawa, M.,
VanderGheynst, J., Fiehn, O., & Arita, M. (2015). MS-DIAL: data-independent MS/MS
deconvolution for comprehensive metabolome analysis. *Nature Methods*, *12*(6), 523–
526. https://doi.org/10.1038/nmeth.3393

Virlouvet, L., & Fromm, M. (2015). Physiological and transcriptional memory in guard cells
during repetitive dehydration stress. *New Phytologist*, *205*(2), 596–607.

Waadt, R., Seller, C. A., Hsu, P.-K., Takahashi, Y., Munemasa, S., & Schroeder, J. I. (2022).
Plant hormone regulation of abiotic stress responses. *Nature Reviews Molecular Cell
Biology*, *23*(10), 680–694. https://doi.org/10.1038/s41580-022-00479-6

Walker, T. W. N., Schrodt, F., Allard, P.-M., Defossez, E., Jassey, V. E. J., Schuman, M. C.,
Alexander, J. M., Baines, O., Baldy, V., & Bardgett, R. D. (2023). Leaf metabolic traits

reveal hidden dimensions of plant form and function. *Science Advances*, *9*(35),
eadi4029.

Wei, T., Gao, H., Yang, D., Zhu, G., Zhang, C., Wang, B., Jia, H., Zhang, C., & Guo, J.
(2024). Effects of methyl jasmonate on physiological characteristics and drought
resistance of alfalfa (Medicago sativa) and ryegrass (Lolium perenne l.). *Journal of*
Plant Growth Regulation, 1–18.

Wei, T., Gao, H., Yang, D., Zhu, G., Zhang, C., Wang, B., Jia, H., Zhang, C., & Guo, J.
(2025). Effects of Methyl Jasmonate on Physiological Characteristics and Drought
Resistance of Alfalfa (Medicago sativa) and Ryegrass (Lolium perenne L.). *Journal of*
Plant Growth Regulation, *44*(1), 335–352. https://doi.org/10.1007/s00344-024-11469-7

Wild, J., Kopecký, M., Macek, M., Šanda, M., Jankovec, J., & Haase, T. (2019). Climate at
ecologically relevant scales: A new temperature and soil moisture logger for long-term
microclimate measurement. *Agricultural and Forest Meteorology*, *268*, 40–47.
https://doi.org/https://doi.org/10.1016/j.agrformet.2018.12.018

You, J., Zhang, Y., Liu, A., Li, D., Wang, X., Dossa, K., Zhou, R., Yu, J., Zhang, Y., Wang, L.,
& Zhang, X. (2019). Transcriptomic and metabolomic profiling of drought-tolerant and
susceptible sesame genotypes in response to drought stress. *BMC Plant Biology*, *19*(1),
267. https://doi.org/10.1186/s12870-019-1880-1

Yu, X., Zhang, W., Zhang, Y., Zhang, X., Lang, D., & Zhang, X. (2018). The roles of methyl
jasmonate to stress in plants. *Functional Plant Biology*, *46*(3), 197–212.

Zhang, D., Liu, Y., Yang, Z., Song, X., Ma, Y., Zhao, J., Wang, X., Liu, H., & Fan, L. (2023).
Widely target metabolomics analysis of the differences in metabolites of licorice under
drought stress. *Industrial Crops and Products*, *202*, 117071.
https://doi.org/https://doi.org/10.1016/j.indcrop.2023.117071

Zhou, H., Zhou, G., He, Q., Zhou, L., Ji, Y., & Zhou, M. (2020). Environmental explanation
of maize specific leaf area under varying water stress regimes. *Environmental and*
Experimental Botany, *171*, 103932.

Treatment	Richness				Shannon				Uniqueness			
	Positive mode		Negative mode		Positive mode		Negative mode		Positive mode		Negative mode	
	P-value	Variation explained	P-value	Variation explained	P-value	Variation explained	P-value	Variation explained	P-value	Variation explained	P-value	Variation explained
D1	0.43	-	0.7	-	0.011	12.29	0.83	-	0.20	-	0.31	-
D2	<0.001	23.6	0.54	-	0.927	-	0.008	14.41	0.001	28.72	0.001	44.07
M1	0.146	-	0.39	-	0.009	12.82	0.24	-	0.25	-	0.094	-
D1×D2	0.038	6.59	0.001	21.34	0.343	-	0.14	-	0.023	5.44	0.014	5.88
D1×M1	0.659	-	0.85	-	0.049	6.63	0.36	-	0.094	-	0.097	-
D2×M1	0.77	-	0.7	-	0.528	-	0.9	-	0.001	11.93	0.02	4.78
D1×D2×M1	0.309	-	0.484	-	0.38	-	0.954	-	0.074	-	0.012	5.39

Table 1: Effects of D1, D2, M1, and their interactions on Richness, Shannon diversity, and Uniqueness in both positive and negative modes. The variation explained is shown for the significant predictors ($p < 0.05$).

843   **Figure 1**: Boxplots showing the effects of interactions of D2, D1, and M1 in positive mode on

844   metabolome (a) Richness, (b) Shannon diversity, and (c, d) Uniqueness in *F. rubra*. The X axis

845   shows the levels of the interacting treatment, with 0 indicating absent and 1 present treatment

846   (drought or MeJa).

Treatment	Metabolome				Physiology	
	Positive Mode		Negative Mode			
	p-value	Var. expl (%)	p-value	Var. expl (%)	p-value	Var. expl (%)
D1	0.004	3.51	0.001	4.14	0.001	18.34
D2	0.001	15.82	0.001	15.57	0.001	23.90
M1	0.001	5.16	0.001	4.92	0.001	31.12
D1×D2	0.151	-	0.098	-	0.146	-
D1×M1	0.005	3.7	0.001	3.66	0.158	-
D2×M1	0.002	3.5	0.004	3.49	0.001	9.83
D1×D2×M1	0.264		0.182		0.014	1.8

Table 2: Summary of Redundancy Analysis (RDA) results showing the effects of drought

memory (D1), current drought (D2), MeJA memory (M1), and their interactions on

metabolomic profiles (in positive and negative ionization modes) and morpho-physiological

traits. The table reports the p-values and percentage of variation explained for each significant

predictor. Only significant values ($p \leq 0.05$) are associated with variation explained (%); dashes

(–) indicate non-significance.

862 **Figure 2**: RDA biplot illustrating the relationships between treatment variables (shown as red

863 arrows) and morpho-physiological traits (shown as grey arrows) in *F. rubra*.

	Positive mode		Negative mode	
	P	Var. expl (%)	P	Var. expl (%)
SLA	0.001	12.26	0.001	11.11
LDMC	0.112	-	0.106	-
RWC	0.003	4.18	0.001	4.07
CCS2	0.015	3.32	0.016	2.92
CFR1	0.135	-	0.136	-

871   **Table 3**: Effects on plant traits on metabolic composition for positive and negative ionization

872   modes assessed using redundancy analysis (RDA).

Figure 3: Heatmap displaying the scaled intensity values of differentially accumulated

metabolites across various treatment groups in *F. rubra* (M1, D1, D2, i.e. MeJA memory,

drought memory and current drought respectively), others are the plants which got combination

M1, D2 and D2 treatment, while missing code in combination indicates that plants where not

treated by the given treatment. C indicates Control, i.e., no drought or MeJa treatment memory,

and also in well-watered conditions in both positive mode (a) and negative mode (b). The

clustering was performed using hierarchical clustering based on Euclidean distance and

complete linkage. The columns represent plants with different memory and current stress

conditions. The orange color in the plot represents a higher accumulation of a metabolite, while

the blue color represents a lower accumulation. Also, the zoomed image of marked clusters is

provided in the supplementary material with upregulated and downregulated metabolite names

in marked clusters.

Supplementary

Treatment	Richness (p-value)		Shannon (p-value)		Hill Even (p-value)		Uniqueness (p-value)		Metabolites-Treatment (p-value)		Traits-Treatment (p-value)
	P	N	P	N	P	N	P	N	P	N	
SM	0.077	0.118	0.45	0.77	0.63	0.67	0.073	0.038	**0.025**	**0.041**	0.131

Table S1: This table presents *p*-values comparing metabolome diversity metrics and metabolomic/trait-level responses for soil memory (SM). Metabolome analysis was performed in both positive (P) and negative (N) ionization modes. Significant results ($p < 0.05$) indicate treatment-induced effects and are in bold.

Text S2: **Experimental design**

The experiment represented a factorial cross of drought and MeJA treatments, each with two levels (yes/no). Each treatment had 10 replicates. The soil utilized in the experiment was sourced from the same meadow where the experimental plants were originally collected. Pots with a volume of 500 ml were filled with the soil, and one *F. rubra* ramet was planted into each pot. All the planted ramets were of similar age and size (approximately 3 weeks old and 3.5 to 4 cm tall). The plants were placed in a growth chamber with long-day conditions, where the temperature ranged from 17 to 20 °C during the day (16 hours) and 7 to 10 °C at night (8 hours). These conditions reflect conditions at the original locality during peak growing season. The plants were allowed to establish for 21 days before starting the experiment. The experiment included cycles of drought and recovery phases, each lasting 30 days (Drought stage 1 - Recovery stage 1 - Drought stage 2 - Recovery stage 2 - Drought stage 3 - Recovery stage 3).

Control watering treatment received regular watering daily, maintaining 1 cm of water above the pot base in the trays throughout the experiment. Drought-stressed plants received 50 ml of water in specific watering intervals during the drought stages: every 7th day during drought stage 1, every 10th day in drought stage 2, and every 15th day in drought stage 3. There was a progressive increase in drought stress among phases because treating plants to an excessive amount of stress initially can result in total plant loss, forcing the experiment to terminate after the first drought stage. In each phase, we attempted to expose the plants to as intense drought stress as possible but made sure they would survive successive drought stages. During the recovery stage, drought-stressed plants received the same watering as the control ones. The soil moisture has been monitored using TMS dataloggers (Wild et al., 2019).MeJA-treated *F. rubra* plants received 5 ml of 10 µM MeJA by pouring it directly onto the soil every 24 hours during the stress phases, but not during recovery. Only drought plants also got 5 ml of distilled water when 5 ml of 10 µM MeJA was given to MeJA-treated plants during drought. Previous studies

predominantly relied on foliar application of MeJA for plant treatment (e.g. Tayyab et al., 2020;

Chungloo et al., 2023)We decided to apply it to the soil as foliar application has various

disadvantages. The volatile nature of MeJA poses a risk of environmental loss when applied

through foliar application, and there is an additional concern of uneven MeJA distribution

among leaves across the plant. Therefore, soil application is a viable solution to address these

challenges effectively. Soil is an efficient absorber, thus minimizing the risk of MeJA loss to

the environment, while root uptake guarantees a more consistent distribution of MeJA

throughout the entire plant.

Figure S1: Correlation heatmap showing the relationship among chlorophyll fluorescence

parameters measured during the stress phase (CFS1, CFS2), recovery phase (CFR1, CFR2),

chlorophyll content measured during the stress phase (CCS1, CCS2), and recovery phase

(CCR1, CCR2). Correlation coefficients are color-coded from dark blue (low correlation) to

dark red (high correlation), with values displayed within each cell.

Treatment	Hill Eveness			
	Positive mode		Negative mode	
	P-value	Variation explained	P-value	Variation explained
D1	0.01	10.66	0.97	-
D2	0.357	-	0.01	14.35
M1	0.007	19.78	0.37	-
D1×D2	0.8	-	0.43	-
D1×M1	0.011	10.47	0.36	-
D2×M1	0.489	-	0.89	-
D1×D2×M1	0.167	-	0.897	-

951 **Table S2:** Table showing effects of M1, D1, D2, i.e., MeJA memory, drought memory, and

952 current drought respectively, and their interactions on Hill Eveness. The variation explained is

953 shown for the significant predictors ($p < 0.05$).

969 **Figure S2**: Boxplots showing the effects of interactions of M1, D1, D2, i.e. MeJA memory,
970 drought memory and current drought, respectively in HillEven for (a) Positive mode and (b)
971 negative mode in *F. rubra*. The X axis shows the levels of the interacting treatment, with 0
972 indicating absent and 1 present treatment (drought or MeJa).

Figure S3: Correlation heatmap showing the relationship among Richness, Shannon diversity,

and HillEvenness. Correlation coefficients are color-coded from dark blue (low correlation) to

dark red (high correlation), with values displayed within each cell.

Figure S4: Boxplots showing the effects of interactions M1, D1, D2, i.e. MeJA memory, drought memory and current drought, respectively, in positive mode for (a) Richness, (b) Shannon diversity, and (c and d) Uniqueness in *F. rubra*. The X axis shows the levels of the interacting treatment, with 0 indicating absent and 1 present treatment (drought or MeJA).

Figure S5: Boxplots show the distribution of metabolic uniqueness scores for each treatment

group under negative ionization mode, representing the triple interaction of drought memory,

current drought, and MeJA memory. In each code, a 1 in the first position indicates the presence

of drought 1 (D1), a 1 in the second position indicates the presence of drought 2 (D2), and a 1

in the third position indicates the presence of the morphological modifier (M1). For example,

1_0_0 represents D1 only, 0_1_0 represents D2 only, and 1_1_1 represents a combination of

D1, D2, and M1. The control group is 0_0_0, where no drought or modifier is applied.

1000

1001

1002

1003 **Figure S6:** The RDA plots both ionization modes (**a**)positive and (**b**) negative, providing a

1004 clear view of the variation in metabolite profiles explained by M1, D1, D2, i.e., MeJA memory,

1005 drought memory and current drought, respectively.

1006

1007

1008

Figure S7: Bar plots representing the number of significant metabolites associated with

drought memory (D1), MeJA memory (M1), current drought (D2), and their interactions

(D1×D2, D1×M1, D2×M1) under positive ionization mode (a) and negative ionization mode

(b). Each treatment or interaction is shown with two bars: shared metabolites (orange) indicate

those common with other conditions, while unique metabolites (green) represent those specific

to a given condition or interaction. The values above each bar indicate the exact number of

metabolites in each category.

1016

1017

1018

1019

Figure S8: Heatmap displaying the scaled intensity values of differentially accumulated

metabolites across various treatment groups in *F. rubra* (M1, D1, D2, i.e. MeJA memory,

drought memory and current drought respectively), others are the plants which got combination

M1, D2 and D2 treatment, while missing code in combination indicates that plants where not

treated by the given treatment. C indicates Control, i.e., no drought or MeJa treatment memory,

and in well-watered conditions. Heatmap a to d shows cluster 1 to 4, respectively for positive

mode, while heatmap e and f shows cluster 1 and 2, respectively for negative mode