Metabolic engineering of *Barbarea vulgaris* identifies β-amyrin synthase LUP5 as a

determinant of saponin-mediated insect resistance

Jincheng Shen[1], Jan Günther[1], Pablo D. Cárdenas[1*], Søren Bak[1*].

[1]Department of Plant and Environmental Sciences, Faculty of Science, University of

Copenhagen, 1871 Frederiksberg C, Denmark.

* Corresponding author: bak@plen.ku.dk

* Co-corresponding author: pdcardenas@plen.ku.dk

Word counts for the main sections: 5748

Number of figures: 5

Supporting information: 2 figures and 9 tables

Summary

• Plant–insect coevolution drives the diversification of specialized metabolites, yet the

genetic basis of ecotypic variation in defensive chemistry remains poorly understood. The

wild crucifer *Barbarea vulgaris* comprises two ecotypes—an insect-resistant G-type and

a susceptible P-type—characterized by distinct triterpenoid saponin profiles.

• To investigate the causal role of saponin composition in insect resistance, we developed

a stable transformation system for *B. vulgaris*. We expressed the G-type β-amyrin

synthase gene (*LUP5*) in both ecotypes and silenced the cytochrome P450 gene

CYP72A552 in the G-type. We combined metabolic profiling with insect feeding assays to

assess functional outcomes.

• Ectopic expression of G-type *LUP5* in the susceptible P-type conferred up to 95%

reduction in *Plutella xylostella* feeding, accompanied by increased accumulation of three

hederagenin-derived monodesmosidic saponins. In contrast, silencing of *CYP72A552*

reduced hederagenin derived saponin levels but did not affect herbivory, suggesting

threshold-dependent resistance.

• We provide direct *in planta* evidence that *LUP5* is a key determinant of ecotypic insect

resistance in *B. vulgaris*. Our findings establish a functional link between oxidosqualene

cyclase activity, saponin structural diversification, and ecological defense, highlighting the

roles of both sapogenin backbone and glycosylation pattern in saponin-mediated

herbivore deterrence.

Keywords: triterpenoid saponins, metabolic engineering, β-amyrin synthase, insect

resistance, *Barbarea vulgaris,* chemical ecology

Introduction

Plants have evolved a wide array of specialized metabolites to deter insect herbivores (Fürstenberg-Hägg, Zagrobelny and Bak, 2013; Yactayo-Chang et al., 2020; Beran and Petschenka, 2022). Among these, triterpenoid saponins, which consist of a hydrophobic sapogenin backbone and one or more hydrophilic sugar moieties, represent a structurally diverse and ecologically important class of defensive compounds. Due to their structural similarity to sterols and steroid hormones, triterpenoid saponins have been hypothesized to interact with eukaryotic cell membranes. In addition, their inherent bitterness may deter herbivores, contributing to their role as effective defense agents (Augustin et al., 2011). Thousands of triterpenoid saponins have been reported to date (Xu, Fazio and Matsuda, 2004; Augustin et al., 2011; Moses, Papadopoulou and Osbourn, 2014; Huang et al., 2020). Importantly, specific saponin structures have been implicated in insect resistance, highlighting the functional significance of their chemical diversity (Stevenson, Nyirenda and Veitch, 2010; Cui et al., 2019; Hussain et al., 2019; Liu et al., 2019; Dolma et al., 2021).

The structure–activity relationships underlying saponin bioactivity are critical to their defensive function. For example, hederagenin 3-*O*-monoglucoside has higher toxicity and antifeedant activity than oleanolic acid 3-*O*-monoglucoside (which lacks C-23 hydroxylation), and gypsogenic acid 3-*O*-monoglucoside (which is further oxidized at C-23 to the ketone) (Liu et al., 2019). Additionally, monodesmosidic hederagenin and oleanolic acid exhibited higher toxicity and antifeedant activity than their bidesmosidic counterparts (Tian et al., 2021). These observations suggest that even subtle modifications to the sapogenin backbone, glycosylation pattern, or sugar linkage can significantly alter saponin bioactivity. This structural diversity makes saponins a valuable system for studying molecular evolutionary mechanisms of specialized metabolism, particularly the structure–activity relationships underlying their ecological functions in insect defense. Despite growing knowledge of these structure–activity relationships, the genetic and biochemical mechanisms underlying saponin diversification remain poorly

understood in most species. One particularly promising ecological model to explore this

question is the wild crucifer *Barbarea vulgaris*, which naturally segregates in two ecotypes

or chemotypes that alter in both saponin profiles and insect resistance.

B. vulgaris has emerged as a powerful ecological model for investigating how

specialized metabolites contribute to herbivore resistance (Kuzina et al., 2011; Hauser,

Toneatto and Nielsen, 2012; Byrne et al., 2017). The species comprises two ecotypes: the

glabrous insect-resistant G-type, and the pubescent insect-susceptible P-type (Agerbirk,

Olsen and Nielsen, 2001). This intraspecific divergence likely arose from isolation in

different Ice Age refugia (Hauser, Toneatto and Nielsen, 2012) and is associated with

marked differences in saponin composition (Kuzina et al., 2009; Khakimov et al., 2012;

Khakimov et al., 2016). Gene duplications events have further contributed to the chemical

divergence and insect resistance (Khakimov et al., 2015; Liu et al., 2019). Insect resistance

in G-type *B. vulgaris* correlates with four oleanane-type defensive saponins, including

oleanolic acid cellobioside, hederagenin cellobioside, gypsogenin cellobioside, and 4-

epihederagenin cellobioside (Agerbirk et al., 2003; Kuzina et al., 2011). Among these,

hederagenin cellobioside is particularly toxic and confers G-type insect resistance

(Agerbirk et al., 2003; Nielsen et al., 2010; Kuzina et al., 2011).

The biosynthetic pathway leading to hederagenin cellobioside in G-type *B. vulgaris*

has been partially elucidated. The oxidosqualene cyclase *LUP5* is highly expressed in the

G-type and primarily produces β-amyrin (Khakimov et al., 2015). Subsequent enzymatic

steps involve CYP716A80, CYP72A552, and UGT73C11, which sequentially convert β-

amyrin into oleanolic acid, hederagenin, and hederagenin 3-*O*-glucoside (Augustin et al.,

2012; Khakimov et al., 2015; Liu et al., 2019). The enzyme responsible for the final step in

forming hederagenin cellobioside remains unidentified (Erthmann, Agerbirk and Bak,

2018). In contrast, *LUP5* is expressed at very low levels in *B. vulgaris* P-type and primarily

produces α-amyrin due to two amino acid substitutions as compared to the *LUP5* in the

G-type (Günther et al., 2021). Despite 98% sequence identity at the amino acid level, the

G- and P-type LUP5 enzymes differ markedly in both their gene expression and product

profiles. The P-type plants preferentially express *LUP2,* an oxidosqualene cyclase over

80% identical at the amino acid level to *LUP5*, but it produces predominantly lupeol (98%)

and only 2% β-amyrin-derived saponins (Khakimov et al., 2015). Notably, the genes

encoding for the enzymes responsible for the biosynthesis of hederagenin 3-*O*-glucoside

(CYP716A80, CYP72A552, and UGT73C11) are present in both P-type and G-type and

exhibit similar substrate and product specificity (Augustin et al., 2012; Khakimov et al.,

2015; Liu et al., 2019). However, the cyclases are differentially regulated suggesting that

upstream regulation and oxidosqualene cyclase product specificity identity may be key

drivers of saponin structural and insect resistance difference. Additionally, CYP72A552,

which catalyzes the C23 hydroxylation of oleanolic acid to hederagenin, the precursor of

the more toxic hederagenin-type saponins, may also be pivotal to defense against insects,

but its precise role in *B. vulgaris* defense remains to be experimentally validated.

Despite these prior insights, a direct causal link between saponin composition and

insect resistance has not yet been established *in planta*. Here, we leverage the natural

variation in *B. vulgaris* to develop a model system that connects specialized metabolism

with ecological function. We provide functional evidence that engineering saponin

biosynthesis through *LUP5* manipulation can substantially enhance insect resistance in

this wild crucifer and explains the molecular basis of the difference in insect resistance

between the G- and P-type *B. vulgaris.*

Materials and methods

Plant and insect materials

Seeds of *Barbarea vulgaris* G-type (glabrous, accession B44) and P-type (pubescent,

accession B4) were provided by Associate professor Niels Agerbirk *(Agerbirk et al., 2021).*

Brassica napus plants were grown in a greenhouse at 19°C under a 16:8 h light-dark cycle.

Plutella xylostella (diamondback moth) was reared on *B. napus* at 20°C under the same

light conditions. Third instar *P. xylostella* larvae were used for the insect feeding assay.

Plasmids construction

For stable transformation, the pJCV51 vector was employed to introduce the G-type

LUP5 gene, while pK7FWG2 was utilized for *CYP72A552* silencing. The *LUP5* gene, its

native promoter (1988 bp), and the *CYP72A552* gene were PCR-amplified from *B. vulgaris*

G-type cDNA and genomic DNA (Table S1, Table S2). G-type *LUP5* expression was driven

either by the 35S promoter or its native promoter, while *CYP72A552* was driven by the

35S promoter.

The pJCV51-p35S::LUP5 and pK7FWG2-CYP72A552 RNAi plasmids were assembled

using Gateway technology (Gateway™ BP Clonase™ II Enzyme mix, Thermo Fisher

Scientific, 11789020; Gateway™ LR Clonase™ II Enzyme mix, Thermo Fisher Scientific,

11791020) (Table S1). pDONR207 was used as the entry clone vector for both constructs

(Thermo Fisher Scientific, 117207-021). Subsequently, the pJCV51-pLUP5::LUP5 plasmid

was generated by replacing the 35S promoter in pJCV51-p35S::LUP5 with the G-type *LUP5*

promoter using the SalI sites. Finally, the plasmids were introduced into *Agrobacterium*

tumefaciens strain AGL1 via electroporation (Hayta et al., 2021).

Stable transformation in wild crucifer *B. vulgaris*

Seed sterilization was achieved by immersing seeds in 70% ethanol for 30 s, followed

by a 12.5 min treatment in 3% sodium hypochlorite (Thermo Fisher Scientific) with 0.1%

Tween20 (Bio-RAD, 1706531) and thorough rinsing with sterile water. Seeds were sown

on germination medium in Magenta boxes and incubated in darkness for 6–8 days (Table

S3) .

Hypocotyl explants (~3 mm) from 6–10-day-old seedlings were incubated for 15 min

with *A. tumefaciens* strains (OD_{600} = 0.5 in 10 mM $MgCl_2$, 100 µM acetosyringone) with

gentle agitation, and then co-cultivated in darkness for three days (Table S3). The pJCV51-

pLUP5::LUP5 and pJCV51-35S::LUP5 constructs were transformed into both *B. vulgaris*

ecotypes, while pK7FWG2-CYP72A552 RNAi was only introduced into the G-type.

After rinsing in Milli-Q water containing 1 µL/mL timentin, the explants were

transferred sequentially to callus induction (one week), shoot induction (two rounds, two

weeks each), and root induction media (two rounds, two weeks each) at 25°C under a

16:8 h light–dark cycle. Throughout these stages, the explants were maintained in a

climate chamber at 25°C under a 16:8 h light-dark cycle. Kanamycin (50 ng/mL) was used

for selecting transformed explants, whereas wild-type explants were cultured without

kanamycin supplementation in the medium. Regenerated plants were confirmed by

DsRed fluorescence (for *LUP5* expression constructs) or NPTII (neomycin

phosphotransferase II) ELISA Kit (Agdia, PSP 73000/0288) for CYP72A552 RNAi.

Subsequently, the confirmed plants were transferred to soil and grown in a greenhouse

at 19°C under a 16:8 h light–dark cycle.

Choice insect feeding assay

Insect resistance was evaluated using at least three individual plants (biological

replicates) per transgenic line. For each transgenic plant, five transgenic and five wild-

type leaf discs (1.57 cm² each) were alternately arranged in a PETRI DISH (9.4 x 1.6 cm)

(Greiner BIO-ONE, 633180) with filter paper moistened with 2 mL of water. After a 3–5

hour starvation period, ten third-instar *P. xylostella* larvae were placed at the center of

the plate, following a modified protocol from Liu et al (Liu et al., 2019). Leaf consumption

was quantified when half of the wildtype leaves were consumed using ImageJ (approx 7.5

h).

Preparation of plant extracts and metabolite analysis

The extraction of saponins and sapogenins was adapted from Khakimov *et al*. (2016).

B. vulgaris leaf samples were ground with liquid nitrogen using a mortar and pestle. For

saponin detection, 100 mg of leaf powder was weighed into 1.5 mL Eppendorf tubes and

extracted with 300 µL of 85% methanol (v/v) via sonication for 30 minutes, followed by

centrifugation at 16,000 g for 10 minutes. Supernatants (200 µL) were filtered through a

0.22-µm filter before LC-qTOF-ESI-MS/MS analysis.

To detect total and free sapogenins, 200 mg of leaf powder was weighed into 2 mL

Eppendorf tubes and extracted with 600 µL of 85% methanol (v/v) at 100°C while mixing

at 1,400 rpm in a thermomixer (Eppendorf, Denmark). The samples were cooled on ice

and centrifuged at 16,000 × g for 3 min. After filtration through a 0.22-μm filter, 100 μL

of supernatant was collected for free sapogenin detection via LC-qTOF-APCI-MS/MS. An

additional 300 μL of supernatant was collected and dried under nitrogen flow. To cleave

off sugar residues, samples were treated with 500 μL of 2 M hydrochloric acid at 100°C

while mixing at 1400 rpm for 1.5 hours in a thermomixer (Eppendorf, Denmark). After

cooling on ice, a double volume of ethyl acetate was added to the acid-water mixture,

followed by vortexing and centrifugation at 3,500 × g for 5 minutes. The samples were

extracted three times with ethyl acetate to collect sapogenins. To remove residual acid,

an equal volume of Milli-Q water was added, the samples was subsequently vortexed and

centrifugated at 3,500 × g for 5 minutes. This washing step was repeated three times. The

final extracts were dried under nitrogen flow, re-solubilized in 300 μL of 100% methanol,

and filtered through a 0.22-μm filter before total sapogenin detection via LC-qTOF-APCI-

MS/MS.

LC-qTOF-ESI-MS/MS was performed according to Liu *et al*. (2019). LC-qTOF-APCI-

MS/MS was performed as described by Patel *et al*. (2020). Untargeted metabolite data

were analyzed using XCMS (Tautenhahn et al., 2012), and targeted saponin and sapogenin

analysis were performed according to Khakimov *et al*. (2016). Hederagenin cellobioside

and oleanolic acid cellobioside standards were used for saponin identification. Lupeol, β-

amyrin, oleanolic acid, and hederagenin standards were employed for sapogenin

identification.

Gene expression analysis

Total RNA was extracted using the Spectrum™ Plant Total RNA Kit (Sigma-Aldrich,

STRN50-1KT) and then synthesized to cDNA by iScript™ cDNA Synthesis Kit (Bio-Rad,

1708891). Relative expression of *LUP5* was quantified via qPCR using kapa SYBR® fast kit

(Merck-Sigma, KK4607) (Khakimov et al., 2015). *Tubulin* (GenBank accession no.

EU555191) was used as reference gene, and *LUP5* expression levels were normalized to

the wild-type P-type control. Primer sequences are provided in Table S1 (Khakimov et al.,
2015).

Results

Establishment of a transformation and regeneration system in wild crucifer *B. vulgaris*

To enable functional studies of insect resistance in *B. vulgaris* as an ecological model
system, we developed a stable transformation and regeneration method for both the G-
type and P-type ecotypes (Fig.1a, Fig. S1, Table S3). To metabolically engineer the saponin
compositions, we introduced the G-type *LUP5* gene into both ecotypes and attempted to
silence *CYP72A552* in the G-type. *LUP5* expression was driven either by the constitutive
35S promoter or the native *LUP5* promoter. Transformation efficiency, calculated as the
percentage of positive transgenic plants to all obtained plants from tissue culture,
reached up to 94% across experiments (Fig. 1b). The native *LUP5* promoter exhibited
approximately two-fold higher transformation efficiency than the 35S promoter in both
the G- and P-type, indicating a selection pressure against overexpression *LUP5* using the
35S-promoter (Fig. 1b).

b Construct for transformation	Ecotype	Total numbers of plants	Positive/total
pJCV51-pLUP5::LUP5	G-Type	31	93.54%
	P-Type	35	60.00%
pJCV51-p35S::LUP5	G-Type	52	40.4%
	P-Type	66	25.8%
pK7FWG2-CYP72A552 RNAi	G-Type	46	28.26%

Fig. 1 Triterpenoid saponin biosynthesis pathways and stable transformation efficiency in *B. vulgaris*. (a) Biosynthetic pathways for lupeol and β–amyrin derived triterpenoid saponins in *B. vulgaris*. Monodesmosidic saponin, a single sugar chain attached at either C-3 (R1) or C-28 (R2); bidesmosidic saponin, sugar chains attached at both C-3 (R1) and C-28 (R2). (b) Transformation efficiency in *B. vulgaris*.

Stably expression of G-type *LUP5* in the P-type confers insect resistance

To evaluate the impact of G-type *LUP5* expression on insect resistance, a choice insect feeding assay was conducted by presenting alternating wildtype and transgenic *B. vulgaris* leaf discs to diamondback moth larvae. Leaf area consumed was measured when approximately half of the total leaf material of the wild type had been eaten (approx. 7.5 hr). *LUP5*-transformed P-type plants exhibited a significant reduction in larval consumption, ranging from 76- 95% (Fig. 2a,b). QPCR on the best performing 35S and LUP5 promoter lines, confirmed that the G-type *LUP5* gene was transformed and expressed in the P-type both under the 35S promoter or the native G-type *LUP5* promoter (Fig. 2c). Interestingly, the 35S promoter expressed the *LUP5* gene at approximately

sixteen-fold higher levels than that in the best pLUP5::LUP5 transformed P-type and the

wildtype P- and G-type (Fig. 2c).

Fig. 2 *B. vulgaris* P-type plants expressing the G-type *LUP5* gene leads to reduced feeding

by diamond back moth larvae. (a) Consumption area (%) after 7.5 h (choice test). (b)

Example of the insect choice feeding assay, and the resulting leaf condition after 7.5 hr

insect exposure. (c) Relative *LUP5* gene expression levels. PWT, wildtype P-type *B. vulgaris*;

GWT, wildtype G-type *B. vulgaris*; p35S::LUP5, p35S::LUP5 transformed P-type *B. vulgaris*;

pLUP5::LUP5, pLUP5::LUP5 transformed P-type *B. vulgaris*, LUP5, P-type *B. vulgaris*

expressing the G-type *LUP5* gene. Statistical significance is indicated by ** ($p < 0.05$) and

*** ($p < 0.005$).

Expression of G-type *LUP5* gene induced β-amyrin-derived saponins in P-type *B.*

***vulgaris*.**

Based on the highest reduction of leaf area consumed, three transgenic lines were

selected for each promoter type (35S and native *LUP5*) and subjected to untargeted and

targeted LCMS analyses to assess metabolite changes. The untargeted metabolite

analysis revealed approximately 9,500 distinct m/z features—each defined as a unique

m/z and retention time (RT) pair. The top 100 significantly changed features (P<0.5) were

identified by comparing *LUP5*-expressing P-type with wild-type. However, extraction of

the m/z features using DataAnalysis software revealed that the majority lacked clear MS²

fragmentation patterns, thus structural identification could not be deduced. This

outcome suggests that many of the significantly altered features may represent in-source

fragments, background ions, or uncharacterized metabolites for which fragmentation

data were inconclusive.

Consequently, a targeted saponin analysis was conducted using DataAnalysis. *B.*

vulgaris saponins were detected by extracted ion chromatograms of our previously

tentatively identified *B. vulgaris* aglycones and their structures were inferred based on

fragmentation patterns. Based on 455 m/z, 469 m/z and 471 m/z, the characteristic

fragmentation patterns of oleanolic acid, gypsogenin and hederagenin were tentatively

identified, respectively. This targeted approach identified ten distinct saponins in P-type

lines and thirty in G-type lines, which all included in the untargeted analysis data (Table

S4, Table S5). All detected saponins were classified as monodesmosidic, based on their

fragmentation patterns. Saponin abundance was subsequently quantified by calculating

peak areas from extracted ion chromatograms (Table S4, Table S5).

The LCMS analysis showed that three hederagenin-derived monodesmosidic

saponins (each with two hexoses) at retention times 10.7, 11.5, 12.3, and one oleanolic

acid monoglucoside (with one hexose) at retention time 9.5 min, accumulated in

significantly higher amount in *LUP5*-transformed P-type as compared to wildtype P-type

(Fig. 3a,b). The most abundant of the three hederagenin-derived saponin eluted at 11.5

min and was increased ~four-fold in the P-type expressing *LUP5* under the native *LUP5*

promoter compared to the wildtype and was ~two-fold more abundant than in P-type

plants transformed with *LUP5* under the 35S promoter (Fig. 3a). Hederagenin cellobioside

(eluted at 12.3 min) was also increased ~four-fold but remained at lower levels in

pLUP5::LUP5 transformed P-type (Fig. 3a). These three hederagenin-derived saponins

were also present in wild G-type plants, with hederagenin cellobioside (eluted at 12.3 min)

being the most abundant hederagenin-derived saponin (Table S5, Table S8). The

increased oleanolic acid monoglucoside (eluted at 9.5 min) was not detected in G-type

(Table S5). The other detected saponins remained unchanged (Table S6).

Fig. 3 Expression of the G-type *LUP5* gene increased glycosylated oleanolic acid and

hederagenin levels in the transgenic P-type *B. vulgaris*. (a) Monodesmosidic hederagenin

(with 2 hexoses) peak area (EIC: m/z = 841.4591 ± 0.2). (b) Oleanolic acid monoglucoside

peak area (EIC: m/z = 617.4059 ± 0.2). (c) Peak area of sapogenins, including total

hederagenin (EIC: m/z = 437.3425 ± 0.2) and total oleanolic acid (EIC: m/z = 439.3593±

0.2). PWT, wildtype P-type *B. vulgaris*; p35S::LUP5, p35S::LUP5 transformed P-type *B.*

vulgaris; pLUP5::LUP5, pLUP5::LUP5 transformed P-type *B. vulgaris*. Statistical

significance is indicated by *(p<0.5), ** (p < 0.05) and *** (p < 0.005).

To evaluate the effect of G-type *LUP5* expression on P-type *B. vulgaris* saponin

aglycone biosynthesis and accumulation, levels of free and total sapogenins were

analyzed. Free sapogenins were measured from untreated samples, representing

aglycones that occur naturally in a non-glycosylated form. Total sapogenins were defined

as the sum of free and sugar-bound sapogenins, the latter released through hydrolysis of

glycosylated saponins. A transgenic line (with three vegetative clones) from both

p35S::*LUP5* and pLUP5::*LUP5*-transformed P-type *B. vulgaris* was selected based on

reduced leaf area consumption and increased accumulation of four saponins (Fig. 2a, Fig.

3a,b). This analysis revealed a significant increase in total hederagenin and oleanolic acid

levels in *LUP5*-transformed P-type plants (Fig. 3c), while total lupeol, β-amyrin, and free

sapogenins remained undetectable (Table S7). Higher accumulation of total hederagenin

was observed in plants expressing *LUP5* under the native promoter compared to those

under the 35S promoter (Fig. 3c).

Introduction of G-type *LUP5* in G-type increased a monodesmosidic hederagenin

(with three hexoses) and oleanolic acid monoglucoside (with two hexoses) in both

p35S::LUP5 and pLUP5::LUP5 transformed G-type plants, retention times 8.5, and 11.0

min, respectively (Fig. 4a-c). Monodesmosidic oleanolic acid (with 3 hexose, retention

time at 10.7 min) increased in all pLUP5::LUP5 transformed G-type lines (Fig. 4c). Twenty-

six other saponins showed no changes (Table S8). The three slightly increased saponins

remained at a low concentration compared to other saponins (Fig. 4 a-c, Table S8).

Fig. 4 Expression of G-type *LUP5* gene increased glycosylated oleanolic acid and

hederagenin levels in G-type in G-type *B. vulgaris*. (a) Monodesmosidic hederagenin (with

3 hexose) peak area (EIC: $m/z = 1003.5115 \pm 0.2$). (b) Monodesmosidic oleanolic acid (with

two hexoses) peak area (EIC: $m/z = 617.4059 \pm 0.2$). (c) Monodesmosidic oleanolic acid

(with 3 hexose) peak area (EIC: $m/z = 987.517 \pm 0.2$). GWT, wildtype G-type *B. vulgaris*;

p35S::LUP5, p35S::LUP5 transformed G-type *B. vulgaris*; pLUP5::LUP5, pLUP5::LUP5

transformed G-type *B. vulgaris*. Statistical significance is indicated by *($p<0.5$), ** ($p < 0.05$)

and *** ($p < 0.005$).

In conclusion, stable transformation of G-type *LUP5* in P-type significantly increased

one oleanolic acid monoglucoside (with one hexose) and three hederagenin-derived

monodesmosidic saponins (each with two hexoses) (Fig. 3a,b). Notably, the oleanolic acid

monoglucoside—eluting at 9.5 minutes—was detected exclusively in the transformed P-

type (Fig. 3b, Table S5). The most abundant hederagenin-derived saponin in P-type eluted

at 11.5 min, whereas hederagenin cellobioside (eluted at 12.3 min) remained the

dominant saponin in G-type (Fig. 3a, Table S6, Table S8). Introduction of G-type *LUP5* in G-

type slightly increased saponins, but their concentrations remained low compared to

other oleanolic acid- and hederagenin-derived compounds (Fig. 4a-c, Table S8).

Expresseion of G-type *LUP5* by its native promoter resulted in higher transformation

efficiency and induced greater accumulation of β-amyrin-derived sapogenins and their

glycosylated saponins compared to the 35S promoter (Fig. 1b, Fig. 3a-c, Fig. 4a-c, Table

S7).

CYP72A552 silencing decreased hederagenin derived saponin accumulation in G-type

B. vulgaris

To evaluate the impact of G-type *CYP72A552* silencing on insect resistance, a choice

insect feeding assay was conducted as described above. However, neither the transgenic

G-type nor the wild-type G-type plants were consumed by diamondback moth larvae (Fig.

5g). To assess the impact of *CYP72A552* downregulation on saponin and sapogenin levels

in G-type *B. vulgaris*, a metabolomic analysis was performed as described above. Silencing

of *CYP72A552* reduced the accumulation of six saponins in the G-type, including

hederagenin monoglucoside and three other hederagenin-derived monodesmosidic

saponins (each with two hexoses), monodesmosidic gypsogenin (with two hexoses), and

one unknown saponin (Fig. 5a-e). Based on MS/MS analysis, the saponin has a sapogenin

with a molecular weight of 457 m/z, along with one hexose and two methylpentoses.

Twenty-three other saponins, including seven glycosylated oleanolic acids, remained

unchanged (Table S9). The sapogenin levels from *CYP72A552* silenced line #13 and

wildtype G-type *B. vulgarism were* compared. Total hederagenin levels decreased by 40%,

while total and free oleanolic acid levels remained unchanged, in agreement with the

saponin data (Fig. 5a-f). Lupeol and β-amyrin remained undetectable in G-type *B. vulgaris*

(Table S7). In conclusion, silencing *CYP72A552* significantly reduced the biosynthesis of

hederagenin and its glycosylated derivatives in G-type *B. vulgaris*.

Fig. 5 Silencing *CYP72A552* decreased hederagenin and its derived saponins in G-type *B. vulgaris*. (a) Hederagenin monoglucoside peak area (EIC: m/z = 679.4063 ± 0.2). (b) Hederagenin cellobioside peak area (EIC: m/z = 841.4591 ± 0.2). (c) Monodesmosidic hederagenin (with two hexoses) peak area (EIC: m/z = 841.4591 ± 0.2). (d) Monodesmosidic gypsogenin (with two hexoses) peak area (EIC: m/z = 839.4409 ± 0.2).

(e) unknown compound-1 peak area (EIC: m/z = 901.2429 ± 0.2). (f) Peak area of

sapogenins, including total hederagenin (EIC: m/z = 437.3425 ± 0.2), free and oleanolic

acid (EIC: m/z = 439.3593± 0.2). GWT, wildtype P-type *B. vulgaris*; CYP72A552::RNAi,

CYP72A552 silenced G-type *B. vulgaris*. Statistical significance is indicated by *($p<0.5$), **

($p < 0.05$) and *** ($p < 0.005$).

Discussion

Triterpenoid saponins are key mediators of plant–insect interactions and have

attracted attention due to their structure-dependent bioactivity and unique modes of

action. Key factors influencing insect resistance include saponin structure, composition of

the sapogenin, the types of sugar moieties, the length and number of glycoside chains,

and the glycosylation sites (Gao et al., 2011; Liu et al., 2019; Tian et al., 2021). Prior

metabolic engineering efforts have focused on modifying saponin composition by

disrupting biosynthetic genes in legumes (Confalonieri et al., 2021; Hodgins et al., 2024).

Confalonieri *et al*. (2021) knocked out *CYP93E2*, which hydroxylates β-amyrin at the C-24

position, in *Medicago truncatula*, resulting in transgenic lines that no longer produced

non-hemolytic soyasapogenols saponins but instead redirecting metabolic flux toward

the synthesis of hemolytic saponins. Hodgins *et al*. (2024) knocked out β-amyrin synthase

BAS1 in pea (*Pisum sativum*), leading to a 99.8% reduction in saponin content in the seeds.

However, these approaches have not addressed how targeted saponin engineering

influences ecological traits such as insect resistance or consumption.

Here, we metabolically engineered the saponin biosynthetic pathway in the wild

crucifer *Barbarea vulgaris* to test whether specific triterpenoid profiles confer resistance

to insect herbivory. We have previously hypothesized—based on quantitative trait loci

analysis and insect assays with saponin-treated leaf disks—that differences in *LUP5*

expression between G- and P-type *B. vulgaris* underlie their distinct saponin profiles and

insect resistance (Augustin et al., 2012; Khakimov et al., 2015; Nielsen et al., 2010; Kuzina

et al., 2011). Our engineering strategy confirms that the susceptibility of the P-type to

diamondback moth feeding is primarily due to its lack of *LUP5* expression. LUP5 encodes

an oxidosqualene cyclase that produces β-amyrin—the precursor of oleanolic acid and
hederagenin-derived saponins—both implicated in insect deterrence. Interestingly, our
data suggest that the glycoside linkage has less impact on feeding behavior than the
triterpenoid backbone itself.

These findings confirm our earlier work showing that saponin toxicity is influenced
by both backbone structure and glycosylation pattern, particularly when hydroxylation
occurs at C23 and glycosylation at C3 (Augustin et al., 2012; Liu et al., 2019). Insect-
susceptible P-type plants predominantly produce lupeol-derived saponins, reflecting their
higher expression of LUP2, while G-type plants express *LUP5* and accumulate β-amyrin-
derived saponins, including oleanolic acid and hederagenin cellobioside—both
contributing to insect resistance (Augustin et al., 2012; Khakimov et al., 2015; Liu et al.,
2019). By combining gene transfer with ecological assays, our study provides a framework
for linking specialized metabolism with ecological function in a non-model plant system.

To test this hypothesis in an *in planta* system, the β-amyrin synthase *LUP5* gene was
expressed in both the insect resistant G-type and insect susceptible P-type *B. vulgaris*. In
the otherwise insect susceptible P-type, this resulted in up to a 95% reduction in leaf
consumption and increased levels of three hederagenin-derived monodesmosidic
saponins (each with two hexoses) and one oleanolic acid monoglucoside (Fig. 2a, Fig.
3a,b). In wildtype *B. vulgaris* G-type, hederagenin cellobioside is the predominant
saponin, while only small amounts of the other two hederagenin-derived saponins were
detected at retention times of 10.7 and 11.5 min (Fig. 4a,b, Table S8). In contrast, in the
LUP5-transformed P-type, the monodesmosidic hederagenin eluting at 11.5 min is more
abundant than the other two (Fig. 3a). This variation highlights differences in saponin
glycosylation patterns between P- and G-types, suggesting that these three hederagenin-
derived monodesmosidic saponins, particularly the one detected at 11.5 min, may
contribute to insect resistance in the transformed P-type plants. Hederagenin 3-*O*-
glucoside exhibits seven-fold stronger feeding reduction to diamondback moth and
tobacco hornworm compared to oleanolic acid 3-*O*-glucoside (Liu et al., 2019). This

enhanced efficacy has been linked to C-23 hydroxylation in hederagenin, which causes
the C-3 glucose moiety to adopt a different orientation relative to the sapogenin
backbone compared to oleanolic acid (Cárdenas, Almeida and Bak, 2019; Liu et al., 2019).
Considering the bioactivity and the relatively low increase of oleanolic acid
monoglucoside compared to hederagenin-derived monodesmosidic saponins in *LUP5*-
transformed P-type suggest that there is a CYP72A552 ortholog in the P-type that
effectively transforms the oleanolic acid to hederagenin, which then is glycosylated by
different UGTs than in the G-type *B. vulgaris* to other linkage types than the 1→4 linkage
cellobiosides that predominantly accumulate in the G-type.

Based on our findings, LUP5 is confirmed as the key determinant for insect resistance
differences between G- and P-type *B. vulgaris*. Furthermore, three hederagenin-derived
monodesmosidic saponins (each with two hexoses) may play a role in insect resistance in
LUP5-transformed P-type, particularly the most abundant saponin at retention time 11.5
min. Future work comparing the insect resistance differences between the elucidated
structures of these three hederagenin-derived monodesmosidic saponins could provide
further insights into how glycosylation patterns affect saponin bioactivity. Glycoside
linkage impacts cytotoxicity of saponins in *in vivo* assays (Chwalek et al., 2006). However,
how glycoside linkage affects insect resistance remains unknown. Unfortunately, due to
their low concentration, isolating and purifying these compounds for bioactivity testing
remains challenging. Future research should focus on identifying candidate
glycosyltransferases that produces a variety of glycoside linkage with the aim to produce
defense-related hederagenin-derived monodesmosidic saponins in planta.

Interestingly, when *LUP5* expression was driven by the constitutive 35S promoter,
LUP5 was expressed at approximately sixteen-fold higher levels than in the best
pLUP5::LUP5 transgenic line and also exceeded expression levels in both wild-type P- and
G-type plants (Fig. 2c). However, this high expression did not translate into significant
increased saponin production (Fig. 2a,b, Fig. 4a-c). In contrast, using the native *LUP5*
promoter not only resulted in about two-fold higher transformation efficiency (Fig. 1b)

but also led to greater accumulation of hederagenin and its glycosides in the P-type background despite the overall much lower expression profile (Fig. 3a,c). These results underscore that promoter choice critically influences metabolic output—not simply through expression level, but likely through coordinated regulation aligned with downstream biosynthetic processes.

While constitutive promoters like 35S often used to drive high transgene expression, they can also impose metabolic burdens and reduce transformation efficiency due to unintended gene expression as shown above. Studies show that although 35S enhances transcript levels higher than native promoters, this does not always result in higher protein accumulation or improved physiological performance than native promoter (Seger et al., 2009; Beringer et al., 2017; Waller et al., 2023). Instead, its broad tissue activity can lead to premature senescence, metabolic depletion, and reduced plant productivity (Seger et al., 2009). Comparative analyses further reveal that native promoters often outperform constitutive promoters in overall plant viability. For example, in maize (*Zea mays*), the *Zm*Ubi1 promoter yielded higher transformation efficiency and better tissue culture survival than the 35S promoter, despite lower expression levels (Beringer et al., 2017). Similarly, *Anthoceros agrestis* expressing *hygromycin resistance* gene under 35S showed growth retardation and smaller leaves, whereas the *Aa*Ef1a promoter avoided these issues (Waller et al., 2023). These examples highlight the value of tissue-specific expression, which prevents unnecessary metabolic drain from non-target tissues and supports healthier development.

In conclusion, although the 35S promoter often drives high levels of transgene expression, its constitutive activity across most tissues can lead to unintended metabolic imbalances, reduced transformation efficiency, and pleiotropic effects on plant development. Furthermore, expression in tissues where the gene is not biologically relevant may not contribute to the desired function. In contrast, native promoters limit gene expression to appropriate tissues and developmental stages, enabling more precise

metabolic control and optimized biosynthetic pathway regulation—factors that better
support the intended biological outcome.

The tissue-specific expression of *LUP5* appears to play a significant role in regulating
the accumulation of hederagenin and its glycosylated saponins in *B. vulgaris*. We
hypothesize that under the 35S promoter, *LUP5* is expressed throughout all leaf tissues,
potentially causing excessive β-amyrin accumulation and contributing to the observed
reduction in transformation efficiency. However, β-amyrin was undetectable in both wild-
type and transgenic *B. vulgaris*, possibly due to rapid metabolic conversion. To further
advance understanding of saponin biosynthesis in *B. vulgaris*, future studies should aim
to identify *LUP5* promoter-driven expression patterns at the tissue level, providing deeper
insight into the spatial regulation of saponin production.

We had anticipated that silencing of *CYP72A552* in G-type would reduce insect
consumption as hederagenin derived saponins are more bioactive than those derived
from oleanolic acid. However, our results show that while silencing of *CYP72A552*
decreased hederagenin biosynthesis by up to 40%, leading to a significant reduction in
hederagenin- and gypsogenin (which is further oxidized at C-23 to the ketone)- derived
monodesmosidic saponins. Specifically, the bioactive hederagenin monoglucoside,
hederagenin cellobioside and monodesmosidic gypsogenin (with two hexoses) were
decreased (Fig. 5a-d). The decrease in monodesmosidic gypsogenin may have been
caused by the reduced hederagenin biosynthesis or off-target effects (Fig. S2), possibly
resulting from the high nucleotide sequence identity of eight tandemly repeated *CYP72A*
homologs (Liu et al., 2019). Previous studies indicate that cellobiosides of hederagenin
and gypsogenin correlate with insect resistance in *B. vulgaris* in a concentration-
dependent manner (Shinoda et al., 2002; Agerbirk et al., 2003; Kuzina et al., 2009; Liu et
al., 2019). The hederagenin cellobioside concentration in G-type *B. vulgaris* leaves is
~eleven-fold higher than the ED50 (50% effective dose) for *P. xylostella* (Shinoda et al.,
2002; Liu et al., 2019). Thus, even with a 40% reduction, the remaining saponin levels still
exceed the toxicity threshold required to deter feeding. Likewise, *LUP5*-transformed P-

type plants did not reach the full resistance potential of G-type plants, likely due to

incomplete accumulation of the most potent hederagenin saponins (Fig. 2a,b, Fig. 3a-c,

Fig. 5a-g). Overexpression of *LUP5* in G-type plants produced only marginal increases in

minor oleanolic acid and hederagenin saponins, suggesting that G-type plants may

already accumulate saponin at or close the acceptable threshold for accumulation (Fig.

5a-c, Table S8).

Conclusion

In summary, this study demonstrates that targeted metabolic engineering of

triterpenoid saponins in the wild crucifer *Barbarea vulgaris* is a robust approach to dissect

the ecological roles of structurally specific saponins and enhance insect resistance.

Expression of the β-amyrin synthase *LUP5* under its native promoter was more effective

than constitutive expression, leading to higher transformation efficiency and increased

accumulation of insect-deterring hederagenin-derived monodesmosidic saponins. This

underscores the importance of promoter choice and precise transcriptional regulation in

optimizing specialized metabolite production.

Our findings establish metabolic engineering as not only a tool for enhancing

resistance traits but also as a powerful approach for functional genomics. We identified

LUP5 as a crucial determinant of insect resistance in *B. vulgaris*, and showed that three

specific hederagenin-derived monodesmosidic saponins, each bearing two hexoses, are

associated for this defense.

Taken together, this work introduces a novel application of metabolic engineering to

unravel ecological functions of plant metabolites and offers promising strategies for crop

protection through fine-tuned biosynthetic pathway manipulation.

Acknowledgements

We thank Thure Hauser providing Plutella *xylostella*, Niels Agerbirk providing

wildtype *Barbarea vulgaris* G- and P-type seeds, Barry Cohen and Hanne Volpin help for

their help in setting up the transformation system, and Jack Olsen and Mariela Alejandra

González Ramírez for their assistance in running analytical instruments. This work was

financially supported by Novo Nordisk Foundation (EcoSap NNF20OC0060298 to SB), the

China Scholarship Council (CSC, 202108330041 to JS) and Marie Sklodowska-Curie

Individual Fellowship (MSCA-IF 752437 to PC).

Conflicts of interest

Authors declare that there are no conflicts of interest.

Author contributions

JS, PC, and SB designed the research and wrote the paper; JS and PC performed the

research; JS, PC, JG and SB analysed the data; JG provided the data analysis strategy and

standards.

References

Agerbirk N, Hansen CC, Olsen CE, Kiefer C, Hauser TP, Christensen S, Jensen KR, Orgaard
M, Pattison DI, Lange CBA, Cipollini D, Koch MA (2021) Glucosinolate profiles and
phylogeny in *Barbarea* compared to other tribe Cardamineae (Brassicaceae) and
Reseda (Resedaceae), based on a library of ion trap HPLC-MS/MS data of
reference desulfoglucosinolates. Phytochemistry **185:** 112658

Agerbirk N, Olsen CE, Bibby BM, Frandsen HO, Brown LD, Nielsen JK, Renwick JAA (2003)
A saponin correlated with variable resistance of Barbarea vulgaris to the
diamondback moth *Plutella xylostella*. Journal of Chemical Ecology **29:** 1417-1433

Agerbirk N, Olsen CE, Nielsen JK (2001) Seasonal variation in leaf glucosinolates and
insect resistance in two types of *Barbarea vulgaris ssp. arcuata*. Phytochemistry
58: 91-100

Augustin JM, Drok S, Shinoda T, Sanmiya K, Nielsen JK, Khakimov B, Olsen CE, Hansen
EH, Kuzina V, Ekstrøm CT, Hauser T, Bak S (2012) UDP-glycosyltransferases from
the UGT73C subfamily in *Barbarea vulgaris* catalyze sapogenin 3-O-glucosylation
in saponin-mediated insect resistance. Plant Physiology **160:** 1881-1895

Augustin JM, Kuzina V, Andersen SB, Bak S (2011) Molecular activities, biosynthesis and
evolution of triterpenoid saponins. Phytochemistry **72:** 435-457

Beran F, Petschenka G (2022) Sequestration of plant defense compounds by insects: from
mechanisms to insect–plant coevolution. Annual Review of Entomology **67:** 163-
180

Beringer J, Chen W, Garton R, Sardesai N, Wang PH, Zhou N, Gupta MJ, Wu HX (2017)
Comparison of the impact of viral and plant-derived promoters regulating
selectable marker gene on maize transformation and transgene expression. Plant
Cell Reports **36:** 519-528

Byrne SL, Erthmann PO, Agerbirk N, Bak S, Hauser TP, Nagy I, Paina C, Asp T (2017) The
genome sequence of *Barbarea vulgaris* facilitates the study of ecological
biochemistry. Scientific Reports **7:** 40728

Cárdenas PD, Almeida A, Bak S (2019) Evolution of structural diversity of triterpenoids.
Frontiers in plant science **10:** 1523

Chwalek M, Lalun N, Bobichon H, Plé K, Voutquenne-Nazabadioko L (2006) Structure-
activity relationships of some hederagenin diglycosides:: Haemolysis, cytotoxicity
and apoptosis induction. Biochimica Et Biophysica Acta-General Subjects **1760:**
1418-1427

Confalonieri M, Carelli M, Gianoglio S, Moglia A, Biazzi E, Tava A (2021) CRISPR/Cas9-
mediated targeted mutagenesis of *CYP93E2* modulates the triterpene saponin
biosynthesis in *Medicago truncatula*. Frontiers in Plant Science **12:** 6090231

Cui CJ, Yang YQ, Zhao TY, Zou KK, Peng CY, Cai HM, Wan XC, Hou RY (2019) Insecticidal
activity and insecticidal mechanism of total saponins from *Camellia oleifera*.
Molecules **24:** 4518

Dolma SK, Suresh PS, Singh PP, Sharma U, Reddy SGE (2021) Insecticidal activity of the
extract, fractions, and pure steroidal saponins of *Trillium govanianum* Wall. ex D.
Don for the control of diamondback moth (*Plutella xylostella* L.) and aphid (*Aphis
craccivora* Koch). Pest Management Science **77:** 956-962

Erthmann PO, Agerbirk N, Bak S (2018) A tandem array of UDP-glycosyltransferases from
the UGT73C subfamily glycosylate sapogenins, forming a spectrum of mono- and
bisdesmosidic saponins. Plant Molecular Biology **97:** 37-55

Fürstenberg-Hägg J, Zagrobelny M, Bak S (2013) Plant defense against insect herbivores.
International Journal of Molecular Sciences **14:** 10242-10297

Gao GC, Lu ZX, Tao SH, Zhang S, Wang FZ (2011) Triterpenoid saponins with antifeedant
activities from stem bark of *Catunaregam spinosa* (Rubiaceae) against *Plutella
xylostella* (Plutellidae). Carbohydrate Research **346:** 2200-2205

Günther J, Erthmann PØ, Khakimov B, Bak S (2021) Reciprocal mutations of two
multifunctional β-amyrin synthases from *Barbarea vulgaris* shift α/β-amyrin
ratios. Plant Physiology **188:** 1483-1495

Hauser TP, Toneatto F, Nielsen JK (2012) Genetic and geographic structure of an insect
resistant and a susceptible type of *Barbarea vulgaris* in western Europe.
Evolutionary Ecology **26:** 611-624

Hayta S, Smedley MA, Clarke M, Forner M, Harwood WA (2021) An efficient
Agrobacterium-mediated transformation protocol for hexaploid and tetraploid
wheat. Current Protocols **1:** e58

**Hodgins CL, Salama EM, Kumar R, Zhao Y, Roth SA, Cheung IZ, Chen JY, Arganosa GC,
Warkentin TD, Bhowmik P, Ham BK, Ro DK** (2024) Creating saponin-free yellow
pea seeds by CRISPR/Cas9-enabled mutagenesis on β-amyrin synthase. Plant
Direct **8:** e563

Huang FQ, Dong XS, Yin XJ, Fan Y, Fan YM, Mao CC, Zhou W (2020) A mass spectrometry
database for identification of saponins in plants. Journal of Chromatography A
1625: 461296

Hussain M, Debnath B, Qasim M, Bamisile BS, Islam W, Hameed MS, Wang L, Qiu D
(2019) Role of saponins in plant defense against specialist herbivores. Molecules
24: 2067

Khakimov B, Amigo JM, Bak S, Engelsen SB (2012) Plant metabolomics: resolution and
quantification of elusive peaks in liquid chromatography–mass spectrometry
profiles of complex plant extracts using multi-way decomposition methods.
Journal of Chromatography a **1266:** 84-94

Khakimov B, Kuzina V, Erthmann PØ, Fukushima EO, Augustin JM, Olsen CE, Scholtalbers
J, Volpin H, Andersen SB, Hauser TP, Muranaka T, Bak S (2015) Identification and
genome organization of saponin pathway genes from a wild crucifer, and their use
for transient production of saponins in *Nicotiana benthamiana*. The Plant Journal
84: 478-490

Khakimov B, Tseng LH, Godejohann M, Bak S, Engelsen SB (2016) Screening for
triterpenoid saponins in plants using hyphenated analytical platforms. Molecules
21: 1614

Kuzina V, Ekstrøm CT, Andersen SB, Nielsen JK, Olsen CE, Bak S (2009) Identification of
defense compounds in Barbarea vulgaris against the herbivore Phyllotreta
nemorum by an ecometabolomic approach. Plant physiology **151:** 1977-1990

Kuzina V, Nielsen JK, Augustin JM, Torp AM, Bak S, Andersen SB (2011) *Barbarea vulgaris*
linkage map and quantitative trait loci for saponins, glucosinolates, hairiness and
resistance to the herbivore *Phyllotreta nemorum*. Phytochemistry **72:** 188-198

Liu Q, Khakimov B, Cárdenas PD, Cozzi F, Olsen CE, Jensen KR, Hauser TP, Bak S (2019)
The cytochrome P450 CYP72A552 is key to production of hederagenin-based
saponins that mediate plant defense against herbivores. New Phytologist **222:**
1599-1609

Moses T, Papadopoulou KK, Osbourn A (2014) Metabolic and functional diversity of
saponins, biosynthetic intermediates and semi-synthetic derivatives. Critical
reviews in biochemistry and molecular biology **49:** 439-462

Nielsen JK, Nagao T, Okabe H, Shinoda T (2010) Resistance in the plant, *Barbarea*
vulgaris, and counter-adaptations in flea beetles mediated by saponins. Journal of
Chemical Ecology **36:** 277-285

Patel VS, Chhalotiya UK, Patel SB, Nuruddin J (2020) Simultaneous quantification of
betulinic acid, lupeol, and β-sitosterol in *Madhuca longifolia* methanolic extract of
bark by liquid chromatography-tandem mass spectrometric method. Journal of
AOAC INTERNATIONAL **104:** 498-505

Seger M, Ortega JL, Bagga S, Sengupta-Gopalan C (2009) Repercussion of mesophyll-
specific overexpression of a soybean cytosolic glutamine synthetase gene in alfalfa
(*Medicago sativa* L.) and tobacco (*Nicotiana tabaccum* L.). Plant Science **176:** 707-
707

Shinoda T, Nagao T, Nakayama M, Serizawa H, Koshioka M, Okabe H, Kawai A (2002)
Identification of a triterpenoid saponin from a crucifer, *Barbarea vulgaris*, as a
feeding deterrent to the diamondback moth, *Plutella xylostella*. Journal of
Chemical Ecology **28:** 587-599

Stevenson PC, Nyirenda SP, Veitch NC (2010) Highly glycosylated flavonoids from the
pods of *Bobgunnia madagascariensis*. Tetrahedron Letters **51:** 4727-4730

Tautenhahn R, Patti GJ, Rinehart D, Siuzdak G (2012) XCMS online: a web-based platform
to process untargeted metabolomic data. Analytical Chemistry **84:** 5035-5039

Tian X, Li Y, Hao N, Su X, Du J, Hu J, Tian X (2021) The antifeedant, insecticidal and insect
growth inhibitory activities of triterpenoid saponins from *Clematis aethusifolia*
Turcz against *Plutella xylostella* (L.). Pest Management Science **77:** 455-463

Waller M, Frangedakis E, Marron AO, Sauret-Güeto S, Rever J, Sabbagh CRR, Hibberd
JM, Haseloff J, Renzaglia KS, Szövényi P (2023) An optimized transformation
protocol for *Anthoceros agrestis* and three more hornwort species. Plant Journal
114: 699-718

Xu R, Fazio GC, Matsuda SPT (2004) On the origins of triterpenoid skeletal diversity.
Phytochemistry **65:** 261-291

Yactayo-Chang JP, Tang HV, Mendoza J, Christensen SA, Block AK (2020) Plant defense
chemicals against insect pests. Agronomy **10:** 1156