1 **Integrative modeling of seasonal influenza evolution via AI-powered antigenic**
2 **cartography**
3

4 Jumpei Ito[1,2,12,13], Shusuke Kawakubo[1,3,12], Hiroaki Unno[1], Adam Strange[1], Spyros
5 Lytras[1,4], Kaho Okumura[1], Alice Lilley[5], Ruth Harvey[5], Nicola Lewis[5], Kei Sato[1,2,3,6,7,8,9,10,11]
6

[1] Division of Systems Virology, Department of Microbiology and Immunology, The Institute
of Medical Science, The University of Tokyo, Tokyo, Japan
[2] International Vaccine Design Center, The Institute of Medical Science, The University of
Tokyo, Tokyo, Japan
[3] Biosystem Diversity Research Group, Molecular Biosystems Research Institute, National
Institute of Advanced Industrial Science and Technology, Tsukuba, Japan
[4] MRC-University of Glasgow Centre for Virus Research, Glasgow, UK
[5] Worldwide Influenza Centre, The Francis Crick Institute, London, United Kingdom
[6] Graduate School of Medicine, The University of Tokyo, Tokyo, Japan
[7] Graduate School of Frontier Sciences, The University of Tokyo, Kashiwa, Japan
[8] International Research Center for Infectious Diseases, The Institute of Medical Science,
The University of Tokyo, Tokyo, Japan
[9] Collaboration Unit for Infection, Joint Research Center for Human Retrovirus infection,
Kumamoto University, Kumamoto, Japan
[10] Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand
[11] Duke-NUS Medical School, Singapore
[12] These authors contributed equally
[13] Lead contact
25

26 Corresponding authors:
27 jampei@g.ecc.u-tokyo.ac.jp (Jumpei Ito)
28 KeiSato@g.ecc.u-tokyo.ac.jp (Kei Sato)
29

30 Lead contact:
31 jampei@g.ecc.u-tokyo.ac.jp (Jumpei Ito)
32

33 X / Twitter:
34 jampei2

36 Short title: AI-powered antigenic cartography
37 Keywords: Seasonal influenza virus; Antigenic cartography; Evolution; Vaccine
38 development; Protein language model
39

ABSTRACT

Seasonal influenza viruses evade host immunity through rapid antigenic evolution.

Antigenicity is assessed by serological assays and typically visualized as antigenic maps,

which represent antigenic differences among virus strains. However, conventional maps

cannot directly infer the antigenicity of unexamined variants from their genotypes. Here, we

present PLANT, a protein language model that projects influenza A/H3N2 viruses onto an

antigenic map using HA protein sequences. Using PLANT-based cartography, we show

that (i) H3N2 antigenic evolution accelerates during periods of disrupted global circulation,

(ii) antigenic novelty accounts for a substantial portion of viral fitness advantage, and (iii)

vaccine strains are often antigenically distant from circulating viruses. We further propose a

PLANT-based framework for selecting vaccine strains with improved antigenic match than

the WHO-recommended strains. This study provides a statistical foundation for integrated

modeling of viral genotype, antigenicity, and fitness, offering quantitative insights into

seasonal influenza virus evolution and supporting rational vaccine design.

Introduction

One of the major challenges in controlling viral infectious diseases is the rapid evolution of viruses. Seasonal influenza viruses undergo continuous antigenic changes, enabling them to evade population immunity established by prior infection and vaccination [1]. The emergence of immune-escape variants poses a significant risk of larger epidemics [2-4]. Furthermore, to develop effective vaccines against circulating strains, it is essential to characterize their antigenicity and to select vaccine strains that closely match them [1,5,6]. Therefore, the antigenicity of emerging seasonal influenza variants is continuously monitored through intensive efforts by multiple research groups, including the Worldwide Influenza Centre [7].

Differences in antigenicity between viral strains can be quantified by antigenic distance, which reflects the extent to which humoral immunity elicited by a reference strain can neutralize a target strain. This distance is measured using serological assays—such as neutralization or hemagglutination inhibition (HI) assays—by testing sera from animals infected with a reference strain against the target strain [8]. Antigenic distance is defined as the difference in the $\log_2$ HI or neutralization titers between homologous and heterologous strain pairs [2,8]. To comprehensively characterize the antigenicity of a given variant, it is informative to measure antigenic distances from multiple reference strains. However, the resulting data are high-dimensional and challenging to interpret intuitively. Antigenic cartography addresses this issue by embedding viral strains into a low-dimensional space using multidimensional scaling (MDS) [8]. Antigenic cartography has been widely used to identify immune-escape variants, guide vaccine strain selection, and track antigenic evolution [1,9]. For instance, vaccine strains have been updated to satisfy the quantitative criterion that their antigenic distance from circulating strains remains within two units [8].

Conventional antigenic cartography cannot be used to predict the antigenicity of strains that have not been experimentally characterized [8]. If the antigenicity of newly emerged variants could be inferred from their genotypes, such as the sequences of viral proteins, it would allow for rapid identification of immune-escape variants based solely on viral genome surveillance data. Also, this capability could partially replace and reduce reliance on labor-intensive serological assays. Most importantly, such a predictive model would enable the construction of comprehensive antigenic maps that encompass all virus strains with available sequence data. When combined with phylogenetic and epidemic dynamics modeling, these maps could offer quantitative insights into the mechanisms of viral evolution and epidemic dynamics, while also supporting more effective vaccine strain selection.

Protein language models (pLMs) are a class of large language models trained not on natural language but on vast collections of protein sequences [10]. The development of pLMs has enabled flexible genotype-to-phenotype modeling of proteins. Notably, pLMs embed protein sequences into a semantic space that captures structural, functional, and evolutionary characteristics of the proteins [10,11]. Previous studies have shown that antigenic viral proteins, such as the hemagglutinin (HA) of influenza, are embedded in ways that

reflect their antigenic similarity [11]. In this study, we developed PLANT (Protein Language

Model for Antigenic cartography), a model that predicts the antigenic map coordinates of

seasonal influenza A/H3N2 viruses from the amino acid sequences of the HA1 domain, by

leveraging the semantic space of a pLM. We further demonstrate the utility of PLANT-

generated antigenic maps for elucidation of viral evolutionary patterns, identification of

immune-escape variants, and improvement of vaccine strain selection.

Results

Training of PLANT

We first developed a novel algorithm we term pLM-based MDS to convert the semantic space of a pLM into an antigenic map (**Figure 1A**). In this framework, a pLM augmented with dimensionality reduction layers embeds the HA1 sequences of viral strains into a low-dimensional semantic space. In this space, pairwise distances between the embedded sequences—referred to as semantic distances—are then computed. The model is trained to minimize the discrepancy between these semantic distances and experimentally measured antigenic distances (termed observed antigenic distances). Through this training, the semantic space is transformed into an antigenic map that reflects antigenic relationships. This method can be regarded as an extension of MDS used in conventional cartography, augmented by the semantic space of a pLM. Importantly, unexamined virus strains can be embedded into the antigenic map constructed using pLM-based MDS, based on their HA1 sequences via the pLM.

PLANT comprises two main components: a cartography module, which constructs the antigenic map using pLM-based MDS, and a systematic error module, which corrects experimental biases arising from differences in assay conditions (**Figure 1B**). The cartography module utilizes the 650-million (M)-parameter version of ESM-2 [12] as its pLM. Given a pair of HA1 sequences for a reference and a target strain, along with metadata describing the experimental settings, PLANT predicts their coordinates on the antigenic map, the Euclidean distance between them (termed cartographic distance), and the systematic error–corrected distance (termed antigenic distance). If an unpaired HA1 sequence is provided, PLANT returns its predicted coordinate. Unlike the conventional method [8], the cartographic distance predicted by PLANT is symmetric; it remains unchanged regardless of which strain is designated as the reference.

For model training, we used HI assay data for seasonal influenza A (H3N2) viruses collected between 1968 and January 2024 by the Worldwide Influenza Centre, along with corresponding HA1 protein sequences obtained from GISAID (https://gisaid.org/). The final dataset comprised 63,118 observed antigenic distances measured between 208 reference strains and 5,271 target strains (**Figure S1A**). Because HI assays are typically conducted between temporally proximate strains, the resulting antigenic distance matrix is highly sparse, with most entries missing (**Figure 1A, bottom left; Figure S1A**) [13]. Moreover, since this dataset was constructed by combining experimental data accumulated over several decades, it inevitably incorporates various sources of systematic error, as mentioned in previous studies [14-16]. We randomly split the dataset along the target virus axis, ensuring that no antigenic distance measurements involving the same target strain appeared in both the training and test sets. This dataset was split without considering the temporal information of virus strains and is hereafter referred to as the "full dataset."

Antigenic map generated by PLANT

When the conventional antigenic cartography tool Racmacs was applied to the full dataset

(prior to train–test splitting), the resulting map failed to preserve the temporal continuity of
antigenic evolution (**Figure 1C**). In sharp contrast, the antigenic map generated by PLANT,
trained on the full dataset (after train–test splitting), successfully captured a continuous
trajectory of antigenic drift spanning the entire evolutionary timeline of human H3N2 viruses
(**Figure 1D, left; Table S1**). This contrast highlights the importance of incorporating viral
genetic information and correcting for systematic error in order to construct effective
antigenic maps from this dataset, in line with previous reports [14]. Details of PLANT's
predictive performance are presented in a later section. Furthermore, because PLANT can
predict antigenic map coordinates even for strains lacking serological measurements, it
enables the construction of a comprehensive map encompassing all 151,784 H3N2 strains
with fully determined HA sequences collected between 1968 and January 2024 (**Figure 1D,
left**).
To characterize the discontinuous nature of H3N2 antigenic evolution, we
performed clustering based on the predicted coordinates and assigned strains to 20
antigenic clusters (**Figure 1D, right**). As expected, these clusters showed strong
concordance with established H3N2 classification systems, such as the Nextclade
subclade designations [17] (**Figure S1B; Table S1**). Since Nextclade designations are
unavailable for older strains (e.g., those sampled before 2010), our downstream analyses
primarily relied on the antigenic cluster classification. We next constructed a distance
matrix representing pairwise cartographic distances between strains. This analysis
confirmed that strains within the same antigenic cluster were tightly co-localized in the
antigenic space (**Figure 1E**). Furthermore, hierarchical clustering based on cluster
centroids revealed a higher-order structure, grouping 19 clusters (excluding c1, which
mainly comprised pre-2000 strains) into four metaclusters: m1 (c2–5), m2 (c6–9), m3 (c10–
17), and m4 (c18–20) (**Figure 1F**). Metaclusters m2, m3, and m4 corresponded to
subclades A, B–F, and G–J in Nextclade, respectively (**Figure S1B**). Together, these
results demonstrate that the PLANT-derived antigenic map effectively reconstructs over
five decades of H3N2 antigenic evolution, capturing both gradual drift and discrete changes
between major antigenic lineages.
We found that antigenic diversity varied substantially across seasons, as
quantified by the cartographic distances from the centroid of epidemic strains in each
season (**Figure 1G, top; Table S1**). Notably, seasons marked by the co-circulation of
multiple antigenic clusters—such as the 2009 Southern Hemisphere (SH; February–
August) season and the 2019 Northern Hemisphere (NH; September–January) season—
exhibited pronounced increases in antigenic diversity, with median distances approaching 2
antigenic units (**Figure 1G, bottom**). These findings suggest that during such antigenically
diverse seasons, selecting a single vaccine strain that is antigenically proximate to all
circulating variants may be inherently difficult. In contrast, antigenic diversity declined
following the emergence of rapidly expanding lineages, which swept through the population
and displaced pre-existing strains.
184

Prediction performance of PLANT

We first evaluated the predictive performance of PLANT trained on the full dataset, which had been trained using a randomly split dataset based on target virus strains, as described earlier. Prediction accuracy, measured by mean absolute error (MAE) and Pearson correlation coefficient, was assessed for each influenza season from 2010SH to 2023NH (**Figure S2A**). In the analyzed dataset, HI assay results showed a mean standard deviation of 0.42 among replicates conducted under identical experimental conditions, reflecting the intrinsic noise of the assay. Nevertheless, PLANT achieved a seasonal average MAE of 0.83 and a correlation coefficient of 0.78, indicating robust predictive performance. Moreover, PLANT's antigenic distance predictions outperformed both the cartographic distances (i.e., without systematic error correction) and the scale-adjusted semantic distances derived from the non-finetuned ESM-2 (**Figure S2A**). These results demonstrate that, under the interpolation condition, PLANT can accurately predict the antigenicity of previously unseen virus variants.

To evaluate the extrapolation performance of PLANT, we next assessed its ability to predict the antigenicity of future variants (**Figure 2**). Based on virus collection dates, the dataset was divided into "past" and "future" season datasets (**Figure 2A**). PLANT was trained exclusively on the past dataset and evaluated on the future dataset. This procedure was repeated for 19 different cutoff seasons, ranging from 2014SH to 2023SH, following the settings in Shah et al.[15]. We compared PLANT's performance against two benchmarks: (i) a PLANT model trained on the full dataset, and (ii) AdaBoost, a state-of-the-art model for antigenicity prediction developed by Shah et al [15]. Like PLANT, AdaBoost predicts antigenic distances from HA1 sequences and experimental metadata, but unlike PLANT, it cannot predict an antigenic map.

As expected, both PLANT and AdaBoost trained only on past data showed reduced accuracy compared to the PLANT model trained on the full dataset, and prediction accuracy generally declined with increasing temporal distance from the cutoff season (**Figures 2B, 2C, S2B, and S2C**). Comparing PLANT and AdaBoost, we found that their MAE values were similar in early post-cutoff seasons. However, after +7 seasons (approximately 3.5 years), AdaBoost's performance declined (i.e., higher MAE) markedly, while PLANT maintained stable accuracy (**Figure 2C**). This decline was particularly evident after the 2021NH season, coinciding with the emergence of an antigenically distinct viral lineage, metacluster m4 (**Figures 2B and 1G**). These findings suggest that PLANT outperforms AdaBoost in MAE-based performance for distant future variants. For the Pearson correlation coefficient, both models performed similarly in early post-cutoff seasons (within +3 seasons). However, in later seasons, PLANT's correlation tended to decline more than AdaBoost's. Taken together, these results indicate that PLANT achieves extrapolation performance comparable to that of the existing state-of-the-art model, at least for several seasons into the future.

Rate of antigenic evolution

Accurately estimating the rate of viral antigenic evolution is crucial for understanding the epidemic factors and mutations driving antigenic change, as well as for optimizing the frequency of vaccine strain updates. To investigate this rate in human H3N2 viruses, we performed an integrative analysis combining PLANT-based antigenic maps with phylogenetic modeling. Specifically, we projected the time-calibrated phylogenetic tree of H3N2 viruses onto the PLANT-based antigenic map and inferred branch-specific rates of antigenic evolution (**Figures 3A and 3B**).

To estimate the average rate of antigenic evolution in H3N2, we calculated the cumulative antigenic change along the branches comprising the trunk of the phylogeny—defined as the set of branches with ≥10 descendant tips—from the root (**Figure 3C**). We then performed a linear regression to model the cumulative cartographic distances of both external and internal nodes as a function of their sampling or inferred dates. This analysis revealed an average antigenic evolution rate of 1.61 antigenic units per year. Although this estimated slope is slightly higher than previously reported estimates (e.g., 1.2 antigenic units in Smith et al. [8]), this discrepancy may stem from differences in the time periods analyzed, as the earlier study focused on an older dataset (i.e., 1968–2003).

Consistent with the pattern observed in **Figure 1E**, the rate of antigenic change during H3N2 evolution appears to vary over time (**Figure 3B**). By assessing antigenic change on the H3N2 phylogeny, we found that external branches tend to accumulate a greater degree of antigenic changes compared to internal branches (**Figure S3A**). This suggests that a substantial proportion of mutations leading to larger antigenic change are not transmitted to subsequent lineages, likely due to their deleterious effects [18]. Furthermore, some of the internal branches exhibited markedly elevated antigenic changes, with 12 branches showing cartographic distances exceeding the three standard deviations over the mean (1.65 units). As expected, some of these corresponded to antigenic cluster transitions (**Figure 3B**). Notably, the rate of antigenic evolution increased substantially at the boundaries between antigenic metaclusters (**Figures 1F, 3C, and 3D**). These transitions occurred during 2009NH–2010NH (m1–m2 boundary), 2013SH–2014SH (m2–m3 boundary), and 2019NH–2020NH (m3–m4 boundary), with corresponding antigenic evolution rates reaching 2.22, 2.39, and 4.82 units per year, respectively (**Figure 3C**). Interestingly, periods of accelerated antigenic evolution coincided with major interruptions in global H3N2 circulation—specifically, the 2009 H1N1 pandemic, the 2013–2014 H1N1 epidemic, and the COVID-19 pandemic (**Figure 3E**). Among these, the most pronounced acceleration occurred during the COVID-19 pandemic, which marked the most severe interruption of global influenza circulation [19]. These observations raise the possibility that such large-scale interruptions in continuous H3N2 circulation may facilitate or accelerate antigenic evolution (see **Discussion**).

Mutations driving antigenic evolution

To identify amino acid substitutions responsible for antigenic changes during H3N2 evolution, we estimated the effect of individual substitutions on cartographic distance by

analyzing a phylogenetic tree embedded in the antigenic map. Cartographic distance is

defined as non-negative, and we assumed that the total antigenic change between a parent

and its descendant strain results from the sum of non-negative contributions of individual

amino acid substitutions, in accordance with Neher et al. [2]. Based on this assumption, we

implemented a Bayesian non-negative linear model in which antigenic change per branch

is explained as a non-negative linear combination of individual mutation effects (**Figure**

S3B). This analysis identified 195 amino acid mutations across 85 sites as having

statistically significant contributions to antigenic change (**Figures 3F and S3C; Table S2**).

As expected, the majority of these mutations were located within antigenic sites

known to be targeted by neutralizing antibodies (**Figures 3F–H; Table S3**). In particular,

antigenic sites A and B showed significant enrichment of impactful mutations, consistent

with a previous study [20]. Furthermore, significant mutations were identified at all seven

positions (145, 155, 156, 158, 159, 189, and 193) previously proposed to drive antigenic

cluster transitions [20]. Moreover, significant mutations were also observed outside these

canonical antigenic sites. For example, positions 1–10 and 223–227—regions

encompassing the undefined epitopes proposed by Shih et al. [21]—were enriched for

significant mutations, supporting a potential role for these regions in antigenic evolution.

Together, these results demonstrate that PLANT can effectively capture the impact of

individual mutations to antigenic changes and thus serves as a powerful tool for dissecting

the genetic basis of antigenic evolution.

Relationship between antigenic novelty and fitness advantage

Using PLANT, we constructed antigenic maps solely from genomic sequence data. This

enabled us to reconstruct both the epidemic dynamics and antigenic evolution of viruses

from the same genome surveillance data, thereby allowing the integrated modeling of viral

genotype, antigenicity, and fitness.

Using this framework, we first examined the relationship between *antigenic novelty*

and *fitness advantage*. Variants (i.e., groups of strains) that are antigenically distinct from

previously circulating strains can generally evade population immunity. Such variants with

higher antigenic novelty are thought to exhibit higher fitness advantage enabling them to

outcompete co-circulating variants [1]. However, the quantitative relationship between

antigenic novelty and fitness advantage—defined as the extent to which a variant's fitness

exceeds that of other concurrently circulating variants—remains poorly understood (**Figure**

4A). Therefore, we applied our integrative modeling approach using PLANT to address this

issue.

The fitness of a variant can be evaluated based on its effective reproduction

number (R_e), which represents the virus's growth potential in a population over time [22-27].

By applying a multinomial logistic model to time-series data of variant frequencies derived

from viral genome surveillance, the relative R_e values between variants can be estimated

[22-27]. Although this method assumes that the relative fitness of variants remains constant

over time—a simplification that may not always reflect real-world dynamics—it is a widely

used approach for comparing the fitness of co-circulating variants based solely on genome
surveillance data [22-27].
We first classified H3N2 strains into *antigenic groups*—variants defined based on
antigenicity—using a Gaussian mixture model (GMM) clustering on the PLANT-derived
antigenic map, resulting in 134 groups (g1–134; **Figure 4B**). The analysis focused on the
span between the 2010SH and 2019NH seasons, before the epidemic interruption by the
COVID-19 pandemic. We then estimated their epidemic dynamics and their relative R_e
using a Bayesian multinomial logistic model (**Figure 4C**). As expected, antigenic groups
that emerged later tended to exhibit higher R_e, reflecting temporal replacement dynamics
and aligning with findings from previous studies [25,26] (**Figure 4D**). An exponential
regression analysis showed that the R_e of H3N2 viruses has increased by approximately
2.22% per year.
As a proxy of antigenic novelty, the median cartographic distance from prior
epidemic strains collected within the preceding 1.5 years was quantified. This time window
was selected to capture strains from the most recent epidemic season in the same
hemisphere. Fitness advantage was defined as the extent to which a variant's fitness
exceeds the expected value at its time of emergence. Accordingly, we used the residual
from the exponential regression curve shown in **Figure 4D** as a proxy for fitness advantage
(**Figure 4E**).
Correlation analysis revealed a positive association between antigenic novelty and
fitness advantage (Spearman's ρ = 0.543; **Figure 4F**). Using a hierarchical Bayesian
model, we then estimated the effect size both as an average across seasons and for each
season individually (**Figure 4G**). The effect size was positive in all seasons except
2014NH, and an overall mean estimate was 1.47. This implies that a one antigenic unit
increase in antigenic novelty corresponds to an average relative R_e increase of 1.47%.
Combining this with the estimated antigenic evolution rate (1.61 antigenic units per year;
Figure 3C), the annual increase in relative R_e attributable to antigenic change is inferred to
be at 2.37% (1.47% per antigenic unit × 1.61 antigenic units per year), which closely aligns
with the directly estimated 2.22% increase in **Figure 4D**. Furthermore, the Bayesian R-
squared of the model was 0.35, suggesting that 35% of the measured variation in fitness
advantage can be explained by antigenic novelty, together with season-specific effects
(**Figure 4H**). These findings collectively suggest that antigenic change is a major driver of
replacement epidemic variants in H3N2 influenza viruses.
341
Automated identification of emerging immune-escape variants
We further applied the integrative framework that combines PLANT with epidemic
dynamics modeling to automatically identify variants with both distinct antigenicity and high
epidemic potential. This method follows the scheme described above: constructing an
antigenic map using PLANT, classifying virus strains into antigenic groups, and estimating
the relative R_e of each group. Because this approach defines variants (i.e., antigenic
groups) based on antigenicity and estimates their fitness accordingly, it eliminates the need

for experts to manually define variants or cross-reference fitness estimates with an
antigenic map.
In this analysis, PLANT was trained on data up to the 2023NH season, and the
antigenic map was constructed using viruses collected through the 2024SH season
(**Figures 5A–C**). This one-season delay in PLANT's training data reflects the typical lag in
the availability of serological assay results. Using this setup, virus strains sampled between
the 2022SH and 2024SH seasons were classified into 30 antigenic groups (g1–30).
The top six antigenic groups ranked by relative R_e exhibited significantly higher R_e
values than the remaining groups (**Figure 5D**). According to the Nextclade subclade
classification, all six high-fitness groups belonged to subclade J.2 (including J.2.1), which
had been rapidly expanding and had become the dominant lineage as of July 2025
(https://nextstrain.org/seasonal-flu/h3n2). Notably, these J.2-related groups exhibited
distinct mutational profiles, with several harboring key amino acid substitutions known to
contribute to antigenic change (**Figure 5E**). For example, antigenic groups g29, g30, and
g26—ranked 1st, 2nd, and 6th in fitness, respectively—contained the substitutions S145N,
N158K, and I160X (X denoting any amino acid). Some strains in g30 carried S145N in
addition to N158K, the hallmark mutation of this group. These mutations are revertant
substitutions that occurred at known hotspots for recurrent mutations during H3N2 virus
evolution [1,20]. On the antigenic map, antigenic group g29 (rank 1) was located near other
J.2 lineage viruses, whereas g30 (rank 2) and g26 (rank 6) were positioned more distantly,
suggesting that they are antigenically distinct from the main J.2 lineage (**Figures 5C and**
5F). In particular, antigenic group g26 was located farthest from the other J.2 strains,
consistent with the fact that g26 carries substitutions at position 160, in which substitutions
showed the highest estimated impact on antigenicity (**Figure 3H**). Also, isoleucine at
position 160 is a hallmark of antigenic cluster c20 and was acquired on the stem branch of
this cluster (**Figure 3C**). Antigenic group g26 comprises strains bearing various amino
acids at position 160, suggesting that the loss of isoleucine at this site plays a key role in
antigenic change, regardless of the substituting residue (**Figures 5G**). Together, these
findings indicate that subclade J.2 encompasses multiple antigenically divergent lineages.
Finally, to assess whether these J.2 descendant strains carrying distinctive
mutations actually expanded, we analyzed viral genomic surveillance data from the
2024NH season, which were used in neither PLANT training nor fitness estimation (**Figure**
5H). The J.2 lineage carrying the S145N substitution (J.2+S145N), corresponding to
antigenic group g29—the variant with the highest estimated fitness—began expanding in
2023NH and accounted for >30% of circulating strains at the beginning of 2024NH. In
contrast, J.2+I160X (g26) and J.2+N158K&S145N (part of g30) remained at low prevalence
throughout the 2023NH and 2024SH seasons, making signs of expansion difficult to detect
visually. However, both lineages subsequently expanded after 2024NH, eventually
comprising approximately 5% of circulating strains at their respective peaks. The
J.2+N158K lineage, which also falls within g30, remained at low prevalence by the end of
2024NH. Collectively, these findings demonstrate that the PLANT-based framework

enables the early and automated identification of antigenically novel variants with the
potential for epidemiological expansion.
392
Evaluation of historical vaccine strains
Vaccine strain selection involves choosing the most appropriate strain—among those
circulating up to a given point in time—to serve as the antigen in upcoming vaccination
campaigns. For seasonal influenza, vaccine strains are updated up to twice per year in
response to antigenic changes in circulating viruses [1,6,7]. A key challenge in this process is
the substantial lead time required for vaccine production and distribution, which
necessitates selecting a strain that is predicted to be antigenically similar to viruses
expected to circulate seven to 12 months in the future. For the Northern and Southern
Hemispheres, vaccine strains are recommended each February and September,
respectively, through WHO-coordinated Vaccine Composition Meetings [1,6,7].
To systematically evaluate the historical vaccine strains, we calculated the
cartographic distance between each virus and its nearest vaccine strain used since 2001,
based on the antigenic map generated by PLANT trained on the full dataset (**Figures 6A**
and 6B; Table S1). This analysis revealed that 97.2% of virus strains were located within 2
antigenic units of their closest vaccine strain. While some strains exceeded this threshold,
recent strains isolated after 2020 were underrepresented among them (**Figure 6B**). These
results suggest that updates to the H3N2 vaccine strain have successfully tracked the
major trajectory of antigenic evolution. In sharp contrast, when cartographic distances were
calculated between each virus and the vaccine strain actually used in the same season,
only 55.1% of viruses fell within 2 antigenic units. Furthermore, even among viruses
isolated after 2020, 20.3% of strains were more than 2 antigenic units away from the
corresponding vaccine strain. These findings indicate that vaccine strain updates have not
consistently kept pace with the rapid antigenic evolution of H3N2, resulting in frequent
mismatches at the time of vaccination.
417
An optimized framework for vaccine strain selection
To address the above issue, we developed a new vaccine strain selection scheme that
integrates PLANT with epidemic dynamics modeling (**Figure 6C**). In this framework,
PLANT is trained annually using genomic and HI assay data available as of January for the
Northern Hemisphere and August for the Southern Hemisphere. Based on the trained
model and sequence data available by those dates, we construct an antigenic map, classify
viruses into antigenic groups, and use a multinomial logistic model to forecast the epidemic
dynamics of each group during the vaccination period. The forecasted prevalence of each
antigenic group during the vaccination period is then computed, and these values are used
to assign weights to virus strains on the antigenic map. A weighted centroid is computed,
and the strain closest to this centroid is selected as the candidate vaccine strain.
We evaluated the performance of our vaccine selection scheme through
retrospective simulations spanning the 2015SH to 2023NH seasons. As a proxy for

antigenic match, we computed the median cartographic distance between the selected
vaccine strain and the viruses that circulated during the vaccination period, based on the
antigenic map by PLANT trained on the full dataset. These distances were compared
across four types of strains: (i) the actual WHO-recommended strains, (ii) the strains
nearest to the centroid of the most recent season, (iii) the strains selected by our proposed
framework, and (iv) hindsight-optimal strains—defined as those closest to future epidemic
strains, selected using future information (**Figure 6D**).
Among the strategies compared, the WHO-recommended strains satisfied the 2-
antigenic unit criterion in 8 out of 18 seasons, with a median antigenic distance of 2.40
units (**Figures 6D and S4A**). The nearest-centroid strains exhibited a similar median
distance. In contrast, our weighted centroid approach met the threshold in 12 seasons and
achieved a substantially lower median distance of 1.64 units. The lowest distance (0.32
units) was observed in the 2022NH season, due to the accurate prediction of the expansion
of the newly emerged antigenic cluster c20 (**Figure S4B**). Although the performance of our
method does not yet match the hindsight-optimal benchmark of 0.75 units, it represents an
average reduction of 0.76 antigenic units compared to the WHO-recommended strains.
Given that a one-unit antigenic distance corresponds to a two-fold change in neutralization
activity [8], this improvement translates to an estimated 1.69-fold increase in vaccine-induced
neutralization. Taken together, these findings demonstrate that integrating our automated
scheme provides a promising strategy for improving the accuracy and effectiveness of
H3N2 vaccine strain selection.
452

Discussion

We here present PLANT, a protein language model that projects influenza A/H3N2 viruses onto an antigenic map using only HA1 sequences. Unlike conventional cartography methods, PLANT enables antigenic mapping for all sequenced strains. This capability allows for the direct integration of antigenic cartography with genomic surveillance through phylogenetic and epidemic modeling. Using this unified framework, we demonstrate that: (i) H3N2 antigenic evolution accelerates during periods of disrupted global circulation (**Figure 3**); (ii) antigenic change is a major driver of variant replacement in H3N2 (**Figure 4**); (iii) emerging immune-escape variants with high epidemic potential can be identified automatically at an early stage (**Figure 5**); and (iv) vaccine strain selection can be made more predictive and effective (**Figure 6**). Together, these results highlight PLANT as a comprehensive platform for jointly modeling viral genotype, antigenicity, and fitness— advancing both our understanding of antigenic evolution and our capacity to control viral infectious diseases.

Although previous studies have developed numerous statistical and machine learning models to predict antigenic distances between pairs of virus strains, these models cannot directly generate antigenic maps [2,3,14-16,28,29]. As an *ad hoc* solution, some studies have used the predicted distances to construct antigenic maps using conventional cartography methods based on MDS [2,15,28,29]. This approach, however, requires reconstructing the entire map whenever new strains are added, leading to inconsistencies in the coordinate system across updates. PLANT avoids these issues by directly predicting antigenic map coordinates from HA1 sequences. This capability makes PLANT a practical and efficient tool for real-time monitoring of viral antigenic evolution in conjunction with routine viral genome surveillance.

The association between antigenic change and fitness advantage among variants has long been recognized in seasonal influenza viruses, particularly H3N2 [1]. However, previous analyses have largely relied on (i) a limited set of variants with available antigenic measurements, (ii) genetic distance as a proxy for antigenic distance, or (iii) season-level analyses based on average values of antigenic change and fitness advantage [2-4,23]. In this study, we conducted the first comprehensive assessment of the relationship between antigenicity and fitness advantage at the variant level (**Figure 4**). We estimated that a one-unit increase in antigenic novelty corresponds to a 1.47% increase in relative fitness, with a Bayesian R-squared value of 0.35. Although this indicates a substantial association, the effect size—as represented by the slope—was modest. This reflects the inherently limited annual fitness increase observed in H3N2, estimated at approximately 2.22% per year (**Figure 4D**). Such limited fitness gains lead to slower replacement of epidemic lineages over time, as previously discussed[1,18,30-32]. Importantly, this result suggests that immune-escape variants are unlikely to trigger exceptionally large-scale epidemic waves in H3N2. For instance, to achieve a 20% increase in fitness—commonly observed every few months in SARS-CoV-2 between 2020 and 2024 [25,26]—a variant would require approximately 13.6 antigenic units of change, equivalent to 8.4 years of antigenic evolution in H3N2. The small

494   effect size further supports the view that while the recurrent emergence of immune-escape

495   variants influences the magnitude of seasonal epidemics, it is not the primary driver of their

496   occurrence [1-4]. These findings demonstrate that PLANT offers a quantitative understanding

497   of the relationship between viral antigenicity and fitness advantage.

498   Global circulation of H3N2 has been intermittently disrupted by major respiratory

epidemics, including the 2009 H1N1 and COVID-19 pandemics, as well as the 2013–2014

H1N1 epidemic [4,19,33]. Our analysis suggests that the rate of antigenic evolution accelerated

during these periods of reduced circulation (**Figures 3C–E**). Previous studies have

proposed that H3N2 evolution is accelerated by population bottlenecks that usually arise

during seasonal transitions, when the dominant epidemic population shifts (e.g., from the

Northern to the Southern Hemisphere) [1]. Our findings support this hypothesis and further

extend it to more extreme scenarios involving global interruptions of viral circulation. The

importance of bottleneck effects in H3N2 evolution likely stems from the relatively modest

fitness advantages conferred by antigenic changes (**Figure 4G**). That is, because even

immune-escape variants require prolonged periods to reach fixation in large populations,

rapid antigenic evolution likely depends on the increased stochasticity introduced by

bottlenecks [30-32]. Also, previous studies proposed that the fixation of advantageous immune

escape mutations in the evolution of H3N2 is slowed down by the accumulation of sublethal

deleterious mutations, increasing fitness variance in the virus population [18,34]. Taking this

into consideration, our findings are consistent with a scenario where population

bottlenecks, resulting from the reduced circulation periods, decrease fitness variance and,

in turn, enable more efficient fixation of antigenic change mutations. Future studies should

aim to elucidate the evolutionary dynamics and specific mechanisms underlying these

patterns.

518   The antigenic map constructed by PLANT enables a systematic assessment of

antigenic similarity between vaccine strains and circulating viruses (**Figure 6**). This

property of our model can be used not only to retrospectively evaluate historical vaccine

choices but also to benchmark alternative selection strategies. Indeed, we demonstrated

that a simple PLANT-based framework can recommend vaccine candidates that are

antigenically closer to circulating strains than those recommended by the WHO. While

logistic regression is widely used to forecast the dynamics of emerging variants [23], our

framework seamlessly integrates this approach with antigenic map predictions by PLANT,

enabling the automated and efficient selection of vaccine strains. While real-world vaccine

selection must also consider practical constraints—such as strain availability and growth

efficiency in egg-based production [5]—our scheme offers a principled statistical framework

for optimizing strain selection. In the future, integrating our scheme with more accurate,

machine learning–based epidemic forecasting methods could further improve vaccine

strain selection.

532   This study has several limitations. First, the extrapolation performance of PLANT

for entirely novel future variants remains limited and does not clearly surpass that of the

state-of-the-art AdaBoost model [15] (**Figure 2**). Nevertheless, the key strength of PLANT lies

in its ability to construct antigenic maps and perform downstream quantitative analyses
under interpolative conditions. Moreover, because PLANT can be retrained within a few
hours on a single GPU, regularly updating the model with the latest data reduces reliance
on extrapolation and ensures adaptability to recent antigenic trends. Second, PLANT
defines cartographic distances symmetrically, meaning that a given genotype is mapped to
the same coordinates regardless of whether it is treated as a reference or a target strain.
While this differs from the original asymmetric implementation [8], we adopted the symmetric
formulation based on preliminary experiments that showed improved extrapolation
performance. Third, although we conducted retrospective validation using temporally split
datasets, further prospective validation with newly emerging viral sequences will be
necessary to rigorously assess PLANT's practical utility in real-time surveillance and
vaccine strain selection.
Despite these limitations, our findings highlight the utility of PLANT not only in
advancing our understanding of antigenic evolution but also in improving the control of viral
infectious diseases. Importantly, the PLANT framework is not limited to seasonal influenza;
it is broadly applicable to other viral pathogens for which serological data are available,
such as SARS-CoV-2, dengue virus, and highly pathogenic avian influenza, and is
extendable to future pandemic threats. By jointly modeling viral genotype, antigenicity, and
fitness, PLANT provides a powerful foundation for understanding, forecasting, and
ultimately mitigating viral epidemics.
555

Methods

Dataset preparation

556
557
We retrieved all available HA sequences of human H3N2 influenza viruses and their
associated metadata from GISAID (https://gisaid.org/) as of January 28, 2025. The
translated amino acid sequences of the HA1 domain were extracted and aligned using
Nextclade v3.14.0 [17], with A/Wisconsin/67/2005-egg (GenBank accession number:
CY163680.1) as the reference. Clade and subclade classifications were also assigned by
Nextclade. We excluded sequences from non-human hosts, sequences lacking collection
date information, and those containing undetermined amino acids in the HA1 domain. Also,
only HA1 sequences with exactly 329 amino acids were retained.

In addition, we collected HI assay data and related experimental metadata from
the annual and interim reports of the Worldwide Influenza Centre
(https://www.crick.ac.uk/research/platforms-and-facilities/worldwide-influenza-
centre/annual-and-interim-reports). The raw data up to the 2021NH season were derived
from Harvey et al. [16], and those through the 2024NH season were generously provided by
the Worldwide Influenza Centre.

HI assays were stratified into experimental batches based on the date of
experiment and the passage type of the reference strain, and observed antigenic distances
were computed within each batch. HI titers (T_{AB}) were measured for combinations of target
strain A and reference strain B, forming a matrix of HI measurements with not assigned
(NA) values, indicating strain pairs for which HI assays were not performed (**Figure S1A**).
The observed antigenic distance between strains A and B was calculated using the
following formula as described in previous studies [2,8]: $D_{AB} = \log_2(T_{AB}) - \log_2(T_{BB})$. The
observed antigenic distance represents the number of two-fold dilution steps by which the
titer for the heterologous strain A differs from that of the homologous strain B, both
measured against sera from ferrets infected with the reference strain B. Observed antigenic
distance is thus an integer value.

In some cases, the observed antigenic distance was negative, indicating a higher
titer for the heterologous strain pair than for the homologous strain pair—likely due to
experimental noise. To avoid potential adverse effects of such noise on model training,
measurements with values below –1 were excluded, while those equal to –1 were set to 0.

Each HI measurement was matched to its corresponding HA1 protein sequence
via strain names. After filtering, the final dataset comprised 63,118 observed antigenic
distances, measured between 208 reference strains and 5,271 target strains (**Figure S1A**).
Observed antigenic distances ranged from 0 to 8, with some values right-censored (e.g.,
>6).

Before training PLANT, the observed antigenic distances were scaled to a range
between 0 and 1. After inference, the predictions were rescaled to the original scale by
multiplying by a scaling factor of 8.

For the unsupervised training of PLANT, we constructed a dataset of HA protein
sequences spanning diverse HA subtypes of influenza A viruses from both human and

animal hosts. HA protein sequences were retrieved from the NCBI Influenza Virus

Database (https://www.ncbi.nlm.nih.gov/genomes/FLU/Database/nph-select.cgi). All unique

type A sequences at the amino acid level were downloaded, resulting in a total of 33,657

sequences (retrieved on April 11, 2024). Metadata including serotype and collection date

were also obtained for all sequences. To remove redundancy, the dataset was clustered

using MMSeqs2 [35] with a minimum sequence identity threshold of 99% (--cov-mode 0), and

a single representative sequence was retained from each cluster. Sequences containing

ambiguous characters (X or J) and those annotated as mixed serotypes were further

excluded, resulting in a final HA-all training dataset comprising 9,890 sequences. The HA1

domain was extracted from each sequence and used for downstream analysis.

Overview of PLANT

PLANT is a model that embeds each virus strain into a Euclidean space based solely on its

HA1 protein sequence. In this space—hereafter referred to as the antigenic map—virus

strains with identical HA1 sequences are assigned the same coordinates, regardless of

whether they serve as reference or target strains. The distance between strains, referred to

as the cartographic distance, is defined as the Euclidean distance between their embedded

coordinates in the antigenic map. Cartographic distance is a symmetric, non-negative value

that depends solely on the HA1 sequence.

616   In contrast, antigenic distances measured by serological assays are influenced not

only by true antigenic differences but also by variation in experimental conditions and

measurement error. Also, true antigenic differences are potentially affected by genotype

differences of viral components other than HA1, such as the HA2 domain or neuraminidase

(NA). In this study, we assume that observed antigenic distances consist of the sum of

three components: a cartographic distance derived from the HA1 sequence; a systematic

error associated with experimental conditions, including antigenic differences arising from

non-HA1 viral components; and a random error component.

624   During training, PLANT learns to match the observed antigenic distance with the

625   sum of a semantic distance (i.e., the distance between sequence embeddings in the

626   dimension-reduced semantic space computed by a pLM) and the estimated systematic

627   error. Through this process, the semantic distance is transformed into a cartographic

628   distance, and the semantic space is converted into an antigenic map space.

629   In addition to supervised learning based on pLM-based MDS, PLANT incorporates

630   a semi-supervised learning framework by utilizing unlabeled HA1 sequences (i.e.,

631   sequences without associated serological data). By imposing structural constraints on the

632   antigenic map through unsupervised losses, this framework enhances the generalization

633   ability of the model (see **Section Unsupervised learning loss**).

634   For inference, PLANT takes as input a pair of HA1 protein sequences (reference

635   and target strains) along with a set of categorical variables representing experimental

636   conditions. It computes antigenic map coordinates for both sequences, calculates the

637   cartographic distance, and adds the estimated systematic error to produce the predicted

638   antigenic distance. If only an unpaired sequence (e.g., a target strain) is provided, the
639   model outputs its coordinate in the antigenic map.
640

Technical discussion on pLM-based MDS
A recent study [36] reported a model that, like PLANT, predicts the antigenic map coordinates
of virus strains from viral sequences. However, this model adopts a learning approach that
targets the coordinates constructed by conventional cartography methods as prediction
outputs. As shown in **Figure 1C**, conventional cartography fails to accurately reconstruct
the overall trajectory of H3N2 antigenic evolution, revealing a structural limitation of this
approach. In contrast, our study introduces a new learning framework, pLM-based MDS,
which embeds virus strains into a low-dimensional space based on pairwise antigenic
distances, without relying on a preconstructed antigenic map. This method transforms the
semantic space of a pLM into a phenotypic space under supervision using antigenic
distances between sequence pairs. pLM-based MDS can be extended beyond antigenic
mapping to construct diverse phenotypic spaces, offering a new direction for flexible
genotype–phenotype modeling using pLMs.
654

Architecture of PLANT
PLANT comprises two modules: (i) a cartography module that computes antigenic map
coordinates from HA1 sequences and outputs the corresponding cartographic distance,
and (ii) a systematic error module that estimates the component of the observed antigenic
distance not attributable to the HA1 genotype, based on experimental metadata.
The cartography module consists of a pre-trained pLM, ESM-2 (650M parameters)
[12], which takes tokenized HA1 sequences as input, followed by three fully connected layers.
The fully connected layers project the per-sequence embedding (CLS token embedding)
from EMS-2 into a 3-dimensional coordinate. The cartographic distance is computed as the
Euclidean distance between the coordinates of the reference and target strains.
The systematic error module receives four categorical variables as one-hot
encoded inputs: the IDs of the target and reference strains, and the passage types (egg,
cell, and unknown) of the target and reference viruses. These metadata are used to
statistically account for sources of antigenic variation not attributable to the HA1 genotype,
including differences in HA2 or NA, variation in virus stock lots, infectivity, hemagglutination
properties, and host animal effects. The selection of metadata follows a prior work [15].
The module systematic error consists of two subnetworks: (i) a three-layer fully
connected subnetwork that takes the reference strain ID and both passage types as input,
and (ii) a single-layer subnetwork that takes only the target strain ID. The outputs of the two
subnetworks are summed to yield the systematic error. This modular design improves
prediction stability for target strains not included in the training data (i.e., those with missing
IDs).
677

678   **Supervised learning loss**

679 To train PLANT on observed antigenic distances between reference–target strain pairs, we

680 designed a custom loss function that addresses three key challenges: (i) right-censoring of

681 observed values, (ii) imbalance in the distribution of observed antigenic distances, and (iii)

682 the need to minimize not only the discrepancy between predicted antigenic and observed

683 distances but also that between cartographic and observed distances.

684  With respect to (i), in HI assays, antigenic distance measurements are subject to

685 an upper bound, and some observations are right-censored [8]. Accordingly, we used a

686 conditional loss function: for uncensored data, the loss was the squared error between

687 predicted and observed values; for censored data, the squared error was applied only if the

688 prediction fell below the observed value, and set to zero otherwise. This design penalizes

689 underestimation while allowing overestimation for censored observations.

690  With respect to (ii), because large antigenic distances are rare, the distribution of

691 values is heavily imbalanced. To mitigate this, we divided the range 0–8 into nine bins,

692 assigned each bin a weight proportional to the inverse of its frequency (capped at 5), and

693 used these weights to compute a weighted squared error across all training samples.

694  With respect to (iii), in PLANT, the primary objective is not to perfectly predict

695 observed antigenic distances but to construct an antigenic map that faithfully reflects

696 underlying antigenic differences. Therefore, we included an additional regularization term

697 that minimizes the squared error between the cartographic distance and the observed

698 antigenic distance. This term was weighted by 0.05 and added to the total loss.

700 **Unsupervised learning loss**

701 To improve the generalizability of PLANT by leveraging information on the diversity and

702 sequence characteristics of HA1 proteins, we incorporated unlabeled HA1 sequences—

703 those lacking serological assay data—into model training. For this purpose, PLANT

704 introduces two regularization losses: the semantic loss and the local-global loss, which

705 impose semantic and geometric constraints on the structure of the antigenic map space. As

706 unlabeled sequences, we used HA1 sequences from human H3N2 viruses, as well as

707 those from other HA subtypes isolated from both human and animal hosts (**see**

708 **Preparation of dataset for unsupervised learning section**).

709  The semantic loss penalizes discrepancies between semantic distances derived

710 from the frozen ESM-2 (650M) model and the cartographic distances predicted by PLANT.

711 To compute semantic distances, we first obtained per-sequence embeddings by applying

712 both mean and max pooling to the final-layer outputs of ESM-2. For each mini-batch—

713 including reference, target, and unlabeled strains—we then calculated pairwise Euclidean

714 distances between embeddings. These distances were rescaled using a learnable scaling

715 parameter and compared to PLANT-predicted cartographic distances via mean squared

716 error. This loss encourages the preservation of the original semantic structure encoded by

717 ESM-2 and helps mitigate catastrophic forgetting during fine-tuning.

718  The local-global loss is designed to promote both local smoothness and global

719 separation within the antigenic map. For each sample in a mini-batch, the model minimizes

the distance to its three nearest neighbors, encouraging antigenically similar sequences to
be placed close together. Conversely, a penalty is applied to pairs of non-neighboring
samples that lie within a margin of 0.125 (equivalent to 1 antigenic unit), promoting clearer
separation between distinct antigenic clusters.
The training dataset consists of both labeled reference–target pairs (with observed
antigenic distances) and unlabeled HA1 sequences. For labeled data, both supervised and
unsupervised losses were applied, whereas for unlabeled data, only the unsupervised
losses were used. The final training loss is a weighted sum of all loss components, with
weights of 0.2 for the semantic loss and 0.01 for the local-global loss.
729

Training with the full dataset
For model training using the full dataset, we split the data into training, validation, and test
sets at a ratio of 0.8 : 0.1 : 0.1. The split was performed at the *target strain* level to ensure
that all samples derived from a given target strain were assigned exclusively to either the
training, validation, or test set. Stratified sampling was applied based on the collection year
of the target strains to preserve temporal diversity.
The HI dataset included multiple biological replicates—that is, samples with
identical virus strain, virus passage type, reference strain, and reference passage type, but
different experimental dates. To reduce redundancy, we applied a downsampling strategy
that randomly selected one sample from each replicate group during every training epoch.
As noted above, the training dataset included both labeled data (reference–target
pairs with observed antigenic distances) and unlabeled data (individual HA1 sequences).
These were randomly sampled and jointly included in each mini-batch.
Training was conducted using a custom implementation of the Hugging Face
Transformers Trainer, with the AdamW optimizer. The parameters of ESM-2 were not
frozen; full fine-tuning was performed. The learning rate was set to 1.0×10^{-4} and updated
with a linear scheduler with a warm-up ratio of 0.1. A batch size of 16 and a total of 20,000
training steps were used. Weight decay was set to 0.005 for the fully connected layers in
the cartography module and 0.01 for the other components. A dropout rate of 0.05 was
applied to all fully connected layers in the cartography and systematic error modules.
Validation loss was evaluated every 1,000 steps, and the model with the lowest validation
loss was selected as the final model.
Training was conducted using a single NVIDIA H100 GPU (CUDA v12.8) on a
high-performance computing cluster.
754

Construction of a comprehensive antigenic map covering all sequenced H3N2
strains
We constructed a comprehensive antigenic map encompassing all H3N2 virus strains
collected between 1968 and January 2024. Using the PLANT model trained on the full
dataset, we predicted antigenic map coordinates for each virus based on its HA1
sequence. Visualization of the resulting map revealed a strong correlation between

antigenic coordinates and collection dates. However, a small number of strains showed
collection dates that were notably inconsistent with their predicted antigenic positions,
suggesting potential metadata errors.
To detect and exclude such outliers, we implemented an outlier detection procedure
(**Figure S1C**). Specifically, we used a generalized additive model (GAM) implemented in
the R package mgcv v1.9 to predict collection dates from antigenic coordinates. Strains
whose documented collection dates deviated by more than 20 years from the GAM-
predicted values were classified as outliers and excluded from downstream analyses. After
filtering, the final antigenic map included 151,784 virus strains.
770
Definition of antigenic clusters
We defined 20 antigenic clusters using a two-stage Gaussian mixture model (GMM)
clustering approach. In the first stage, virus strains were clustered based on their antigenic
coordinates using the GMM implementation in the R package mclust v6.1.1. The number
and shape of clusters were automatically determined based on the Bayesian Information
Criterion (BIC), with the number of clusters explored in the range of 200 to 1000 (in
increments of 100).
In the second stage, the centroid coordinates of the first-stage clusters were subjected to
another round of GMM clustering, with the number of clusters explored in the range of 20 to
40 (in increments of 5). The resulting clusters from this second stage were defined as
antigenic clusters and used in subsequent analyses.
782
Evaluation of PLANT's predictive performance for future strains
To assess PLANT's ability to predict the antigenicity of novel, future-emerging variants, we
split the dataset based on virus collection dates, constructing a training dataset composed
of past variants and a test dataset comprising future variants.
Given that seasonal influenza vaccine strain selection meetings are typically held
in February and September, we defined influenza seasons as follows: the Southern
Hemisphere (SH) season spans February to August, and the Northern Hemisphere (NH)
season spans September to January of the following year. For each of the 19 seasons from
2014SH to 2023SH, virus strains collected up to the cutoff season were included in the
train-validation dataset, while those collected after the cutoff were assigned to the test
dataset. For instance, when the cutoff is 2014SH, the train-validation dataset excludes any
data with a virus or reference strain collection date on or after September 1, 2014. This
temporal split was applied consistently to both labeled data (i.e., reference–target pairs with
observed antigenic distances) and unlabeled data (i.e., HA1 sequences without serological
measurements).
Within the train-validation dataset, we further divided data into training and
validation subsets at a 0.9 : 0.1 ratio. The split was stratified by the collection year of the
target strains and grouped at the target strain level, following the same procedure as used
in full dataset training.

Training of the AdaBoost model

To provide a comparative benchmark for evaluating PLANT's predictive performance, we

trained the AdaBoost-based model for H3N2 antigenicity prediction proposed by Shah et al.

[15] using the same neutralization assay dataset as PLANT. Like PLANT, the AdaBoost

model takes as input the HA1 protein sequences of the reference and target strains, along

with categorical variables representing experimental conditions, to predict the antigenic

distance under those conditions. However, unlike PLANT, it neither predicts antigenic map

coordinates nor enables the construction of an antigenic map.

Model training involved a two-step hyperparameter optimization process

consisting of an initial broad grid search followed by a refined Bayesian search. Both

searches were conducted using Scikit-learn's GridSearchCV and BayesSearchCV modules

on the 2014SH season dataset. In the grid search, we explored combinations of four

hyperparameters: the number of estimators (n_estimators) was set to 50, 100, or 200; the

learning rate (learning_rate) was set to 0.001, 0.01, 0.1, or 1.0; the maximum tree depth

(estimator__max_depth) was set to 500, 1000, or 2000; and the maximum number of

features considered at each split (estimator__max_features) was set to "auto", "sqrt", or

"log2". The grid search used five-fold cross-validation with negative mean absolute error as

the scoring metric and was parallelized using 14 jobs. The best-performing configuration

from the grid search was: n_estimators = 200, learning_rate = 0.001,

estimator__max_depth = 500, and estimator__max_features = "sqrt".

The Bayesian optimization then fine-tuned the search space based on these

results. It sampled n_estimators as an integer between 150 and 600, learning_rate as a

real number between 0.0005 and 0.005, and estimator__max_depth as an integer between

100 and 750, while fixing estimator__max_features to "sqrt". This second-stage search ran

for 50 iterations with five-fold cross-validation and used the same scoring metric. It was

parallelized with 5 jobs. The best parameters identified were: n_estimators = 361,

learning_rate = 0.0005, estimator__max_depth = 728, and estimator__max_features =

"sqrt".

All computations were performed on Ubuntu Linux 20.04 using Python 3.10.13.

The code used for AdaBoost hyperparameter tuning and model training is available in the

PLANT code repository.

Identifying phylogenetic branches with elevated antigenic change

We analyzed HA1 protein sequences of human H3N2 influenza viruses collected between

1968 and 2024, as described in the Dataset preparation section. To reduce redundancy

while preserving genetic diversity, the sequences were clustered using CD-HIT v4.8.1 [37]

with a 99% identity threshold, yielding 2,748 representative sequences. A maximum-

likelihood phylogenetic tree was then reconstructed using IQ-TREE2 v2.4.0 [38] with the

LG+F+G4 amino acid substitution model. This tree was subsequently rooted and time-

calibrated using TreeTime v0.11.4 [39], which also performed ancestral sequence
reconstruction for all internal nodes under default settings.

Antigenic map coordinates were predicted for both terminal and internal HA1
sequences using the PLANT model trained on the full dataset. For each phylogenetic
branch, we computed the cartographic distance between its parent and child nodes.
Branches with significantly elevated antigenic change were identified as those whose
cartographic distance exceeded the mean across all internal branches by more than three
standard deviations.

To compare antigenic change between internal and external branches while
minimizing sampling bias, we repeated the entire analysis pipeline—from tree construction
to antigenic change inference—using all 24,166 unique HA1 sequences (i.e., without
applying 99% clustering).

To estimate the average rate of antigenic evolution in H3N2, we calculated the
cumulative antigenic change along the phylogenetic trunk, defined as the set of branches
with ≥10 descendant tips, starting from 1997. A linear regression was then performed to
model cumulative cartographic distance of both internal and external nodes as a function of
their sampling or inferred dates.

Clusters and metaclusters were assigned to internal nodes based on their
predicted antigenic map coordinates, using the same GMM described earlier. Monthly
proportions of each metacluster were calculated. U.S. H3N2 influenza case numbers were
obtained from sentinel surveillance data available via FluNet
(https://www.who.int/tools/flunet).

All phylogenetic tree processing was performed using ete3 v3.1.3 [40], and
visualizations were created with ggtree v3.14.0 [41].

Estimating the effects of individual mutations on antigenic change
To identify mutations contributing to the antigenic evolution of H3N2 viruses, we developed
a Bayesian non-negative linear model. Cartographic distance is inherently non-negative
and defined as zero when the HA1 protein sequences of two strains are identical. Based on
this, we assumed that mutations either have no effect or increase this distance. To reflect
this biological constraint, the model imposes non-negativity on all regression coefficients
(i.e., the effect sizes of individual mutations).

In this model, amino acid substitutions acquired on each branch are encoded as
one-hot vectors and used as explanatory variables, while the response variable is the
cartographic distance between the parent and child nodes as predicted by PLANT. The
effect size of each mutation is assigned a prior distribution defined by a non-negative half-t
distribution with zero mean and a scale parameter σ, ensuring that the posterior distribution
remains non-negative. Additionally, because strains with identical HA1 sequences should
have zero antigenic distance, the model does not include an intercept term. This model
does not account for interactions between mutations and assumes, like standard linear
regression, that mutation effects are additive.

883   To evaluate model validity, we compared the estimated mutation effect sizes from

884   the Bayesian non-negative linear model with those obtained from a standard

885   (unconstrained) linear regression (**Figure S3B**). The two estimates were broadly

886   consistent, especially for mutations with larger effects, and the non-negative model yielded

887   coefficients that were all zero or greater, as expected.

888   Parameter estimation was carried out using cmdstan v2.36.0 and cmdstanpy

889   v1.2.5, employing Markov Chain Monte Carlo (MCMC) sampling via the No-U-Turn

890   Sampler (NUTS) implemented in Stan. Four independent MCMC chains with 3,000 steps

891   were run. The warm-up period was set to 1,000 steps. Before using the MCMC outputs in

892   downstream analyses, we verified convergence criteria for all parameters: R-hat ≤ 1.05,

893   and both bulk and tail effective sample sizes ≥ 200.

894   To evaluate antigenic effects at the site level rather than for individual mutations,

895   we calculated the site-wise effect for each HA1 position by aggregating the inferred

896   antigenic effects of all mutations occurring at the same site. Specifically, we summed the

897   posterior mean effect of each mutation at a given position and scaled the total by a factor of

898   1/20, corresponding to the number of possible amino acid types.

899   The HA trimer structure [PDB ID: 4FNK] was visualized using ChimeraX v1.8 [42].

Antigenic sites were annotated based on amino acid positions defined in a previous study

[43], and the computed site-wise antigenic effects were mapped onto the structure for spatial

interpretation.

Integrating PLANT-based antigenic cartography with epidemic modeling

Antigenic groups were defined based on the positions of virus strains in the antigenic map.

To remove redundancy, we first selected a single representative from each set of viruses

sharing an identical HA1 protein sequence. Clustering was then performed using a GMM

implemented in the R package mclust v6.1.1, with the number of clusters automatically

determined according to the BIC. For the fitness advantage analysis (**Figure 4**), we

evaluated cluster numbers ranging from 100 to 500 in increments of 100. In contrast, for

immune-escape variant detection (**Figure 5**) and vaccine strain selection (**Figure 6**), we

used a range of 10 to 50 in increments of 10. These differences reflect variations in the

number of influenza seasons analyzed—18 seasons in **Figure 4** and five seasons in

Figures 5 and 6. The resulting antigenic group classification was applied to all virus

strains, including those with redundant sequences, so that every strain was assigned an

antigenic group for downstream analyses.

917   In the vaccine strain selection analysis (**Figure 6**), we merged antigenic groups

that were excessively close in the antigenic map to achieve robust, rather than overly

sensitive, groupings. For this purpose, we used the Mahalanobis distance, which accounts

for correlations among dimensions and differences in scale, providing a reliable measure of

multivariate distance [44]. Specifically, for each pair of antigenic groups, we calculated the

Mahalanobis distance between their centroids. If this distance was less than or equal to

1.0, the two groups were merged into a single antigenic group. This threshold of 1.0
corresponds to approximately one standard deviation in multivariate space.
925   For all downstream analyses, we retained only antigenic groups containing at
least 20 virus strains. In the vaccine strain selection analysis (**Figure 6**), we further limited
the set to those with a prevalence of at least 0.5% across the entire dataset. The
emergence date of each antigenic group was defined as the 5th percentile of the collection
dates of its constituent strains. Group IDs were assigned in ascending order of these
emergence dates. Importantly, group IDs were dynamically assigned in each analysis and
do not represent fixed or shared labels across analyses.
932   To estimate the epidemic dynamics and associated parameters of each antigenic
933   group, we applied a Bayesian multinomial logistic regression model to weekly detection
934   frequency data. This analysis was performed without country- or region-level stratification,
935   using all available virus strains in aggregate.
936   Parameter estimation was carried out using cmdstan v2.34.1 and cmdstanr
937   v2.34.1, employing MCMC sampling via NUTS implemented in Stan. Four independent
938   MCMC chains were run for each analysis: 6,000 steps for **Figure 4** and 2,000 steps for
939   **Figures 5 and 6**. The warm-up period was set to 500 steps in all cases. Before using the
940   MCMC outputs in downstream analyses, we verified convergence criteria for all
941   parameters: R-hat ≤ 1.05, and both bulk and tail effective sample sizes ≥ 200.
942   The relative R_e of the antigenic group g, r_g, was calculated according to the slope
943   parameter β_g of the antigenic group as $r_g = \exp(\gamma \beta_g)$, where γ is the average viral
944   generation time (3.0 days) [45]. In the vaccine strain selection analysis (**Figure 6**), epidemic
945   trajectories for each antigenic group were projected up to two seasons ahead.
946
Definition of fitness advantage and its association with antigenic novelty
To estimate the fitness advantage of each antigenic group, we first classified virus strains
from 2010SH to 2019NH into 134 antigenic groups using the method described above and
estimated their emergence dates and relative R_e. To capture the temporal trend in relative
R_e, we fitted an exponential regression model to the relationship between the emergence
date and relative R_e, thereby estimating the annual rate of increase in relative R_e. The
exponential regression was implemented as a linear model in which the natural logarithm of
relative R_e served as the response variable and the emergence date as the explanatory
variable. The residuals from this regression model quantify how much a group's relative R_e
deviates from the expected value based on its emergence date. Therefore, these residuals
reflect the extent to which a given antigenic group exhibits a higher (or lower) fitness
relative to contemporaneously emerging antigenic groups. This means that this residual
can be interpreted as a proxy for fitness advantage.
960   As a proxy of antigenic novelty, the median cartographic distance from prior
epidemic strains collected within the preceding 1.5 years was quantified. This time window
was selected to capture strains from the most recent epidemic season in the same
hemisphere.

To quantitatively assess the relationship between fitness advantage and
cartographic distance, we constructed a hierarchical Bayesian linear model. In this model,
we introduced season-specific random effects for both the intercept and the slope of
cartographic distance, allowing us to statistically model seasonal variation. Specifically, the
slope for each season was assumed to follow a normal distribution with a mean of zero and
a standard deviation of σ, thereby enabling the estimation of inter-seasonal variability in a
unified framework. To evaluate the explanatory power of the model, we computed the
Bayesian R-squared, as defined by Gelman et al [46].
972
Framework for automated identification of immune-escape variants
Following the procedure described above, we classified virus strains that emerged during
the 2022SH and 2024SH seasons into 30 antigenic groups based on their coordinates in
the antigenic map. For each group, we estimated the relative R_e. To identify mutations
potentially associated with immune escape, we used Nextclade to detect amino acid
substitutions in each strain relative to the reference strain A/Darwin/6/2021 instead of
A/Wisconsin/67/2005-egg.
To conduct a retrospective analysis based on virus genome surveillance data up
to the 2024NH season, we downloaded HA genome sequences and associated metadata
for viruses collected during this period from GISAID on June 30, 2025.
983
Evaluation of historical vaccine strains and development of an automated framework
for vaccine strain selection
Information on historical vaccine strains was obtained from GISAID
(https://gisaid.org/resources/human-influenza-vaccine-composition/) for strains
recommended from the 2010NH season onward, and from the WHO website
(https://www.who.int/teams/global-influenza-programme/vaccines/who-
recommendations/recommendations-for-influenza-vaccine-composition-archive) for strains
from 2001SH to 2010SH. This study focused on WHO-recommended strains for egg-based
inactivated influenza vaccines, which represent the most widely used modality for seasonal
influenza vaccination. These WHO-recommended strains are annotated in **Table S1**.
For the labeled data used to train PLANT (i.e., observed antigenic distances
derived from HI assays), we partitioned the data based on the date the HI assay was
performed, rather than the collection date of the virus. For example, to train the model for
predicting the vaccine strain for the 2014NH season, we included only HI assay data
generated before August 31, 2013. In contrast, unlabeled HA1 sequence data were split
based on the collection date of the virus.
For each cutoff date, an antigenic map was constructed using a PLANT model
trained exclusively on data available up to that time point. As input, we used HA1 sequence
data from viruses collected within the five preceding seasons.
Vaccine strain selection was conducted using the epidemic projection framework
described earlier. Specifically, we projected the prevalence of each antigenic group up to

two seasons into the future from the cutoff date and estimated the posterior mean and 50%
credible interval of group prevalence during the vaccination period.
Each virus strain was then assigned a weight according to the projected
prevalence (posterior mean) of its corresponding antigenic group. Using these weights, we
calculated the weighted median centroid in the antigenic map and selected the virus strain
closest to this centroid as the candidate vaccine strain.
To evaluate the antigenic match between the candidate vaccine strain and the
circulating viruses during the vaccination period, we used the cartographic distances output
by a PLANT model trained on the full dataset.
Additionally, we defined a "hindsight-optimal strain" as the virus strain that existed
prior to the cutoff date and exhibited the smallest median cartographic distance to the
circulating strains during the vaccination period. This allowed us to benchmark our
automated selection method against the theoretical optimal choice under hindsight
conditions.
1019
1020   **Code availability**
1021   The computational code used in this study is available at the GitHub repository
1022   (https://github.com/TheSatoLab/PLANT).
1023
Data availability
HI assay data and associated experimental metadata used in this study are available in the
annual and interim reports of the Worldwide Influenza Centre
(https://www.crick.ac.uk/research/platforms-and-facilities/worldwide-influenza-
centre/annual-and-interim-reports). Genomic sequences of H3N2 viruses analyzed in this
study are available from the GISAID database (https://gisaid.org/). The GISAID
acknowledgment table is provided in the GitHub repository
(https://github.com/TheSatoLab/PLANT).
1032

Acknowledgments
We gratefully acknowledge all data contributors, i.e. the Authors and their Originating
laboratories responsible for obtaining the specimens, and their Submitting laboratories for
generating the genetic sequence and metadata and sharing via the GISAID Initiative, on
which this research is based. This study was carried out using the TSUBAME4.0
supercomputer at Institute of Science Tokyo.
We gratefully acknowledge the valuable discussions and constructive feedback
provided by Dr. Junna Kawasaki (Chiba University, Japan), Mr. Takuya Hemmi (Kyoto
University, Japan), Mr. Keno Strotjohann (Swiss Tropical and Public Health Institute,
Switzerland), Mr. Kent Mori (Kyushu University, Japan), Mr. Moonjong Kang (The
University of Tokyo, Japan), Dr. June Sese (Humanome Lab), Dr. Keietsu Abe (Tohoku
University), and Dr. Ryuichi Koga (National Institute of Advanced Industrial Science and
Technology).

This study was supported in part by JST PRESTO (JPMJPR22R1, to Jumpei Ito);
JSPS KAKENHI Grant-in-Aid for Scientific Research B (JP25K00116, to Jumpei Ito); AMED
SCARDA Japan Initiative for World-leading Vaccine Research and Development Centers
"UTOPIA" (JP223fa627001, to Jumpei Ito, Kei Sato, Shusuke Kawakubo, Spyros Lytras);
SHIONOGI Infectious Disease Research Promotion Foundation (to Jumpei Ito); JSPS
KAKENHI Fund for the Promotion of Joint International Research (International Leading
Research) (JP23K20041, to Kei Sato); AMED ASPIRE Program (25jf0126002, to Kei Sato);
AMED SCARDA Program on R&D of new generation vaccine including new modality
application (253fa727002, to Kei Sato); AMED Research Program on Emerging and Re-
emerging Infectious Diseases (24fk0108907, 25fk0108690, to Kei Sato); AMED Japan
Program for Infectious Diseases Research and Infrastructure (Collaborative Research via
Overseas Research Centers) (25wm0225041, to Kei Sato); JSPS KAKENHI Grant-in-Aid for
Scientific Research A (JP24H00607, to Kei Sato); the Platform Project for Supporting Drug
Discovery and Life Science Research (Basis for Supporting Innovative Drug Discovery and
Life Science Research (BINDS)) from AMED (JP24ama121012, supporting numbers
S02820001 and S02820002, to Kei Sato), AIST KAKUSEI project (FY2024 and FY2025 to
Shusuke Kawakubo).

1063

Declaration of interest

Jumpei Ito has consulting fees and honoraria for lectures from Takeda Pharmaceutical Co.
Ltd and AstraZeneca. Kei Sato has consulting fees from Moderna Japan Co., Ltd. and
Takeda Pharmaceutical Co. Ltd., and honoraria for lectures from Moderna Japan Co., Ltd.,
Shionogi & Co., Ltd and AstraZeneca. The other authors declare no competing interests.
Conflicts that the editors consider relevant to the content of the manuscript have been
disclosed.

1071

Supplemental Tables

Table S1. Information on virus strains used in this study.

The coordinates (X, Y, Z) represent the antigenic map positions predicted by PLANT
trained on the full dataset. Clade and Subclade were assigned using Nextclade. The
columns "is WHO vaccine strain", "Closest vaccine strain", "Distance from closest vaccine
strain", "Season matched vaccine strain", and "Distance from season matched vaccine
strain" are assigned only to virus strains collected from 2001SH onward.

Table S2 Summary of predicted antigenic effects for individual mutations.

The positions correspond to amino acid residues in the HA1 protein. The column
'isSignificant' indicates statistical significance, with 1 representing true and 0 representing
false.

**Table S3 Summary of predicted antigenic effects aggregated by position across
mutation types.**

1087   The predicted antigenic effect of each mutation type was aggregated at each site using the
1088   sum values. The column 'Site-wise effect' represents the sum of effects divided by 20,
1089   corresponding to the total number of possible amino acid types at each site.
1090
1091
1092

Author Contributions

Project design and management: Jumpei Ito

Development of PLANT: Jumpei Ito

Improvement of PLANT: Jumpei Ito, Adam Strange, and Shusuke Kawakubo

AdaBoost-based analysis: Adam Strange

PLANT-integrated phylogenetic analyses: Shusuke Kawakubo

Development of a Bayesian non-negative linear model: Hiroaki Unno

Immune escape variant identification analysis: Jumpei Ito and Hiroaki Unno

Other analyses: Jumpei Ito

Dataset preparation: Shusuke Kawakubo, Hiroaki Unno, Adam Strange, Spyros Lytras,

Kaho Okumura, and Jumpei Ito

Serological data provision: Alice Lilley, Ruth Harvey, and Nicola Lewis

Initial draft writing: Jumpei Ito and Shusuke Kawakubo

Improving the initial draft: Jumpei Ito, Spyros Lytras, and Kei Sato

Figure preparation: Jumpei Ito, Shusuke Kawakubo, and Kaho Okumura

Scientific illustration: Jumpei Ito, Kaho Okumura

Naming of PLANT and logo design: Shusuke Kawakubo

Project administration: Kei Sato

All authors reviewed and approved the final manuscript.

1112

1113

| 1114 | | **Reference** |

1. Petrova, V.N., and Russell, C.A. (2018). The evolution of seasonal influenza viruses. Nat Rev Microbiol *16*, 47-60. 10.1038/nrmicro.2017.118.

2. Neher, R.A., Bedford, T., Daniels, R.S., Russell, C.A., and Shraiman, B.I. (2016). Prediction, dynamics, and visualization of antigenic phenotypes of seasonal influenza viruses. Proc Natl Acad Sci U S A *113*, E1701-1709. 10.1073/pnas.1525578113.

3. Bedford, T., Suchard, M.A., Lemey, P., Dudas, G., Gregory, V., Hay, A.J., McCauley, J.W., Russell, C.A., Smith, D.J., and Rambaut, A. (2014). Integrating influenza antigenic dynamics with molecular evolution. Elife *3*, e01914. 10.7554/eLife.01914.

4. Perofsky, A.C., Huddleston, J., Hansen, C.L., Barnes, J.R., Rowe, T., Xu, X., Kondor, R., Wentworth, D.E., Lewis, N., Whittaker, L., et al. (2024). Antigenic drift and subtype interference shape A(H3N2) epidemic dynamics in the United States. Elife *13*. 10.7554/eLife.91849.

5. Chen, J.R., Liu, Y.M., Tseng, Y.C., and Ma, C. (2020). Better influenza vaccines: an industry perspective. J Biomed Sci *27*, 33. 10.1186/s12929-020-0626-6.

6. Gerdil, C. (2003). The annual production cycle for influenza vaccine. Vaccine *21*, 1776-1779. 10.1016/s0264-410x(03)00071-9.

7. Hay, A.J., and McCauley, J.W. (2018). The WHO global influenza surveillance and response system (GISRS)-A future perspective. Influenza Other Respir Viruses *12*, 551-557. 10.1111/irv.12565.

8. Smith, D.J., Lapedes, A.S., de Jong, J.C., Bestebroer, T.M., Rimmelzwaan, G.F., Osterhaus, A.D., and Fouchier, R.A. (2004). Mapping the antigenic and genetic evolution of influenza virus. Science *305*, 371-376. 10.1126/science.1097211.

9. Fouchier, R.A., and Smith, D.J. (2010). Use of antigenic cartography in vaccine seed strain selection. Avian Dis *54*, 220-223. 10.1637/8740-032509-ResNote.1.

10. Bepler, T., and Berger, B. (2021). Learning the protein language: Evolution, structure, and function. Cell Syst *12*, 654-669.e653. 10.1016/j.cels.2021.05.017.

11. Hie, B., Zhong, E.D., Berger, B., and Bryson, B. (2021). Learning the language of viral evolution and escape. Science *371*, 284-288. 10.1126/science.abd7331.

12. Lin, Z., Akin, H., Rao, R., Hie, B., Zhu, Z., Lu, W., Smetanin, N., Verkuil, R., Kabeli, O., Shmueli, Y., et al. (2023). Evolutionary-scale prediction of atomic-level protein structure with a language model. Science *379*, 1123-1130. 10.1126/science.ade2574.

13. Cai, Z., Zhang, T., and Wan, X.-F. (2010). A Computational Framework for Influenza Antigenic Cartography. PLOS Computational Biology *6*, e1000949. 10.1371/journal.pcbi.1000949.

14. Han, L., Li, L., Wen, F., Zhong, L., Zhang, T., and Wan, X.F. (2019). Graph-guided multi-task sparse learning model: a method for identifying antigenic variants of influenza A(H3N2) virus. Bioinformatics *35*, 77-87. 10.1093/bioinformatics/bty457.

15. Shah, S.A.W., Palomar, D.P., Barr, I., Poon, L.L.M., Quadeer, A.A., and McKay, M.R.
(2024). Seasonal antigenic prediction of influenza A H3N2 using machine learning.
Nat Commun *15*, 3833. 10.1038/s41467-024-47862-9.
16. Harvey, W.T., Davies, V., Daniels, R.S., Whittaker, L., Gregory, V., Hay, A.J.,
Husmeier, D., McCauley, J.W., and Reeve, R. (2023). A Bayesian approach to
incorporate structural data into the mapping of genotype to antigenic phenotype of
influenza A(H3N2) viruses. PLoS Comput Biol *19*, e1010885.
10.1371/journal.pcbi.1010885.
17. Aksamentov, I., Roemer, C., Hodcroft, E.B., and Neher, R.A. (2021). Nextclade:
clade assignment, mutation calling and quality control for viral genomes. Journal of
Open Source Software *6*, 3773. 10.21105/joss.03773.
18. Luksza, M., and Lässig, M. (2014). A predictive fitness model for influenza. Nature
507, 57-61. 10.1038/nature13087.
19. Chen, Z., Tsui, J.L., Gutierrez, B., Busch Moreno, S., du Plessis, L., Deng, X., Cai, J.,
Bajaj, S., Suchard, M.A., Pybus, O.G., et al. (2024). COVID-19 pandemic
interventions reshaped the global dispersal of seasonal influenza viruses. Science
386, eadq3003. 10.1126/science.adq3003.
20. Koel, B.F., Burke, D.F., Bestebroer, T.M., van der Vliet, S., Zondag, G.C., Vervaet,
G., Skepner, E., Lewis, N.S., Spronken, M.I., Russell, C.A., et al. (2013).
Substitutions near the receptor binding site determine major antigenic change during
influenza virus evolution. Science *342*, 976-979. 10.1126/science.1244730.
21. Shih, A.C., Hsiao, T.C., Ho, M.S., and Li, W.H. (2007). Simultaneous amino acid
substitutions at antigenic sites drive influenza A hemagglutinin evolution. Proc Natl
Acad Sci U S A *104*, 6283-6288. 10.1073/pnas.0701396104.
22. Abousamra, E., Figgins, M., and Bedford, T. (2024). Fitness models provide accurate
short-term forecasts of SARS-CoV-2 variant frequency. PLoS Comput Biol *20*,
e1012443. 10.1371/journal.pcbi.1012443.
23. Castro, L.A., Bedford, T., and Ancel Meyers, L. (2020). Early prediction of antigenic
transitions for influenza A/H3N2. PLoS Comput Biol *16*, e1007683.
10.1371/journal.pcbi.1007683.
24. Suzuki, R., Yamasoba, D., Kimura, I., Wang, L., Kishimoto, M., Ito, J., Morioka, Y.,
Nao, N., Nasser, H., Uriu, K., et al. (2022). Attenuated fusogenicity and pathogenicity
of SARS-CoV-2 Omicron variant. Nature *603*, 700-705. 10.1038/s41586-022-04462-
1.
25. Obermeyer, F., Jankowiak, M., Barkas, N., Schaffner, S.F., Pyle, J.D., Yurkovetskiy,
L., Bosso, M., Park, D.J., Babadi, M., MacInnis, B.L., et al. (2022). Analysis of 6.4
million SARS-CoV-2 genomes identifies mutations associated with fitness. Science
376, 1327-1332. 10.1126/science.abm1208.
26. Ito, J., Strange, A., Liu, W., Joas, G., Lytras, S., and Sato, K. (2025). A protein
language model for exploring viral fitness landscapes. Nat Commun *16*, 4236.
10.1038/s41467-025-59422-w.

27.   Kikawa, C., Loes, A.N., Huddleston, J., Figgins, M.D., Steinberg, P., Griffiths, T.,
Drapeau, E.M., Peck, H., Barr, I.G., Englund, J.A., et al. (2025). High-throughput
neutralization measurements correlate strongly with evolutionary success of human
influenza strains. bioRxiv. 10.1101/2025.03.04.641544.
28.   Yao, Y., Li, X., Liao, B., Huang, L., He, P., Wang, F., Yang, J., Sun, H., Zhao, Y., and
Yang, J. (2017). Predicting influenza antigenicity from Hemagglutintin sequence data
based on a joint random forest method. Sci Rep *7*, 1545. 10.1038/s41598-017-
01699-z.
29.   Li, X., Li, Y., Shang, X., and Kong, H. (2024). A sequence-based machine learning
model for predicting antigenic distance for H3N2 influenza virus. Front Microbiol *15*,
1345794. 10.3389/fmicb.2024.1345794.
30.   Cox, N.J., and Bender, C.A. (1995). The molecular epidemiology of influenza viruses.
Seminars in Virology *6*, 359-370. https://doi.org/10.1016/S1044-5773(05)80013-7.
31.   Plotkin, J.B., Dushoff, J., and Levin, S.A. (2002). Hemagglutinin sequence clusters
and the antigenic evolution of influenza A virus. Proceedings of the National
Academy of Sciences *99*, 6263-6268. 10.1073/pnas.082110799.
32.   Strelkowa, N., and Lässig, M. (2012). Clonal Interference in the Evolution of
Influenza. Genetics *192*, 671-682. 10.1534/genetics.112.143396.
33.   Chen, Z., Tsui, J.L., Cai, J., Su, S., Viboud, C., du Plessis, L., Lemey, P., Kraemer,
M.U.G., and Yu, H. (2025). Disruption of seasonal influenza circulation and evolution
during the 2009 H1N1 and COVID-19 pandemics in Southeastern Asia. Nat Commun
16, 475. 10.1038/s41467-025-55840-y.
34.   Koelle, K., and Rasmussen, D.A. (2015). The effects of a deleterious mutation load
on patterns of influenza A/H3N2's antigenic evolution in humans. Elife *4*, e07361.
10.7554/eLife.07361.
35.   Steinegger, M., and Söding, J. (2017). MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. Nat Biotechnol *35*, 1026-1028.
10.1038/nbt.3988.
36.   Durazzi, F., Koopmans, M.P.G., Fouchier, R.A.M., and Remondini, D. (2025).
Language models learn to represent antigenic properties of human influenza A(H3)
virus. Scientific Reports *15*, 21364. 10.1038/s41598-025-03275-2.
37.   Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: accelerated for clustering
the next-generation sequencing data. Bioinformatics *28*, 3150-3152.
10.1093/bioinformatics/bts565.
38.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von
Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods
for Phylogenetic Inference in the Genomic Era. Mol Biol Evol *37*, 1530-1534.
10.1093/molbev/msaa015.
39.   Sagulenko, P., Puller, V., and Neher, R.A. (2018). TreeTime: Maximum-likelihood
phylodynamic analysis. Virus Evol *4*, vex042. 10.1093/ve/vex042.

40.   Huerta-Cepas, J., Serra, F., and Bork, P. (2016). ETE 3: Reconstruction, Analysis,
and Visualization of Phylogenomic Data. Mol Biol Evol *33*, 1635-1638.
10.1093/molbev/msw046.
41.   Yu, G., Lam, T.T., Zhu, H., and Guan, Y. (2018). Two Methods for Mapping and
Visualizing Associated Data on Phylogeny Using Ggtree. Mol Biol Evol *35*, 3041-
3043. 10.1093/molbev/msy194.
42.   Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H.,
and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure building and analysis.
Protein Sci *32*, e4792. 10.1002/pro.4792.
43.   Broecker, F., Liu, S.T.H., Sun, W., Krammer, F., Simon, V., and Palese, P. (2018).
Immunodominance of Antigenic Site B in the Hemagglutinin of the Current H3N2
Influenza Virus in Humans and Mice. J Virol *92*. 10.1128/jvi.01100-18.
44.   Etherington, T.R. (2019). Mahalanobis distances and ecological niche modelling:
correcting a chi-squared probability error. PeerJ *7*, e6678. 10.7717/peerj.6678.
45.   Carrat, F., Vergu, E., Ferguson, N.M., Lemaitre, M., Cauchemez, S., Leach, S., and
Valleron, A.J. (2008). Time lines of infection and disease in human influenza: a
review of volunteer challenge studies. Am J Epidemiol *167*, 775-785.
10.1093/aje/kwm375.
46.   Gelman, A., Goodrich, B., Gabry, J., and Vehtari, A. (2019). R-squared for Bayesian
Regression Models. The American Statistician *73*, 307-309.
47.   Ekiert, D.C., Kashyap, A.K., Steel, J., Rubrum, A., Bhabha, G., Khayat, R., Lee, J.H.,
Dillon, M.A., O'Neil, R.E., Faynboym, A.M., et al. (2012). Cross-neutralization of
influenza A viruses mediated by a single antibody loop. Nature *489*, 526-532.
10.1038/nature11414.
1259

 Figure Legend

Figure 1 Construction of the antigenic map by PLANT

A) Overview of PLANT training. Each virus strain is embedded in a low-dimensional

semantic space derived from a pLM using its HA1 sequence. The model is trained so that

the distances between target and reference strains in this space, referred to as semantic

distances, reflect experimentally measured antigenic distances. This training transforms the

semantic space into an antigenic map. In the bottom left, the structure of the HI assay

dataset is shown. Each point represents a virus pair tested in an HI assay, colored by the

observed antigenic distance. Gray indicates pairs without available measurements. Target

and reference strains are ordered by collection date.

B) Model architecture of PLANT. The model takes HA1 sequences of target and reference

strains, along with experimental conditions, as input. It outputs coordinates in the antigenic

map, the cartographic distance (Euclidean distance on the map), and antigenic distance

adjusted for estimated systematic error.

C) Antigenic map reconstructed using a conventional method, Racmacs

(https://acorg.github.io/Racmacs/).

D) Comprehensive antigenic map covering 151,784 H3N2 strains with HA1 sequences

collected from 1968 to January 2024. Points are colored by collection date (left) or by 20

antigenic clusters (i.e., c1–20; right).

E) Cartographic distance matrix. For each antigenic cluster, 10% of strains (or at least 20 if

fewer) were randomly sampled. Strains are ordered by antigenic cluster and collection

date. Antigenic metaclusters defined in (**F**) are labeled.

F) Hierarchical clustering of antigenic clusters. Based on centroid coordinates, clusters

were grouped into four metaclusters (i.e., m1–4) using Ward's method with a defined cutoff

(dotted line). Antigenic cluster c1, which consists primarily of pre-2000 strains, was

excluded. Nextclade subclade classifications [17] corresponding to each metacluster are also

indicated.

G) Distributions of cartographic distances from seasonal centroids (top) and seasonal

detection frequencies of antigenic clusters (bottom). Dotted lines indicate the 2-antigenic

unit criterion (top) and metacluster boundaries (bottom). SH: Southern Hemisphere season;

NH: Northern Hemisphere season.

1291

Figure 2 Evaluation of the extrapolation performance of PLANT

A) Dataset splitting based on virus collection date. For each cutoff (e.g., end of 2014SH),

virus strains were split into past and future subsets. PLANT was trained on past data and

evaluated on future data. Nineteen such datasets were prepared using cutoffs from

2014SH to 2023SH.

B) Summary of prediction performance across all cutoff points. For each later season, MAE

(top) and Pearson correlation (bottom) are shown for PLANT and an AdaBoost model [15].

Results from PLANT trained on the full dataset are overlaid for comparison. Only seasons

with HI assay data for ≥20 target strains are included.

C) Average prediction performance stratified by the number of seasons since the cutoff.

Error bars indicate standard deviations.

1303

Figure 3 Rate of antigenic evolution and genetic drivers of antigenic change in H3N2

A) Workflow for estimating antigenic changes along phylogenetic branches. Following

ancestral sequence reconstruction, PLANT (trained on the full dataset) was used to predict

antigenic map coordinates for all tip and internal node sequences. Cartographic distances

between parent and child nodes were then calculated for each branch.

B) Time-scaled maximum-likelihood phylogenetic tree of H3N2 HA1, with branches colored

according to the magnitude of antigenic change. A total of 2,748 HA1 sequences,

downsampled via 99% sequence identity clustering, were included. Antigenic clusters and

metaclusters are annotated at the tip nodes. Internal nodes exhibiting significant antigenic

changes are labeled with their corresponding acquired substitutions.

C) Antigenic evolution along the phylogenetic trunk, defined as the set of internal branches

with ≥10 descendant tips. The y-axis indicates the cumulative cartographic distance along

each node from the root. Antigenic cluster c1 was excluded. Branch color indicates

antigenic metaclusters. Internal nodes associated with major antigenic transitions are

annotated with the acquired substitutions. The periods corresponding to the 2009 H1N1

pandemic, the 2013–2014 H1N1 epidemic, and the COVID-19 pandemic are highlighted.

Evolutionary rates of antigenic change for the indicated periods, estimated by linear

regression, are shown.

D) Epidemic dynamics of antigenic metaclusters. The proportion of each metacluster in

viral genome surveillance is shown for each monthly time bin.

E) H3N2 epidemic curve (monthly reported cases) based on FluNet data from the US

(https://www.who.int/tools/flunet).

F) Estimated antigenic effects of individual mutations using a Bayesian non-negative linear

model. The lower panel shows the posterior means and 94% highest density intervals

(HDIs). Mutations whose 94% HDI lower bounds exceed 0.05 are considered to have

statistically significant effects and are shown in black. Colored regions represent antigenic

sites A–E [43]. The P values for each antigenic site were calculated by comparing the effect

sizes of mutations within the site to those outside the site using a two-sided Mann–Whitney

U-test, followed by multiple testing correction with the Benjamini–Hochberg method. The

upper panel shows the rolling sum of estimated effects across sequence positions (window

size = 5, step size = 3).

G) HA trimer structure [PDB ID: 4FNK] [47], visualized by known antigenic sites (left) and by

site-wise effect sizes. Site-wise effects were calculated as the sum of mutational effects at

each site, scaled by a factor of 1/20.

H) Ranking of amino acid positions based on site-wise effects.

1339

Figure 4 Relationship between antigenic novelty and fitness advantage

A) Antigenic novelty and fitness advantage. Left) Antigenic novelty, using cartographic

distance from recently circulating strains as a proxy. Right) Fitness advantage, using
relative fitness compared to co-circulating variants as a proxy.
B) Definition of antigenic groups by GMM clustering on the antigenic map. Virus strains
from 2010SH to 2019NH were classified into 134 antigenic groups (g1–134).
C) Epidemic dynamics of antigenic groups estimated using a Bayesian multinomial logistic
model. Lines indicate posterior means.
D) Emergence date and relative effective reproduction number (R_e) of each antigenic
group. Emergence date is defined as the 5th percentile of strain collection dates. Relative
R_e values—normalized to the dominant 2010NH strain—are shown as posterior means.
Point size reflects peak prevalence. The fitted exponential regression curve and the
estimated average annual increase in relative R_e are also indicated.
E) Emergence date and fitness advantage of each antigenic group. Fitness advantage was
defined as the residual from the exponential regression in (**D**) and expressed as a
percentage.
F) Association between antigenic novelty and fitness advantage. The x-axis represents
antigenic novelty, defined as the median cartographic distance from strains that circulated
within the previous 1.5 years.
G) Effect size of cartographic distance on fitness advantage estimated via Bayesian
hierarchical modeling. Posterior means and 90% credible intervals are shown for each
season and overall (highlighted in light green). Gray indicates seasons when the lower 5%
of the posterior is below zero.
H) Bayesian R^2 values [46] for the model. Posterior means and 95% credible intervals are
shown. As a control, the R^2 value for a model that does not incorporate cartographic
distance effects is also included.
1366	
Figure 5 Automated identification of upcoming immune escape variants with PLANT
A) Antigenic group classification. PLANT trained on data through 2023NH was applied to
viruses collected between 2022SH and 2024SH (note: 2024SH was not included in
training). GMM clustering was used to define 30 antigenic groups (i.e., g1–30).
B) Antigenic map colored by estimated relative R_e of each antigenic group.
C) Visualization of antigenic groups of interest.
D) Relative R_e values of antigenic groups. R_e is scaled to 1 for g6 (dominant in 2022SH).
Posterior means and 95% credible intervals are shown.
E) Amino acid substitution patterns for the top six antigenic groups ranked by estimated
fitness. Amino acid substitutions are relative to A/Darwin/6/2021. Only substitutions with
differing presence or absence patterns across the antigenic groups are displayed.
F) Cartographic distance matrix among centroids of clusters in the J.2 subclade.
G) Antigenic map colored by amino acid residues at position 160. Only J.2 subclade strains
are shown.
H) Observed epidemic dynamics of J.2 descendants carrying specific mutations, based on
viral genome surveillance data up to 2024NH. Proportions were calculated in monthly bins.

The period during which data were collected for PLANT training and fitness estimation is annotated.

Figure 6 Evaluation of historical vaccine strains and vaccine strain recommendation with PLANT

A) Cartographic distance from each virus strain to the nearest historical vaccine strain (left) and to the season-matched vaccine strain (right). Analysis includes viruses from 2001SH onward.

B) Histogram of cartographic distances to the nearest historical vaccine strain. The proportion within 2 antigenic units is indicated.

C) Vaccine strain recommendation framework. 1) Construct an antigenic map for the last five seasons and assign viruses to antigenic groups. 2) Predict the prevalence of antigenic groups during the vaccination period using a multinomial logistic model. 3) Compute a weighted centroid and select the nearest strain as the vaccine strain. The predicted prevalence of antigenic groups during the vaccination period was used as the weight.

D) Comparison of selected vaccine strains based on antigenic match, assessed by the median cartographic distance to strains circulating during the vaccination period. This panel compares (i) WHO-recommended strains, (ii) strains nearest to the centroid of the most recent season, (iii) strains selected by our proposed framework (weighted centroid), and (iv) hindsight-optimal strains. The orange line and accompanying numbers indicate the overall median across all seasons. The difference between WHO-selected strains and those chosen by our framework is highlighted in red. The dotted line denotes the 2-antigenic unit threshold.

Supplemental Figures

Figure S1. Description of datasets (related to Figure 1)

A) Structure of the HI assay dataset. Each point represents a virus pair tested in an HI assay, colored by observed antigenic distance. Gray color indicates virus pairs without observed antigenic distance. Target and reference strains are ordered by collection date.

B) Proportions of Nextclade subclades within each antigenic cluster. Note that subclade assignments were unavailable (i.e., NA) for most strains in clusters c1–8. Also, corresponding antigenic metaclusters are indicated.

C) Detection of virus strains with outlier collection dates. Collection dates were predicted using a generalized additive model (GAM) based on the antigenic map coordinates of each strain. Strains exhibiting a discrepancy of more than 20 years between their documented and predicted collection dates were removed as outliers. The left and right panels show the data before and after filtering, respectively.

Figure S2. Prediction performance of PLANT (related to Figure 2)

A) Interpolation performance of PLANT. Prediction accuracy was evaluated using random splits along the virus strain axis. Mean absolute error (MAE) and Pearson correlation are shown for antigenic distance, cartographic distance, and scaled semantic distance from the un-finetuned ESM-2, across individual seasons. Median values across seasons are shown on the right side of the plots. Only seasons with HI assay data for ≥20 target strains are included.

B and C) Evaluation of the extrapolation performance of PLANT in each epidemic season. Unlike **Figure 2B**, results are shown separately for 19 datasets with distinct cutoffs. For each later season, MAE (**B**) and Pearson correlation (**C**) are shown for PLANT, an AdaBoost model [15], and PLANT trained on the full dataset.

Figure S3. Rate of antigenic evolution and genetic drivers of antigenic change in H3N2 (related to Figure 3)

A) Comparison of cartographic distances between external and internal branches in the phylogenetic tree. For minimizing sampling bias, we reconstructed the phylogenetic tree of H3N2 using all 24,166 unique HA1 sequences (i.e., without applying 99% clustering).

B) Comparison of estimated substitution effects between the Bayesian non-negative linear model and the unconstrained linear model. Posterior means (dots) and 94% HDIs (lines) are shown.

C) Ranking plot of estimated substitution effects. Top 40 mutations are annotated with text labels. Mutations that are located in antigenic sites are labelled with color according to the inset color key. Significant and non-significant mutations are shown as pink and gray, respectively.

Figure S4. Vaccine strain recommendation with PLANT (related to Figure 6)

A) Comparison of vaccine strains selected by four different methods. Violin plots show the

distribution of cartographic distances to epidemic strains during the vaccination period, with

dots indicating median values.

B) Projected epidemic dynamics of antigenic groups. Results are shown for the 2022NH

vaccination season. Posterior mean (line) and 50% credible interval (ribbon) are indicated.

The plot includes the top five antigenic groups based on forecasted epidemic frequency

during the vaccination period. For each antigenic group, the most frequent antigenic cluster

and Nextclade subclade among its constituent strains are displayed alongside the group

name.

1456

A

H3N2's HA1 sequences

MVCLFLVPLV**TT**
MVCLF**Q**VPLVSS
MVCLF**Q**V**R**LVSS
MVCLF**R**VP**R**VS**T**

pLM →

Semantic space

Semantic distance

pLM-based MDS ▶▶▶

Antigenic map

Antigenic distance

■ Reference
● Target

Systematic HI assay

Antigenic distance

Gray: NA

Reference

Target

Ferret infection

Serum →

HI assay

Reference strain

Target strain

B

HA1 protein sequences

Target   Reference

Tokenizing

Cartography module

ESM-2 (650M)

Embedding

Linear

Coordinates*
(x, y, z)   (x, y, z)

Cartographic distance*

Experimental conditions
- Target ID
- Reference ID
- Target passage
- Reference passage

One-hot encoding

Systematic error module

Linear

Systematic error

Antigenic distance*   *Output

C

Racmacs map

1968   2024
Collection date

D

PLANT map

Antigenic clusters

c1
c2
c3
c4
c5
c6
c7
c8
c9
c10
c11
c12
c13
c14
c15
c16
c17
c18
c19
c20

1968   2024
Collection date

Recent →

E

Strains

Strains

c1
c2
c3 ─┐
c4   │ m1
c5 ─┘
c6 ─┐
c7   │ m2
c8   │
c9 ─┘
c10 ─┐
c11   │
c12   │
c13   │ m3
c14   │
c15   │
c16   │
c17 ─┘
c18 ─┐
c19   │ m4
c20 ─┘

0   4
Cartographic distance

F

c2 ─┐
c3   │ m1 (c2–5)
c4   │
c5 ─┘
c6 ─┐
c7   │ m2 (c6–9)
c8   │ Subclade A
c9 ─┘
c10 ─┐
c11   │
c12   │
c14   │ m3 (c10–17)
c13   │ Subclades B–F
c15   │
c17   │
c16 ─┘
c18 ─┐
c19   │ m4 (c18–20)
c20 ─┘ Subclades G–J

G

Distance from season's centroid

Antigenic cluster

Proportion

A

Antigenic novelty:
Cartographic distance from the recent variants

Variants in
the past 1.5 years

Far

Near

Contemporaneous
variants

Antigenic map

Fitness advantage:
Fitness relative to contemporaneous variants

Frequency

Same emergence time

High

Low

Time

B

Antigenic group
g1
g134

Z

X

Y

C

Estimated frequency

Antigenic group
g1
g134

2010   2015   2020

D

Relative fitness (R_e)

Fitness elevation:
2.22% annually

Antigenic group
g1
g134

Prevalence
at peak
0.05
0.50

2010   2015   2020

Emergence date

E

Fitness advantage (residual)

Antigenic group
g1
g134

Prevalence
at peak
0.05
0.50

2010   2015   2020

Emergence date

F

Fitness advantage (residual)

Spearman r:
0.543

Antigenic novelty
(median distance from recent variants)

2011NH
2012SH
2012NH
2013SH
2013NH
2014SH
2014NH
2015SH
2015NH
2016SH
2016NH
2017SH
2017NH
2018SH
2018NH
2019SH
2019NH

G

Estimated slope

1.47

Overall
2011NH
2012SH
2012NH
2013SH
2013NH
2014SH
2014NH
2015SH
2015NH
2016SH
2016NH
2017SH
2017NH
2018SH
2018NH
2019SH
2019NH

H

Bayesian R-squared

Full   Slope null

A Antigenic group

B Relative fitness (R_e)

C Antigenic groups in the subclade J.2

Antigenic group: g1 → g30

Relative R_e: High → Low

g26, g30, g29

J.2: g25, J.2: g26, J.2: g27, J.2: g28, J.2: g29, J.2: g30, J, Others

D

J, J.2, J.2.1, Others

Subclade J.2

Antigenic cluster: g6, g3, g10, g5, g1, g11, g4, g14, g15, g12, g18, g8, g7, g16, g19, g2, g22, g13, g23, g20, g21, g24, g26, g27, g25, g28, g30, g29

+I160X, +N158K +S145N

E

g29, g30, g28, g25, g27, g26

F79L, S145N, N158K, I160X, P239S

Proportion: 0 → 1

F

g29, g30, g28, g25, g27, g26

g29, g30, g28, g25, g27, g26

Cartographic distance: 0 → 2.0

G

g26

g26 (160:K), g26 (160:L), g26 (160:M), g26 (160:R), g26 (160:T), g26 (160:V), Other J.2 (160:I)

H

2024NH

J.2 decendants

S145N (g29), N158K (g30), I160X (g26), N158K & S145N (g30)

Fitness estimation (up to 2024SH)

PLANT training (up to 2023NH)

2022, 2023, 2024, 2025

A

Distance from the closest vaccine strain

Distance from the season-matched vaccine strain

Cartographic distance

B

Distance from the closest vaccine strain

Within 2 units: 97.2%

Frequency

Cartographic distance

Distance from the season-matched vaccine strain

Within 2 units: 55.1%

Frequency

Cartographic distance

Season

2001SH

2023NH

C

1, Cartography & clustering

Antigenic map

2, Forecasting epidemic dynamics

Past   Future

Vaccination

Frequency

Time

3, Calculating weighted centroid & selecting its nearest strain

Antigenic map

Forecasted frequency (weight)

High

Low

+ Centroid
✕ Weighted centroid
○ Selected strain

D

Median distance

0.76

2.26   1.64

2.40   0.75

WHO-selected   Centroid   Weighted centroid (proposed method)   Hindsight optimal

■ Median

Season

2015SH

2023NH