1   **Systemic Translocation of *S. aureus* Drives Anti-CD4 Autoimmunity in Treated**

2   **HIV Infection**

4   Short title: Autoimmunity in HIV immunopathogenesis

6   Da Cheng[1], Zhenwu Luo[1], Wangbin Ning[1], Sonya L. Heath[2], Magnus Gisslen[3,4,5],

7   Richard W Price[6], Ruth Adekunle[7,8], Tabinda Salman[1], Douglas Johnson[1], John E

8   McKinnon[7,8], Lishomwa C. Ndhlovu[9], Wenhui Hu[10], Wei Jiang[1,7,8*] (ORCiD 0000-

9   0002-1374-053X)

10
11
[1]Department of Pharmacology and Immunology, Medical University of South Carolina,

Charleston, SC, 29425, USA.

[2]Division of Infectious Diseases, Department of Medicine, University of Alabama at

Birmingham, Birmingham, AL, 35294, USA.

[3]Department of Infectious Diseases, Institute of Biomedicine, Sahlgrenska Academy,

University of Gothenburg, 41645, Gothenburg, Sweden

[4]Department of Infectious Diseases, Sahlgrenska University Hospital, Region Västra

Götaland,Gothenburg, Sweden

[5]Public Health Agency of Sweden, Solna, Sweden

[6]Department of Neurology, University of California, San Francisco, San Francisco

General Hospital, San Francisco, CA, 94110, USA

[7]Department of Medicine, Division of Infectious Diseases, Medical University of South

Carolina, Charleston, SC, 29425, USA.

[8]Ralph H. Johnson VA Medical Center, Charleston, SC, USA

[9]Department of Medicine, Division of Infectious Diseases, Weill Cornell Medicine, New

York, NY, 10021, USA

[10]Department of Anatomy and Neurobiology, Virginia Commonwealth

University, Richmond, Virginia, 23298-0709, USA

36   *Corresponding author: Wei Jiang, 173 Ashley Ave. Charleston, SC, 29425, email:

37   jianw@musc.edu

Abstract

Background: Anti-CD4 autoantibodies in people with HIV (PWH) receiving suppressive antiretroviral therapy (ART) appear to prevent CD4+ T cell reconstitution, yet the mechanisms underlying their production remain unclear. Emerging evidence implicates *Staphylococcus aureus* and its peptidoglycan (PGN) in autoimmunity.

Methods: Plasma from 32 ART-naive PWH, 53 ART-treated PWH, and 32 HIV-negative controls was analyzed for IgG autoantibodies and markers of *S. aureus* translocation. Causality was examined in EcoHIV-infected mice administered PGN from *S. aureus* or *Bacillus subtilis*. PGN structure was analyzed via mass spectrometry.

Results: Among 87 autoantibodies, 40% were elevated in ART-naive PWH and largely normalized by ART; however, anti-CD4 IgGs remained elevated in PWH on ART. Anti-CD4 IgG levels inversely correlated with CD4+ T cell counts in ART-treated PWH and positively with *S. aureus* translocation. In mice, *S. aureus* PGN induced anti-CD4 IgGs, reduced gut CD4+ T cells, and promoted surface IgG binding and apoptosis in CD4+ T cells.

Conclusion: *S. aureus* and its PGN translocation may contribute to anti-CD4 autoimmunity and hinder immune recovery in ART-treated PWH, representing a potential therapeutic target.

Keywords: HIV, autoimmunity, anti-CD4 autoantibodies, *Staphylococcus aureus* PGN, failed immune reconstitution, microbiome

Introduction

Recent studies from us and others reveal that autoimmunity contributes to pathogenesis in infectious diseases without clinical autoimmune diseases (Bastard *et al*, 2020; Koning *et al*, 2021; Manry *et al*, 2022; Ning *et al*, 2022b). Approximately 20% of people with HIV (PWH) on antiretroviral therapy (ART) fail to restore CD4+ T cell counts despite plasma viral suppression, leading to increased risks of comorbidities and mortality (Gazzola *et al*, 2009). We demonstrate that anti-CD4 autoantibodies mediate CD4+ T cell death via antibody-mediated cytotoxicity (ADCC) and contributes to poor immune reconstitution following ART (Luo *et al*, 2017a). The effects of anti-CD4 IgG on CD4+ T cell depletion in HIV were further reported by Lisco et al. (Lisco *et al*, 2019). However, the mechanism driving anti-CD4 IgG production during HIV infection remains unknown.

Our previous study revealed that *Staphylococcus aureus* PGN promoted pathogenic autoantibody production and kidney damage in systemic lupus erythematosus (SLE) (Ning *et al*, 2022a). Intraperitoneal (i.p.) injection of heat-killed *S. aureus* or its peptidoglycan (PGN) induced pathogenic autoantibody production in C57BL/6J and autoimmune-prone mice (Luo *et al*, 2019; Ning *et al.*, 2022a). PGN is a major component of the bacterial cell wall and is highly antigenic. PGNs from different bacteria can lead to different immune regulatory activities (Wolf & Underhill, 2018). *S. aureus* nasal colonization is more common in PWH (Rossetti *et al*, 2019). Thus, we

82   investigated the role of *S. aureus* or its PGN in anti-CD4 IgG production and found that

83   *S. aureus* translocation contributes to anti-CD4 IgG production and poor immune

84   recovery in PWH from ART.

86
87
88
89

Results

HIV-associated IgG autoantibody landscape and the unique characteristics of anti-CD4 IgG

We assessed 87 autoantibody IgGs (autoIgGs) from healthy controls, untreated PWH (HIV+/ART-naive), and PWH on suppressive ART (HIV+/ART+). Levels of 35 autoIgGs (40%) were increased and 3 autoIgGs were decreased in HIV+/ART+ compared to controls, while the rest showed similar levels (Figure 1A). ART was associated with normalized autoIgGs except for anti-CD4 and anti-prothrombin protein IgGs, which remained elevated in PWH on ART (HIV+/ART+, Figures 1A-1B). Anti-CD8 IgG levels were unexpectedly lower in HIV+/ART-naive versus controls (Figure 1C). Notably, an inverse correlation was observed between plasma anti-CD4 IgG levels and CD4+ T cell counts in HIV+/ART+ but not in the other two groups (Figure 1D). Moreover, anti-CD4 IgM and IgA levels were similar across the groups. Among anti-CD4 IgGs, IgG1 and IgG3, but not IgG2 or IgG4, were elevated in HIV+/ART+ compared to controls, linking them to persistent low CD4+ T cell counts and poor recovery after ART (Figure S1A-S1B) (Luo *et al.*, 2017a; Song *et al*, 2020).

Elevated anti-CD4 IgG levels are associated with serologic and cellular markers of *S. aureus* translocation in PWH on suppressive ART

Next, we found that plasma levels of IgGs against *S. aureus* antigens were positively correlated with plasma anti-CD4 IgGs in the HIV+/ART+ group (Figure 2A-2B). In the

B-cell RNA sequencing analysis, the *S. aureus* signaling pathway differed between the

HIV+/high versus HIV+/low anti-CD4 IgG groups as well as the HIV+/high anti-CD4

IgG group versus controls (Figure 2C).

Administration of *S. aureus* PGN led to anti-CD4 IgG production in EcoHIV mice

We established and confirmed EcoHIV infection in the mouse model and virus dose

titration (Figure S2). In the current study, serum levels of anti-CD4 IgG, but not total

IgG, were increased in the *S. aureus* PGN group compared to controls (Figure 3A-3B),

indicating that *S. aureus* PGN-mediated antibody production has limited polyclonal B

cell activation. The main subtypes of anti-CD4 IgG increased in HIV+/ART+ were IgG1

and IgG3, supporting its ADCC activity (de Taeye *et al*, 2020; Rozsnyay *et al*, 1989).

In mice, IgG2a and IgG2b, but not IgG1, mediate ADCC activity (Kipps *et al*, 1985).

Notably, anti-CD4 IgG2b and IgG3 were increased among *S. aureus* PGN-elevated

subclasses (Figure 3C).

Administration of *S. aureus* PGN decreased the percentages of gut CD4+ T cells

likely indirectly via increased highly apoptotic IgG binding CD4+ T cells in

EcoHIV mice

To investigate whether *S. aureus* PGN induces pathogenic anti-CD4 IgGs that mediate

CD4+ T cell death *in vivo*, we evaluated the percentage of CD4+ T cells in the gut,

blood, and spleen. Notably, the percentages of CD3+CD4+ T cells among total CD3+

T cells decreased (Figure 3D-3E), and the frequencies of autoreactive IgG surface

binding on CD4+ T cells increased (Figure 4A-4B) in mesenteric lymph nodes and not

in the blood or spleens of *S. aureus* PGN-treated mice compared to the control. In all

three sites tested, IgG+CD4+ T cells were highly apoptotic, but not IgG−CD4+ T cells

(Figure 4C). However, in contrast to gut CD4+CD3+ T cells, *S. aureus* PGN did not

decrease the percentage of gut CD4−CD3+ T cells among total CD3+ T cells; also, we

found that gut CD4−CD3+ T cells rarely had surface autoIgG binding (Figure S3).

Moreover, *S. aureus* PGN did not directly induce gut CD4+ T cell apoptosis *in vitro*

(Figure S4). These results indicate that *S. aureus* PGN may induce gut CD4+ T cell

apoptosis indirectly through autoIgG binding to the CD4+ T cell surface.

***S. aureus* PGN drove CSR via TLR2**

As indicated in Figure 3C, serum anti-CD4 IgG2b and IgG3 subclasses increased in

response to *S. aureus* PGN. CSR is a crucial biological process for antibody maturation

in response to both foreign antigens and self-antigens in the germinal center (Mietzner

et al, 2008). Here, *S. aureus* PGN, but not *B. subtilis* PGN, promoted CSR to IgG2b

and IgG3 via TLR2 *in vitro* (Figure 5A-5B; Figure S6) in response to the optimal PGN

concentration (Figure S5A-S5B). Notably, after denaturing any contaminating proteins,

heat-inactivated *S. aureus* PGN exhibited similar CSR induction compared with the

untreated PGN (Figure S5C). Furthermore, *B. subtilis* PGN induced greater TLR2

signaling activity compared to *S. aureus* PGN (Figure S5D), suggesting that *S. aureus*

PGN-mediated CSR did not exclusively account for its ability to activate TLR2.

PGN structural analysis of *S. aureus* and *B. subtilis*

The length of the sugar polymers of PGN varies among bacterial species; *S. aureus*

consists of ~18 disaccharides, while *Bacillus* has 80-250 disaccharides (Wolf &

Underhill, 2018). In this study, *S. aureus* PGN exhibited the total hydrolysate containing

muramic acid (Mur) and the amino acids lysine (Lys), alanine (Ala), glycine (Gly) and

glutamic acid (Glu). Quantification of amino acids by GC/MS of N-heptafluorobutyric

amino acid isobutylesters shows the molar ratio: 1.4 Ala: 2.9 Gly: 1.0 Glu: 1.1 Lys. *B.*

subtilis PGN exhibited the total hydrolysate containing Mur and the amino acids

diaminopimelic acid (Dpm), Ala, and Glu. The molar ratio is 1.0 Ala: 1.0 Glu: 0.8 Dpm.

The analysis of enantiomers of the total hydrolysate showed the presence of meso-

Dpm and DGlu. In summary, *S. aureus* PGN was A3α (L-Lys-Gly5-6) L-Lys -Gly3

whereas *B. subtilis* PGN was A1γ meso-Dpm-direct, providing the foundation of their

differences in immunomodulatory activities.

Discussion

Autoimmunity plays a pathogenic role in infectious diseases in the absence of any

clinical autoimmune disease (Bastard *et al.*, 2020; Koning *et al.*, 2021; Manry *et al.*,

2022). During HIV infection, CD4+ T cell depletion characterizes immune non-

responders as their CD8+ T cell counts are increased compared to healthy controls

(Luo *et al.*, 2017a). Despite extensive research on poor CD4+ T cell recovery post

ART, no study has proposed a CD4-specific mechanism (Zhang *et al*, 2021) until we

first reported anti-CD4 IgG autoimmunity in HIV (Luo *et al.*, 2017a). The effects of anti-

CD4 IgG on CD4+ T cell depletion in HIV were further reported by other colleagues

(Ouyang *et al*, 2025) (Lisco *et al.*, 2019). Here, we reveal that anti-CD4 IgG

autoimmunity is driven by a specific microbiome *S. aureus*. Notably, PWH from Africa

show a higher failure rate of CD4+ T cell recovery following suppressive ART (Lawn

et al, 2006) as well as a higher rate of *S. aureus* infection or colonization; thus, there

may be a causal link between *S. aureus* translocation, anti-CD4 IgG autoimmunity,

and immune reconstitution failure in HIV (Olalekan *et al*, 2012).

Although anti-CD4 IgG was studied in the pre-ART era, its pathogenic activity was not

proven (Chams *et al*, 1991; Sekigawa *et al*, 1991) until 2017 when we reported

evidence of anti-CD4 IgG-mediated CD4+ T cell death and poor immune recovery

following ART (Luo *et al.*, 2017a). Recently, a study revealed that plasma anti-CD4 IgG

levels increased during HIV infection prior to ART initiation and decreased after ART,

however, levels remained higher in some PWH on ART (Song *et al.*, 2020). Moreover,

an inverse association between plasma anti-CD4 IgG levels and CD4+ T cell counts

after 48 and 96 weeks of ART, but not during pre-ART or after 24 weeks of ART were

reported (Song *et al.*, 2020). Low-affinity nonfunctional or natural autoantibodies can

be induced under persistent inflammation (Hunziker *et al*, 2003). The development of

a pathogenic autoantibody may occur over many years due to disordered somatic

hypermutation and CSR (Elkon & Casali, 2008). Indeed, we reveal that anti-CD4 IgGs

from PWH on ART but not from ART-naive induce ADCC (Cheng *et al*, 2023). Thus,

differing from HIV+/ART+, anti-CD4 IgGs from untreated PWH are likely natural

autoantibodies without cytotoxicity and are generated by high viremia and cytokine

storms. Autoimmune features in HIV often develop after ART initiation (Iordache *et al*,

2014; Zandman-Goddard & Shoenfeld, 2002). These results imply that pathogenic

autoantibodies develop post-ART.

S. aureus colonization and antigen translocation may not be sufficient to induce

pathogenic autoantibody production unless a compromised barrier, host immune

perturbations, or a genetic background susceptible to autoimmunity. Differing from

autoantibodies in autoimmune diseases, pathogenic anti-CD4 IgG in PWH on

suppressive ART may result from the following factors: 1) remaining higher rates of

apoptotic CD4+ T cells lead to increased CD4 self-antigen, especially in the lymph

nodes (Luo *et al*, 2017b); 2) viral proteins are constantly released despite suppressive

ART (Folkvord *et al*, 2005; Heesters *et al*, 2015; Ramratnam *et al*, 2004; Rothenberger

et al, 2015); HIV gp120 increases the binding of CD4 to anti-CD4 antibodies of PWH

(Sekigawa *et al.*, 1991). Consequently, HIV gp120-CD4 accumulates in lymph nodes

and promotes anti-CD4 IgG production (Lederman *et al*, 2011); 3) most PWH do not

have the genetic background of autoimmune diseases.

Although CD4+ T cell apoptosis was observed only in the mesenteric lymph nodes and

not in the spleen after 3-month administration of *S. aureus* PGN, evidence is known

that SIV/HIV infection leads to CD4+ T cell depletion starting from the gut during acute

infection, and develops to peripheral CD4+ T cell depletion during chronic infection due

to exhaustion (Veazey, 2019). We hypothesize that long-term (e.g., one year)

administration of *S. aureus* PGN may eventually deplete peripheral CD4+ T cells.

Notably, *S. aureus* PGN increased the frequencies of autoIgG binding to the surface

of CD4+ T cells in the gut but not in the spleen and blood; the IgG+CD4+ T cells were

highly apoptotic. Mesenteric CD4+ T cell death was not induced by *S. aureus* PGN

directly *in vitro*. These findings suggest that *S. aureus* PGN may induce mesenteric

CD4+ T cell apoptosis indirectly through autoIgG binding on CD4+ T cells and

subsequently exert an ADCC effect. Nonetheless, the mechanisms of *S. aureus* PGN-

mediated gut autoIgG production related to CD4+ T cell decline deserve further

investigation.

The main limitation is the potential contamination in the *S. aureus* PGN we

purchased, which could contain other components (e.g., *Staphylococcal* protein A,

surface protein A, and lipoteichoic acid [LTA]). However, strong evidence supports the

role of *S. aureus* PGN in autoantibody production, as follows: 1) a TLR2 inhibitor

blocked the PGN effects; 2) PGNs can be detected in the circulation in healthy humans

without clinical infections (Huang *et al*, 2019; Molinaro *et al*, 2019; Wolf & Underhill,

2018); 3) only non-protein bacterial antigens can be detected in the blood in humans

(Azzouz *et al*, 2019; Brenchley *et al*, 2006); 4) heat inactivation did not reduce CSR;

5) Staph Protein A and *Staphylococcus* Enterotoxin B reduced SLE severity in lupus-

prone mice (Braun *et al*, 2000; Kim *et al*, 1991); 6) *S. aureus* LTA predominately

induced pro-inflammatory cytokines in myeloid cells, but failed to induce

autoantibodies (Danforth *et al*, 1995; Ning *et al.*, 2022a; Skinner *et al*, 2005; Standiford

et al, 1994). In this translational study, we revealed fundamental findings *in vivo*

through autoantibody screening using human samples and verified causality using an

HIV mouse model.

In summary, this is the first study to show the role of *S. aureus* translocation in inducing

pathogenic anti-CD4 IgGs and CD4+ T cell depletion in HIV, which may provide novel

therapies targeting *S. aureus* colonization or translocation (e.g., receptor inhibitors and

probiotics), and help improve CD4+ T cell recovery and reduce morbidity in PWH on

suppressive ART.

Materials and methods

Human subjects

This study included 32 uninfected controls, 32 untreated PWH (HIV+/ART-naive), and 53 PWH on ART (HIV+/ART+, $\geq$ 2 years of undetectable plasma HIV-1 RNA). All participants were recruited from the Sahlgrenska University Hospital, San Francisco General Hospital, and the Medical University of South Carolina (MUSC) (Gisslen *et al*, 2021; Jiang *et al*, 2021). This study was approved by the local institutional review board. The clinical information is shown in Table 1. PWH were stratified into low and high anti-CD4 IgG subgroups based on a cutoff of 35 ng/ml, determined from 95[th] percentile in healthy controls.

Autoantigen protein array, B-cell gene profiles, and plasma levels of anti-CD4 IgG and subclasses, IgM, and IgA

In recent studies, we described the methods of protein array containing 87 self-antigens, microarray for B-cell gene expression profiles, and plasma anti-CD4 autoantibody levels (Luo *et al.*, 2019; Luo *et al.*, 2017a).

Plasma *S. aureus* antigen translocation

S. aureus lysate was prepared using the BugBuster Master Mix and coated onto plates at 2 µg/100 µL/well at 4°C overnight. The plate was blocked with PBS containing 3% BSA. Diluted plasma samples (1:4000 for IgA, 1:400000 for IgG) were added and

incubated under RT for 60 minutes. Detection was performed using horseradish

peroxidase-labeled anti-human IgG and IgA via 405 nm absorbance with ABTS.

Bacterial PGN structural analysis using mass spectrometry

PGNs from *S. aureus* (ATCC, 6538P) and *B. subtilis* (NR-607, BEI, Manassas, VA)

were separated with an Agilent HPLC system and analyzed using mass spectrometry

(Creative Proteomics, Shirley, NY, USA).

Animal Experiments

Six-week-old female C57BL/6J mice (Jackson Laboratory, Bar Harbor, ME) were

housed at a specific pathogen-free (SPF) facility at MUSC. EcoHIV plasmid (GFP

labeled) was obtained from Dr. Hu from Virginia Commonwealth University (Yin *et*

al, 2016; Yin *et al*, 2017). Briefly, we administered mixed antibiotics (Gibco™ Antibiotic

Mixture) in the drinking water (1mg/mL) for 3 days, and stopped for 2 days. Next,

EcoHIV virus was i.p. injected into B6 mice (50 μL of virus 3.45 x 10^8 IU/mL, n=6 per

group) once. Two weeks post-infection, the mice were i.p. injected with PBS, *S.*

aureus, or *B. subtilis* PGN (InvivoGen, San Diego, CA) twice per week for 12 weeks.

Mouse serum anti-CD4 IgG levels

The 96-well plate was coated with 1 μg/100 μL/well of recombinant mouse CD4 protein

(Sino Biological, Wayne, PA) at 4°C overnight. After blocking with 3% BSA, diluted

serum samples were added and horseradish peroxidase-labeled anti-mouse IgG or its

subclasses were used. Absorbance was measured at 405 nm with ASBT.

B cell class switch DNA recombination (CSR)

B cells were isolated from the spleen of 8-week-old unmanipulated mice using the B

Cell Negative Isolation Kit (STEMCELL, Vancouver, BC). B cells were cultured with

medium or SA-PGN or BS-PGN (10 µg/mL), +/- anti-TLR2 neutralizing antibody or

isotype IgG2a (InvivoGen, 100 ng/mL), or a positive control (LPS from *Escherichia coli*

055:B5 [3 µg/mL, InvivoGen] plus IFN-γ [50 ng/mL, BioLegend] or TGF-β1 [2 ng/mL,

R&D Systems, Minneapolis, MN]). CSR was analyzed by the percentages of

proliferating IgG3 and IgG2b cells using a BD FACSVerse™ Cell Analyzer (BD, San

Jose, CA) after surface staining with anti-B220 (RA3-6B2), anti-IgG2b (RMG2b-1),

anti-IgG3 (R40-82), anti-CD138 (281-2), and anti-IgM (RMM-1) from BioLegend.

IgG surface binding on CD4+ T cells

At the end of the study, cells from blood, spleens, and mesenteric lymph nodes were

collected, stained with anti-CD3 (17A2), anti-CD4 (GK1.5), anti-IgG (Poly4053,

BioLegend), Annexin V (BD Pharmingen), anti-B220 (RA3-6B2), anti-CD19 (1D3),

anti-F4/80 (BM8), anti-IgM (RMM-1), and streptavidin (1:1000, BD Pharmingen), and

analyzed using a BD FACSVerse™ flow cytometer and FlowJo software.

PGN-mediated TLR2 activity

TLR2 activity was detected using the HEK-Blue™ mTLR2 cells (InvivoGen) following

the manufacturer's instruction.

Statistics

 Data were analyzed using Student's t-test or Mann-Whitney test for comparisons

between two groups, and one-way ANOVA with the Kruskal-Wallis test for comparisons

involving more than two groups. Associations were evaluated using Spearman's

correlation test. $P < 0.05$ was considered statistically significant.

Authors' contributions

D.C. wrote the first version of the manuscript. D.C., Z.L., T.S., D.J., and W.N. performed experiments and analyzed data. M.G., R.W., S.H., and Z.L. recruited the study participants. D.C., Z.L., W.J., L.N., R.P., W. H., M.G., JEM, and R.A. were involved in critically revising the manuscript.

Conflicts of Interest

The authors declare no competing interests.

Acknowledgments

We would like to thank John D. Dinolfo, Ph.D. from MUSC for English Language editing. This work was supported by Ralph H. Johnson VA Medical Center, Merit Review Award Number (CX002422, Jiang) from the United States (U.S.) Department of Veterans Affairs (VA) Office of Research and Development (CSR&D) Service, the VA Southeast Network 7 Research Development Award (RDA, Adekunle), the National Institute on Drug of Abuse (R01DA059854, Jiang), National Institute of Allergy and Infectious Diseases (P30AI027767, Saag/Health), National Institute of Health (R01NS094067, Price), the Swedish state, under an agreement between the Swedish government and the county councils (ALF agreement ALFGBG-965885), and the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number UL1TR001450 (the pilot grant, Jiang), TL1TR002382 (Adekunle), and UL1TR002378 (TL1 grant, Adekunle). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Availability of data and materials

Microarray data are available in GEO (Gene Expression Omnibus, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE109837) under accession number GSE109837.

 Figure legends

Figure 1. **HIV-associated IgG autoantibodies.** (A) Z-scored plasma levels of 87

autoantibodies in controls (n = 32), untreated PWH (n = 32), and ART-suppressed

PWH (n = 53). (B) Anti-CD4 and anti-prothrombin IgG levels. (C) Anti-CD8 IgG levels.

(D) Correlation between anti-CD4 IgG levels and CD4+ T cell counts. ANOVA and

Spearman correlation tests were applied.

Figure 2. Association of *S. aureus* serological and cellular markers with anti-CD4

IgG in PWH on ART. (A) Positive correlation between plasma anti-CD4 IgG and anti-

S. aureus IgG levels. (B) Subgroup of HIV+/ART+ individuals shows elevated anti-*S.*

aureus IgG. (C–D) B cell gene expression analysis reveals enriched immune pathways

in PWH vs. controls. Spearman correlation and ANOVA were applied; $P < 0.05$.

Figure 3. *S. aureus* PGN induces anti-CD4 IgG and reduces gut CD4+ T Cells in

EcoHIV mice. C57BL/6J mice received i.p. injections of PBS, *S. aureus* PGN, or *B.*

subtilis PGN twice weekly for 12 weeks (n = 8/group). (A–C) Serum levels of anti-CD4

IgG, total IgG, and anti-CD4 IgG subclasses. (D–E) Gating strategy and CD4+ T cell

percentages in mesenteric lymph nodes, spleen, and blood. Student's t-test and

ANOVA were applied. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$.

Figure 4. *S. aureus* PGN increases apoptotic IgG+CD4+ T Cells in gut-Associated

tissues in EcoHIV mice. (A) Annexin V binding on IgG+CD4+ vs. IgG–CD4+ T cells.

(B–D) Frequencies of IgG+CD4+ T cells in mesenteric lymph nodes, spleen, and

blood. (E–G) Annexin V binding on IgG+CD4+ and IgG−CD4+ T cells in the same

tissues. Student's t-test and one-way ANOVA were applied.

Figure 5. *S. aureus* **PGN induced CSR to IgG2b and IgG3** *in vitro*. Splenic B cells

from unmanipulated C57BL/6J werecultured with the medium (control), *S. aureus*

(SA) or *B. subtilis* (BS) PGN (10 µg/mL) plus TLR2-mAb or isotype Ab (1µg/mL) for 96

hours. The percentages of proliferating IgG2b+ and IgG3+ B cells

(%CFSE^{low}IgG2b/IgG3+ in IgG2b/Ig3+B220+ cells) (A-B) are shown. ANOVA tests.

Figure S1. **Elevated plasma levels of anti-CD4 IgG1 and IgG3 subclasses in PWH**

on ART. (A) The median plasma levels of anti-CD4 IgG, IgM, and IgA were evaluated

in healthy controls (n = 15) and HIV+/ART+ subjects (n = 23). (B) The median plasma

levels of anti-CD4 IgG subclasses (IgG1-IgG4) were determined in controls and

HIV+/ART+ subjects. Mann-Whitney tests.

Figure S2. Confirmation of established EcoHIV infection in the mouse model.

EcoHIV virus was i.p. injected into B6 mice with EcoHIV and verified using qPCR in

blood samples.

Figure S3. *S. aureus* **PGN did not decrease the percent of gut CD4−CD3+ T cells**

in total CD3+ T cells, and rare cells from gut CD4−CD3+ T cells had surface IgG

binding. (A) Gating strategies and (B) percentages of CD4−CD3+ T cells in total CD3+

T cells and IgG+CD4−CD3+ T cells in the mesenteric lymph nodes. ANOVA tests.

Figure S4. ***S. aureus* PGN did not induce mesenteric CD4+ T cell apoptosis**

directly *in vitro.* Lymphocytes from mesenteric lymph nodes were treated with *S.*

aureus PGN (0.1-10 µg/mL) for 4 days. (A) Annexin V binding on the surface of CD4+

T cells, percentages of CD4+ T cells (B) and annexin V binding on CD4+ T cells were

calculated. ANOVA tests.

Figure S5. **Dose titration of PGN for CSR and TLR2 activity**. Different doses of *S.*

aureus PGN (0.1-10 µg/mL) for the induction of the percentages of total IgG2b+ B cells

(A) and proliferating IgG2b+ B cells (B). (C) Comparison of CSR induction by heat-

inactivated vs. untreated *S. aureus* PGN. (D) mTLR2 reporter cell line was treated with

PBS and different doses of *S. aureus* PGN or *B. subtilis* PGN (0.1-10 µg/mL) for 16

hours. The mTLR2 activation was determined using a spectrophotometer at 620 nm.

ANOVA tests.

Figure S6. *S. aureus* PGN induced CSR to IgG2b and IgG3 in plasmablast cells

in vitro. Splenic B cells from unmanipulated mice were cultured with *S. aureus* (SA),

or *B. subtilis* (BS) PGN (10 µg/mL) for 96 hours.Percentages of IgG2b+ and IgG3+

plasmablast cells (%CD138+CFSE^{low}IgG2b/IgG3+ in B220+ cells) (A-B) are shown.

ANOVA tests.

468
469
470

472   **Table 1 Clinical characteristics of study subjects**

473
474

	HIV-negative (n = 32)	HIV+/ART naive (n = 32)	HIV+/ART+ (n = 53)	P value (two HIV+ groups)
Age (years)	49 (34-58)	49 (35-55)	42 (36-52)	0.16
Sex ratio, male: female	21:11	22:10	37:16	0.92
CD4+ T cell counts	797 (554-916)	255 (98-490)	570 (456-724)	< 0.0001
Nadir CD4+ T cell counts		215 (97-490)	294 (164-464)	0.37
Duration of ART (years)			5.5 (4-6)	

475
476

477   ART, antiretroviral therapy; CD4+ T cell count (cells/µL)

478   Data are means (25th percentile, 75th percentile)

References

References

Azzouz D, Omarbekova A, Heguy A, Schwudke D, Gisch N, Rovin BH, Caricchio R,

Buyon JP, Alekseyenko AV, Silverman GJ (2019) Lupus nephritis is linked to disease-

activity associated expansions and immunity to a gut commensal. *Ann Rheum Dis*

78: 947-956

Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann HH, Zhang Y, Dorgham K,

Philippot Q, Rosain J, Beziat V *et al* (2020) Autoantibodies against type I IFNs in

patients with life-threatening COVID-19. *Science* 370

Braun N, Erley C, Klein R, Kotter I, Saal J, Risler T (2000) Immunoadsorption onto

protein A induces remission in severe systemic lupus erythematosus. *Nephrol Dial*

Transplant 15: 1367-1372

Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, Kazzaz Z,

Bornstein E, Lambotte O, Altmann D *et al* (2006) Microbial translocation is a cause of

systemic immune activation in chronic HIV infection. *Nat Med* 12: 1365-1371

Chams V, Idziorek T, Klatzmann D (1991) Biological properties of anti-CD4

autoantibodies purified from HIV-infected patients. *AIDS* 5: 565-569

Cheng D, Luo Z, Fitting S, Stoops W, Heath SL, Ndhlovu LC, Jiang W (2023) The link

between chronic cocaine use, B cell perturbations, and blunted immune recovery in

HIV-infected individuals on suppressive ART. *NeuroImmune Pharm Ther* 2: 71-79

Danforth JM, Strieter RM, Kunkel SL, Arenberg DA, VanOtteren GM, Standiford TJ

(1995) Macrophage inflammatory protein-1 alpha expression in vivo and in vitro: the

role of lipoteichoic acid. *Clin Immunol Immunopathol* 74: 77-83

de Taeye SW, Bentlage AEH, Mebius MM, Meesters JI, Lissenberg-Thunnissen S,

Falck D, Senard T, Salehi N, Wuhrer M, Schuurman J *et al* (2020) FcgammaR

Binding and ADCC Activity of Human IgG Allotypes. *Frontiers in immunology* 11: 740

Elkon K, Casali P (2008) Nature and functions of autoantibodies. *Nat Clin Pract*

Rheumatol 4: 491-498

Folkvord JM, Armon C, Connick E (2005) Lymphoid follicles are sites of heightened

human immunodeficiency virus type 1 (HIV-1) replication and reduced antiretroviral

effector mechanisms. *AIDS Res Hum Retroviruses* 21: 363-370

Gazzola L, Tincati C, Bellistri GM, Monforte A, Marchetti G (2009) The absence of

CD4+ T cell count recovery despite receipt of virologically suppressive highly active

antiretroviral therapy: clinical risk, immunological gaps, and therapeutic options. *Clin*

Infect Dis 48: 328-337

Gisslen M, Keating SM, Spudich S, Arechiga V, Stephenson S, Zetterberg H, Di

Germanio C, Blennow K, Fuchs D, Hagberg L *et al* (2021) Compartmentalization of

cerebrospinal fluid inflammation across the spectrum of untreated HIV-1 infection,

central nervous system injury and viral suppression. *PLoS One* 16: e0250987

Heesters BA, Lindqvist M, Vagefi PA, Scully EP, Schildberg FA, Altfeld M, Walker BD,

Kaufmann DE, Carroll MC (2015) Follicular Dendritic Cells Retain Infectious HIV in

Cycling Endosomes. *PLoS Pathog* 11: e1005285

Huang Z, Wang J, Xu X, Wang H, Qiao Y, Chu WC, Xu S, Chai L, Cottier F, Pavelka
N *et al* (2019) Antibody neutralization of microbiota-derived circulating peptidoglycan
dampens inflammation and ameliorates autoimmunity. *Nat Microbiol* 4: 766-773
Hunziker L, Recher M, Macpherson AJ, Ciurea A, Freigang S, Hengartner H,
Zinkernagel RM (2003) Hypergammaglobulinemia and autoantibody induction
mechanisms in viral infections. *Nat Immunol* 4: 343-349
Iordache L, Launay O, Bouchaud O, Jeantils V, Goujard C, Boue F, Cacoub P,
Hanslik T, Mahr A, Lambotte O *et al* (2014) Autoimmune diseases in HIV-infected
patients: 52 cases and literature review. *Autoimmun Rev* 13: 850-857
Jiang W, Luo Z, Stephenson S, Li H, Di Germanio C, Norris PJ, Fuchs D, Zetterberg
H, Gisslen M, Price RW (2021) Cerebrospinal Fluid and Plasma Lipopolysaccharide
Levels in Human Immunodeficiency Virus Type 1 Infection and Associations With
Inflammation, Blood-Brain Barrier Permeability, and Neuronal Injury. *J Infect Dis* 223:
1612-1620
Kim C, Siminovitch KA, Ochi A (1991) Reduction of lupus nephritis in MRL/lpr mice
by a bacterial superantigen treatment. *J Exp Med* 174: 1431-1437
Kipps TJ, Parham P, Punt J, Herzenberg LA (1985) Importance of immunoglobulin
isotype in human antibody-dependent, cell-mediated cytotoxicity directed by murine
monoclonal antibodies. *J Exp Med* 161: 1-17
Koning R, Bastard P, Casanova JL, Brouwer MC, van de Beek D, with the
Amsterdam UMCC-BI (2021) Autoantibodies against type I interferons are associated
with multi-organ failure in COVID-19 patients. *Intensive Care Med* 47: 704-706
Lawn SD, Myer L, Bekker LG, Wood R (2006) CD4 cell count recovery among HIV-
infected patients with very advanced immunodeficiency commencing antiretroviral
treatment in sub-Saharan Africa. *BMC Infect Dis* 6: 59
Lederman MM, Calabrese L, Funderburg NT, Clagett B, Medvik K, Bonilla H,
Gripshover B, Salata RA, Taege A, Lisgaris M *et al* (2011) Immunologic failure
despite suppressive antiretroviral therapy is related to activation and turnover of
memory CD4 cells. *The Journal of infectious diseases* 204: 1217-1226
Lisco A, Wong CS, Lage SL, Levy I, Brophy J, Lennox J, Manion M, Anderson MV,
Mejia Y, Grivas C *et al* (2019) Identification of rare HIV-1-infected patients with
extreme CD4+ T cell decline despite ART-mediated viral suppression. *JCI Insight* 4
Luo Z, Li M, Wu Y, Meng Z, Martin L, Zhang L, Ogunrinde E, Zhou Z, Qin S, Wan Z *et*
al (2019) Systemic translocation of Staphylococcus drives autoantibody production in
HIV disease. *Microbiome* 7: 25
Luo Z, Li Z, Martin L, Wan Z, Meissner EG, Espinosa E, Wu H, Yu X, Fu P, Julia
Westerink MA *et al* (2017a) Pathological Role of Anti-CD4 Antibodies in HIV-Infected
Immunologic Nonresponders Receiving Virus-Suppressive Antiretroviral Therapy. *J*
Infect Dis 216: 82-91
Luo Z, Zhou Z, Ogunrinde E, Zhang T, Li Z, Martin L, Wan Z, Wu H, Qin Z, Ou T *et al*
(2017b) The effect of plasma auto-IgGs on CD4+ T cell apoptosis and recovery in
HIV-infected patients under antiretroviral therapy. *J Leukoc Biol*

Manry J, Bastard P, Gervais A, Le Voyer T, Rosain J, Philippot Q, Michailidis E,

Hoffmann HH, Eto S, Garcia-Prat M *et al* (2022) The risk of COVID-19 death is much

greater and age dependent with type I IFN autoantibodies. *Proc Natl Acad Sci U S A*

119: e2200413119

Mietzner B, Tsuiji M, Scheid J, Velinzon K, Tiller T, Abraham K, Gonzalez JB, Pascual

V, Stichweh D, Wardemann H, Nussenzweig MC (2008) Autoreactive IgG memory

antibodies in patients with systemic lupus erythematosus arise from nonreactive and

polyreactive precursors. *Proc Natl Acad Sci U S A* 105: 9727-9732

Molinaro R, Mukherjee T, Flick R, Philpott DJ, Girardin SE (2019) Trace levels of

peptidoglycan in serum underlie the NOD-dependent cytokine response to

endoplasmic reticulum stress. *J Biol Chem* 294: 9007-9015

Ning W, Cheng D, Howe PH, Bian C, Kamen DL, Luo Z, Fu X, Ogunrinde E, Yang L,

Wang X *et al* (2022a) Staphylococcus aureus peptidoglycan (PGN) induces

pathogenic autoantibody production via autoreactive B cell receptor clonal selection,

implications in systemic lupus erythematosus. *J Autoimmun* 131: 102860

Ning W, Xu W, Cong X, Fan H, Gilkeson G, Wu X, Hughes H, Jiang W (2022b)

COVID-19 mRNA vaccine BNT162b2 induces autoantibodies against type I

interferons in a healthy woman. *J Autoimmun* 132: 102896

Olalekan AO, Schaumburg F, Nurjadi D, Dike AE, Ojurongbe O, Kolawole DO, Kun

JF, Zanger P (2012) Clonal expansion accounts for an excess of antimicrobial

resistance in Staphylococcus aureus colonising HIV-positive individuals in Lagos,

Nigeria. *Int J Antimicrob Agents* 40: 268-272

Ouyang Y, Wu K, Fu L, Yi P, Cheng D, Fu X (2025) Effect of anti-CD4 mAb induced

by inhibiting B cell disorder on immune reconstruction of HIV-infected immunological

non-responders. *Mol Med* 31: 244

Ramratnam B, Ribeiro R, He T, Chung C, Simon V, Vanderhoeven J, Hurley A, Zhang

L, Perelson AS, Ho DD, Markowitz M (2004) Intensification of antiretroviral therapy

accelerates the decay of the HIV-1 latent reservoir and decreases, but does not

eliminate, ongoing virus replication. *J Acquir Immune Defic Syndr* 35: 33-37

Rossetti B, Lombardi F, Belmonti S, D'Andrea MM, Tordini G, D'Avino A, Borghetti A,

Moschese D, De Luca A, Montagnani F (2019) Nasopharyngeal bacterial and fungal

colonization in HIV-positive versus HIV-negative adults. *New Microbiol* 42: 37-42

Rothenberger MK, Keele BF, Wietgrefe SW, Fletcher CV, Beilman GJ, Chipman JG,

Khoruts A, Estes JD, Anderson J, Callisto SP *et al* (2015) Large number of

rebounding/founder HIV variants emerge from multifocal infection in lymphatic

tissues after treatment interruption. *Proc Natl Acad Sci U S A* 112: E1126-1134

Rozsnyay Z, Sarmay G, Walker M, Maslanka K, Valasek Z, Jefferis R, Gergely J

(1989) Distinctive role of IgG1 and IgG3 isotypes in Fc gamma R-mediated functions.

Immunology 66: 491-498

Sekigawa I, Groopmen JE, Allan JD, Ikeuchi K, Biberfield G, Takatsuki K, Byrn RA

(1991) Characterization of autoantibodies to the CD4 molecule in human

immunodeficiency virus infection. *Clinical immunology and immunopathology* 58:

617 145-153

618 Skinner NA, MacIsaac CM, Hamilton JA, Visvanathan K (2005) Regulation of Toll-like
619 receptor (TLR)2 and TLR4 on CD14dimCD16+ monocytes in response to sepsis-
620 related antigens. *Clin Exp Immunol* 141: 270-278

621 Song A, Li Z, Luo Z, Lu X, Wang R, Liu L, Xia W, Wan Z, Zhang T, Su B *et al* (2020)
622 Effects of Early and Delayed Antiretroviral Therapy on Plasma Anti-CD4 Autoreactive
623 IgG and Its Association With CD4(+) T-Cell Recovery in Acute HIV-Infected
624 Individuals. *Front Pharmacol* 11: 449

625 Standiford TJ, Arenberg DA, Danforth JM, Kunkel SL, VanOtteren GM, Strieter RM
626 (1994) Lipoteichoic acid induces secretion of interleukin-8 from human blood
627 monocytes: a cellular and molecular analysis. *Infect Immun* 62: 119-125

628 Veazey RS (2019) Intestinal CD4 Depletion in HIV / SIV Infection. *Curr Immunol Rev*
629 15: 76-91

630 Wolf AJ, Underhill DM (2018) Peptidoglycan recognition by the innate immune
631 system. *Nat Rev Immunol* 18: 243-254

632 Yin C, Zhang T, Li F, Yang F, Putatunda R, Young WB, Khalili K, Hu W, Zhang Y
633 (2016) Functional screening of guide RNAs targeting the regulatory and structural
634 HIV-1 viral genome for a cure of AIDS. *AIDS* 30: 1163-1174

635 Yin C, Zhang T, Qu X, Zhang Y, Putatunda R, Xiao X, Li F, Xiao W, Zhao H, Dai S *et*
636 *al* (2017) In Vivo Excision of HIV-1 Provirus by saCas9 and Multiplex Single-Guide
637 RNAs in Animal Models. *Mol Ther* 25: 1168-1186

638 Zandman-Goddard G, Shoenfeld Y (2002) HIV and autoimmunity. *Autoimmun Rev* 1:
639 329-337

640 Zhang Y, Jiang T, Li A, Li Z, Hou J, Gao M, Huang X, Su B, Wu H, Zhang T, Jiang W
641 (2021) Adjunct Therapy for CD4(+) T-Cell Recovery, Inflammation and Immune
642 Activation in People Living With HIV: A Systematic Review and Meta-Analysis. *Front*
643 *Immunol* 12: 632119

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

674
675

Figure S2

A

B

Mesenteric lymph nodes

%CD4−CD3+ in total T cells

%IgG+ on CD4−CD3+ T cells

Figure S3

Figure S4

Figure S5

Figure S6