Systemic translocation of *Staphylococcus aureus* promotes autoimmunity:

implications in autoantibody-mediated poor immune reconstitution from

antiretroviral therapy in HIV

Short title: Autoimmunity in HIV immunopathogenesis

Da Cheng[1], Zhenwu Luo[1], Wangbin Ning[1], Sonya L. Heath[2], Magnus Gisslen[3,4,5],

Richard W Price[6], Ruth Adekunle[7,8], Tabinda Salman[1], Douglas Johnson[1], John E

McKinnon[7,8], Lishomwa C. Ndhlovu[9], Reafa Hossain[10], Wenhui Hu[10], Wei Jiang[1,7,8*]

(ORCiD 0000- 0002-1374-053X)

[1]Department of Pharmacology and Immunology, Medical University of South

Carolina, Charleston, SC, 29425, USA.

[2]Division of Infectious Diseases, Department of Medicine, University of Alabama at

Birmingham, Birmingham, AL, 35294, USA.

[3]Department of Infectious Diseases, Institute of Biomedicine, Sahlgrenska Academy,

University of Gothenburg, 41645, Gothenburg, Sweden

[4]Department of Infectious Diseases, Sahlgrenska University Hospital, Region Västra

Götaland, Gothenburg, Sweden

[5]Public Health Agency of Sweden, Solna, Sweden

[6]Department of Neurology, University of California, San Francisco, San Francisco

General Hospital, San Francisco, CA, 94110, USA

[7]Department of Medicine, Division of Infectious Diseases, Medical University of

South Carolina, Charleston, SC, 29425, USA.

[8]Ralph H. Johnson VA Medical Center, Charleston, SC, USA

[9]Department of Medicine, Division of Infectious Diseases, Weill Cornell Medicine,

New York, NY, 10021, USA

[10]Department of Anatomy and Neurobiology, Virginia Commonwealth

University, Richmond, Virginia, 23298-0709, USA

35 *Corresponding author: Wei Jiang, 173 Ashley Ave. Charleston, SC, 29425, email:

36 jianw@musc.edu

Abstract

Background: In 2017, our group first demonstrated that autoimmunity contributes to HIV pathogenesis, even without autoimmune disease. This concept is now broadly recognized, exemplified by the role of autoimmunity in severe COVID-19. In people with HIV (PWH) on suppressive ART, anti-CD4 autoantibodies may impair CD4+ T cell recovery, though the mechanisms driving their production remain unclear. Building on evidence from our group and others that *Staphylococcus aureus* and its peptidoglycan (PGN) promote autoimmunity, we investigated their contribution to anti-CD4 IgG in HIV.

Methods: Plasma from 32 ART-naive PWH, 53 ART-treated PWH, and 32 HIV-negative controls was analyzed for IgG autoantibodies and markers of *S. aureus* translocation using protein array, ELISA, and microarray. EcoHIV mice were injected intraperitoneally with saline, *S. aureus* PGN, or *Bacillus subtilis* PGN. PGN structures were compared by mass spectrometry.

Results: Among 87 autoantibodies, 40% were elevated in ART-naive PWH and largely normalized by ART; however, anti-CD4 IgGs remained elevated in PWH on ART. Anti-CD4 IgG levels inversely correlated with CD4+ T cell counts in ART-treated PWH and positively with *S. aureus* translocation. In mice, *S. aureus* PGN induced anti-CD4 IgGs, reduced gut CD4+ T cells, and promoted surface IgG binding and apoptosis in CD4+ T cells.

Conclusion: *S. aureus* and its PGN translocation may drive anti-CD4 autoimmunity

and hinder immune recovery in PWH on suppressive ART, highlighting *S. aureus*

colonization as a therapeutic target and supporting the development of competitive

probiotic interventions such as *Bacillus subtilis*.

Keywords: HIV, autoimmunity, anti-CD4 autoantibodies, failed immune reconstitution,

microbiome, *Staphylococcus aureus*

Introduction

Recent studies, including ours, show that autoimmunity contributes to pathogenesis in infectious diseases without clinical autoimmune conditions (1). About 20% of PWH on ART fail to restore CD4+ T cell counts despite viral suppression, increasing comorbidity and mortality risks (2). We demonstrated that anti-CD4 autoantibodies drive CD4+ T cell death via ADCC, contributing to poor immune reconstitution post-ART (3), a finding supported by other researchers (4), though outcomes vary (5). The mechanisms driving anti-CD4 IgG production in HIV remain unclear.

Our prior work (6) demonstrated that S. aureus PGN induces pathogenic autoantibodies and kidney injury in SLE. Intraperitoneal injection of heat-killed S. aureus or its PGN elicited autoantibody responses in both C57BL/6J and autoimmune-prone mice (6, 7). As a major antigenic cell wall component, PGN exerts distinct immune effects depending on its bacterial source (8). Because S. aureus colonization is more frequent in PWH (9), we examined links between autoantibodies and CD4+ T cell counts and tested the role of S. aureus PGN in anti-CD4 IgG production. Our findings show that S. aureus PGN promotes anti-CD4 IgG generation, leading to CD4+ T cell loss and impaired immune recovery in PWH on suppressive ART.

Results

Autoantibody landscape and the unique characteristics of anti-CD4 IgGs

Among the 87 autoantibody IgGs (autoIgGs) in HIV+/ART+, 35 autoIgGs (40%) were

elevated, 3 were reduced, and the rest were similar to controls (Figure 1A). ART

normalized most autoIgGs, but anti-CD4 and anti-prothrombin IgGs remained

elevated in HIV+/ART+ (Figure 1A-1B). Anti-CD8 IgG was lower in HIV+/ART-naive

versus controls (Figure 1C). Plasma anti-CD4 IgG levels inversely correlated with

CD4+ T cell counts in HIV+/ART+ only (Figure 1D). Anti-CD4 IgM and IgA levels

were comparable across groups. In HIV+/ART+, anti-CD4 IgG1 and IgG3, but not

IgG2 or IgG4, were elevated (Figure S1A-S1B) (3, 10). IgG1 and IgG3 in humans

support ADCC (11, 12). In addition, plasma levels of anti-prothrombin IgG were

directly correlated with anti-CD4 IgGs (r = 0.22, P = 0.01), but not correlated with

CD4+ T cell counts (r = -0.16, P = 0.22).

Elevated anti-CD4 IgG levels are associated with plasma levels of *S. aureus*

translocation in PWH on suppressive ART

Our previous study in systemic lupus erythematosus (SLE) demonstrated that

Staphylococcus aureus contributes to the production of pathogenic lupus-associated

anti–double-stranded DNA IgG antibodies and to kidney damage (6). Based on these

findings, we evaluated the potential role of *S. aureus* in the generation of anti-CD4

autoantibodies in HIV infection. To assess systemic exposure to *S. aureus*, we

measured plasma IgG levels against *S. aureus* antigens as a surrogate of bacterial

translocation. We found that plasma IgG levels against *S. aureus* antigens were

positively correlated with anti-CD4 IgG levels in HIV+/ART+ individuals. (Figure 2A-

2B).

***S. aureus* PGN induced anti-CD4 IgG production in EcoHIV mice**

EcoHIV mouse study design is shown in Figure 3A and infection was verified by

qPCR (Figure S2). S. aureus PGN increased serum anti-CD4 IgG without affecting

total IgG (Figure 3B), indicating limited polyclonal B cell activation. In mice, IgG2a

and IgG2b mediate ADCC (13), and S. aureus PGN selectively increased anti-CD4

IgG2b and IgG3 (Figure 3C). By contrast, B. subtilis PGN, a probiotic (14), did not

induce anti-CD4 IgG (Figure 3B–3C).

S. aureus PGN reduced gut CD4+ T cells through apoptosis of IgG-bound CD4+

T cells in EcoHIV mice

Furthermore, S. aureus PGN decreased CD3+CD4+ T cells (Figure 3D) and

increased surface-bound autoIgG on CD4+ T cells in the gut, but not in blood or

spleen (Figure 4A–4D). IgG+CD4+ T cells were highly apoptotic at all sites, in

contrast to IgG–CD4+ T cells (Figure 4E-4G). CD4–CD3+ T cell frequencies were

unaffected, with minimal IgG binding (Figure S3). In vitro, S. aureus PGN did not

directly induce gut CD4+ T cell apoptosis (Figure S4). These results indicate that S.

aureus PGN indirectly drives gut CD4+ T cell apoptosis through autoIgG binding.

S. aureus PGN drives class switch DNA recombination (CSR) via TLR2 in vitro.

Both native and heat-inactivated S. aureus PGN promoted CSR through TLR2, unlike

B. subtilis PGN (Figures 5A–5B, S5A–S5C, S6). Notably, B. subtilis PGN induced

stronger TLR2 signaling than S. aureus PGN (Figure S5D), suggesting that S. aureus

PGN–mediated CSR involves additional mechanisms beyond TLR2 activation.

PGN structural analysis

S. aureus PGN (~18 disaccharide units) contains muramic acid (Mur) and the amino

acids Lys, Ala, Gly, and Glu, with a molar ratio of 1.4 Ala: 2.9 Gly: 1.0 Glu: 1.1 Lys

(GC/MS). In contrast, B. subtilis PGN (80–250 disaccharide units) contains Mur,

Dpm, Ala, and Glu, with a molar ratio of 1.0 Ala: 1.0 Glu: 0.8 Dpm, and includes

meso-Dpm and D-Glu enantiomers. Structurally, S. aureus PGN is classified as A3α

(L-Lys–Gly$_{5-6}$, L-Lys–Gly$_3$), while B. subtilis PGN is A1γ (meso-Dpm–direct),

accounting for their distinct immunomodulatory properties.

Discussion

Autoimmunity contributes to infectious disease pathogenesis, even in the absence of

clinical autoimmune disease (1). We first reported anti-CD4 IgG autoimmunity in HIV

in 2017, linking it to CD4+ T cell depletion and poor immune recovery post-ART (3), a

finding later supported by others (15, 16). Here, we show that S. aureus translocation

drives anti-CD4 IgG production, potentially explaining higher CD4+ T cell recovery

failure in African PWH with prevalent S. aureus colonization (17).

Anti-CD4 IgG was first described in the pre-ART era (18), but its pathogenicity

remained unclear until our 2017 study demonstrated its role in poor CD4+ T cell

recovery post-ART (3). Recent studies show that anti-CD4 IgG levels rise during

untreated HIV infection, decline with ART, yet persist in a subset of PWH on ART; an

inverse correlation with CD4+ T cell counts was observed at 48 and 96 weeks post-

ART, but not before ART or at 24 weeks (10), while other reports found no

association after prolonged ART (5). Low-affinity, non-cytotoxic natural

autoantibodies may emerge from chronic inflammation (19), whereas pathogenic

autoantibodies require disordered somatic hypermutation and CSR over time. Unlike

pre-ART anti-CD4 IgGs, which are non-cytotoxic natural autoantibodies likely

induced by viremia and inflammation, pathogenic anti-CD4 IgGs in PWH on ART

mediate ADCC and develop post-ART (15, 20-22). Anti-prothrombin IgG (aPT IgG) is

an antiphospholipid antibody directed against prothrombin; our study reveals that

plasma levels of anti-prothrombin IgG were directly correlated with anti-CD4 IgG

levels, but not with CD4+ T cell counts, suggesting that these two autoantibodies

may be generated through a similar mechanism and anti-prothrombin IgG does not

contribute to CD4+ T cell counts. Previous studies (23, 24) have shown increased

levels of anti-prothrombin IgG in HIV-infected, predominantly ART-untreated

individuals; however, these antibodies were typically transient and not strongly

associated with thrombotic events or overt antiphospholipid syndrome.

S. aureus colonization and antigen translocation alone may not trigger pathogenic

autoantibody production without a compromised barrier, immune perturbations, or

genetic predisposition to autoimmunity. Unlike autoantibodies in autoimmune

diseases, pathogenic anti-CD4 IgG in PWH on suppressive ART may arise from: 1)

elevated apoptotic CD4+ T cells increasing CD4 self-antigen in lymph nodes (25); 2)

persistent viral protein release, including HIV gp120, enhancing CD4 binding to anti-

CD4 IgG and promoting its production (26); 3) the absence of autoimmune-prone

genetic backgrounds in most PWH differentiates these responses from classical

autoimmune diseases.

CD4+ T cell apoptosis was detected only in mesenteric lymph nodes, not in the

spleen or blood, after S. aureus PGN administration. Although IgG+CD4+ T cells

exhibited higher levels of apoptosis than IgG−CD4+ T cells across all sites (Figure

4E-4G), increased IgG surface binding on CD4+ T cells following *S. aureus* PGN

intraperitoneal injection was observed only in mesenteric lymph nodes, and not in the

spleen or blood (Figure 4B-4D). The site-specific effects likely reflect differences in

local exposure to SA-PGN following intraperitoneal administration. Mesenteric lymph

nodes are expected to encounter higher concentrations of SA-PGN than the spleen

or peripheral blood. Since SIV/HIV infection initiates CD4+ T cell loss in the gut

during acute infection and progresses to peripheral depletion in chronic stages (27),

we hypothesize that prolonged exposure to S. aureus PGN (e.g., 1 year) may also

deplete peripheral CD4+ T cells. Importantly, mesenteric CD4+ T cell death was not

induced by PGN directly in vitro, suggesting an indirect mechanism via autoIgG

binding and cell apoptosis. The mechanisms driving S. aureus PGN-mediated gut

autoIgG production and CD4+ T cell decline require further studies.

Previous studies reveal that bacterial lipoteichoic acid (LTA), a TLR2 ligand, primarily

induces TLR2-dependent pro-inflammatory cytokines rather than autoantibodies (6,

28). Generally, TLR2 signaling in B cells more often supports cytokine secretion,

activation marker upregulation, or differentiation than direct mitogenesis, comparable

to strong mitogens such as CpG (TLR9) or anti-IgM (29). In addition, the downstream

pathways and outcomes of TLR2 activation can differ in cytokine production and

proliferation. S. aureus PGN exhibited structural differences compared with B. subtilis

PGN. An additional mechanism may account for S. aureus PGN-mediated B cell

CSR, which warrants further investigation.

The limitations of this study are: 1) there is no figure reported by the PGN structure

analysis from the company (Creative Proteomics, Shirley, NY, USA); 2) the potential

contamination of protein A or LTA in S. aureus PGN. Nonetheless, several lines of

evidence support PGN as a driver of autoantibody production: (1) a TLR2 inhibitor

blocked PGN effects; (2) PGN, but not the other Gram+ bacterial products, is

detectable in human circulation without a clinical bacterial infection (8); (3) S. aureus

protein A and enterotoxin B reduce lupus autoimmunity in mice (30, 31); and (5) LTA,

a TLR2 ligand, primarily induces pro-inflammatory cytokines rather than

autoantibodies (6, 28).

Currently, no treatment is available for improving CD4+ T cell recovery in PWH on

suppressive ART. Up to 20% of PWH on ART fail to restore peripheral CD4+ T cell

counts to the levels of healthy controls; increased morbidity and mortality have been

demonstrated in these patients, which is the main challenge in the HIV clinic. This

study reveals that *S. aureus* and its PGN translocation may drive anti-CD4

autoimmunity and hinder immune recovery in PWH on suppressive ART, highlighting

S. aureus colonization as a therapeutic target and supporting the development of

competitive probiotic interventions such as *Bacillus subtilis* (32).

Materials and methods

Human subjects

This study enrolled 32 controls (without HIV), 32 HIV+/ART-naïve, and 53

HIV+/ART+ individuals (>2 years undetectable plasma HIV-1 RNA) from Sahlgrenska

University Hospital (Gothenburg, Sweden), San Francisco General Hospital (San

Francisco, CA, USA), Ralph H. Johnson Veteran Health Care System (RHJ VAHCS,

SC, USA), and the Medical University of South Carolina (MUSC, SC, USA). The local

institutional review board approved the study. Demographics and clinical details are

provided in Table 1. PWH were divided into low and high anti-CD4 IgG subgroups

using a 35 ng/ml cutoff, based on the 95^{th} percentile in controls.

Autoantigen protein array, B-cell gene profiles, and plasma levels of anti-CD4

autoantibodies

We reported the methods of protein array containing 87 self-antigens, B cell gene

microarray, and plasma anti-CD4 autoantibody levels (3, 7).

Plasma *S. aureus* antigen translocation

S. aureus lysate was prepared using the BugBuster Master Mix and coated onto

high-binding plates (Sino Biological, Wayne, PA) at 2 µg/100 µL/well. The plate was

blocked with 3% BSA and added with diluted plasma samples (1:4000 for IgA,

1:400000 for IgG). Detection was performed using horseradish peroxidase-labeled

anti-human IgG and IgA.

Bacterial PGN structural analysis

PGNs from *S. aureus* (ATCC, 6538P) and *B. subtilis* (NR-607, BEI, Manassas, VA)

were separated with an Agilent HPLC system and analyzed using mass spectrometry

(Creative Proteomics, Shirley, NY, USA).

Mice Experiment

Mouse study design was shown in Figure 3A. EcoHIV plasmid was obtained from

Dr. Hu (Virginia Commonwealth University) (33). Six-week-old female C57BL/6J mice

(Jackson Laboratory, Bar Harbor, ME) received mixed antibiotics (Gibco™, 1 mg/mL)

in drinking water for 3 days and a 2-day break. Mice were injected i.p. with EcoHIV

(50 µL, 3.45×10^8 IU/mL; n = 6/group), followed by i.p. of PBS, S. aureus PGN, or

B. subtilis PGN (InvivoGen, San Diego, CA) twice weekly for 12 weeks. Whole blood

was drawn via the tail vein method from EcoHIV-injected mice under aseptic

conditions. To verify ecoHIV infection, the blood was used to isolate DNA via the

Quick-DNA/RNA Miniprep Plus Kit (Zymo; Irvine, CA, Cat # D7003). The standard

curve was generated from viral titer and threshold cycles using the ABM qPCR

lentivirus titer kit (LV900). qPCR for HIV-LTR was performed for the unknown and

standard samples via QuantaBio PerfecTa Toughmix (QuantaBio, Beverly, MA),

supplemented with the following primers and a probe: LTRF, 5'-

CTGGCTAATTAGGGAACCCACTG-3';   LTRR,   5'-

GGACTAAACGGATCTGAGGGATCTC-3';   and   the   LTRP,   5'-(FAM)-

TTACCAGAGTCACACAACAGACGGGCA-(MGBNFQ)-3'. The thermal program used

was 3 min at 55°C, 10 min at 95°C, and 40 cycles of 15 sec at 95°C and 1 min at

60°C. qPCR was performed at the Bio-Rad CFX96 qPCR instrument (Bio-Rad

Laboratories, Hercules, CA), and data analysis was done with the built-in Bio-Rad

software. The number of proviral copies in the unknown sample is the number of the

total number of copies minus the number of copies in the control reaction. All

samples were run in triplicate.

Mouse serum anti-CD4 IgG levels

Serum anti-CD4 IgG was detected using recombinant mouse CD4 protein (1 μg/100

μL/well) and horseradish peroxidase-labeled anti-mouse IgG or its subclasses.

B cell class switch DNA recombination (CSR)

B cells, isolated from spleens of 8-week-old mice (STEMCELL B Cell Negative

Isolation Kit), were cultured with medium, S. aureus PGN, or B. subtilis PGN (10

μg/mL), ± anti-TLR2 or isotype IgG2a (100 ng/mL, InvivoGen), or positive controls

(LPS from E. coli 055:B5 [3 μg/mL, InvivoGen] plus IFN-γ [50 ng/mL, BioLegend] or

TGF-β1 [2 ng/mL, R&D Systems]). CSR was measured by proliferating IgG3 and

IgG2b B cell percentages using a BD FACSVerse™ Cell Analyzer after staining with

anti-B220, anti-IgG2b, anti-IgG3, anti-CD138, and anti-IgM antibodies (BioLegend).

IgG surface binding on CD4+ T cells

Cells from blood, spleen, and mesenteric lymph nodes were stained with anti-CD3,

anti-CD4, anti-IgG, Annexin V, and streptavidin, then analyzed on a BD

FACSVerse™ flow cytometer using FlowJo software.

PGN-mediated TLR2 activity

TLR2 activity was detected using the HEK-Blue™ mTLR2 cells (InvivoGen) following

the manufacturer's instruction.

Statistics

Data were analyzed using Student's t-test or Mann–Whitney test for two-group

comparisons and one-way ANOVA for multiple groups. Associations were assessed

by Spearman correlation. A P value < 0.05 was considered significant.

Authors' contributions

D.C. wrote the first version of the manuscript. D.C., Z.L., T.S., D.J., and W.N. performed experiments and analyzed data. M.G., R.W., S.H., and Z.L. recruited the study participants. D.C., Z.L., W.J., L.C.N., R.P., R.H., W. H., M.G., JEM, and R.A. were involved in critically revising the manuscript.

Conflicts of Interest

L.C.N. reports grants from the NIH and has received consulting fees from work as a scientific advisor for AbbVie, ViiV Healthcare, Ledidi AS, serves on the Board of CytoDyn and has financial interests in Ledidi AS all for work outside of the submitted work. All other authors declare no competing interests.

Acknowledgments

We thank John D. Dinolfo, Ph.D. (MUSC), for English language editing. This work was supported by the Ralph H. Johnson VA Health Care System I01CX002422 (Jiang) from the U.S. Department of Veterans Affairs; the VA Southeast Network 7 Research Development Award (Adekunle); NIH grants R01DA059854 (Jiang), R01DA059538 (Jiang), R01DA056876 (Hu), P30AI027767 (Saag/Health), R01NS094067 (Price), UL1TR001450, TL1TR002382 (Adekunle), and UL1TR002378 (Adekunle, TL1). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Availability of data and materials

Data are available.

Figure legends

Figure 1. HIV-associated IgG autoantibodies. (A) Z-scored plasma levels of 87 autoantibodies in controls (n = 32), untreated PWH (n = 32), and ART-suppressed PWH (n = 53). (B) Anti-CD4 and anti-prothrombin IgG. (C) Anti-CD8 IgG. (D) Correlation between anti-CD4 IgG and CD4+ T cell counts. ANOVA and Spearman correlation. Samples were analyzed for autoantibodies using a protein array and collected from four institutions: Sahlgrenska University Hospital, San Francisco General Hospital, RHJ VAHCS, and MUSC.

Figure 2. Association of S. aureus markers with anti-CD4 IgG in PWH on ART. (A) Correlation of plasma anti-CD4 IgG with anti-S. aureus IgG. (B) Elevated plasma anti-S. aureus IgG in HIV+/ART+. Spearman correlation and ANOVA. PWH were divided into low and high anti-CD4 IgG subgroups using a 35 ng/ml cutoff, based on the 95th percentile in controls. Samples from MUSC were analyzed for anti-CD4 IgG and anti-S. aureus IgG by ELISA.

Figure 3. S. aureus PGN induces anti-CD4 IgG and reduces gut CD4+ T cells in EcoHIV mice. C57BL/6J mice received i.p. PBS, S. aureus PGN, or B. subtilis PGN. (A) Mouse study design. (B–C) Serum anti-CD4 IgG, total IgG, and IgG subclasses. (D) Gating strategy and CD4+ T cell frequencies in the three sampling sites. Student's t-test and ANOVA.

Figure 4. S. aureus PGN increases apoptotic IgG+CD4+ T cells in gut-associated tissues. (A) Annexin V binding on IgG+ vs. IgG– CD4+ T cells. (B–D)

Frequencies of IgG+CD4+ T cells in mesenteric lymph nodes, spleen, and blood. (E–

G) Annexin V binding on IgG+ vs. IgG– CD4+ T cells. Student's t-test and ANOVA.

Figure 5. S. aureus PGN induces CSR to IgG2b and IgG3 in vitro. Splenic B cells

from naive C57BL/6J mice were cultured with medium, and PGNs (10 µg/mL) ±

TLR2-mAb or isotype Ab (1 µg/mL) for 96 h. Percentages of proliferating IgG2b+ and

IgG3+ B cells (%CFSE^{low}IgG2b/IgG3+ in B220+ cells) are shown (A–B). ANOVA.

Figure S1. **Elevated plasma anti-CD4 IgG1 and IgG3 subclasses in PWH on**

ART. (A) The median plasma levels of anti-CD4 IgG, IgM, and IgA in controls (n =

15) and HIV+/ART+ subjects (n = 23). (B) The median plasma levels of anti-CD4 IgG

subclasses (IgG1-IgG4) between the two groups. Mann-Whitney tests.

Figure S2. Confirmation of established EcoHIV infection in mice. The EcoHIV

virus was verified by qPCR in blood samples collected one month after infection.

Figure S3. S. aureus PGN does not affect gut CD4–CD3+ T cells in vivo. (A)

Gating strategy. (B) Percentages of CD4–CD3+ T cells among total CD3+ T cells

and IgG+CD4–CD3+ T cells in mesenteric lymph nodes. ANOVA.

Figure S4. S. aureus PGN does not directly induce gut CD4+ T cell apoptosis in

vitro. Lymphocytes from mesenteric lymph nodes were treated with S. aureus PGN

(0.1–10 µg/mL) for 4 days. (A-B) Percentages of CD4+ T cells and Annexin V+CD4+

T cells. ANOVA.

Figure S5. PGN dose titration for CSR and TLR2 activity. (A–B) S. aureus PGN

induced total (A) and proliferating (B) IgG2b+ B cells. (C) CSR induction by heat-

inactivated vs. native S. aureus PGN. (D) mTLR2 reporter cell responses after 16 h

treatment. ANOVA.

Figure S6. S. aureus PGN induces CSR in vitro. Splenic B cells from naive mice

were cultured with PGN (10 μg/mL) for 96 h. Percentages of CD138+ IgG2b+ and

IgG3+ plasmablasts (%CD138+CFSE^lowIgG2b/IgG3+ in B220+ cells) are shown

(A–B). ANOVA.

425 **Table 1 Clinical characteristics of study participants**

426
427

	HIV-negative/Controls (n = 32)	HIV+/ART naive (n = 32)	HIV+/ART+ (n = 53)	P value (two HIV+ groups)

Age (years)	49 (34-58)	49 (35-55)	42 (36-52)	0.16
Sex ratio, male: female	21:11	22:10	37:16	0.92
CD4+ T cell counts	797 (554-916)	255 (98-490)	570 (456-724)	< 0.0001
Nadir CD4+ T cell counts		215 (97-490)	294 (164-464)	0.37
Duration of ART (years)			5.5 (4-6)	

430   ART, antiretroviral therapy; CD4+ T cell count (cells/µL)

431   Data are means (25th percentile, 75th percentile)

References

1.   Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann HH, Zhang Y, Dorgham K, Philippot
Q, Rosain J, Beziat V, Manry J, Shaw E, Haljasmagi L, Peterson P, Lorenzo L, Bizien L,
Trouillet-Assant S, Dobbs K, de Jesus AA, Belot A, Kallaste A, Catherinot E, Tandjaoui-
Lambiotte Y, Le Pen J, Kerner G, Bigio B, Seeleuthner Y, Yang R, Bolze A, Spaan AN,
Delmonte OM, Abers MS, Aiuti A, Casari G, Lampasona V, Piemonti L, Ciceri F, Bilguvar K,
Lifton RP, Vasse M, Smadja DM, Migaud M, Hadjadj J, Terrier B, Duffy D, Quintana-Murci L,

van de Beek D, Roussel L, Vinh DC, Tangye SG, et al. 2020. Autoantibodies against type I
IFNs in patients with life-threatening COVID-19. Science 370.

2.   Gazzola L, Tincati C, Bellistri GM, Monforte A, Marchetti G. 2009. The absence of CD4+ T
cell count recovery despite receipt of virologically suppressive highly active antiretroviral
therapy: clinical risk, immunological gaps, and therapeutic options. Clin Infect Dis 48:328-37.

3.   Luo Z, Li Z, Martin L, Wan Z, Meissner EG, Espinosa E, Wu H, Yu X, Fu P, Julia Westerink
MA, Kilby JM, Wu J, Huang L, Heath SL, Li Z, Jiang W. 2017. Pathological Role of Anti-
CD4 Antibodies in HIV-Infected Immunologic Nonresponders Receiving Virus-Suppressive
Antiretroviral Therapy. J Infect Dis 216:82-91.

4.   Lisco A, Wong CS, Lage SL, Levy I, Brophy J, Lennox J, Manion M, Anderson MV, Mejia Y,
Grivas C, Mystakelis H, Burbelo PD, Perez-Diez A, Rupert A, Martens CA, Anzick SL, Morse
C, Chan S, Deleage C, Sereti I. 2019. Identification of rare HIV-1-infected patients with
extreme CD4+ T cell decline despite ART-mediated viral suppression. JCI Insight 4.

5.   Bowler SA, Premeaux TA, Ratzan L, Friday C, Gianella S, Landay AL, Ndhlovu LC, NWCS
ACTG. 2025. Plasma anti-CD4 IgG levels are associated with poor immune recovery in
people with HIV initiating antiretroviral therapy. AIDS 39:208-210.

6.   Ning W, Cheng D, Howe PH, Bian C, Kamen DL, Luo Z, Fu X, Ogunrinde E, Yang L, Wang
X, Li QZ, Oates J, Zhang W, White D, Wan Z, Gilkeson GS, Jiang W. 2022. Staphylococcus
aureus peptidoglycan (PGN) induces pathogenic autoantibody production via autoreactive B
cell receptor clonal selection, implications in systemic lupus erythematosus. J Autoimmun
131:102860.

7.   Luo Z, Li M, Wu Y, Meng Z, Martin L, Zhang L, Ogunrinde E, Zhou Z, Qin S, Wan Z,
Westerink MAJ, Warth S, Liu H, Jin P, Stroncek D, Li QZ, Wang E, Wu X, Heath SL, Li Z,
Alekseyenko AV, Jiang W. 2019. Systemic translocation of Staphylococcus drives
autoantibody production in HIV disease. Microbiome 7:25.

8.   Wolf AJ, Underhill DM. 2018. Peptidoglycan recognition by the innate immune system. Nat
Rev Immunol 18:243-254.

9.   Rossetti B, Lombardi F, Belmonti S, D'Andrea MM, Tordini G, D'Avino A, Borghetti A,
Moschese D, De Luca A, Montagnani F. 2019. Nasopharingeal bacterial and fungal
colonization in HIV-positive versus HIV-negative adults. New Microbiol 42:37-42.

10.   Song A, Li Z, Luo Z, Lu X, Wang R, Liu L, Xia W, Wan Z, Zhang T, Su B, Jiang W, Wu H.
2020. Effects of Early and Delayed Antiretroviral Therapy on Plasma Anti-CD4 Autoreactive
IgG and Its Association With CD4(+) T-Cell Recovery in Acute HIV-Infected Individuals.
Front Pharmacol 11:449.

11.   de Taeye SW, Bentlage AEH, Mebius MM, Meesters JI, Lissenberg-Thunnissen S, Falck D,
Senard T, Salehi N, Wuhrer M, Schuurman J, Labrijn AF, Rispens T, Vidarsson G. 2020.
FcgammaR Binding and ADCC Activity of Human IgG Allotypes. Front Immunol 11:740.

12.   Rozsnyay Z, Sarmay G, Walker M, Maslanka K, Valasek Z, Jefferis R, Gergely J. 1989.
Distinctive role of IgG1 and IgG3 isotypes in Fc gamma R-mediated functions. Immunology
66:491-8.

13.   Kipps TJ, Parham P, Punt J, Herzenberg LA. 1985. Importance of immunoglobulin isotype in
human antibody-dependent, cell-mediated cytotoxicity directed by murine monoclonal

antibodies. J Exp Med 161:1-17.

14.   Piewngam P, Khongthong S, Roekngam N, Theapparat Y, Sunpaweravong S, Faroongsarng D,
Otto M. 2023. Probiotic for pathogen-specific Staphylococcus aureus decolonisation in
Thailand: a phase 2, double-blind, randomised, placebo-controlled trial. Lancet Microbe
4:e75-e83.

15.   Ouyang Y, Wu K, Fu L, Yi P, Cheng D, Fu X. 2025. Effect of anti-CD4 mAb induced by
inhibiting B cell disorder on immune reconstruction of HIV-infected immunological non-
responders. Mol Med 31:244.

16.   Epling BP, Lisco A, Manion M, Laidlaw E, Galindo F, Anderson M, Roby G, Sheikh V,
Migueles SA, Poole A, Perez-Diez A, Liu X, Rao VK, Burbelo PD, Sereti I. 2024. Impact of
Anti-CD4 Autoantibodies on Immune Reconstitution in People With Advanced Human
Immunodeficiency Virus. Clin Infect Dis doi:10.1093/cid/ciae562.

17.   Lawn SD, Myer L, Bekker LG, Wood R. 2006. CD4 cell count recovery among HIV-infected
patients with very advanced immunodeficiency commencing antiretroviral treatment in sub-
Saharan Africa. BMC Infect Dis 6:59.

18.   Chams V, Idziorek T, Klatzmann D. 1991. Biological properties of anti-CD4 autoantibodies
purified from HIV-infected patients. AIDS 5:565-9.

19.   Hunziker L, Recher M, Macpherson AJ, Ciurea A, Freigang S, Hengartner H, Zinkernagel RM.
2003. Hypergammaglobulinemia and autoantibody induction mechanisms in viral infections.
Nat Immunol 4:343-9.

20.   Zandman-Goddard G, Shoenfeld Y. 2002. HIV and autoimmunity. Autoimmun Rev 1:329-37.

21.   Cheng D, Luo Z, Fitting S, Stoops W, Heath SL, Ndhlovu LC, Jiang W. 2023. The link
between chronic cocaine use, B cell perturbations, and blunted immune recovery in HIV-
infected individuals on suppressive ART. NeuroImmune Pharm Ther 2:71-79.

22.   Olalekan AO, Schaumburg F, Nurjadi D, Dike AE, Ojurongbe O, Kolawole DO, Kun JF,
Zanger P. 2012. Clonal expansion accounts for an excess of antimicrobial resistance in
Staphylococcus aureus colonising HIV-positive individuals in Lagos, Nigeria. Int J
Antimicrob Agents 40:268-72.

23.   Loizou S, Singh S, Wypkema E, Asherson RA. 2003. Anticardiolipin, anti-beta(2)-
glycoprotein I and antiprothrombin antibodies in black South African patients with infectious
disease. Ann Rheum Dis 62:1106-11.

24.   Becker AC, Libhaber E, Sliwa K, Singh S, Stewart S, Tikly M, Essop MR. 2011.
Antiphospholipid antibodies in black south africans with hiv and acute coronary syndromes:
prevalence and clinical correlates. BMC Res Notes 4:379.

25.   Luo Z, Zhou Z, Ogunrinde E, Zhang T, Li Z, Martin L, Wan Z, Wu H, Qin Z, Ou T, Zhang J,
Ma L, Liao G, Heath S, Huang L, Jiang W. 2017. The effect of plasma auto-IgGs on CD4+ T
cell apoptosis and recovery in HIV-infected patients under antiretroviral therapy. J Leukoc
Biol doi:10.1189/jlb.5A0617-219R.

26.   Ramratnam B, Ribeiro R, He T, Chung C, Simon V, Vanderhoeven J, Hurley A, Zhang L,
Perelson AS, Ho DD, Markowitz M. 2004. Intensification of antiretroviral therapy accelerates
the decay of the HIV-1 latent reservoir and decreases, but does not eliminate, ongoing virus
replication. J Acquir Immune Defic Syndr 35:33-7.

546 27. Veazey RS. 2019. Intestinal CD4 Depletion in HIV / SIV Infection. Curr Immunol Rev 15:76-
547   91.

548 28. Danforth JM, Strieter RM, Kunkel SL, Arenberg DA, VanOtteren GM, Standiford TJ. 1995.
549   Macrophage inflammatory protein-1 alpha expression in vivo and in vitro: the role of
550   lipoteichoic acid. Clin Immunol Immunopathol 74:77-83.

551 29. Bekeredjian-Ding I, Jego G. 2009. Toll-like receptors--sentries in the B-cell response.
552   Immunology 128:311-23.

553 30. Braun N, Erley C, Klein R, Kotter I, Saal J, Risler T. 2000. Immunoadsorption onto protein A
554   induces remission in severe systemic lupus erythematosus. Nephrol Dial Transplant 15:1367-
555   72.

556 31. Kim C, Siminovitch KA, Ochi A. 1991. Reduction of lupus nephritis in MRL/lpr mice by a
557   bacterial superantigen treatment. J Exp Med 174:1431-7.

558 32. Williams RE. 1963. Healthy carriage of Staphylococcus aureus: its prevalence and importance.
559   Bacteriol Rev 27:56-71.

560 33. Yin C, Zhang T, Qu X, Zhang Y, Putatunda R, Xiao X, Li F, Xiao W, Zhao H, Dai S, Qin X,
561   Mo X, Young WB, Khalili K, Hu W. 2017. In Vivo Excision of HIV-1 Provirus by saCas9 and
562   Multiplex Single-Guide RNAs in Animal Models. Mol Ther 25:1168-1186.

Figure 1

Figure 2

Figure 3

A

Figure 4

Figure 5

Figure S1

Figure S2

A

B

Mesenteric lymph nodes

Figure S3

Figure S4

Figure S5

Figure S6