Evolutionary rescue in a consumer-resource system depends on the affected ecological traits and the population's resident life-history

Abstract

With evolutionary rescue, a population that is declining due to an environmental change adapts to its environment, avoiding extinction. Previous theoretical work has explored the effects of negative density-dependence on rescue, showing that it may aid or hinder persistence. However, these models typically assume that it is only population intrinsic growth rates, r, or carrying capacity, K, that are negatively affected, and do not model density-dependence explicitly. Here, we analyze evolutionary rescue in a consumer-resource species following an abrupt environmental change, characterizing how rescue is dependent on the ecological effects of the environmental change on the consumer, which differently affect r and K through subsequent interactions with an explicit non-substitutable resource species. We derive approximate analytical predictions for the fixation probabilities of beneficial alleles, mutational supply, and times to extinction, which work well when selection is weak and individual turnover rates are low. We demonstrate that consumer rescue is dependent on the ecological effect of the environmental change, the resident life-history of the population, and the genetic architectures of evolving traits (monogenic versus polygenic). Our model suggests that measurements of intrinsic growth rates alone will be insufficient to predict rescue probabilities. This work extends our understanding of the interplay between ecology and evolution in influencing population persistence.

Introduction

Understanding the factors underlying the ability of populations to persist in response to

environmental change is of utmost importance, particularly in light of the rapidly accelerating

loss of biodiversity due to climate change (Berg et al. 2010, Bellard et al. 2012). One such

mechanism of persistence is through evolutionary adaptation, or specifically evolutionary rescue

(see Carlson et al. 2014; Bell 2017 for reviews), the process by which a population that is

declining due to an environmental perturbation subsequently evolves and adapts to its new

environment, avoiding local extinction. Theoretical models of evolutionary rescue (hereafter

'rescue') are characterized by the concurrent modelling of both evolutionary and demographic

processes, whereby successful rescue of a population is contingent on its rate of adaptation

outpacing its rate of demographic decline (Gomulkiewicz & Holt 1995). Classical models of

rescue have typically considered a single isolated species evolving to avoid extinction (e.g.

Gomulkiewicz & Holt 1995; Burger & Lynch 1995; Lande & Shannon 1996). This work has

recovered many important insights into the factors underlying persistence, including the positive

roles of additive genetic variance in fitness and of initial population size. Later work has

extended this theory to consider in detail the effects of population structure (Uecker et al. 2014),

recombination (Uecker & Hermisson 2016), generation time (Draghi et al. 2024), and various

other factors (Orr & Unckless 2008, 2014; Uecker 2017; Osmond et al. 2020; Xu 2023). The

simplifying assumption of a single species existing in isolation, however, fails to capture

ecological interactions among individuals such as predation and competition.

Ecological interactions may drastically influence demographic and evolutionary

outcomes, and therefore rescue probabilities. Predation typically imposes a demographic cost

on prey species, accelerating demographic decline. However, predation may also positively

influence prey rescue if predators preferentially consume maladapted individuals–accelerating

the rate of prey adaptation–and the selective benefit outweighs the demographic cost (Jones

2008; Osmond et al. 2017). A similar concept also applies if maladapted individual experience

greater interspecific competition (Osmond & de Mazancourt 2013). If selection and density-

dependence act on prey birth rates, prey evolution may also be accelerated via the effect of

predation to reduce prey density-dependence, reducing prey generation time (Osmond 2017;

also see Draghi et al. 2024). Predators may also experience "indirect evolutionary rescue,"

whereby rescue in a declining predator population is instead driven by prey adaptive evolution, if

the prey exhibit trade-offs between growth and defense against predation (Yamamichi & Miner

2015; Cortez & Yamamichi 2019; Hermann & Becks 2022; van Velzen 2023). It is also possible

for a species with lower population size to exhibit a higher probability of rescue than its more

abundant competitor, if the resource distribution evolves towards its optimum and away from its

competitor (Van Den Elzen et al. 2017).

Theoretical work has also suggested that negative density-dependence, particularly

intra-specific competition, also reduces rescue probabilities (Chevin & Lande 2010, Osmond &

de Mazancourt 2013, Nordstrom et al. 2023; Draghi et al. 2024), although the magnitude of this

effect remains controversial in the literature (Bell 2017). While the models of negative density-

dependence analyzed by these authors differ in certain mathematical specifications, all of them

share a number of simplifying assumptions that obscure the relationship between demography

and evolution. It is often either assumed that selection acts solely on intrinsic growth rates, r,

(e.g. Chevin & Lande 2010; Nordstrom et al. 2023) or population carrying capacity, K (Osmond

& de Mazancourt 2013), representing opposites on the continuum of r- and K-selection.

However, these two parameters are likely not independent in reality (Pianka 1970, Mallet 2012,

Travis et al. 2023), and there is much empirical evidence to suggest that r and K may co-vary

positively or negatively due to varying life-history strategies in yeast (Wei & Zhang 2019),

bacteria (Reding-Roman et al. 2017; Marshall et al. 2023), and spider mites (Bisschop et al.

2021). For example, selection acting to increase individual resource acquisition rates may

increase r, but decrease K by reducing total resource availability among conspecifics at high

densities (Agrawal & Whitlock 2012), thus intensifying density-dependence. However, selection

acting to increase the efficiency of resource use would lead to a greater r as well as K by

lowering the per-capita resource requirement for maintenance (Travis et al. 2023). In contrast,

r-selected models of rescue assume that the intensity of negative density-dependence is solely

a function of population size/density, and itself doesn't evolve (since K is held constant), thus

neglecting the above relationships. Currently, we lack a theoretical framework to understand

evolutionary rescue through the evolution of ecological traits that differentially influence both r

and K, representing a significant gap in the literature.

In this paper, we use a consumer-resource model with overlapping generations to

investigate how evolutionary rescue of a consumer species is influenced by the ecological

effects of an abrupt environmental change that, in the absence of evolution, would lead to

extinction. Briefly, consumer-resource models explicitly track both consumer and resource

population dynamics (see Sakarchi & Germain 2024 for a review). Consumer growth is a

function of various ecological factors: resource acquisition rates, the efficiency of resource

conversion into offspring, resource abundance, and mortality rates, whereas resource growth is

typically modelled using logistic growth minus consumption by consumers. Consumer density-

dependence arises via decreasing resource availability, settling on an equilibrium between

resource consumption and replenishment. Using this framework, we characterize how

evolutionary rescue in a consumer species is influenced by the ecological trait affected by the

environmental change. We consider both a one-locus and polygenic basis to the affected

ecological traits, exploring the interplay between demography, selection, resource competition,

and generation time. We begin our analysis by deriving analytical predictions for the fixation

probabilities of rescue alleles that influence each ecological trait, the expected number of rescue

mutations to occur during the process of population decline, and the mean time to consumer

extinction in the absence of evolution, which work well under weak selection and low individual

turnover rates (i.e. slow-growing populations with long generation times). Given the complexity

of our eco-evolutionary model, we then use forward-time simulations to explore areas of the

parameter space where our theoretical approximations are not met, that is, for populations that

exhibit rapid growth at low densities and have short generation times.

Ecological model

107 We modelled the ecological dynamics of the system using a discrete-time consumer-

108 resource model (MacArthur 1970; Abrams 2019). Consumer ecological dynamics are described

109 by the following recurrence relation:

$$N_{t+1} = N_t(aCR_t + S), \tag{1}$$

where N_t is consumer abundance at time-step t, a is the resource acquisition rate, C is the

conversion efficiency of acquired resources into offspring, R_t is resource abundance at time-

step t, and S is the survival rate. Our life-cycle proceeds with consumer births and resource

consumption occurring simultaneously, followed by deaths. We consider overlapping

generations using a simple form of age-structure. Individuals are able to reproduce and are

susceptible to mortality beginning the time-step following their birth, and individual fecundity

(aCR_t) remains constant with age. An individual's absolute fitness at time-step t, W_t, is therefore

$W_t = aCR_t + S$. The quantity W_t also describes the geometric growth rate of a lineage with trait

values a, C, and S, in a population with resource abundance R_t. The ecological dynamics of the

resource are described by the following recurrence relation,

$$R_{t+1} = R_t(1 + r_R(1 - \frac{R_t}{K_R}) - aN_t), \tag{2}$$

122 where r_R is the intrinsic growth rate of the resource and K_R is its carrying capacity.

Simulation methods

We ran individual-based forward-time simulations using SLiM v4.01 (Haller & Messer

2022). Briefly, we initialized our simulations with consumers and resources at demographic

equilibrium. An abrupt environmental change was introduced at time-step t_* such that the

population would go extinct in the absence of evolutionary rescue. A population was recorded

as having gone extinct if its population size equaled 0. Conversely, a population was recorded

as having rescued if it returned to a non-zero equilibrium, either through the fixation of a single

beneficial allele (one-locus simulations) or through a return to its phenotypic optimum

(quantitative trait simulations). Downstream analyses were performed using the programming

language R 4.4.2 (R Core Team 2024).

We modelled the total number of births to be Poisson distributed with a mean of $N_t a C R_t$

for the consumers and a mean of $R_t r_R (1 - \frac{R_t}{K_R})$ for the resources. Once the total number of

offspring was established, parentage was retroactively assigned for consumer offspring by

drawing from a multinomial distribution, with the probability of an individual being assigned

parentage proportional to its fecundity. Consumer deaths were modelled by drawing from a

binomial distribution with the probability of survival equal to the individual's survivorship. We did

not explicitly model resource death as its net growth is already captured by the r_R term, which

we allocate to births for brevity. Resource consumption was modelled using a binomial

distribution with R_t trials and probability of success $a N_t$, the per-capita resource consumption

rate.

One locus rescue

Consider a monomorphic consumer population at demographic equilibrium, $\widehat{N}_0$ with

ecological trait values a_0, C_0, and S_0, consuming a resource population with intrinsic growth rate

r_R and carrying capacity K_R. An abrupt environmental change occurs at time t_* reducing the

mean growth rate of the resident population below unity (i.e. $W_{t_*} < 1$) either via a reduction in

individual resource acquisition rates, resource conversion efficiency, or survivorship. In the

absence of evolution, the resident population, now exhibiting trait values a_{res}, C_{res}, and S_{res} will

go extinct even at low density when the resource is at its carrying capacity (i.e., $a_{res}C_{res}K_R +$

$S_{res} < 1$, such that $\widehat{N}_{res} \leq 0$, where $\widehat{N}_{res}$ is the equilibrium consumer population size following the

environmental change). Evolutionary rescue may occur if and only if a mutant lineage that may

persist in this novel environment ($\widehat{N}_{mut} > 0$) invades. Note that the mutant lineage need not

return to $\widehat{N}_0$. However, without loss of generality, we will assume in our simulations that all

beneficial mutations have the effect of return the ecological trait to its value prior to the

environmental change. We also assume for now that the consumer population is haploid and

asexual to simplify the mathematics; we later relax this assumption when considering polygenic

selection.

160   The probability of evolutionary rescue at time-step t, P_t, is equal to one minus the

161   probability that all rescue mutants arising at time-step t go extinct: $P_t = 1 - (1 - \psi_t)^{N_t U_t}$, where

162   ψ_t and U_t refer to the probability of fixation of a rescue mutation and the expected number of

163   rescue mutations per individual at time-step t, respectively (Orr & Unckless 2008). Following

164   this logic, the total probability of rescue is equal to $P = 1 - \prod_{t=0}^{\infty}(1 - \psi_t)^{N_t U_t}$. Assuming ψ_t and

165    $N_t U_t$ are small for any given t, we approximate the above formula as an exponential decay:

166
$$P = 1 - \exp\left[-\sum_{t=0}^{\infty} \psi_t N_t U_t\right] \tag{3}$$

167   We solve U_t by considering that the number of mutations that arise at any time-point is simply a

168   function of an individual's fecundity multiplied by its mutation rate μ. Thus, the total mutation rate

169   per individual at time-point t is

170
$$U_t = aCR_t\mu. \tag{4}$$

From eq. 4 we observe that the per-individual mutational supply will be decreased if an

environmental perturbation reduces consumer fecundity instead of survivorship, all else

remaining equal.

We now derive analytical predictions of ψ_t for mutant lineages. Consider a mutant allele

that exhibits ecological trait values $a_{\text{res}}(1 + \delta_a), C_{\text{res}}(1 + \delta_C)$, and $S_{\text{res}}(1 + \delta_S)$. The fixation

probability of such a mutant allele will depend on its absolute fitness. If the individual first

carrying the mutant allele has an absolute fitness below replacement ($a_{\text{res}}(1 + \delta_a)C_{\text{res}}(1 +$

$\delta_C)R_t + S_{\text{res}}(1 + \delta_S) < 1$), its lineage will deterministically go extinct. Fixation is only possible if

the allele is able to increase its absolute representation in the population ($a_{\text{res}}(1 + \delta_a)C_{\text{res}}(1 +$

$\delta_C)R_t + S_{\text{res}}(1 + \delta_S) > 1$), although it must also escape stochastic loss when rare. Following the

logic of branching processes, the probability that a new mutant allele experiences stochastic

loss is equal to the probability that the individual in which it arises has some number of

offspring, all of whose lineages eventually go extinct, and–with non-overlapping generations–

that the individual itself fails to survive (Fisher 1922; Haldane 1927; Wright 1931; Otto &

Whitlock 1997). The fixation probability of the mutant allele that occurs in a single individual at

time t is thus equal to one minus its probability of stochastic loss. Assuming the number of

offspring produced per individual in a time-step follows a Poisson distribution, we derive the

following formula:

189
$$\psi_t = 1 - \left(\sum_{0}^{\infty} \frac{e^{f_t} f_t^k}{k!} (1 - \psi_{t+1})^k \right)(1 - S(1 + \delta_S)\psi_{t+1}), \tag{5}$$

where $f_t = a_{\text{res}}(1 + \delta_a)C_{\text{res}}(1 + \delta_C)R_t$ is the fecundity of a mutant individual at time-step t. Given

that fecundity is density-dependent, $\psi_t \neq \psi_{t+1}$ in a population of changing size. Note that the

fitness of the mutant allele is a function of the resource abundance, R_t. Thus, in a declining

consumer population (and increasing resource population), the individual carrying the mutant

allele will experience an additional boost in fecundity and, consequently, an increased fixation

probability the later its occurrence (i.e. $\psi_{t+1} > \psi_t$). Assuming $a_{res}, \delta_a, C_{res}, \delta_C, S_{res}, \delta_S,$ and ψ_t

are small and a separation of time-scales between consumer and resource growth, we derive a

continuous-time approximation for the change in ψ as a function of N, which we solve to get the

fixation probability as a function of the current population size, $\psi(N_t)$ (see *Appendix A* for the

derivation). Ignoring terms of order $O(\psi^2)$ or above, the fixation probability of a mutant allele that

increases fecundity for a given population size, $\psi_f(N_t)$, either via increasing a or C is:

201
$$\psi_f(N_t) \approx \frac{\delta_f r_R \big(a_{res} C_{res} R_t (1 + \delta_f) + S_{res} - 1\big)}{a_{res} S_{res}(1 + \delta_f)(C_{res} K_R r_R \delta_f + N_t(1 - S_{res}))}, \qquad (6a)$$

202   where $\delta_f = \delta_a$ when the mutant allele increases a and $\delta_f = \delta_C$ when it increases C (we do not

203   consider the case where a mutant allele increases both). Conversely, a mutant allele that

204   increases survivorship, $\psi_S(N_t)$, will experience fixation probability:

205
$$\psi_S(N_t) \approx \frac{\delta_S r_R (a_{res} C_{res} R_t + S_{res}(1 + \delta_S) - 1)}{a_{res} C_{res} K_R (1 + \delta_S)(r_R S_{res} \delta_S + a_{res} N_t(1 - S_{res}(1 + \delta_S)))}, \qquad (6b)$$

A quick analysis of eqs. 6a and 6b reveals that fixation probabilities of both fecundity and

survival mutants are maximized as $N_t \rightarrow 0$, as mutant individuals benefit from increased

resource availability ($R_t \rightarrow K_R$), thus increasing their number of offspring.

Our last step is to derive a general solution for N_t and R_t in the resident population. To

do so, we must make two key assumptions. First, we assume that resource abundance is at

equilibrium with respect to consumer abundance and trait values ("separation of time-scales")

and, second, that few consumer birth and death events occurs per time-step such that

consumer growth is approximately continuous (i.e. low individual turnover). Dropping subscripts,

we find that $\frac{dN}{dt} = N(aCK_R \left(1 - \frac{aN}{r_R}\right) + S - 1)$, where $R = K_R \left(1 - \frac{aN}{r_R}\right)$. Here consumer population

growth is logistic, and $\frac{dN}{dt}$ can be solved to get $N_t = K_N \frac{N_0}{N_0 + e^{-r_N t}(K_N - N_0)}$, where $r_N = a\,C\,K_R + S -$

1and $K_N = \frac{r_N r_R}{a^2 C K_R}$. Note that $r_N = W_{R_t = K_R} - 1$, the growth rate of the consumer when resource

availability is maximal. When $r_N > 0$, the population is able to grow at low density. This general

solution for N_t can then be used to approximate R_t with the separation of time-scales

assumption. We use these forms for N_t and R_t to estimate ψ_t and U_t, from which we

numerically estimate the total probability of rescue, P, using eq. 3. Mean times to extinction, t_e,

may be approximated by solving for the time-step at which there is only one consumer individual

remaining in the population ($N_{t_e} = 1$) given the resident ecological trait values following the

environmental change. Note that our quasi-deterministic approximation of N_t and R_t here will

generally over-estimate t_e as it neglects demographic stochasticity and tends to over-estimate

the resource abundance.

We also cross-examine our analytical predictions of ψ_t and U_t with numerical

predictions. We numerically predict ψ_t using an iterative method whereby we solve for ψ_{t_*} at

equilibrium ($\psi_t = \psi_{t+1} = \psi_{t_*}$), which occurs when $N_t = N_{t+1} = 0$. We then retroactively solve for

ψ_t as a function of ψ_{t+1} (see eq. 4) by moving backwards in time from the point at which the

consumer population went extinct. We numerically predict U_t by calculating it from observed

consumer and resource abundances in the simulations, thus giving its expected value.

We use forward-time simulations to test the accuracy of our approximate theory and

characterize how the probability of evolutionary rescue depends on whether acquisition rates,

conversion efficiencies, or survivorships are negatively impacted. We simulate the scenario

where an abrupt environmental change affects one ecological trait such that the population will

go extinct in the absence of evolution, and we draw comparisons among these three scenarios

with respect to their intrinsic growth rates following the environmental change, r_{N*} (where $r_{N*} <$

0). We model rescue mutations to occur *de novo* at constant rate μ per offspring following the

environmental change, with rescue mutations returning the ecological trait to its value prior to

environmentally induced maladaptation.

We simulate evolutionary rescue of a "slow-growing" consumer population (low

fecundity, high survivorship) with an intrinsic growth rate of $r_N = 0.03$ and $K_N = 2084$ and a

"fast-growing" consumer population (high fecundity, low survivorship) with an intrinsic growth

rate of $r_N = 0.4$ and $K_N = 2230$ prior to the environmental change (see fig. 1 caption for

parameters). Despite the number of assumptions made in our approximate theory, it performs

remarkably well in the slow-growing case (Fig. 1A squares), when birth and death rates are low

and selection weak. In the fast-growing case, the approximate theory substantially over-predicts

rescue probabilities (Fig. 1B squares). Here our assumptions of small fixation probabilities and

continuous-time population growth are not met due to strong selection and rapid consumer

growth. This results in over-estimation of fixation probabilities and, more importantly, substantial

over-estimates of the resource abundance and therefore mutational supply (see eq. 4). In this

parameter space, consumer and resource intrinsic growth rates are the same order of

magnitude, such that the resource does not reach a quasi-equilibrium with respect to consumer

density.

We correct these discrepancies to some degree by instead using numerical estimates of

fixation probabilities and estimated mutational supply from observed data, although we still

observe some over-estimation of rescue probabilities for slower rates of decline (Fig. 1B

triangles). In this regime, our theoretical framework will always over-predict fixation probabilities

and ultimately rescue because it neglects several sources of stochasticity (birth, consumption,

death) from our simulations that are most pertinent during intermediate and high consumer

densities. Overall, our analytical theory provides a useful starting point to develop intuition for

our rescue scenario, but is nonetheless limited in scope due to the complexity of our eco-

evolutionary model, which incorporates aspects of density-dependent demography, selection,

and overlapping generations with two demographic variables (N_t and R_t).

Figure 1. Probabilities of one-locus rescue as a function of population intrinsic growth rates and the affected ecological trait following an abrupt environmental change in (A) a slow-growing population ($a_0 = 9.5 \times 10^{-5}, C_0 = 0.0095, S_0 = 0.95, r_R = 0.475, K_R = 95,000$) and (B) a fast-growing population ($a_0 = 9.5 \times 10^{-5}, C_0 = 0.1\ S_0 = 0.5, r_R = 0.475, K_R = 95,000$). The mutation rate is 10^{-4}. Small dots connected by a line represent data from at least 1000 replicate simulations. Squares represent rescue predictions using eqs. 6a & 6b and our quasi-deterministic theory applied to eq. 3. Triangles represent numerical predictions of fixation probability using our iterative numerical method alongside estimates of expected mutation supply using simulation data.

The central result from our simulations is that the probability of evolutionary rescue depends on the affected (and subsequently evolving) ecological trait, and the ecological trait most favourable to rescue via *de novo* mutations depends on the population's resident life-history (Fig. 1). Resident populations exhibiting the slow-growing strategy experience the greatest probability of rescue from declines in survivorship, followed by acquisition rates and then conversion efficiencies (Fig. 1A). On the other hand, populations exhibiting the fast-growing resident strategy exhibit the greatest probability of rescue following declines in

acquisition rates, with declines in survivorship and conversion efficiency resulting in similar

rescue probabilities (Fig. 1B).

When fecundity is negatively impacted, consumer populations always experience higher

probabilities of rescue when their resource acquisition rates are reduced rather than their ability

to convert acquired resources into offspring. Despite a and C mutants of equal effect size ($\delta_a =$

$\delta_C = \delta_f$) experiencing equivalent fixation probabilities for any given consumer and resource

density (eq. 6a), individuals carrying the rescue allele in a population experiencing reductions in

acquisition rates will benefit from greater resource availability earlier on during the process of

population decline (Fig. 2A), hastening its spread in the population (Fig. 2B).

Figure 2. (A) Resource abundance and (B) rescue allele frequency as a function of time
following an environmental change that reduces acquisition rates (orange) and conversion
efficiencies (green). Observed data was averaged from 3000 replicate simulations of the slow-
growing population with $a_0 = 9.5 \times 10^{-5}, C_0 = 0.0095, S_0 = 0.95, r_R = 0.475, K_R = 95{,}000,$ and
$N_0 = 2084,$ where $a_* = 3.5 \times 10^{-5}$ (orange) or $C_* = 0.0035$ (green). In both cases, $r_{N*} = -0.018.$
The grey dashed line in (A) represents the resource's carrying capacity, $K_R = 95{,}000.$
301
Whether rescue via *de novo* mutation is more favourable following declines in fecundity

or survivorship is variable with respect to the population's resident life-history strategy, and this

result is primarily driven by differences in the supply of rescue mutations. Our simulation results

suggest that rescue probabilities are approximately linear with respect to mutational supply

(Figs. 3A & D) because we are only considering rescue via *de novo* mutations. In simulations of

the slow-growing population, the affected ecological trait does not significantly influence time to

extinction, t_e (Fig. 3C), although it is worth noting a slight discrepancy between the traits.

Instead, decreases in survivorship result in a much greater total supply of rescue mutations than

decreases in fecundity for any given t_e (Fig. 3B). Population mutation rates are a function of

both total population size and birth rates. Thus, while all populations experience lower

populational mutation input following the environmental change due to smaller population size,

populations with reduced fecundity suffer additional reductions in mutation rates due to fewer

births.

In the fast-growing population, on the other hand, it is following reductions in resource

acquisition rates that the mutational supply, and therefore the probability of rescue, is

maximized (Fig. 3D & E), and this result corresponds to a slower time to extinction (Fig. 2F).

Interestingly, reductions in survivorship lead to a mutational supply almost indistinguishable, if

not lower, from reductions in conversion efficiency (Fig. 3E). This result can be explained by

considering the direct effect of survivorship on generation time, particularly since we model

fecundity to be constant with age. The average generation time is $\frac{1}{1-S}$, a function of survivorship.

Reduced survival accelerates individual turnover and reduces generation time, thus leading to

extinction in fewer time-steps than when fecundity is reduced, in the absence of evolution. This

effect is marginal in the slow-growing population (where $S_{0,slow} = 0.95$), but occurs in much

greater magnitude in the fast-growing population ($S_{0,fast} = 0.5$), where negative intrinsic growth

rates via decreases in survivorship are only possible when $S_{*,fast} < 0.0975$ (for $r_{N*} < 0$),

resulting in substantially reduced generation times and rapid individual turn over. The similar

results between the survivorship and conversion efficiency cases are thus primarily driven by

trade-offs in mutation supply per time-step and extinction time. These trade-offs also occur

when comparing acquisition rates to survivorship, although populations with lessened

acquisition rates additionally benefits from greater resource abundance at an earlier rate due to

lower per-capita resource consumption rates, leading to the greatest probability of rescue here.

Figure 3. Probability of rescue, mutational supply, and times to extinction for a slow-growing
population (A) – (C) and fast-growing population (D) – (F). The circles connected by lines
represent observed mutation supply and times to extinction from at least 1000 replicate
simulations, and the squares represent numerical estimates from our quasi-deterministic theory.
338
Although rescue via *de novo* mutation appears to be primarily driven by differences in

total mutational supply, regardless of the ecological trait, it is worth briefly describing the

mechanics of selection and fixation probabilities in our model, which indeed differ among

ecological traits. An interesting result of our analysis is that the fixation probabilities, ψ, of

rescue alleles at any time-point do not necessarily correspond to their selective advantage, s,

defined as the fitness of individuals carrying the rescue allele relative to the declining resident

allele (see Fig. 4 caption). Instead, the fixation probability of a rescue allele will be additionally

dependent on the rate of consumer population decline as well as resource abundance, which

are differentially influenced by the three ecological traits.

Figure 4. Selection coefficients, fixation probabilities, consumer and resource abundance over
time in the slow-growing population. (A) Selection coefficients were calculated using the formula
$s = \frac{W_{\text{mut}}}{W_{\text{res}}} - 1$, where W_{mut} is the fitness of the mutant type and W_{res} the fitness of the resident
type in a population in need of rescue ($W_{\text{res}} < 1$). Data was averaged from at least 1000
replicate simulations of the slow-growing population with $r_{N*} = -0.027$ from $a_* = 2.5 \times 10^{-5}$
(orange), $C_* = 0.0025$ (green), or $S_* = 0.8868$ (blue). (B) Fixation probabilities were estimated
using eqs. 6a and 6b, which perform well for these simulations (see Fig. 1A). (C–D) Mean
consumer and resource abundance over time.

Selection on mutant alleles that increase fecundity, either via increasing a or C, are

selectively favoured at low consumer density, whereas selection on increased survivorship is

maximized at high consumer density (Fig. 4A). However, fixation probabilities of rescue mutants

increase monotonically with declining consumer population sizes, albeit at different rates for

each ecological trait, ultimately converging onto a single maximum as consumer population size

declines towards extinction (Fig. 4B). This apparent contradiction between s and ψ may be

resolved simply by considering that, despite simulations being matched for r_{N*}, such that their

fixation probabilities once $N_t = 0$ are equal, they differ in their rates of decline and effects on

resource abundance at intermediate densities (Fig. 4C & D). For example, when resource

acquisition rates are reduced, the rate of consumer population decline is marginally slower, and

the rate of resource growth marginally faster than a survivorship mutant, such that, despite

having a lower selective advantage (*cf.* Fig. 4A blue & orange lines), a rescue allele that

increases acquisition rates will experience a higher fixation probability at intermediate to high

consumer population sizes. Conversely, despite decreases in survivorship leading to more rapid

declines in consumer density than decreases in conversion efficiency (*cf.* Fig. 4A blue & green

lines), the selective advantage of a survivorship mutant exceeds that of a conversion efficiency

mutant, such that it experiences higher fixation probabilities for intermediate consumer

population sizes (Fig. 4B).

Polygenic rescue

Our analysis has until now been restricted to the case of evolutionary rescue via the *de*

novo occurrence and fixation of a single beneficial allele in a haploid population, from which we

develop some analytical theory and derive general intuition for the mechanisms underlying

rescue. However, it is also possible, and indeed likely, that the genetic architecture underlying

consumer ecological traits is polygenic, with standing genetic variation in ecological trait values

prior to the environmental change. We explore this scenario by modelling a sexual, diploid

consumer population with polygenic phenotypes underlying ecological performance. Our

polygenic model slightly differs from those typically used in the classical quantitative genetics

literature in that the phenotype instead determines an individual's ecological trait value (a, C, or

S), which then influences fitness (recall that $W = aCR_t + S$) instead of the phenotype directly

influencing fitness (as in Gomulkiewicz & Holt 1995, Burger & Lynch 1995, etc.). Briefly, we

consider an arbitrary phenotypic trait (e.g. height, size) of the consumer underlying a single

ecological trait ϕ (where ϕ may refer to a, C, or S), holding all other traits constant. The

phenotypic trait value z of an individual is determined by the sum of the genetic contribution of

100 loci with additive phenotypic effects plus a normally distributed environmental effect with

mean 0 and variance 1. The ecological trait value of an individual with phenotypic trait value z is

described by a Gaussian performance function:

394
$$\phi(z) = \phi_{max} \exp\left[-\frac{(z - z_{max})^2}{2\omega^2}\right],$$
(7)

where ϕ_{max} refers to the maximum ecological trait value, z_{max} refers to the phenotypic value

which maximizes ϕ (when $z = z_{max}$, $\phi(z) = \phi_{max}$), and ω^2 describes the function's width.

Following a burn-in period where genetic variation is introduced via mutation-selection balance,

we model rescue by imposing an abrupt environmental change that modifies z_{max} of one

ecological trait to z_{max*} such that $r_{N*} < 0$ and the population will deterministically go extinct in

the absence of evolution. The population must now evolve toward its new phenotypic optimum

to increase ϕ, and therefore mean fitness, $\overline{W}$, above one. At the time of the environmental

change, the population experiences some additional reduction in fitness due to segregating

variation around its optimum (variance load), but also experiences a head-start in the adaptive

process following an environmental change due to some individuals carrying alleles that may

move their phenotype towards the new optimum.

We again simulate the case of a slow-growing population ($\bar{r}_N \approx 0.03$, where the

approximation is due to environmental variation in phenotype) with low fecundity relative to its

survival, and a fast-growing population with high fecundity relative to its survival ($\bar{r}_N \approx 0.28$),

although with slightly different parameters. We recover a qualitatively similar result to our one-

locus simulations, with rescue probabilities being contingent on both the affected ecological trait

and the resident life-history of the population (Fig. 5). Decreases in survivorship are heavily

favourable to rescue in slow-growing populations already exhibiting high survivorship, followed

by acquisition rates and conversion efficiencies, a result almost identical to the one-locus case

(*cf.* Fig. 1A & Fig. 5A). However, we observe some differences in the fast-growing case. Here,

decreases in survivorship substantially hinder rescue, but rescue probabilities following

decreases in acquisition rates or conversion efficiencies are similar.

Figure 5. Rescue probabilities via evolution of a polygenic trait as a function of intrinsic growth
rates and the affected ecological trait following an abrupt environmental change in (A) a slow-
growing population ($a_{max} = 10^{-4}, C_{max} = 0.01, S_{max} = 0.99, r_{R,max} = 0.5,$ and $K_{R,max} = 10^5$) and
(B) fast-growing population ($a_{max} = 10^{-4}, C_{max} = 0.093, S_{max} = 0.5, r_{R,max} = 0.5,$ and $K_{R,max} = 10^5$). The dots connected by lines represent observed data from at least 100 replicate
simulations, and error bars represent 95% confidence intervals. The evolving ecological trait
was determined by 100 loci evenly distributed among 10 chromosomes of map length 30cM.
Mutations were drawn at per base-pair rate 10^{-4} from a normal distribution with mean 0 and
variance 0.5.

In this scenario, *de novo* mutation plays a diminished role in the probability of rescue

because of the occurrence of segregating variation. Thus, it follows that the advantage of

greater mutational supply following decreased survivorship relative to decreased conversion

efficiency is diminished, but populations with reduced conversion efficiency still benefit from

slower times to extinction than populations with reduced survivorship (*cf*. Fig. 1B & 5B).

Additionally, phenotypic change is no longer contingent on the establishment of large-effect

beneficial mutations as in the one-locus case, but allele frequency change at many small effect

loci resulting in the evolution of phenotypic trait values toward the new optimum. Thus, under a

polygenic architecture, selective advantages experienced by favoured alleles play a relatively

greater role in rescue than in the one-locus case due to the diminished role of mutational input.

Selection

439   We now describe selection coefficients for mutant individuals relative to residents

440   immediately following the environmental change and at low density. Immediately after an abrupt

441   environmental change that would result in extinction of the resident type, the selection

442   coefficient, s, experienced by a mutant individual is

443
$$s = \frac{(1 - S_0)a_{\mathrm{mut}}C_{\mathrm{mut}} + a_0 C_0 S_{\mathrm{mut}}}{(1 - S_0)a_{\mathrm{res}}C_{\mathrm{res}} + a_0 C_0 S_{\mathrm{res}}} - 1, \tag{8}$$

where a_0, C_0, and S_0 are ecological trait values of the population prior to the environmental

change, a_{res}, C_{res}, and S_{res} are ecological trait values of the resident allele following the

environmental change, and a_{mut}, C_{mut}, and S_{mut} are ecological trait values of individuals carrying

a mutant allele (see *Appendix B* for a derivation). From eq. 8 it is visible through the numerator

that, at high density, the selective advantage of a fecundity mutant is proportional to $1 - S_0$, the

mortality rate of the resident population prior to the environmental change, whereas the

selective advantage of a survival mutant is proportional to $a_0 C_0$, the total number of offspring

produced per resource individual.

Consider the case where survivorship is reduced by an environmental change while

fecundity remains unaffected, and there is the occurrence of a mutant allele that increases

survivorship by factor δ_S. In this case, $a_{\mathrm{mut}}C_{\mathrm{mut}} = a_{\mathrm{res}}C_{\mathrm{res}} = a_0 C_0$, $S_{\mathrm{mut}} = S_{\mathrm{res}}(1 + \delta_S)$, and eq. 8

reduces to $s = \frac{\delta_S S_{\mathrm{res}}}{1 - S_0 + S_{\mathrm{res}}}$. Note that this formula does not depend on fecundity-related traits and

increases monotonically with S_0. If instead an environmental change reduces fecundity, the

mutant allele increases either acquisition rates or conversion efficiencies by factor δ_f, and

survivorship remains constant, eq. 8 reduces to $s = \frac{(1-S_0)a_{\text{res}}C_{\text{res}}\delta_f}{a_0 C_0 S_0 + (1-S_0)a_{\text{res}}C_{\text{res}}}$, a monotonically

decreasing function of S_0. Thus, at high density, the selective advantage of a survivorship

mutant will increase as a function of S_0, whereas the selective advantage of a fecundity mutant

will decrease as a function of S_0. Additionally, given that the strength of selection is dependent

on the evolving trait and resident life-history trait values, the equilibrium genetic variance

available to the consumer population, due to mutation-selection balance, immediately following

an abrupt environmental stressor will therefore also depend on the ecological trait under

selection.

At low density, the selection coefficient is:

467
$$s = \frac{(a_{\text{mut}}C_{\text{mut}} - a_{\text{res}}C_{\text{res}})K_R + S_{\text{mut}} - S_{\text{res}}}{a_{\text{res}}C_{\text{res}}K_R + S_{\text{res}}}. \tag{9}$$

If selection acts on fecundity, then $a_{\text{mut}}C_{\text{mut}} = a_{\text{res}}C_{\text{res}}(1 + \delta_f)$ and $S_{\text{mut}} = S_{\text{res}}$, eq. B4 reduces to

$s = \frac{\delta_f K_R}{a_{\text{res}}C_{\text{res}}K_R + S_{\text{res}}}$. If we instead consider selection on survivorship $a_{\text{mut}}C_{\text{mut}} = a_{\text{res}}C_{\text{res}}$ and $S_{\text{mut}} =$

$S_{\text{res}}(1 + \delta_S)$, eq. B4 reduces to $s = \frac{\delta_S}{a_{\text{res}}C_{\text{res}}K_R + S_{\text{res}}}$. Thus, the selective benefit of a fecundity

mutant will be scaled by the total resource availability at low density, such that its selective

benefit will be orders of magnitude greater than that of a survivorship mutant at low density if

their phenotypic effects are similar to one another ($\delta_f \approx \delta_S$). Additionally, at low density,

selection does not depend on the ecological trait values of the population prior to the

environmental change.

Discussion

In this article we have shown that the probability of evolutionary rescue of a consumer

species following an abrupt environmental change that reduces its intrinsic growth rate below

zero is dependent on both the affected ecological trait and the population's resident life-history

strategy. In a slow-growing population, characterized by low fecundity and high survivorship, our

analysis suggests that rescue is most favourable following reductions in survivorship, followed

by resource acquisition rates and then conversion efficiencies. In a fast-growing population,

characterized by high fecundity and low survivorship, it is instead following reductions in

resource acquisition rates that rescue is most probable, followed by survivorship and resource

conversion efficiency. These results are driven by the interplay between various factors,

including the effects ecological traits have on the mutational supply, resource competition (i.e.

density-dependence), selection, and times to extinction. We derive analytical formulae to predict

the fixation probabilities of mutant alleles, rescue probabilities at any given time-point, and times

to extinction, which work well when selection is weak and individual turnover rates are low.

There have been a number of previous studies on whether effects on either birth or

death rates influence evolutionary rescue (Osmond et al. 2017, Czuppon et al. 2023, Raatz &

Traulsen 2023, Draghi et al. 2024). In theoretical explorations of the evolution of antibiotic

resistance, it has been suggested that rescue is most likely following increases in death rates

versus birth rates due to reductions in competition experience by resistant strains (Czuppon et

al. 2023; Raatz & Traulsen 2023) as well as increased mutational supply. Draghi et al. (2024)

recover a similar result, finding that increased death rates lead to faster rates of adaptation, and

therefore rescue, due to reduced generation time. In this paper we show that this is not

necessarily the case; instead, rescue may be favoured following decreases in birth rates in

populations with high fecundity. This is because, while increases in death rates may result in

lower reductions in mutational supply per time-step and faster adaptation than reductions in

birth rates, they also accelerate times to extinction, perhaps leading to a lower overall

mutational supply. Moreover, the probability of rescue is affected by whether birth rates are

reduced via reductions in resource acquisition rates versus their efficiency of conversion to

offspring. In populations with already high fecundity and low survivorship, reductions in resource

acquisition rates–but not resource conversion efficiency–lead to greater resource availability to

rescue mutants, facilitating their establishment and therefore rescue to a greater degree than

when mortality rates are increased.

Another novel aspect of our model is the analysis of the interplay between genetic

architecture and the affected ecological trait on the probability of rescue in a two-species model.

When rescue occurs via *de novo* mutation at a single locus, ecological variables such as

resource availability and consumer vital rates substantially influence rescue through their effects

on the establishment probability of beneficial mutations and the mutational supply. Conversely,

when rescue occurs via both *de novo* mutation and standing genetic variation at many loci,

differences in rescue more closely reflect differences in selective pressures, leading to a

reduced benefit to increased mutation rates per time-step as in the case of increased mortality.

Given that selection is indeed dependent on resident ecological trait values, this also suggests

that the genetic variation available to selection for differing traits may indeed be dependent on

the population's resident life-history strategy, which may significantly influence rescue. For

instance, Yamamichi et al. (2019) have shown, in a model of non-overlapping generations with

a fluctuating environment, that increased survival may produce reservoirs of genetic variation

(e.g. harbored by older or dormant individuals), although much of this variation would be

deleterious due to those individuals phenotypically lagging behind fitter types. The analysis of

evolutionary rescue under fluctuating environments instead of an abrupt environmental change

may thus present an interesting extension to our work.

A central result of our simulations is that intrinsic growth rates alone are insufficient to

predict the probability of rescue, because negative intrinsic growth rates may be the result of

different ecological factors, each with a potentially different effect on factors such as the

mutational supply, time to extinction, and density-dependence. To a first approximation, intrinsic

growth rates may be a useful metric of population vulnerability, but our simulations suggest that

the probability of rescue for a given intrinsic growth rate following an abrupt environmental

change may be heavily dependent on the underlying ecological mechanism. However, it may be

possible to predict these differences if empiricists are able to measure fecundity and viability

directly.

One open question raised by our model may be what biological mechanisms may lead to

decreases in traits such as acquisition rates, conversion efficiencies, and survivorship rates, and

whether these have been observed. Regarding survivorship, there are various well-

characterized examples of evolutionary rescue following antibiotic treatments in bacteria,

herbicide treatment in weeds, and fungicide treatment for fungal diseases of economically

important crops (see Bell 2017). Regarding acquisition rates and conversion efficiencies, Wei &

Zhang (2019) show that quantitative trait loci (QTLs) in the yeast *S. cerevisiae* tend to show

concordant effects on r and K (r/K trade-ups) in lower quality environments, but negative

effects (r/K trade-offs) in higher-quality environments. They propose an explanatory model

characterized by trade-offs between ATP production rates and utilization efficiency in the

context of cell maintenance and division. Under low quality environments, r is low, and the

evolution of increased r increases resource use efficiency, thus increasing both r and K.

Conversely, in a high quality environment where r is already high, conversion efficiency

decreases with increasing r. Here, the population is instead dominated by competition for the

acquisition of resources. It is also well known that antibiotics may be distinguished into two

classes according to mode of selection: drugs that act to increase mortality rates (bacteriocidal)

versus decrease birth rates (bacteriostatic) (Czuppon et al. 2023). For instance, doxycycline, a

bacteriostatic agent, acts by inhibiting protein production, and it has been shown that the

evolution of doxycycline resistance in *E. coli* results in both increased intrinsic growth rates and

carrying capacities (Reding-Roman et al. 2017). It may thus be possible to characterize the

evolution of doxycycline resistance, and other chemically similar agents, in terms of the

evolution of increased resource conversion efficiency. In general, empirical studies of

evolutionary rescue often appear to focus on net rates of change rather than characterizing the

specific ecological effects of environmental stressors. We thus highly encourage empiricists to

pay greater attention to teasing out the effects of environmental stressors on birth and death

rates when feasible, as this will lead to richer insights into the factors determining persistence in

declining populations.

A major simplifying assumption of our theoretical analysis has been that ecological traits

are uncorrelated, and only one trait is influenced by an abrupt environmental change and may

subsequently evolve. This simple scenario is not intended to be a reflection of biological reality,

but is a useful starting to point to characterize the core dynamics of the system. Survival and

reproduction are indeed often correlated (e.g. Franco & Silvertown 2004), and evidence

suggests that these correlations are altered by environmental change (Iles et al. 2019). We also

do not consider a resource distribution, where competition for resource may lead to adaptive

divergence among competing types (Dieckmann & Doebeli 1999). This would present an

interesting extension to our model, as it is possible that adaptive divergence may relax

competition, thus aiding rescue, but may result in less resource availability for divergent types,

and therefore increased susceptibility to extinction over longer time-scales. Strong selective

pressures imposed by an abrupt environmental change are also likely to bias the evolution of

life-history traits to divergence between conspecifics, potentially promoting diversification

(Chaparro-Pedraza & Bank 2025).

We have here provided an improved theoretical framework to understand how

evolutionary rescue is influenced by the complex interplay between evolution and demography.

The use of mechanistic models that provide increased biological realism is an important step

forward in determining our treatment of populations at risk of extinction (Urban et al. 2024), an

increasingly relevant problem given the rate of climate change (Berg 2010). The work presented

here extends our understanding of the conditions under which we expect populations to adapt to

environmental conditions that would otherwise lead to extinction.

Appendix A. Derivation of $\psi(N_t)$

Here we outline the derivation of the approximate fixation probability of a rescue mutant as a function of the current population size, $\psi(N_t)$, following Otto & Whitlock (1997). We evaluate the sum in eq. 5 from the main text to get:

$$\psi_t = 1 - e^{-a(1+\delta_a)C(1+\delta_C)R_t\psi_{t+1}}(1 - S(1+\delta_S)\psi_{t+1}), \tag{A1}$$

Assuming $a(1+\delta_a)C(1+\delta_C)R_t$ and ψ_{t+1} are small, we perform a first-order Taylor series expansion of the exponential around 0 to get:

$$\psi_t \approx 1 - (1 - a(1+\delta_a)C(1+\delta_C)R_t\psi_{t+1})(1 - S(1+\delta_S)\psi_{t+1}), \tag{A2}$$

Since birth rates are density-dependent, and we are considering a scenario with variable demography, this is a time-inhomogeneous branching process (i.e. $\psi_t \neq \psi_{t+1}$). In a declining consumer population, for example, surviving individuals will benefit from greater resource availability in the future, leading to increased fecundity in future time-steps. Assuming the difference in fixation probabilities between any given time-step is small, we use a continuous-time approximation for $\Delta\psi = \psi_{t+\Delta t} - \psi_t$. Substituting in the right hand side of eq. A2 in place of ψ_t and then taking the limit of $\frac{\Delta\psi}{\Delta t}$ as $\Delta t \to 0$, we get $\frac{d\psi}{dt}$. Dropping subscripts,

$$\frac{d\psi}{dt} = \psi + (1 - a(1+\delta_a)C(1+\delta_C)R_t\psi)(1 - S(1+\delta_S)\psi) - 1. \tag{A3}$$

We can use the chain rule to re-write ψ as a function of population size N, $\frac{d\psi}{dN} = \frac{d\psi}{dt}\frac{dt}{dN}$. We derive $\frac{dt}{dN}$ by assuming a separation of time-scales between consumer and resource growth. The logic behind this assumption is that if resource growth rates are much greater than consumer growth rates, then the resource will remain at a "quasi-equilibrium" state with respect to the consumer density at any time-step. Under these assumptions, consumer population dynamics are well represented by the logistic growth formula, $\frac{dN}{dt} = N\left(r_N\left(1 - \frac{N}{K_N}\right)\right)$ (Travis et al. 2023). We can solve for r_N by considering that the intrinsic growth rate of the consumer is simply its growth rate

607   at low density, when the resource is at its carrying capacity. Specifically, when $N_t = 0$, $R_t = K_R$

608   and $r_N = aCK_R + S - 1$. We find K_N by solving eq. 2 in the main text for $\hat{N}$, the consumer

609   population's demographic equilibrium. After some algebraic rearrangement, we get $K_N = \frac{r_R r_N}{a^2 C K_R}$.

610   Plugging r_N and K_N into the logistic growth formula and simplifying, we get:

$$611 \qquad \frac{dN}{dt} = N\left(aCK_R\left(1 - \frac{aN}{r_R}\right) + S - 1\right). \qquad (A4)$$

612   We can then solve for $\frac{dt}{dN}$ by taking the inverse of eq. A4:

$$613 \qquad \frac{dt}{dN} = \left(\frac{dN}{dt}\right)^{-1} = \frac{1}{N\left(aCK_R\left(1 - \frac{aN}{r_R}\right) + S - 1\right)}. \qquad (A5)$$

614   Note that $R_t = K_R\left(1 - \frac{aN}{r_R}\right)$, which can be found by solving eq. 2 in the main text for $R_t = R_{t+1} =$

615     $\hat{R}$. Since most individuals at any given time-point will be of the resident type, population

616   dynamics will be a function of resident ecological trait values in a declining population.

617   Combining eqs. A3 and A5 using the chain rule and re-writing R as a function of N,

$$618 \qquad \frac{d\psi}{dN} = \frac{\psi + (1 - a(1 + \delta_a)C(1 + \delta_C)K_R\left(1 - \frac{aN}{r_R}\right)\psi)(1 - S(1 + \delta_S)\psi) - 1}{N\left(aCK_R\left(1 - \frac{aN}{r_R}\right) + S - 1\right)}. \qquad (A5)$$

619   Eq. A5 can be solved directly and simplified, depending on which ecological trait evolves, to get

620   eqs. 6a and 6b.

Appendix B. Selection at high and low density

622   Here we describe selection coefficients for a mutant allele that increases ecological trait values

623   immediately following an abrupt environmental change that reduces fitness such that the

624   consumer population will deterministically go extinct, and at low density, when the consumer

625   population is nearly extinct. The fitness of the ancestral genotype after the environmental

626   change is given by

$$627 \qquad W_{res,t} = a_{res}C_{res}R_t + S_{res}, \qquad (B1)$$

where a_{res}, C_{res}, and S_{res} refer to ecological trait values as changed by the new environment and

R_t is the size of the resource population at time t. In order for the population to require

evolutionary rescue, this fitness value must be less than one when the population is at low

density; i.e., $a_{\text{res}}C_{\text{res}}K_R + S_{\text{res}} < 1$.

The invasion fitness of a rare mutant allele that occurs at time-step t is simply equal to

its geometric growth rate,

$$W_{\text{mut},t} = a_{\text{mut}}C_{\text{mut}}R_t + S_{\text{mut}}. \tag{B2}$$

635 Therefore, the selection coefficient for the mutant allele at time t is given by

$$s = \frac{a_{\text{mut}}C_{\text{mut}}R_t + S_{\text{mut}}}{a_{res}C_{res}R_t + S_{res}} - 1. \tag{B3}$$

637 Immediately after the environmental change, resource abundance, R_0, is still at equilibrium with

638 respect to the ecological trait values prior to the environmental change, a_0, C_0, and S_0, where

639 $R_0 = \frac{1-S_0}{a_0 C_0}$ (which is derived from rearrangement of eq. 2 in the main text). Plugging this into our

640 formula for s and after some rearrangement, we get

$$s = \frac{(1 - S_0)a_{\text{mut}}C_{\text{mut}} + a_0 C_0 S_{\text{mut}}}{(1 - S_0)a_{\text{res}}C_{\text{res}} + a_0 C_0 S_{\text{res}}} - 1, \tag{B4}$$

642 At low density, on the other hand, $R_t = K_R$, from which eq. B2 simplifies to

$$s = \frac{(a_{\text{mut}}C_{\text{mut}} - a_{\text{res}}C_{\text{res}})K_R + S_{\text{mut}} - S_{\text{res}}}{a_{\text{res}}C_{\text{res}}K_R + S_{\text{res}}} = \frac{r_{N,\text{mut}} - r_{N,\text{res}}}{r_{N,\text{res}}}. \tag{B5}$$

644 Here, the selective difference between the mutant and resident fecundity types is scaled by the

645 carrying capacity of the resource.

646 References:

647
648   1. Abrams, P. A. Will small population sizes warn us of impending extinctions? *Am. Nat.*
649   **160**, 293–305 (2002).
650   2. Abrams, P. A. How does the evolution of universal ecological traits affect population
651   size? Lessons from simple models. *Am. Nat.* **193**, 814–829 (2019).

3. Agrawal & Whitlock. Mutation load: the fitness of individuals in populations where deleterious alleles are abundant. *Annu. Rev. Ecol. Syst.* **43,** 115–135 (2012).

4. Bell, G. Evolutionary Rescue. *Annu. Rev. Ecol. Evol. Syst.* **48**, 605–627 (2017).

5. Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W. & Courchamp, F. Impacts of climate change on the future of biodiversity. *Ecology Letters* **15**, 365–377 (2012).

6. Berg, M. P. *et al.* Adapt or disperse: understanding species persistence in a changing world. *Glob. Chang. Biol.* **16**, 587–598 (2010).

7. Bisschop, K., Alzate, A., Bonte, D. & Etienne, R. S. The demographic consequences of adaptation: Evidence from experimental evolution. *Am. Nat.* **199**, 729–742 (2022).

8. Boulding, E. G. & Hay, T. Genetic and demographic parameters determining population persistence after a discrete change in the environment. *Heredity* **86**, 313–324 (2001).

9. Burger, R. & Lynch, M. Evolution and Extinction in a Changing Environment: A Quantitative-Genetic Analysis. *Evolution* vol. 49 151 (1995).

10. Carlson, S. M., Cunningham, C. J. & Westley, P. A. H. Evolutionary rescue in a changing world. *Trends Ecol. Evol.* **29**, 521–530 (2014).

11. Chaparro-Pedraza, P. C. & Bank, C. Evolving life-history traits promote biodiversity via eco-evolutionary feedback mechanisms. *PLoS Biol.* **23**, e3003492 (2025).

12. Chevin, L.-M. & Lande, R. When do adaptive plasticity and genetic evolution prevent extinction of a density-regulated population? *Evolution* **64**, 1143–1150 (2010).

13. Cortés, E. Perspectives on the intrinsic rate of population growth. *Methods Ecol. Evol.* **7**, 1136–1145 (2016).

14. Cortez, M. H. & Yamamichi, M. How (co)evolution alters predator responses to increased mortality: extinction thresholds and hydra effects. *Ecology* **100**, e02789 (2019).

15. Czuppon, P., Day, T., Débarre, F. & Blanquart, F. A stochastic analysis of the interplay between antibiotic dose, mode of action, and bacterial competition in the evolution of antibiotic resistance. *PLoS Comput. Biol.* **19**, e1011364 (2023).

16. Dieckmann, U. & Doebeli, M. On the origin of species by sympatric speciation. *Nature* **400**, 354–357 (1999).

17. Draghi, J. A., McGlothlin, J. W. & Kindsvater, H. K. Demographic feedbacks during evolutionary rescue can slow or speed adaptive evolution. *Proc. Biol. Sci.* **291**, 20231553 (2024).

18. Fisher, R. A. On the dominance ratio. *Proc Roy Soc Edinb* **42**, 321–341 (1922).

19. Fisher, R. A. *The Genetical Theory of Natural Selection*. Clarendon Press (1930).

20. Franco, M. & Silvertown, J. A comparative demography of plants based upon elasticities of vital rates. *Ecology* **85**, 531–538 (2004).

21. Gomulkiewicz, R. & Holt, R. D. When does evolution by natural selection prevent extinction? *Evolution* **49**, 201–207 (1995).

22. Haldane, J. B. S. A mathematical theory of natural and artificial selection. V. Selection and mutation. *Proc Camb Phil Soc* **23**, 838–844 (1927).

23. Haller, B. C. & Messer, P. W. SLiM 4: Multispecies eco-evolutionary modeling. *Am. Nat.* (2022) doi:10.1086/723601.

24. Hermann, R. J. & Becks, L. Change in prey genotype frequency rescues predator from extinction. *R. Soc. Open Sci.* **9**, 220211 (2022).

25. Iles, D. T., Rockwell, R. F. & Koons, D. N. Shifting vital rate correlations alter predicted population responses to increasingly variable environments. *Am. Nat.* **193**, E57–E64 (2019).

26. Jones, A. G. A theoretical quantitative genetic study of negative ecological interactions and extinction times in changing environments. *BMC Evol. Biol.* **8**, 119 (2008).

27. Lachapelle, J. & Bell, G. Evolutionary rescue of sexual and asexual populations in a deteriorating environment. *Evolution* **66**, 3508–3518 (2012).

28. Lande, R. & Shannon, S. The Role of Genetic Variation in Adaptation and Population Persistence in a Changing Environment. *Evolution* vol. 50 434 (1996).

29. MacArthur, R. Species packing and competitive equilibrium for many species. *Theor. Popul. Biol.* **1**, 1–11 (1970).

30. MacArthur R. Geographical Ecology. New York: Harper and Row (1972).

31. Mallet, J. The struggle for existence. How the notion of carrying capacity, K, obscures the links between demography, Darwinian evolution and speciation. *Evol. Ecol. Res.* (2012).

32. Marshall, D. J., Cameron, H. E. & Loreau, M. Relationships between intrinsic population growth rate, carrying capacity and metabolism in microbial populations. *ISME J.* **17**, 2140–2143 (2023).

33. Nordstrom, S. W., Hufbauer, R. A., Olazcuaga, L., Durkee, L. F. & Melbourne, B. A. How density dependence, genetic erosion and the extinction vortex impact evolutionary rescue. *Proc. Biol. Sci.* **290**, 20231228 (2023).

34. Orr, H. A. & Unckless, R. L. Population extinction and the genetics of adaptation. *Am. Nat.* **172**, 160–169 (2008).

35. Orr, H. A., Allen Orr, H. & Unckless, R. L. The Population Genetics of Evolutionary Rescue. *PLoS Genetics* vol. 10 e1004551 (2014).

36. Osmond, M. M. & de Mazancourt, C. How competition affects evolutionary rescue. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368**, 20120085 (2013).

37. Osmond, M. M., Otto, S. P. & Martin, G. Genetic Paths to Evolutionary Rescue and the Distribution of Fitness Effects Along Them. *Genetics* **214**, 493–510 (2020).

38. Osmond, M. M., Otto, S. P. & Klausmeier, C. A. When Predators Help Prey Adapt and Persist in a Changing Environment. *Am. Nat.* **190**, 83–98 (2017).

39. Pianka, E. R. On r- and K-Selection. *Am. Nat.* **104**, 592–597 (1970).

40. Raatz, M. & Traulsen, A. Promoting extinction or minimizing growth? The impact of treatment on trait trajectories in evolving populations. *Evolution* **77**, 1408–1421 (2023).

41. Reding-Roman, C. *et al.* The unconstrained evolution of fast and efficient antibiotic-resistant bacterial genomes. *Nat. Ecol. Evol.* **1**, 50 (2017).

42. R Core Team (2024). _R: A Language and Environment for Statistical Computing_. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

43. Sakarchi, J. & Germain, R. M. MacArthur's consumer-resource model: a 'Rosetta Stone' for competitive interactions. *Am. Nat.* (2024) doi:10.1086/733516.

44. Sheth, S. N. & Angert, A. L. Demographic compensation does not rescue populations at a trailing range edge. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 2413–2418 (2018).

45. Travis, J., Bassar, R. D., Coulson, T., Reznick, D. & Walsh, M. Density-dependent selection. *Annu. Rev. Ecol. Evol. Syst.* **54**, 85–105 (2023).

46. Uecker, H. Evolutionary rescue in randomly mating, selfing, and clonal populations. *Evolution* **71**, 845–858 (2017).

47. Uecker, H. & Hermisson, J. The Role of Recombination in Evolutionary Rescue. *Genetics* **202**, 721–732 (2016).

48. Uecker, H., Otto, S. P. & Hermisson, J. Evolutionary rescue in structured populations. *Am. Nat.* **183**, E17-35 (2014).

49. Urban, M. C. *et al.* When and how can we predict adaptive responses to climate change? *Evol. Lett.* **8**, 172–187 (2024).

50. Van Den Elzen, C. L., Kleynhans, E. J. & Otto, S. P. Asymmetric competition impacts evolutionary rescue in a changing environment. *Proc. Biol. Sci.* **284**, (2017).

51. van Velzen, E. High importance of indirect evolutionary rescue in a small food web. *Ecol. Lett.* **26**, 2110–2121 (2023).

52. Villellas, J., Doak, D. F., García, M. B. & Morris, W. F. Demographic compensation among populations: what is it, how does it arise and what are its implications? *Ecol. Lett.* **18**, 1139–1152 (2015).

53. Wei, X. & Zhang, J. Environment-dependent pleiotropic effects of mutations on the maximum growth rate r and carrying capacity K of population growth. *PLoS Biol.* **17**, e3000121 (2019).

54. Wright, S. Evolution in mendelian populations. *Genetics* **16**, 97–159 (1931).

55. Xu, K. Population Rescue through an Increase in the Selfing Rate under Pollen Limitation: Plasticity versus Evolution. *Am. Nat.* **202**, 337–350 (2023).

56. Yamamichi, M. & Miner, B. E. Indirect evolutionary rescue: prey adapts, predator avoids extinction. *Evol. Appl.* **8**, 787–795 (2015).

57. Yamamichi, M., Hairston, N. G., Jr, Rees, M. & Ellner, S. P. Rapid evolution with generation overlap: the double-edged effect of dormancy. *Theor. Ecol.* **12**, 179–195 (2019).