Main Manuscript for

Temporal orchestration of transcriptional and epigenomic programming
underlying maternal embryonic diapause in a cricket model

Kosuke Kataoka[1,2*†], Yuta Shimizu[3†], Ryuto Sanno[4], Yuichi Koshiishi[5], Ken Naito[6], Kei Yura[7,8],
Toru Asahi[2,4,8], Shin G. Goto[3*]

1. Division of Biotechnology and Life Science, Institute of Engineering, Tokyo University of
Agriculture and Technology, 2-24-16 Naka-cho, Koganei-shi, Tokyo 184-8588, Japan.
2. Comprehensive Research Organization, Waseda University, 2-2 Wakamatsu-cho, Shinjuku-
ku, Tokyo 162-8480, Japan.
3. Graduate School of Science, Osaka Metropolitan University, 3-3-138 Sugimoto, Sumiyoshi-
ku, Osaka 558-8585, Japan.
4. Graduate School of Advanced Science and Engineering, Waseda University, 2-2
Wakamatsu-cho, Shinjuku-ku, Tokyo 162-8480, Japan.
5. NODAI Genome Research Center, Tokyo University of Agriculture, 1-1-1 Sakuragaoka,
Setagaya-ku, Tokyo 156-8502, Japan.
6. Research Center of Genetic Resources, National Agriculture and Food Research
Organization, 2-1-2 Kannondai, Tsukuba-shi, Ibaraki 305-8602, Japan.
7. Graduate School of Humanities and Sciences, Ochanomizu University, 2-1-1 Otsuka,
Bunkyo-ku, Tokyo 112-8610, Japan.
8. Department of Life Science and Medical Bioscience, Waseda University, 513 Waseda
Tsurumaki-cho, Shinjuku-ku, Tokyo 162-0041, Japan.

† These authors equally contributed to this work.

* Kosuke Kataoka: kataokak@go.tuat.ac.jp
* Shin G. Goto: shingoto@omu.ac.jp

Author Contributions: K.K., Y. S., and S.G.G. designed research; K.K., Y.S., R.S., Y.K., K.N.,
K.Y., and T.A. performed research; K.K. analyzed data; and K.K., Y.S., and S.G.G. wrote the
paper.

Competing Interest Statement: The authors have no competing interests to disclose.

Abstract

Maternal perception of environmental conditions can direct offspring developmental trajectories, providing adaptive flexibility across taxa. In the band-legged ground cricket *Dianemobius nigrofasciatus*, maternal exposure to short days induces embryonic diapause at the cellular blastoderm stage in offspring. Here, we investigate molecular mechanisms underlying this transgenerational adaptation through genome assembly (1.45 Gbp) and time-series transcriptomic analyses of diapause and non-diapause eggs from 12 to 72 hours post-oviposition. Despite morphological similarity, diapause-destined eggs show early upregulation of ATP-dependent chromatin remodeling genes at 24 hours. ATAC-seq reveals reduced chromatin accessibility at neural and cell cycle-related genes. Time-series clustering identifies precocious shifts in RNA processing machinery (peaking at 24 versus 40 hours in non-diapause eggs), followed by metabolic regulation toward amino acid catabolism and gluconeogenesis sustaining long-term survival during developmental arrest. Our findings reveal diapause as actively coordinated molecular programming involving epigenetic, transcriptional, and metabolic remodeling, providing insights into transgenerational environmental adaptation.

Main Text

Introduction

Across the tree of life, organisms have evolved remarkable strategies to survive temporarily hostile environments through dormancy, a physiological state characterized by dramatically reduced metabolic activity that enhances survival during adverse conditions [1,2]. Among the most widespread environmental challenges is seasonality, which poses significant physiological demands on organisms inhabiting temperate regions. To survive periods unsuitable for growth and reproduction, diverse taxa have independently evolved dormancy strategies: hibernation and torpor in mammals and birds, as well as embryonic developmental arrest in vertebrates and developmental pauses in numerous invertebrates, notably insects [3].

Among these strategies, diapause emerges as a programmed form of dormancy employed by many insect species in anticipation of seasonal stressors. Diapause can occur at any developmental stage—egg, larva, pupa, or adult—depending on the species-specific life cycle strategy [4]. A number of common physiological events are associated with diapause, including metabolic suppression and developmental arrest [5], enhanced resistance to cold and desiccation [6,7], and accumulation of energy reserves [8].

In many species, individuals enter diapause upon receiving a token stimulus. However, a particularly intriguing mechanism involves maternal influence, whereby the mother receives environmental signals and transfers this information to her progeny. This transgenerational effect is referred to as the maternal effect [9]. Maternal diapause induction is particularly valuable for embryonic diapause, a situation in which an individual has limited access to or time to independently evaluate seasonal conditions. This transgenerational adaptation has been well-documented across diverse taxa, including mites [10], crickets [11–13], rice leaf bug [14], flies [15], mosquitoes [16], and parasitic wasps [17,18].

Despite the growing interest in maternal diapause induction, the underlying molecular mechanisms remain poorly understood. Recent omics studies have highlighted the importance of alterations in endocrine signaling, cell proliferation, metabolism, and energy production in

82   maternal diapause induction [19,20]; however, these investigations have not comprehensively
83   addressed how maternal influences impact specific molecular pathways initiating diapause, due
84   to limited sampling periods. A comprehensive understanding of the molecular processes through
85   which maternal environmental signals redirect the developmental trajectory towards diapause
86   requires investigation of multiple time points throughout early development.
87
Dianemobius nigrofasciatus, the band-legged ground cricket, serves as a prominent model for
investigating the maternal effect on embryonic diapause (Fig. 1a, b) [13,21]. Adult females reared
under long-day conditions produce eggs that develop directly into nymphs without undergoing
diapause (hereafter called long-day eggs), whereas those exposed to short-day conditions lay
diapause-destined eggs (hereafter called short-day eggs) (Fig. 1b) [12,13]. A previous morphological
study [22] established that long- and short-day eggs remain visually indistinguishable until 56 hours
post-oviposition (hpo). During the first 40 hpo, energids (early embryonic cells) proliferate and
migrate to the egg surface in both egg types. Subsequently, between 40 and 56 hpo, both types
of eggs undergo cellular blastoderm formation. However, by 72 hpo, a clear morphological
divergence emerges: long-day eggs proceed to form a germ band at the posterior pole, while
short-day eggs arrest at the cellular blastoderm stage without germ band formation. Although the
involvement of suppressed cell cycle regulators, small silencing RNAs, and segment patterning
genes has been suggested [23], the precise molecular mechanisms underlying maternal diapause
induction remain unknown.
102
In this study, we assembled a high-quality genome of *D. nigrofasciatus* and conducted a
comprehensive time-series transcriptomic analyses of long- and short-day eggs to elucidate the
molecular events occurring before and after the onset of embryonic diapause. Based on the
morphological milestones, we strategically selected timepoints that would capture molecular
changes before (12 and 24 hpo), during (40 and 56 hpo), and after (72 hpo) the critical
developmental decision point where diapause and non-diapause trajectories diverge. Our findings
suggest that this process is orchestrated not by a straightforward developmental switch, but
rather by a sophisticated temporal orchestration of molecular programs.
111

Results
Chromosome-scale genome assembly and gene annotation of *D. nigrofasciatus*

To investigate the molecular mechanisms of maternal diapause induction, we generated a
chromosome-scale genome assembly of *D. nigrofasciatus* using short-read, long-read, and Omni-
C sequencing technologies. Flow cytometry revealed genome size differences between sexes,
with males (1496.9 Mbp) having smaller genomes than females (1612.6 Mbp) (Fig. 1c;
Supplementary Table 1), consistent with the XO/XX sex determination system in crickets [24]. *K*-
mer analysis estimated a haploid genome size of 1.24 Gbp, which is smaller than the flow
cytometry estimates, likely due to underrepresentation of repetitive sequences in short-read data
(Supplementary Fig. 1; Supplementary Table 2).
122
The hybrid assembly strategy integrated 100.84× Illumina, 19.87× Oxford Nanopore, and 78.82×
Omni-C coverage (Supplementary Table 3), resulting in a 1.45 Gbp assembly across 482
scaffolds with a scaffold N50 of 154 Mbp and minimal contamination (Supplementary Fig. 2).
87.1% of sequences were anchored into 10 linkage groups (Fig. 1d; Supplementary Fig. 3;
Supplementary Tables 4, 5). Coverage analysis of male genomic sequences revealed that LG8
and LG9 constitute the X chromosome (Supplementary Fig. 4; Supplementary Table 6).
129

Assembly quality assessment using BUSCO (Benchmarking Universal Single-Copy Orthologs) [25]
demonstrated high completeness, with 90.0% complete single-copy and 6.2% duplicated
orthologs for the Arthropoda dataset (n=1,013) (Fig. 1e; Supplementary Table 4). When
evaluated against the more specific Insecta dataset (n=1,367), the assembly showed even higher
completeness with 93.2% single-copy and 3.0% duplicated orthologs (Fig. 1e; Supplementary
Table 4). Gene annotation using BRAKER3 [26–28], StringTie2 [29], and GeMoMa [30] yielded 20,812
protein-coding genes (Supplementary Table 7). Phylogenetic analysis of 334 single-copy
orthologs placed *D. nigrofasciatus* within Trigonidiidae, sister to Gryllidae (Supplementary Fig.
5),consistent with previous studies [31–33].
139
140   This high-quality genome assembly and gene annotation provide crucial resources for elucidating
141   the mechanisms of transgenerational diapause programming.
142

Gradual molecular divergence in maternal diapause induction
To elucidate the molecular mechanisms underlying maternal diapause induction, we performed
comprehensive time-series RNA-seq analysis of both long- and short-day eggs at five
developmental timepoints: 12, 24, 40, 56, and 72 hpo (Fig. 2a). Principal component analysis
(PCA) of the transcriptome data revealed intriguing patterns in the temporal dynamics of
developmental divergence (Fig. 2b). Given that maternal diapause induction presumably involves
signaling from mother to offspring, we initially expected substantial transcriptomic differences
even at the earliest timepoint. However, molecular differences between long- and short-day eggs
were minimal at 12-24 hpo, despite both egg types having been exposed to different maternal
environments. The transcriptomic distance between the two egg types then expanded
progressively from 24 to 72 hpo, with the most dramatic separation occurring between 56 and 72
hpo, coinciding with morphological divergence.
155

Differential expression analysis substantiated these observations, revealing a progressive
increase in the number of differentially expressed genes (DEGs) between long- and short-day
eggs over time ($|\log_2$ fold change$| > 1$, adjusted *p*-value < 0.05) (Fig. 2c, left panel;
Supplementary Table 8). Remarkably, at 12 hpo, merely 31 genes showed differential expression
(16 upregulated, 15 downregulated in short-day eggs). This number increased modestly to 120
DEGs at 24 hpo, then expanded more substantially to 371 DEGs at 40 hpo, 486 DEGs at 56 hpo,
and culminated in 1,994 DEGs at 72 hpo (1,078 upregulated, 916 downregulated in short-day
eggs).
164

Analysis of temporal gene expression changes between consecutive timepoints revealed three
distinct developmental phases (Fig. 2c, right panel; Supplementary Table 9). The first major
transcriptional transition (24-40 hpo) coincided with energid proliferation and migration, showing
3,316 DEGs in long-day eggs and 3,104 DEGs in short-day eggs. Transcriptional activity then
decreased during cellular blastoderm formation (40-56 hpo), with 858 and 722 DEGs in long- and
short-day eggs, respectively. The final phase (56-72 hpo) showed divergent patterns, particularly
in long-day eggs (2,922 DEGs) compared to short-day eggs (2,031 DEGs), reflecting the initiation
of germ band formation in long-day eggs while short-day eggs undergo molecular changes
associated with diapause [22].
174

Temporal coordination and divergence of gene expression patterns reveal
diapause preparation dynamics

To comprehensively characterize these temporal dynamics and identify genes with similar
expression patterns across developmental time, we performed fuzzy c-means clustering using
Mfuzz [34] on the time-series transcriptome data. This soft clustering assigns genes to multiple
clusters with varying membership degrees. This feature makes it particularly suitable for capturing
the continuous nature of developmental processes. Following independent clustering of each
dataset (long- and short-day eggs), we identified cluster pairs between the two egg types that
showed significant gene overlap using upper-tail hypergeometric tests (Fig. 3a; Supplementary
Fig. 6, 7). These overlapping cluster pairs were further classified based on their temporal
expression patterns: "homochronic clusters" displayed synchronized expression dynamics
between egg types, while "heterochronic clusters" exhibited temporal shifts in peak expression
timing, revealing potential differences in developmental timing between the two trajectories
(Supplementary Fig. 8; Supplementary Table 10).
189
To understand which developmental processes remain conserved, we first examined
homochronic gene clusters that maintained synchronized expression between egg types. The
majority of homochronic cluster pairs exhibited peak expression during the early developmental
periods (12-24 hpo), suggesting that fundamental embryonic processes are largely conserved
before the diapause trajectory diverges (Supplementary Fig. 9; Supplementary Table 11).
Analysis of these early-peaking homochronic clusters revealed common activation of essential
cellular machinery: genes encoding components of clathrin-coated vesicles and membrane
trafficking systems showed high initial expression followed by continuous decline, likely reflecting
the early embryo's use of maternally supplied membrane and vesicle pools required for
cellularization, together with maternally provided trafficking proteins and maternally deposited
mRNAs that are translated during early development (Fig. 3b) [35–37]. Both egg types also showed
coordinated activation of autophagy-related genes, peptidases, and catabolic processes at 12-24
hpo, suggesting a conserved program for recycling maternal components during early
embryogenesis (Fig. 3b) [37–40].
204
In contrast, heterochronic clusters unveiled critical timing differences in key molecular pathways
(Supplementary Fig. 10; Supplementary Table 12). Notably, genes involved in RNA processing
and spliceosome function exhibited marked temporal advancement in short-day eggs (Fig. 3c).
While these genes peaked at 40 hpo in long-day eggs, corresponding to the onset of cellular
blastoderm formation, they reached maximum expression already at 24 hpo in short-day eggs.
This precocious activation of RNA processing machinery coincides with the upregulation of
chromatin remodeling factors, as shown in the subsequent epigenetic analyses (Fig. 4a, b),
suggesting a coordinated early establishment of the molecular foundation required for
developmental arrest.
214
Early epigenetic remodeling drives diapause preparation
Differentially expressed genes unveiled the molecular mechanisms initiating diapause
(Supplementary Table 13). At 12 hpo, the sole significantly enriched pathway was
glycolysis/gluconeogenesis, showing downregulation in short-day eggs (Fig. 4a). However, by 24
hpo, this initial metabolic shift had expanded into comprehensive molecular programming, with
multiple pathways emerging as significantly enriched between egg types. Most strikingly, ATP-
dependent chromatin remodeling genes showed specific upregulation in short-day eggs at this
timepoint (Fig. 4a). Detailed examination revealed increased expression of core subunits of the
Brahma (SWI/SNF) complex, including *Bap111*, *pont*, and *Bap60* (Fig. 4b). The Brahma complex
is a critical and conserved regulator of chromatin accessibility and gene expression [41–43],
suggesting that epigenetic remodeling may serve as a molecular switch for diapause induction.

To test this hypothesis, we performed ATAC-seq analysis on 24 hpo eggs to assess genome-
wide chromatin accessibility. Short-day eggs exhibited reduced chromatin accessibility
specifically near genes associated with neural development and cell cycle regulation
(Supplementary Fig. 11; Supplementary Table 14, 15). Notable examples include *Kr-h1*, encoding
a juvenile hormone-responsive transcription factor essential for developmental progression [44],
and *crol*, an ecdysone-inducible transcription factor that recruits Polycomb group proteins [45,46].
Additionally, chromatin closure was observed at *Tlk* (Tousled-like kinase), which triggers S-phase
entry and histone assembly [47], and *Ssrp*, a FACT complex subunit required for transcription
initiation [48]. Neural signaling genes also showed reduced accessibility, including synaptic
components such as *Tollo* (Toll-6) and *nAChRα1*. This selective chromatin closure likely
contributes to the developmental arrest characteristic of diapause. The timing of these chromatin
changes before morphological divergence suggests that epigenetic remodeling is not a
consequence but rather a driver of diapause preparation.

Metabolic regulation and developmental signaling orchestrate diapause
preparation
Following the early establishment of epigenetic and transcriptional regulatory programs, a
comprehensive metabolic and developmental restructuring emerged as the next major phase of
diapause preparation. While initial changes were subtle, a dramatic transformation unfolded from
40 hpo onwards, intensifying through the morphological divergence point.

The metabolic shift was characterized by coordinated activation of multiple amino acid
degradation pathways. By 56-72 hpo, valine, leucine, and isoleucine degradation, tryptophan
metabolism, lysine degradation, and beta-alanine metabolism were significantly upregulated in
short-day eggs (Fig. 4a). This was accompanied by increased expression of key gluconeogenic
enzymes, with phosphoenolpyruvate carboxykinase 1 (*Pepck1*) showing marked upregulation
from 40 hpo and pyruvate carboxylase (*PCB*) increasing progressively from 56 hpo onwards (Fig.
4c). Genes encoding malate-aspartate shuttle components (*Got2* and *Mdh1/2*) also showed
elevated expression in short-day eggs at 56-72 hpo (Supplementary Fig. 12), suggesting
enhanced mitochondrial-cytoplasmic metabolite exchange. These enzymes enable the
conversion of amino acid-derived carbon skeletons into glucose [8,49], suggesting a metabolic
strategy to maintain energy homeostasis during impending developmental arrest.

Complementing this metabolic regulation, protein homeostasis pathways showed coordinated
adjustment. From 56 hpo onwards, coinciding with peak metabolic restructuring, proteasome,
protein processing in the endoplasmic reticulum, and spliceosome pathways were significantly
suppressed in short-day eggs (Fig. 4a). This comprehensive downregulation of both protein
synthesis and degradation machinery effectively pauses the cellular protein economy to preserve
resources for extended diapause.

Concurrent with these metabolic changes, key developmental signaling pathways diverged
between diapause and non-diapause trajectories. At 40 hpo, we observed significant
downregulation of TGF-beta signaling in short-day eggs (Fig. 4a). As a master regulator of cell
proliferation and differentiation [50,51], suppression of TGF-beta signaling effectively applies
molecular brakes on developmental progression, occurring before morphological divergence.

Discussion

Our comprehensive molecular analysis reveals that maternal diapause induction in *D.*
nigrofasciatus proceeds through a precisely orchestrated temporal cascade of molecular events
(Fig. 4d). Both long- and short-day eggs initially share common early developmental processes,
including endocytosis and autophagy for recycling maternal components (12-24 hpo). However,
short-day eggs diverge through a distinctive molecular program: early activation of chromatin
remodeling machinery and precocious RNA processing (24-40 hpo), followed by progressive
metabolic restructuring with upregulation of amino acid catabolism and gluconeogenesis
pathways that continues through 72 hpo. In contrast, long-day eggs maintain sustained protein
synthesis machinery, RNA processing at normal developmental timing, and active cell cycle
progression leading to cellular differentiation. This temporal progression demonstrates that
diapause preparation is not a simple developmental arrest but rather an active programming
process requiring substantial molecular investment before the morphological divergence point.
285
An important consideration in interpreting this temporal cascade is the timing of the maternal-to-
zygotic transition (MZT), during which developmental control shifts from maternally deposited
transcripts to embryo genome–derived transcription [52]. Although the precise timing of MZT has
not been formally characterized in *D. nigrofasciatus*, previous analyses in this species
demonstrated clear temporal increases in multiple developmental regulators, including
segmentation-related genes, between 12 and 40 hpo, suggesting the onset of zygotic
transcription during this period [23]. In line with these observations, our RNA-seq data show
minimal photoperiod-dependent differences at 12 hpo, followed by progressive divergence from
24 hpo onward, suggesting that embryo genome transcription is already active by 12–24 hpo.
Under this framework, maternal diapause induction is unlikely to operate solely through
differential deposition of maternal mRNAs. Instead, maternal photoperiodic experience likely
establishes distinct epigenetic states in the egg, such as chromatin accessibility, that
subsequently bias embryo-driven transcriptional programs toward diapause or direct
development. A similar principle has been demonstrated in *Drosophila melanogaster*, in which
MZT occurs well before cellularization and early chromatin and transcriptional states critically
shape later developmental trajectories [52].
302
The temporal advancement of components of RNA processing in short-day eggs, peaking at 24
hpo versus 40 hpo in long-day eggs, represents a striking example of heterochronic gene
expression. This precocious activation likely serves multiple adaptive functions. First, the early
investment in RNA processing capacity could enable the preparation of stable mRNA pools that
persist throughout the diapause period [53,54]. Second, this enhanced RNA processing machinery
may facilitate rapid transcriptional responses upon receiving diapause-terminating signals,
allowing quick resumption of development [53,54]. Third, it may establish diapause-specific
alternative splicing programs that generate protein isoforms optimized for dormancy [55]. The
sustained elevation of these components throughout diapause preparation suggests they play a
central role in maintaining the molecular readiness for eventual developmental resumption.
313
The early upregulation of Brahma complex components at 24 hpo, preceding any morphological
changes, suggests that chromatin remodeling serves as the primary molecular switch for
diapause induction. This finding aligns with emerging evidence that epigenetic mechanisms
underlie phenotypic plasticity across diverse organisms [56–59]. The selective reduction in chromatin
accessibility at developmental regulators such as *Kr-h1* and *crol* effectively locks cells into a pre-
differentiation state, preventing the normal progression of embryonic development [44,45]. Both
genes are central components of juvenile hormone (JH) and ecdysteroid signaling cascades,
which are essential endocrine pathways governing insect growth and developmental progression.

Notably, however, our RNA-seq data did not reveal clear differences in juvenile hormone or
ecdysteroid signaling pathways between long- and short-day eggs at 24 hpo. Consistent with this,
prior physiological assays in *D. nigrofasciatus* did not detect egg JH III or photoperiod-dependent
differences in egg 20-hydroxyecdysone during early development [60]. These observations suggest
that early chromatin closure at endocrine-responsive developmental regulators does not reflect
active hormonal signaling at this stage, but rather a preemptive epigenetic mechanism that
suppresses the embryo's capacity to respond to later endocrine cues required for developmental
progression. Within this framework, rather than being directly specified by maternally deposited
Brahma transcripts, we propose that maternal photoperiodic perception establishes distinct
epigenetic contexts in the egg, within which the Brahma complex might function as an executor
that stabilizes and amplifies these maternally primed regulatory states. This epigenetic regulation
occurs during a critical window (within 24 hpo), demonstrating how maternal environmental
signals can be translated into stable epigenetic modifications that fundamentally alter offspring
developmental trajectory. Intriguingly, this 24-hour window coincides with the period during which
temperature pulses can still reverse diapause fate decisions [61], suggesting that chromatin
remodeling may define developmental fate plasticity.

In *D. nigrofasciatus*, parental RNAi is technically feasible and could, in principle, enable functional
validation of the Brahma complex [62]. However, knockdown of Brahma has been reported to
severely impair fertility and fecundity in other insect species, including coleopteran and
hemipteran models [63], rendering such approaches impractical for maintaining sufficient egg
production in this system. Importantly, diapause fate in *D. nigrofasciatus* eggs can be
experimentally induced or prevented by temperature manipulation [63], and diapause termination
occurs spontaneously after prolonged arrest and is accelerated by winter-like chilling [64].
Therefore, examining Brahma complex expression and chromatin states under experimentally
altered diapause induction and termination trajectories provides a realistic and informative
framework for assessing its functional relevance *in vivo*.

The progressive metabolic restructuring observed from 56 hpo onwards in short-day eggs reveals
a sophisticated survival strategy distinct from simple developmental arrest. The coordinated
upregulation of amino acid catabolism pathways and gluconeogenic enzymes indicates a
fundamental shift from anabolic growth metabolism to a catabolic maintenance mode. This
metabolic flexibility enables short-day eggs to utilize maternally deposited amino acids as an
energy source while preserving essential proteins needed for eventual development [53,54]. The
timing of these changes, beginning before morphological divergence and intensifying through 72
hpo, indicates that metabolic programming is an integral preparatory component rather than a
consequence of arrest. This preemptive metabolic adjustment, coupled with the suppression of
protein synthesis and proteasome pathways, creates a molecular economy optimized for long-
term survival while maintaining developmental competence [13,21].

The remarkable plasticity of *D. nigrofasciatus*, which can facultatively switch between diapause
and direct development based on maternal photoperiod experience, makes it a unique model for
understanding how environmental information is translated into developmental outcomes [64,65].
Our findings illuminate how subtle maternal signals can be progressively amplified into
dramatically different developmental outcomes through self-reinforcing molecular cascades that
combine epigenetic, transcriptional, and metabolic regulation. This temporal orchestration
framework likely extends beyond insects to other organisms that employ maternal programming
of developmental dormancy, including delayed implantation in mammals [65,66], suggesting that
coordinated molecular timing represents a fundamental mechanism for transgenerational

environmental adaptation across diverse organisms. This cricket model system provides a
powerful framework for understanding the universal principles governing how organisms across
the tree of life integrate environmental information to orchestrate developmental decisions
through conserved molecular mechanisms.

Methods

Sample collection

The *D. nigrofasciatus* individuals used in this study was collected in a grassy field on the campus
of Osaka Metropolitan University (34°35′N, 135°30′E) and the bank of the Yamato-gawa River
(34°35′N, 135°30′E), Japan, from June to November 2022 and 2023. The rearing temperature
(25.0 ± 1.0 °C) was chosen to approximate ambient temperatures commonly experienced during
the natural oviposition seasons of this species in the Osaka region. Approximately 40–60
individuals were reared in a plastic case (29.5 cm width, 19.0 cm depth, and 17.0 cm height)
under long-day conditions (16-h light:8-h dark, 25.0 ± 1.0 °C) with a piece of moist cotton as a
water source and oviposition site. They were fed an artificial diet (Masu Chigyo Super EPC-2,
Marubeni Nisshin Feed Co., Ltd., Tokyo, Japan) and fresh carrot. The collected eggs were placed
on a piece of moist cotton in plastic dishes (50 mm diameter, 12 mm depth) and continuously
maintained under long-day conditions [66]. For the experiments, first instar nymphs were kept in
long-day conditions or were transferred to short-day conditions (12-h light:12-h dark, 25.0 ±
1.0 °C). Adult females reared under long-day conditions lay eggs destined to direct development
(non-diapause), whereas those under short-day conditions lay eggs destined to enter diapause [67].
In the present study, females reared under long- and short-day conditions are referred to as long-
and short-day females, respectively. Eggs laid by long- and short-day females are referred to as
long- and short-day eggs, respectively. Eggs deposited within a 2 h period were collected and
used for experiments. These eggs were continuously maintained under long-day conditions until
use for experiments. Under these conditions, the incidence of diapause was 14.6% in eggs laid
by long-day females and 95.7% in eggs laid by short-day females, confirming that diapause fate
was determined by maternal photoperiodic history rather than by environmental conditions
experienced by the eggs.

Genome and RNA sequencing

Alive *D. nigrofasciatus* was used to extract its genomic DNA. Total genomic DNA was extracted
from 10 whole male bodies after removal of the digestive tract, as well as from the head and hind
legs of a single male and female *D. nigrofasciatus*, using NucleoBond® HMW DNA (Macherey-
Nagel, Germany) according to the manufacturer's instructions. The resulting genomic DNA was
size-selected using a Short Read Eliminator Kit (PacBio, CA, USA). Oxford Nanopore
Technologies (ONT) sequencing libraries were then constructed and sequenced on the
PromethION platform (Oxford Nanopore Technologies, UK) with the ligation sequencing kit SQK-
LSK112 and flow cell R10.4. Base-calling was performed using Guppy basecalling software
v6.3.8 with the model dna_r10.4_e8.1_sup. Finally, we obtained 30.40 Gbp ONT sequencing
data; average and N50 read lengths were 19.11 Kbp and 31.18 Kbp, respectively. Additionally,
using the same DNA sample, we prepared a whole genome sequencing library using the TruSeq
DNA PCR-Free Library Prep Kit (Illumina, CA, USA) following the manufacturer's protocol. This
library was sequenced on the Illumina NovaSeq 6000 platform with 150 bp paired-end reads
(PE150), generating 152.7 Gbp of short-read data. For chromosome-scale scaffolding by the
Dovetail™ Omni-C™ Kit (Dovetail Genomics, CA, USA), the head and hind legs of another single
male *D. nigrofasciatus* were used according to the manufacturer's instructions. The Hi-C

sequencing library was built on the Illumina NovaSeq 6000 platform with PE150 and generated
127.1 Gbp raw data. DNA purity and concentrations were measured by spectrometry using
NanoPhotometer NP80-TOUCH (Implen, Germany) and fluorometry using Qubit 4 (Thermo
Fisher Scientific, MA, USA).
423
Total RNA was extracted from whole eggs 12, 24, 40, 56, and 72 hours post-oviposition (hpo)
using TRIzol Reagent with the PureLink RNA Mini Kit and PureLink DNase Set (Thermo Fisher
Scientific) after homogenizing with the BioMasher II tissue grinder (Nippi, Inc., Tokyo, Japan)
(Supplementary Table 16). Total RNA of 12, 24, 40, and 56 hpo eggs was purified with ssDNA /
RNA Clean & Concentrator (ZYMO Research, CA, USA). RNA integrity was confirmed using a
NanoDrop spectrophotometer (Thermo Fisher Scientific). For 12, 24, 40, and 56 hpo eggs,
libraries were constructed using the NEBNext® Single Cell/Low Input RNA Library Prep Kit (New
England Biolabs, MA, USA) and sequenced on an Illumina NextSeq 1000 platform with PE150.
For 72 hpo eggs, libraries were prepared using the SMART-Seq® HT PLUS Kit (Takara Bio,
Japan) and sequenced on an Illumina NovaSeq 6000 platform with PE150. To improve gene
annotation and provide tissue-specific expression profiles, we also performed RNA-seq on adult
tissues. Ovaries and neural tissues (brains and suboesophageal ganglions) were dissected from
10-day-old adult females at ZT 9-10, maintained under either long- or short-day photoperiod
conditions. For these adult tissue samples, mRNA was enriched using the NEBNext® Poly(A)
mRNA Magnetic Isolation Module, and libraries were prepared using the NEBNext® Ultra™ II
Directional RNA Library Prep Kit (New England Biolabs). Both brain and ovary libraries were
sequenced on an Illumina NovaSeq 6000 platform with PE150.
441
442 **Genome size estimation**
443
444 Genome size was computationally estimated using k-mer frequency distribution analysis. Raw
445 Illumina sequencing reads were processed using KMC v3 (K-mer Counter) [68] with k-mer length of
446 21. K-mers were counted. The resulting k-mer frequency histogram was generated using
447 kmc_tools with a maximum coverage cutoff of 10,000. The k-mer frequency distribution was
448 analyzed using GenomeScope2 [69] with k=21 and ploidy set to 2 (diploid).
449
Genome size was also estimated using flow cytometry to provide an independent validation of the
computational estimates. The 1C (haploid) genome size of males and females of *D.
nigrofasciatus* was estimated with reference to the method previously described [70]. Uniform
suspensions of single nuclei were extracted from a single *D. nigrofasciatus* head, a single
Drosophila virilis female head (1C =328 Mbp) and a portion of brain from a *Periplaneta americana*
male (1C =3300 Mbp) using the BD Cycletest Plus DNA Reagent Kit (BD Biosciences, CA, USA).
After trypsin and RNase A treatments, the samples were incubated with propidium iodide for 10
min in the dark at 4 °C. Finally, the sample was filtered through a 50-μm nylon mesh. Cellular
DNA content was assessed on the BD Accuri C6 Flow Cytometer (BD Biosciences). The obtained
data were analyzed using the FCS Express software (De Novo Software, CA, USA). Haploid (1C)
DNA quantity was calculated as (2C sample mean fluorescence/2C standard mean fluorescence)
times 328 Mbp for the *D. virilis* standard and times 3300 Mbp for the *P. Americana* standard. The
estimates based on the two standards were averaged to produce a 1C estimate for each sample.
463
464 *De novo* **genome assembly**
465
466 Initial genome assembly was performed using MaSuRCA v4.1.0 [71] with a hybrid approach
467 combining Illumina paired-end reads and Oxford Nanopore long reads. Illumina paired-end reads
468 (153 bp mean insert size, 102 bp standard deviation) were provided as input along with the

469 parameters FLYE_ASSEMBLY=0 to use CABOG assembly algorithm. The resulting primary
470 assembly scaffolds were used for subsequent processing.
471
To remove redundant haplotigs from the assembly, we employed Purge Haplotigs v1.1.2 [72].
Illumina paired-end reads were mapped to the primary assembly using minimap2 v2.24-r1122 [73]
with the -ax sr option. Read-depth histogram was generated using the purge_haplotigs hist
command to identify haploid and diploid peaks. Based on the histogram analysis showing an
average coverage of 94.5×, cutoff values were set as: low=5, midpoint=70, high=190, with junk
coverage threshold at 101. Coverage statistics were calculated using purge_haplotigs cov,
followed by the actual purging step using purge_haplotigs purge.
479
Potential contamination in the assembly was removed using BlobTools v1.1.127 [74], which
analyzes unexpected coverage, GC content, or similarity to bacterial and other contaminant
sequences. Sequence coverage was determined by mapping Illumina reads with bwa v0.7.17-
r1188. Similarity analysis was performed using BLASTn v2.13.0+ [75] against NCBI NT database
v5 (options: -task megablast -culling_limit 10 -evalue 1e-25 -outfmt '6 qseqid staxids bitscore std
sscinames sskingdoms stitle'). Mitochondrial genomes were identified through gene prediction
using the MITOS2 webserver (accessed on November 28, 2023) and subsequently removed. As
a result, one contig was removed as a bacterial genome and another as a mitochondrial genome.
488
The purged and decontaminated assembly was further scaffolded using Hi-C data. The input *de*
novo assembly and Dovetail OmniC library reads were processed using HiRise [76], a software
pipeline designed specifically for using proximity ligation data to scaffold genome assemblies.
Dovetail OmniC library sequences were aligned to the draft input assembly using bwa [77]. The
separations of Dovetail OmniC read pairs mapped within draft scaffolds were analyzed by HiRise
to produce a likelihood model for genomic distance between read pairs. This model was used to
identify and break putative misjoins, score prospective joins, and make joins above a threshold.
496
A Circos plot illustrating the distribution of genomic elements was generated using Circos v0.69-9
[78]. The plot displays the major scaffolds arranged in a circular layout with tracks showing gene
density, GC content, repeat element distribution, and other genomic features. The completeness
of the genome assembly was evaluated using BUSCO (Benchmarking Universal Single-Copy
Orthologs) v5.1.2 [78] implemented in gVolante [79]. Assembly completeness was assessed against
both the arthropoda_odb10 and insecta_odb10 lineage-specific ortholog datasets to evaluate the
representation of conserved genes at different taxonomic levels.
504
505 **Prediction of repeat regions and annotation of protein-coding genes**
506
507 For *de novo* repeat prediction, RepeatModeler v2.0.1 [80] was first used to identify novel repetitive
508 elements in the genome. This *de novo* library was supplemented with known repeat sequences
509 from closely related species to create a comprehensive repeat database. The combined repeat
510 library was then used to identify and softmask repetitive elements in the genome using
511 RepeatMasker v4.1.2 [81].
512
513 Structural annotation of protein-coding genes was performed on the soft-masked genome using
514 three complementary approaches: *ab initio* prediction, homology-based prediction, and RNA-seq-
515 based prediction. Each prediction method incorporated RNA-seq data from eggs, ovaries, and
516 brains as supporting evidence.
517

The *ab initio* prediction was carried out using BRAKER v3.0.2 [26–29,82,83], incorporating protein data
from OrthoDB 10's arthropoda dataset and the mapping data of the filtered RNA-seq reads. This
BRAKER prediction served as the foundation for our gene set. To complement and improve this
base set, we employed two additional approaches. GeMoMa v1.9 [29] was used for homology-
based prediction with gene sets from four model species (*Apis mellifera*, *Drosophila*
melanogaster, and *Tribolium castaneum*), while StringTie2 v2.1.7 [83] was used for RNA-seq-based
transcript assembly and gene prediction. The predictions from GeMoMa and StringTie2 were
used to identify and add genes that BRAKER3 had missed. These combined predictions were
merged and duplicate genes were removed using GffCompare v0.12.6 [84] with transcript
classification codes set to u, i, and y to form a final, comprehensive consensus gene set.
528
Gene functional annotation was conducted using eggNOG-mapper online (http://eggnog-
mapper.embl.de/) and BLASTp-based methods. For the BLASTp-based annotation, we searched
against databases including *Homo sapiens* (GRCh38), *Mus musculus* (GRCm39), *Caenorhabditis*
elegans (WBcel235), *Drosophila melanogaster* (BDGP6.32), and UniProt Swiss-Prot to identify
the best hits for annotation (E-value < 1.0×10^{-5}). The completeness of the genome assembly
was also evaluated using BUSCO v5.1.2 [78] implemented in gVolante [79].
535
536 **Phylogenetic analysis**
537
538 To reconstruct the phylogenetic relationships among cricket species, we analyzed protein
539 sequences from eight species: *Acheta domesticus* [85], *Apteronemobius asahinai* [86], *Dianemobius*
540 *nigrofasciatus*, *Gryllus bimaculatus* [87], *Laupala kohalensis* [87], *Locusta migratoria* [88], *Schistocerca*
541 *gregaria* (GCF_023897955.1), and *Teleogryllus occipitalis* [89]. Translated coding sequences
542 (CDS) from each species were used as input for ortholog identification.
543
Orthologous gene groups were identified using OrthoFinder v2.5.4 [90] with default parameters.
Single-copy orthologous genes (SCOs) were extracted for phylogenetic reconstruction, resulting
in 334 SCOs shared across all eight species. Each SCO was aligned using MAFFT v7.505 [91] with
the --auto option, which selected the L-INS-i strategy. The aligned sequences were subsequently
trimmed using trimAl v1.4.rev15 [92] with the -automated1 option to remove poorly aligned regions
and gaps.
550
The trimmed alignments were concatenated, and a partition file was generated to define gene
boundaries within the concatenated alignment. Maximum likelihood phylogenetic analysis was
performed using IQ-TREE v2.2.0.3 [93] with the following parameters: -bb 1000 for ultrafast
bootstrap analysis with 1000 replicates, -alrt 1000 for SH-like approximate likelihood ratio test
with 1000 replicates, -m MFP for automatic model selection using ModelFinder [94], and partition
model optimization (-sp).
557
558 Divergence times were obtained from TimeTree (https://timetree.org/) [95] and mapped onto the
559 resulting phylogeny to provide temporal context for the evolutionary relationships among the
560 analyzed species.
561
562 **RNA-seq analysis**
563
To prepare the reference transcriptome for quantification, the longest isoform for each gene was
extracted from the gene models. Transcript abundance estimation was performed using Salmon
v1.10.1 [96]. First, a quasi-mapping index was built from the longest isoform cDNA sequences

derived from the *D. nigrofasciatus* genome annotation. RNA-seq data from eggs collected at five
developmental time points (12, 24, 40, 56, and 72 hpo) under both diapause (short-day) and non-
diapause (long-day) conditions were analyzed. Paired-end reads were mapped using Salmon
quant with automatic library type detection (-l A). Transcript-level estimates were imported and
processed using the tximport package [97] in R. Transcripts per million (TPM) values and estimated
read counts were extracted for downstream analysis.
573
574 Differential expression analysis was performed using DESeq2 v1.38.0 [98]. Low-expressed genes
575 were filtered by removing genes with total read counts less than 10 across all samples. Library
576 size normalization and dispersion estimation were performed using default DESeq2 parameters,
577 followed by the Wald test for differential expression. Results were ordered by adjusted *p*-value
578 (Benjamini-Hochberg correction) and exported for further analysis.
579
580 Gene Ontology (GO) and KEGG pathway enrichment analyses were performed using g:Profiler
581 (https://biit.cs.ut.ee/gprofiler/) [99]. Differentially expressed genes were mapped to their *Drosophila*
582 *melanogaster* orthologs based on the BLASTp annotation results, and the *D. melanogaster* gene
583 IDs from the BDGP6.32 reference genome were used as input for enrichment analysis.
584
585 **Principal component analysis**
586
587 To visualize the overall transcriptional differences between diapause and non-diapause
588 conditions across developmental time points, three-dimensional principal component analysis
589 (PCA) was performed. TPM values from all samples were used. Genes with zero variance across
590 all samples were removed prior to analysis. PCA was performed using the prcomp function in R
591 with scaling (scale. = TRUE).
592
593 The first three principal components were extracted and visualized using the rgl package for 3D
594 plotting. Samples were colored according to their diapause status (diapause vs. non-diapause) to
595 assess the separation of conditions in the reduced dimensional space. The interactive 3D
596 visualization was generated using plot3d functions.
597
598 **Time-series clustering analysis, cluster comparison and classification**
599
600 To identify genes with similar temporal expression patterns during embryonic development, time-
601 series clustering was performed using the Mfuzz package [98]. First, normalized count data from
602 DESeq2 were obtained for each developmental time point (12h, 24h, 40h, 56h, and 72h). The
603 mean expression values were calculated separately for diapause and non-diapause conditions,
604 and genes with zero standard deviation were removed.
605
606 To focus on genes with substantial expression changes across time points, genes with a
607 coefficient of variation less than 1.0 were filtered out. This filtering step ensures that the clustering
608 analysis captures biologically meaningful expression dynamics rather than noise from low-
609 variance genes.
610
611 The filtered expression data were converted to ExpressionSet objects and standardized using the
612 Mfuzz standardize function. The fuzziness parameter (m) was estimated using the mestimate
613 function. Based on the elbow method analysis and practical considerations, the number of
614 clusters was set to 20 for both diapause and non-diapause datasets.
615

To identify relationships between diapause and non-diapause expression patterns, cosine
similarity was calculated between all pairwise combinations of cluster centroids. Cluster pairs
were classified as "homochronic" (similar temporal dynamics) or "heterochronic" (distinct
temporal dynamics) based on a cosine similarity threshold of 0.8. This classification allows for the
identification of genes that maintain similar expression patterns across conditions versus those
that show condition-specific temporal regulation. Gene overlap significance between clusters was
assessed using upper-tail hypergeometric tests. Cluster pairs with p-values < 10^{-10} were
considered significantly overlapping, indicating non-random association of genes between
diapause and non-diapause conditions.
625
626   For functional interpretation, genes from cluster pairs in the top and bottom 10th percentiles of
627   cosine similarity were extracted. These gene sets, representing the most homochronic and
628   heterochronic patterns respectively, were subjected to GO and KEGG enrichment analysis using
629   g:Profiler [99] to identify biological processes associated with conserved versus divergent temporal
630   expression dynamics during diapause induction.
631

ATAC-seq library preparation and sequencing

632
633
634   Chromatin accessibility profiling was performed using ATAC-seq on 24 hpo eggs. For either long-
635   or short-day conditions, two biological replicates were prepared (Supplementary Table 17).
636
637   Libraries were prepared using the ATAC-seq Kit (Active Motif, CA, USA). Frozen egg samples
638   were lysed in 100 µL ice-cold ATAC Lysis Buffer and mechanically disrupted using scissors to
639   release nuclei. Cell debris and eggshell fragments were removed by filtration through
640   pluriStrainer-Mini 40 µm filters (pluriSelect, Germany). Subsequent library preparation steps were
641   performed according to the manufacturer's protocol.
642
643   Library quality was assessed using NanoDrop One (Thermo Fisher Scientific) for concentration
644   measurement and DNA ScreenTape (Agilent Technologies, CA, USA) for fragment size
645   distribution analysis. Qualified libraries were sequenced on an Illumina NovaSeq 6000 platform
646   with 150PE. The sequencing generated 11.15-11.60 Gbp of data per sample.
647

ATAC-seq data analysis

648
649
650   Raw sequencing reads were quality-trimmed using Skewer v0.2.2 [100] with the following
651   parameters: maximum allowed error rate of 0.1, maximum allowed indel error rate of 0.03, and
652   minimum read length of 18 bp after trimming. Paired-end mode was enabled for all samples.
653
654   Trimmed reads were aligned to the *D. nigrofasciatus* reference genome using Bowtie2 v2.3.4.2
655   [101] with parameters -N 1 (maximum mismatch of 1) and -X 2000 (maximum fragment length of
656   2000 bp). Peak calling was performed using MACS2 v2.1.2 [102] with a q-value threshold of 0.05 (-
657   q 0.05) and parameters --nomodel --shift -75 --extsize 150, which defines a 150 bp region
658   centered on the read start position.
659
660   To create a unified peak set, detected peaks from all samples were sorted and merged using
661   bedtools v2.27.1 [103] with default settings. The resulting consensus peak regions were formatted
662   into SAF format for downstream analysis.
663

Genomic annotation of peaks was performed by categorizing each peak into the following
genomic regions: 1-5 kb upstream (1to5kb), <1 kb upstream (promoter), 5'UTR, exon, intron,
3'UTR, exon/intron boundary (defined as 100 bp surrounding exon-intron junctions), and
intergenic regions. When peaks overlapped multiple features, priority was given in the following
order: 5'-/3'-UTR > exon/intron boundary > exon/intron. Genes located within 1 Mbp of each peak
were assigned as nearby genes.
670
Read counts for consensus peaks were quantified using featureCounts v1.6.3 [104]. Differential
accessibility analysis between diapause-inducing and non-diapause conditions was performed
using edgeR v3.26.8 [105] with TMM normalization. Log2 fold changes were calculated for each
peak region. Peaks with uncorrected *p*-values < 0.05 were considered differentially accessible,
and their associated nearby genes from different genomic contexts were extracted for functional
analysis. Genomic contexts included: 1-5 kb upstream regions (1-5 kbp upstream of transcription
start sites, TSS), promoter regions (<1 kbp upstream of TSS), 5'UTR, exon, intron, 3'UTR,
exon/intron boundaries (within 100 bp of exon-intron junctions), and intergenic regions (areas not
classified into any of the above categories).

Data availability

Raw read data used in this article (RNA-seq, ATAC-seq, Omni-C and WGS) have been submitted
at DNA Data Bank of Japan (DDBJ) repository under accession no. PRJDB18895 and are
publicly available as of the date of publication. The datasets are available in following link;
https://ddbj.nig.ac.jp/resource/bioproject/PRJDB18895. The assembled genome and annotation
datasets are available in https://doi.org/10.6084/m9.figshare.29665742.v1. The scripts used for
the analyses in this study are available on GitHub (https://github.com/Kataoka-K-
Lab/Dnigrofasciatus_EggDiapause.git).
688

Acknowledgments

This work was supported by the Cooperative Research Grant of the Genome Research for
BioResource (NODAI Genome Research Center, Tokyo University of Agriculture), Cross-
ministerial Moonshot Agriculture, Forestry and Fisheries Research and Development Program,
"Technologies for Smart Bio-industry and Agriculture" (Bio-oriented Technology Research
Advancement Institution, BRAIN) [Grant Number: JPJ009237], JSPS KAKENHI Grant-in-Aid for
Scientific Research (C) [Grant Number: 21K05614], and Grant-in-Aid for JSPS Fellows [Grant
Number: 21J23478/22KJ2609]. We thank Dr. Tarô Adati (Department of International Agricultural
Development, Faculty of International Agriculture and Food Studies, Tokyo University of
Agriculture) for his valuable advice and support.

 Figures legends

Figure 1. Life cycle and genomic characterization of the band-legged ground cricket,
***Dianemobius nigrofasciatus*. (a)** Adult female *D. nigrofasciatus*. **(b)** Annual life cycle diagram in
Osaka, Japan, illustrating photoperiod-dependent developmental transitions and diapause
regulation. The outer ring represents the developmental stages (egg, nymph, and adult) and the
diapause egg state. The inner circle shows seasonal progression with photoperiod-dependent
phases: the pink area indicates the period with the longer critical day length, whereas the blue
area indicates the period with the shorter critical day length. The first-generation adults emerge in
September-October under short-day conditions and produce predominantly diapause-destined
eggs that overwinter in the cellular blastoderm stage, whereas the second-generation adults
appear in June-August under long-day conditions and produce non-diapause eggs. **(c)** Flow
cytometric analysis of brain cells from adult females and males showing genome size estimation.
Box plots show the distribution of genome size measurements, with blue numbers indicating
mean ± 95% confidence interval for each sex. **(d)** Circos plot depicting genomic features across
the ten linkage groups (LG1-10). From outer to inner tracks: gene density (I), repetitive element
content (II), and GC content distribution (III). **(e)** BUSCO completeness assessment of the
genome assembly and gene annotation against Arthropoda and Insecta databases via the
gVolante web server. Colors indicate complete single-copy (solid green), completely duplicated
(blue with diagonal stripes), fragmented (yellow crosshatched pattern), and missing (red with
black dot pattern) genes.
720
Figure 2. Transcriptome dynamics during early embryogenesis in long- and short-day
eggs of *D. nigrofasciatus*. (a) Schematic representation of developmental progression from 0 to
72 hours post-oviposition (hpo) at 25°C. Adults reared under short-day conditions produce
diapause-destined eggs (short-day eggs), while those under long-day conditions produce eggs
that undergo direct development (long-day eggs). The morphological features of early embryonic
stages were characterized by Tanigawa et al. (2009). In both egg types, nuclei migrate from the
yolk mass to the egg surface by 24 hpo. Cellular blastoderm formation occurs between 40-56 hpo
in both trajectories. While long-day eggs proceed to form a germ band by 72 hpo, short-day eggs
arrest development at the cellular blastoderm stage, maintaining this state throughout the
diapause period. **(b)** Three-dimensional principal component analysis of transcriptome profiles
across developmental time points. Red and blue marks indicate long- and short-day eggs,
respectively. Dashed line circles indicate clustering of samples corresponding to each
developmental time point. **(c)** *Left panel*: Number of differentially expressed genes (DEGs)
between long- and short-day eggs at each developmental timepoint (12, 24, 40, 56, and 72 hpo).
DEGs were defined as genes with $|\log_2$ fold change (FC)$| > 1$ and adjusted *p*-value < 0.05. Green
bars with crosshatched pattern indicate genes upregulated in short-day eggs, while yellow bars
with diagonal stripes represent genes downregulated in short-day eggs. *Right panels*: Temporal
dynamics of gene expression changes within each egg type, comparing later versus earlier
developmental timepoints. For long- (left) and short-day eggs (right), bars show the number of
genes upregulated (green with crosshatched pattern) or downregulated (yellow with diagonal
stripes) at the later timepoint relative to the earlier timepoint in each comparison ($|\log_2$ FC$| > 1$,
adjusted *p*-value < 0.05).
743
Figure 3. Comparative analysis of temporal gene expression dynamics between long- and
short-day eggs. (a) Heat map showing the significance of gene overlap between temporal
expression clusters in long- and short-day eggs across five developmental timepoints (12, 24, 40,
56, and 72 hpo) as derived from an upper-tail hypergeometric test. Gene expression profiles were

clustered using the Mfuzz soft clustering algorithm. Numbers on the right and bottom axes
represent cluster IDs for long-day and short-day eggs, respectively. Color intensity represents -
$\log_{10}$(adjusted *p*-value), with darker red indicating more significant overlap between clusters.
Homochronic clusters (highlighted in blue boxes) show conserved temporal expression patterns
between the two egg types. Heterochronic clusters (highlighted in green boxes) exhibit shifted
temporal dynamics between long- and short-day eggs. **(b1, b2, c1, and c2)** Expression
trajectories show representative homochronic **(b)** or heterochronic **(c)** cluster pairs, where color
intensity reflects membership scores calculated by Mfuzz (magenta indicating high membership
and green indicating low membership). Bar graphs display functional enrichment analysis results
using overlapping genes, with terms categorized as GO Biological Process (GO:BP, orange), GO
Cellular Component (GO:CC, green), GO Molecular Function (GO:MF, blue), or KEGG pathways
(red). For each category, the top five most significantly enriched terms (ranked by adjusted *p*-
value) are shown; when fewer than five terms were significant, all enriched terms are displayed.
Bar length represents -$\log_{10}$(adjusted *p*-value).
762
Figure 4. Temporal coordination of biological processes during early embryogenesis in
long- and short-day eggs. (a) KEGG pathway enrichment analysis of differentially expressed
genes (DEGs) identified between long-day and short-day conditions at each time point (adjusted
p-value < 0.05). Dot size represents the percentage of DEGs annotated to each pathway, and
color intensity indicates the adjusted *p*-value. **(b and c)** Temporal expression patterns (TPM
values) of genes involved in chromatin remodeling **(b)** and gluconeogenesis **(c)**. The red frames
in (b) highlight the critical periods of expression divergence between long- and short-day eggs.
(d) Schematic representation of key biological processes and their temporal dynamics from 0 to
72 hpo. Both long- and short-day eggs show similar timing of membrane trafficking processes
(endocytosis and autophagy) during early embryogenesis (12-24 hpo). Chromatin remodeling
occurs within a specific time window around 24 hpo. Prior to diapause entry at the cellular
blastoderm stage, short-day eggs show advanced activation of RNA processing and protein
synthesis compared to long-day eggs. During later development (>56 hpo), short-day eggs
activate amino acid metabolism and gluconeogenesis pathways, while long-day eggs proceed
with neural development and cell cycle progression. Blue-shaded processes indicate molecular
programs that are activated in both short-day (diapause) and long-day (non-diapause) eggs,
whereas red-shaded processes indicate pathways that are specifically activated in only one
developmental trajectory.
781

References

1. Lennon, J. T., den Hollander, F., Wilke-Berenguer, M. & Blath, J. Principles of seed banks and the emergence of complexity from dormancy. *Nat. Commun.* **12**, 4807 (2021).
2. Özgüldez, H. Ö. & Bulut-Karslioğlu, A. Dormancy, quiescence, and diapause: Savings accounts for life. *Annu. Rev. Cell Dev. Biol.* **40**, 25–49 (2024).
3. Wilsterman, K., Ballinger, M. A. & Williams, C. M. A unifying, eco‐physiological framework for animal dormancy. *Funct. Ecol.* **35**, 11–31 (2021).
4. Denlinger, D. L. *Insect Diapause*. (Cambridge University Press, Cambridge, England, 2022).
5. J., M., Tauber, C. A. & Masaki, S. *Seasonal Adaptations of Insects*. (Oxford University Press, New York, NY, 1985).
6. Rinehart, J. P. *et al.* Up-regulation of heat shock proteins is essential for cold survival during insect diapause. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11130–11137 (2007).
7. Benoit, J. B. Water management by dormant insects: comparisons between dehydration resistance during summer aestivation and winter diapause. *Prog. Mol. Subcell. Biol.* **49**, 209–229 (2010).
8. Hahn, D. A. & Denlinger, D. L. Energetics of insect diapause. *Annu. Rev. Entomol.* **56**, 103–121 (2011).
9. Mousseau, T. A. & Fox, C. W. The adaptive significance of maternal effects. *Trends Ecol. Evol.* **13**, 403–407 (1998).
10. Oku, K., Yano, S. & Takafuji, A. Different maternal effects on diapause induction of tetranychid mites, *Tetranychus urticae* and *T. kanzawai* (Acari: Tetranychidae). *Appl. Entomol. Zool. (Jpn.)* **38**, 267–270 (2003).
11. Huestis, D. L. & Marshall, J. L. Interaction between maternal effects and temperature affects diapause occurrence in the cricket *Allonemobius socius*. *Oecologia* **146**, 513–520 (2006).
12. Masaki, S. Climatic adaptation and photoperiodic response in the band-legged ground cricket. *Evolution* **26**, 587–600 (1972).
13. Shiga, S. & Numata, H. Seasonal changes in the incidence of embryonic diapause in the band-legged ground cricket, *Dianemobius nigrofasciatus*. *Zool. Sci.* **14**, 1015–1018 (1997).
14. Shintani, Y. Effect of seasonal variation in host‐plant quality on the rice leaf bug, *Trigonotylus caelestialium*. *Entomol. Exp. Appl.* **133**, 128–135 (2009).
15. Rockey, S., Miller, B. B. & Denlinger, D. A diapause maternal effect in the flesh fly, *Sarcophaga bullata*: Transfer of information from mother to progeny. *Journal of Insect Physiology* **35**, 553–558 (1989).
16. Bova, J., Soghigian, J. & Paulson, S. The prediapause stage of *Aedes japonicus japonicus* and the evolution of embryonic diapause in Aedini. *Insects* **10**, 222 (2019).
17. Schneiderman, H. A. & Horwitz, J. The induction and termination of facultative diapause in the chalcid wasps *Mormoniella vitripennis* (Walker) and *Tritneptis klugii* (Ratzeburg). *J. Exp. Biol.* **35**, 520–551 (1958).
18. Saunders, D. S. Larval diapause of maternal origin: Induction of diapause in *Nasonia vitripennis* (Walk.) (Hymenoptera: Pteromalidae). *J. Exp. Biol.* **42**, 495–508 (1965).
19. Poelchau, M. F., Reynolds, J. A., Elsik, C. G., Denlinger, D. L. & Armbruster, P. A. Deep sequencing reveals complex mechanisms of diapause preparation in the invasive mosquito, *Aedes albopictus*. *Proc. Biol. Sci.* **280**, 20130143 (2013).
20. Gong, J. *et al.* Early molecular events during onset of diapause in silkworm eggs revealed by transcriptome analysis. *Int. J. Mol. Sci.* **21**, 6180 (2020).

21. Kidokoro, T. & Masaki, S. Photoperiodic response in the relation to variable voltinism in the
ground cricket, *Pteronemobius fascipes* Walker (Orthoptera: Gryllidae). *Jpn. J. Ecol* **28**,
291–298 (1978).

22. Tanigawa, N., Matsumoto, K., Yasuyama, K., Numata, H. & Shiga, S. Early embryonic
development and diapause stage in the band-legged ground cricket *Dianemobius*
nigrofasciatus. *Dev. Genes Evol.* **219**, 589–596 (2009).

23. Shimizu, Y., Tamai, T. & Goto, S. G. Cell cycle regulator, small silencing RNA, and
segmentation patterning gene expression in relation to embryonic diapause in the band-
legged ground cricket. *Insect Biochem. Mol. Biol.* **102**, 75–83 (2018).

24. Blackmon, H., Ross, L. & Bachtrog, D. Sex determination, sex chromosomes, and karyotype
evolution in insects. *J. Hered.* **108**, 78–93 (2017).

25. Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M. BUSCO update:
Novel and streamlined workflows along with broader and deeper phylogenetic coverage for
scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.* **38**, 4647–4654
(2021).

26. Stanke, M., Diekhans, M., Baertsch, R. & Haussler, D. Using native and syntenically
mapped cDNA alignments to improve *de novo* gene finding. *Bioinformatics* **24**, 637–644
(2008).

27. Stanke, M., Schöffmann, O., Morgenstern, B. & Waack, S. Gene prediction in eukaryotes
with a generalized hidden Markov model that uses hints from external sources. *BMC*
Bioinformatics **7**, 62 (2006).

28. Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M. & Stanke, M. TSEBRA: transcript selector
for BRAKER. *BMC Bioinformatics* **22**, 566 (2021).

29. Kovaka, S. *et al.* Transcriptome assembly from long-read RNA-seq alignments with
StringTie2. *Genome Biol.* **20**, 278 (2019).

30. Keilwagen, J., Hartung, F. & Grau, J. GeMoMa: Homology-based gene prediction utilizing
intron position conservation and RNA-seq data. *Methods Mol. Biol.* **1962**, 161–177 (2019).

31. Ito, Y. *et al.* Chromosome-scale whole genome assembly and annotation of the Jamaican
field cricket *Gryllus assimilis*. *Sci. Data* **12**, 826 (2025).

32. Murata, K. *et al.* Complete mitochondrial genome sequences of two ground crickets,
Dianemobius fascipes nigrofasciatus and *Polionemobius taprobanensis* (Orthoptera:
Grylloidea: Trigonidiidae). *Mitochondrial DNA Part B* **8**, 1311–1315 (2023).

33. Sanno, R. *et al.* Comparative analysis of mitochondrial genomes in Gryllidea (Insecta:
Orthoptera): Implications for adaptive evolution in ant-loving crickets. *Genome Biol. Evol.*
13, evab222 (2021).

34. Kumar, L. & E Futschik, M. Mfuzz: a software package for soft clustering of microarray data.
Bioinformation **2**, 5–7 (2007).

35. Pelissier, A., Chauvin, J.-P. & Lecuit, T. Trafficking through Rab11 endosomes is required
for cellularization during *Drosophila* embryogenesis. *Curr. Biol.* **13**, 1848–1857 (2003).

36. Lecuit, T. Junctions and vesicular trafficking during *Drosophila* cellularization. *J. Cell Sci.*
117, 3427–3433 (2004).

37. Lefebvre, F. A. & Lécuyer, É. Flying the RNA nest: *Drosophila* reveals novel insights into the
transcriptome dynamics of early development. *J. Dev. Biol.* **6**, 5 (2018).

38. Kuhn, H., Sopko, R., Coughlin, M., Perrimon, N. & Mitchison, T. The Atg1-Tor pathway
regulates yolk catabolism in *Drosophila* embryos. *Development* **142**, 3869–3878 (2015).

39. Harrison, M. M. & Eisen, M. B. Transcriptional activation of the zygotic genome in
Drosophila. *Curr. Top. Dev. Biol.* **113**, 85–112 (2015).

40. Laver, J. D., Marsolais, A. J., Smibert, C. A. & Lipshitz, H. D. Regulation and function of
maternal gene products during the maternal-to-zygotic transition in *Drosophila*. *Curr. Top.*
Dev. Biol. **113**, 43–84 (2015).
41. Shi, J. *et al. Drosophila* Brahma complex remodels nucleosome organizations in multiple
aspects. *Nucleic Acids Res.* **42**, 9730–9739 (2014).
42. Möller, A., Avila, F. W., Erickson, J. W. & Jäckle, H. *Drosophila* BAP60 is an essential
component of the Brahma complex, required for gene activation and repression. *J. Mol. Biol.*
352, 329–337 (2005).
43. Papoulas, O. *et al.* The HMG-domain protein BAP111 is important for the function of the
BRM chromatin-remodeling complex *in vivo*. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 5728–5733
(2001).
44. Beck, Y., Pecasse, F. & Richards, G. Krüppel-homolog is essential for the coordination of
regulatory gene hierarchies in early *Drosophila* development. *Dev. Biol.* **268**, 64–75 (2004).
45. Mitchell, N., Cranna, N., Richardson, H. & Quinn, L. The ecdysone-inducible zinc-finger
transcription factor Crol regulates Wg transcription and cell cycle progression in *Drosophila*.
Development **135**, 2707–2716 (2008).
46. Erokhin, M. *et al.* Crol contributes to PRE-mediated repression and Polycomb group
proteins recruitment in *Drosophila*. *Nucleic Acids Res.* **51**, 6087–6100 (2023).
47. Carrera, P. *et al.* Tousled-like kinase functions with the chromatin assembly pathway
regulating nuclear divisions. *Genes Dev.* **17**, 2578–2590 (2003).
48. Chen, J. *et al.* Piwi regulates the usage of alternative transcription start sites in the
Drosophila ovary. *Nucleic Acids Res.* **53**, gkae1160 (2025).
49. Izadi, H. Endocrine and enzymatic shifts during insect diapause: a review of regulatory
mechanisms. *Front. Physiol.* **16**, 1544198 (2025).
50. Zhang, X.-S., Wang, Z.-H., Li, W.-S. & Xu, W.-H. FoxO induces pupal diapause by
decreasing TGFβ signaling. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2210404119 (2022).
51. Li, H.-Y., Lin, X.-W., Geng, S.-L. & Xu, W.-H. TGF-β and BMP signals regulate insect
diapause through Smad1-POU-TFAM pathway. *Biochim. Biophys. Acta Mol. Cell Res.* **1865**,
1239–1249 (2018).
52. Hamm, D. C. & Harrison, M. M. Regulatory principles governing the maternal-to-zygotic
transition: insights from Drosophila melanogaster. *Open Biol.* **8**, 180183 (2018).
53. Stadler, M. & Fire, A. Conserved translatome remodeling in nematode species executing a
shared developmental transition. *PLoS Genet.* **9**, e1003739 (2013).
54. Clouser, P. R., Riggs, C. L., Romney, A. L. T. & Podrabsky, J. E. Diapause and anoxia-
induced quiescence are unique states in embryos of the annual killifish, *Austrofundulus*
limnaeus. *Biomolecules* **15**, 515 (2025).
55. Steward, R. A., Pruisscher, P., Roberts, K. T. & Wheat, C. W. Genetic constraints in genes
exhibiting splicing plasticity in facultative diapause. *Heredity (Edinb.)* **132**, 142–155 (2024).
56. Li, B. *et al.* DNA methylation is correlated with gene expression during diapause termination
of early embryonic development in the silkworm (*Bombyx mori*). *Int. J. Mol. Sci.* **21**, 671
(2020).
57. Tian, Z. *et al.* The PBAP chromatin remodeling complex mediates summer diapause via
H3K4me3-driven juvenile hormone regulation in *Colaphellus bowringi*. *Proc. Natl. Acad. Sci.*
U. S. A. **122**, e2422328122 (2025).
58. Singh, P. P. *et al.* Evolution of diapause in the African turquoise killifish by remodeling the
ancient gene regulatory landscape. *Cell* **187**, 3338–3356 (2024).
59. Wrona, M. *et al.* The BRAHMA-associated SWI/SNF chromatin remodeling complex
controls *Arabidopsis* seed quality and physiology. *Plant Physiol.* **197**, kiae642 (2024).

60. Shimizu, Y. & Goto, S. G. No evidence for the involvement of juvenile hormone III and 20-
hydroxyecdysone in maternal decisions for embryonic diapause and diapause entry in the
band-legged ground cricket *Dianemobius nigrofasciatus* (Orthoptera: Trigonidiidae). *Appl.*
Entomol. Zool. (2023) doi:10.1007/s13355-023-00853-6.

61. Fukumoto, E., Numata, H. & Shiga, S. Effects of temperature of adults and eggs on the
induction of embryonic diapause in the band‑legged ground cricket, *Dianemobius*
nigrofasciatus. *Physiol. Entomol.* **31**, 211–217 (2006).

62. Goto, S. G. & Nagata, M. The circadian clock gene (Clock) regulates photoperiodic time
measurement and its downstream process determining maternal induction of embryonic
diapause in a cricket. *Eur. J. Entomol.* **119**, 12–22 (2022).

63. Fishilevich, E. *et al.* Use of chromatin remodeling ATPases as RNAi targets for parental
control of western corn rootworm (*Diabrotica virgifera virgifera*) and Neotropical brown stink
bug (*Euschistus heros*). *Insect Biochem. Mol. Biol.* **71**, 58–71 (2016).

64. Masaki, S. Thermal relations of diapause in the eggs of certain crickets (Orthoptera:
Gryllidae). *Bull. Fac. Agric. Hirosaki Univ.* **6**, 5–20 (1960).

65. Ptak, G. E. *et al.* Embryonic diapause is conserved across mammals. *PLoS One* **7**, e33027
(2012).

66. McGowen, M. R., Erez, O., Romero, R. & Wildman, D. E. The evolution of embryo
implantation. *Int. J. Dev. Biol.* **58**, 155–161 (2014).

67. Goto, S. G., Doi, K., Nakayama, S. & Numata, H. Maternal control of cold and desiccation
tolerance in eggs of the band-legged ground cricket *Dianemobius nigrofasciatus* in relation
to embryonic diapause. *Entomol. Res.* **38**, 17–23 (2008).

68. Kokot, M., Dlugosz, M. & Deorowicz, S. KMC 3: counting and manipulating k-mer statistics.
Bioinformatics **33**, 2759–2761 (2017).

69. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot
for reference-free profiling of polyploid genomes. *Nat. Commun.* **11**, 1432 (2020).

70. Johnston, J. S., Bernardini, A. & Hjelmen, C. E. Genome size estimation and quantitative
cytogenetics in insects. *Methods Mol. Biol.* **1858**, 15–26 (2019).

71. Zimin, A. V. *et al.* The MaSuRCA genome assembler. *Bioinformatics* **29**, 2669–2677 (2013).

72. Roach, M. J., Schmidt, S. A. & Borneman, A. R. Purge Haplotigs: Allelic contig
reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 460 (2018).

73. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

74. Laetsch, D. R. & Blaxter, M. L. BlobTools: Interrogation of genome assemblies. *F1000Res.*
6, (2017).

75. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421
(2009).

76. Putnam, N. H. *et al.* Chromosome-scale shotgun assembly using an *in vitro* method for
long-range linkage. *Genome Res.* **26**, 342–350 (2016).

77. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv [q-bio.GN] (2013).

78. Krzywinski, M. *et al.* Circos: an information aesthetic for comparative genomics. *Genome*
Res. **19**, 1639–1645 (2009).

79. Nishimura, O., Hara, Y. & Kuraku, S. Evaluating genome assemblies and gene models
using gVolante. *Methods Mol. Biol.* **1962**, 247–256 (2019).

80. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable
element families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457 (2020).

81. Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in
genomic sequences. *Curr. Protoc. Bioinformatics* **Chapter 4**, 4.10.1-4.10.14 (2009).
82. Gabriel, L. *et al.* BRAKER3: Fully automated genome annotation using RNA-seq and
protein evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res.* **34**, 769–
777 (2024).
83. Brůna, T., Lomsadze, A. & Borodovsky, M. GeneMark-ETP significantly improves the
accuracy of automatic annotation of large eukaryotic genomes. *Genome Res.* **34**, 757–768
(2024).
84. Pertea, G. & Pertea, M. GFF utilities: GffRead and GffCompare. *F1000Res.* **9**, 304 (2020).
85. Dossey, A. T. *et al.* Genome and genetic engineering of the house cricket (*Acheta*
domesticus): A resource for sustainable agriculture. *Biomolecules* **13**, 589 (2023).
86. Satoh, A., Takasu, M., Yano, K. & Terai, Y. *De novo* assembly and annotation of the
mangrove cricket genome. *BMC Res. Notes* **14**, 387 (2021).
87. Ylla, G. *et al.* Insights into the genomic evolution of insects from cricket genomes. *Commun*
Biol **4**, 733 (2021).
88. Wang, X. *et al.* The locust genome provides insight into swarm formation and long-distance
flight. *Nat. Commun.* **5**, 2957 (2014).
89. Kataoka, K. *et al.* The draft genome dataset of the Asian cricket *Teleogryllus occipitalis* for
molecular research toward entomophagy. *Front. Genet.* **11**, 470 (2020).
90. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).
91. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).
92. Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973
(2009).
93. Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for phylogenetic inference
in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).
94. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,
587–589 (2017).
95. Kumar, S. *et al.* TimeTree 5: An expanded resource for species divergence times. *Mol. Biol.*
Evol. **39**, msac174 (2022).
96. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and
bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).
97. Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Res.* **4**, 1521 (2015).
98. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
99. Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and
conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191–W198 (2019).
100. Jiang, H., Lei, R., Ding, S.-W. & Zhu, S. Skewer: a fast and accurate adapter trimmer for
next-generation sequencing paired-end reads. *BMC Bioinformatics* **15**, 182 (2014).
101. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**,
357–359 (2012).
102. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).
103. Quinlan, A. R. BEDTools: The Swiss-army tool for genome feature analysis: BEDTools: The
Swiss-army tool for genome feature analysis. *Curr. Protoc. Bioinformatics* **47**, 11.12.1-34
(2014).

104. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).
105. Chen, Y., Chen, L., Lun, A. T. L., Baldoni, P. L. & Smyth, G. K. edgeR v4: Powerful
differential analysis of sequencing data with expanded functionality and improved support
for small counts and larger datasets. *Nucleic Acids Res.* **53**, gkaf018 (2025).

Figure 1

Figure 2

Figure 3

Figure 4

a
Up-regulated and down-regulated pathways in short-day eggs

b
Brahma complex core subunits

c
Gluconeogenesis

d