Drugging the Medullary Thyroid Cancer Surfaceome with T Cell Engagers Targeting CEA, GFRA4 and DLL3 as Monotherapy and In Combination with Tyrosine Kinase Inhibitors

Tim Andrew Erickson

Abstract

Medullary thyroid cancer (MTC) is a rare form of thyroid cancer, and the definitive treatment is surgical resection. For patients who are not cured by surgery or for patients who present with distant metastatic disease, no curative therapy exists. T cell engagers (TCEs) are an emerging class of biologics that simultaneously bind to tumor cell surface antigens and the CD3 domain of T cells, thereby redirecting endogenous T cells to lyse tumor cells. TCEs can mediate durable tumor regression, including complete responses, and are now approved to treat multiple cancer types outside of MTC. RNASeq analysis of 30 MTC tumors, cross-referencing with public databases and an extensive literature review were used to identify CEA, DLL3 and GFRA4 as promising tumor antigens to target with TCEs. Expression of these antigens was then validated on MTC cell lines. Herein, I describe the discovery, application and development of MTC-targeted TCEs (MTCEs) recognizing the MTC tumor antigens CEA, DLL3 and GFRA4. *In* vitro, MTCEs mediate potent target-dependent and T cell-dependent cytotoxicity against MTC cell lines at concentrations of just 10 ng/mL. MTCEs also induce target-dependent IFNg secretion from T cells, with no measurable IFNg secretion observed in the presence of antigen-negative cell lines. Further, MTCEs are functionally compatible with the FDA-approved tyrosine kinase inhibitors selpercatinib and cabozantinib, and combination therapy numerically enhances cytotoxicity. These preclinical data provide strong rationale for continued development of MTCEs, which may one day revolutionize the treatment of metastatic MTC.

Introduction

Medullary thyroid cancer (MTC) is a relatively rare thyroid cancer that originates from the neuroendocrine parafollicular C-cells of the thyroid [1]. While localized MTC can sometimes be cured surgically, there is presently no cure for metastatic MTC [2]. While high-dose stereotactic body radiotherapy (SBRT) can provide local control of individual tumors [3] and Y-90 radioembolization can mediate durable control of liver metastases [4], there is no highly effective adjuvant therapy for MTC capable of eliminating all sites of metastatic disease. Unlike differentiated thyroid cancers, MTC does not concentrate radioactive iodine [5] and MTC is generally resistant to cytotoxic chemotherapy, as evidenced by low response rates and the transient nature of responses [6].

In the setting of progressive metastatic MTC, the daily oral tyrosine kinase inhibitors (selpercatinib, cabozantinib and vandetanib) can prolong survival for MTC patients harboring RET mutations, at the expense of side effects and cumulative toxicities [7-9]. In particular, selpercatinib treatment resulted in high response rates and a robust multi-year overall survival (OS) benefit in RET mutant MTC [9]. However, no randomized trial to date has demonstrated any OS benefit in RET wild-type patients [7]. Thus, there is a medical need to move the field forward and explore additional therapeutic strategies with the moonshot aim of curing metastatic MTC.

T cell engagers (TCEs) are an emerging class of biologics that simultaneously bind to tumor-associated surface antigens and to CD3 expressed on T cells, forming an immunological synapse, resulting in T cell activation and tumor cell lysis [10]. During the last decade, multiple TCEs targeting CD19, CD20 and

BCMA have been approved to treat hematological malignancies [11]. The 2024 FDA approval of the anti-DLL3 TCE tarlatamab in small cell lung cancer (SCLC) marked the first approval of a TCE in any solid tumor type [12]. In recent clinical trials, TCEs targeting STEAP1 and PSMA have shown robust antitumor activity in treatment-refractory prostate cancer, including durable responses and 99% PSA declines [13,14]. Further, TCEs targeting CEACAM5 (CEA) [15] and EGFR [16] have mediated objective responses in clinical trials in GI malignancies.

The successful application of TCEs in MTC would represent a new treatment paradigm, enabling the direct eradication of MTC tumor cells by endogenous T cells, in a tumor type which has historically proven resistant to immunotherapies, including immune checkpoint inhibitors and tumor-antigen vaccines [17,18]. While TCEs have been evaluated in multiple tumor types, PubMed and Google Scholar searches did not yield any publications evaluating TCEs in MTC, to date. Herein, the discovery, development and *in vitro* preclinical activity of TCEs targeting the MTC tumor antigens CEA, DLL3 and GFRA4 is described. These MTC-targeting TCEs are termed MTCEs. The identification of CEA, DLL3 and GFRA4 as addressable target antigens based on patient data is first discussed followed by target validation on MTC tumor cells lines using flow cytometry. Then, the development of MTCEs is described, and their independent binding to both T cells and target antigens is quantified using flow cytometry. Lastly, the *in vitro* activity of MTCEs alone and in combination with the FDA-approved kinase inhibitors cabozantinib and selpercatinib is characterized through live-cell imaging, cytotoxicity assays and measurement of MTCE-mediated IFNg secretion in tumor cell: T cell co-cultures.

RESULTS

Identification of Target Antigens for MTCEs

30 MTC patient tumors (26 primary; 4 tumor infiltrated lymph nodes (TILN)) were analyzed by bulk RNA sequencing, as previously described [19]. An analysis of the RNASeq dataset in conjunction with Human Protein Atlas curation and a literature review was used to identify target antigens for MTCEs. To conduct the analysis, the MTC patient dataset was truncated to exclude genes coding for intracellular proteins, leaving a set of 3,412 genes which code for the extracellular protein surfaceome [20]. MTC tumor surfaceome genes were then ranked by median expression level. Next, to identify genes specifically upregulated in MTC tumors, and to exclude genes with global elevated gene expression, the surfaceome of the TRON cell line database [21] was analyzed by dividing the expression value of each gene in the MTC TT cell line by the corresponding median expression value of all other 1,081 non-MTC cell lines in the TRON database (TT cell line/TRON cell line median). Based on the differential gene expression analysis, the top 40 relatively overexpressed TT surfaceome genes were selected for further analysis. Then analysis was undertaken to rank genes based on both high differential expression and high expression in the MTC patient dataset, noting that this approach filtered out non-MTC genes present in the MTC dataset, such as TPO and TSHR due to the presence of residual thyroid tissue.

The product of differential gene expression and median MTC patient expression was computed and candidate genes were ranked accordingly (Table 1). Lastly, each gene was annotated according to normal tissue expression in the Human Protein Atlas [22], in order to exclude genes with broad expression in normal tissues and simultaneous high expression in MTC, such as RET. An exception was made for CEACAM5 (CEA) because of the reported aberrant polarization (basal membrane expression) in tumor cells, which should theoretically limit on-target, off-tumor toxicity as T cells can not readily access the lumen-facing apical membrane, where CEA exhibits restricted expression in normal epithelial tissues

[23]. Per the aforementioned analysis, CEA, GFRA4, DLL3, GHSR, APLP1, PCDHA5 and PCDHA8 were identified as candidate target genes (Figure 1), as they showed high relative expression and differential expression in MTC and restricted normal tissue expression.

Figure 1: RNA expression in FPKM of candidate target antigens from a dataset comprised of 30 MTC patient tumors (26 primary tumor and 4 lymph-node infiltrated tumors). Paraffin-embedded tumor tissue blocks were used as input material.

As indicated in Figure 1, the well-established MTC tumor marker CEA [24] showed the highest expression, while the RET co-receptor GFRA4 (GDNF family receptor alpha-4) showed the highest differential gene expression in the TRON cell line database, in agreement with the previous reports indicating that GFRA4 expression is reported to be highly restricted to thyroid C cells and MTC [25]. Notably, both anti-CEA and anti-GFRA4 chimeric antigen receptor (CAR) T cells have demonstrated robust preclinical activity in MTC [26,27], and a clinical trial is currently evaluating anti-GFRA4 CAR-T cells in MTC patients [28]. Further, a clinical trial of the anti-CEA TCE cibisatamab in colon cancer mediated objective responses without dose limiting toxicities [15].

DLL3, GHSR, APLP1, PCDHA5 and PCDHA8, all show normal tissue expression restricted to the regions of the brain, which may preclude these antigens from being targeted due to the potential for neural toxicity. Surprisingly, DLL3 has proven to be a safe target [29], potentially due to aberrant extracellular trafficking in tumor cells [30]. Recent reports have indicated that DLL3 is abundantly expressed at both RNA and protein levels in aggressive metastatic MTC, which often harbors a transcriptomic signature (DLL3, SOX2, ASCL1) akin to SCLC [31]. Further, DLL3 has been detected in up to 100% of tumor cells in lymph node metastases by IHC [32]. Based on the totality of data, including high expression, relative overexpression, validated expression and indications of safety, CEA, DLL3 and GFRA4 were selected as target antigens from MTCEs. Of the remaining 4 candidate genes in the curated list, APLP1 shows the highest and most uniform expression and may be a suitable target although it was not selected for

further development due lack of protein expression data in the literature and RNA expression in the human brain.

Expression of CEA, DLL3 and GFRA4 on MTC Cell Lines

For proof-of-concept studies, flow cytometry was used to measure the expression of CEA, DLL3 and GFRA4 in the TT and MZ-CRC-1 (MZ) MTC cell lines and non-MTC cell lines (SHP-77, LNCaP, Raji, SKOV3, CHO-wt). The SCLC cell line SHP-77 shares a neuroendocrine origin with MTC and is known to express DLL3 and CEA [21], whereas the other cell lines are negative for all three targets. As shown in Figure 2, TT cells highly and broadly express CEA, DLL3 and GFRA4 with nearly 100% of cells expressing all three targets. In contrast, MZ cells show almost no detectable expression of GFRA4, moderate CEA (30%) expression and broad DLL3 expression. SHP-77 cells were found to broadly express DLL3 [33], and also show moderate CEA expression. The low, but detectable levels of GFRA4 on MZ and SHP-77 cell lines be may be a staining artifact from staining antibody cross-reactivity with another ligand, as SHP-77 cells are not reported to express GFRA4 at the RNA level [21]. CEA, DLL3 and GFRA4 were not detectable by flow on any of the other cell lines (data not shown).

Figure 2: Expression of CEA, DLL3 and GFRA4 on MZ (MTC), TT (MTC) and SHP-77 (SCLC) cell lines as measured by flow cytometry. TT cells show high expression of all three antigens, whereas MZ and SHP-77 cells do not express significant amounts of GFRA4.

Engineering and Synthesis of MTCEs.

Humanized monoclonal antibodies recognizing CEA and GFRA4 were generated through murine immunization, followed by AI-based affinity maturation and humanization by grafting the light chain and heavy chain CDRs onto homologous human antibody frameworks. Single chain variable fragment (scFv) targeting CEA and GFRA4 were then generated by linking the heavy and light chain variable regions of each monoclonal antibody using a (GGGGS)₃ linker. In order to generate MTCEs targeting CEA (anti-CEA) and GFRA4 (anti-GFRA4), each respective scFv was in turn linked to a proprietary anti-CD3 scFv/half-life extension domain. Specifically, the anti-CD3 domain was derived from a humanized scFv of the SP34 anti-CD3 antibody [34], and the half-life extension domain was derived from an Fc-silenced constant region of human IgG4. For proof-of-concept studies, tarlatamab was selected as the candidate anti-DLL3 MTCE because it is already FDA approved in SCLC and could be readily used to treat MTC patients off-label.

Anti-CEA and anti-GFRA4 MTCEs were expressed via transient transfection of CHO cells with plasmids encoding each MTCE followed by Protein A and size exclusion chromatography purification. Purity exceeded 90% as measured by SDS-PAGE and SEC-HPLC with yields of approximately 300 mg/L. The anti-DLL3 MTCE tarlatamab, cibisatamab, and gresonitamab were all purchased directly from a commercial vendor (MedChemExpress). All MTCEs were tested for endotoxin and confirmed to have levels of <0.1 EU/mg. The general structure of each MTCE, expressed as a continuous single-chain protein, is depicted in Figure 3.

Figure 3: Generalized structure of tarlatamab and anti-CEA and anti-GFRA4 MTCEs. Each TCE is comprised of an antigen binding domain, a CD3 binding domain and a Fc-based half-life extension domain.

Functional Binding of MTCEs to T Cells and Target Antigens

Flow cytometry-based assays were used to characterize the functional binding of MTCEs to T cells and tumor antigens. For T cell binding, T cells were first isolated from peripheral blood PBMCs using negative selection. Purity was checked by flow and found to exceed 97%. Purified T cells were then incubated

with varying concentrations (100 pg/mL to 10 ug/mL) of MTCEs for 20 minutes at room temperature, then washed and counterstained with FITC-conjugated CD8 mAb, PerCp-conjugated CD4 mAb and 1 ug/mL of PE-conjugated goat anti-human IgG antibody fragment, which binds to the Fc-domain of the MTCEs. Cells were washed to remove unbound antibodies, and flow data was acquired on a BD Accuri cytometer. As demonstrated in Figure 4, all MTCEs bind to CD8 T cells with similar affinity, and binding increases monotonically with MTCE concentration with detectable binding observed at concentrations as low as 10 ng/mL. Similar binding for each MTCE was observed for CD4 T cells, whereas no binding was observed for CD20+ B cells, CD14+ monocytes, or CD16+CD56+ NK cells (data not shown).

Figure 4: Binding of MTCEs to CD8 T Cells

For tumor antigen binding analysis, CHO cells were transduced with lentiviral vectors encoding human CEA (NP_004354.3) , DLL3 (NP_058637.1) or GFRA4 (NP_665705.1). Limiting dilution was used to select clonal populations with high and homogenous expression for each antigen. CHO-wt, CHO-CEA, CHO-DLL3 and CHO-GFRA4 cells were then incubated with varying concentrations (100 pg/mL to 10 ug/mL) of MTCEs for 20 minutes at room temperature, washed and then counterstained with PE-conjugated goat anti-human IgG, washed again and then analyzed on a BD Accuri flow cytometer. No detectable binding was observed for any MTCEs to CHO-wt cells and no cross-reactive binding was observed for MTCEs against non-target transduced CHO cells, i.e. binding of anti-CEA MTCE to GFRA4 transduced CHO cells. Figure 5 shows the on-target binding data for each MTCE, which was normalized to account for differences in target-antigen expression levels on CHO cells. Whereas anti-DLL3 and anti-CEA MTCEs showed similar binding affinity, with binding clearly detectable at concentrations as low as 10 ng/mL, there was an approximate log-shift in binding affinity for anti-GFRA4 where binding could only be clearly discerned at a concentration of 100 ng/mL.

Figure 5: Normalized binding of MTCEs to target-transduced CHO cells. The data shows binding of anti-GFRA4 MTCE to CHO-GFRA4 (green circles), binding of anti-CEA MTCE to CHO-CEA (black squares) and binding of anti-DLL3 MTCE to CHO-DLL3 (red triangles).

MTCEs Mediate Cytotoxicity Against MTC Cell Lines

MTCEs were capable of mediating potent cytotoxicity against MTC cells in the presence of T cells. The XTT cytotoxicity assay and endpoint live-cell imaging were used to assess MTCE-dependent and T-cell dependent cytotoxicity against MTC cell lines *in vitro.* Briefly, the XTT assay is based on the conversion of the tetrazolium salt XTT into an orange formazan product by metabolically active adherent tumor cells, and the colorimetric shift is quantified by absorption spectroscopy. Separate experiments were conducted to characterize the influence of both MTCE concentration and E:T ratio on cytotoxicity. Each experimental condition was run in triplicate using T cells from an otherwise healthy MTC patient donor with limited metastatic disease.

In the MTCE titration experiment, the E:T ratio was fixed at 1:1. Briefly, 15,000 eGFP-transduced MTC cells were plated overnight in each well of a 96-well flat bottom plate, and the following day, 15,000 T cells were added to each well along with varying concentrations of each MTCE. Cells were co-cultured for 48 hours, and then endpoint live-cell imaging was conducted on a fluorescence microscope before performing the XTT assay. The results in Figure 6 demonstrate how each MTCE mediated cytotoxicity against MZ and TT cells in a concentration-dependent manner. MTCEs targeting DLL3 and CEA exhibited potent cytotoxicity against both cell lines, whereas anti-GFRA4 was only effective against the TT cell line, consistent with the lack of expression of GFRA4 on the MZ cells. In general, anti-DLL3 (tarlatamab) mediated the broadest and most potent cytotoxicity across both cell lines with an approximate EC50 of ~2 ng/mL for TT cells and 0.2 ng/mL for MZ cells. Anti-CEA MTCE induced potent cytotoxicity in TT cells with an EC50 similar to tarlatamab, but showed reduced cytotoxicity in the MZ cell line, consistent with the reduced expression of CEA in the MZ cell line. While anti-GFRA4 induced potent cytotoxicity against TT cells at high concentrations, it had the highest EC50 value (EC50$_{TT}$ ~ 20 ng/mL and EC50$_{MZ}$ ~ 10 ug/mL, which may be a function of reduced binding affinity compared to anti-CEA and anti-DLL3 (Figure 5) or potentially lower GFRA4 protein expression relative to DLL3 and CEA.

Figure 6: Concentration-dependent cytotoxicity of MTCEs against the MZ and TT MTC cell lines at a 1:1 E:T ratio. Tumor cells were co-cultured with T cells for 48 hours in the presence of MTCEs as indicated, and then cytotoxicity was measured using the XTT assay.

For the T cell titration experiment, 15,000 tumor cells were again plated overnight in 96-well plates, and the following day T cells were added at E:T ratios ranging from 1:1 to 1:32 in the presence of 1 ug/mL of each MTCE. Both media only and the anti-CLDN18.2 TCE gresonitamab were used as negative controls, as CLDN18.2 is not expressed on MTC cell lines. After 24 hours, live-cell imaging was performed on an inverted fluorescence microscope for all TCEs at the 1:2 E:T ratio condition, in order to visually confirm cytotoxicity. As shown in Figure 7, all three MTCEs mediated potent cytotoxicity against TT cells as evidenced by reduced GFP surface area and clumping of GFP+ tumor cells. By contrast, T cells alone or T cells in the presence of gresonitamab did not mediate appreciable cytotoxicity, as the appearance is similar to TT cells in media only. For MZ cells, only tarlatamab induced high, visually discernible cytotoxicity, whereas anti-CEA cytotoxicity was discernible, but much less pronounced, and neither anti-GFRA4, gresonitamab nor T cells alone mediated visually detectable cytotoxicity.

Figure 7: Live-cell imaging reveals lysis of MTC cells mediated by T cells in the presence of MTCEs. All three MTCEs mediated observable cytotoxicity against the TT cell lines, whereas only anti-DLL3 and anti-CEA mediated observable clearance of MZ cells.

XTT assay results (Figure 8) demonstrate that MTCEs exhibited T cell-dependent cytotoxicity, with cytotoxicity beginning to fall off at E:T ratios of 1:4 and lower. Notably, tarlatamab demonstrated the highest and most consistent cytotoxicity at the lowest E:T ratio (1:32), and no detectable cytotoxicity was observed in the absence of MTCEs, nor in the absence of T cells or with gresonitamab at any E:T ratio (data not shown).

Figure 8: MTCE-mediated cytotoxicity requires T cells and depends on the effector to target (E:T) ratio.

MTCEs Mediate Cytotoxicity in a Target-Dependent Manner

To determine the impact of target antigen expression on cytotoxicity, a 48-hour XTT co-culture assay was conducted with 10 ug/mL of each MTCE at 1:1 E:T ratio (15,000 T cells, 15,000 tumor cells in a 96-well

plate) for 48-hours. Antigen-positive (MZ (CEA+, DLL3+), TT (triple positive), SHP-77 (CEA+, DLL3+), CHO (single positive)) and antigen-negative (LNCaP, SKOV3, Raji, CHO-wt) cell lines were used in the assay. Gresonitamab was used as a negative control, and the CEA-reactive TCE cibisatamab was used as a positive control for CEA+ cell lines and a negative control for CEA- cell lines. As shown in Figure 9, MTCEs exhibited cytotoxicity in an antigen-dependent manner. Notably, MTCEs only mediated cytotoxicity against CHO cells transduced to express their target antigen. Further, while both anti-CEA and cibisatamab mediated cytotoxicity against CEA+ cells lines, anti-CEA MTCE exhibited more potent cytotoxicity than cibisatamab in MZ cells, TT cells, SHP-77 cells and CHO-CEA cells. Critically, MTCEs did not mediate detectable cytotoxicity against antigen-negative cell lines at the highest E:T ratio and highest MTCE concentration tested, and no appreciable cytotoxicity was observed for cells exposed to gresonitamab.

Figure 9: MTCEs mediate cytotoxicity in a target-dependent manner with no off-target cytotoxicity observed in target antigen-negative cell lines.

Combining MTCEs with Tyrosine Kinase Inhibitors (TKIs) Increases Cytotoxicity

The RET-specific tyrosine kinase inhibitor selpercatinib and the multikinase inhibitor cabozantinib are both approved to treat metastatic MTC, and intriguingly both have been reported to enhance immunotherapy by reducing immunosuppression in the tumor microenvironment (TME) by upregulating MHC Class I, increasing T cell infiltration, and reducing the presence and function of immunosuppressive cellular subsets [35-37]. As TKIs may further potentiate the activity of MTCEs *in vivo* through several mechanisms, both cytotoxicity and IFNg secretion assays were used to assess the *in vitro* activity and compatibility of MTCEs with TKIs at clinically relevant concentrations of 5 uM for selpercatinib and 3 uM for cabozantinib [38,39]. To model a patient undergoing TKI treatment followed by MTCE therapy, 15,000 MZ and TT cells were plated overnight and then exposed to each TKIs (or media only) for 2 days, after which the media was refreshed with the same concentration of TKIs and then 7,500 T cells were added with the 1 ug/mL of each MTCE. The co-culture was then carried out for 48 hours, after which IFNg secretion and cytotoxicity were measured. As shown in Figure 10, the addition of selpercatinib or cabozantinib generally decreased IFNg secretion, but did not abrogate it entirely. This phenomenon may be a function of reduced tumor cell volume, which in turn my reduce the total surface area available for

T cell binding and activation. Interestingly, the reduction in the IFNg did not lead to reduced cytotoxicity. Rather, trends toward increased cytotoxicity were observed for all MTCEs in the presence of selpercatinib and cabozantinib. These data indicate that TKIs are likely functionally compatible with MTCEs, as they do not hinder cytotoxicity, and that TKIs may reduce unwanted side effects, such as cytokine release by reducing tumor surface area. Notably, while anti-GFRA4 MTCE was effective against the TT cell line, it was totally ineffective in the MZ cell line, as cytotoxicity was driven entirely by the addition of TKIs. This mirrors a situation where a single MTCE is used and tumors show resistance via antigen loss, highlighting the importance of targeting multiple antigens simultaneously.

Figure 10: Effect of Combining TKIs with MTCEs. TKIs moderate IFNg secretion, while numerically increasing cytotoxicity.

Combining Multiple MTCEs

The anti-GFRA4 resistant MZ cells were used as a model system of tumor resistance mediated by target-antigen loss. As a countermeasure against antigen escape, all three MTCEs were combined at a concentration of 333 ng/mL each vs. individual MTCEs at a concentration of 1 ug/mL. The concentration was adjusted to keep the total amount of MTCEs in the cell culture constant. For the experiment, 15,000 MZ cells were plated overnight and then co-cultured for 24 hours with 7,500 T cells in the presence of MTCEs as indicated. Cytotoxicity was measured via XTT assay. As shown in Figure 11, the combination of all three MTCEs at reduced doses could mediate cytotoxicity and potentially overcome tumor escape mediated by loss of a single target antigen. Interestingly, the triple combo yielded increased cytotoxicity over anti-CEA alone, which was likely driven by anti-DLL3, consistent with prior experiments showing that anti-DLL3 is the most potent MTCE against MZ cells. Triple MTCE treatment at reduced doses yielded comparable cytotoxicity to anti-DLL3 monotherapy at a higher dose, highlighting the feasibility of simultaneously targeting all three antigens to prevent antigen-loss escape.

Figure 11: MTCEs can be combined to prevent antigen-loss escape in the anti-GFRA4 resistant MZ cell line.

Methods

RNASeq and Gene Expression Data

MTC whole exome RNASeq data was acquired as previously described [19], and the entire dataset is fully disclosed in this manuscript for the first time. Tumor Cell line RNASeq data was downloaded from the TRON Cell Line Database (https://github.com/TRON-Bioinformatics/TCLP). Surfaceome genes were acquired from Table S2 in [20]. The Human Protein Atlas online portal (https://www.proteinatlas.org/) was used to evaluate protein expression and distribution in normal tissues. Analysis was performed using Excel software and all graphs were generated using Graphpad Prism software.

Synthesis of anti-CEA and anti-GFRA4 MTCEs

MTCEs were synthesized at GenScript using their proprietary TurboCHO™ transient transfection and purification process. Protein sequences including a designated signal peptide were disclosed to GenScript. The protein sequence was then codon optimized, and the corresponding gene was then cloned into pcDNA3.4 plasmids. After plasmid amplification and purification, CHO-S cells were transiently transfected with plasmids encoding each MTCE. After 5 days, the supernatant containing each MTCE was harvested and purified using Protein A chromatography followed by size exclusion chromatography followed by buffer exchange. Each MTCE was then molecularly characterized using SDS-Page and SEC-HPLC with both methods indicating purity exceeding 90% for anti-CEA MTCE and anti-GFRA4 MTCE. MTCE concentration was measured using A280 absorbance spectroscopy.

Tumor Cell Lines, T cells and Cell Culture

With the exception of the MZ cell line, all cell lines were purchased from ATCC. The MZ cell was kindly received as a gift from Dr. Jena French (UC-Denver). Tumor cell lines were generally cultured in RPMI (CHO, Raji, LNCaP), DMEM (MZ, SKOV3, SHP-77) or Ham's F12K (TT) media supplemented with 10% FCS and antibiotics. Cells were maintained with Normocin (InvivoGen) to prevent fungal and mycoplasma contamination. CHO-CEA, CHO-DLL3 and CHO-GFRA4 were generated by transducing CHO cells with lentiviral vectors encoding each target gene under control of an EF1-a promoter (Vectorbuilder). Cells

were transduced with polybrene (8 ug/mL, MOI=10) before single cell cloning through limiting dilution, in order to establish pure monoclonal population of 100% transduced cells.

T cells used in experiments were generated as follows. First, peripheral blood PBMCs were isolated from whole blood using Ficoll-Paque centrifugation. Then cells were cultured in T cell media (AIM-V supplemented with 5% FCS, antibiotics and 100 IU/mL of IL-2) for 2 days in GREX 24-well plates at a concentration of 2e6/cells in 4 mL of media. After 2 days, CD8 T cells or bulk T cells were isolated using "untouched" commercial negative selection kits (#130-096-495, #130-096-535) from Miltenyi Biotech, prior to being used in assays and T cell purity exceeding 97% was confirmed by flow cytometry.

Flow Cytometry

Flow staining antibodies were acquired from commercial vendors, as indicated in Table 2. For surface antigen staining of tumor cell lines, cells were gently detached using Accutase for 5-10 minutes, in order to avoid Trypsin-mediated protein cleavage [40]. Then cells were neutralized using ice-cold cell culture media. Cells were then spun down and resuspended in ice-cold media prior to staining 200,000 cells in 200 uL of media for 30 minutes at 4C. Staining antibodies were generally used at 1:100 dilution. After staining, cells were washed twice in cold FACS buffer, and then immediately analyzed on a BD Accuri flow cytometer.

IFNg Measurement

IFNg was measured using an ELISA kit from Biolegend (#430101) in accordance with the manufacturer's instructions. Prior to measurement, samples were diluted 1:5 in assay buffer, and this dilution factor was then corrected in the final results. ELISA plates were read on a Biotek EL808 plate reader at 450 and 650 nm.

Cell Co-culture and XTT Cytotoxicity Assays

For each assay, tumor cells and T cells were cultured in a 50/50 blend of T cell media and tumor cell media in the presence of MTCEs as indicated. Wells were run in triplicate. To mitigate cytotoxicity from alloreactivity, each well was supplemented with the anti-MHC Class I blocking antibody W6/32 (Biolegend) at a concentration of 40 ug/mL. After 24 or 48 hours, the plate was flicked to remove non-adherent cells (dead cells or T cells). The plate was then washed once with warm media, flicked again and then 100 uL of media was added to each well, after which 50 uL of XTT reagent was added to each well, in accordance with instructions provided in the assay kit (Biotium, #30007). After XTT reagent addition, cells were then cultured for an additional 8 hours. Then plates were read on a Biotek EL808 ELISA plate reader at 490 nm and 650 nm. For analysis, the background was subtracted from each well ($A_{490} - A_{650}$) and then denoted as A_{sample}. Then MTCE-mediated cytotoxicity was computed as follows. The T cells + media only condition ($A_{baseline}$) was used as a baseline. The MTCE-specific cytotoxicity was calculated as $[(A_{baseline} - A_{sample})/A_{baseline}] \times 100\%$. Under this calculation, when sample and baseline wells show equivalent absorbance ($A_{baseline} = A_{sample}$) due to wells harboring the same number of metabolically active tumor cells, cytotoxicity is zero. Conversely, when A_{sample} is zero due to elimination of all metabolically active cells, then cytotoxicity is 100%.

Discussion

The application of T cell engagers in MTC is a novel therapeutic strategy. MTCEs enable endogenous T cells to robustly lyse MTC cells expressing the targeted antigens CEA, DLL3 or GFRA4 *in vitro*. These antigens were identified by leveraging data from an MTC patient dataset in conjunction with differential gene expression analysis of large cell line database, and their expression was confirmed on MTC cell lines. While anti-CEA and anti-GFRA4 MTCEs are novel molecular entities, tarlatamab (anti-DLL3) is already FDA approved. Importantly, both anti-CEA and anti-GFRA exhibited robust *in vitro* activity at doses (~10 ng/mL) well below the average and trough concentrations of tarlatamab measured in clinical trials (C_{avg} = 1040 ng/mL and C_{trough} = 495 ng/mL) [40]. In terms of monotherapy, tarlatamab consistently generated the most consistent cytotoxicity across both cell lines, even at low concentrations, and is thus recommended for investigational off-label treatment in MTC patients whose tumors highly express DLL3.

While tarlatamab has demonstrated robust clinical activity, it is notable that the reported complete response rate in SCLC is only 2% [29]. As tumor heterogeneity and plasticity are hallmarks of solid tumors [42-44], it is likely that the rarity of complete responses could be due to loss or downregulation of DLL3 expression. Targeting more than a single antigen may be necessary for tumor eradication in the majority of MTC patients. Thus, the utilization of MTCEs targeting three independent antigens (CEA, DLL3, GFRA4) represents a compelling strategy to overcome antigen-loss escape [45]. Recent scRNA analysis has demonstrated that multiple MTC subtypes with distinct transcriptional programs can exist within the same tumor, indicating that MTC cells can exist in multiple states during the time-course of evolution from C cell, to differentiated tumor cells, to dedifferentiated and aggressive malignant cells [44]. The transcriptional changes can influence antigen expression, as observed in the MZ cell line, which shows minimal GFRA4 expression, moderate CEA expression and high DLL3 expression. However, it is notable that both MTC cell lines express a triple-negative population by flow, which could lead to tumor recurrence even under optimal MTCE therapy. It is unclear if antigen-expression is cell-cycle dependent, or if cells may temporally recover expression at later timepoints. Further analysis of this population by flow sorting, followed by re-culturing and epigenetic methylation analysis would be particularly illuminating.

A prerequisite for MTCEs to function is the presence of T cells in close proximity to tumor cells. While MTC is generally thought to be a cold tumor lacking significant T cell infiltration, T cells have been found in the majority of MTC primary tumors although data on distant metastases is lacking. However, T cell engagers are active in other immunologically cold tumor types, such as prostate cancer and pancreatic cancer, indicating that T cell may exist in sufficient quantities to generate an antitumor response. Notably, the lymph nodes are often the first site of metastases in MTC [46], and therefore lymph node tumors may be particularly amenable to MTCE therapy, as the tumor would inherently be in close proximity to T cells. Tertiary lymphoid structures could be another source of T cells, and it is possible that initial killing by sparse T cells could lead to inflammation and trafficking of additional T cells to the tumor site. Notably, MTCEs were observed to be functionally combinable with TKIs, which can relieve immunosuppression in the TME through a variety of mechanisms. These T cells could further be supported by costimulatory agonist antibodies and/or novel IL-2 analogs, in order to overcome the immunosuppressive TME. As MTC is a rare tumor type, there is still an incomplete picture on the immune composition at common distant metastatic sites, including the lungs, liver and bones. Data on both the immune infiltrate and target antigen expression in distant metastases would be particularly helpful in identifying patients who could respond to MTCE therapy. However, the observation that higher

tumor burden correlates with higher serum CEA indicates that many distant metastases retain CEA expression.

Our data indicates a more favorable ability to eradicate tumor cells at higher effector to target ratios. In the clinical setting, it is therefore likely that patients with minimal residual disease would have higher response and greater chances of a cure than patients with bulky metastatic disease. However, clinical data would be needed to confirm the veracity of this hypothesis. As there is no effective adjuvant therapy for MTC, MTCEs might represent a compelling treatment to eradicate residual metastatic disease and ultimately cure patients who would otherwise recur. This would be analogous to radioactive iodine therapy in differentiated thyroid cancer patients. For patients with bulky metastatic disease, our data indicates that MTCEs are functionally compatible with standard-of-care TKIs, and that the two treatment modalities may be synergistic in the clinic. Reduced tumor burden would correspond to fewer activated T cells, and therefore less cytokine release, which is a dose-limiting toxicity of T cell engagers. Additionally, combination therapy may further shift the equilibrium toward tumor eradication over tumor growth.

The proof-of-concept data presented herein demonstrate MTCEs as a novel off-the-shelf strategy for redirecting the immune system to eradicate MTC. Additional preclinical experiments to assess potential immunogenicity [47-48] and off-target binding of anti-GFRA4 and anti-CEA are warranted. Validation of target expression in distant metastases along with a comprehensive analysis of the tumor-immune microenvironment in such metastases would be helpful in identifying patients who are likely to benefit from MTCE therapy. These studies will be the subject of future preclinical efforts, with the ultimate aim of bringing MTCEs into the clinic and radically improving the treatment of metastatic MTC.

Surfaceome Genes	MTC Patient Average of Surface Gene	TT Cell Line/Tron Cell Line Average	Normal Tissue Expression
CEACAM5	350.6	20.8	*restricted in basal membrane, broad apical membrane
BCAM	91.4	29.0	broad
RET	54.3	98.2	broad
GAP43	47.0	16.2	brain and peripheral nerves
APLP1	33.7	17.6	brain
NFASC	28.4	14.4	broad
TPO	27.4	15.1	normal thyrocytes, non-MTC tissue from thyroid gland
PTPRN	23.1	63.2	brain, adrenals, endocrine cells
CASR	22.4	271.9	parathyroid, pancreatic endocrine cells
ABCC8	21.1	161.9	brain, pancreas, adrenals
TMPRSS6	16.1	16.1	liver, brain, pituitary, testis
GHSR	15.7	15.8	restricted to pituitary
GABRB3	15.2	13.3	broad
CAMK2B	14.7	19.5	broad
CNTN1	12.9	42.6	brain, adrenals, retina
GP2	12.0	28.3	pancreas
PCDHB11	10.5	14.2	broad
KDR (VEGFR2)	9.9	101.0	broad
KCNK9	8.3	47.5	brain, adrenals, pituitary
PCDHA5	8.2	23.7	brain
PTPRN2	7.3	48.4	brain, adrenal, pituitary, endocrine cells
GRM8	7.0	45.6	brain, eye
MRAP2	6.8	21.8	broad
GLRA1	6.8	13.0	brain, eye
LGR5	6.4	14.8	broad
PCDHA8	6.3	23.5	brain
SCN9A	6.3	25.3	broad
KCNK3	6.2	19.9	adrenals
PCDHA3	6.1	19.3	broad
SUSD4	5.9	16.1	broad
SEMA6D	5.8	16.8	broad
GFRA4	5.8	426.5	highly restricted to thyroid C cells
DLL3	5.5	20.6	brain
HCN1	5.3	40.4	broad
PTPRO	4.8	20.9	broad
SEZ6L2	4.7	12.3	broad
PCDHA10	4.6	34.8	broad
PCDHAC2	4.5	23.9	broad
SLC6A17	4.5	20.6	brain, pituitary, retina

Table 1: MTC tumor RNASeq analysis, differential gene expression analysis and normal tissue expression analysis used to identify MTCE target antigens

Antibody	Clone	Supplier	Catalog Number
CD8-FITC	HIT8a	Biolegend	500906
CD4-FITC	OKT4	Sony	2187040
Human IgG-PE	N/A	Jackson ImmunoResearch	109-116-170
CD3-PerCp	UCHT1	Biolegend	300428
CEA-APC	CI-P83-1	Santa Cruz Bio	sc-23928
DLL3-PE	tarlatamab	MedChemExpress	HY-P99575
GFRA4-PE	polyclonal	Saphire Bioscience	BS-0168R-PE
CD20-FITC	2H7	Biolegend	302203
CD16-FITC	3G8	Biolegend	980112
CD56-APC	QA17A16	Biolegend	985906
CD14-FITC	M5E2	Biolegend	982502

Table 2: Antibodies used in experiments

References

[1] Hazard JB, HAWK WA, CRILE JR GE. Medullary (solid) carcinoma of the thyroid—a clinicopathologic entity. The Journal of Clinical Endocrinology & Metabolism. 1959 Jan 1;19(1):152-61.

[2] Chen H, Roberts JR, Ball DW, Eisele DW, Baylin SB, Udelsman R, Bulkley GB. Effective long-term palliation of symptomatic, incurable metastatic medullary thyroid cancer by operative resection. Annals of surgery. 1998 Jun 1;227(6):887-95.

[3] Kukulska A, Krajewska J, Kołosza Z, Grządziel A, Gajek M, Paliczka-Cieślik E, Syguła D, Ficek K, Kluczewska-Gałka A, Jarząb B. Stereotactic radiotherapy is a useful treatment option for patients with medullary thyroid cancer. BMC Endocrine Disorders. 2021 Dec;21:1-6.

[4] Puleo L, Agate L, Bargellini I, Boni G, Piaggi P, Traino C, Depalo T, Lorenzoni G, Bianchi F, Volterrani D, Brogioni S. Yttrium-90 transarterial radioembolization for liver metastases from medullary thyroid cancer. European thyroid journal. 2022 Dec 1;11(6).

[5] Meijer JA, Bakker LE, Valk GD, De Herder WW, De Wilt JH, Netea-Maier RT, Schaper N, Fliers E, Lips PT, Plukker JT, Links TP. Radioactive iodine in the treatment of medullary thyroid carcinoma: a controlled multicenter study. European journal of endocrinology. 2013 May;168(5):779-86.

[6] Hadoux J, Pacini F, Tuttle RM, Schlumberger M. Management of advanced medullary thyroid cancer. The lancet Diabetes & endocrinology. 2016 Jan 1;4(1):64-71.

[7] Schoffski P, Elisei R, Müller S, Brose MS, Shah MH, Licitra LF, Jarzab B, Medvedev V, Kreissl M, Niederle B, Cohen EE. An international, double-blind, randomized, placebo-controlled phase III trial (EXAM) of cabozantinib (XL184) in medullary thyroid carcinoma (MTC) patients (pts) with documented RECIST progression at baseline.

[8] Wells Jr, Samuel A., et al. "Vandetanib in patients with locally advanced or metastatic medullary thyroid cancer: a randomized, double-blind phase III trial." *Journal of clinical oncology* 30.2 (2012): 134-141.

[9] Hadoux J, Elisei R, Brose MS, Hoff AO, Robinson BG, Gao M, Jarzab B, Isaev P, Kopeckova K, Wadsley J, Führer D. Phase 3 trial of selpercatinib in advanced RET-mutant medullary thyroid cancer. New England Journal of Medicine. 2023 Nov 16;389(20):1851-61.

[10] Huehls, Amelia M., Tiffany A. Coupet, and Charles L. Sentman. "Bispecific T-cell engagers for cancer immunotherapy." *Immunology and cell biology* 93.3 (2015): 290-296.

[11] Cech P, Skórka K, Dziki L, Giannopoulos K. T-Cell Engagers—The Structure and Functional Principle and Application in Hematological Malignancies. Cancers. 2024 Apr 20;16(8):1580.

[12] Alamgir I, Alamgir U, Alamgir E, Qureshi AA, Siddiqui MO, Jaber MH, Motwani J. Tarlatamab's FDA approval: shaping the future of cancer therapy. Annals of Medicine and Surgery. 2024 Oct 1;86(10):5676-9.

[13] Kelly WK, Danila DC, Lin CC, Lee JL, Matsubara N, Ward PJ, Armstrong AJ, Pook D, Kim M, Dorff TB, Fischer S. Xaluritamig, a STEAP1× CD3 XmAb 2+ 1 immune therapy for metastatic castration-resistant

prostate cancer: results from dose exploration in a first-in-human study. Cancer discovery. 2024 Jan 12;14(1):76-89.

[14] Clarke H. JANX007 demonstrates encouraging clinical activity in metastatic CRPC. Urology Times. 2025 Jan 1;53(1).

[15] Segal NH, Melero I, Moreno V, Steeghs N, Marabelle A, Rohrberg K, Rodriguez-Ruiz ME, Eder JP, Eng C, Manji GA, Waterkamp D. CEA-CD3 bispecific antibody cibisatamab with or without atezolizumab in patients with CEA-positive solid tumours: results of two multi-institutional Phase 1 trials. Nature Communications. 2024 May 15;15(1):4091.

[16] CytomX Therapeutics announces positive initial phase 1a dose escalation data for monotherapy CX-904 (EGFR/CD3 PROBODY T-cell engager). News release. CytomX Therapeutics, Inc. May 8, 2024. Accessed May 9, 2024. https://www.globenewswire.com/news-release/2024/05/08/2878255/37704/en/CytomX-Therapeutics-Announces-Positive-Initial-Phase-1a-Dose-Escalation-Data-for-Monotherapy-CX-904-EGFRxCD3-PROBODY-T-Cell-Engager.html

[17] Bilusic M, Heery CR, Arlen PM, Rauckhorst M, Apelian D, Tsang KY, Tucker JA, Jochems C, Schlom J, Gulley JL, Madan RA. Phase I trial of a recombinant yeast-CEA vaccine (GI-6207) in adults with metastatic CEA-expressing carcinoma. Cancer Immunology, Immunotherapy. 2014 Mar;63:225-34.

[18] Private communication with Dr. Ravi Madan, Principal Investigator or Phase II clinical trial evaluating pembrolizumab in MTC. (https://clinicalstudies.info.nih.gov/protocoldetails.aspx?id=17-C-0061&&query=Type%20Iv)

[19] Pozdeyev N, Erickson TA, Zhang L, Ellison K, Rivard CJ, Sams S, Hirsch FR, Haugen BR, French JD. Comprehensive immune profiling of medullary thyroid cancer. Thyroid. 2020 Sep 1;30(9):1263-79.

[20] Hu Z, Yuan J, Long M, Jiang J, Zhang Y, Zhang T, Xu M, Fan Y, Tanyi JL, Montone KT, Tavana O. The Cancer Surfaceome Atlas integrates genomic, functional and drug response data to identify actionable targets. Nature cancer. 2021 Dec;2(12):1406-22.

[21] Scholtalbers J, Boegel S, Bukur T, Byl M, Goerges S, Sorn P, Loewer M, Sahin U, Castle JC. TCLP: an online cancer cell line catalogue integrating HLA type, predicted neo-epitopes, virus and gene expression. Genome medicine. 2015 Dec;7:1-7.

[22] Pontén F, Jirström K, Uhlen M. The Human Protein Atlas—a tool for pathology. The Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland. 2008 Dec;216(4):387-93.

[23] Bacac M, Fauti T, Sam J, Colombetti S, Weinzierl T, Ouaret D, Bodmer W, Lehmann S, Hofer T, Hosse RJ, Moessner E. A novel carcinoembryonic antigen T-cell bispecific antibody (CEA TCB) for the treatment of solid tumors. Clinical Cancer Research. 2016 Jul 1;22(13):3286-97.

[24] Turkdogan S, Forest VI, Hier MP, Tamilia M, Florea A, Payne RJ. Carcinoembryonic antigen levels correlated with advanced disease in medullary thyroid cancer. Journal of Otolaryngology-Head & Neck Surgery. 2018 Jan;47(1):55.

[25] Lindahl M, Timmusk T, Rossi J, Saarma M, Airaksinen MS. Expression and alternative splicing of mouse Gfra4 suggest roles in endocrine cell development. Molecular and Cellular Neuroscience. 2000 Jun 1;15(6):522-33.

[26] Erickson TA, Shih YP, Fass J, Jang M, Tran E. T cells engineered to express immunoreceptors targeting the frequently expressed medullary thyroid cancer antigens calcitonin, CEA, and RET M918T. Thyroid. 2022 Jul 1;32(7):789-98.

[27] Bhoj VG, Li L, Parvathaneni K, Zhang Z, Kacir S, Arhontoulis D, Zhou K, McGettigan-Croce B, Nunez-Cruz S, Gulendran G, Boesteanu AC. Adoptive T cell immunotherapy for medullary thyroid carcinoma targeting GDNF family receptor alpha 4. Molecular Therapy-Oncolytics. 2021 Mar 26;20:387-98.

[28] https://clinicaltrials.gov/study/NCT04877613?term=gfra4&rank=1

[29] Paz-Ares LG, Felip E, Ahn MJ, Blackhall FH, Borghaei H, Cho BC, Johnson ML, Ramalingam SS, Reck M, Zhang A, Ju CH. Randomized phase 3 study of tarlatamab, a DLL3-targeting bispecific T-cell engager (BiTE), compared to standard of care in patients with relapsed small cell lung cancer (DeLLphi-304).

[30] Wang H, Zheng T, Xu D, Sun C, Huang D, Liu X. Targeting DLL3: Innovative Strategies for Tumor Treatment. Pharmaceutics. 2025 Apr 16;17(4):520.

[31] Ruz-Caracuel I, Caniego-Casas T, Alonso-Gordoa T, Carretero-Barrio I, Ariño-Palao C, Santón A, Rosas M, Pian H, Molina-Cerrillo J, Luengo P, Palacios J. Transcriptomic differences in medullary thyroid carcinoma according to grade. Endocrine Pathology. 2024 Sep;35(3):207-18.

[32] Ingenwerth M, Brandenburg T, Führer-Sakel D, Goetz M, Weber F, Dralle H, Schildhaus HU, Schmid KW, Theurer S. DLL3 (delta-like protein 3) expression correlates with stromal desmoplasia and lymph node metastases in medullary thyroid carcinomas. Endocrine connections. 2021 Mar 1;10(3):283-9.

[33] Giffin MJ, Cooke K, Lobenhofer EK, Estrada J, Zhan J, Deegen P, Thomas M, Murawsky CM, Werner J, Liu S, Lee F. AMG 757, a half-life extended, DLL3-targeted bispecific T-cell engager, shows high potency and sensitivity in preclinical models of small-cell lung cancer. Clinical Cancer Research. 2021 Mar 1;27(5):1526-37.

[34] Yang SY, Chouaib S, Dupont B. A common pathway for T lymphocyte activation involving both the CD3-Ti complex and CD2 sheep erythrocyte receptor determinants. Journal of immunology (Baltimore, Md.: 1950). 1986 Aug 15;137(4):1097-100.

[35] Kwilas AR, Ardiani A, Donahue RN, Aftab DT, Hodge JW. Dual effects of a targeted small-molecule inhibitor (cabozantinib) on immune-mediated killing of tumor cells and immune tumor microenvironment permissiveness when combined with a cancer vaccine. Journal of translational medicine. 2014 Dec;12:1-5.

[36] Oh CY, Klatt MG, Bourne C, Dao T, Dacek MM, Brea EJ, Mun SS, Chang AY, Korontsvit T, Scheinberg DA. ALK and RET inhibitors promote HLA class I antigen presentation and unmask new antigens within the tumor immunopeptidome. Cancer immunology research. 2019 Dec 1;7(12):1984-97.

[37] Ruan X, Yi J, Hu L, Zhi J, Zeng Y, Hou X, Huang J, Gu P, Hao W, Gao M, Pan Y. Reduced MHC class II expression in medullary thyroid cancer identifies patients with poor prognosis. Endocrine-Related Cancer. 2022 Feb 1;29(2):87-98.

[38] Lacy SA, Miles DR, Nguyen LT. Clinical pharmacokinetics and pharmacodynamics of cabozantinib. Clinical pharmacokinetics. 2017 May;56:477-91.

[39] https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/213246s008lbl.pdf

[40] Chen S, So EC, Strome SE, Zhang X. Impact of detachment methods on M2 macrophage phenotype and function. Journal of immunological methods. 2015 Nov 1;426:56-61.

[41] https://www.accessdata.fda.gov/drugsatfda_docs/nda/2024/761344Orig1s000MultidisciplineR.pdf

[42] Sayar E, Patel RA, Coleman IM, Roudier MP, Zhang A, Mustafi P, Low JY, Hanratty B, Ang LS, Bhatia V, Adil M. Reversible epigenetic alterations mediate PSMA expression heterogeneity in advanced metastatic prostate cancer. JCI insight. 2023 Apr 10;8(7):e162907.

[43] Yang D, Denny SK, Greenside PG, Chaikovsky AC, Brady JJ, Ouadah Y, Granja JM, Jahchan NS, Lim JS, Kwok S, Kong CS. Intertumoral heterogeneity in SCLC is influenced by the cell type of origin. Cancer discovery. 2018 Oct 1;8(10):1316-31.

[44] Wang LF, Zhou WW, Yuan F, Fu KW, He Y, Chen R. Medullary thyroid cancer: single-cell transcriptome and tumor evolution. Holistic Integrative Oncology. 2024 Aug 30;3(1):43.

[45] Gonzalez-Exposito R, Semiannikova M, Griffiths B, Khan K, Barber LJ, Woolston A, Spain G, von Loga K, Challoner B, Patel R, Ranes M. CEA expression heterogeneity and plasticity confer resistance to the CEA-targeting bispecific immunotherapy antibody cibisatamab (CEA-TCB) in patient-derived colorectal cancer organoids. Journal for immunotherapy of cancer. 2019 Dec;7:1-4.

[46] Scollo C, Baudin E, Travagli JP, Caillou B, Bellon N, Leboulleux S, Schlumberger M. Rationale for central and bilateral lymph node dissection in sporadic and hereditary medullary thyroid cancer. The Journal of Clinical Endocrinology & Metabolism. 2003 May 1;88(5):2070-5.

[47] Harris CT, Cohen S. Reducing immunogenicity by design: approaches to minimize immunogenicity of monoclonal antibodies. BioDrugs. 2024 Mar;38(2):205-26.

[48] Kroenke MA, Milton MN, Kumar S, Bame E, White JT. Immunogenicity risk assessment for multi-specific therapeutics. The AAPS journal. 2021 Nov 5;23(6):115.

Acknowledgements

The author would like to personally thank Dr. Jena French (CU-Anschutz Medical Center) for many helpful discussions and note that the MTC patient RNASeq data extensively analyzed for MTCE development *de novo* in this manuscript was generated during a prior study and was the subject of the publication titled, "Comprehensive immune profiling of medullary thyroid cancer" which is reference [19]. The author would also like to thank Dr. Eric Tran (Providence Earle A. Chiles Research Institute) for countless helpful discussions, being a wonderful mentor and allowing me to use his fluorescence microscope to procure images presented in this manuscript.