The dynamic evolution of panarthropod germ cell specification mechanisms

Jonchee A. Kao[1,2†], Emily L. Rivard[1,3†], Rishabh R. Kapoor[4], Cassandra G. Extavour[1,5,6]*

1. Department of Molecular and Cellular Biology, Harvard University, Cambridge MA, USA
2. Current address: Laboratory of Insect Physiology, School of Agriculture, Kyoto University, Kyoto, Japan
3. Current address: Department of Molecular Biology and Genetics, Cornell University, Ithaca NY, USA
4. Program for Systems, Synthetic and Quantitative Biology, Harvard University, Boston MA, USA
5. Department of Organismic and Evolutionary Biology, Harvard University, Cambridge MA, USA
6. Howard Hughes Medical Institute, Chevy Chase MD, USA

† equal contribution

* correspondence: extavour@oeb.harvard.edu

Abstract

Germ cells enable the reproduction of an organism and the continuity of its lineage.

Across animals, these crucial cells are segregated from the soma at different times and

places and via distinct mechanisms. Understanding the evolution of germ cell

specification across animals is complicated by the difficulty of making meaningful

comparisons of embryonic development between diverse animal species. Here, we

characterize germ cell specification in Panarthropoda, an ancient clade that

encompasses massive animal biodiversity, within which we can conduct meaningful

comparative embryology. We amass data from centuries of studies describing the

timing and mechanisms of germ cell formation, and apply ancestral state reconstruction

to these data to propose novel hypotheses about the trajectory of evolution in this

process. Furthermore, we speculate about the mechanisms underlying these

evolutionary dynamics by considering the relationships among germ cell specification,

concurrent developmental processes and the germ line gene network. Collectively, this

Review derives new insights from a rich historical database of embryological

observations, offering broad implications for understanding the evolution of metazoan

germ cells.

Keywords: germ plasm, pole cells, inductive signaling, BMP signaling pathway,

cytoplasmic determinant

Introduction

During animal development, the germ line comprises a collection of cells specified early in embryonic development that later give rise to the next generation via eggs or sperm. Whereas historical researchers relied primarily on observation of unperturbed embryos to understand germ cell specification and development (e.g. histological analysis of fixed, sectioned embryos; Bounoure, 1939; Wolff, 1962), we can now investigate these processes using functional genetic analysis. This allows us not only to identify germ cells with more certainty, but also to gain more insight into the molecular mechanisms of germ cell specification. In this Review, we have used "specification" to mean the process that imbues embryonic cells with the ultimate fate of generating eggs and sperm when the animal is sexually mature, and "differentiation" to mean the processes that make germ cells distinct from somatic cells in one or more of their morphology, behavior, or gene expression. Several hypotheses have been proposed to explain evolutionary transitions between the mechanisms underlying specification and differentiation across animals, including potential changes in cis-regulatory mechanisms, changes in evolvability or body plan diversity, or impacts on germ line mutation rate and ultimately fitness (Buss, 1982; Buss, 1983; Crother et al., 2007; Crother et al., 2016; Evans et al., 2014; Extavour, 2007; Johnson et al., 2011; Whittle and Extavour, 2016; Whittle and Extavour, 2017). However, such hypotheses are difficult to test because embryonic development of animals is too different across vast evolutionary time scales to be meaningfully compared. Moreover, there is disagreement in the field about whether existing well-studied mechanisms of specification (discussed below) are truly the only mechanisms, and whether they are

discrete, mutually exclusive mechanisms. Furthermore, there are conflicting reports

about the impact of germ cell specification mode on the evolution of genomes, body

plans and fitness (Evans et al., 2014; Otto and Hastings, 1998; Queller, 2000; Whittle

and Extavour, 2016; Whittle and Extavour, 2017). Here, we examine variation in germ

cell specification across the most diverse clade of animals, the Panarthropoda, wherein

the general features of embryogenesis can nevertheless be meaningfully compared.

This clade displays striking diversity in the temporal and spatial origin of germ cells, as

well as multiple transitions between specification mechanisms, offering an ideal system

to study evolution and regulation of germ cell specification.

Here, we first review the existing observational data regarding embryonic germ

line development across panarthropods. We are primarily interested in the first

embryonic appearance of germ cells before their arrival in the gonad, during which time

we refer to them as primordial germ cells (PGCs). We identify three categories of

temporally distinct developmental context in which PGCs are first detectable during

embryogenesis, and we analyze the phylogenetic distribution of these three categories.

We then summarize perturbation-based experimental studies aimed at uncovering the

mechanisms of germ cell specification. Finally, we discuss the implications of these data

on our understanding of the evolution of germ cell specification in panarthropods, and

more broadly across Metazoa.

Identification of primordial germ cells

Historically, embryologists studied early development using histological

preparations of sectioned embryos (His, 1868). With light microscopy, PGCs can be

readily distinguished from surrounding somatic cells by their rounder and larger nuclei

(Nieuwkoop and Sutasurya, 1979). The chromatin of germ cells often appears more

diffuse than that of somatic cells, and the cytoplasm, especially cortical or perinuclear

cytoplasm, may contain electron-dense, non-membrane bound granular material (Eddy,

1975). However, tracking these cells throughout development using fixed samples and

morphological features alone is not always technically possible (reviewed in XXXX).

Since the late 20^{th} century, studies assessing gene expression patterns unique to

or diagnostic of germ cells (henceforth referred to as molecular markers) have been

instrumental in identifying germ cells at earlier stages than previously recognized. The

most commonly used markers are the mRNA transcript or protein products (detectable

by *in situ* hybridization or antibody staining respectively) of the genes *vasa*, *nanos*, *tudor*

and *piwi* (Ewen-Campen et al., 2010). The use of molecular markers can enable germ

cell identification at earlier developmental stages, since some germ cells express the

genes that encode these molecules before their morphological differentiation becomes

unambiguously detectable (e.g. Ewen-Campen et al., 2013a). Additionally, these

markers facilitate tracking of germ cells over the course of development. However,

using molecular markers is a biased approach, as it relies on previous knowledge of

germ line gene expression from other species. Furthermore, interpretation of molecular

marker data can be complicated by pleiotropy, as the genes that encode these

molecules may also be expressed by somatic cells (e.g. Yajima and Wessel, 2011).

Germin cell formation in panarthropods

In our review of the embryological literature from 1860 to the present, we

identified three categories of PGC specification in panarthropods, based on the timing of

specification during embryogenesis (Fig. 1). We describe these categories below, along

with relevant details of embryogenesis for context.

Category 1: PGC specification at early cleavage and blastoderm stages

After fertilization, embryogenesis begins with cleavage, defined as a series of

mitotic divisions of the zygote (Gilbert, 1997). In most arthropods, cleavage leads to the

formation of the *blastoderm*, a relatively uniform layer of cells at the egg cortex,

surrounding a central yolk mass populated by a small number of yolk cells (Anderson,

1973; Scholtz and Wolff, 2013). The first category of PGC specification in

panarthropods occurs during or before the formation of the blastoderm (Fig. 1A).

In insects

In most insects, the blastoderm is initially a syncytium, which is a single cell

containing multiple nuclei generated by incomplete cleavage division cycles that do not

include cytokinesis (Anderson, 1973; Johannsen and Butt, 1941). In some insects, the

PGCs are the first cells to form, appearing at the posterior pole before the rest of the

blastoderm is cellularized (Fig. 1A1), and therefore often referred to as pole cells. While

pole cells are most commonly seen in species of Diptera (flies, midges and mosquitoes),

Coleoptera (beetles), and Hymenoptera (bees, ants and wasps), they have also been

reported in Thysanoptera (thrips) and Dermaptera (earwigs) (references in Table S1).

Pole cells were first reported in a species of the midge *Chironomus* by Weismann

(1863), although Metschnikow (1866) was the first to report their migration into the

gonad. In many insects, the cytoplasm at the posterior pole of the embryo is

morphologically distinct from that of the rest of the embryo, often with visible granules

(Hegner, 1914; Zissler, 1992). The close association of this granular cytoplasm with

pole cells led researchers to speculate that the granular material contains germ line

determinants (e.g. Hegner, 1909a; Zissler, 1992). It was long believed that pole cells

gave rise to somatic cell types in addition to the germ cells (Anderson, 1962a; Anderson,

1962b; Poulson, 1947; Poulson, 1950). However, careful lineage tracing in the fruit fly

Drosophila melanogaster showed that all pole cells are determined as germ cells, and

any that become lost during their migration die (Underwood et al., 1980).

While the function of pole cells as PGCs appears conserved across insects, the

specific details of their formation can vary considerably across species. For example,

pole cells differ in which cleavage cycle they cellularize: they form during the ninth

mitotic cycle of *D. melanogaster* (Huettner, 1923) and during the sixth mitotic cycle in

the parasitic wasp *Nasonia vitripennis* (Buchta et al., 2013). To our knowledge, the

earliest observed formation of pole cells is during the third cleavage cycle in the midges

Miastor metraloas and *Chironomus sp.* (Hegner, 1914; Weismann, 1863). The number

of pole cells formed is also variable: a single pole cell forms in *Chironomus* embryos

(Hegner, 1914; Weismann, 1863), whereas four appear in the fruit fly *Bactrocera tryoni*

(Anderson, 1962a) and sixteen pole cells form in the beetle *Calligrapha multipunctata*

(Hegner, 1909b).

After the pole cells form, they enter the egg and migrate to the location of the

future embryonic gonad, which is quite far from the posterior pole (Santos and Lehmann,

2004). Several different migration pathways have been described. In *D. melanogaster*

and many other Diptera, the pole cells are carried into the egg during posterior midgut

invagination, then migrate through the epithelium of the posterior midgut to enter the

abdominal mesoderm and associate with the developing somatic gonad (Metschnikow,

1866; Sonnenblick, 1941). In Hymenoptera, pole cells are internalized with the rest of

the mesoderm during gastrulation (Gatenby, 1917). Yet another variation is observed in

Chrysomelidae (leaf beetles) (Hegner, 1909a), where germ cells are internalized with

the rest of the germ band as the extraembryonic membranes fold over the germ band.

In other arthropods

PGCs are also specified early in Collembola (springtails), a clade of hexapods

that are sister to insects (Du et al., 2024). Embryogenesis in these animals begins with

a period of holoblastic cleavage, and certain blastomeres on the basal (yolk) side of the

blastoderm can be recognized as likely PGCs very early in development (Jura, 1967;

Klag, 1982). The cytoplasm of these cells contains granular material visible by light and

electron microscopy (Klag, 1982), resembling the cytoplasm containing germ line

determinants observed in insects (Hegner, 1914; Zissler, 1992). Rather than

maintaining contact with the germ band as in insects, collembolan germ cells migrate

into the center of the yolk after their specification until eventually migrating to associate

with the abdominal mesoderm, from which the embryonic gonad develops (Jura, 1967).

Among non-hexapod Pancrustacea, embryos of Copepoda (copepods),

Amphipoda (beach hoppers), Decapoda (crabs, lobsters and shrimp), and some

Cladocera (water fleas) undergo stereotypical *holoblastic cleavage*, allowing precise

tracking of cell lineages and the identification of a specific cleavage blastomere as the

progenitor of PGCs (Gerberding et al., 2002; Hertzler, 2005; Kühn, 1913; McClendon,

1907). In amphipods and decapods, a putative germ plasm has been identified in the

freshly laid egg, termed the RNA-containing body (Gupta and Extavour, 2013) or

intracellular body (Biffis et al., 2009; Chen et al., 2021). During cleavage, this region of

cytoplasm is inherited asymmetrically by the blastomeres that give rise to the PGCs

(Gupta and Extavour, 2013) (Fig. 1A2).

In chelicerates (spiders, scorpions, ticks and mites), candidate PGCs are first

detectable with molecular markers in the yolk of blastoderm-stage embryos in the mite

Tetranychus urticae (Dearden et al., 2003). As the germ band forms, the PGCs

associate with the posterior germ band near the primordia of the fourth walking legs

(Dearden et al., 2003). We are not aware of any other chelicerate or myriapod

(centipedes and millipedes) species in which PGC specification has been proposed to

occur before gastrulation.

Category 2: PGC specification at gastrulation

After formation of the blastoderm, a portion of the blastoderm is specified as the

presumptive embryonic region, variably referred to as the *germ band*, *germ rudiment*,

embryonic rudiment or *germ disc* depending on its initial form (Anderson, 1973;

Johannsen and Butt, 1941). The rest of the blastoderm differentiates into one or more

extraembryonic membranes (Johannsen and Butt, 1941). Soon after this specification,

gastrulation occurs, whereby some of the blastoderm cells are internalized and become

the mesoderm (Stern, 2004). In insects, mesoderm internalization primarily occurs

along the ventral midline of the germ band (Leptin, 2004; Roth, 2004). In other

arthropods, mesoderm may be internalized through a *blastopore* near the germ

rudiment (Chaw et al., 2007; Gerberding and Patel, 2004), or by epiboly (Edgar et al.,

2015). The second category of PGC differentiation occurs at or very close to the time of

gastrulation (Fig. 1B).

In insects

Among insects, PGC specification during gastrulation is the most commonly

reported timing in members of Hemiptera (true bugs), Psocodea (lice) and Lepidoptera

(moths and butterflies), although it is also reported in species of Dermaptera, Blattodea

(cockroaches), Hymenoptera, Megaloptera (alderflies), Strepsiptera (twisted-wing flies),

Coleoptera, Mecoptera (scorpion flies) and Diptera. Three different spatial patterns of

germ cell specification have been reported around the time of gastrulation: before germ

band formation, at the posterior of the germ band after its formation, or at the ventral

midline of the blastoderm just before gastrulation. In Hemiptera, Psocodea and

Tenebrionidae (darkling beetles; Coleoptera), germ cells are first detectable at the

ventral posterior region of the cellular blastoderm in an invaginating structure called the

posterior pit (Fig. 1B1) (Butt, 1949; Goss, 1952; Mellanby, 1935; Schröder, 2006; Shinji,

1919). These cells appear to form on the side of the blastoderm epithelium facing the

yolk, rather than on the outside of the blastoderm, such as the pole cells described in

Category 1. In Hemiptera and Psocodea, the invagination continues into the egg

towards the anterior, forming a definitive germ band with germ cells at its posterior tip

(Fig. 1B1). Germ cells in tenebrionid beetles move further inside the embryo once the

amnion (ventral extraembryonic membrane) encloses the entire ventral surface of the

germ band, similar to the internalization seen in coleopteran species that specify their

germ cells as pole cells (e.g. the weevils *Callosobruchus maculatus* (Brauer, 1925;

Brauer, 1949; Quan, 2018) and *Acanthoscelides obtectus* (Jung, 1966; Lynch et al.,

2011)). The newly formed germ cells then migrate along the dorsal side of the germ

band toward the anterior, eventually settling into the somatic gonad in the abdominal

mesoderm.

In several species that appear to specify germ cells around the time of

gastrulation, PGCs were first identified at the posterior tip of the germ band after its

detachment from the blastoderm [*e.g., Periplaneta orientalis* (Blattodea) (Heymons,

1895), *Sialis mitsuhashii* (Megaloptera) (Suzuki et al., 1981), *Nematus ribesii*

(Hymenoptera) (Singh, 1967), and Mecoptera (Suzuki, 1990)]. Early studies of

tenebrionid beetles described this pattern of germ cell differentiation as well (Rempel

and Church, 1969; Ullmann, 1964). However, later work using molecular markers

suggests that PGCs in at least one such beetle, *Tribolium castaneum*, appear earlier,

during the cellular blastoderm stage in a posterior pit much like the Hemiptera (Schröder,

2006). This raises the possibility that in some other tenebrionid species, PGCs may

likewise differentiate from the blastoderm, with the reported later appearance in the

germ band simply reflecting limitations in detection methods.

In contrast to the posterior pit, germ cells differentiate along the ventral midline of

the blastoderm in Lepidoptera, separating from the blastoderm before gastrulation to lie

on the dorsal side of the presumptive germ band in the presumptive abdominal region

(Ando and Tanaka, 1980; Miya, 1958a; Presser and Rutschky, 1957; Tanaka, 1987;

Woodworth, 1889). Unlike when germ cells form at the posterior pit or at the posterior

tip of the germ band, in Lepidoptera there does not appear to be long-range germ cell

migration, since the germ cells differentiate in the same abdominal location as the

gonad.

In other panarthropods

Among non-insect Pancrustacea, PGCs are the first cells to be internalized from

the blastopore lip in Mysida (opossum shrimp) (Manton, 1928), Anomopoda (water

fleas) (Cannon, 1921), and the copepod orders Siphonostomatoida (sea lice)

(McClendon, 1907), Harpacticoida (Witschi, 1934) and Cyclopoida (Schimkewitsch,

1896). In Isopoda (woodlice and pill bugs), PGCs differentiate from the mesendoderm

soon after internalization (Fig. 1B2) (Goodrich, 1939; Needham, 1942; Strömberg, 1965;

Wolff, 2009). Thus, PGC differentiation seems to be tied to gastrulation in these

crustacean orders.

In chelicerates, PGCs have been identified based on morphological criteria at the

blastopore in Opiliones (harvestmen) (Fig. 1B3) (Faussek, 1888; Faussek, 1892; Holm,

1947) and Scorpiones (Brauer, 1894), and a recent study using molecular markers

supported this interpretation (Gainett et al., 2022). Putative PGCs have also been

identified at the blastopore in the centipedes *Scolopendra cingulata* and *Strigamia*

maritima (Myriapoda) (Green and Akam, 2014; Heymons, 1901).

In Tardigrada (water bears), germ cells are the first cells to enter the blastopore

during gastrulation, based on data from both morphological characteristics (Hejnol and

Schnabel, 2005) and molecular markers (Heikes et al., 2023) (Fig. 1B4). Lineage

tracing in these embryos suggests that the germ cells are derived from both

blastomeres of the two-cell embryo (Gabriel et al., 2007; Hejnol and Schnabel, 2005).

Finally, onychophoran (velvet worm) germ cells have been described only with

classic histological methods, to our knowledge (Fig. 1B5). They emerge either at the

blastopore lip during gastrulation (Manton, 1949) or at the posterior end of the germ

band during elongation (Manton, 1949; Mayer and Tait, 2009; Sedgwick, 1887).

Category 3: PGC specification from mesoderm

After gastrulation, the germ band elongates and becomes segmented, and the

mesoderm becomes organized into pairs of somites in each segment. In most living

arthropods, each somite forms a small *coelom* enclosing a *coelomic cavity* (Anderson,

1973; Johannsen and Butt, 1941). The coelomic cavities are eventually lost as the

mesoderm differentiates into different cell types and reorganizes into the primordia of

the internal organ systems, including the somatic gonad (Anderson, 1973; Johannsen

and Butt, 1941). The final category of PGC differentiation timing encompasses species

in which PGCs differentiate from the mesoderm at this comparatively late stage of

embryogenesis (Fig. 1C).

In insects

In insects, PGC specification from the mesoderm has been reported in members

of Archaeognatha (jumping bristletails), Zygentoma (silverfish and firebrats), Orthoptera

(crickets and grasshoppers) (Fig. 1C1), Mantodea (praying mantises), Blattodea,

Trichoptera (caddisflies), Coleoptera and Lepidoptera. PGCs may differentiate before

formation of the coelomic cavities [e.g., Orthoptera (Ewen-Campen et al., 2013b),

Mantodea (Görg, 1959)], after formation of the coelomic cavities [e.g., Archaeognatha

(Larink, 1969), Zygentoma (Woodland, 1957)], or after the coelomic cavities are no

longer recognizable [e.g., Trichoptera (Miyakawa, 1974), *Odontotaenius disjunctus*

(Coleoptera) (Krause, 1947), *Apis mellifera* (Hymenoptera) (Dearden, 2006; Nelson,

1915)]. In this developmental timing category, PGCs are usually specified in the same

abdominal segments where the somatic gonad forms, so there is no long-range

movement of PGCs after their specification.

In other arthropods

Differentiation of PGCs from the mesoderm has also been described in

Leptostraca (mud shrimp) (Figure 1C2) (Manton, 1934), Anaspidacea (mountain

shrimp) (Hickman, 1936) and Branchiopoda (fairy, clam and tadpole shrimp, and water

fleas) (Anderson, 1967; Mitsumoto and Makioka, 2002; Sudler, 1899). However, data

for stages of crustacean development late enough to include differentiation of the trunk

mesoderm from which primordial germ cells might arise are sparse. This relative

scarcity of early embryonic descriptions may be in part because many crustaceans

hatch as a nauplius, comprising only the head segments (Scholtz, 2000). In these

animals, germ cells might not be specified until later larval stages, after formation of the

trunk. Many studies of naupliar and larval development in crustaceans are focused on

organogenesis of the gonad as a whole, without specific reference to the origin of PGCs

(Cannon, 1924; Ikuta and Makioka, 1994; Mitsumoto and Makioka, 2002).

Classical studies on myriapods suggest a late differentiation of PGCs from the

mesoderm in three of the four myriapod classes [Pauropoda (Tiegs, 1947), Symphyla

(pseudocentipedes) (Tiegs, 1940) and Diplopoda (millipedes) (Dohle, 1964)], although

only one species was reported from each class in these studies. Germ cells in these

myriapod classes have not been investigated with molecular markers to our knowledge.

Among chelicerates, PGCs differentiate from the mesoderm of the opisthosomal

segments in spiders (Fig. 1C3) (Gough, 1902; Kautzsch, 1910; Schimkewitsch, 1903),

consistent with data from molecular markers are (Schwager et al., 2015). In

pycnogonids (sea spiders), PGCs are first detectable in the mesoderm of the prosoma

(Alexeeva and Tamberg, 2021; Alexeeva et al., 2018; Miyazaki and Makioka, 2012).

Summary: evolutionary trends from late to early germ cell specification

In his monograph on embryogenesis in Dermaptera and Orthoptera, Richard

Heymons (1895) proposed that 'the formation of sex cells at the rear end of the

embryonic primordium, which has now been clearly demonstrated for representatives of

very different groups, is probably generalizable for all insects'. Rather than making

broad statements such as this, we summarize the data regarding the timing of germ cell

specification (Figs. 2–3; Table S1) in the context of currently accepted panarthropod

phylogeny (Ballesteros et al., 2022; Fernández et al., 2018; Giribet and Edgecombe,

2019; Misof et al., 2014; Schwentner et al., 2017). This shows that in euarthropod

orders that are close to the base of the tree, PGCs tend to be first observed later in

embryogenesis (discussed above as Category 3). This pattern suggests that the timing

of first germ cell appearance evolved to be earlier in development in multiple lineages

independently, as seen in Dermaptera, Condylognatha (true bugs and thrips) and

Holometabola (undergoing complete metamorphosis) within insects; Copepoda,

Malacostraca (the largest class of crustaceans) and Cladocera within non-hexapod

Pancrustacea; Chilopoda (centipedes) within myriapods; and Opiliones, scorpions and

mites within chelicerates. These data suggest the possibility that differentiation from the

mesoderm is the ancestral mode of PGC specification in Euarthropoda.

Conflicting interpretations in the literature about the timing of germ cell

specification have sometimes resulted from technical advances enabling identification of

germ cells at stages earlier than previously detected. For example, early work on the

silk moth *Bombyx mori* (Lepidoptera) reported PGCs emerging from the mesoderm of

coelomic cavities in late embryogenesis (discussed above as Category 3) (Toyama,

1902), while later studies identified PGCs on the posterior side of the ventral midline of

the early germ band (as in Category 2) (Miya, 1953). The onset of PGC specification

before gastrulation is further supported by data from molecular markers, which showed

accumulation of *vasa* mRNA in a cluster of presumptive PGCs at the ventral midline of

the early germ band (Nakao, 1999; Nakao et al., 2006). Therefore, it is possible that

species which we classified here in Category 3, may be reclassified in Categories 1 or 2

by future work deploying molecular markers. This caveat notwithstanding, we note that

molecular markers and functional genetic studies often corroborate late germ cell

specification [see for example *Gryllus bimaculatus* (Orthoptera, Insecta) (Ewen-Campen

et al., 2013b), *A. mellifera* (Hymenoptera, Insecta) (Dearden, 2006), *Parasteatoda*

tepidariorum (Araneae, Chelicerata) (Schwager et al., 2015)]. We, therefore, do not

assume that the hypothesis of ancestral late PGC specification is an artifact of exclusive

use of morphological identification methods.

Going forward, data from key understudied lineages should be gathered to

further test our hypothesis about the ancestral timing of PGC specification. For example,

among the hexapods, data from members of the Palaeoptera (mayflies, dragonflies and

damselflies), the ametabolous Archaeognatha and Zygentoma, or the non-insect

hexapod Diplura (two-pronged bristletails) could be especially useful. Similarly, data

about PGC specification are missing from Oligostraca, an understudied clade of

Pancrustacea that includes Ostracoda (seed shrimp) and Branchiura (fish lice).

Outside of Pancrustacea, the field has yet to achieve clear consensus on the

phylogenetic relationships of the groups within Chelicerata (Sharma and Gavish-Regev,

2025), and there is too little embryological data available for us to suggest evolutionary

trends in PGC specification within this subphylum. Data regarding PGC specification are

even sparser for myriapods. Specifically, we are aware of relevant data from only five

species across the four myriapod classes (Dohle, 1964; Green and Akam, 2014;

Heymons, 1901; Tiegs, 1940; Tiegs, 1947). Studies of PGCs in a broader range of both

chelicerates and myriapods, as well as detailed embryological studies in unexplored

taxa, will therefore be needed to fully understand the developmental context and

evolutionary trends of germ cell specification in these subphyla.

Molecular and genetic mechanisms of germ cell specification in panarthropods

Beyond studying where and when germ cells first appear during arthropod

embryogenesis, researchers have sought to understand how they are specified. Germ

cell specification in animals has historically been classified into two mechanisms, often

referred to as maternal inheritance and zygotic induction (Extavour and Akam, 2003;

Nieuwkoop and Sutasurya, 1981). In maternal inheritance, germ cell determinants

accumulate in a specific region of the oocyte cytoplasm, forming a germ plasm. Cells

that inherit sufficient germ plasm components after the embryo divides become PGCs.

In embryos that undergo zygotic induction, all cells are in principle competent to

become germ cells at the beginning of embryogenesis. Certain cells later receive

signals from neighboring cells, produced by the activity of the zygotic genome, which

induce them to take on germ cell identity. In the following section, we describe how

historical observational data and perturbational experimental approaches have offered

increasingly robust evidence to distinguish between these mechanisms in several

panarthropod species.

Germ cell specification through maternal inheritance

In the early twentieth century, researchers speculated that the electron-dense,

membraneless granules frequently found in the cytoplasm of PGCs were responsible for

their specification. Experiments in chrysomelid beetles provided early evidence that the

posterior cytoplasm of the oocyte or early embryo contains germ cell determinants.

Burning or puncturing the posterior poles of beetle eggs with needles to damage or

remove posterior cytoplasm reduced or eliminated PGCs (Hegner, 1908; Hegner, 1911),

demonstrating that this cytoplasm does not simply correlate with PGC specification, but

rather plays a causal role in PGC specification. Studies in *D. melanogaster* offered

further evidence for the necessity of the germ plasm in PGC specification. Removing or

destroying the posterior cytoplasm of *D. melanogaster* eggs via ultraviolet irradiation

demonstrated that this region is required for germ cell formation (Geigy, 1931).

Subsequent centrifugation experiments in *D. melanogaster* embryos revealed that

shifting the granular structures in posterior cytoplasm away from the posterior pole

prevented germ cell formation or led to germ cells forming in a different part of the

embryo (Jazdowska-Zagrodzińska, 1966).

Experiments transplanting specific regions of cytoplasm between *D.*

melanogaster embryos showed that the contents of the posterior cytoplasm were

sufficient to specify germ cells. Transplanting posterior cytoplasm from undamaged *D.*

melanogaster embryos into sterile ones in which the posterior pole had been damaged

with ultraviolet irradiation rescued fertility, while transplanting anterior cytoplasm did not

(Okada et al., 1974). Further transplantation experiments demonstrated that posterior

pole plasm introduced into the anterior of another embryo caused formation of ectopic

pole cells that gave rise to functional germ cells when transplanted again to the

posterior of a third, sterile embryo (Illmensee and Mahowald, 1974). Similar

experiments transferring the posterior cytoplasm from late-stage oocytes into embryos

revealed that the oocyte pole plasm could specify functional PGCs (Illmensee et al.,

1976), demonstrating that the necessary germ cell determinants in *D. melanogaster* are

deposited during oogenesis. These foundational studies showed that maternally derived

cytoplasmic granules are both necessary and sufficient for germ cell formation in some

species.

Contemporary studies testing for maternal inheritance often ask whether known

germ line gene products accumulate during late oogenesis or early embryogenesis in

the region where PGCs first arise. For example, antibody staining or *in situ* hybridization

experiments to detect germ line molecules including Vasa protein or *nanos* mRNA have

suggested the presence of germ plasm in mosquitoes (Juhn and James, 2006; Juhn et

al., 2008), scuttle flies (Wotton et al., 2014), ants (Khila and Abouheif, 2008; Lynch et al.,

2011; Rafiqi et al., 2020), wasps (Kharel et al., 2024; Lynch et al., 2011), weevils (Lynch

et al., 2011), aphids (Chang et al., 2006; Chang et al., 2007; Chen et al., 2009; Lin et al.,

2014), amphipods (Gupta and Extavour, 2013) and prawns (Chen et al., 2021).

Transcriptomic analyses of different portions of embryos and oocytes (Carter et al.,

2015; Klomp et al., 2015; Quan and Lynch, 2016; Yoon et al., 2019) have also revealed

asymmetric localization of molecules associated with PGC specification.

However, expression data showing asymmetric localization of putative germ

plasm components are not sufficient to demonstrate that these molecules function in

PGC specification. To support such claims, functional genetic techniques are needed to

manipulate gene activity and test the effect on germ plasm assembly and germ line

development. The function of germ plasm components in panarthropods has been most

thoroughly studied in *D. melanogaster* (Lehmann, 2016; Mahowald, 2001; Trcek and

Lehmann, 2019). The membraneless ribonucleoprotein granules at the posterior of *D.*

melanogaster oocytes and early embryos are nucleated by the protein Oskar, which is

both necessary and sufficient for germ plasm assembly (Ephrussi and Lehmann, 1992;

Kistler et al., 2018). Oskar-mediated localization of core germ line molecules such as

Vasa, Tudor, Aubergine and *nanos* to germ granules is essential for PGC specification

and maintenance of germ cell identity during cell migration and gonad formation in

these fruit flies (Lehmann, 2016; Mahowald, 2001).

Analysis of germ plasm composition and localization in species beyond *D.*

melanogaster has revealed both conserved and rapidly evolving features. Within the

Drosophila genus, germ granule size, morphology (Counce, 1963; Mahowald, 1968;

Rivard et al., 2025), and the relative proportions of different granule components vary

widely (Doyle et al., 2023). The wasp *N. vitripennis* also has a well-characterized germ

plasm in the form of a single large structure called the oosome (Bull, 1982). Akin to *D.*

melanogaster, N. vitripennis requires *oskar* to assemble a germ plasm containing many

of the same molecules (e.g., Vasa, Tudor, and *nanos*) (Lynch and Desplan, 2010;

Lynch et al., 2011). However, while *D. melanogaster* has many small germ granules

(Mahowald, 1962), the *N. vitripennis* oosome has a distinct protein organization and

transcript composition compared to that observed in *D. melanogaster* (Kemph et al.,

2025; Kharel et al., 2024; Lynch et al., 2011). In several crustacean species, the germ

plasm appears as a large, round granule in the cortex of embryos and contains

conserved germ plasm markers (Chen et al., 2021; Grattan et al., 2013; Gupta and

Extavour, 2013). Studying germ plasm in additional species is needed to clarify whether

and how variation in germ plasm makeup and organization influences the post-

transcriptional regulation and function of germ line determinants.

Germ cell specification through zygotic induction

Experimentally demonstrating inductive germ cell specification, requiring physical,

molecular or genetic perturbation methods, is more technically challenging than

detecting germ plasm accumulation, for which in situ hybridization techniques or protein

detection may suffice. Consequently, most studies have inferred or hypothesized

zygotic induction based on absence of evidence of germ plasm. For example, *in situ*

hybridization experiments fail to reveal asymmetric localization of putative germ line

gene transcripts in oocytes and early embryos of various species, including milkweed

bugs (Ewen-Campen et al., 2013a), crickets (Ewen-Campen et al., 2013b), honeybees

(Dearden, 2006), mites (Dearden et al., 2003), spiders (Schwager et al., 2015),

centipedes (Green and Akam, 2014) and tardigrades (Heikes et al., 2023). Instead, key

germ line genes are expressed in a subset of cells later in embryogenesis, coincident

with reports of the first PGCs based on morphological criteria (Table S1). However, we

note that even in the well-accepted case of mouse PGC induction (Saitou and Yamaji,

2010), it remains formally possible that some unknown, and therefore untested, germ

line determinant or factor that makes cells competent to respond to later germ line

induction signals, is asymmetrically localized and preferentially inherited by germ cells.

This caveat notwithstanding, there are multiple examples of experimental support

for induction rather than inheritance across panarthropods. Maternal knockdown of

putative germ plasm components in milkweed bugs and crickets reveals that the

maternal products of these genes are dispensable for PGC formation (Ewen-Campen et

al., 2013a; Ewen-Campen et al., 2013b). Laser ablation of single cells in 2- and 4-cell

stage embryos of the tardigrade *Thuliana stephaniae* does not prevent PGC

specification (Hejnol and Schnabel, 2005). This observation suggests that this species

lacks a cytoplasmic region with inherited germ line determinants, and that it might

instead specify PGCs through inductive pathways.

Germ cell formation in embryos with tissue duplications can also be interpreted

as evidence against germ plasm and consistent with induction. For example, in the

drain fly *Clogmia albipunctata* and the flour beetle *T. castaneum*, genetic manipulations

producing double abdomen embryos lead to germ cell formation at both anterior and

posterior ends (Ansari et al., 2018; Yoon et al., 2019). Similarly, naturally occurring

"twins" of the stick insect *Carausius morosus* are duplicated embryos fused along their

anterior–posterior (A–P) axis, which both possess germ cells (Cavallin, 1971). In

contrast, double abdomen embryos of *D. melanogaster* only produce germ cells at the

true posterior end, since germ plasm is absent at the ectopic posterior end (Lasko and

Ashburner, 1990).

Among panarthropods, experiments in the cricket *G. bimaculatus* have provided

the most direct evidence for zygotic induction of PGCs. Knockdown of the mesoderm

specification factor *twist* leads to loss of PGCs, suggesting that the mesoderm

contributes to germ cell specification and/or maintenance (Ewen-Campen et al., 2013b)

and consistent with morphological evidence that orthopteran PGCs first arise from the

embryonic mesoderm (Wheeler, 1893). The bone morphogenetic protein (BMP)

signaling pathway, which induces germ cells in mice (Hopf et al., 2011; Lawson et al.,

1999; Ying and Zhao, 2001; Ying et al., 2000), is also required for cricket germ cell

specification. Specifically, BMP signaling and the downstream transcription factor Blimp-

1 induce a subset of mesodermal cells to become PGCs in crickets (Donoughe et al.,

2014; Nakamura and Extavour, 2016).

The shared use of both BMP and Blimp-1 in mice and crickets may have

convergently evolved, or could suggest that this signaling mechanism is ancestral in

bilaterian inductive PGC specification (Lochab and Extavour, 2017). Determining

whether BMP and its downstream target genes are required for germ line specification

in more species will be necessary to distinguish between these possibilities. Only a

small number of signaling pathways contribute to developmental fate decisions in

animals (Pires-daSilva and Sommer, 2003), increasing the likelihood of evolutionary

convergence and limiting the number of candidates to evaluate for roles in PGC

induction in different species.

Intermediate and flexible strategies of germ cell specification

Evidence from some animal species suggests that there may not be a strict

dichotomy between inheritance and induction (discussed by Seervai and Wessel, 2013).

These two mechanisms might instead represent ends of a spectrum that are not

necessarily mutually exclusive. Below, we discuss cases of panarthropod species for

which data suggest that their mechanisms of germ cell specification may fall

somewhere along this continuum.

In the silk moth *B. mori*, historical studies have generated conflicting evidence

about how germ cells are specified. Germ granules were not detected in early light

microscopy experiments, but germ cells failed to form after cauterization of the ventral

side of cleavage stage embryos, suggesting this region could contain germ cell

determinants (Miya, 1958a). Subsequent studies found that Vasa protein is not

asymmetrically localized before cellularization, again suggesting a lack of germ plasm,

or at least a germ plasm lacking Vasa protein (Nakao et al., 2006). However, the mRNA

of one of the four *B. mori nanos* homologs, *nanosO*, localizes along the ventral midline

of syncytial embryos immediately following egg-laying, and is inherited by the PGCs

(Nakao et al., 2008) suggesting the possibility that a germ plasm containing *nanosO*

transcript specifies PGCs. Intriguingly, researchers found that maternal and zygotic

nanosO act redundantly in PGC specification; loss of one or the other reduces but does

not completely eliminate germ cells, whereas loss of both prevents germ cell formation

(Nakao and Takasu, 2019). The possibility that both inheritance and induction

mechanisms contribute to *B. mori* PGC specification suggests that this species could

offer insight into transitions in germ cell specification mechanisms within Lepidoptera or

across insects more broadly.

Although in *D. melanogaster*, PGC specification is traditionally characterized as

driven by germ plasm, zygotic signaling is also suggested to contribute to germ cell

specification in this species (Colonnetta et al., 2022). Loss-of-function mutations in the

BMP family morphogen Dpp lead to partial loss of transcriptional quiescence and

mislocalization of germ plasm in the newly formed pole cells (Colonnetta et al., 2022).

Furthermore, loss of BMP signaling leads to activation of the normally suppressed

terminal signaling pathway in pole cells, which in turn disrupts their migration

(Colonnetta et al., 2022). These observations illustrate a universal conceptual challenge

in the field of developmental biology: distinguishing between, or even consistently

defining, cell type specification and maintenance. Although pole cells can form upon

disruption of BMP signaling, they do not contain the proper molecular components to

act as PGCs (Colonnetta et al., 2022). We suggest that it may be most useful to define

PGC specification mechanisms as those that ensure both the physical formation of

germ cell precursors (including adoption of specific morphologies and activation of

characteristic gene regulatory states), as well as the competence of those cells to take

the next developmental steps toward joining the gonad. Under this definition, we would

indeed concede that both germ plasm and zygotic inductive mechanisms play roles in *D.*

melanogaster germ cell specification.

The ability to post-embryonically regenerate a germ line initially specified via

maternal inheritance has also been reported in multiple invertebrate species, including

the sea squirt *Ciona intestinalis* (Takamura et al., 2002), the earthworm *Enchytraeus*

japonensis (Tadokoro et al., 2006) and the marine worm *Capitella teleta* (Dannenberg

and Seaver, 2018) (see also Özpolat, 2024 for a review of this phenomenon in

Annelida). The wasp *Pimpla turionellae* (Hymenoptera) has an oosome (Meng, 1968)

that is required for germ cell specification. Damaging the oosome results in no pole cells,

which are the PGCs of this species (Achtelig and Krause, 1971). However, oosome-

deficient embryos lacking pole cells develop into larvae with germ cells in their gonad,

indicating that an additional mechanism must regenerate germ cells later in

development (Achtelig and Krause, 1971). Evidence of germ line regeneration has also

been reported in the amphipod *Parhyale hawaiensis* (Kaczmarczyk, 2014), which has a

germ plasm that is asymmetrically segregated to the *g* micromere and its lineage

(Extavour, 2005; Gerberding et al., 2002; Gupta and Extavour, 2013). Ablation of the *g*

micromere eliminates PGCs in the embryo (Extavour, 2005), but the germ line is

replaced post-embryonically with germ cells that appear to be derived from the

mesoderm (Kaczmarczyk, 2014). The authors of this study speculated that these

replacement germ cells specifically derive from cells of the somatic gonad

(Kaczmarczyk, 2014), reminiscent of a previously popular theory that primordial germ

cells in most animals derived from the inner epithelium of the somatic gonad (Everett,

1944; Hargitt, 1919; Heys, 1931). Species capable of regenerating PGCs present the

opportunity to investigate the relative contributions of and potential interactions between

inheritance- and induction-based mechanisms within a single species. Furthermore,

these species may represent living examples of the intermediates that are hypothesized

to exist in evolutionary transitions between modes of germ line specification (Extavour,

2007).

Evolutionary drivers of panarthropod germ cell specification

In the following section, we discuss the evolutionary dynamics of germ cell specification

mechanisms across Panarthropoda, and propose potential developmental and genetic

factors that might contribute to these dynamics.

Convergent evolution of panarthropod germ plasm

The phylogenetic distribution of germ cell specification mechanisms across

panarthropods reveals a complex evolutionary history. Data for tardigrades, chelicerates

and myriapods are consistent with induction as the ancestral mechanism of

panarthropod primordial germ cell specification (Figure 3), and this hypothesis is

strongly supported by statistical tests described below (Figs. S1-S2). In Pancrustacea,

the most heavily represented group in our dataset (Table S1), the late origin of

primordial germ cells in several species suggests induction may be widely used.

However, there appear to be independent origins of germ plasm in Malacostraca and

Branchiopoda, a hypothesis that is also supported by formal statistical testing described

below (Figs. S1-S2; Table S3). The use of induction and germ plasm in different

decapod species suggests that further sampling could reveal patterns in the evolution of

germ cell formation within this order (Biffis et al., 2009; Chen et al., 2021; Terao, 1929).

However, functional experiments and broader taxonomic sampling will be necessary to

confirm mechanisms and reconstruct their evolutionary trajectories across

Panarthropoda as a whole.

Reminiscent of a broader pattern that we and others previously reported across

animals (Extavour, 2007; Extavour and Akam, 2003; Johnson et al., 2003; Whittle and

Extavour, 2017), in some insect orders we noted that lineages branching closer to the

base of a clade deployed induction, while later branching lineages appeared to use

germ plasm (Figs. 2–3). This pattern is well exemplified in Coleoptera, (Kobayashi et al.,

2006; Komatsu and Kobayashi, 2012; Krause, 1947; Niikura et al., 2017; Schröder,

2006; Ullmann, 1964), with pole cells forming only in Chrysomeloidea (including long-

horned and leaf beetles) and Curculionidae (Brauer, 1925; Hegner, 1908; Lynch et al.,

2011). These observations raise the possibility that germ plasm may not be ancestral to

Coleoptera. When considered alongside the absence of germ plasm in most

Lepidoptera, the reports of germ cells that first arise in the mesoderm (Category 3) in

Trichoptera (Miyakawa, 1974) and of germ cells specified at gastrulation (Category 2) in

Mecoptera (Suzuki, 1990), and the distinct morphologies of *N. vitripennis* and *D.*

melanogaster germ plasm (see *Molecular and genetic mechanisms of germ cell*

specification in panarthropods above), are consistent with a novel hypothesis: germ

plasm evolved independently multiple times across insects, rather than arising only

once in the Holometabola as we and others have previously proposed (Ewen-Campen

et al., 2012; Ewen-Campen et al., 2013b; Lynch et al., 2011). Our previous work

suggested the independent evolution of germ plasm at the phylum level (Extavour and

Akam, 2003), but here, we hypothesize widespread convergent evolution of maternal

inheritance-based germ cell specification within a phylum.

To test this hypothesis, we performed ancestral character reconstruction using

both maximum parsimony and maximum likelihood approaches (details in

Supplementary Information, Tables S2-S7 Figs. S1-S2). Both a parsimony analysis (Fig.

S1) and marginal ancestral reconstructions of a maximum likelihood approach (Fig. S2;

see Table S7 for marginal probabilities) provided strong statistical support for induction

as the ancestral specification mechanism in Panarthropoda, Arthropoda and Insecta,

with multiple independent instances of germ plasm evolution and reversion to induction

(see Table S3 for a comparison of the two approaches).

We further find strong evidence that induction was the ancestral specification

mechanism in Holometabola (probability 80.6%), followed by independent transitions to

germ plasm in the common ancestors of Diptera and Mecoptera (or independently

within Mecoptera, Siphonaptera (fleas) and Diptera); derived lineages of Lepidoptera, in

the common ancestor of Chrysomelidae and Curculionidae (Coleoptera); and in the

common ancestor of Hymenoptera. Outside of Holometabola, we infer independent

transitions to germ plasm in Thysanoptera, Hemiptera, Dermaptera, Collembola (which

may have undergone multiple independent transitions within the class), Decapoda, the

last common ancestor of Hyalidae and Talitridae (Amphipoda), and in the chelicerate

Tetranychus urticae. Collectively, these qualitative and quantitative analyses suggest a

previously unappreciated dynamism in the evolutionary history of germ cell specification

within Panarthropoda.

The relationship between germ cells and somatic germ layers

In arthropods with late formation of PGCs (Category 3), these cells appear to

differentiate directly from the mesoderm (Evans, 1901; Hickman, 1936; Nelson, 1915).

Consistent with this hypothesis, abolishing mesoderm prevents germ cell specification

in the cricket *G. bimaculatus* (Ewen-Campen et al., 2013b). Germ cell differentiation

occurs concurrently with the differentiation of many other mesodermal cell types,

including the somatic gonad, musculature, heart and fat body (Johannsen and Butt,

1941). We suggest that the types of mechanisms that specify germ cells are not distinct

in principle from those specifying any other mesodermal cell type in these animals.

The germ layer origin of germ cells is less clear in arthropods whose germ cells

appear before or very close to the time of gastrulation (Category 2). For example, PGCs

in many crustaceans are described as being the first to enter the blastopore during

gastrulation, and are usually considered by researchers to be derived from the

mesendoderm (e.g. Cannon, 1921; Goodrich, 1939; Manton, 1928). In contrast, germ

cells in Category 2 insects arise before gastrulation (Mellanby, 1935; Ullmann, 1964),

consistent with the hypothesis that germ cells are distinct from the mesoderm and may

not derive from any germ layer.

Crustaceans with putative germ plasm typically have stereotypical embryonic

cleavage patterns and embryonic cell lineages show that the germ cell lineage is sister

to the mesoderm or the mesendoderm [e.g. Amphipoda (Extavour, 2005; Gerberding et

al., 2002; Scholtz and Wolff, 2002; Wolff and Scholtz, 2002), Euphausiacea (krill)

(Alwes and Scholtz, 2014), Decapoda (Hertzler, 2005)], but never to the ectoderm. In

addition, removing the putative germ plasm from embryos of the amphipod *P.*

hawaiensis makes the *g* micromere morphologically indistinguishable from the other

micromeres, which give rise to mesoderm or endoderm in unperturbed embryos (Gupta

and Extavour, 2013). This result suggests that in the absence of germ cell determinants,

the *g* micromere, normally the germ line progenitor, could instead adopt

mesendodermal fate. This hypothesis has not been directly tested in this amphipod

because gastrulation fails after ablation of the putative germ plasm (Gupta and Extavour,

2013). Together, these data suggest a close association between the germ cell and

mesendodermal lineages in these crustaceans. This relationship is apparent beyond

Pancrustacea, since germ cells induced late in embryogenesis arise from the

mesoderm (Category 3). We thus speculate that germ cells in panarthropods may be

ancestrally derived from the mesodermal lineage, consistent with our inference that

inductive germ cell specification is ancestral to panarthropods (also discussed earlier).

Evolutionary shifts in the spatial and temporal origin of PGCs

We have seen significant variation in the spatial and temporal origin of PGCs

across panarthropods (Figs. 2–3; Table S1). However, regardless of when PGCs first

appear, we find a common stage across many panarthropods in which PGCs are

situated in bilateral segmental clusters across multiple trunk segments. Soon after this

stage, the segmental clusters on each side fuse together into bilaterally symmetrical

genital ridges, corresponding to the primordial embryonic gonad (Miya, 1958b; Nelsen,

1934; Seidel, 1924; Ullmann, 1964).

Most arthropods undergo "short germ" development, wherein the blastoderm

tissue generates only anterior segments, with posterior segments being added

sequentially to the posterior end of the germ band later in development (Krause, 1939).

In contrast, the "long germ" mode of development seen in *D. melanogaster*, wherein the

blastoderm tissue immediately generates all body segments, is an evolutionary novelty

found only in some insects (Clark et al., 2019; Krause, 1939). In Category 1 or 2

arthropods with short germ development, germ cells are often specified before the

formation of the segments that include the future embryonic gonad. In Category 2

insects, germ cells often associate with the posterior end of the germ band and then

migrate toward the anterior as the germ band elongates (Goss, 1953; Hegner, 1909b;

Heming and Huebner, 1994; Paz, 1958; Suzuki, 1990; Ullmann, 1964). This migratory

path is so common among insects that Johanssen and Butt included it in their

description of embryogenesis in a stereotypical insect (Johannsen and Butt, 1941).

The association of germ cells with the posterior end of the embryo may have had

other evolutionary consequences in insects. Many insects, both long and short germ,

specify the body plan early in development via maternal deposition of axial patterning

determinants, independent of the mechanism of germ cell specification (reviewed in

Lynch, 2019; Rosenberg et al., 2009). For example, the short germ grasshopper

Schistocerca americana exhibits asymmetric localization of the posterior patterning

determinant *nanos* in the absence of a germ plasm (Lall et al., 2003). However, insects

with germ plasm that includes axial patterning determinants, such as *D. melanogaster*,

often exhibit long germ development. Based on these observations, we employed

phylogenetic logistic regression and Pagel's test of evolutionary rate covariation (Pagel,

1994) to test the hypothesis that the acquisition of long germ band development

influences germ cell mechanism transitions or vice versa, and found statistically

significant evidence of this association using both methods (Figs. S3-S4; Table S8).

Overall, this suggests that long germ band development may drive the evolution of germ

plasm, which in turn may also drive transitions to long germ band development.

We further hypothesize that a spatial shift in germ cell specification to the

posterior end of the embryo in Category 2 insects created a condition in which germ line

determinants are localized to the same embryonic region as posterior patterning

determinants (Fig. 4B). Subsequently, germ line determinants may have hitchhiked onto

the asymmetric localization machinery, coupling germ cell specification to posterior

patterning and thus promoting the evolution of germ plasm (Fig. 4C). Under this

hypothesis, the mechanism that specifies the germ plasm of *D. melanogaster* would

represent an advanced stage of this evolutionary scenario, as its germ granules include

both germ line and abdominal patterning determinants (Ephrussi and Lehmann, 1992;

Trcek et al., 2015; Wang and Lehmann, 1991).

N. vitripennis is an intriguing case to study the relationship between A–P

patterning and germ cell specification. The posterior cytoplasm of this wasp has two

populations of *nanos* mRNA: one that is incorporated into the oosome and eventually

into pole cells, and another that functions in axial patterning (Lynch and Desplan, 2010).

The localization of *nanos* to the oocyte posterior in this wasp depends on actin-

dependent anchoring (Olesnicky and Desplan, 2007), as seen in *D. melanogaster*

(Forrest and Gavis, 2003), suggesting a conserved localization mechanism between

these two species. We speculate that the PGC specification mechanism used in *N.*

vitripennis might be an evolutionary intermediate between the mechanism used in

species where A–P patterning is entirely independent of germ cell formation (Fig. 4A),

and that used in species where these two processes are fully linked (Fig. 4C). Our

evolutionary hypothesis could be tested further by examining the mechanisms of

localization of posterior patterning determinants and germ line determinants in more

insect species to understand how existing cytoskeletal machinery for one

developmental process may have been coopted for use in another.

Potential genetic basis of evolutionary shifts in the molecular mechanism of PGC

specification

Beyond the timing and position of germ cell origin, evolutionary changes in the

localization of gene products and timing of gene expression may also underlie

transitions in the mechanism of specification (Extavour, 2007; Srouji and Extavour,

2010). Many of the genes important for specifying germ cells are conserved across

metazoans (Ewen-Campen et al., 2010), but they exhibit distinct temporal and spatial

expression patterns across species. In species that use inheritance, the germ line gene

products are expressed during oogenesis and persist in the early embryonic cytoplasm

to be inherited by the PGCs. In species that induce germ cells, these genes are either

not expressed in oocytes or the gene products are neither asymmetrically nor

preferentially inherited by PGCs. As we have suggested previously (Extavour, 2007;

Srouji and Extavour, 2010), relatively simple regulatory modifications affecting post-

transcriptional regulation, the cytoskeleton, or RNA or protein localization could cause

germ line gene product retention in oocytes, thereby enabling the transition from

induction to inheritance.

The insect-specific gene *oskar* offers a potential example of gene network

evolution driving shifts in germ cell specification mechanism. Since *oskar* is likely a

nucleator of germ plasm in multiple insects (Ephrussi et al., 1991; Jones and Macdonald,

2007; Kemph and Lynch, 2022; Lynch et al., 2011; Rivard et al., 2025) and has been

lost multiple times since its origin before the divergence of Pterogyta from Ametabola

(Blondel et al., 2021), we hypothesized that *oskar* gene presence in the genome might

be correlated with germ line specification mode in a given taxon (Fig. S3). A related

hypothesis, that the evolutionary advent of the *oskar* gene facilitated the evolution of

holometabolous germ plasm, was previously proposed based on the co-occurrence of

oskar transcript and germ plasm in *N. vitripennis* and *D. melanogaster*, as well as the

absence of both *oskar* homologs and germ plasm in *T. castaneum*, *A. mellifera*, *C.*

albipunctata, most Lepidoptera, and the hemimetabolous genomic resources available

at the time (Lynch et al., 2011). The results of applying phylogenetic logistic regression

to this hypothesis suggest a strong and statistically significant association between

oskar presence and maternal inheritance (Fig. S3; Table S9). We emphasize that this

observed association does not imply that *oskar* is necessary or sufficient for germ plasm

formation; indeed, there are insect species without *oskar* that nevertheless have germ

plasm and species with *oskar* that lack germ plasm (Fig. S3), as we discuss below.

In contrast to prior hypotheses from our group and others of a single origin of

germ plasm at the base of the Holometabola (Ewen-Campen et al., 2012; Ewen-

Campen et al., 2013b; Kemph and Lynch, 2022; Lynch et al., 2011), our current

likelihood and parsimony reconstruction of induction as ancestral to Holometabola (Fig.

5; Figs. S1–S2) implies that *oskar* has been independently recruited as a germ plasm

nucleator at least twice, in Hymenoptera (Lynch et al., 2011) and in Diptera (Ephrussi

and Lehmann, 1992). The strong association between *oskar* presence and germ plasm

(Fig. S3) implies that it may have been co-opted for germ plasm nucleation even more

times across insects. We note that such "gene reuse" is a common feature of

convergent evolution across multiple biological processes (Chan et al., 2010; Martin and

Orgogozo, 2013; Morris, 2009; Reed et al., 2011; Satterlee et al., 2024). Since *oskar* is

co-expressed with germ plasm components in nervous systems of hemimetabolous and

holometabolous insects (Ewen-Campen et al., 2012; Kulkarni et al., 2023; Xu et al.,

2013), we posit that independent acquisitions of germ plasm in distinct lineages may

have proceeded via parallel cooption of a pre-existing gene interaction network. Future

studies should aim to determine whether each independent origin of germ plasm in the

Holometabola arose by co-option of *oskar* to the germ line, or by *oskar*-independent

mechanisms of germ plasm assembly.

The results of our analysis (Fig. S3; Table S9) suggest that *oskar*'s functional

evolution may have contributed to a transition from induction to inheritance, but

precisely when and how *oskar* acquired a germ line role remains unclear. In the

genomes of species of Carabidae (ground beetles), a beetle family that lacks pole cells,

oskar orthologs have been identified, decoupling the presence of *oskar* from germ

plasm in Holometabola (Blondel et al., 2021; Kobayashi et al., 2013). Morphological

data suggest that thrips use germ plasm (Haga, 1985; Heming, 1979), and *oskar*

homologs have been found in this order (Blondel et al., 2021). If *oskar* plays a role in

thrips germ cell specification, it could support the entirely novel hypothesis that *oskar*

evolved its germ line determinant properties before the evolution of complete insect

metamorphosis.

The case for novel panarthropod germ plasm nucleators

Recruitment of *oskar* to the germ line is not the only mechanism explaining the

evolution of germ plasm in arthropods. The genome of the hemimetabolous pea aphid

Acyrthosiphon pisum lacks an *oskar* ortholog but their embryos have a germ plasm

containing Vasa and Nanos proteins (Chang et al., 2006; Chen et al., 2009; Lin et al.,

2014), which are asymmetrically localized by unknown mechanisms. Vasa is also

asymmetrically localized to a putative germ plasm in amphipod and decapod species

(Chen et al., 2021; Gupta and Extavour, 2013), which is likely to have evolved via

distinct mechanisms since *oskar* is an insect-specific gene (Blondel et al., 2021).

Outside of arthropods, germ plasm nucleators have convergently evolved in diverse

animal phyla (Aoki et al., 2016; Bontems et al., 2009; Kulkarni and Extavour, 2017; Lee

et al., 2020; Marlow and Mullins, 2008; Scholl et al., 2024). Similarly, we posit that germ

plasm nucleators other than *oskar* likely also exist across the panarthropods, awaiting

discovery.

Concluding remarks

By reinterpreting the meticulous work of generations of classical embryologists

through the lens of modern developmental genetics and an improved panarthropod

phylogeny, we have generated four major, novel and testable evolutionary hypotheses

about the mechanisms that specify the germ line. First, inductive mechanisms likely

specified PGCs ancestrally in Panarthropoda, and differentiation from the mesoderm is

the likely ancestral mode of PGC specification in Euarthropoda. Second, germ plasm

arose independently multiple times across Panarthropoda, in many different crustacean

and insect orders. Third, in contrast to the *Drosophila* paradigm of PGC specification via

germ plasm, induction was likely the ancestral PGC specification mechanism in the

Holometabola, and germ plasm likely arose independently in some lineages of flies,

fleas, beetles, wasps, ants, and moths. Finally, the *oskar* gene was likely co-opted as a

germ plasm nucleator at least twice independently in flies and wasps, and many

panarthropods are likely to use something other than *oskar* to nucleate their germ plasm.

Together, these hypotheses have implications for the genetic basis of convergence in

evolution that extend beyond the specific problem of germ cell formation. The advent of

new genetic tools in an increasing number of panarthropod model species places us in

an exciting era to test the new hypotheses we propose here, tackle age-old questions

about the evolution of panarthropod germ cell specification, and understand the impact

of these mechanisms of genome evolution and body patterning in the most successful

group of animals on earth.

Acknowledgements

We thank members of the Extavour Lab for helpful discussions, Kira Heikes for the image of tardigrade germ cells (Fig. 1B4), and Prashant Sharma for the image of harvestmen germ cells (Fig. 1B3).

Data Availability

All data sources are listed in the accompanying Supplementary Material and/or are available at

https://github.com/rishabhrajkapoor/panarthropoda_gc_specification_evolution, commit ID ff9945c.

Author Contributions

All authors gathered data and performed data analysis and interpretation. JAK, ELR and RRK wrote the first draft of the manuscript. CGE conceived of the study, obtained funding for the study, supervised its execution, and reviewed and edited the manuscript.

Funding

This study was supported by NSF award IOS-2220747, funds from Harvard University, and from the Howard Hughes Medical Institute awarded to CGE. Additional support was provided by the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard (award number #1764269), the Harvard Quantitative Biology Initiative, a Herchel Smith Graduate Fellowship awarded to ELR, and an NSF Graduate Research Training Fellowship to RRK. CGE is an Investigator of the Howard Hughes Medical Institute.

Conflicts of Interest

The authors declare no conflicts of interests.

 Figure Legends

Figure 1. Primordial germ cell (PGC) origin in the context of panarthropod
embryogenesis. In all schematics, A/a, anterior; P/p, posterior; D/d, dorsal; V/v,
ventral.lower case letters are shown in cases where egg and embryonic axes are not
the same, and/or indicate embryonic axes in embryos that do not occupy the entire egg
volume; if D/V is not indicated, schematics show ventral view; magenta, approximate
location of germ plasm or PGCs; gray, yolk or extraembryonic tissue; blue box in
schematic, stage/location where germ cells are detectable in the micrographs and
drawings on the right. (**A**) Description of Category 1: PGC specification at early
cleavage and blastoderm stages. (**A1**) In the fruit fly *Drosophila melanogaster*, germ
plasm accumulates at the posterior of syncytial embryos and is inherited by pole cells
that migrate to the ultimate site of the embryonic gonad (Illmensee and Mahowald,
1974). Right: TEM micrograph displaying pole cells budding in *D. melanogaster*
embryos. The pole cell nuclei (N) have characteristically diffuse chromatin, and polar
granules (P) are distributed throughout the cytoplasm. Image reproduced with
permission from (Mahowald, 1968). M: mitochondria; N: nuclei; P: polar granules. (**A2**)
In the amphipod *Parhyale hawaiensis*, germ plasm accumulates in a region of the
embryo called the RNA-containing body, which is inherited by the *g* micromere at the 8-
cell stage (Extavour, 2005; Gupta and Extavour, 2013). All descendants of *g* are PGCs.
Right: PGCs visualized by immunohistochemistry with a cross-reactive antibody raised
against recombinant Vasa protein from both *D. melanogaster* and the grasshopper
Schistocerca americana (Chang et al., 2002). (**B**) Description of Category 2: PGC
specification at gastrulation. (**B1**) In the milkweed bug *Oncopeltus fasciatus*, PGCs first
appear in the ventral posterior blastoderm in an invaginating structure called the
posterior pit (Ewen-Campen et al., 2013a). PGCs remain at the posterior tip of the germ
band as it invaginates into the yolk. The PGCs migrate along the germ band toward the
anterior to segments A4–A6. Right: Hand drawing of a light micrograph of a sectioned
late blastoderm stage *O. fasciatus* embryo showing germ cells in the posterior pit.
Image reproduced with permission from (Butt, 1949). gb, germ band; gc, germ cell. (**B2**)
In the mysid *Hemimysis lamornæ,* PGCs differentiate in the center of the blastopore
area and are internalized at gastrulation (Manton, 1928). Right: Drawing of sagittal view
of an *H. lamornæ* embryo at gastrulation. Image reproduced with permission from
(Manton, 1928). B, blastopore area; E, ectodermal teloblast; en, endoderm cell; G,
genital rudiment; M, mesodermal teloblast; m, head mesoderm band cell. (**B3**) In the
harvestman *Phalangium opilio*, PGCs are first identifiable at the posterior end of the
germ band soon after the germ band forms (Faussek, 1888; Gainett et al., 2022). PGCs
remain at the posterior end of the germ band at least until the fourth opisthosomal
segment is formed, after which the germ cells can be found in that segment. Right:
Chromatin (visualized with the nucleic acid dye Hoechst 33342) is more diffuse in PGCs
than in surrounding cells. Image kindly provided by Prashant Sharma (University of
Wisconsin, Madison). Green, alpha-tubulin. (**B4**) In the tardigrade *Hypsibius exemplaris*,
PGCs are first identifiable in the 32-cell stage and are the first cells to be internalized
during gastrulation. Right: PGCs visualized by *in situ* hybridization against *He-vasa*.
Image kindly provided by Kira Heikes (Duke University). (**B5**) In the velvet worm
Peripatopsis moseleyi, PGCs are first identifiable near the blastopore at the beginning

of gastrulation. As the embryo elongates and more segments form, the PGCs migrate to
more anterior segments. Right: Hand drawing of a light micrograph of a *P. moseleyi*
embryo showing primordial germ cells soon after their internalization during gastrulation,
distinguished by their lighter-staining nuclei and cytoplasm. Image reproduced with
permission from (Manton, 1949). G, germ cells; en, endoderm; b.a., blastoporal area.
(**C**) Description of Category 3: PGC specification from mesoderm. (**C1**) In the cricket
Gryllus bimaculatus, PGCs are first detectable among the mesoderm in segments A2–
A4 at embryonic stage 6, after most abdominal segments have been specified (Ewen-
Campen et al., 2013b). Right: PGCs visualized by immunohistochemistry with an
antibody against *Gb*-Piwi. (**C2**) In the leptostracan *Nebalia bipes*, PGCs are first
identifiable in the mesoderm in the thorax and abdomen during the extended germ band
stage (Manton, 1934). Right: hand drawing of a light micrograph of a sectioned *N. bipes*
embryo showing PGCs among the mesoderm in the extended germ band, distinguished
by their lighter-staining cytoplasm and larger nuclei. Image reproduced with permission
from (Manton, 1934). G, genital rudiment; l.c.m., circular muscle round liver lobe; th.8,
eighth thoracic segment. (**C3**) In the house spider *Parasteatoda tepidariorum*, PGCs are
first identifiable in segmental clusters in the opisthosomal mesoderm after all segments
have formed (Schwager et al., 2015). Right: PGCs visualized by immunohistochemistry
using an antibody raised against *Pt*-Vasa.
951
Figure 2. Germ cell specification across Hexapoda. Phylogeny of studied hexapod
lineages indicating the PGC origin timing category and the inferred mechanism of PGC
specification. Phylogenetic relationships in the cladogram are derived from published
relationships listed in Table S2; branch lengths are not to scale. Data on the timing of
germ cell specification are shown in the first column as follows: dark purple, Category 1
(cleavage/blastoderm); medium purple, Category 2 (gastrulation); light purple, Category
3 (mesoderm differentiation). Data on the mechanism of germ cell specification, inferred
as described in Supplementary Information, are shown in the second column: dark pink,
experimental evidence supports the use of germ plasm; light pink, morphological
evidence of germ plasm; blue, experimental evidence of induction. Data and data
sources are described in more detail in Table S1.
963
Figure 3. Panarthropod germ cell specification outside of Hexapoda. Phylogeny of
studied non-hexapod panarthropod lineages indicating the PGC origin timing category
and the inferred mechanism of PGC specification. Phylogenetic relationships in the
cladogram are derived from published relationships listed in Table S2; branch lengths
are not to scale. Data on the timing of germ cell specification are shown in the first
column as follows: dark purple, Category 1 (cleavage/blastoderm); medium purple,
Category 2 (gastrulation); light purple, Category 3 (mesoderm differentiation). Data on
the mechanism of germ cell specification, inferred as described in Supplementary
Information, are shown in the second column: dark pink, experimental evidence
supports the use of germ plasm; light pink, morphological evidence of germ plasm; blue,
experimental evidence of induction. Data and data sources are described in more detail
in Table S1.
976

Figure 4. Hypothesized links between spatial and temporal dynamics of PGC
origination and the mechanism of germ cell specification. (**A**) In species that induce
PGCs later in embryogenesis (Category 3), the asymmetrically localized anterior–
posterior (A–P) patterning determinants (blue) and germ cell determination factors
(pink) are spatially and temporally separate. (**B**) Proposed shift to earlier germ cell
specification at the embryonic posterior in Category 2 insects enabled overlap between
the patterning and PGC determinants. (**C**) Subsequent hitchhiking of the asymmetric
localization machinery by germ line determinants (indicated by mixture of blue and pink
at the embryo posterior) could have coupled PGC specification and posterior patterning
and thus promoted the evolution of germ plasm. Anterior is left and ventral is facing the
reader in all schematics.
988
Figure 5. Evolutionary dynamics of panarthropod germ cell specification.
Phylogenetic relationships among all study species (Table S1) are derived from
published relationships listed in Table S2; branch lengths are not to scale. Additional
taxonomic information is provided in Fig. S1. Colored lines to the right of each branch in
the cladogram designate the mechanism of germ cell specification (inferred as
described in Supplementary Materials) as follows: pink, germ plasm; blue, induction.
Pink asterisks (*) denote proposed instances of germ plasm evolution (based on
maximum parsimony) and pink pluses (+) mark ambiguous cases of germ plasm gain.
Blue asterisks (*) denote proposed instances of germ plasm loss and blue pluses (+)
mark ambiguous cases of germ plasm loss.
999

References

1000

1001

Abbassy, M. M., Helmy, N., Osman, M., Cope, S. E. and Presley, S. M. (1995a).
Embryogenesis of the sand fly Phlebotomus papatasi (Diptera: Psychodidae): Cell
cleavage, blastoderm formation, and gastrulation. *Ann. Entomol. Soc. Am.* **88**, 809–
814.

Abbassy, M. M., Helmy, N., Osman, M., Cope, S. E. and Presley, S. M. (1995b).
Embryogenesis of the sand fly Phlebotomus papatasi (Diptera: Psychodidae):
Organogenesis including segmentation, blastokinesis, mouthparts, and alimentary
canal. *Ann. Entomol. Soc. Am.* **88**, 815–820.

Abbassy, M. M., Helmy, N., Osman, M., Cope, S. E. and Presley, S. M. (1995c).
Embryogenesis of the sand fly Phlebotomus papatasi (Diptera: Psychodidae):
Organogenesis of the nervous system, tracheal system, muscular system, heart,
and gonad rudiments. *Ann. Entomol. Soc. Am.* **88**, 821–826.

Achtelig, M. and Krause, G. (1971). Experimente am ungefurchten Ei von Pimpla
turionellae L. (Hymenoptera) zur Funktionsanalyse des Oosombereichs. *Wilhelm
Roux Arch. Dev. Biol.* **167**, 164–182.

Ajidagba, P., Pitts, C. W. and Bay, D. E. (1983). Early embryogenesis in the stable
fly (Diptera: Muscidae). *Ann. Entomol. Soc. Am.* **76**, 616–623.

Alexeeva, N. and Tamberg, Y. (2021). Anatomical changes in postembryonic
development of Pycnogonum litorale. *J. Morphol.* **282**, 329–354.

Alexeeva, N., Tamberg, Y. and Shunatova, N. (2018). Postembryonic
development of pycnogonids: A deeper look inside. *Arthropod Struct. Dev.* **47**, 299–
317.

Alwes, F. and Scholtz, G. (2004). Cleavage and gastrulation of the euphausiacean
Meganyctiphanes norvegica (Crustacea, Malacostraca). *Zoomorphology* **123**, 125–
137.

Alwes, F. and Scholtz, G. (2014). The early development of the onychopod
cladoceran Bythotrephes longimanus (Crustacea, Branchiopoda). *Front. Zool.* **11**,
10.

Amy, R. L. (1961). The embryology of Habrobracon juglandis (Ashmead). *J.
Morphol.* **109**, 199–217.

Anderson, D. T. (1962). The embryology of Dacus tryoni (Frogg.) [Diptera,
Trypetidae (= Tephritidae)], the Queensland fruit-fly. *Development* **10**, 248–292.

Anderson, D. T. (1962b). The Epigenetics of the Larva in Diptera. *Acta Zool.* **43**,
221–228.

Anderson, D. T. (1967). Larval development and segment formation in the
branchiopod crustaceans Limnadia stanleyana King (Conchostraca) and Artemia
salina (L.) (Anostraca). *Aust. J. Zool.* **15**, 47–91.

Anderson, D. T. (1973). *Embryology and Phylogeny in Annelids and Arthropods.*
Oxford, UK: Pergamon Press.

Anderson, D. T. and Wood, E. C. (1968). The morphological basis of embryonic
movements in the light brown apple moth, Epiphyas postvittana (Walk.)
(Lepidoptera: Torticidae). *Aust. J. Zool.* **16**, 763–793.

Ando, H. (1962). *The comparative embryology of Odonata with special reference to
a relic dragonfly, Epiophlebia superstes Selys.* Tokyo: Japan Society for the
Promotion of Science.

Ando, H. and Tanaka, M. (1980). Early embryonic development of the primitive
moths, Endoclyta signifier Walker and E. excrescens Butler (Lepidoptera:
Hepialidae). *Int. J. Ins. Morph. Embryol.* **9**, 67–77.

Ansari, S., Troelenberg, N., Dao, V. A., Richter, T., Bucher, G. and Klingler, M.
(2018). Double abdomen in a short-germ insect: Zygotic control of axis formation
revealed in the beetle Tribolium castaneum. *Proc. Natl. Acad. Sci.* **115**, 1819–1824.

Aoki, S. T., Kershner, A. M., Bingman, C. A., Wickens, M. and Kimble, J. (2016).
PGL germ granule assembly protein is a base-specific, single-stranded RNase. *Proc
Natl Acad Sci U S A* **113**, 201524400.

Auten, M. (1934). The early embryological development of Phormia regina: Diptera
(Calliphoridae). *Ann. Entomol. Soc. Am.* **27**, 481–506.

Ayers, H. (1884). On the development of Oecanthus niveus, and its parasite,
Teleas. *Mem. Boston Soc. Nat. Hist.* **3**, 225–282.

Baker, C. M., Buckman-Young, R. S., Costa, C. S. and Giribet, G. (2021).
Phylogenomic analysis of velvet worms (Onychophora) uncovers an evolutionary
radiation in the neotropics. *Mol. Biol. Evol.* **38**, 5391–5404.

Baldass, F. von (1937). Entwicklung von Holopedium gibberum. *Zool. Jahrb. Anat.*
63, 399–454.

**Ballesteros, J. A., Santibáñez-López, C. E., Baker, C. M., Benavides, L. R.,
Cunha, T. J., Gainett, G., Ontano, A. Z., Setton, E. V. W., Arango, C. P., Gavish-
Regev, E., et al.** (2022). Comprehensive Species Sampling and Sophisticated

Algorithmic Approaches Refute the Monophyly of Arachnida. *Mol Biol Evol* **39**,
msac021.

Berg, G. J. and Gassner, G. (1978). Fine structure of the blastoderm embryo of the
pink bollworm, Pectinophora gossypiella (Saunders) (Lepidoptera: Gelechiidae). *Int.*
J. Insect Morphol. Embryol. **7**, 81–105.

Bernot, J. P., Owen, C. L., Wolfe, J. M., Meland, K., Olesen, J. and Crandall, K.
A. (2023). Major revisions in pancrustacean phylogeny and evidence of sensitivity to
taxon sampling. *Mol. Biol. Evol.* **40**, msad175.

Biffis, C., Alwes, F. and Scholtz, G. (2009). Cleavage and gastrulation of the
dendrobranchiate shrimp Penaeus monodon (Crustacea, Malacostraca, Decapoda).
Arthropod Struct. Dev. **38**, 527–540.

Blaimer, B. B., Santos, B. F., Cruaud, A., Gates, M. W., Kula, R. R., Mikó, I.,
Rasplus, J.-Y., Smith, D. R., Talamas, E. J., Brady, S. G., et al. (2023). Key
innovations and the diversification of Hymenoptera. *Nat. Commun.* **14**, 1212.

Blondel, L., Besse, S., Rivard, E. L., Ylla, G. and Extavour, C. G. (2021).
Evolution of a cytoplasmic determinant: Evidence for the biochemical basis of
functional evolution of the novel germ line regulator Oskar. *Mol. Biol. Evol.* **38**, 5491–
5513.

Bontems, F., Stein, A., Marlow, F., Lyautey, J., Gupta, T., Mullins, M. C. and
Dosch, R. (2009). Bucky Ball Organizes Germ Plasm Assembly in Zebrafish. *Curr.*
Biol. **19**, 414–422.

Bounoure, L. (1939). *L'origine des cellules reproductrices et le problème de la*
lignée germinale. Paris, France: Gauthier-Villars.

Brauer, A. (1894). Beiträge zur Kenntnis der Entwicklungsgeschichte des
Skorpions. *Z wiss. Zool.* **57**, 402–432.

Brauer, A. (1925). Studies on the embryology of Bruchus quadrimaculatus, Fabr.
Ann. Entomol. Soc. Am. **18**, 283–312.

Brauer, A. (1949). Localization of presumptive areas in the blastoderm of the pea
beetle Callosobruchus maculatus Fabr., as determined by ultraviolet (2537 Å)
irradiation injury. *J. Exp. Zool.* **112**, 165–193.

Bronskill, J. F. (1959). Embryology of Pimpla turionellae (L.) (Hymenoptera:
Ichneumonidae). *Can. J. Zool.* **37**, 655–688.

Buchta, T., Özüak, O., Stappert, D., Roth, S. and Lynch, J. A. (2013). Patterning
the dorsal–ventral axis of the wasp Nasonia vitripennis. *Dev Biol* **381**, 189–202.

Bull, A. L. (1982). Stages of Living Embryos in the Jewel Wasp Mormoniella
(Nasonia) vitripennis (Walker) (Hymenoptera: Pteromalidae). *Int. J. Insect Morphol.*
Embryol. **11**, 1–23.

Buss, L. W. (1982). Somatic cell parasitism and the evolution of somatic tissue
compatibility. *Proc. Natl. Acad. Sci.* **79**, 5337–5341.

Buss, L. W. (1983). Evolution, development, and the units of selection. *Proc. Natl.*
Acad. Sci. **80**, 1387–1391.

Butt, F. H. (1934). Embryology of Sciara (Sciaridae: Diptera). *Ann. Entomol. Soc.*
Am. **27**, 565–579.

Butt, F. H. (1936). The early embryological development of the parthenogenetic
alfalfa snout beetle, Brachyrhinus ligustici L. *Ann. Entomol. Soc. Am.* **29**, 1–14.

Butt, F. H. (1949). Embryology of the Milkweed Bug, Oncopeltus fasciatus
(Hemiptera). *Cornell Experiment Station Memoir* 3–43.

Cannon, H. G. (1921). The Early Development of the summer egg of a Cladoceran
(Simocephalus vetulus). *J. Cell Sci.* **s2-65**, 627–642.

Cannon, H. G. (1924). On the development of an estherid crustacean. *Philos.*
Trans. R. Soc. Lond. Ser. B **212**, 395–430.

Carter, J.-M., Baker, S. C., Pink, R., Carter, D. R., Collins, A., Tomlin, J., Gibbs,
M. and Breuker, C. J. (2013). Unscrambling butterfly oogenesis. *BMC Genomics*
14, 283.

Carter, J.-M., Gibbs, M. and Breuker, C. J. (2015). Divergent RNA Localisation
Patterns of Maternal Genes Regulating Embryonic Patterning in the Butterfly
Pararge aegeria. *PLoS ONE* **10**, e0144471.

Cavallin, M. (1970). Développement embryonnaire de l'appareil génital chez le
phasme, Carausius morosus Br. *Bull. Biol. Fr. Belg.* **104**, 343–66.

Cavallin, M. (1971). ["Substitutive polyembryony" and the problem of germinal line
origin in the phasma Carausius morosus Br]. *C. R. Hebd. des seances l'Acad. des*
Sci. Ser. D: Sci. Nat. **272**, 462–5.

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S.
D., Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., et al. (2010).
Adaptive Evolution of Pelvic Reduction in Sticklebacks by Recurrent Deletion of a
Pitx1 Enhancer. *Science* **327**, 302–305.

Chang, C., Dearden, P. and Akam, M. (2002). Germ Line Development in the Grasshopper Schistocerca gregaria: vasa As a Marker. *Dev. Biol.* **252**, 100–118.

Chang, C., Lee, W.-C., Cook, C. E., Lin, G.-W. and Chang, T. (2006). Germ-plasm specification and germline development in the parthenogenetic pea aphid Acyrthosiphon pisum: Vasa and Nanos as markers. *Int. J. Dev. Biol.* **50**, 413–421.

Chang, C., Lin, G., Cook, C. E., Horng, S., Lee, H. and Huang, T. (2007). Apvasa marks germ-cell migration in the parthenogenetic pea aphid Acyrthosiphon pisum (Hemiptera: Aphidoidea). *Dev. Genes Evol.* **217**, 275–287.

Chang, H., Qiu, Z., Yuan, H., Wang, X., Li, X., Sun, H., Guo, X., Lu, Y., Feng, X., Majid, M., et al. (2020). Evolutionary rates of and selective constraints on the mitochondrial genomes of Orthoptera insects with different wing types. *Mol. Phylogenet. Evol.* **145**, 106734.

Chaw, R. C., Vance, E. and Black, S. D. (2007). Gastrulation in the spider Zygiella x-notata involves three distinct phases of cell internalization. *Dev. Dynam.* **236**, 3484–3495.

Chen, R. P.-Y., Shih, C.-L., Lin, G.-W., Cook, C. E., Huang, T.-Y. and Chang, C.-C. (2009). Developmental expression of Apnanos during oogenesis and embryogenesis in the parthenogenetic pea aphid Acyrthosiphon pisum. *Int. J. Dev. Biol.* **53**, 169–176.

Chen, Y., Fang, X., Tian, X.-Q., Cui, Z., Feng, H.-Y. and Qiu, G.-F. (2021). Germ plasm and the origin of the primordial germ cells in the oriental river prawn, Macrobrachium nipponense. *Cell Tissue Res.* **386**, 559–569.

Choban, R. G. and Gupta, A. P. (1972). Meiosis and early embryology of Blissus leucopterus hirtus, Montandon (Heteroptera: Lygaeidae). *Int. J. Insect Morphol. Embryol.* **1**, 301–314.

Church, N. S. and Rempel, J. G. (1971). The embryology of Lytta viridana LeConte (Coleoptera: Meloidae). VI. The appendiculate, 72-h embryo. *Can. J. Zool.* **49**, 1563–1570.

Clark, E., Peel, A. D. and Akam, M. (2019). Arthropod segmentation. *Development* **146**, dev170480.

Claypole, A. M. (1898). The embryology and oögenesis of Anurida maritima (Guér.). *J. Morphol.* **14**, 219–300.

Colonnetta, M. M., Goyal, Y., Johnson, H. E., Syal, S., Schedl, P. and Deshpande, G. (2022). Preformation and epigenesis converge to specify primordial germ cell fate in the early Drosophila embryo. *PLoS Genet.* **18**, e1010002.

Copilaş-Ciocianu, D., Borko, Š. and Fišer, C. (2020). The late blooming amphipods: Global change promoted post-Jurassic ecological radiation despite Palaeozoic origin. *Mol. Phylogenet. Evol.* **143**, 106664.

Counce, S. J. (1963). Developmental morphology of polar granules in Drosophila. Including observations on pole cell behavior and distribution during embryogenesis. *J. Morphol.* **112**, 129–145.

Crother, B. I., White, M. E. and Johnson, A. D. (2007). Inferring developmental constraint and constraint release: Primordial germ cell determination mechanisms as examples. *J Theor Biol* **248**, 322–330.

Crother, B. I., White, M. E. and Johnson, A. D. (2016). Diversification and Germ-Line Determination Revisited: Linking Developmental Mechanism with Species Richness. *Frontiers Ecol Evol* **4**, 26.

Dannenberg, L. C. and Seaver, E. C. (2018). Regeneration of the germline in the annelid Capitella teleta. *Dev. Biol.* **440**, 74–87.

daSilva, A. F., Machado, L. C., Paula, M. B. de, Vieira, C. J. da S. P., Bronzoni, R. V. de M., Santos, M. A. V. de M. and Wallau, G. L. (2020). Culicidae evolutionary history focusing on the Culicinae subfamily based on mitochondrial phylogenomics. *Sci. Rep.* **10**, 18823.

Davis, C. (1967). A comparative study of larval embryogenesis in the mosquito Culex fatigans Wiedemann (Diptera⬚: Culicidae) and the sheep-fly, Lucilia sericata Meigen (Diptera⬚: Calliphoridae). *Aust. J. Zool.* **15**, 547–579.

Dearden, P. K. (2006). Germ cell development in the Honeybee (Apis mellifera); Vasa and Nanos expression. *BMC Dev. Biol.* **6**, 6.

Dearden, P., Grbic, M. and Donly, C. (2003). Vasa expression and germ-cell specification in the spider mite Tetranychus urticae. *Dev. Genes Evol.* **212**, 599–603.

Demotte, P., Panchaksaram, M., Kumarasinghe, H., Ly-Trong, N., Reis, M. dos and Minh, B. Q. (2025). IQ2MC: A New Framework to Infer Phylogenetic Time Trees Using IQ-TREE 3 and MCMCTree with Mixture Models. *EcoEvoRxiv* **10.32942/x2cd2x**, 1–30.

Dogiel, V. (1913). Embryologische Studien an Pantopoden. *Z. wiss. Zool.* **107**, 575–741.

Dohle, W. (1964). Die Embryonalentwicklung von Glomeris marginata (Villers) im Vergleich zur Entwicklung anderer Diplopoden. *Zool. Jahrb. Anat.* **81**, 241–310.

Donnell, D. M., Corley, L. S., Chen, G. and Strand, M. R. (2004). Caste
determination in a polyembryonic wasp involves inheritance of germ cells. *Proc. Nat.*
Acad. Sci. **101**, 10095–10100.

Donoughe, S., Nakamura, T., Ewen-Campen, B., Green, D. A., Henderson, L.
and Extavour, C. G. (2014). BMP signaling is required for the generation of
primordial germ cells in an insect. *Proc. Natl. Acad. Sci.* **111**, 4133–4138.

Doyle, D. A., Burian, F. N., Aharoni, B., Klinder, A. J., Menzel, M. M., Nifras, G.
C. C., Shabazz-Henry, A. L., Palma, B. U., Hidalgo, G. A., Sottolano, C. J., et al.
(2023). Germ Granule Evolution Provides Mechanistic Insight into Drosophila
Germline Development. *Mol. Biol. Evol.* **40**, msad174.

Du, S., Tihelka, E., Yu, D., Chen, W.-J., Bu, Y., Cai, C., Engel, M. S., Luan, Y.-X.
and Zhang, F. (2024). Revisiting the four Hexapoda classes: Protura as the sister
group to all other hexapods. *Proc. Natl. Acad. Sci.* **121**, e2408775121.

DuBois, A. M. (1932). A contribution to the embryology of Sciara (Diptera). *J.*
Morphol. **54**, 161–195.

Eastham, L. (1931). The Embryology of Pieris rapae - Organogeny. *Phil. Trans.*
Roy. Soc. B **219**, 1–50.

Eddy, E. M. (1975). Germ Plasm and the Differentiation of the Germ Cell Line.
International Review of Cytology **43**, 229–280.

Eddy, S. (2023). HMMER User's Guide: Biological sequence analysis using profile
hidden Markov models. http://eddylab.org/software/hmmer/Userguide.pdf

Edgar, A., Bates, C., Larkin, K. and Black, S. (2015). Gastrulation occurs in
multiple phases at two distinct sites in Latrodectus and Cheiracanthium spiders.
EvoDevo **6**, 33.

Edgar, R. C. (2004). MUSCLE: a multiple sequence alignment method with reduced
time and space complexity. *BMC Bioinform.* **5**, 1–19.

Elmer, K. R. and Clobert, J. (2025). Dollo's law of irreversibility in the post-genomic
age. *Trends Ecol. Evol.* **40**, 136–146.

Ephrussi, A. and Lehmann, R. (1992). Induction of germ cell formation by oskar.
Nature **358**, 387–392.

Ephrussi, A., Dickinson, L. K. and Lehmann, R. (1991). oskar organizes the germ
plasm and directs localization of the posterior determinant nanos. *Cell* **66**, 37–50.

Evans, R. (1901). On the Malayan Species of Onychophora. Part II - The
Development of Eoperipatus weldoni. *Q. J. Microscop. Sci.* **45**, 41–88.

Evans, T., Wade, C. M., Chapman, F. A., Johnson, A. D. and Loose, M. (2014).
Acquisition of Germ Plasm Accelerates Vertebrate Evolution. *Science* **344**, 200–203.

Everett, N. B. (1944). The present status of the germ-cell problem in vertebrates.
Biol Rev **20**, 45–55.

Ewen-Campen, B., Donoughe, S., Clarke, D. N. and Extavour, C. G. (2013b).
Germ cell specification requires zygotic mechanisms rather than germ plasm in a
basally branching insect. *Curr. Biol.* **23**, 835–842.

Ewen-Campen, B., Jones, T. E. M. and Extavour, C. G. (2013a). Evidence against
a germ plasm in the milkweed bug Oncopeltus fasciatus, a hemimetabolous insect.
Biol. Open **2**, 556–568.

Ewen-Campen, B., Schwager, E. E. and Extavour, C. G. M. (2010). The molecular
machinery of germ line specification. *Mol. Reprod. Dev.* **77**, 3–18.

Ewen-Campen, B., Srouji, J. R., Schwager, E. E. and Extavour, C. G. (2012).
oskar Predates the Evolution of Germ Plasm in Insects. *Curr Biol* **22**, 2278–2283.

Extavour, C. G. (2005). The fate of isolated blastomeres with respect to germ cell
formation in the amphipod crustacean Parhyale hawaiensis. *Dev. Biol.* **277**, 387–
402.

Extavour, C. G. and Akam, M. (2003). Mechanisms of germ cell specification
across the metazoans: epigenesis and preformation. *Development* **130**, 5869–5884.

Extavour, C. G. M. (2007). Evolution of the bilaterian germ line: lineage origin and
modulation of specification mechanisms. *Integr. Comp. Biol.* **47**, 770–785.

Eyun, S.-I. (2017). Phylogenomic analysis of Copepoda (Arthropoda, Crustacea)
reveals unexpected similarities with earlier proposed morphological phylogenies.
BMC Evol. Biol. **17**, 23.

Faussek, V. (1888). Ueber die embryonale Entwicklung der Geschlechtsorgane bei
der Afterspinne (Phalangium). *Biol. Zentralbl.* **8**, 359–363.

Faussek, V. (1892). Zur Anatomie und Embryologie der Phalangiden. *Biol.*
Zentralbl. **12**, 1–8.

Fernández, R., Edgecombe, G. D. and Giribet, G. (2018). Phylogenomics
illuminates the backbone of the Myriapoda Tree of Life and reconciles morphological
and molecular phylogenies. *Sci. Rep.* **8**, 83.

Fernando, W. (1934). The early embryology of a viviparous psocid. *J. Cell Sci.* **s2-**
77, 99–119.

Fish, W. A. (1947a). Embryology of Lucilia sericata Meigen (Diptera: Calliphoridae)
Part I. Cell cleavage and early embryonic development. *Ann. Entomol. Soc. Am.* **40**,
15–28.

Fish, W. A. (1947b). Embryology of Lucilia sericata Meigen (Diptera: Calliphoridae)
Part II. The blastoderm, yolk cells and germ cells. *Ann. Entomol. Soc. Am.* **40**, 677–
687.

Fish, W. A. (1949). Embryology of Phaenicia sericata (Meigen) Diptera:
Calliphoridae Part III. The gastrular tube. *Ann. Entomol. Soc. Am.* **42**, 121–133.

Fish, W. A. (1952). Embryology of Phaenicia sericata (Meigen) (Diptera:
Calliphoridae). Part IV. The inner layer and mesenteron rudiments. *Ann. Entomol.*
Soc. Am. **45**, 1–22.

Fleig, R. and Sander, K. (1986). Embryogenesis of the honeybee Apis mellifera L.
(Hymenoptera: Apidae): An SEM Study. *Int. J. Insect Morphol. Embryol.* **15**, 449–
462.

Forrest, K. M. and Gavis, E. R. (2003). Live Imaging of Endogenous RNA Reveals
a Diffusion and Entrapment Mechanism for nanos mRNA Localization in Drosophila.
Curr. Biol. **13**, 1159–1168.

Gabriel, W. N., McNuff, R., Patel, S. K., Gregory, T. R., Jeck, W. R., Jones, C. D.
and Goldstein, B. (2007). The tardigrade Hypsibius dujardini, a new model for
studying the evolution of development. *Dev. Biol.* **312**, 545–559.

Gainett, G., Crawford, A. R., Klementz, B. C., So, C., Baker, C. M., Setton, E. V.
W. and Sharma, P. P. (2022). Eggs to long-legs: embryonic staging of the
harvestman Phalangium opilio (Opiliones), an emerging model arachnid. *Front. Zool.*
19, 11.

Gambrell, F. L. (1933). The embryology of the black fly, Simulium pictipes Hagen.
Ann. Entomol. Soc. Am. **26**, 641–471.

Gatenby, J. B. (1917). The embryonic development of Trichogramma evanescens,
Westw., monembryonic egg parasite of Donacia simplex, Fab. *J. Cell Sci.* **s2-62**,
149–187.

Gatenby, J. B. (1918). The segregation of the germ-cells in Trichogramma
evanescens. *J. Cell Sci.* **s2-63**, 161–174.

Geigy, R. (1931). Action de l'ultra-violet sur le pôle germinal dans l'oeuf de
Drosophila melanogaster (Castration et mutabilité). *Rev. Suisse Zool.* **38**, 187–285.

Gerberding, M. and Patel, N. H. (2004). Chapter 6. Gastrulation in Crustaceans:
germ layers and cell lineages. In *Gastrulation: From Cells to Embryo* (ed. Stern, C.),
p. Cold Spring Harbor Laboratory Press.

Gerberding, M., Browne, W. E. and Patel, N. H. (2002). Cell lineage analysis of the
amphipod crustacean Parhyale hawaiensis reveals an early restriction of cell fates.
Development **129**, 5789–5801.

Gilbert, S. (1997). *Embryology: Constructing the Organism.* Sunderland, MA, USA:
Sinauer Associates Inc.

Giribet, G. (2018). Current views on chelicerate phylogeny—A tribute to Peter
Weygoldt. *Zool. Anz.* **273**, 7–13.

Giribet, G. and Edgecombe, G. D. (2019). The Phylogeny and Evolutionary History
of Arthropods. *Curr. Biol.* **29**, R592–R602.

Goodrich, A. L. (1939). The origin and fate of the entoderm elements in the
embryogeny of Porcellio laevis Latr. and Armadillidium nasutum B.L. (Isopoda). *J.*
Morphol. **64**, 401–426.

Görg, I. (1959). Untersuchungen am Keim von Hierodula (Rhombodera) crassa
Giglio Tos, ein Beitrag zur Embryologie der Mantiden. *Dtsch. Entomol. Z.* **6**, 389–
450.

Goss, R. J. (1952). The early embryology of the book louse, Liposcelis divergens
Badonnel (Psocoptera; Liposcelidae). *J. Morphol.* **91**, 135–167.

Goss, R. J. (1953). The advanced embryology of the book louse, Liposcelis
divergens Badonnel (Psocoptera; Liposcelidae). *J. Morphol.* **92**, 157–205.

Gough, L. H. (1902). The development of Admetus pumilio, Koch: a contribution to
the embryology of the Pedipalps. *J. Cell Sci.* **s2-45**, 595–630.

Grattan, R. M., McCulloch, R. J., Sellars, M. J. and Hertzler, P. L. (2013).
Ultrastructure of putative germ granules in the penaeid shrimp Marsupenaeus
japonicus. *Arthropod Struct. Dev.* **42**, 153–164.

Grbic', M. (2003). Polyembryony in parasitic wasps: evolution of a novel mode of
development. *Int. J. Dev. Biol.* **47**, 633–642.

Green, J. E. and Akam, M. (2014). Germ cells of the centipede Strigamia maritima
are specified early in embryonic development. *Dev. Biol.* **392**, 419–430.

Grobben, C. (1879). Die Entwickelungsgeschichte der Moina rectirostris. *Arb. Zool.*
Ins. Univ. Wien **2**, 203–226.

Grobben, C. (1881). Die Entwicklungsgeschichte von Cetochilus septentrionalis
Goodsir. *Arb. Zool. Ins. Univ. Wien* **3**, 243–282.

Gupta, T. and Extavour, C. G. (2013). Identification of a putative germ plasm in the
amphipod Parhyale hawaiensis. *EvoDevo* **4**, 34.

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden,
J., Couger, M. B., Eccles, D., Li, B., Lieber, M., et al. (2013). De novo transcript
sequence reconstruction from RNA-seq using the Trinity platform for reference
generation and analysis. *Nat. Protoc.* **8**, 1494–1512.

Hafiz, H. A. (1935). The embryological development of Cheyletus eruditus (a mite).
Proc. Roy. Soc. Lond. Ser. B **117**, 174–201.

Haga, K. (1985). Oogenesis and embryogenesis of the idolothripine thrips,
Bactrothrips brevitubus (Thysanoptera, Phlaeothripidae). In *Recent Advances in*
Insect Embryology in Japan, pp. 45–106. Tsukuba: ISEBU.

Hammerschmidt, J. (1910). Beiträge zur Entwicklung der Phasmatiden. *Z. wiss.*
Zool. **95**, 221–242.

Harber, P. A. and Mutchmor, J. A. (1970). The early embryonic development of
Culiseta inornata (Diptera: Culicidae). *Ann. Entomol. Soc. Am.* **63**, 1609–1614.

Hargitt, G. T. (1919). Germ Cells of Coelenterates. VI. General considerations,
discussion, conclusions. *Journal of Morphology* **33**, 1–59.

Hegner, R. W. (1908). Effects of removing the germ-cell determinants from the eggs
of some chrysomelid beetles. Preliminary report. *Biol. Bull.* **16**, 19–26.

Hegner, R. W. (1909a). The effects of centrifugal force upon the eggs of some
chrysomelid beetles. *J. Exp. Zool.* **6**, 507–552.

Hegner, R. W. (1909b). The origin and early history of the germ-cells in some
chrysomelid beetles. *J. Morphol.* **20**, 231–296.

Hegner, R. W. (1911). Experiments with chrysomelid beetles. III. The effects of
killing parts of the eggs of Leptinotarsa decemlineata. *Biol. Bull.* **20**, 237–251.

Hegner, R. W. (1914). Studies on germ cells. I. The history of the germ cells in
insects with special reference to the Keimbahn-determinants. II. The origin and
significance of the Keimbahn-determinants in animals. *J. Morphol.* **25**, 375–509.

Heider, K. (1889). *Die Embryonalentwicklung von Hydrophilus piceus, L.* Jena: Verlag von Gustav Fischer.

Heikes, K. L., Game, M., Smith, F. W. and Goldstein, B. (2023). The embryonic origin of primordial germ cells in the tardigrade Hypsibius exemplaris. *Dev. Biol.* **497**, 42–58.

Hejnol, A. and Schnabel, R. (2005). The eutardigrade Thulinia stephaniae has an indeterminate development and the potential to regulate early blastomere ablations. *Development* **132**, 1349–1361.

Heming, B. S. (1979). Origin and fate of germ cells in male and female embryos of Haplothrips verbasci (Osborn) (Insecta, Thysanoptera, Phlaeothripidae). *J. Morphol.* **160**, 323–343.

Heming, B. S. and Huebner, E. (1994). Development of the germ cells and reproductive primordia in male and female embryos of Rhodnius prolixus Stål (Hemiptera: Reduviidae). *Can. J. Zool.* **72**, 1100–1119.

Hertzler, P. L. (2002). Development of the mesendoderm in the dendrobranchiate shrimp Sicyonia ingentis. *Arthropod Struct. Dev.* **31**, 33–49.

Hertzler, P. L. (2005). Cleavage and gastrulation in the shrimp Penaeus (Litopenaeus) vannamei (Malacostraca, Decapoda, Dendrobranchiata). *Arthropod Struct. Dev.* **34**, 455–469.

Heymons, R. (1895). *Die Embryonalentwickelung von Dermapteren und Orthopteren unter Besonderer Berücksichtigung der Keimblätterbildung.* Jena: Jena Verlag von Gustav Fischer.

Heymons, R. (1897). Entwicklungsgeschichtliche Untersuchungen an Lepisma saccharina L. *Z wiss Zool* 583–631.

Heymons, R. (1901). Die Entwicklungsgeschichte der Scolopender. *Zoologica* **33**, 1–244.

Hickman, V. (1936). The embryology of the Syncarid Crustacean, Anaspides tasmaniae. *Pap. Proc. R. Soc. Tasman.* **1936**, 1–36.

His, W. (1868). *Untersuchungen über die erste Anlage des Wirbelthierleibes: die erste Entwickelung des Hünchens im Ei.* Leipzig: F. C. W. Vogel.

Hodson, A. C. (1934). The origin and differentiation of the sex organs of Tribolium confusum Duval. *Ann. Entomol. Soc. Am.* **27**, 278–291.

Holm, Å. (1947). On the development of Opilio parietinus Deg. *Zool. bidr. Uppsala* 409–422.

Hopf, C., Viebahn, C. and Püschel, B. (2011). BMP signals and the transcriptional repressor BLIMP1 during germline segregation in the mammalian embryo. *Dev. Genes Evol.* **221**, 209.

Howard, R. J., Giacomelli, M., Lozano-Fernandez, J., Edgecombe, G. D., Fleming, J. F., Kristensen, R. M., Ma, X., Olesen, J., Sørensen, M. V., Thomsen, P. F., et al. (2022). The Ediacaran origin of Ecdysozoa: integrating fossil and phylogenomic data. *J. Geol. Soc.* **179**, jgs2021-107.

Huerta-Cepas, J., Serra, F. and Bork, P. (2016). ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic Data. *Mol. Biol. Evol.* **33**, 1635–1638.

Huettner, A. F. (1923). The origin of the germ cells in Drosophila melanogaster. *J. Morphol.* **2**, 385–423.

Hughes, T. E. (1950). The embryonic development of the mite Tyroglyphus farinae Linnaeus 1758. *Proc. Zool. Soc. London* **119**, 873–886.

Ikuta, K. and Makioka, T. (1994). Notes on the postembryonic development of the ovary in Argulus japonicus (Crustacea: Branchiura). *Proc. Arthropod. Embryol. Soc. Jpn.* **29**, 15–17.

Illmensee, K. and Mahowald, A. P. (1974). Transplantation of posterior polar plasm in Drosophila. Induction of germ cells at the anterior pole of the egg. *Proc. Natl. Acad. Sci.* **71**, 1016–1020.

Illmensee, K., Mahowald, A. P. and Loomis, M. R. (1976). The ontogeny of germ plasm during oogenesis in Drosophila. *Dev. Biol.* **49**, 40–65.

Ives, A. R. and Theodore, Jr. G. (2010). Phylogenetic logistic regression for binary dependent variables. *Syst. Biol.* **59**, 9–26.

Jazdowska-Zagrodzińska, B. (1966). Experimental studies on the role of "polar granules" in the segregation of pole cells in Drosophila melanogaster. *J. Embryol. Exp. Morph.* **16**, 391–9.

Jiménez-Guri, E., Wotton, K. R., Gavilán, B. and Jaeger, J. (2014). A staging scheme for the development of the moth midge Clogmia albipunctata. *PLoS ONE* **9**, e84422.

Johannsen, O. A. (1929). Some phases in the embryonic development of Diacrisia virginica fabr. (Lepidoptera). *J. Morphol.* **48**, 493–541.

Johannsen, O. A. and Butt, F. H. (1941). *Embryology of Insects and Myriapods:*
The developmental history of insects, centipedes, and millepedes from egg
deposition to hatching. New York: McGraw-Hill, Inc.

Johnson, A. D., Drum, M., Bachvarova, R. F., Masi, T., White, M. E. and Crother,
B. I. (2003). Evolution of predetermined germ cells in vertebrate embryos:
implications for macroevolution. *Evol Dev* **5**, 414–431.

Johnson, A. D., Richardson, E., Bachvarova, R. F. and Crother, B. I. (2011).
Evolution of the germ line–soma relationship in vertebrate embryos. *Reproduction*
141, 291–300.

Jones, J. R. and Macdonald, P. M. (2007). Oskar controls morphology of polar
granules and nuclear bodies in Drosophila. *Development* **134**, 233–236.

Juhn, J. and James, A. A. (2006). oskar gene expression in the vector mosquitoes,
Anopheles gambiae and Aedes aegypti. *Insect Mol. Biol.* **15**, 363–372.

Juhn, J., Marinotti, O., Calvo, E. and James, A. A. (2008). Gene structure and
expression of nanos (nos) and oskar (osk) orthologues of the vector mosquito, Culex
quinquefasciatus. *Insect Mol. Biol.* **17**, 545–552.

Jung, E. (1966). Untersuchungen am Ei des Speisebohnenkäfers Bruchidius
obtectus Say (Coleoptera). *Z. Morphol. Ökol. Tiere* **56**, 444–480.

Jura, C. (1967). Origin of germ cells and gonads formation in embryogenesis of
Tetrodontophora bielanensis (Collembola). *Acta Biol. Crac. Zool.* **10**, 97–103.

Kaczmarczyk, A. N. (2014). Germline maintenance and regeneration in the
amphipod crustacean, Parhyale hawaiensis. PhD Dissertation, University of
California, Berkeley.

Kao, J. A., Ewen-Campen, B. and Extavour, C. G. (2025). Divergence of germ
cell-less roles in germ line development across insect species. *Dev. Biol.* **525**, 93–
101.

Kautzsch, G. (1910). Über die Entwicklung von Agelena labyrinthica Clerck. II. Teil.
Zool. Jahrb. Anat. **30**, 535–602.

Kawahara, A. Y., Plotkin, D., Espeland, M., Meusemann, K., Toussaint, E. F. A.,
Donath, A., Gimnich, F., Frandsen, P. B., Zwick, A., Reis, M. dos, et al. (2019).
Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths.
Proc. Natl. Acad. Sci. **116**, 22657–22663.

Kelly, G. M. and Huebner, E. (1989). Embryonic development of the hemipteran
insect Rhodnius prolixus. *J. Morphol.* **199**, 175–196.

Kemph, A. and Lynch, J. A. (2022). Evolution of germ plasm assembly and
function among the insects. *Curr. Opin. Insect Sci.* **50**, 100883.

Kemph, A., Kharel, K., Tindell, S. J., Arkov, A. L. and Lynch, J. A. (2025). Novel
structure and composition of the unusually large germline determinant of the wasp
Nasonia vitripennis. *Mol. Biol. Cell* **36**, ar55.

Kessel, E. L. (1939). *The embryology of fleas.* Washington: Smithsonian Institution.

Kharel, K., Tindell, S. J., Kemph, A., Schmidtke, R., Alexander, E., Lynch, J. A.
and Arkov, A. L. (2024). Dynamic protein assembly and architecture of the large
solitary membraneless organelle during germline development in the wasp Nasonia
vitripennis. *Development* **151**, dev202877.

Khila, A. and Abouheif, E. (2008). Reproductive constraint is a developmental
mechanism that maintains social harmony in advanced ant societies. *Proc. Natl.*
Acad. Sci. **105**, 17884–17889.

Khila, A. and Abouheif, E. (2010). Evaluating the role of reproductive constraints in
ant social evolution. *Philos. Trans. R. Soc. B: Biol. Sci.* **365**, 617–630.

Kingsley, J. S. (1893). The embryology of Limulus. — part II. *J. Morphol.* **8**, 195–
268.

Kistler, K. E., Trcek, T., Hurd, T. R., Chen, R., Liang, F.-X., Sall, J., Kato, M. and
Lehmann, R. (2018). Phase transitioned nuclear Oskar promotes cell division of
Drosophila primordial germ cells. *eLife* **7**, e37949.

Klag, J. (1982). Germ line of Tetrodontophora bielanensis (Insecta, Collembola).
Ultrastructural study on the origin of primordial germ cells. *Development* **72**, 183–
195.

Klag, J. and Bilinski, S. (1993). Oosome formation in two ichneumonid wasps.
Tissue Cell **25**, 121–128.

Klag, J. and Światek, P. (1999). Differentiation of primordial germ cells during
embryogenesis of Allacma fusca (L.) (Collembola: Symphypleona). *Int. J. Insect*
Morphol. Embryol. **28**, 161–168.

Kobayashi, H. and Ando, H. (1985). Early Embryogenesis of Fireflies, Luciola
cruciata, L. lateralis and Hotaria parvula (Coleoptera, Lampyridae). *Recent*
Advances in Insect Embryology in Japan.

Kobayashi, Y. and Ando, H. (1984). Mesodermal organogenesis in the embryo of
the primitive moth, Neomicropteryx nipponensis Issiki (Lepidoptera,
Micropterygidae). *J. Morphol.* **181**, 29–47.

Kobayashi, Y., Niikura, K., Oosawa, Y. and Takami, Y. (2013). Embryonic
development of Carabus insulicola (Insecta, Coleoptera, Carabidae) with special
reference to external morphology and tangible evidence for the subcoxal theory. *J.*
Morphol. **274**, 1323–1352.

Kobayashi, Y., Watanabe, T. and Suzuki, H. (2006). Embryonic development of
the firefly Pyrocoelia rufa Olivier (Insecta: Coleoptera, Lampyridae), with special
reference to its hibernal diapause. *Proc. Arthropod. Embryol.Soc. Jpn.* **41**, 47–53.

Komatsu, S. and Kobayashi, Y. (2012). Embryonic development of a whirligig
beetle, Dineutus mellyi, with special reference to external morphology (Insecta:
Coleoptera, Gyrinidae). *J. Morphol.* **273**, 541–560.

Krause, G. (1939). Die Eitypen der Insekten. *Biol. Zentralbl.* **59**, 495–536.

Krause, J. B. (1947). The development of the gonads of the wood-eating beetle,
Passalus cornutus Fabricius. *Ann. Entomol. Soc. Am.* **40**, 172–202.

Kühn, A. (1913). Die Sonderung der Keimesbezirke in der Entwicklung der
Sommereier von Polyphemus pediculus de Geer. *Zool. Jahrb. Anat.* **35**, 243–340.

Kulkarni, A. and Extavour, C. G. (2017). Convergent evolution of germ granule
nucleators: A hypothesis. *Stem Cell Res.* **24**, 188–194.

Kulkarni, A., Ewen-Campen, B., Terao, K., Matsumoto, Y., Li, Y., Watanabe, T.,
Kao, J. A., Parhad, S. S., Ylla, G., Mizunami, M., et al. (2023). oskar acts with the
transcription factor Creb to regulate long-term memory in crickets. *Proc. Natl. Acad.*
Sci. **120**, e2218506120.

Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M.,
Stecher, G. and Hedges, S. B. (2022). TimeTree 5: An expanded resource for
species divergence times. *Mol. Biol. Evol.* **39**, msac174.

Küthe, H.-W. (1966). Das Differenzierungszentrum als selbstregulierendes
Faktorensystem für den Aufbau der Keimanlage im Ei von Dermestes frischi
(Coleoptera). *Wilhelm Roux Arch. Entwickl. Mech. Org.* **157**, 212–302.

Lall, S., Ludwig, M. Z. and Patel, N. H. (2003). Nanos Plays a Conserved Role in
Axial Patterning outside of the Diptera. *Curr. Biol.* **13**, 224–229.

Lane, R. L. (1977). A development investigation of the reproductive systems of
Armadillidium vulgare (Latreille) and Porcellionides pruinosus (Brandt) (Isopoda).
Crustaceana **33**, 237–248.

Larink, O. (1969). Zur Entwicklungsgeschichte von Petrobius brevistylis (Thysanura,
Insecta). *Helgoländer Wiss. Meeresunters.* **19**, 111–155.

Lasko, P. F. and Ashburner, M. (1990). Posterior localization of vasa protein
correlates with, but is not sufficient for, pole cell development. *Genes Dev.* **4**, 905–
921.

Lassmann, G. W. P. (1936). The early embryological development of Melophagus
ovinus L., with special reference to the development of the germ cells. *Ann. Entomol.*
Soc. Am. **29**, 397–414.

Lawson, K. A., Dunn, N. R., Roelen, B. A. J., Zeinstra, L. M., Davis, A. M.,
Wright, C. V. E., Korving, J. P. W. F. M. and Hogan, B. L. M. (1999). Bmp4 is
required for the generation of primordial germ cells in the mouse embryo. *Genes*
Dev. **13**, 424–436.

Lee, C.-Y. S., Putnam, A., Lu, T., He, S., Ouyang, J. P. T. and Seydoux, G.
(2020). Recruitment of mRNAs to P granules by condensation with intrinsically-
disordered proteins. *eLife* **9**, e52896.

Lehmann, R. (2016). Germ Plasm Biogenesis - An Oskar-Centric Perspective. *Curr.*
Top. Dev. Biol. **116**, 679–707.

Leptin, M. (2004). Chapter 7: Gastrulation in Drosophila. In *Gastrulation: From Cells*
to Embryo (ed. Stern, C.), p. New York, NY, USA: Cold Spring Harbor Laboratory
Press.

Letunic, I. and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to
the phylogenetic tree display and annotation tool. *Nucleic Acids Res.* **52**, W78–W82.

Lin, G., Cook, C. E., Miura, T. and Chang, C. (2014). Posterior localization of
ApVas1 positions the preformed germ plasm in the sexual oviparous pea aphid
Acyrthosiphon pisum. *EvoDevo* **5**, 18.

Lin, G., Cook, C. E., Miura, T. and Chang, C. (2014). Posterior localization of
ApVas1 positions the preformed germ plasm in the sexual oviparous pea aphid
Acyrthosiphon pisum. *EvoDevo* **5**, 18.

Lochab, A. K. and Extavour, C. G. (2017). Bone Morphogenetic Protein (BMP)
signaling in animal reproductive system development and function. *Dev. Biol.* **427**,
258–269.

Luginbill, P. (1953). A Contribution to the Embryology of the May Beetle. *Ann.*
Entomol. Soc. Am. **46**, 505–528.

Lynch, J. A. (2019). Evolution of maternal control of axial patterning in insects. *Curr.*
Opin. Insect Sci. **31**, 37–42.

Lynch, J. A. and Desplan, C. (2010). Novel modes of localization and function of
nanos in the wasp Nasonia. *Development* **137**, 3813–3821.

Lynch, J. A., Özüak, O., Khila, A., Abouheif, E., Desplan, C. and Roth, S. (2011).
The Phylogenetic Origin of oskar Coincided with the Origin of Maternally Provisioned
Germ Plasm and Pole Cells at the Base of the Holometabola. *PLoS Genet.* **7**,
e1002029.

Ma, K.-Y., Tian, X.-Q., Liu, Z.-Q. and Qiu, G.-F. (2019). Observations on the
embryonic development of the oriental river prawn Macrobrachium nipponense (De
Haan, 1849) (Decapoda: Caridea: Palaemonidae). *J. Crustac. Biol.* **39**, 261–266.

Maddison, W. P. and Maddison, D. R. (2025). *Mesquite: A modular system for*
phylogenetic workflows and analyses. Version 4.01.

Mahowald, A. P. (1962). Fine structure of pole cells and polar granules in
Drosophila melanogaster. *J. Exp. Zool.* **151**, 201–215.

Mahowald, A. P. (1968). Polar granules of Drosophila. II. Ultrastructural changes
during early embryogenesis. *J. Exp. Zool.* **167**, 237–261.

Mahowald, A. P. (1975). Ultrastructural Changes in the Germ Plasm during the Life
Cycle of Miastor (Cecidomyidae, Diptera). *Wilhelm Roux Arch. Entwickl. Mech. Org.*
176, 223–240.

Mahowald, A. P. (2001). Assembly of the Drosophila germ plasm. *Int. Rev. Cytol.*
203, 187–213.

Mahowald, A. P., Illmensee, K. and Turner, F. R. (1976). Interspecific
transplantation of polar plasm between Drosophila embryos. *J. Cell Biol.* **70**, 358–
373.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. and Zdobnov, E. M. (2021).
BUSCO update: Novel and streamlined workflows along with broader and deeper
phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.
Mol. Biol. Evol. **38**, 4647–4654.

Manton, S. M. (1928). On the Embryology of a Mysid Crustacean, Hemimysis
lamornae. *Philos. Trans. R. Soc. B.* **216**, 363–463.

Manton, S. M. (1934). On the embryology of the crustacean Nebalia bipes. *Philos.*
Trans. R. Soc. B. **223**, 163–238.

Manton, S. M. (1949). Studies on the Onychophora VII. The early embryonic stages
of Peripatopsis, and some general considerations concerning the morphology and
phylogeny of the Arthropoda. *Philos. Trans. R. Soc. B.* **233**, 483–580.

Marlow, F. L. and Mullins, M. C. (2008). Bucky ball functions in Balbiani body
assembly and animal–vegetal polarity in the oocyte and follicle cell layer in
zebrafish. *Dev. Biol.* **321**, 40–50.

Martin, A. and Orgogozo, V. (2013). The loci of repeated evolution: A catalog of
genetic hotspots of phenotypic variation. *Evolution* **67**, 1235–1250.

Mayer, G. and Tait, N. N. (2009). Position and development of oocytes in velvet
worms shed light on the evolution of the ovary in Onychophora and Arthropoda.
Zool. J. Linn. Soc. Lond. **157**, 17–33.

McClendon, J. F. (1907). On the development of parasitic copepods. Part II. *Biol.*
Bull. **12**, 53–88.

McGowan, J. (2024). *BUSCO_phylogenomics*.
https://github.com/jamiemcg/BUSCO_phylogenomics

Mellanby, H. (1935). The Early Embryonic Development of Rhodnius prolixus
(Hemiptera, Heteroptera). *Q. J. Microscop. Sci.* **309**, 71–90.

Meng, C. (1968). Strukturwandel und histochemische Befunde insbesondere am
Oosom während der Oogenese und nach der Ablage des Eies von Pimpla
turionellae L. (Hymenoptera, Ichneumonidae). *Wilhelm Roux' Arch. für*
Entwicklungsmechanik Org. **161**, 162–208.

Metcalfe, M. E. (1935). The germ-cell cycle in Phytophaga destructor Say. *J. Cell*
Sci. **s2-77**, 585–604.

Metschnikow, E. (1866). *Embryologische studien an insecten*. Leipzig, Germany:
Verlag von Wilhelm Engelmann.

Meusemann, K., Trautwein, M., Friedrich, F., Beutel, R. G., Wiegmann, B. M.,
Donath, A., Podsiadlowski, L., Petersen, M., Niehuis, O., Mayer, C., et al. (2020).
Are fleas highly modified Mecoptera? Phylogenomic resolution of Antliophora
(Insecta: Holometabola). *bioRxiv* 2020.11.19.390666.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D.,
Haeseler, A. von and Lanfear, R. (2020). IQ-TREE 2: New models and efficient
methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–
1534.

Misof, B., Liu, S., Meusemann, K., Peters, R. S., Donath, A., Mayer, C.,
Frandsen, P. B., Ware, J., Flouri, T., Beutel, R. G., et al. (2014). Phylogenomics
resolves the timing and pattern of insect evolution. *Science* **346**, 763–767.

Mitsumoto, H. and Makioka, T. (2002). Preliminary notes on the structure and
postembryonic development of gonads in two tadpole shrimps Triops longicaudatus
and T. numidicus (Crustacea: Branchiopoda, Notostraca). *Proc. Arthropod. Embryol.*
Soc. Jpn. **37**, 13–17.

Mitsumoto, H. and Makioka, T. (2003). Postembryonic development of gonads in
Triops numidicus (Grube) (Crustacea: Branchiopoda, Notostraca). *Proc. Arthropod.*
Embryol. Soc. Jpn. **38**, 9–12.

Miura, T., Braendle, C., Shingleton, A., Sisk, G., Kambhampati, S. and Stern, D.
L. (2003). A comparison of parthenogenetic and sexual embryogenesis of the pea
aphid Acyrthosiphon pisum (Hemiptera: Aphidoidea). *J. Exp. Zool. B* **295B**, 59–81.

Miya, K. (1953). The presumptive genital region at the blastoderm stage of the
silkworm egg. *J. Fac. Agr. Iwate Univ.* **1**, 223–227.

Miya, K. (1958). Studies on the Embryonic Development of the Gonad in the
Silkworm, Bombyx mori L. Part I. Differentiation of Germ Cells. *J. Fac. Agr. Iwate*
Univ. 436–467.

Miya, K. (1958a). Studies on the Embryonic Development of the Gonad in the
Silkworm, Bombyx mori L. Part I. Differentiation of Germ Cells. *J. Fac. Agr. Iwate*
Univ. 436–467.

Miya, K. (1958b). Studies on the Embryonic Development of the Gonad in the
Silkworm, Bombyx mori L. Part II. Formation of the Gonad. *J. Fac. Agr. Iwate Univ.*
126–151.

Miyakawa, K. (1974). The embryology of the caddisfly Stenopsyche griseipennis
MacLachlan (Trichoptera, Stenopsychidae): IV. Organogenesis: mesodermal
derivatives. *Kontyû* **42**, 451–466.

Miyazaki, K. and Makioka, T. (2012). Postembryonic development of the female
reproductive system in the pycnogonid, Propallene longiceps (Pycnogonida,
Callipallenidae). *Invertebr. Reprod. Dev.* **56**, 287–292.

Morris, S. C. (2009). The predictability of evolution: glimpses into a post-Darwinian
world. *Die Naturwissenschaften* **96**, 1313–37.

Nair, K. B. (1939). The reproduction, oogenesis and development of Mesopodopsis
orientalis Tatt. *Proc. Indian Acad. Sci. - Sect. B* **9**, 175–223.

Nair, K. B. (1949). The embryology of Caridina laevis Heller. *Proc. Indian Acad. Sci.*
- Sect. B **29**, 211–288.

Nakamura, T. and Extavour, C. G. (2016). The transcriptional repressor Blimp-1
acts downstream of BMP signaling to generate primordial germ cells in the cricket
Gryllus bimaculatus. *Development* **143**, 255–263.

Nakao, H. (1999). Isolation and characterization of a Bombyx vasa-like gene. *Dev.*
Genes Evol. **209**, 312–316.

Nakao, H. and Takasu, Y. (2019). Complexities in Bombyx germ cell formation
process revealed by Bm-nosO (a Bombyx homolog of nanos) knockout. *Dev. Biol.*
445, 29–36.

Nakao, H., Hatakeyama, M., Lee, J. M., Shimoda, M. and Kanda, T. (2006).
Expression pattern of Bombyx vasa-like (BmVLG) protein and its implications in
germ cell development. *Dev. Genes Evol.* **216**, 94–99.

Nakao, H., Matsumoto, T., Oba, Y., Niimi, T. and Yaginuma, T. (2008). Germ cell
specification and early embryonic patterning in Bombyx mori as revealed by nanos
orthologues. *Evol. Dev.* **10**, 546–554.

Needham, A. E. (1937). Some points in the development of Neomysis vulgaris. *J.*
Cell Sci. **s2-79**, 559–588.

Needham, A. E. (1942). The structure and development of the segmental excretory
organs of Asellus aquaticus (Linné). *J. Cell Sci.* **s2-83**, 205–243.

Nelsen, O. E. (1934). The segregation of the germ cells in the grasshopper,
Melanoplus differentialis (Acrididae; Orthoptera). *J. Morphol.* **55**, 545–575.

Nelson, J. A. (1915). *The Embryology of the Honey Bee.* Princeton: Princeton
University Press.

Nie, R.-E., Andújar, C., Gómez-Rodríguez, C., Bai, M., Xue, H.-J., Tang, M.,
Yang, C.-T., Tang, P., Yang, X.-K. and Vogler, A. P. (2020). The phylogeny of leaf
beetles (Chrysomelidae) inferred from mitochondrial genomes. *Syst. Entomol.* **45**,
188–204.

Nieuwkoop, P. D. and Sutasurya, L. A. (1979). *Primordial Germ Cells in the*
Chordates: embryogenesis and phylogenesis. Cambridge University Press.

Nieuwkoop, P. D. and Sutasurya, L. A. (1981). *Primordial Germ Cells in the*
Invertebrates: from epigenesis to preformation. Cambridge: Cambridge University
Press.

Niikura, K., Hirasawa, K., Inoda, T. and Kobayashi, Y. (2017). Embryonic
Development of a Diving Beetle, Hydaticus pacificus Aubé (Insecta: Coleoptera;

Dytiscidae): External Morphology and Phylogenetic Implications. *Proc. Arthropod.*
Embryol. Soc. Jpn. **48**, 19–32.

Noskiewicz, J. and Poluszyński, G. (1927). Embryologische Untersuchungen an
Strepsipteren. 1. Teil. Embryogenesis der Gattung Stylops Kirby. *Bull. Acad. Pol.*
Sci. Lett. B **6**, 1093–1227.

Okada, M., Kleinman, I. A. and Schneiderman, H. A. (1974). Restoration of fertility
in sterilized Drosophila eggs by transplantation of polar cytoplasm. *Dev. Biol.* **37**,
43–54.

Olesnicky, E. C. and Desplan, C. (2007). Distinct mechanisms for mRNA
localization during embryonic axis specification in the wasp Nasonia. *Dev Biol* **306**,
134–142.

OpenTreeOfLife, Redelings, B., Reyes, L. L. S., Cranston, K. A., Allman, J.,
Holder, M. T. and McTavish, E. J. (2019). Open Tree of Life Synthetic Tree (12.3).
Zenodo.

Otto, S. P. and Hastings, I. M. (1998). Mutation and selection within the individual.
Genetica **102–103**, 507.

Özhan-Kizil, G., Havemann, J. and Gerberding, M. (2009). Germ cells in the
crustacean Parhyale hawaiensis depend on Vasa protein for their maintenance but
not for their formation. *Dev. Biol.* **327**, 230–239.

Özpolat, B. D. (2024). Annelids as models of germ cell and gonad regeneration. *J.*
Exp. Zool. Part B **342**, 126–143.

Pagel, M. (1994). Detecting correlated evolution on phylogenies: a general method
for the comparative analysis of discrete characters. *Proc. Roy. Soc. London, Ser. B,*
Biol. Sci **255**, 37–45.

Paterson, N. F. (1931). A contribution to the embryological development of Euryope
terminalis Baly. *S. Afr. J. Sci.* **28**, 344–371.

Paterson, N. F. (1935). Observations on the Embryology of Corynodes pusis
(Coleoptera, Chrysomelidae). *J Cell Sci.* **s2-78**, 91–131.

Pawlak, J. B., Sellars, M. J., Wood, A. and Hertzler, P. L. (2010). Cleavage and
gastrulation in the Kuruma shrimp Penaeus (Marsupenaeus) japonicus (Bate): A
revised cell lineage and identification of a presumptive germ cell marker. *Dev.*
Growth Differ. **52**, 677–692.

Paz, J. de la (1958). Studies on the embryogeny of Oncopeltus fasciatus (Dallas)
(Hemiptera: Lygaeidae).

Phalle, B. de S. and Sullivan, W. (1996). Incomplete sister chromatid separation is
the mechanism of programmed chromosome elimination during early Sciara
coprophila embryogenesis. *Development* **122**, 3775–3784.

Pires-daSilva, A. and Sommer, R. J. (2003). The evolution of signalling pathways
in animal development. *Nat. Rev. Genet.* **4**, 39–49.

Poulson, D. F. (1947). The pole cells of Diptera, their fate and significance. *Proc.*
Natl. Acad. Sci. **33**, 182–4.

Poulson, D. F. (1950). Histogenesis, organogenesis, and differentiation in the
embryo of Drosophila melanogaster Meigen. In *Biology of Drosophila* (ed. Demerec,
M.), pp. 168–274. New York: Hafner Pub. Co.

Presser, B. D. and Rutschky, C. W. (1957). The embryonic development of the
corn earworm, Heliothis zea (Boddie) (Lepidoptera, Phalaenidae). *Ann. Entomol.*
Soc. Am. **50**, 133–164.

Qiu, G.-F., Chen, Y., Cui, Z. and Zhu, X.-L. (2013). Localization of germline marker
vasa homolog RNA to a single blastomere at early cleavage stages in the oriental
river prawn Macrobrachium nipponense: Evidence for germ cell specification by
preformation. *Gene* **513**, 53–62.

Quan, H. (2018). Evolution of insect germline specification. PhD Dissertation,
University of Illinois at Chicago.

Quan, H. and Lynch, J. A. (2016). The evolution of insect germline specification
strategies. *Curr. Opin. Insect Sci.* **13**, 99–105.

Quan, H., Arsala, D. and Lynch, J. A. (2019). Transcriptomic and functional
analysis of the oosome, a unique form of germ plasm in the wasp Nasonia
vitripennis. *BMC Biol.* **17**, 78.

Queller, D. C. (2000). Relatedness and the fraternal major transitions. *Philos. Trans.*
R. Soc. Lond. Ser. B: Biol. Sci. **355**, 1647–1655.

Rafiqi, Ab. M., Rajakumar, A. and Abouheif, E. (2020). Origin and elaboration of a
major evolutionary transition in individuality. *Nature* **585**, 239–244.

Raminani, L. N. and Cupp, E. W. (1975). Early embryology of Aedes aegypti (L.)
(Diptera: Culicidae). *Int. J. Insect Morphol. Embryol.* **4**, 517–528.

Reed, R. D., Papa, R., Martin, A., Hines, H. M., Counterman, B. A., Pardo-Diaz,
C., Jiggins, C. D., Chamberlain, N. L., Kronforst, M. R., Chen, R., et al. (2011).
optix drives the repeated convergent evolution of butterfly wing pattern mimicry.
Science **333**, 1137–41.

Rempel, J. G. and Church, N. S. (1969). The embryology of Lytta viridana Le
Conte (Coleoptera: Meloidae). V. The blastoderm, germ layers, and body segments.
Can. J. Zool. **47**, 1157–1171.

Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic
comparative methods (and other things). *PeerJ* **12**, e16505.

Revell, L. J. and Harmon, L. J. (2024). A discrete character evolution model for
phylogenetic comparative biology with Γ-distributed rate heterogeneity among
branches of the tree. *bioRxiv* 2024.05.25.595896.

Rivard, E. L., Srouji, J., Repouliou, A. and Extavour, C. G. (2025). Protein
sequence evolution underlies interspecies incompatibility of a cell fate determinant.
bioRxiv **10.1101/2025.08.02.668269**, 1–53.

Roonwal, M. L. (1937). Studies on the embryology of the African migratory locust,
Locusta migratoria migratorioides Reiche and Frm. (Orthoptera, Acrididae) II -
Organogeny. *Philos. Trans. R. Soc. Lond. B* **227**, 175–244.

Rosenberg, M. I., Lynch, J. A. and Desplan, C. (2009). Heads and tails: Evolution
of antero-posterior patterning in insects. *BBA* **1789**, 333–342.

Roth, S. (2004). Chapter 8: Gastrulation in Other Insects. In *Gastrulation: From*
Cells to Embryo (ed. Stern, C.), p. New York, NY, USA: Cold Spring Harbor
Laboratory Press.

Ryan, R. B. (1963). Contribution to the embryology of Coeloides brunneri
(Hymenoptera: Braconidae). *Ann. Entomol. Soc. Am.* **56**, 639–648.

Sabroux, R., Corbari, L. and Hassanin, A. (2023). Phylogeny of sea spiders
(Arthropoda: Pycnogonida) inferred from mitochondrial genome and 18S ribosomal
RNA gene sequences. *Mol. Phylogenet. Evol.* **182**, 107726.

Sagawa, K., Yamagata, H. and Shiga, Y. (2005). Exploring embryonic germ line
development in the water flea, Daphnia magna, by zinc-finger-containing VASA as a
marker. *Gene Expr. Patterns* **5**, 669–678.

Saito, S. (1937). On the development of the tusser, Antheraea pernyi Guèrin-
Meneville, with special reference to the comparative embryology of insects. *J. Fac.*
Agr. Hokkaido Imp. Univ. **40**, 35–109.

Saitou, M. and Yamaji, M. (2010). Germ cell specification in mice: signaling,
transcription regulation, and epigenetic consequences. *Reproduction* **139**, 931–942.

Santos, A. C. and Lehmann, R. (2004). Germ Cell Specification and Migration in
Drosophila and beyond. *Curr Biol* **14**, R578–R589.

Satterlee, J. W., Alonso, D., Gramazio, P., Jenike, K. M., He, J., Arrones, A.,
Villanueva, G., Plazas, M., Ramakrishnan, S., Benoit, M., et al. (2024).
Convergent evolution of plant prickles by repeated gene co-option over deep time.
Science **385**, eado1663–eado1663.

Schimkewitsch, V. M. (1903). Über die Entwicklung von Thelyphonus caudatus
(L.). *Zool. Anz.* **26**, 665–685.

Schimkewitsch, W. (1896). Studien über parasitische Copepoden. *Z. wiss. Zool.*
61, 339–362.

Scholl, A., Liu, Y. and Seydoux, G. (2024). Caenorhabditis elegans germ granules
accumulate hundreds of low translation mRNAs with no systematic preference for
germ cell fate regulators. *Development* **151**, dev202575.

Scholl, G. (1963). Embryologische Untersuchungen an Tanaidaceen (Heterotanais
oerstedi Kröyer). *Zool. Jahrb. Anat.* **80**, 500–554.

Scholtz, G. (1990). The formation, differentiation and segmentation of the post-
naupliar germ band of the amphipod Gammarus pulex L. (Crustacea, Malacostraca,
Peracarida). *Proc. R. Soc. Lond. B* **239**, 163–211.

Scholtz, G. (2000). Evolution of the nauplius stage in malacostracan crustaceans. *J.*
Zool. Syst. Evol. Res. **38**, 175–187.

Scholtz, G. and Wolff, C. (2002). Cleavage, gastrulation, and germ disc formation
of the amphipod Orchestia cavimana (Crustacea, Malacostraca, Peracarida).
Contrib. Zool. **71**, 9–28.

Scholtz, G. and Wolff, C. (2013). Chapter 4: Arthropod Embryology: Cleavage and
Germ Band Development. In *Arthropod Biology and Evolution* (ed. Minelli, A.), pp.
63–89.

Schrader, F. (1922). The sex ratio and oogenesis of Pseudococcus citri. *Z. für*
Indukt. Abstamm.- Vererbungslehre **30**, 163–182.

Schröder, R. (2006). vasa mRNA accumulates at the posterior pole during
blastoderm formation in the flour beetle Tribolium castaneum. *Dev. Genes Evol.* **216**,
277-283.

Schwager, E. E., Meng, Y. and Extavour, C. G. (2015). vasa and piwi are required
for mitotic integrity in early embryogenesis in the spider Parasteatoda tepidariorum.
Dev. Biol. **402**, 276–290.

Schwentner, M., Combosch, D. J., Nelson, J. P. and Giribet, G. (2017). A
Phylogenomic Solution to the Origin of Insects by Resolving Crustacean-Hexapod
Relationships. *Curr. Biol.* **27**, 1818-1824.e5.

Sedgwick, A. (1887). The development of the Cape species of Peripatus: Part III.
On the changes from stage A to stage F. *J. Cell Sci.* **s2-27**, 467–550.

Seervai, R. N. H. and Wessel, G. M. (2013). Lessons for inductive germline
determination. *Mol. Reprod. Dev.* **80**, 590–609.

Seidel, F. (1924). Die geschlechtsorgane in der Embryonalen Entwicklung von
Pyrrhocoris apterus L. *Z. Morphol. Ökol. Tiere* **1**, 429–506.

Shafiq, S. A. (1954). A Study of the Embryonic development of the Gooseberry
Sawfly, Pteronidea ribesii. *Q. J. Microscop. Sci.* **95**, 93–113.

Sharma, P. P. and Gavish-Regev, E. (2025). The Evolutionary Biology of
Chelicerata. *Annu. Rev. Entomol.* **70**, 143–163.

Shinji, G. O. (1919). Embryology of coccids, with especial reference to the formation
of the ovary, origin and differentiation of the germ cells, germ layers, rudiments of
the midgut, and the intracellular symbiotic organisms. *J. Morphol.* **33**, 72–167.

Singh, J. P. (1967). Early embryonic development of gonads in Labidura riparia
(Pallas) (Labiduridae: Dermaptera). *Agra Univ. J. Res.* **1621**, 67–76.

Song, N., Wang, M.-M., Huang, W.-C., Wu, Z.-Y., Shao, R. and Yin, X.-M. (2024).
Phylogeny and evolution of hemipteran insects based on expanded genomic and
transcriptomic data. *BMC Biol.* **22**, 190.

Sonnenblick, B. P. (1941). Germ cell movements and sex differentiation of the
gonads in the Drosophila embryo. *Proc. Natl. Acad. Sci.* **27**, 484–9.

Srouji, J. and Extavour, C. G. (2010). Redefining Stem Cells and Assembling
Germ Plasm: Key Transitions in the Evolution of the Germ Line. In *Key Innovations*
in Animal Evolution (ed. DeSalle, R.) and Schierwater, B.), pp. 360–397. Enfield,
NH: CRC Press.

Stanke, M., Steinkamp, R., Waack, S. and Morgenstern, B. (2004). AUGUSTUS:
a web server for gene finding in eukaryotes. *Nucleic Acids Res.* **32**, W309-12.

Stanley, M. S. M. and Grundmann, A. W. (1970). The Embryonic Development of
Tribolium confusum. *Ann. Entomol. Soc. Am.* **63**, 1248–1256.

Stern, C. D. (2004). *Gastrulation: From Cells to Embryo.* (ed. Stern, C. D.) New
York, USA: Cold Spring Harbor Laboratory Press.

Strömberg, J.-O. (1965). On the embryology of the isopod Idotea. *Ark. Zool.* **2**,
421–473.

Strömberg, J.-O. (1968). Segmentation and organogenesis in Limnoria lignorum
(Rathke) (Isopoda). *Ark. Zool.* **20**, 91.

Strömberg, J.-O. (1971). Contribution to the embryology of bopyrid isopods with
special reference to Bopyroides, Hemiarthrus, and Peseudione (Isopoda,
Epicaridea). *Sarsia* **47**, 1–46.

Strömberg, J.-O. (1972). Cyathura polita (Crustacea, Isopoda), some embryological
notes. *Bull. Mar. Sci.* **22**, 463–482.

Sucena, É., Vanderberghe, K., Zhurov, V. and Grbić, M. (2014). Reversion of
developmental mode in insects: evolution from long germband to short germband in
the polyembryonic wasp Macrocentrus cingulum Brischke. *Evol. Dev.* **16**, 233–246.

Sudler, M. T. (1899). The development of Penilia schmackeri Richard. *Proc. Boston
Soc. Nat. Hist.* **29**, 109–131.

Sun, X. and Cheng, J. (2023). Conflicts in mitochondrial phylogenomics of
Branchiopoda, with the first complete mitogenome of Laevicaudata (Crustacea:
Branchiopoda). *Curr. Issues Mol. Biol.* **45**, 820–837.

Suvorov, A., Kim, B. Y., Wang, J., Armstrong, E. E., Peede, D., D'Agostino, E.
R. R., Price, D. K., Waddell, P. J., Lang, M., Courtier-Orgogozo, V., et al. (2022).
Widespread introgression across a phylogeny of 155 Drosophila genomes. *Curr.
Biol.* **32**, 111-123.e5.

Suzuki, N. (1990). Embryology of the Mecoptera (Panorpidae, Panorpodidae,
Bittacidae and Boreidae). *Bull. Sugadaira Montane Res. Cent.* **11**, 1–87.

Suzuki, N., Shimizu, S. and Ando, H. (1981). Early embryology of the alderfly,
Sialis mitsuhashii Okamoto (Megaloptera: Sialidae). *Int. J. Insect Morphol. Embryol.*
10, 409–418.

Tadokoro, R., Sugio, M., Kutsuna, J., Tochinai, S. and Takahashi, Y. (2006).
Early Segregation of Germ and Somatic Lineages during Gonadal Regeneration in
the Annelid Enchytraeus japonensis. *Curr. Biol.* **16**, 1012–1017.

Takamura, K., Fujimura, M. and Yamaguchi, Y. (2002). Primordial germ cells
originate from the endodermal strand cells in the ascidian Ciona intestinalis. *Dev.
Genes Evol.* **212**, 11–18.

Tanaka, M. (1985). Early embryonic development of Amata fortunei (Lepidoptera,
Amatidae). In *Recent Advances in Insect Embryology in Japan*, pp. 139–155.
Tsukuba: ISEBU.

Tanaka, M. (1987). Differentiation and Behaviour of Primordial Germ Cells during
the Early Embryonic Development of Parnassius glacialis Butler, Luehdorfia japonica
Leech and Byasa (Atrophaneura) alcinous alcinous Klug (Lepidoptera: Papilionidae).
Recent Advances in Insect Embryology in Japan and Poland 255–266.

Taube, E. (1909). Beiträge zur Entwicklungsgeschichte der Euphausiden. I. Die
Furchung des Eies bis zur Gastrulation. *Z wiss Zool* **92**, 427–464.

Taube, E. (1915). Beiträge zur Entwicklungsgeschichte der Euphausiden. II. Von der
Gastrula bis zum Furciliastadium. *Z. wiss. Zool.* **114**, 577–656.

Terao, A. (1929). On the embryonic development of the spiny lobster, Panulirus
japonicus (v. Siebold). *Jpn. J. Zool.* **2**, 387–449.

Terao, A. and Cheng, T. Z. (1926). On the rudiments of the gonad in an isopod,
Ligyda exotica (Roux). *Annot. Zool. Japon.* **11**, 29–32.

Thorpe, J. A. T. (2024). Phylogenomics supports a single origin of terrestriality in
isopods. *Proc. Roy. Soc. B* **291**, 20241042.

Tiegs, O. W. (1940). The embryology and affinities of the Symphyla, based on a
study of Hanseniella agilis. *J. Cell Sci.* **s2-82**, 1–208.

Tiegs, O. W. (1947). The development and affinities of the Pauropoda, based on a
study of Pauropus silvaticus, Part I. *J. Cell Sci.* **s3-88**, 165–267.

Tiegs, O. W. and Murray, F. V. (1938). The Embryonic Development of Calandra
oryzae. *J. Cell Sci.* **s2-80**, 159–273.

Tihelka, E., Giacomelli, M., Huang, D.-Y., Pisani, D., Donoghue, P. C. J. and Cai,
C.-Y. (2020). Fleas are parasitic scorpionflies. *Palaeoentomology* **3**, 641–653.

Tomizuka, S. and Machida, R. (2015). Embryonic development of a collembolan,
Tomocerus cuspidatus Börner, 1909: With special reference to the development and
developmental potential of serosa (Hexapoda: Collembola, Tomoceridae). *Arthropod*
Struct. Dev. **44**, 157–172.

Toyama, K. (1902). Contributions to the Study of Silk-Worms. I. On the Embryology
of the Silk-Worms. *Bull. Coll. Agric. Tokyo Imp. Univ.* **5**, 73–118.

Trcek, T. and Lehmann, R. (2019). Germ granules in Drosophila. *Traffic* **20**, 650–
660.

Trcek, T., Grosch, M., York, A., Shroff, H., Lionnet, T. and Lehmann, R. (2015).
Drosophila germ granules are structured and contain homotypic mRNA clusters.
Nat. Commun. **6**, 7962.

Tsutsumi, K. and Machida, R. (2006). Embryonic development of a snakefly,
Inocellia japonica Okamoto: an outline (Insecta: Neuroptera, Raphidiodea). *Proc.*
Arthropod Embryol. Soc. Jpn. **41**, 37–45.

Uemiya, H. and Ando, H. (1991). Mesoderm formation in a springtail, Tomocerus
ishibashii Yosii (Collembola□: Tomoceridae). *Int. J. Insect Morphol. Embryol.* **20**,
283–290.

Ullmann, S. L. (1964). The origin and structure of the mesoderm and the formation
of the coelomic sacs in Tenebrio molitor L. [Insecta, Coleoptera]. *Philos. Trans. R.*
Soc. Lond., B **248**, 245–277.

Underwood, E. M., Caulton, J. H., Allis, C. D. and Mahowald, A. P. (1980).
Developmental fate of pole cells in Drosophila melanogaster. *Dev. Biol.* **77**, 303–
314.

Vincent, K. A. and Hertzler, P. L. (2018). Putative germ granular material is
inherited by the primordial germ cell in Marsupenaeus japonicus (Spence Bate,
1888) (Decapoda: Dendrobranchiata: Penaeidae). *J. Crustac. Biol.* **38**, 267–277.

Wang, C. and Lehmann, R. (1991). Nanos is the localized posterior determinant in
Drosophila. *Cell* **66**, 637–647.

Weismann, A. (1863). Die Entwickelung der Dipteren im Ei, nach Beobachtungen
an Chironomous spec., Musca vomitoria und Pulex canis. *Zeit. wiss. Zool.* **13**, 107–
220.

Wenck, W. (1914). Entwicklungsgeschichtliche Untersuchungen an Tardigraden
(Macrobiotus lacustris Duj.). *Zool. Jahrb. Anat.* **37**, 465–514.

West, J. A., Cantwell, G. E. and Shortino, T. J. (1968). Embryology of the House
Fly, Musca domestica (Diptera: Muscidae), to the Blastoderm Stage. *Ann. Entomol.*
Soc. Am. **61**, 13–17.

Weygoldt, P. (1960). Embryologische Untersuchungen an Ostrakoden: Die
Entwicklung von Cyprideis litoralis (G.S. Brady) (Ostracoda, Podocopa, Cytheridae).
Zool. Jahrb. Anat. **78**, 369–426.

Weygoldt, P. (1964). Vergleichend-embryologische Untersuchugen an
Pseudoscorpionen (Chelonethi). *Z. Morphol. Ökol. Tiere* **54**, 1–106.

Weygoldt, P. (1975). Untersuchungen zur Embryologie und Morphologie der Geißelspinne, Tarantula marginemaculata C. L. Koch (Arachnida, Amblypygi, Tarantulidae). *Zoomorphologie* **82**, 137–199.

Wheeler, W. M. (1893). A Contribution to Insect Embryology. *J. Morphol.* **8**, 1–160.

Whittle, C. A. and Extavour, C. G. (2016). Refuting the hypothesis that the acquisition of germ plasm accelerates animal evolution. *Nat. Commun.* **7**, 12637.

Whittle, C. A. and Extavour, C. G. (2017). Causes and evolutionary consequences of primordial germ-cell specification mode in metazoans. *Proc. Natl. Acad. Sci.* **114**, 5784–5791.

Wiegmann, B. M., Trautwein, M. D., Winkler, I. S., Barr, N. B., Kim, J.-W., Lambkin, C., Bertone, M. A., Cassel, B. K., Bayless, K. M., Heimberg, A. M., et al. (2011). Episodic radiations in the fly tree of life. *Proc. Natl. Acad. Sci.* **108**, 5690–5695.

Wiesmann, R. (1926). Entwicklung und Organogenese der Cölomblasen. In *Zur Kenntnis der Anatomie und Entwicklungsgeschichte der Stabheuschrecke Carausius morosus Br.*, pp. 124–321. Jena: Verlag von Gustav Fischer.

Witschi, E. (1934). On determinative cleavage and yolk formation in the harpactid copepod Tisbe furcata (Baird). *Biol. Bull.* **67**, 335–340.

Wolfe, J. M., Breinholt, J. W., Crandall, K. A., Lemmon, A. R., Lemmon, E. M., Timm, L. E., Siddall, M. E. and Bracken-Grissom, H. D. (2019). A phylogenomic framework, evolutionary timeline and genomic resources for comparative studies of decapod crustaceans. *Proc. R. Soc. B* **286**, 20190079.

Wolff, C. (2009). The embryonic development of the malacostracan crustacean Porcellio scaber (Isopoda, Oniscidea). *Dev. Genes Evol.* **219**, 545–564.

Wolff, C. and Scholtz, G. (2002). Cell Lineage, Axis Formation, and the Origin of Germ Layers in the Amphipod Crustacean Orchestia cavimana. *Dev. Biol.* **250**, 44–58.

Wolff, E. (1962). *L' origine de la lignée germinale chez les vertébrés et chez quelques groupes d'invertébrés*. Paris, France: Éditions Hermann.

Wong, T. K. F., Ly-Trong, N., Ren, H., Baños, H., Roger, A. J., Susko, E., Bielow, C., Maio, N. D., Goldman, N., Hahn, M. W., et al. (2025). IQ-TREE 3: Phylogenomic Inference Software using Complex Evolutionary Models. *EcoEvoRxiv* **10.32942/x2p62n**,.

Woodland, J. T. (1957). A contribution to our knowledge of lepismatid development.
J. Morphol. **101**, 523–577.

Woodworth, C. W. (1889). Studies on the embryological development of Euvanessa
antiopa. In *The Butterflies of the Eastern United States and Canada, with special*
reference to New England, pp. 95–104. Cambridge, MA: Published by the Author.

Wotton, K. R., Jiménez-Guri, E., Matheu, B. G. and Jaeger, J. (2014). A staging
scheme for the development of the scuttle fly Megaselia abdita. *PLoS ONE* **9**,
e84421.

Wray, D. L. (1937). The embryology of Calendra callosa Olivier; The Southern corn
billbug (Coleoptera, Rhynchophoridae). *Ann. Entomol. Soc. Am.* **30**, 361–409.

Xu, X., Brechbiel, J. L. and Gavis, E. R. (2013). Dynein-Dependent Transport of
nanos RNA in Drosophila Sensory Neurons Requires Rumpelstiltskin and the Germ
Plasm Organizer Oskar. *J. Neurosci.* **33**, 14791–14800.

Yajima, M. and Wessel, G. M. (2011). The DEAD-box RNA helicase Vasa functions
in embryonic mitotic progression in the sea urchin. *Development* **138**, 2217–2222.

Yang, Z. (1997). PAML: a program package for phylogenetic analysis by maximum
likelihood. *Bioinformatics* **13**, 555–556.

Yang, Z. (2007). PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol. Biol.*
Evol. **24**, 1586–1591.

Ying, Y. and Zhao, G.-Q. (2001). Cooperation of Endoderm-Derived BMP2 and
Extraembryonic Ectoderm-Derived BMP4 in Primordial Germ Cell Generation in the
Mouse. *Dev. Biol.* **232**, 484–492.

Ying, Y., Liu, X. M., Marble, A., Lawson, K. A. and Zhao, G. Q. (2000).
Requirement of Bmp8b for the generation of primordial germ cells in the mouse. *Mol.*
Endocrinol. **14**, 1053–63.

Yoon, Y., Klomp, J., Martin-Martin, I., Criscione, F., Calvo, E., Ribeiro, J. and
Schmidt-Ott, U. (2019). Embryo polarity in moth flies and mosquitoes relies on
distinct zold genes with localized transcript isoforms. *eLife* **8**, e46711.

Yu, D., Du, S., Wei, X., Zhu, J., Ding, Y., Hu, F., Liu, M. and Zhang, F. (2024).
Whole-genome-based phylogenetic analyses provide new insights into the evolution
of springtails (Hexapoda: Collembola). *Mol. Phylogenet. Evol.* **200**, 108169.

Zhang, S.-Q., Che, L.-H., Li, Y., Dan, L., Pang, H., Ślipiński, A. and Zhang, P.
(2018). Evolutionary history of Coleoptera revealed by extensive sampling of genes
and species. *Nat. Commun.* **9**, 205.

Zhurov, V., Terzin, T. and Grbic, M. (2004). Early blastomere determines embryo
proliferation and caste fate in a polyembryonic wasp. *Nature* **432**, 764–769.

Zissler, D. (1992). From egg to pole cells: Ultrastructural aspects of early cleavage
and germ cell determination in insects. *Microsc. Res. Tech.* **22**, 49–74.

Zissler, D. and Sander, K. (1973). The cytoplasmic architecture of the egg cell of
Smittia spec. (Diptera, Chironomidae). I. Anterior and Posterior Pole Regions.
Wilhelm Roux Arch. Entwickl. Mech. Org. **172**, 175–186.

2037

A. Category 1: PGC specification at blastoderm & early cleavage stages

A1 *Drosophila melanogaster* (Diptera)

A2 *Parhyale hawaiensis* (Amphipoda)

Vasa

B. Category 2: PGC specification at gastrulation

B1 *Oncopeltus fasciatus* (Hemiptera)

B2 *Hemimysis lamornae* (Mysida)

B3 *Phalangium opilio* (Opiliones)

nuclei
tubulin

B4 *Hypsibius exemplaris* (Tardigrada)

piwi
nuclei

B5 *Peripatopsis moseleyi* (Onychophora)

C. Category 3: PGC specification from mesoderm

C1 *Gryllus bimaculatus* (Orthoptera)

C2 *Nebalia bipes* (Leptostraca)

C3 *Parasteatoda tepidariorum* (Araneae)

Vasa
nuclei

Clade

| Tardigrada |
| Onychophora |
| Chelicerata |
| Myriapoda |
| Pancrustacea |

Arthropoda

Timing category
- Category 1
- Category 2
- Category 3

Evidence of mechanism
- Germ plasm (experimental)
- Germ plasm (observational)
- Induction (experimental)

A AP patterning determinants and germ cell determinants are spatially and temporally separate

B AP patterning determinants and germ cell determinants overlap spatially and temporally

C Germ plasm contains both AP patterning determinants and germ cell determinants

Inferred mechanism

Germ plasm

Induction

* Germ plasm gain

+ Germ plasm gain (ambiguous)

* Germ plasm loss

+ Germ plasm loss (ambiguous)

Tardigrada
Onychophora
Chelicerata
Myriapoda
Non-insect pancrustaceans
Collembola
Ametabola
Hemimetabola
Holometabola
Insecta